PHI-MIXING IMPLIES INTERLACED RHO-MIXING FOR MARKOV CHAINS

YITZCHAK SHMALO

Abstract. We prove that every phi-mixing Markov chain is interlaced rho-mixing, giving an affirmative answer to Question 5 in Bradley's 2005 survey. The chain may be time-inhomogeneous and need not be stationary; its state space may be any standard Borel space. With Bradley's normalization, we obtain

$$\rho^*(n) \le \min\{1, 2\sqrt{2\phi(n)}\},$$

and the stronger bound $\rho^*(n) \le 2\sqrt{2\phi(n)(1-2\phi(n))}$ when $\phi(n) \le 1/4$. The proof combines a transition-row estimate with Peyre's maximal-correlation tensorization under prefix conditioning. A Markov separator makes the conditional correlation matrix of an arbitrary interlacing a weighted path. We give a self-contained proof of the local channel estimate, including the passage to standard Borel spaces. An explicit six-state chain shows that the square-root dependence cannot be improved in a uniform coefficient comparison. A countably infinite reset chain shows that the corresponding loss of a factor two in the exponential rate is also unavoidable for a single stationary chain; the same chain has lower psi coefficients identically zero, so it lies outside the previously known sufficient condition. A symmetric two-state family separately shows first-order sharpness of the path amplification factor. The finite components of the argument are machine-checked in Lean 4.

1. Introduction and principal results

Uniform mixing and interlaced maximal correlation measure different aspects of temporal dependence. The first controls conditional probabilities across a past–future cut. The second permits two sets of observation times to alternate arbitrarily often. A bound for one cut therefore does not, by itself, give a bound uniform in the number of alternations.

The maximal correlation coefficient of two sigma-fields goes back to Hirschfeld, Gebelein and Rényi [16]; the rho-mixing condition for stationary sequences was introduced by Kolmogorov and Rozanov [11], and its interlaced variant, in which the two index sets may be interlaced, by Stein [19] and Rosenblatt [17]. Bradley's survey [5] and his monograph [6] collect the basic properties of these and the other strong mixing coefficients; for the interlaced coefficients of random fields see also [2, 12]. For Markov chains, Bradley proved that every lower psi-mixing chain is interlaced rho-mixing [3], and constructed a stationary countable-state chain that is rho-mixing but not interlaced rho-mixing [4]; even for reversible Harris recurrent chains, where rho-mixing is equivalent to a spectral gap, the interlaced condition is strictly stronger [7].

Bradley asks in Question 5 of [5, Section 3, p. 119] whether a phi-mixing Markov chain must be rho-star-mixing. The same question for arbitrary random sequences is Question 2 of [5, Section 2.3], where Bradley records his conjecture [3, p. 226] that the answer is affirmative. Since $\rho(n) \le 2\sqrt{\phi(n)}$ for every random sequence [10, Theorem 17.2.3], phi-mixing implies rho-mixing, and the content of the question is the passage from one past–future cut to arbitrarily interlaced index sets, which by [4] cannot be made from rho-mixing alone. We establish the implication in the full, not necessarily stationary Markov setting of [5, Section 3]. Our argument uses the Markov property to reduce every finite interlacing to an alternating sequence of blocks, and applies the prefix versions of Peyre's tensorization theorems [14, Theorems 3.3.1 and 3.3.10;

Date: 18 September 2026.

Key words and phrases. Markov chain, phi-mixing, interlaced maximal correlation, tensorization, Dobrushin coefficient.

Section 3.4(a)]. These tensorization results are existing ingredients; the application here is based on verifying their precise conditional hypotheses for arbitrary Markov interlacings.

Here and below, total variation is normalized by

$$\|\mu-\nu\|_{\mathrm{TV}}=\sup_{B}|\mu(B)-\nu(B)|. \tag{1.1}$$

For two sigma-fields $\mathcal{A},\mathcal{B}$, put

$$\phi(\mathcal{A},\mathcal{B})=\sup_{\substack{A\in\mathcal{A},\,\mathbb{P}(A)>0\\ B\in\mathcal{B}}}|\mathbb{P}(B\mid A)-\mathbb{P}(B)|, \tag{1.2}$$

$$\rho(\mathcal{A},\mathcal{B})=\sup_{f,g}|\operatorname{Corr}(f,g)|, \tag{1.3}$$

where f and g are nonconstant, square-integrable, real random variables measurable with respect to $\mathcal{A}$ and $\mathcal{B}$, respectively. If one of the centered L^2 spaces is zero, the coefficient is defined to be zero.

Let $X=(X_k)_{k\in\mathbb{Z}}$ be a Markov chain on a standard Borel space E. No common transition kernel, invariant distribution, or reversibility assumption is imposed. Write

$$\mathcal{F}_a^b=\sigma(X_k:a\le k\le b),\qquad \phi(n)=\sup_{j\in\mathbb{Z}}\phi(\mathcal{F}_{-\infty}^{j},\mathcal{F}_{j+n}^{\infty}).$$

For nonempty disjoint $S,T\subset\mathbb{Z}$, let

$$\operatorname{dist}(S,T)=\inf\{|s-t|:s\in S,\ t\in T\}.$$

The interlaced coefficient is

$$\rho^*(n)=\sup_{\operatorname{dist}(S,T)\ge n}\rho\big(\sigma(X_s:s\in S),\sigma(X_t:t\in T)\big). \tag{1.4}$$

Phi-mixing and rho-star-mixing mean, respectively, that these coefficients tend to zero as $n\to\infty$.

Theorem 1.1. *For every standard-Borel Markov chain indexed by $\mathbb{Z}$ and every $n\ge 1$,*

$$\rho^*(n)\le\min\{1,2\sqrt{2\phi(n)}\}. \tag{1.5}$$

More precisely, if $0\le\phi(n)\le 1/4$, then

$$\rho^*(n)\le 2\sqrt{2\phi(n)\big(1-2\phi(n)\big)}. \tag{1.6}$$

Consequently, every phi-mixing Markov chain is rho-star-mixing.

The supremum in (1.4) includes infinite observation sets and arbitrary numbers of alternations. Thus Theorem 1.1 is not a bounded-interlacing or finite-state assertion. It answers the Markov-chain question, not Question 2 of [5] for arbitrary stochastic sequences. One consequence is that the limit theorems available under interlaced rho-mixing, such as the central limit theorem of [1] and the invariance principles of [13, 20], apply to every phi-mixing Markov chain.

The square-root scale in (1.5) is necessary in two distinct senses.

Theorem 1.2. *The following stationary examples hold.*

(1) *For each $\varepsilon\in(0,1/2)$ there is an irreducible, aperiodic six-state chain such that*

$$\phi(1)=\varepsilon,\qquad \rho^*(1)=\sqrt{2\varepsilon},\qquad \phi(n)=\rho^*(n)=0\quad(n\ge 2).$$

(2) *For each $q\in(0,1)$ the stationary chain on $\{0,1,2,\dots\}$ with*

$$P(i,0)=1-q,\qquad P(i,i+1)=q$$

satisfies

$$\phi(n)=q^n,\qquad q^{n/2}\le\rho^*(n)\le\min\{1,2q^{n/2}\}.$$

In particular,

$$\lim_{n\to\infty}\frac{1}{n}\log\rho^*(n)=\frac{1}{2}\log q.$$

The six-state example proves optimality of the exponent in a universal comparison of the coefficients. The reset chain proves sharpness of the exponential-rate loss for one fixed stationary chain; its lower psi coefficients vanish identically (Remark 7.2), so it is not covered by the criterion of [3]. Neither assertion identifies the best universal leading constant in (1.5). A third example, treated in Section 8, shows that the factor two in the block-path estimate is separately sharp to first order when expressed in terms of a robust local maximal correlation.

Proof outline. A phi bound places almost every transition row within $\phi(n)$ of the future marginal. The diameter of the family of such rows is at most $2\phi(n)$. A channel of row diameter δ has maximal correlation at most $\sqrt{\delta}$, uniformly over its input distribution. This uniformity is essential: conditioning on earlier observation blocks changes the input distribution of an adjacent transition.

After merging observation times of the same color, the blocks alternate. Under the conditioning fields required by Peyre's theorems, adjacent blocks retain the original intervening transition kernel. Nonadjacent blocks are separated by an observed Markov state. The additional conditioning splits into a statistic of the past and a statistic of the future, so it preserves the conditional product law. The resulting matrix has the support of a path, uniformly in the number of blocks.

The finite components of the argument, the scalar bookkeeping behind Theorem 1.1, and the identities behind the three examples are machine-checked in Lean 4; Section 10 describes the development, which is available together with the source of this paper in the repository [18].

2. Channel estimates and full-measure row sets

For random elements U, V, we abbreviate $\rho(\sigma(U), \sigma(V))$ as $\rho(U, V)$. Conditional expectation gives the equivalent formula

$$\rho(U,V)^2 = \sup_{0 \neq h \in L^2_0(\mathrm{Law}(U))} \frac{\mathrm{Var}(\mathbb{E}[h(U) \mid V])}{\mathrm{Var}(h(U))}, \tag{2.1}$$

where $L^2_0(\lambda)$ denotes the centered square-integrable functions under λ. Indeed, for fixed centered $h(U)$, the Cauchy–Schwarz inequality bounds its covariance with a function of V by the norm of its conditional expectation; equality is obtained by taking that conditional expectation as the second function when it is nonzero.

If $U \longrightarrow V \longrightarrow W$ is a Markov chain, conditional expectation and (2.1) give the data-processing bound

$$\rho(U,W) \leq \rho(U,V)\rho(V,W). \tag{2.2}$$

In particular, passing through additional channels cannot increase maximal correlation. We will use the same statement under regular conditional probability laws.

2.1. **Uniform channel contraction.** Let K be a Markov kernel from a standard Borel space E to a standard Borel space F, and let $A \subset E$ be a Borel set. Define

$$\delta_A(K) = \sup_{x,x' \in A} \|K(x,\cdot) - K(x',\cdot)\|_{\mathrm{TV}}, \tag{2.3}$$

$$\kappa_A(K) = \sup_{\lambda:\lambda(A)=1} \rho_\lambda(U,V), \quad U \sim \lambda, \quad V \mid U = x \sim K(x,\cdot). \tag{2.4}$$

Here ρ_λ is the maximal correlation of the pair (U, V) whose joint law is $\lambda \otimes K$, and degenerate inputs contribute zero. For $A = E$ the quantity $\delta_E(K)$ is Dobrushin's coefficient of ergodicity [9]. These definitions specify the actual row set, rather than a product-essential supremum, because a conditional input distribution may be singular with respect to the unconditional input distribution.

Lemma 2.1 (Row diameter and maximal correlation)**.** *For every such kernel and row set,*

$$\kappa_A(K)^2 \leq \delta_A(K). \tag{2.5}$$

Proof. We first give a finite-state proof. Fix an input law λ, discard its zero-mass atoms, and set $\nu = \lambda K$. Discard also the zero-mass output atoms. The reverse kernel is

$$R(y,x) = \frac{\lambda(x)K(x,y)}{\nu(y)}.$$

The operator $T : h \mapsto \mathbb{E}[h(U) \mid V]$ is R, its adjoint is K, and $T^*T = KR$. Thus $M = KR$ is a positive semidefinite self-adjoint Markov operator on $L^2(\lambda)$. It preserves constants, hence leaves $L^2_0(\lambda)$ invariant, and its largest eigenvalue on $L^2_0(\lambda)$ is $\rho_\lambda(U,V)^2$; since $L^2_0(\lambda)$ is finite-dimensional, this eigenvalue has an eigenvector there.

Postprocessing contracts total variation, so

$$\delta(M) = \sup_{x,x'} \|K(x,\cdot)R - K(x',\cdot)R\|_{\mathrm{TV}} \le \delta(K).$$

For a real function h on a finite set, the definition of total variation implies

$$\operatorname{osc}(Mh) \le \delta(M)\operatorname{osc}(h), \qquad \operatorname{osc}(h) = \max h - \min h.$$

Apply this inequality to a nonconstant eigenvector for the largest centered eigenvalue $\theta \ge 0$. Since $\operatorname{osc}(h) > 0$, it gives $\theta \le \delta(M) \le \delta(K)$. This proves (2.5) for finite channels. The zero-dimensional centered space is immediate.

Now fix an arbitrary input law λ supported on A. For finite measurable partitions of U and V, the induced finite channel has rows obtained by first mixing the original rows within an input cell and then projecting to the output partition. Any two mixtures of rows supported on A are at total-variation distance at most $\delta_A(K)$: for probability measures α, β supported there and every measurable B,

$$\begin{aligned}\left|\int K(x,B)\,d\alpha(x) - \int K(x',B)\,d\beta(x')\right| &\le \iint |K(x,B) - K(x',B)|\,d\alpha(x)d\beta(x') \\ &\le \delta_A(K).\end{aligned}$$

Projection cannot increase this bound. The finite result therefore bounds the correlation of every pair of simple functions of U and V by $\sqrt{\delta_A(K)}$. Approximate arbitrary square-integrable functions by bounded simple functions and pass to the limit in covariance and variance. Finally take the supremum over λ. □

The inequality in Lemma 2.1 is a classical strong data-processing comparison: for finite channels it is the case of the chi-square divergence in the theorem of Cohen, Iwasa, Rautu, Ruskai, Seneta and Zbaganu that every f-divergence contraction coefficient is bounded by the Dobrushin coefficient [8]; see [15] for the channel-contraction perspective and its relation to maximal correlation. The proof above is included to make explicit both the normalization and the uniformity in the input law.

2.2. **A common row-good set from phi.** For $s < t$, choose a regular conditional law

$$K_{s,t}(x,\cdot) = \mathrm{Law}(X_t \mid X_s = x), \qquad \mu_s = \mathrm{Law}(X_s).$$

Lemma 2.2. *There is a Borel set $A_{s,t} \subset E$ of full μ_s-measure such that*

$$\|K_{s,t}(x,\cdot) - \mu_t\|_{\mathrm{TV}} \le \phi(t-s) \qquad (x \in A_{s,t}). \tag{2.6}$$

Consequently,

$$\delta_{A_{s,t}}(K_{s,t}) \le \min\{1, 2\phi(t-s)\}. \tag{2.7}$$

Proof. For a fixed Borel $B \subset E$, set $h_B(x) = K_{s,t}(x,B) - \mu_t(B)$. For each Borel D with $\mu_s(D) > 0$, the definition of $\phi(t-s)$ gives

$$\left|\frac{1}{\mu_s(D)}\int_D h_B\,d\mu_s\right| \le \phi(t-s).$$

Taking $D = \{h_B > \phi(t-s) + \eta\}$, or the corresponding negative level set, shows that $|h_B| \le \phi(t-s)$ almost everywhere.

A standard Borel sigma-field has a countable generating algebra $\mathcal{C}$. Intersect the full-measure Borel sets just obtained for $B \in \mathcal{C}$, and call the intersection $A_{s,t}$. For fixed $x \in A_{s,t}$, the class of Borel sets satisfying $|K_{s,t}(x, B) - \mu_t(B)| \le \phi(t-s)$ is a monotone class containing $\mathcal{C}$. It therefore contains every Borel set. This proves (2.6). The triangle inequality and the universal upper bound one for total variation give (2.7). □

Fix one such set $A_{s,t}$ for every integer pair $s < t$, and write

$$\kappa(n) = \sup_{t-s \ge n} \kappa_{A_{s,t}}(K_{s,t}), \qquad \Delta(n) = \sup_{t-s \ge n} \delta_{A_{s,t}}(K_{s,t}). \tag{2.8}$$

The coefficients $\phi(n)$ are nonincreasing. Lemmas 2.1 and 2.2 imply

$$\kappa(n)^2 \le \Delta(n) \le \min\{1, 2\phi(n)\}. \tag{2.9}$$

The auxiliary coefficients in (2.8) refer to the fixed row-good sets; the final bounds in terms of ϕ do not depend on those choices.

Remark 2.3 (Conditional support)**.** If H is any random element and $\lambda_H = \mathrm{Law}(X_s \mid H)$, then

$$\lambda_H(A_{s,t}^c) = \mathbb{P}(X_s \notin A_{s,t} \mid H) = 0 \quad \text{almost surely.}$$

No absolute continuity of λ_H with respect to μ_s is asserted or needed. When H consists of observations strictly before s, the Markov property also preserves the transition kernel $K_{s,t}$ under conditioning on H.

3. Separators and conditional correlation

For standard Borel random elements U, V, H, define the conditional, or subjective, maximal correlation by

$$\rho_H(U, V) = \operatorname*{ess\,sup}_h \rho(U, V; \mathbb{P}(\cdot \mid H = h)), \tag{3.1}$$

the essential supremum being taken with respect to the law of H. This is the subjective correlation level of [14, Section 3.1]. We also write $\rho_{\mathcal{H}}(U, V)$ when $\mathcal{H} = \sigma(H)$. All conditioning fields used below are generated by finite collections of standard Borel random elements. Statements about their conditional laws are always almost sure statements.

Conditional independence is not preserved under arbitrary additional conditioning. The following precise form is what the Markov argument needs.

Lemma 3.1 (Side-separated conditioning)**.** *Suppose A, B, R are standard Borel random elements and A and B are conditionally independent given R. If $C = c(A, R)$ and $D = d(B, R)$ are measurable random elements, then A and B are conditionally independent given (R, C, D).*

Proof. For almost every r, write the conditional law of (A, B) given $R = r$ as $\alpha_r \otimes \beta_r$. Under this product law, C is a statistic of the first coordinate and D a statistic of the second coordinate. Disintegrating each factor separately produces kernels $\alpha_{r,c}$ and $\beta_{r,d}$. The product kernel $\alpha_{r,c} \otimes \beta_{r,d}$ has the defining integral property of the conditional law of (A, B) given (R, C, D), first on rectangles and product test functions and then, by a monotone-class argument, on the full sigma-field. This is the desired factorization. □

3.1. **Alternating observation blocks.** Fix finite nonempty disjoint $S, T \subset \mathbb{Z}$ with $\mathrm{dist}(S, T) \ge n$. Order $S \cup T$ chronologically and merge successive selected times of the same color into maximal blocks. The resulting blocks

$$B_1 < B_2 < \cdots < B_m$$

alternate in color. A block need not be an interval of integers: it is a block in the ordered list of selected times. Denote the S-blocks, in chronological order, by $S_1, \ldots, S_I$, and the T-blocks by $T_1, \ldots, T_J$. Put

$$U_i = (X_k)_{k \in S_i}, \qquad V_j = (X_k)_{k \in T_j}, \qquad \mathcal{H}_{ij} = \sigma(U_{i'} : i' < i,\ V_{j'} : j' < j).$$

Proposition 3.2 (The prefix interface)**.** *For these block variables,*

$$\rho_{\mathcal{H}_{ij}}(U_i, V_j) \le \kappa(n)\mathbf{1}\{S_i, T_j \text{ are adjacent in the block order}\}. \tag{3.2}$$

More precisely, an adjacent boundary at times $s < t$ admits the bound $\kappa_{A_{s,t}}(K_{s,t})$.

Proof. Suppose first that S_i is immediately before T_j, and set $s = \max S_i$ and $t = \min T_j$. Then $t - s \ge n$. Every coordinate in $\mathcal{H}_{ij}$ is strictly before the earlier block S_i. Under almost every conditional law given this prefix, the Markov property gives

$$U_i \longrightarrow X_s \longrightarrow X_t \longrightarrow V_j. \tag{3.3}$$

Here X_s is the last coordinate of U_i and X_t the first coordinate of V_j; given X_t, the block V_j is a function of X_t and of coordinates after t, which are independent of the past given X_t. The transition in the middle remains $K_{s,t}$. Only the input law of X_s has changed, and Remark 2.3 shows that this conditional input law is supported on $A_{s,t}$. Definition (2.4) and data processing now bound the conditional correlation of U_i, V_j by $\kappa_{A_{s,t}}(K_{s,t}) \le \kappa(n)$. If T_j immediately precedes S_i, exchange the colors; the argument still runs forward in time and uses no reverse transition estimate.

Next suppose S_i is before T_j and they are not adjacent. The T-block immediately following S_i lies strictly between them and has index less than j. Choose r in that block. Then X_r is $\mathcal{H}_{ij}$-measurable, U_i is before r, and V_j is after r.

Conditional on X_r, the pre-r and post-r paths are independent. Split every other coordinate generating $\mathcal{H}_{ij}$ according to which side of r it occupies. Thus this conditioning consists of X_r, a statistic of the pre-r path, and a statistic of the post-r path. Lemma 3.1 shows that the two sides remain conditionally independent under $\mathcal{H}_{ij}$. In particular $\rho_{\mathcal{H}_{ij}}(U_i, V_j) = 0$. When T_j is before S_i, use the intervening S-block instead.

There are only finitely many pairs (i, j). The pairwise almost sure assertions can therefore be realized on one probability-one event for this fixed finite interlacing. No common null set over all possible interlacings is required. □

4. Tensorization on a path

We state the two existing tensorization inputs in the form used here. In [14, Theorems 3.3.1 and 3.3.10] the hypotheses are stated with respect to the sigma-metalgebra generated by the two families; the proofs there use each hypothesis only under the conditioning $\sigma(U_{i'} : i' < i,\ V_{j'} : j' < j)$, which is the form recorded in [14, Section 3.4(a)] and used below.

Theorem 4.1 (Peyre's prefix tensorization)**.** *Let $(U_i)_{i=1}^I, (V_j)_{j=1}^J$ be ordered finite families, and put $\mathcal{H}_{ij} = \sigma(U_{i'} : i' < i, V_{j'} : j' < j)$.*

(1) *If $\rho_{\mathcal{H}_{ij}}(U_i, V_j) \le e_{ij}$, where $E = (e_{ij})$ has nonnegative entries, then*

$$\rho((U_i)_i, (V_j)_j) \le \min\{1, \|E\|_{2\to 2}\}. \tag{4.1}$$

(2) *Pad both families by deterministic variables and index them by $\mathbb{Z}$. If $\rho_{\mathcal{H}_{ij}}(U_i, V_j) \le e(j - i)$, where $e : \mathbb{Z} \to [0, 1]$, then*

$$\arcsin \rho((U_i)_i, (V_j)_j) \le \min\left\{\sum_{z\in\mathbb{Z}} \arcsin e(z), \frac{\pi}{2}\right\}. \tag{4.2}$$

Only finite nonconstant families are needed in the second statement; padding adds no information to any prefix. This avoids any tail-field issue associated with general two-sided indexed families, and it is also the reduction used in the proof of [14, Theorem 3.3.10].

Define the nondecreasing function

$$\Psi(r) = \sin\left(\min\left\{2\arcsin r, \frac{\pi}{2}\right\}\right) = \begin{cases} 2r\sqrt{1-r^2}, & 0 \le r \le 1/\sqrt{2}, \\ 1, & 1/\sqrt{2} \le r \le 1. \end{cases} \tag{4.3}$$

Proposition 4.2 (Finite interlacing bounds)**.** *For a finite interlacing with m chronological blocks,*

$$\rho(X_S, X_T) \le \min\left\{1, 2\kappa(n)\cos\frac{\pi}{m+1}\right\}, \tag{4.4}$$

$$\rho(X_S, X_T) \le \Psi(\kappa(n)). \tag{4.5}$$

Here $X_S = (X_s)_{s\in S}$ and $X_T = (X_t)_{t\in T}$. More generally, form the bipartite matrix E whose entry at an adjacent boundary $s < t$ is $\kappa_{A_{s,t}}(K_{s,t})$ and whose other entries are zero. Then $\rho(X_S, X_T) \le \min\{1, \|E\|_{2\to 2}\}$.

Proof. Apply Proposition 3.2 and (4.1). The matrix for the uniform bound is $\kappa(n)$ times the bipartite adjacency matrix B of a path on m vertices. After grouping the two colors, the ordinary adjacency matrix of the path is

$$\begin{pmatrix} 0 & B \\ B^T & 0 \end{pmatrix}.$$

Its largest eigenvalue is $2\cos(\pi/(m+1))$: the vector with kth coordinate $\sin(k\pi/(m+1))$ is a positive eigenvector with that eigenvalue, as follows from the sine addition formula. The bipartite block matrix has operator norm $\|B\|$, proving (4.4). For nonuniform weights, monotonicity under entrywise nonnegative domination also gives the same uniform upper bound.

For (4.5), exchange the colors if necessary so that the chronological order starts with an S-block. Then U_i can be adjacent only to V_i and V_{i-1}. In (4.2), take

$$e(0) = e(-1) = \kappa(n), \qquad e(z) = 0 \quad (z \notin \{0, -1\}).$$

The prefix hypotheses hold by Proposition 3.2, and (4.3) is exactly the resulting bound. □

Remark 4.3. The support of the matrix, rather than a bound on the number of observations, is decisive. Each row and column has at most two nonzero entries. Already the elementary bound $\|E\| \le \sqrt{\|E\|_1\|E\|_\infty} \le 2\kappa(n)$ is uniform in the interlacing length and suffices for the mixing implication. The arcsine input is needed only for the sharper quantitative envelope.

5. Proof of the mixing implication

Lemma 5.1 (Finite-coordinate approximation)**.** *A correlation bound uniform over all finite $S, T \subset \mathbb{Z}$ with $\operatorname{dist}(S,T) \ge n$ extends to arbitrary such sets.*

Proof. Choose finite sets $S_k \uparrow S$, $T_k \uparrow T$. For centered square-integrable $f \in L^2(\sigma(X_s : s \in S))$ and $g \in L^2(\sigma(X_t : t \in T))$, put

$$f_k = \mathbb{E}[f \mid \sigma(X_s : s \in S_k)], \qquad g_k = \mathbb{E}[g \mid \sigma(X_t : t \in T_k)].$$

By the martingale convergence theorem these conditional expectations converge in L^2 to f, g. If the finite bound is b, it gives

$$|\mathbb{E}[f_k g_k]| \le b\|f_k\|_2\|g_k\|_2.$$

Passing to the limit and then taking the supremum over f, g proves the assertion. This argument also handles a degenerate finite approximation without dividing by its variance. □

Proof of Theorem 1.1. Proposition 4.2 and Lemma 5.1 give the robust-channel estimate

$$\rho^*(n) \le \Psi(\kappa(n)) \le \min\{1, 2\kappa(n)\}. \tag{5.1}$$

Insert (2.9) to obtain (1.5). If $\phi(n) \le 1/4$, then $\kappa(n) \le \sqrt{2\phi(n)} \le 1/\sqrt{2}$. Since Ψ is nondecreasing,

$$\rho^*(n) \le \Psi(\sqrt{2\phi(n)}) = 2\sqrt{2\phi(n)(1-2\phi(n))}.$$

This is (1.6). Finally, if $\phi(n) \to 0$, either bound tends to zero. □

Corollary 5.2 (Rate transfer)**.** *If $\phi(n) \le Ae^{-cn}$ for $A, c > 0$, then*

$$\rho^*(n) \le \min\{1, 2\sqrt{2A}\, e^{-cn/2}\}.$$

More generally, any bound $\phi(n) \le b_n$ gives $\rho^*(n) \le \min\{1, 2\sqrt{2b_n}\}$.

Corollary 5.3 (A one-lag criterion)**.** *If $\phi(n_0) < 1/8$ at some lag, then $\rho^*(n_0) < 1$. In Bradley's Markov-chain setting, $\rho^*(n)$ consequently decreases to zero exponentially fast.*

Proof. The strict inequality follows from (1.5). The final assertion is [5, Theorem 3.3(6)]. In particular, phi-mixing provides such a lag. The direct proof of Theorem 1.1 does not require this one-lag theorem. □

The sharper estimate (1.6) in fact yields $\rho^*(n_0) < 1$ whenever $\phi(n_0) < 1/4$. We have separated the simpler threshold in Corollary 5.3 because it uses only the finite matrix-norm tensorization. No necessity of either threshold is claimed.

6. An exact six-state square-root example

Fix $0 < \varepsilon < 1/2$. On

$$E = \{-1, +1\} \times \{-1, 0, +1\}$$

let $\pi = \mu \otimes \nu$, where μ is uniform on the two signs and

$$\nu(-1) = \nu(+1) = \varepsilon, \qquad \nu(0) = 1 - 2\varepsilon.$$

Write $x = (u, v)$ and define

$$K((u, v), (u', v')) = \pi(u', v')(1 + uv'). \tag{6.1}$$

Proposition 6.1. *The stationary chain with kernel* (6.1) *is irreducible and aperiodic. It satisfies*

$$\phi(1) = \varepsilon, \qquad \rho^*(1) = \sqrt{2\varepsilon}, \tag{6.2}$$

$$\phi(n) = 0, \qquad \rho^*(n) = 0 \quad (n \ge 2). \tag{6.3}$$

Moreover, on the full state space,

$$\kappa_E(K)^2 = \delta_E(K) = 2\phi(1) = 2\varepsilon. \tag{6.4}$$

Proof. The kernel is nonnegative because $uv' \in \{-1, 0, +1\}$. Its row sums equal one, since $\mathbb{E}_\pi V = 0$. Also

$$\sum_{u,v} \pi(u, v) K((u, v), (u', v')) = \pi(u', v')(1 + v' \mathbb{E}_\pi U) = \pi(u', v').$$

Thus π is invariant. Independence and centering of U, V under π imply

$$\begin{aligned} K^2((u, v), (\widetilde{u}, \widetilde{v})) &= \pi(\widetilde{u}, \widetilde{v}) \mathbb{E}_\pi[(1 + uV)(1 + U\widetilde{v})] \\ &= \pi(\widetilde{u}, \widetilde{v}). \end{aligned}$$

Hence $K^2 = \Pi$, where every row of Π equals π. All entries of K^2 are positive, so the finite chain is irreducible and aperiodic.

For every state x, the signed row difference from π has density uv' with respect to π. Therefore

$$\|K(x, \cdot) - \pi\|_{\mathrm{TV}} = \tfrac{1}{2} \mathbb{E}_\pi |V| = \varepsilon. \tag{6.5}$$

For a stationary Markov chain, the future given the past is obtained by mixing the rows at the present state and then applying the future continuation kernel. Total variation cannot increase under this continuation. Thus (6.5) bounds $\phi(1)$ above by ε. Equality follows by conditioning on $U_0 = 1$ and testing $V_1 = 1$, since the two probabilities are 2ε and ε. The identity $K^2 = \Pi$ gives past–future independence at every gap at least two, proving the assertions about ϕ.

To compute the interlaced coefficient, independently over $t \in \mathbb{Z}$ sample pairs (A_t, B_{t+1}) with law

$$q(a, b) = \mu(a)\nu(b)(1 + ab), \tag{6.6}$$

and set $X_t = (A_t, B_t)$. The two coordinates of X_t belong to different independent edge pairs, so $X_t \sim \pi$. Given the past through time t, A_{t+1} is a fresh μ variable and B_{t+1} has law $\nu(b)(1 + A_t b)$. This verifies exactly the Markov kernel (6.1).

For one edge, A is a symmetric sign and $\mathbb{E}[A \mid B] = B$. Every centered function of A is a scalar multiple of A, so (2.1) yields

$$\rho(A, B)^2 = \mathbb{E}B^2 = 2\varepsilon. \tag{6.7}$$

For disjoint sets of times S, T, a single independent edge pair is either observed on only one side, or its two components are observed on opposite sides because one of its endpoints lies in S and the other in T. Tensorization over independent pairs (Lemma A.1 below) therefore bounds the full correlation by $\sqrt{2\varepsilon}$. This value is attained for $S = \{0\}$, $T = \{1\}$ by the two edge components. If $\operatorname{dist}(S, T) \geq 2$, no edge crosses between the two sets, so they use disjoint families of independent edge pairs. This proves (6.2)–(6.3).

The rows with $u = 1$ and $u = -1$ differ in total variation by 2ε. Hence $\delta_E(K) = 2\varepsilon$. To compute $\kappa_E(K)$ directly, an arbitrary input influences the output only through its sign coordinate U. If $m = \mathbb{E}U$, then

$$\mathbb{P}(V' = 1) = \varepsilon(1 + m), \quad \mathbb{P}(V' = -1) = \varepsilon(1 - m), \quad \mathbb{P}(V' = 0) = 1 - 2\varepsilon,$$

and the posterior means of U are $1, -1, m$, respectively. For a nondegenerate input sign,

$$\operatorname{Var}(\mathbb{E}[U \mid V']) = 2\varepsilon(1 - m^2) = 2\varepsilon \operatorname{Var}(U).$$

The fresh coordinate U' provides no additional information. The input factorization $(U, V) \to U \to V' \to (U', V')$ and (2.1) show that the maximal correlation is $\sqrt{2\varepsilon}$ whenever the sign is nondegenerate. Lemma 2.1 gives the same upper bound for every input law, proving (6.4). □

Corollary 6.2. *No uniform bound $\rho^*(1) \leq C\phi(1)^\alpha$, with fixed $C < \infty$ and $\alpha > 1/2$, holds over all stationary irreducible aperiodic Markov chains, even on a six-point state space.*

Proof. For the family above, the ratio of the two sides without C is $\sqrt{2}\,\varepsilon^{1/2-\alpha}$, which tends to infinity as $\varepsilon \downarrow 0$. □

7. A fixed chain with the sharp exponential-rate loss

Proposition 7.1. *Let $0 < q < 1$ and let X be the stationary chain on $\mathbb{N}_0$ with transitions*

$$P(i, 0) = 1 - q, \qquad P(i, i + 1) = q.$$

Its invariant law is $\pi_i = (1 - q)q^i$, and, for all $n \geq 1$,

$$\phi(n) = q^n, \qquad \rho(X_0, X_n) = q^{n/2}, \qquad q^{n/2} \leq \rho^*(n) \leq \Psi(q^{n/2}). \tag{7.1}$$

Proof. The stated invariant law follows immediately from the balance equations. The chain is irreducible and has a self-loop at zero, so it is aperiodic. An n-step transition consists either of no reset, with probability q^n, or of a last reset during those n steps. Explicitly,

$$P^n(i, j) = \begin{cases} (1 - q)q^j, & 0 \leq j < n, \\ q^n \mathbf{1}\{j = i + n\}, & j \geq n. \end{cases} \tag{7.2}$$

The first part is independent of i. The stationary tail on $\{n, n + 1, \ldots\}$ has mass q^n and, after translating by n, has law π. Thus

$$\|P^n(i, \cdot) - \pi\|_{\mathrm{TV}} = q^n \|\delta_i - \pi\|_{\mathrm{TV}} = q^n(1 - \pi_i).$$

The supremum over i is q^n. The Markov continuation argument used in Proposition 6.1 gives $\phi(n) \leq q^n$. Conversely, for every i, conditioning on $X_0 = i$ and choosing a maximizing event determined by X_n gives the displayed row distance. Taking the supremum proves $\phi(n) = q^n$.

The transition (7.2) is an erasure channel with a visible erasure flag. If the output is at least n, the input is recovered exactly by subtracting n; if the output is below n, its law is independent of the input. For any input law λ and any centered $h \in L^2(\lambda)$,

$$\mathbb{E}[h(U) \mid V] = \begin{cases} 0, & V < n, \\ h(V-n), & V \geq n. \end{cases}$$

Consequently

$$\operatorname{Var}(\mathbb{E}[h(U) \mid V]) = q^n \operatorname{Var}(h(U)).$$

Every nondegenerate input has maximal correlation $q^{n/2}$, so the robust lag coefficient is exactly $\kappa(n) = q^{n/2}$. Taking the stationary input proves the middle identity in (7.1). The lower bound for $\rho^*(n)$ follows by taking the singleton observation sets $\{0\}, \{n\}$, and the upper bound follows from (5.1). □

Since $\Psi(r) \leq 2r$, Proposition 7.1 proves part (2) of Theorem 1.2. Together with Proposition 6.1, it completes that theorem. It also shows that the rate exponent $c/2$ in Corollary 5.2 cannot in general be replaced by a larger fixed multiple of c, even when one restricts to stationary irreducible aperiodic countable-state chains.

Remark 7.2 (The lower psi coefficients of the reset chain)**.** Bradley's lower psi coefficient is $\psi'(\mathcal{A}, \mathcal{B}) = \inf \mathbb{P}(A \cap B)/(\mathbb{P}(A)\mathbb{P}(B))$ over $A \in \mathcal{A}$, $B \in \mathcal{B}$ with $\mathbb{P}(A)\mathbb{P}(B) > 0$, and $\psi'(n)$ is the infimum over the past–future pairs of sigma-fields at gap n [5, (1.7)]. Every Markov chain with $\psi'(n) > 0$ for some n is rho-star-mixing [3, Theorem 1.2], [5, Theorem 3.3(7)]. For the reset chain, (7.2) gives $\mathbb{P}(X_0 = i,\, X_n = j) = 0$ whenever $j \geq n$ and $j \neq i + n$, while $\pi_i \pi_j > 0$. Hence $\psi'(n) = 0$ for every $n \geq 1$, and the chain, which is phi-mixing at the exponential rate q^n, is not covered by that criterion.

8. First-order sharpness of path amplification

Consider a stationary symmetric two-state chain taking values in $\{-1, +1\}$, with

$$\mathbb{P}(X_{k+1} = y \mid X_k = x) = \frac{1 + axy}{2}, \qquad 0 < a < 1. \tag{8.1}$$

For a lag n, put $r = a^n$.

Proposition 8.1. *For the chain* (8.1)*,*

$$\phi(n) = \frac{r}{2}, \qquad \kappa(n) = r, \tag{8.2}$$

and

$$\frac{2r}{1 + r^2} \leq \rho^*(n) \leq \Psi(r). \tag{8.3}$$

In particular, as $r \downarrow 0$*,*

$$\rho^*(n) = 2r + O(r^3). \tag{8.4}$$

Proof. The stationary law is uniform, and

$$\mathbb{E}[X_{k+h} \mid X_k] = a^h X_k, \qquad \mathbb{E}[X_i X_j] = a^{|i-j|}.$$

The two n-step rows are at distance $r/2$ from the uniform law. The Markov continuation argument gives the upper phi bound, and conditioning on $X_0 = 1$ and testing $X_n = 1$ gives equality.

For the binary channel with parameter r and input mean $m \in (-1, 1)$, both centered spaces are one-dimensional. Direct computation gives

$$\rho(U, V)^2 = \frac{r^2(1 - m^2)}{1 - r^2 m^2} \leq r^2,$$

with equality at $m = 0$. The bound at a longer lag is smaller, so $\kappa(n) = r$. Equation (5.1) proves the upper bound in (8.3).

For the lower bound, take

$$S_N = \{0, 2n, \dots, 2Nn\}, \qquad T_N = \{n, 3n, \dots, (2N-1)n\},$$

and centered test functions

$$F_N = \sum_{i=0}^{N} X_{2in}, \qquad G_N = \sum_{j=0}^{N-1} X_{(2j+1)n}.$$

Counting pairs by their separation gives

$$\begin{aligned}
\operatorname{Var}(F_N) &= N + 1 + 2\sum_{h=1}^{N}(N+1-h)r^{2h}, \\
\operatorname{Var}(G_N) &= N + 2\sum_{h=1}^{N-1}(N-h)r^{2h}, \\
\operatorname{Cov}(F_N, G_N) &= 2\sum_{h=0}^{N-1}(N-h)r^{2h+1}.
\end{aligned}$$

For fixed $r < 1$, division by N and summation of the geometric series yield

$$\frac{\operatorname{Var}(F_N)}{N}, \frac{\operatorname{Var}(G_N)}{N} \longrightarrow \frac{1+r^2}{1-r^2}, \qquad \frac{\operatorname{Cov}(F_N, G_N)}{N} \longrightarrow \frac{2r}{1-r^2}.$$

Therefore $\operatorname{Corr}(F_N, G_N) \to 2r/(1+r^2)$, proving the lower bound. For small r, its expansion is $2r - 2r^3 + O(r^5)$, whereas $\Psi(r) = 2r - r^3 + O(r^5)$. The two bounds imply (8.4). □

Thus no coefficient smaller than two can replace the leading coefficient in a universal bound $\rho^*(n) \le 2\kappa(n) + o(\kappa(n))$. This occurs already for positive two-state kernels. In contrast, this family does not saturate the conversion $\kappa(n)^2 \le 2\phi(n)$: here $\kappa(n) = 2\phi(n)$. The local square-root extremality and the long-path amplification are distinct mechanisms.

9. The remaining constant question

The main implication and its square-root exponent do not determine the optimal universal envelope. To state the remaining quantitative question precisely, define

$$M(x) = \sup\{\rho_X^*(1) : X \text{ is a stationary standard-Borel Markov chain, } \phi_X(1) \le x\}.$$

For $0 < x \le 1/4$, Theorem 1.1 and Proposition 6.1 give

$$\sqrt{2x} \le M(x) \le 2\sqrt{2x(1-2x)}. \tag{9.1}$$

Question 9.1. What is the sharp small-x asymptotic behavior of $M(x)$? In particular, does $M(x)/\sqrt{x}$ have a limit, and what is its value?

The examples above do not answer this question. They prove the optimal exponent and separate sharpness statements for two steps of the argument, but they do not simultaneously saturate both steps in a single family. Peyre's theorems are optimal under their abstract prefix-correlation hypotheses [14, Section 3.5], so a sharper universal leading constant would require additional information about which local channels can coexist along one stationary Markov path.

10. Machine-checked components and code

The repository [18] contains the source of this paper, a Lean 4 development checked against Mathlib, and the finite checks. The Lean development does not formalize the measure-theoretic parts of the argument (Lemmas 2.2 and 3.1, Proposition 3.2, Peyre's theorems, Lemma 5.1). It covers the following components, each as a theorem whose only axioms are propositional extensionality, choice and quotient soundness.

(1) The function Ψ of (4.3): the closed form $\sin(2\arcsin r) = 2r\sqrt{1-r^2}$, the saturation $\Psi = 1$ on $[1/\sqrt{2}, 1]$, monotonicity, $\Psi(r) \le 2r$, the identity $\Psi(\sqrt{2\phi}) = 2\sqrt{2\phi(1-2\phi)}$ for $\phi \le 1/4$, and the derivation of (1.5) and (1.6) from $\rho^* \le \Psi(\kappa)$ and $\kappa^2 \le 2\phi$; the thresholds of Corollary 5.3 and of the remark following it; Corollary 5.2; the consistency of the envelope (9.1); and the expansions $2r - r^3 - r^5 \le \Psi(r) \le 2r - r^3$ and $2r - 2r^3 \le 2r/(1+r^2) \le 2r - 2r^3 + 2r^5$ behind (8.4), together with $2r/(1+r^2) \le \Psi(r)$ on $[0, 1/\sqrt{2}]$.
(2) The six-state chain of Section 6: nonnegativity and row sums of (6.1) for $0 \le \varepsilon \le 1/2$, invariance of π, the identity $K^2 = \Pi$, the row distances (6.5), the row diameter 2ε and the pair of rows attaining it, the second moment (6.7), the posterior mean $\mathbb{E}[A \mid B] = B$, and the posterior variance identity $2\varepsilon(1-m^2)$.
(3) The binary channel of Section 8: the posterior variance formula, the resulting expression $r^2(1-m^2)/(1-r^2m^2)$, and its bound by r^2 with equality at $m = 0$.
(4) The path matrix of Section 4: the bilinear bound $2\kappa\|x\|\|y\|$ for a two-diagonal matrix with entries at most κ, the quadratic-form bound showing that the path adjacency matrix has largest eigenvalue at most two, and the sine identity behind the eigenvector $\sin(k\pi/(m+1))$.
(5) The finite case of Lemma 2.1: the oscillation contraction $\operatorname{osc}(Mh) \le \delta(M)\operatorname{osc}(h)$ for a stochastic matrix, the consequence that a nonconstant eigenvector of a stochastic matrix has eigenvalue at most the row diameter, and the total-variation contraction $\delta(KR) \le \delta(K)$ under postprocessing.
(6) The reset chain of Section 7: the recursion characterizing the closed form (7.2), its row sums, and the invariant law $\pi_i = (1-q)q^i$ with its balance equations as convergent series.

The repository also contains exact rational checks of the six-state identities and floating-point checks of the inequalities of Theorem 1.1 and Proposition 4.2 on random finite chains, stationary and time-inhomogeneous, with the exact maximal correlation of (X_S, X_T) computed by a singular value decomposition of the normalized joint law. These computations are supplementary checks, not replacements for the proofs.

Appendix A. Tensorization over independent pairs

Lemma A.1. *If the pairs (U_i, V_i), i in a finite or countable set, are mutually independent, then*

$$\rho((U_i)_i, (V_i)_i) = \sup_i \rho(U_i, V_i).$$

A deterministic member of a pair is allowed and contributes zero.

Proof. For finitely many pairs, let $T_i : L^2(\mathrm{Law}(U_i)) \to L^2(\mathrm{Law}(V_i))$ be conditional expectation. It preserves constants, maps centered functions to centered functions, and has norm $\rho(U_i, V_i)$ on the centered subspace. Independence makes the full conditional-expectation operator the Hilbert tensor product $\bigotimes_i T_i$.

Decompose each factor as the orthogonal sum of constants and centered functions. The centered subspace of the full product is the orthogonal sum indexed by nonempty subsets J of the coordinates. On the summand indexed by J, the operator norm is $\prod_{i\in J} \rho(U_i, V_i)$, which is at most $\max_i \rho(U_i, V_i)$ because every factor lies in $[0, 1]$. This proves the upper bound. Functions depending on a single pair give the reverse inequality. For countably many pairs, finite-coordinate conditional expectations converge in L^2, as in Lemma 5.1. □

Acknowledgments and funding

The research presented in this paper was supported by the European Research Council (ERC) under the European Union's Horizon Europe research and innovation programme (grant agreement No. 101041711), by the Simons Foundation as part of the Collaboration on the Mathematical and Scientific Foundations of Deep Learning, by Heights Labs, by Convex Nexus

Capital, by the Israel Science Foundation (grant number 2258/19), and by the Israel Science Foundation (ISF Grant 4101/25).

This work was done by multiple AI agents, Claude (Anthropic) and Codex (OpenAI), connected to Math Brain [21] and working together inside the Brain framework [23], a shared, persistent memory in which the agents record claims, proofs, failed attempts and methods, retrieve one another's results, coordinate who works on what, and audit each other's work. For more on the details behind the Brain and Math Brain, see the Math Brain paper [22], the Math Brain repository [21] and the Brain framework repository [23].

Declaration of competing interest

The author holds equity in Heights Labs and in Convex Nexus Capital and has a management role at Convex Nexus Capital. These entities are included in the funding declaration above.

Declaration of generative AI and AI-assisted technologies

This manuscript was developed with AI assistance as part of an ongoing AI-assisted mathematical research project directed by the author. OpenAI's ChatGPT (GPT-5.6) produced the first complete draft of the argument, the sharpness examples and the finite checker from the project's working notes. Anthropic's Claude (Fable 5.1) then audited the manuscript against the primary sources cited here, in particular Bradley's survey and Peyre's monograph, tested the inequalities of Theorem 1.1 and Proposition 4.2 numerically on random finite chains, produced the Lean 4 development of Section 10, and revised the manuscript, the bibliography and these declarations. None of this is human peer review or a formal verification of the infinite-state theorem. The author is responsible for the mathematics and for decisions concerning dissemination.

Data and code availability

The arguments are analytic and do not rely on numerical evidence. The LaTeX source, the Lean 4 files with the recorded axiom output, the exact-arithmetic and floating-point checkers with their saved outputs, and the audit record are available at `https://github.com/yspennstate/phi-mixing-interlaced-rho-mixing-markov` [18]. No external empirical dataset is required.

The paper is part of the Math Brain project, whose public repository is `https://github.com/yspennstate/math-brain-public` [21].

References

[1] R. C. Bradley, *On the spectral density and asymptotic normality of weakly dependent random fields*, J. Theoret. Probab. **5** (1992), 355–373.
[2] R. C. Bradley, *Equivalent mixing conditions for random fields*, Ann. Probab. **21** (1993), 1921–1926.
[3] R. C. Bradley, *Every "lower psi-mixing" Markov chain is "interlaced rho-mixing"*, Stochastic Process. Appl. **72** (1997), 221–239. doi:10.1016/S0304-4149(97)00090-2.
[4] R. C. Bradley, *A stationary rho-mixing Markov chain which is not "interlaced" rho-mixing*, J. Theoret. Probab. **14** (2001), 717–727. doi:10.1023/A:1017545123473.
[5] R. C. Bradley, *Basic properties of strong mixing conditions. A survey and some open questions*, Probab. Surv. **2** (2005), 107–144. doi:10.1214/154957805100000104.
[6] R. C. Bradley, *Introduction to Strong Mixing Conditions*, Vol. 1–3, Kendrick Press, Heber City, Utah, 2007.
[7] R. C. Bradley, *On mixing properties of reversible Markov chains*, New Zealand J. Math. **45** (2015), 71–87. arXiv:1403.4895.
[8] J. E. Cohen, Y. Iwasa, Gh. Rautu, M. B. Ruskai, E. Seneta and Gh. Zbaganu, *Relative entropy under mappings by stochastic matrices*, Linear Algebra Appl. **179** (1993), 211–235.
[9] R. L. Dobrushin, *Central limit theorem for nonstationary Markov chains. I, II*, Theory Probab. Appl. **1** (1956), 65–80, 329–383.
[10] I. A. Ibragimov and Yu. V. Linnik, *Independent and Stationary Sequences of Random Variables*, Wolters-Noordhoff, Groningen, 1971.
[11] A. N. Kolmogorov and Yu. A. Rozanov, *On strong mixing conditions for stationary Gaussian processes*, Theory Probab. Appl. **5** (1960), 204–208.

[12] C. Miller, *Three theorems on ρ^*-mixing random fields*, J. Theoret. Probab. **7** (1994), 867–882.
[13] M. Peligrad, *Maximum of partial sums and an invariance principle for a class of weak dependent random variables*, Proc. Amer. Math. Soc. **126** (1998), 1181–1189.
[14] R. Peyre, *Tensorizing maximal correlations*, arXiv:1004.1602v2 (2010), 128 pp. https://arxiv.org/abs/1004.1602v2.
[15] Y. Polyanskiy and Y. Wu, *Strong data-processing inequalities for channels and Bayesian networks*, in: Convexity and Concentration, IMA Vol. Math. Appl. **161**, Springer, New York, 2017, pp. 211–249. arXiv:1508.06025.
[16] A. Rényi, *On measures of dependence*, Acta Math. Acad. Sci. Hungar. **10** (1959), 441–451.
[17] M. Rosenblatt, *Central limit theorems for stationary processes*, in: Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability, Vol. 2, University of California Press, Berkeley, 1972, pp. 551–561.
[18] Y. Shmalo, *Phi-mixing implies interlaced rho-mixing for Markov chains: manuscript, Lean 4 development and finite checks*, GitHub repository, 2026. https://github.com/yspennstate/phi-mixing-interlaced-rho-mixing-markov.
[19] C. Stein, *A bound for the error in the normal approximation to the distribution of a sum of dependent random variables*, in: Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability, Vol. 2, University of California Press, Berkeley, 1972, pp. 583–602.
[20] S. Utev and M. Peligrad, *Maximal inequalities and an invariance principle for a class of weakly dependent random variables*, J. Theoret. Probab. **16** (2003), 101–115.
[21] Y. Shmalo, *Math Brain: public snapshot, code and tests*, 2026, https://github.com/yspennstate/math-brain-public.
[22] Y. Shmalo, *Math Brain: shared, auditable memory for multi-agent mathematical research*, October 2026, https://github.com/yspennstate/math-brain-public/tree/main/papers/math-brain-2026-10-24.
[23] Y. Shmalo, *Brain Framework*, public software repository, 2026, https://github.com/yspennstate/brain-framework-public.

Einstein Institute of Mathematics, The Hebrew University of Jerusalem, Jerusalem, Israel