ARC-TRANSITIVE GRAPHS THAT MIMIC THE CYCLE

YITZCHAK SHMALO

Abstract. Aldous and Fill ask (Open Problem 7.27 of *Reversible Markov Chains and Random Walks on Graphs*) whether, over arc-transitive graphs other than the cycles, the maximal mean commute time of simple random walk is $O(n \log n)$, the relaxation time is $O(n)$, and the ratio of the maximal mean commute time to twice the average hitting time tends to one. The answer to all three questions is no. Replacing every vertex of the m-cycle by $r \geq 2$ pairwise non-adjacent copies gives a connected arc-transitive graph of degree $2r$ on $n = rm$ vertices whose mean hitting times are $j(m-j)+(r-1)m$ at cycle-distance $j \geq 1$ and rm between copies of the same vertex; its maximal mean commute time and its relaxation time are of order n^2, and the ratio tends to $3/2$. More generally, for any connected regular graph H on N vertices and $r \geq 2$ the blow-up $H[\overline{K}_r]$ has mean hitting times $\mathbb{E}_u T_v + (r-1)N$, average hitting time $\tau_0(H) + (r-1)N$ and relaxation time $\max(\tau_2(H), 1)$, and every arc-transitive graph is such a blow-up of a twin-free arc-transitive graph. For bounded r the three properties therefore pass between a graph and its twin-free quotient, so the question of Aldous and Fill, with the blow-ups of cycles excluded, is equivalent to its restriction to twin-free arc-transitive graphs, where it remains open. The arithmetic identities behind the formulas are checked in Lean 4.

1. Introduction

All graphs are finite and simple. Let G be a connected regular graph on n vertices and let (X_t) be simple random walk on G, in discrete time or in its continuous-time version with unit rate; the quantities below do not depend on the choice. Write $T_w = \inf\{t \geq 0 : X_t = w\}$ for the hitting time of w, $\mathbb{E}_v T_w$ for its mean when the walk starts at v, and π for the stationary distribution, which is uniform. Aldous and Fill [1] study, among others, the parameters

$$\tau^* = \max_{v,w} \big(\mathbb{E}_v T_w + \mathbb{E}_w T_v\big), \qquad \tau_0 = \sum_{v,w} \pi_v \pi_w \, \mathbb{E}_v T_w, \qquad \tau_2 = \frac{1}{1-\lambda_2}, \tag{1}$$

the maximal mean commute time, the average hitting time, and the relaxation time, λ_2 being the second largest eigenvalue of the transition matrix P. As in [1], τ_2 is defined through λ_2 alone; it is the relaxation time of the continuous-time chain and is unaffected by periodicity of the discrete-time walk.

A graph is *vertex-transitive* if its automorphism group acts transitively on vertices, and *arc-transitive* if it acts transitively on arcs, that is, for any two ordered pairs of adjacent vertices (v_1, w_1) and (v_2, w_2) there is an automorphism γ with $\gamma(v_1) = v_2$ and $\gamma(w_1) = w_2$ [3, 7]. Chapter 7 of [1] is devoted to walks on such graphs. On a vertex-transitive graph $\mathbb{E}_v T_w = \mathbb{E}_w T_v$ [2], so $\tau^* = 2 \max_{v,w} \mathbb{E}_v T_w$; in particular $\tau^* \geq 2\tau_0$, a maximum being at least an average, and $\tau^* \leq 4\tau_0$ by Lemma 7.2 of [1]. On an arc-transitive graph on n vertices, $\mathbb{E}_v T_w = n - 1$ for every edge (v, w) and $\mathbb{E}_v T_w \geq n - 2 + d(v, w)$ in general (Lemma 7.19 of [1]).

For the n-cycle (Example 5.7 of [1]), $\tau^* = 2\lfloor n^2/4 \rfloor$, $\tau_0 = (n^2-1)/6$, $\tau_2 = 1/(1-\cos(2\pi/n)) \sim n^2/(2\pi^2)$, and $\tau^*/(2\tau_0) \to 3/2$. Example 7.14 of [1] is the Cayley graph of $\mathbb{Z}_m \times \mathbb{Z}_2$ with generators $(\pm 1, 0)$ and $(0, 1)$, a ladder closed into a ring. It is vertex-transitive, its parameters τ^* and τ_2 are of order n^2 like those of the cycle (its transition matrix has the eigenvalues

Date: September 15, 2026.

$(2\cos(2\pi k/m)\pm 1)/3)$, but the mean hitting time of a neighbour along a rung differs from that along the ring, so it is not arc-transitive. Aldous and Fill ask:

> **Open Problem 7.27** [1]. Is it true that, over arc-transitive graphs excluding the n-cycles, $\tau^* = O(n\log n)$, $\tau_2 = O(n)$ and $\tau^*/(2\tau_0) = 1 + o(1)$?

The answer to each of the three questions is no. For a graph H and $r \geq 1$ let $H[\overline{K}_r]$ be the lexicographic product of H with the edgeless graph on r vertices: its vertex set is $V(H)\times\{1,\dots,r\}$, and (u,a) is adjacent to (v,b) if and only if u is adjacent to v in H. We call $H[\overline{K}_r]$ the r-fold blow-up of H; it replaces every vertex of H by r pairwise non-adjacent copies, which we call *twins*. When $H = C_m$ is the m-cycle and $r = 2$, the graph $C_m[\overline{K}_2]$ is the ladder of Example 7.14 with its rungs removed and the two diagonals of every square added.

Theorem 1.1. *Let $m \geq 4$ and $r \geq 2$, and let $G = C_m[\overline{K}_r]$, a connected graph of degree $2r$ on $n = rm$ vertices. Then G is arc-transitive, and for vertices v, w of G whose cycle coordinates are at distance j,*

$$\mathbb{E}_v T_w = \begin{cases} j(m-j) + (r-1)m, & 1 \leq j \leq m-1, \\ rm, & j = 0,\ v \neq w. \end{cases}$$

Consequently

$$\tau^* = 2\left\lfloor \frac{m^2}{4} \right\rfloor + 2(r-1)m, \qquad \tau_0 = \frac{m^2-1}{6} + (r-1)m, \qquad \tau_2 = \frac{1}{1-\cos(2\pi/m)},$$

so that for fixed r, as $m \to \infty$, $\tau^* \sim n^2/(2r^2)$, $\tau_2 \sim n^2/(2\pi^2 r^2)$ and $\tau^*/(2\tau_0) \to 3/2$.

The graph $C_m[\overline{K}_r]$ is the lexicographic product of a cycle with an empty graph [11, 8]; it is the wreath graph $W(m,r)$ of [9] and the graph $C(r,m,1)$ of Praeger and Xu [10], and its arc-transitivity is classical. For $m = 4$ it is the complete bipartite graph $K_{2r,2r}$. The content of Theorem 1.1 is what simple random walk does on it. The mechanism is not special to cycles.

Theorem 1.2. *Let H be a connected regular graph on $N \geq 2$ vertices, $r \geq 1$, and $G = H[\overline{K}_r]$.*

(i) *For $u \neq v$ in H and any a, b, $\mathbb{E}_{(u,a)} T_{(v,b)} = \mathbb{E}_u T_v + (r-1)N$; for $a \neq b$, $\mathbb{E}_{(u,a)} T_{(u,b)} = rN$.*

(ii) *If $r \geq 2$, then $\tau^*(G) = \max\left(\tau^*(H) + 2(r-1)N,\ 2rN\right)$, $\tau_0(G) = \tau_0(H) + (r-1)N$, and $\tau_2(G) = \max(\tau_2(H), 1)$.*

(iii) *If H is arc-transitive, so is G. Conversely, every connected arc-transitive graph G is isomorphic to $H[\overline{K}_r]$ for a unique $r \geq 1$ and a twin-free arc-transitive graph H, unique up to isomorphism, namely the quotient of G by the relation "same neighbourhood".*

Two vertices of a graph are twins if they have the same neighbourhood; a graph is twin-free if no two distinct vertices are twins. Twin-free graphs are also called point-determining [13], R-thin [8], irreducible [12] and worthy [9]. The quotient by the twin relation is a classical device: Sabidussi [12] used it for vertex-transitive graphs, and the fact that a vertex-transitive graph is arc-transitive if and only if its twin quotient is appears as Corollary 7 of [4]. Part (iii) records this in the form needed here, with a proof for completeness; the spectral part of (ii) is the classical spectrum of a lexicographic product [5, Section 2.5]. Part (iii) says that the arc-transitive graphs with twins are exactly the blow-ups of twin-free arc-transitive graphs, and part (ii) says how the three parameters of (1) change under a blow-up. Combining the two:

Corollary 1.3. *Let (G_k) be a sequence of connected arc-transitive graphs on $n_k \to \infty$ vertices with twin classes of bounded size, and let H_k be the twin-free quotient of G_k. If only finitely many H_k are cycles, then (G_k) satisfies the three statements of Open Problem 7.27 if and only if (H_k) does. If H_k is a cycle for infinitely many k, then (G_k) satisfies none of them along that subsequence.*

So, for bounded twin classes, the cycle is the only obstruction that twins can introduce, and Open Problem 7.27 with the blow-ups of cycles excluded is equivalent to its restriction to twin-free graphs:

Question 1.4. Is it true that, over twin-free arc-transitive graphs excluding the n-cycles, $\tau^* = O(n \log n)$, $\tau_2 = O(n)$ and $\tau^*/(2\tau_0) = 1 + o(1)$?

Question 1.4 is open. The ladder of Example 7.14 is twin-free but not arc-transitive; the complete graphs and the hypercubes are twin-free and arc-transitive and satisfy the three statements. For Cayley graphs of moderate growth in the sense of [6] the relaxation time is bounded above and below by constant multiples of the square of the diameter, the constants depending on the growth parameters, so an affirmative answer would bound the diameter of every twin-free arc-transitive Cayley graph of moderate growth, other than a cycle, by a constant multiple of $\sqrt{n}$. Unbounded twin classes behave differently: if $r/m \to \infty$ the graph $C_m[\overline{K}_r]$ satisfies all three statements, since the term $2(r-1)m$ then dominates both τ^* and $2\tau_0$ while $\tau_2 \le m^2 \le n$, so the family $C_m[\overline{K}_r]$ interpolates between the cycle and the complete bipartite graph $K_{r,r} = K_2[\overline{K}_r]$.

Section 2 proves Theorem 1.2, Section 3 proves Theorem 1.1 and Corollary 1.3, and Section 4 describes the exact numerical checks and the Lean verification of the arithmetic.

2. Random walk on a blow-up

Throughout, H is a connected d-regular graph on N vertices, $r \ge 1$, and $G = H[\overline{K}_r]$, which is connected and rd-regular on $n = rN$ vertices. Write a vertex of G as (u, a) with $u \in V(H)$ and $a \in [r] = \{1, \dots, r\}$, and call $\{u\} \times [r]$ the fibre over u.

Lemma 2.1 (the walk on a blow-up)**.** *Let $X_t = (Y_t, \xi_t)$ be simple random walk on G. Then (Y_t) is simple random walk on H, and conditionally on $(Y_t)_{t \ge 0}$ the coordinates $\xi_1, \xi_2, \dots$ are independent and uniform on $[r]$.*

Proof. From (u, a) the walk moves to a uniform element of $N_H(u) \times [r]$, so Y_{t+1} is uniform on $N_H(Y_t)$ and ξ_{t+1} is uniform on $[r]$, independently of each other and of the past. □

Proof of Theorem 1.2(i). Fix the target (v, b) and a starting vertex $(u, a) \ne (v, b)$. Let $\sigma_1 < \sigma_2 < \cdots$ be the successive times $t \ge 1$ at which $Y_t = v$, and let $K = \min\{k : \xi_{\sigma_k} = b\}$, so that $T_{(v,b)} = \sigma_K$. By Lemma 2.1, K is geometric with success probability $1/r$ and independent of the sequence (σ_k); in particular $\mathbb{E}K = r$. By the strong Markov property of Y at the times σ_k, the increments $\sigma_{k+1} - \sigma_k$ are independent copies of the return time of Y to v, whose mean is $1/\pi_H(v) = N$. Hence

$$\mathbb{E}T_{(v,b)} = \mathbb{E}\sigma_1 + (\mathbb{E}K - 1)\, N = \mathbb{E}\sigma_1 + (r-1)N.$$

If $u \ne v$ then σ_1 is the hitting time of v by Y started at u, with mean $\mathbb{E}_u T_v$; if $u = v$ then σ_1 is the return time to v, with mean N, and the formula gives rN. □

Proof of Theorem 1.2(ii). Let $r \ge 2$. The stationary distribution of G is uniform. By (i) the commute time between (u, a) and (v, b) is $\mathbb{E}_u T_v + \mathbb{E}_v T_u + 2(r-1)N$ when $u \ne v$ and $2rN$ when $u = v$, $a \ne b$, which gives $\tau^*(G)$. For τ_0,

$$\begin{aligned}\tau_0(G) &= \frac{1}{r^2 N^2}\Big[\sum_u r(r-1)\, rN + \sum_{u \ne v} r^2\big(\mathbb{E}_u T_v + (r-1)N\big)\Big] \\ &= (r-1) + \tau_0(H) + (r-1)(N-1) = \tau_0(H) + (r-1)N.\end{aligned}$$

For τ_2, identify functions on $V(G)$ with functions $f(u,a)$; then

$$(P_G f)(u,a) = \frac{1}{d}\sum_{v\sim u}\frac{1}{r}\sum_{b=1}^{r} f(v,b),$$

that is, $P_G = P_H \otimes \frac{1}{r}J_r$ with J_r the all-ones $r\times r$ matrix. Since $\frac{1}{r}J_r$ has eigenvalues 1 (once) and 0 ($r-1$ times), the spectrum of P_G is the spectrum of P_H together with 0 with multiplicity $(r-1)N \geq 1$. Hence the second largest eigenvalue of P_G is $\max(\lambda_2(H),0)$ and $\tau_2(G) = \max(\tau_2(H),1)$. □

Proof of Theorem 1.2(iii). Suppose H is arc-transitive. The maps $(u,a)\mapsto(\gamma u,a)$ for $\gamma\in \mathrm{Aut}(H)$ and $(u,a)\mapsto(u,\sigma_u(a))$ for any family of permutations σ_u of $[r]$ are automorphisms of G. Given arcs $((u,a),(v,b))$ and $((u',a'),(v',b'))$ of G, choose γ with $\gamma u = u'$, $\gamma v = v'$, and then permutations in the fibres over u' and v' (distinct fibres, since $u'\neq v'$) sending $a\mapsto a'$ and $b\mapsto b'$. So G is arc-transitive.

Conversely let G be arc-transitive, and call $x,y\in V(G)$ twins if $N_G(x) = N_G(y)$. This is an equivalence relation, twins are never adjacent (as $x\notin N_G(x)$), and if x,y are twins then every neighbour of x is a neighbour of y, so adjacency is a well-defined relation between twin classes. Automorphisms preserve the twin relation and hence permute the classes; as G is vertex-transitive, all classes have the same size r. Let H be the graph on the classes, with $[x]\sim[y]$ when $x\sim y$. Then $G\cong H[\overline{K}_r]$: choosing a bijection of each class with $[r]$, $(u,a)\sim(v,b)$ in G exactly when $u\sim v$ in H. The graph H is twin-free, since two classes with the same neighbourhood in H would form a single twin class of G, and it is arc-transitive, since an automorphism of G carrying the arc (x,y) to (x',y') induces an automorphism of H carrying $([x],[y])$ to $([x'],[y'])$. Uniqueness holds because in $H[\overline{K}_r]$ with H twin-free the twin classes are exactly the fibres, so r and H are determined by G up to isomorphism. □

Remark 2.2. The formula in (i) has a check built in. If H is arc-transitive on N vertices, Lemma 7.19 of [1] gives $\mathbb{E}_u T_v = N-1$ for every edge of H, and then (i) gives $\mathbb{E}_{(u,a)}T_{(v,b)} = N-1+(r-1)N = rN-1 = n-1$ for every edge of G, as Lemma 7.19 applied to the arc-transitive graph G requires.

3. The cycle

Proof of Theorem 1.1. Apply Theorem 1.2 with $H = C_m$, $N = m$, $d = 2$. The cycle is arc-transitive (its automorphism group is the dihedral group, which is transitive on arcs), so $G = C_m[\overline{K}_r]$ is arc-transitive by (iii); it is connected, $2r$-regular, and not a cycle since $2r\geq 4$.

For the cycle, $\mathbb{E}_u T_v = j(m-j)$ when u and v are at distance j: the function $h(j) = j(m-j)$ satisfies $h(0) = h(m) = 0$ and $h(j) = 1 + \frac{1}{2}(h(j-1)+h(j+1))$ for $1\leq j\leq m-1$, and the mean hitting times are the unique solution of this system. Part (i) of Theorem 1.2 gives the two hitting-time formulas of the theorem. Part (ii) gives

$$\tau^* = \max\Big(2\max_j j(m-j) + 2(r-1)m,\ 2rm\Big) = 2\Big\lfloor\frac{m^2}{4}\Big\rfloor + 2(r-1)m,$$

since $\max_j j(m-j) = \lfloor m^2/4\rfloor \geq m$ for $m\geq 4$. The cycle has $\tau_0(C_m) = \frac{1}{m}\sum_{j=0}^{m-1} j(m-j) = \frac{m^2-1}{6}$, so $\tau_0 = \frac{m^2-1}{6} + (r-1)m$. The transition matrix of the cycle has eigenvalues $\cos(2\pi k/m)$, $0\leq k<m$, with $\lambda_2 = \cos(2\pi/m)\geq 0$ for $m\geq 4$, so $\tau_2 = \max(\tau_2(C_m),1) = 1/(1-\cos(2\pi/m))$.

With $n = rm$ and r fixed, $\tau^*\sim m^2/2 = n^2/(2r^2)$, $\tau_2\sim m^2/(2\pi^2) = n^2/(2\pi^2 r^2)$, and

$$\frac{\tau^*}{2\tau_0} = \frac{\lfloor m^2/4\rfloor + (r-1)m}{(m^2-1)/6 + (r-1)m} \longrightarrow \frac{1/4}{1/6} = \frac{3}{2}.$$

□

For $r = 2$ and m odd the formulas read $\tau^* = (m^2-1)/2 + 2m \sim n^2/8$, $\tau_0 = (m^2-1)/6 + m$, $\tau_2 \sim n^2/(8\pi^2)$; Table 1 lists the first values, and Figure 1 the ratio.

m	n	τ^*	τ_0	$\tau^*/(2\tau_0)$	τ_2
5	10	22	9	1.2222	1.4472
7	14	38	15	1.2667	2.6560
9	18	58	67/3	1.2985	4.2743
11	22	82	31	1.3226	6.2994
15	30	142	157/3	1.3567	11.5668
21	42	262	283/3	1.3887	22.5087
31	62	542	191	1.4188	48.8518

TABLE 1. The parameters of $C_m[\overline{K}_2]$; τ^* and τ_0 are exact, $\tau_2 = 1/(1-\cos(2\pi/m))$.




FIGURE 1. $\tau^*/(2\tau_0)$ for $C_m[\overline{K}_2]$, m odd, $5 \le m \le 61$. The limit is $3/2$, the value for the cycle itself.

Proof of Corollary 1.3. Let G_k have n_k vertices, twin classes of size $r_k \le R$, and twin-free quotient H_k on $N_k = n_k/r_k$ vertices; by Theorem 1.2(iii) H_k is arc-transitive, and by (ii)

$$\tau^*(G_k) = \max\big(\tau^*(H_k) + 2(r_k-1)N_k,\, 2r_kN_k\big),$$
$$\tau_0(G_k) = \tau_0(H_k) + (r_k-1)N_k, \qquad \tau_2(G_k) = \max(\tau_2(H_k), 1).$$

Since $N_k \le n_k \le RN_k$, the terms $2(r_k-1)N_k$ and $2r_kN_k$ are $O(n_k)$, and $O(N_k \log N_k)$ and $O(n_k \log n_k)$ are the same condition; so $\tau^*(G_k) = O(n_k \log n_k)$ if and only if $\tau^*(H_k) = O(N_k \log N_k)$, and likewise $\tau_2(G_k) = O(n_k)$ if and only if $\tau_2(H_k) = O(N_k)$. For the ratio, write $a_k = \tau^*(H_k)$, $b_k = 2\tau_0(H_k)$ and $c_k = 2(r_k-1)N_k$. On a vertex-transitive graph $\tau^* \ge 2\tau_0$, so $a_k \ge b_k$, and $b_k \ge 2(N_k-1)^2/N_k$ by Lemma 7.19 of [1], so $c_k = O(b_k)$. When $a_k + c_k \ge 2r_kN_k$,

$$\frac{\tau^*(G_k)}{2\tau_0(G_k)} - 1 = \frac{a_k - b_k}{b_k + c_k},$$

which tends to 0 if and only if $(a_k - b_k)/b_k$ does. The other case, $a_k + c_k < 2r_kN_k$, means $\tau^*(H_k) < 2N_k$, which by Lemma 7.19(ii) of [1] forces H_k to be complete; then G_k is the complete multipartite graph with N_k parts of size r_k, for which $\tau^* \le 2n_k$, $\tau_2 \le 1$, and $\tau^*/(2\tau_0) - 1 = (2N_k-1)/\big((N_k-1)^2 + (r_k-1)N_k^2\big)$ for $r_k \ge 2$ and $1/(N_k-1)$ for $r_k = 1$; since $n_k = r_kN_k \to \infty$

these tend to 0, so all three statements hold for both G_k and H_k along such k. If H_k is a cycle for infinitely many k, then, since C_3 is complete and C_4 has twins, $H_k = C_{m_k}$ with $m_k \geq 5$, and $m_k = n_k/r_k \to \infty$; the cycle itself has $\tau^* \sim m_k^2/2$, $\tau_2 \sim m_k^2/(2\pi^2)$ and $\tau^*/(2\tau_0) \to 3/2$, and Theorem 1.1 gives the same for $G_k = C_{m_k}[\overline{K}_{r_k}]$ when $r_k \geq 2$, so G_k fails all three statements along that subsequence. □

4. Numerical checks and formal verification

Exact rational computation of the mean hitting times of simple random walk on $C_m[\overline{K}_r]$, by solving the linear system $(I - P)h = \mathbf{1}$ off the target, confirms the formulas of Theorem 1.1, including $\mathbb{E}_v T_w = n - 1$ on edges, for $r = 2$ and $m \in \{5, 6, 7, 9, 11, 15, 21, 31\}$ and for $(m, r) \in \{(5, 3), (7, 3), (7, 4), (9, 2), (9, 3), (11, 4)\}$. Arc-transitivity of $C_m[\overline{K}_2]$ for $m \in \{5, 6, 7, 9, 11, 15\}$ is confirmed by computing the orbit of one arc under the group generated by a rotation, a reflection and the transpositions of the two vertices of each fibre, which is the whole set of $8m$ arcs. The identities of Theorem 1.2(i) and (ii) hold, exactly for the hitting times and to 10^{-9} for the relaxation times, on the blow-ups $H[\overline{K}_2]$ and $H[\overline{K}_3]$ of K_2, K_4, $K_{3,3}$, the 3-cube, the Petersen graph and C_6. Table 1 and Figure 1 come from the same computation.

The arithmetic behind Theorem 1.1 is checked in Lean 4 (version 4.33.1) against Mathlib. For all integers m, r and j the files prove that $g(j) = j(m - j) + (r - 1)m$ satisfies the harmonic equation $2g(j) = 2 + g(j - 1) + g(j + 1)$, that $1 + g(1) = rm$ (the twin value) and $g(1) = rm - 1$ (Lemma 7.19), that $4j(m - j) \leq m^2$ and that the value at $j = \lfloor m/2 \rfloor$ is $\lfloor m^2/4 \rfloor$, the closed form $\sum_{j=0}^{m-1} j(m - j) = m(m^2 - 1)/6$ by induction, the resulting formula $\tau_0 = (m^2 - 1)/6 + (r - 1)m$, and the identity $2 \cdot 3\tau^* - 3 \cdot 6\tau_0 = -6(r - 1)m$ for odd m, which shows that $\tau^*/(2\tau_0) \to 3/2$. Every theorem depends only on the axioms `propext`, `Classical.choice` and `Quot.sound`. The probabilistic inputs, the uniqueness of the solution of the hitting-time system and the tensor decomposition of P_G, are not formalised.

Funding

The research presented in this paper was supported by the European Research Council (ERC) under the European Union's Horizon Europe research and innovation programme (grant agreement No. 101041711), by the Simons Foundation as part of the Collaboration on the Mathematical and Scientific Foundations of Deep Learning, by Heights Labs, by Convex Nexus Capital, by the Israel Science Foundation (grant number 2258/19), and by the Israel Science Foundation (ISF Grant 4101/25).

Declaration of competing interest

The author holds equity in Heights Labs and in Convex Nexus Capital and has a management role at Convex Nexus Capital. These entities are included in the funding declaration above.

Declaration of generative AI and AI-assisted technologies

This paper is a product of an ongoing project on the use of AI systems for research mathematics, provisionally called Math Brain, which studies how to make such systems better at solving open problems; the project and its tooling will be released publicly in due course. The mathematics, the literature search, the computations, the Lean formalisation and the text were produced by AI systems (Codex with GPT-6 Astra, OpenAI; Claude Opus 5 and Claude Fable 5.1, Anthropic), working from the list of open problems in [1]; three independent adversarial reviews by Claude Fable 5.1 re-derived every claim. The arguments have not been refereed by a human mathematician. The author set the goals of the project, commissioned and directed the

work and approved the release; the author did not carry out the mathematics, the verification, the computations or the writing, and takes responsibility for the decision to publish.

Data and code availability

The numerical code, the Lean files with their build logs and axiom reports, and the LaTeX source are available at

https://github.com/yspennstate/arc-transitive-graphs-that-mimic-the-cycle

References

[1] D. Aldous and J. A. Fill, Reversible Markov Chains and Random Walks on Graphs, unfinished monograph, 2002 (recompiled 2014; the numbering is that of the online edition), https://www.stat.berkeley.edu/~aldous/RWG/book.html.
[2] D. Aldous, Hitting times for random walks on vertex-transitive graphs, *Math. Proc. Cambridge Philos. Soc.* 106 (1989), 179–191.
[3] N. Biggs, Algebraic Graph Theory, 2nd ed., Cambridge University Press, 1993.
[4] S. Cockburn, K. Hatavets and E. Swartz, Twins and co-twins in circulant graphs, arXiv:2605.14903 (2026).
[5] D. Cvetković, M. Doob and H. Sachs, Spectra of Graphs: Theory and Applications, 3rd ed., Johann Ambrosius Barth, 1995.
[6] P. Diaconis and L. Saloff-Coste, Moderate growth and random walk on finite groups, *Geom. Funct. Anal.* 4 (1994), 1–36.
[7] C. Godsil and G. Royle, Algebraic Graph Theory, Graduate Texts in Mathematics 207, Springer, 2001.
[8] R. Hammack, W. Imrich and S. Klavžar, Handbook of Product Graphs, 2nd ed., CRC Press, 2011.
[9] P. Potočnik and S. Wilson, Recipes for edge-transitive tetravalent graphs, *Art Discrete Appl. Math.* 3 (2020), P1.08.
[10] C. E. Praeger and M.-Y. Xu, A characterization of a class of symmetric graphs of twice prime valency, *European J. Combin.* 10 (1989), 91–102.
[11] G. Sabidussi, The composition of graphs, *Duke Math. J.* 26 (1959), 693–696.
[12] G. Sabidussi, Vertex-transitive graphs, *Monatsh. Math.* 68 (1964), 426–438.
[13] D. P. Sumner, Point determination in graphs, *Discrete Math.* 5 (1973), 179–187.

Einstein Institute of Mathematics, The Hebrew University of Jerusalem, Jerusalem, Israel
Email address: yitzchak.shmalo@gmail.com