THE TIME-STATIONARY AND PALM INTER-ARRIVAL TIMES OF A MARKOV-MODULATED POISSON PROCESS NEED NOT BE ORDERED

YITZCHAK SHMALO

Abstract. For an irreducible Markov-modulated Poisson process, let T^π be the time from a stationary instant to the next arrival and T^α the time from an arrival to the next one in the event-stationary version. Asanjarani and Nazarathy conjectured that T^π is stochastically larger than T^α, and He listed the same property among open questions on the MMPP class of phase-type distributions. The conjecture is false. The order is equivalent to the hazard rate of T^π never exceeding the arrival rate. For a Cox process driven by a non-constant periodic intensity with uniform phase this fails on a set of positive measure. For the MMPP on the directed n-cycle with arrival rates $\varepsilon(1+b\cos(2\pi j/n))$ it fails for all small $\varepsilon>0$ whenever $n\geq 12$, and an explicit twelve-state MMPP with positive rational rates has $\mathbb{P}(T^\pi>19/2)-\mathbb{P}(T^\alpha>19/2)<-1/300000$, proved in exact rational arithmetic. For reversible modulators the order holds. We also show that the mean of He's MMPP-type distribution need not be convex in the speed of modulation.

1. Introduction

A Markov-modulated Poisson process (MMPP) is given by an irreducible generator Q on a finite set $\{1,\dots,m\}$ and a nonzero diagonal matrix $D=\operatorname{diag}(d_1,\dots,d_m)$ with $d_i\geq 0$: a Markov chain J with generator Q runs in the background, and arrivals occur at rate d_{J_t}. Write $C=Q-D$, let π be the stationary distribution of Q, let $\lambda=\pi D\mathbf{1}$ be the mean arrival rate, and let $\alpha=\pi D/\lambda$ be the law of the phase at an arrival in the event-stationary version. The time T^π from time 0 to the first arrival in the time-stationary process and the time T^α from an arrival to the next one in the event-stationary process are phase-type, with

$$S_\pi(t)=\mathbb{P}(T^\pi>t)=\pi e^{Ct}\mathbf{1},\qquad S_\alpha(t)=\mathbb{P}(T^\alpha>t)=\alpha e^{Ct}\mathbf{1}. \tag{1}$$

Asanjarani and Nazarathy [1] study four properties of such processes: $d^2\geq 1$ for the index of dispersion of counts, a decreasing hazard rate for T^α, $c^2\geq 1$ for the squared coefficient of variation of T^α, and the stochastic order $T^\pi\geq_{\mathrm{st}}T^\alpha$, which is property (IV), equation (2.9) of [1]:

$$\pi e^{Ct}\mathbf{1}\ \geq\ \alpha e^{Ct}\mathbf{1}\qquad\text{for all } t\geq 0. \tag{2}$$

They prove all four for Markov-switched Poisson processes and for MMPPs of order two, observe that the hazard-rate property fails for general MMPPs, and state (2) for every irreducible MMPP as Conjecture 3.9 of [1, p. 65] (Conjecture 4.2 in the preprint [2]). Their search over 10^6 random MMPPs of each order $3,4,5,6$ and over 10^6 random cyclic generators of order 4 found no violation. An earlier preprint of the same authors [3, Theorem 3.1] states (2) for every MMPP; it has since been withdrawn. He [4, Section 4.3] lists the same property, in the form "X_x is stochastically larger than $X_{D,x}$", among the properties that all MMPP-type phase-type distributions have numerically. The squared-coefficient conjecture is a theorem of He [4, Theorem 3.2].

2020 *Mathematics Subject Classification.* 60G55, 60K15, 60E15.

Key words and phrases. Markov-modulated Poisson process, Markovian arrival process, Palm distribution, stochastic order, Cox process, phase-type distribution.

We show that (2) fails. The first observation is a reformulation. Since $\pi Q = 0$, $S'_\pi(t) = \pi C e^{Ct}\mathbf{1} = -\pi D e^{Ct}\mathbf{1} = -\lambda S_\alpha(t)$, so

$$S_\pi(t) - S_\alpha(t) = S_\pi(t) + \frac{S'_\pi(t)}{\lambda} = \frac{e^{-\lambda t}}{\lambda}\, g'(t), \qquad g(t) = e^{\lambda t} S_\pi(t). \tag{3}$$

Thus (2) says exactly that g is nondecreasing, that is, that the hazard rate $-S'_\pi/S_\pi$ of T^π never exceeds λ. It always starts at λ, since $g'(0) = 0$. Our counterexamples are processes in which this hazard overshoots λ.

Theorem 1.1. *Let ℓ be a bounded measurable function on $\mathbb{R}$ that is periodic and not almost everywhere constant, let $T > 0$ be its least period (Lemma 2.1), let θ be uniform on $[0, T)$, and let N be the Cox process with intensity $s \mapsto \ell(\theta + s)$. Put $\lambda = T^{-1}\int_0^T \ell$, $L_\vartheta(t) = \int_0^t \ell(\vartheta + s)\, ds$,*

$$S(t) = \mathbb{P}(N(0,t] = 0) = \frac{1}{T}\int_0^T e^{-L_\vartheta(t)} d\vartheta, \qquad S^0(t) = \frac{1}{\lambda T}\int_0^T \ell(\vartheta)\, e^{-L_\vartheta(t)} d\vartheta.$$

Then $-S' = \lambda S^0$ almost everywhere, $g(t) = e^{\lambda t} S(t)$ satisfies $g(0) = g(T) = 1 < g(t)$ for $0 < t < T$, and $S(t) < S^0(t)$ for all t in a set of positive measure in $(0, T)$.

Here S^0 is the survival function of the interval after a typical arrival: the phase seen at an arrival has density proportional to ℓ, exactly as the phase of an MMPP seen at an arrival has law $\alpha = \pi D/\lambda$. Asanjarani and Nazarathy suggest Cox-process versions of their conjectures [1, Section 4]; Theorem 1.1 says the order fails for every non-constant periodic intensity. A long cycle of phases traversed at a steady pace imitates a periodic intensity, and this is the source of the MMPP counterexamples. The cleanest family is a small cosine modulation of a directed cycle.

Theorem 1.2. *Let $n \geq 12$ and $0 < b \leq 1$. Let Q be the generator of the directed n-cycle, $Q_{j,j+1} = 1$ and $Q_{jj} = -1$ (indices mod n), and for $\varepsilon > 0$ let $D_\varepsilon = \varepsilon\,\mathrm{diag}(1 + b\cos(2\pi j/n))_{j=0}^{n-1}$. Put $t_n = 3\pi/(2\sin(2\pi/n))$. There is $\varepsilon_0 > 0$ such that for $0 < \varepsilon < \varepsilon_0$ the MMPP (Q, D_ε) satisfies $\mathbb{P}(T^\pi > t_n) < \mathbb{P}(T^\alpha > t_n)$. For $b < 1$ all arrival rates are positive.*

The threshold 12 is where the first-order term changes sign: for $3 \leq n \leq 11$ the first-order term is positive at every $t > 0$ (Remark 3.1). Rounding the case $n = 12$, $b = 1/2$, $\varepsilon = 10^{-3}$ to rationals gives an explicit instance.

Theorem 1.3. *Let Q be the directed 12-cycle of Theorem 1.2 and $D = 10^{-6}\,\mathrm{diag}(p_0, \ldots, p_{11})$ with*

$$(p_0, \ldots, p_{11}) = (1500, 1433, 1250, 1000, 750, 567, 500, 567, 750, 1000, 1250, 1433).$$

Then $\mathbb{P}(T^\pi > 19/2) - \mathbb{P}(T^\alpha > 19/2) < -1/300000$. Numerically

$$\mathbb{P}(T^\pi > 19/2) = 0.990546031181966, \qquad \mathbb{P}(T^\alpha > 19/2) = 0.990549434945316,$$

and the difference is $-3.40376335 \times 10^{-6}$.

The reformulation (3) also shows where to look. The order cannot fail for reversible modulators (Proposition 4.1); this contains the case of symmetric C mentioned in [1, Section 4]. The directed cycle is as far from reversible as possible.

The same list [4, Section 4.3] contains the property that $E[X_x]$ is convex and non-increasing in $x \geq 0$, where X_x has phase-type representation $(\pi, xQ - D)$. Proposition 5.1 shows that convexity fails, and with it the convexity of the squared coefficient of variation of $X_{D,x}$. We do not address monotonicity.

2. Periodic Cox processes

Lemma 2.1. *Let $\ell \in L^\infty(\mathbb{R})$ satisfy $\ell(\cdot+\tau)=\ell$ almost everywhere for some $\tau>0$, and suppose ℓ is not almost everywhere constant. Then $P=\{s\in\mathbb{R}:\ell(\cdot+s)=\ell \text{ a.e.}\}=T\mathbb{Z}$ for some $T>0$.*

Proof. P is a subgroup of $\mathbb{R}$, and it is closed because $s\mapsto\ell(\cdot+s)$ is continuous into L^1_{loc}. A closed subgroup of $\mathbb{R}$ is $\{0\}$, $T\mathbb{Z}$ or $\mathbb{R}$. It contains τ, so it is not $\{0\}$. If $P=\mathbb{R}$, then $\ell*\phi$ is constant for every continuous ϕ with compact support, and letting ϕ run through an approximate identity shows that ℓ is constant almost everywhere. □

Proof of Theorem 1.1. Conditionally on $\theta=\vartheta$ the process is Poisson with intensity $\ell(\vartheta+\cdot)$, which gives the formula for S. Since ℓ is bounded, $t\mapsto e^{-L_\vartheta(t)}$ is Lipschitz uniformly in ϑ, so by Fubini's theorem S is Lipschitz and for almost every t

$$-S'(t)=\frac{1}{T}\int_0^T \ell(\vartheta+t)\,e^{-L_\vartheta(t)}d\vartheta.$$

Fix such a t and put $H(\vartheta)=L_\vartheta(t)=\Lambda(\vartheta+t)-\Lambda(\vartheta)$, where $\Lambda(x)=\int_0^x\ell$. Then H is Lipschitz and T-periodic with $H'(\vartheta)=\ell(\vartheta+t)-\ell(\vartheta)$ for almost every ϑ, so

$$\int_0^T\big(\ell(\vartheta+t)-\ell(\vartheta)\big)e^{-H(\vartheta)}d\vartheta=\big[-e^{-H}\big]_0^T=0,$$

and $-S'(t)=\lambda S^0(t)$.

Next, $\mathbb{E}L_\theta(t)=\int_0^t T^{-1}\int_0^T\ell(\vartheta+s)\,d\vartheta\,ds=\lambda t$, so Jensen's inequality gives $S(t)\ge e^{-\lambda t}$, that is $g(t)\ge1$, with equality only if $\vartheta\mapsto L_\vartheta(t)$ is almost everywhere constant on $[0,T)$. Since it is continuous, it would then be constant, its derivative $\ell(\vartheta+t)-\ell(\vartheta)$ would vanish for almost every ϑ, and t would lie in $P=T\mathbb{Z}$. Hence $g(t)>1$ for $0<t<T$. At $t=T$ the exposure $L_\vartheta(T)=\lambda T$ does not depend on ϑ, so $g(T)=1$; and $g(0)=1$.

Pick $t_1\in(0,T)$. The function g is Lipschitz and $\int_{t_1}^T g'=g(T)-g(t_1)<0$, so $g'<0$ on a subset of (t_1,T) of positive measure. At almost every point of it, $S-S^0=S+S'/\lambda=\lambda^{-1}e^{-\lambda t}g'<0$. □

For the square wave $\ell=2$ on $[0,\frac12)$ and 0 on $[\frac12,1)$, with $\lambda=1$, one finds $g(\frac12)=1.04219$ and a hazard rate $-S'/S=1.1237$ at $t=\frac34$. From a typical instant, the phases that survive to time t are those that have not yet met the high part of the intensity. Near a full period the cycle brings the high part around to all of them at once, and the hazard of T^π overshoots the mean rate. An arrival happens inside the high part, so the Palm interval has already paid that cost.

3. Cosine modulation of a directed cycle

Proof of Theorem 1.2. Write $\vartheta=2\pi/n$, $c_j=\cos(\vartheta j)$ and $f=\mathbf{1}+bc$, so $D_\varepsilon=\varepsilon\,\mathrm{diag}(f)$. The cycle generator is doubly stochastic, so π is uniform. Since $\sum_j c_j=0$ for $n\ge2$, the mean rate is $\lambda=\varepsilon$ and $\alpha_j=f_j/n$, so $\pi-\alpha=-(b/n)c$ does not depend on ε. Let

$$F_\varepsilon(t)=S_\pi(t)-S_\alpha(t)=-\frac{b}{n}\,c^\top e^{(Q-\varepsilon\,\mathrm{diag}\,f)t}\mathbf{1}.$$

For fixed t this is an entire function of ε with $F_0(t)=-\frac{b}{n}\sum_j c_j=0$. By Duhamel's formula,

$$\frac{d}{d\varepsilon}e^{(Q-\varepsilon\,\mathrm{diag}\,f)t}\Big|_{\varepsilon=0}\mathbf{1}=-\int_0^t e^{(t-s)Q}\,\mathrm{diag}(f)\,e^{sQ}\mathbf{1}\,ds=-\int_0^t e^{uQ}f\,du,$$

using $e^{sQ}\mathbf{1}=\mathbf{1}$. For the vector $e_\vartheta=(e^{i\vartheta j})_j$ we have $Qe_\vartheta=(e^{i\vartheta}-1)e_\vartheta$, and $e^{i\vartheta}-1=-\delta+i\omega$ with $\delta=1-\cos\vartheta$, $\omega=\sin\vartheta$. Taking real parts, $(e^{uQ}c)_j=e^{-\delta u}\cos(\vartheta j+\omega u)$, so $e^{uQ}f=$

$\mathbf{1}+b\,e^{-\delta u}\big(\cos(\vartheta j+\omega u)\big)_j$. For $n\ge 3$, $\sum_j \cos(2\vartheta j+\omega u)=0$, hence $\sum_j c_j\cos(\vartheta j+\omega u)=\frac{n}{2}\cos(\omega u)$ and

$$\frac{\partial F_\varepsilon(t)}{\partial\varepsilon}\Big|_{\varepsilon=0}=\frac{b}{n}\int_0^t c^\top e^{uQ}f\,du=\frac{b^2}{2}J_n(t),\qquad J_n(t)=\int_0^t e^{-\delta u}\cos(\omega u)\,du. \tag{4}$$

Explicitly,

$$J_n(t)=\frac{\delta-e^{-\delta t}(\delta\cos\omega t-\omega\sin\omega t)}{\delta^2+\omega^2},\qquad J_n(t_n)=\frac{\delta-\omega e^{-\delta t_n}}{\delta^2+\omega^2},$$

because $\omega t_n=3\pi/2$. Put $x=\delta/\omega=\tan(\pi/n)$. Then $\delta t_n=\frac{3\pi}{2}x$, and $J_n(t_n)<0$ if and only if $x\,e^{3\pi x/2}<1$. The left side is increasing in x, and x decreases in n. At $n=11$ it equals 1.1714 and at $n=12$ it equals 0.9472, so $J_n(t_n)<0$ exactly for $n\ge 12$. For such n, $F_\varepsilon(t_n)=\varepsilon\frac{b^2}{2}J_n(t_n)+O(\varepsilon^2)$ is negative for all small $\varepsilon>0$. □

Remark 3.1. The minima of J_n on $(0,\infty)$ sit at $\omega t=2\pi k+3\pi/2$, $k\ge 0$, where $J_n=(\delta-\omega e^{-\delta t})/(\delta^2+\omega^2)$ increases with k. So for $3\le n\le 11$ the first-order term (4) is positive for every $t>0$, and this family gives no counterexample at small ε. This fits the experiments of [1], whose cyclic generators had order four. Whether (2) holds for every MMPP of order at most 11 is open.

Proof of Theorem 1.3. Here π is uniform, $\lambda=10^{-3}$ and $\alpha_j=p_j/12000$. Let $r=2003/2000=1+\max_j d_j$ and $B=I+C/r$. Then $B_{jj}=(1500-p_j)/M$ and $B_{j,j+1}=10^6/M$ with $M=1001500$, so B is nonnegative with row sums at most one, and $0\le B^k\mathbf{1}\le\mathbf{1}$ for all k. With $a=rt=38057/4000$ at $t=19/2$, uniformization gives

$$e^a\big(S_\pi(t)-S_\alpha(t)\big)=\sum_{k\ge 0}\frac{a^k}{k!}\,wB^k\mathbf{1},\qquad w=\pi-\alpha.$$

Let $N=80$, let Σ be the sum over $k\le N$, $E=\sum_{k\le N}a^k/k!$, and

$$R=\frac{a^{N+1}}{(N+1)!}\Big(1-\frac{a}{N+2}\Big)^{-1}\ \ge\ \sum_{k>N}\frac{a^k}{k!}.$$

Then $E\le e^a\le E+R$, and the tail of the series has absolute value at most $\|w\|_1R$, where $\|w\|_1=3732/12000$. If $\Sigma+\|w\|_1R<0$, then $S_\pi(t)-S_\alpha(t)\le(\Sigma+\|w\|_1R)/(E+R)$. The quantities Σ, E and R are rational, and an exact computation (Section 6) gives

$$300000\,(\Sigma+\|w\|_1R)+E+R<0,$$

which is the claim. The same computation encloses $S_\pi(t)-S_\alpha(t)$ in an interval of width below 10^{-45} around $-3.40376335044947\times10^{-6}$. □

Remark 3.2. A cruder imitation of a square wave also works. On the directed 24-cycle with rate $2/3$ on states $0,\dots,11$ and 0 on states $12,\dots,23$, so that $\lambda=1/3$, one has $\mathbb{P}(T^\pi>79/4)=0.0142536155$ and $\mathbb{P}(T^\alpha>79/4)=0.0154255223$; the same uniformization bound with $N=140$ encloses the difference around -1.1719068×10^{-3}.

4. Reversible modulators

Proposition 4.1. *If Q is reversible with respect to π, then* (2) *holds.*

Proof. Let $\langle u,v\rangle_\pi=\sum_i\pi_iu_iv_i$. Reversibility makes Q, hence $A=D-Q=-C$, self-adjoint for this inner product. Let $v_1,\dots,v_m$ be an orthonormal eigenbasis of A with eigenvalues μ_k, and $a_k=\langle\mathbf{1},v_k\rangle_\pi^2$. Then $a_k\ge0$, $\sum_k a_k=\langle\mathbf{1},\mathbf{1}\rangle_\pi=1$, and

$$S_\pi(t)=\langle\mathbf{1},e^{-At}\mathbf{1}\rangle_\pi=\sum_k a_ke^{-\mu_kt},\qquad\lambda=\langle\mathbf{1},A\mathbf{1}\rangle_\pi=\sum_k a_k\mu_k,$$

the second because $Q\mathbf{1} = 0$. Let M take the value μ_k with probability a_k. By (3), $S_\alpha = -S'_\pi/\lambda$, so

$$S_\pi(t) - S_\alpha(t) = \mathbb{E}e^{-Mt} - \frac{\mathbb{E}[Me^{-Mt}]}{\mathbb{E}M} = -\frac{\mathrm{Cov}(M, e^{-Mt})}{\mathbb{E}M} \geq 0,$$

since the covariance of an increasing and a decreasing function of the same random variable is nonpositive. □

He [4, Theorem 3.4] uses reversibility in the same way to show that $E[X_x]$ is non-increasing in x.

5. Convexity in the speed of modulation

He [4] writes X_x for the phase-type variable with representation $(\pi, xQ - D)$, and $X_{D,x}$ for the one with representation $(\pi D/\lambda, xQ - D)$; for $x = 1$ these are T^π and T^α. He proves $E[X_x] = \pi(D - xQ)^{-1}\mathbf{1} \geq 1/\lambda$ and $\mathrm{SCV}(X_{D,x}) = 2\lambda E[X_x] - 1$ [4, Theorems 3.1 and 3.2]. Theorem 1.3 shows that the first property in [4, Section 4.3] fails at $x = 1$. The next proposition concerns the third.

Proposition 5.1. *Let Q be the directed 6-cycle and $D = \mathrm{diag}(\frac{1}{3}, \frac{2}{5}, \frac{2}{3}, 1, \frac{2}{3}, \frac{2}{5})$, and $f(x) = E[X_x] = \pi(D - xQ)^{-1}\mathbf{1}$ for $x \geq 0$. Then $f''(0) = -1$. Hence f, and $x \mapsto \mathrm{SCV}(X_{D,x})$, are strictly concave on $[0, x_0]$ for some $x_0 > 0$.*

Proof. D is invertible, so $D - xQ$ is invertible near $x = 0$ and f is smooth there, with $f''(0) = 2\pi D^{-1}QD^{-1}QD^{-1}\mathbf{1}$. With π uniform this is a finite computation: $D^{-1}\mathbf{1} = (3, \frac{5}{2}, \frac{3}{2}, 1, \frac{3}{2}, \frac{5}{2})$, and two further applications of $D^{-1}Q$, where $(Qv)_j = v_{j+1} - v_j$, followed by averaging and doubling give -1. The second claim follows from $\mathrm{SCV}(X_{D,x}) = 2\lambda f(x) - 1$ and continuity of f''. □

For this example $f(0) = 2$ and $f'(0) = -\frac{1}{4}$, and at positive speeds the exact values give $f(10^{-3}) + f(3 \cdot 10^{-3}) - 2f(2 \cdot 10^{-3}) = -9.739 \times 10^{-7} < 0$.

6. Computations

The script `code/checks.py` in [5] carries out the certificate of Theorem 1.3 in exact rational arithmetic, twice, with the series $wB^k\mathbf{1}$ computed once as $w(B^k\mathbf{1})$ and once as $(wB^k)\mathbf{1}$; it compares the result with a 40-digit matrix exponential and checks the opposite sign at $t = 1/2$. It does the same for Remark 3.2. It checks the sign of $J_n(t_n)$ for $3 \leq n \leq 400$, compares $F_\varepsilon/\varepsilon$ with $\frac{b^2}{2}J_n$ for $n = 12, 20$, and checks that $F_\varepsilon > 0$ on a grid of times for $n = 11$, $b = \frac{1}{2}$, $\varepsilon = 10^{-4}$. It computes the square-wave example of Section 2, tests Proposition 4.1 on 1763 random reversible MMPPs of orders 2 to 7, and computes $f''(0)$ and the convexity gap of Proposition 5.1 exactly. The file `lean/MMPP.lean` contains the inequality of Theorem 1.3, multiplied out to an inequality between integers, and the value $f''(0) = -1$ of Proposition 5.1, both checked in Lean 4 by evaluation in the kernel.

Acknowledgements and funding

The research presented in this paper was supported by the European Research Council (ERC) under the European Union's Horizon Europe research and innovation programme (grant agreement No. 101041711), by the Simons Foundation as part of the Collaboration on the Mathematical and Scientific Foundations of Deep Learning, by Heights Labs, by Convex Nexus Capital, by the Israel Science Foundation (grant number 2258/19), and by the Israel Science Foundation (ISF Grant 4101/25).

This work was done by multiple AI agents, Claude (Anthropic) and Codex (OpenAI), connected to Math Brain [6] and working together inside the Brain framework [8], a shared, persistent

memory in which the agents record claims, proofs, failed attempts and methods, retrieve one another's results, coordinate who works on what, and audit each other's work. For more on the details behind the Brain and Math Brain, see the Math Brain paper [7], the Math Brain repository [6] and the Brain framework repository [8].

Declaration of competing interest

The author holds equity in Heights Labs and in Convex Nexus Capital and has a management role at Convex Nexus Capital. These entities are included in the funding declaration above.

Declaration of generative AI and AI-assisted technologies

The paper comes out of Math Brain, an ongoing project on the use of AI systems for research mathematics. On 8 September 2026 a Claude agent (Anthropic), working in Claude Code, found the reformulation (3), Theorem 1.1 and the 24-state example of Remark 3.2; on the same day a Codex agent (OpenAI) found the cosine profile behind Theorems 1.2 and 1.3, the rational certificate of Theorem 1.3, the argument of Proposition 4.1 and the example of Proposition 5.1. On 18 October 2026 Claude Opus 5.5, working in Claude Code, checked the proofs line by line, wrote Theorem 1.2 and its proof, the computer checks, the Lean file and the paper. The author directed the work and is responsible for the content of the paper and for decisions concerning its dissemination.

Data availability

No data were used. The code is available at
`https://github.com/yspennstate/mmpp-stationary-order`.

The paper is part of the Math Brain project, whose public repository is `https://github.com/yspennstate/math-brain-public` [6].

References

[1] A. Asanjarani and Y. Nazarathy, *Stationary Markovian arrival processes: results and open problems*, ANZIAM J. **64** (2022), 54–68.

[2] A. Asanjarani and Y. Nazarathy, *Stationary Markovian arrival processes, results and open problems*, arXiv:1905.01736 (2019).

[3] A. Asanjarani and Y. Nazarathy, *The squared coefficient of variation for MMPP is greater than unity*, arXiv:1802.08400v1 (2018), withdrawn.

[4] Q.-M. He, *Bounds on the mean and squared coefficient of variation of phase-type distributions*, J. Appl. Probab. **58** (2021), 880–889.

[5] Y. Shmalo, *Code and Lean files for this paper*, `https://github.com/yspennstate/mmpp-stationary-order`.

[6] Y. Shmalo, *Math Brain: public snapshot, code and tests*, 2026, `https://github.com/yspennstate/math-brain-public`.

[7] Y. Shmalo, *Math Brain: shared, auditable memory for multi-agent mathematical research*, October 2026, `https://github.com/yspennstate/math-brain-public/tree/main/papers/math-brain-2026-10-24`.

[8] Y. Shmalo, *Brain Framework*, public software repository, 2026, `https://github.com/yspennstate/brain-framework-public`.

Einstein Institute of Mathematics, The Hebrew University of Jerusalem, Jerusalem, Israel
Email address: `yitzchak.shmalo@gmail.com`