A COUNTEREXAMPLE TO A CONJECTURE OF CERF ON SECOND-ORDER PIVOTS

YITZCHAK SHMALO

ABSTRACT. For an increasing subset A of $\{0,1\}^n$, Cerf considers the number $\phi_2(A)$ of pairs (ω, i) with $\omega \in A$ and i a positive pivot of order at most two at ω, and the analogous count $\phi_{2\setminus 1}(A)$ of pivots of order exactly two. He conjectured that every maximizer of either functional contains all configurations with at least $n/2+1$ ones and no configuration with fewer than $n/2-2$ ones, which would give $\phi_2(A) \le 6\sqrt{n}\,2^n$. We show that the conjecture fails. For $n = 10^{14}$, and more generally for every $n = h^2$ with $10 \mid h$ and $h > 2{,}592{,}000$, there is an increasing set with $\phi_{2\setminus 1}(A) \ge 2^n n/259200$, while every increasing set inside the conjectured band has $\phi_2(A) \le 10\sqrt{n}\,2^n$. Consequently no maximizer of either functional lies in the band. The set is a random monotone CNF of the kind used by Talagrand, and the proof is a first and second moment count over the clauses.

1. INTRODUCTION

Let $\Omega_n = \{0,1\}^n$ with the coordinatewise order, and identify a configuration ω with its support $\{i : \omega(i) = 1\}$. For $\omega \in \Omega_n$ and $i \in \{1,\dots,n\}$ write ω_i and ω^i for the configurations obtained from ω by setting the i-th coordinate to 0 and to 1. A set $A \subset \Omega_n$ is increasing if $\omega \in A$ and $\omega \le \eta$ imply $\eta \in A$. The positive pivots of A at ω are

$$P^+(A,\omega) = \{i :\ \omega_i \notin A,\ \omega^i \in A\}.$$

In his monograph on the percolation probability at criticality [1], Cerf defines positive pivots of higher order: for $k \ge 1$ and $I \subset \{1,\dots,n\}$ let ω_I be ω with every coordinate in I set to 0, and

$$P_k^+(A,\omega) = \big\{i :\ \exists I,\ |I| \le k-1,\ \omega(j) = 1 \text{ for all } j \in I,\ i \in P^+(A,\omega_I)\big\} \tag{1.1}$$

[1, (49.1)]. Thus $P_1^+ = P^+$, and $P_2^+(A,\omega)$ consists of the first pivots at ω together with the first pivots at the configurations obtained by switching off one occupied coordinate. Cerf studies the functionals

$$\phi_{2\setminus 1}(A) = \sum_{\omega\in A} \big|P_2^+(A,\omega) \setminus P_1^+(A,\omega)\big|, \qquad \phi_2(A) = \sum_{\omega\in A} \big|P_2^+(A,\omega)\big|$$

and their maxima $M_{2\setminus 1}(n)$ and $M_2(n)$ over increasing sets [1, (51.1), (52.1)]. For first-order pivots the corresponding maximum is attained by small perturbations of the majority set $\{\omega : |\omega| > n/2\}$ and is of order $\sqrt{n}\,2^n$ [1, §50]. Exhaustive computations for $n \le 7$ led Cerf to the following statement [1, Conjecture 53.1].

Conjecture 1.1 (Cerf)**.** *Let $n \ge 1$. If A^* is an increasing subset of Ω_n which realizes the maximum of $\phi_{2\setminus 1}$ or of ϕ_2, then*

$$\{\omega :\ |\omega| \ge n/2+1\} \subset A^* \subset \{\omega :\ |\omega| \ge n/2-2\}.$$

Date: 5 November 2026.

2020 *Mathematics Subject Classification.* 06E30, 05D40, 60K35, 60C05.

Key words and phrases. increasing sets, Boolean functions, pivotal coordinates, second-order pivots, random CNF, Talagrand's construction.

The conjecture would give $\max(M_{2\setminus 1}(n), M_2(n)) \le 6\sqrt{n}\,2^n$, that is, $\mathbb{E}|P_2^+(A)| \le 6\sqrt{n}$ under the uniform measure for every increasing A [1, (53.2)], the analogue of the first-order bound. We show that it is false.

Theorem 1.2. *Let $n = h^2$ where h is divisible by 10 and $h > 2{,}592{,}000$; for instance $n = 10^{14}$. There is an increasing set $A \subset \Omega_n$ with*

$$2^{-n}\phi_{2\setminus 1}(A) \;\ge\; \frac{n}{259200} \;>\; 10\sqrt{n}.$$

Every increasing set B with $\{|\omega| \ge n/2+1\} \subset B \subset \{|\omega| \ge n/2-2\}$ satisfies $2^{-n}\phi_2(B) \le 10\sqrt{n}$. Consequently, for this n, no maximizer of $\phi_{2\setminus 1}$ and no maximizer of ϕ_2 satisfies the inclusions of Conjecture 1.1.

Since $\phi_2 \ge \phi_{2\setminus 1}$, the last sentence follows from the two inequalities: a maximizer A^* of either functional has $2^{-n}F(A^*) \ge 2^{-n}F(A) > 10\sqrt{n}$, which no set in the band attains.

The set A is a random monotone CNF with 2^r clauses of r coordinates each, $r = h/10$. This is the construction Talagrand used to exhibit an increasing set whose first-order pivot count is of order $\sqrt{n}$ on a set of probability bounded below [2]. In Section 62 of [1], Cerf adapts Talagrand's set to second-order pivots and argues that $\mathbb{P}(|P_2^+(A^\circ)| \ge cn) \ge c$ for large n; his Proposition 62.1 records a lower bound on $M_2(n)$ of order n. A bound of the form $\mathbb{E}|P_2^+(A)| \ge cn$ is incompatible with the estimate (53.2) of [1] for large n, so Section 62 already points to the failure of the conjecture, although it does not return to Conjecture 53.1. The present note gives a complete argument with explicit constants and an explicit dimension. Two points need care. Coordinates are sampled with replacement, so a clause may contain a coordinate several times, and a clause whose positive part is a single repeated coordinate still produces a first pivot; and a pivot of order two must be certified through a partner coordinate whose removal keeps the configuration in A. The pair witness of Lemma 2.1 and the clause graph of Section 5 handle both.

There is also a small ambiguity in (1.1). Read literally, the definition allows a coordinate i with $\omega(i) = 0$ to belong to $P_2^+(A,\omega)$: take j occupied with $\omega_j \notin A$ but $(\omega_j)^i \in A$. The tables in [1, §§51–52] agree instead with the restriction of $P_2^+(A,\omega)$ to occupied coordinates. For instance, for $A = \{10, 01, 11\}$ the literal reading gives $\phi_2(A) = 6$ and the occupied reading gives 4; for $n \le 5$ the occupied reading reproduces the maxima $1, 4, 9, 24, 62$ of ϕ_2 and $0, 2, 6, 12, 38$ of $\phi_{2\setminus 1}$ (Section 7). Theorem 1.2 holds under both readings: the lower bound counts only occupied pivots and the upper bound only uses $|P_2^+(A,\omega)| \le n$.

2. Pivots of order two

Let A be increasing and $\omega \in A$. For $i \notin \omega$ we have $\omega_i = \omega \in A$, so

$$P_1^+(A,\omega) = \{i \in \omega :\ \omega \setminus \{i\} \notin A\}. \tag{2.1}$$

Lemma 2.1. *Let A be increasing, $\omega \in A$, and let $i \ne j$ be occupied coordinates of ω such that $\omega \setminus \{j\} \in A$ and $\omega \setminus \{i,j\} \notin A$. Then $i \in P_2^+(A,\omega)$. If moreover $\omega \setminus \{i\} \in A$, then $i \notin P_1^+(A,\omega)$.*

Proof. Take $I = \{j\}$ in (1.1); j is occupied. With $\eta = \omega_j = \omega \setminus \{j\}$ we have $\eta^i = \eta \in A$ because $i \in \eta$, and $\eta_i = \omega \setminus \{i,j\} \notin A$. So $i \in P^+(A,\eta) \subset P_2^+(A,\omega)$. The second statement is (2.1). □

Conversely, an occupied $i \in P_2^+(A,\omega) \setminus P_1^+(A,\omega)$ always has such a partner j, but only the direction of Lemma 2.1 is used below. The program in Section 7 checks on every increasing set with $n \le 5$ that the occupied part of $P_2^+(A,\omega)$ is exactly the set of first pivots together with the coordinates having a partner as in the lemma.

3. The bound inside the band

Lemma 3.1. *Let $n = 2d \geq 2$ and let B be increasing with $\{|\omega| \geq d+1\} \subset B \subset \{|\omega| \geq d-2\}$. Then $2^{-n}\phi_2(B) \leq 10\sqrt{n}$, under either reading of* (1.1).

Proof. Let $\omega \in B$ with $|\omega| \geq d+3$. Switching off at most one occupied coordinate in (1.1), and then testing one more coordinate at 0, leaves a configuration with at least $d+1$ ones, which lies in B. Hence $P_2^+(B,\omega) = \emptyset$. Every $\omega \in B$ has $|\omega| \geq d-2$, so only the five levels $d-2 \leq |\omega| \leq d+2$ contribute. Each level has at most $\binom{2d}{d}$ elements and each contributes at most n, so $\phi_2(B) \leq 5n\binom{2d}{d}$. Put $b_d = \binom{2d}{d}4^{-d}$. Then $b_0 = 1$ and $b_{d+1}/b_d = (2d+1)/(2d+2)$, and $b_d \leq (d+1)^{-1/2}$ follows by induction from $(2d+1)^2(d+2) \leq 4(d+1)^3$, whose difference is $3d+2$. Since $n \leq 4(d+1)$, we get $b_d \leq 2/\sqrt{n}$ and $2^{-n}\phi_2(B) \leq 5n \cdot 2/\sqrt{n} = 10\sqrt{n}$. □

The constant is cruder than the $6\sqrt{n}$ of [1, (53.2)] because five levels can contribute rather than four, and we use the elementary bound on b_d.

4. A random monotone CNF

Fix h divisible by 10 with $h \geq 30$, and put

$$n = h^2, \qquad r = h/10, \qquad M = 2^r,$$

so that $n = 100r^2$. Let $Y = (Y_{a,b})_{1\leq a\leq r,\, 1\leq b\leq M}$ be independent and uniform on $\{1,\dots,n\}$; clause b is the list $(Y_{1,b},\dots,Y_{r,b})$, repetitions allowed. Let

$$A_Y = \{\omega : \text{ for every } b \text{ there is } a \text{ with } Y_{a,b} \in \omega\}.$$

For each Y this is an increasing set. All probabilities below are over Y on a finite space; no array is constructed explicitly.

Fix ω with $s = |\omega|$ and $|s - n/2| \leq h$, and put $z = 1 - s/n$. Then $|2z-1| \leq 2/h$, and $n/3 \leq s-1 < s \leq 2n/3$. A clause misses ω with probability $q_0 = z^r$. The events that the individual clauses meet ω are independent, so conditionally on $\omega \in A_Y$ the clauses remain independent, each distributed as a clause conditioned to meet ω. We write $\mathbb{P}_\omega$ and $\mathbb{E}_\omega$ for this conditional law.

We use two elementary inequalities: $(1+u)^k \leq (1-ku)^{-1}$ for $u \geq 0$ and $ku < 1$, from $\binom{k}{j} \leq k^j$; and Bernoulli's inequality $(1-u)^k \geq 1-ku$ for $0 \leq u \leq 1$. Since $r \cdot 2/h = 1/5$ and $r(2/h + 2/n) < 1/4$,

$$(2z)^a \leq \tfrac{5}{4} \;\; (0 \leq a \leq r), \qquad (2z)^{r-2} \geq \tfrac{4}{5}, \qquad (2z + 2/n)^{r-1} \leq \tfrac{4}{3}. \tag{4.1}$$

In particular $Mq_0 = (2z)^r \leq 5/4$ and $q_0 \leq 1/2$, so, using $\log(1-u) \geq -2u$ on $[0,1/2]$ and $e^4 < 81$,

$$\mathbb{P}(\omega \in A_Y) = (1-q_0)^M \geq e^{-2Mq_0} \geq e^{-4} > \tfrac{1}{81}. \tag{4.2}$$

5. The clause graph

Fix Y with $\omega \in A_Y$. For a clause b, its positive positions are the a with $Y_{a,b} \in \omega$. Let S be the set of coordinates i such that, for some clause, the set of values at its positive positions is $\{i\}$; the clause may contain i several times. By (2.1), $S = P_1^+(A_Y,\omega)$: removing i from ω empties some clause exactly when such a clause exists.

Let G be the multigraph on ω with one edge $\{i,j\}$ for each clause that has exactly two positive positions, carrying two distinct values $i \neq j$. Let C be the set of vertices covered by edges, call an edge bad if one of its endpoints lies in S, and let B be the number of bad edges counted with multiplicity.

Lemma 5.1. $|P_2^+(A_Y,\omega)\setminus P_1^+(A_Y,\omega)| \geq |C| - 2B$.

Proof. Let V be the set of endpoints of bad edges; $|V| \leq 2B$. Let $i \in C \setminus V$ and choose an edge $\{i,j\}$, coming from clause b. Neither i nor j is in S, so $\omega\setminus\{i\}$ and $\omega\setminus\{j\}$ lie in A_Y. The positive part of clause b is $\{i,j\}$, so $\omega\setminus\{i,j\} \notin A_Y$. By Lemma 2.1, $i \in P_2^+ \setminus P_1^+$. □

6. Proof of Theorem 1.2

We keep the notation of Section 4, with ω fixed in the central range.

Coverage. For $i \in \omega$, a clause supplies an edge at i with probability

$$a_i = r(r-1)\,\frac{s-1}{n^2}\,z^{r-2}:$$

choose the two positive positions, which of them carries i, and the other value among the $s-1$ remaining coordinates of ω, the other $r-2$ positions falling outside ω. Under $\mathbb{P}_\omega$ the probability is $t_i = a_i/(1-q_0) \in [a_i, 2a_i]$. By (4.1), $r(r-1) \geq r^2/2$ and $(s-1)/n \in [1/3, 2/3]$,

$$Mt_i \geq Ma_i = 4r(r-1)\frac{s-1}{n^2}(2z)^{r-2} \geq \frac{8}{15}\frac{r^2}{n}, \qquad Mt_i \leq \frac{32}{3}\frac{r^2}{n} = \frac{8}{75} < 1.$$

The clauses being independent under $\mathbb{P}_\omega$, the Bonferroni inequality gives $\mathbb{P}_\omega(i \in C) \geq Mt_i - \binom{M}{2}t_i^2 \geq Mt_i(1 - Mt_i/2) \geq Mt_i/2 \geq r^2/(4n)$, and summing over the $s \geq n/3$ coordinates of ω,

$$\mathbb{E}_\omega|C| \geq r^2/12. \tag{6.1}$$

Bad edges. The probability that a clause has exactly one distinct positive value, and that value is a given $i \in \omega$, is $(z+1/n)^r - z^r \leq (r/n)(z+1/n)^{r-1}$. These events are disjoint over i, so the probability q_S that a clause is a singleton clause satisfies $q_S \leq (sr/n)(z+1/n)^{r-1}$, while a clause supplies an edge with probability $q_E = \binom{r}{2}(s/n)^2(1-1/s)z^{r-2}$. By (4.1), with $Q_S = q_S/(1-q_0) \leq 2q_S$ and $Q_E = q_E/(1-q_0) \leq 2q_E$,

$$MQ_S \leq 2\cdot\frac{2}{3}r\cdot 2\cdot\frac{4}{3} = \frac{32}{9}r < 6r, \qquad MQ_E \leq 2\cdot\frac{r^2}{2}\cdot\frac{4}{9}\cdot 4\cdot\frac{5}{4} < 4r^2.$$

A clause that supplies an edge is not a singleton clause. Fix an edge clause b and one of its two endpoints e; every other clause is independent of b under $\mathbb{P}_\omega$ and, by symmetry among the coordinates of ω, is a singleton clause at e with probability Q_S/s. A union bound over the edge clause, its two endpoints and the other $M-1$ clauses gives

$$\mathbb{E}_\omega B \leq MQ_E\cdot 2(M-1)\frac{Q_S}{s} \leq 4r^2\cdot 6r\cdot\frac{2\cdot 3}{n} = \frac{144r^3}{n}. \tag{6.2}$$

No independence between different pivot indicators is used, and repeated edges are counted in B with multiplicity.

Conclusion. By Lemma 5.1, (6.1) and (6.2),

$$\mathbb{E}_\omega\big|P_2^+(A_Y,\omega)\setminus P_1^+(A_Y,\omega)\big| \geq \frac{r^2}{12} - \frac{288r^3}{n} \geq \frac{r^2}{24},$$

the last step because $n = 100r^2 \geq 6912r$. Now let ω be uniform on Ω_n and independent of Y. Its size has mean $n/2$ and variance $n/4 = h^2/4$, so by Chebyshev's inequality $\mathbb{P}(||\omega| - n/2| \leq h) \geq 3/4$. Combining with (4.2),

$$\mathbb{E}_Y\big[2^{-n}\phi_{2\setminus 1}(A_Y)\big] = \mathbb{E}_\omega\mathbb{E}_Y\Big[\mathbf{1}_{\omega\in A_Y}\big|P_2^+(A_Y,\omega)\setminus P_1^+(A_Y,\omega)\big|\Big] \geq \frac{3}{4}\cdot\frac{1}{81}\cdot\frac{r^2}{24} = \frac{r^2}{2592} = \frac{n}{259200}.$$

All sums are finite. Hence some array Y_0 satisfies $2^{-n}\phi_{2\setminus 1}(A_{Y_0}) \geq n/259200$. Finally, $n/259200 > 10\sqrt{n}$ exactly when $h >$ 2,592,000, and Lemma 3.1 applies because n is even. This proves Theorem 1.2. □

Remark 6.1. The argument gives an increasing set with $\mathbb{E}|P_2^+(A) \setminus P_1^+(A)| \geq cn$ for an absolute $c > 0$ along $n = h^2$, $10 \mid h$, so $M_{2\setminus 1}(n)$ and $M_2(n)$ are of order $n2^n$ along this sequence, the trivial upper bound being $n2^n$. The constants have not been optimized, and we do not know the smallest dimension at which the conjecture fails; it holds for $n \leq 7$ [1].

7. Computations

The program `code/checks.py` in the repository [3] uses only the Python standard library and runs in a few seconds. It enumerates the $3, 6, 20, 168, 7581$ increasing subsets of Ω_n for $n = 1, \ldots, 5$ and computes both functionals under both readings of (1.1). The literal reading gives the maxima $\phi_2 = 1, 6, 15, 36, 100$ and $\phi_{2\setminus 1} = 0, 4, 12, 24, 76$; the occupied reading gives $1, 4, 9, 24, 62$ and $0, 2, 6, 12, 38$, the values in the tables of [1]. In every case every maximizer lies in the band, in agreement with Cerf's verification for $n \leq 7$. The program also checks Lemma 5.1 on all 729 arrays with $n = 3$, $r = 2$, $M = 3$, which give 2640 pairs (Y, ω) with $\omega \in A_Y$, and checks the rational inequalities used above for $h = 10^7$ and $h = 2{,}592{,}010$. The Lean 4 file `lean/Certificate.lean` proves the deterministic step of Lemmas 2.1 and 5.1 for arbitrary clause and coordinate types and the numerical margins at $h = 10^7$ with Mathlib's `norm_num`; the probabilistic estimates are not formalized.

Acknowledgements and funding

The research presented in this paper was supported by the European Research Council (ERC) under the European Union's Horizon Europe research and innovation programme (grant agreement No. 101041711), by the Simons Foundation as part of the Collaboration on the Mathematical and Scientific Foundations of Deep Learning, by Heights Labs, by Convex Nexus Capital, by the Israel Science Foundation (grant number 2258/19), and by the Israel Science Foundation (ISF Grant 4101/25). Computational resources were provided in part by Google Cloud through its education credits program (credit 530550103).

Declaration of competing interest

The author holds equity in Heights Labs and in Convex Nexus Capital and has a management role at Convex Nexus Capital. These entities are included in the funding declaration above.

Declaration of generative AI and AI-assisted technologies

The paper comes out of Math Brain, an ongoing project on the use of AI systems for research mathematics. In October 2026 OpenAI GPT models, running as research tasks of that project, gave the proof of Theorem 1.2 at $n = 10^{14}$, including the clause graph and the treatment of repeated coordinates, and wrote the Lean file and the first version of the verification programs. Claude (Anthropic) checked the proof independently in October 2026; in November 2026 Claude Opus 5.5, working in Claude Code, extended the statement to the dimensions $n = h^2$ of Theorem 1.2, rewrote the verification program and wrote the paper. The author directed the work and is responsible for the content of the paper and for decisions concerning its dissemination.

Data availability

No data were used. The code is available at
`https://github.com/yspennstate/second-order-pivots-cerf-conjecture`.

References

[1] R. Cerf, $\theta(p_c, \mathbb{Z}^d) = 0$?, arXiv:2608.23661v1, 2026.
[2] M. Talagrand, How much are increasing sets positively correlated?, *Combinatorica* **16** (1996), 243–258.
[3] Y. Shmalo, Code for "A counterexample to a conjecture of Cerf on second-order pivots", `https://github.com/yspennstate/second-order-pivots-cerf-conjecture`, 2026.

Einstein Institute of Mathematics, The Hebrew University of Jerusalem, Jerusalem, Israel
Email address: `yitzchak.shmalo@gmail.com`