THE CONJECTURED DOMINATING PAIR FOR DP-SGD WITH SHUFFLED BATCHES IS NOT DOMINATING

YITZCHAK SHMALO

Abstract. Chua, Ghazi, Kamath, Kumar, Manurangsi, Sinha and Zhang conjectured that the pair of Gaussian mixtures $P_{\mathcal{S}} = \frac{1}{T}\sum_t \mathcal{N}(2e_t, \sigma^2 I)$, $Q_{\mathcal{S}} = \frac{1}{T}\sum_t \mathcal{N}(e_t, \sigma^2 I)$ tightly dominates the adaptive batch linear query mechanism with shuffled batches, for every adaptive query method. We show that the conjecture fails for every number of steps $T \geq 2$ and every batch size. The query method that breaks it answers as the conjectured worst case at every step but the last, and at the last step changes the query when the earlier outputs suggest that the differing record was in an early batch. At $T = 2$, $\sigma = 1$ and $\varepsilon = \frac{1}{2}$ its hockey-stick divergence is 0.22323 against 0.22127 for the conjectured pair, and an interval-arithmetic certificate bounds the excess below by $1.04 \cdot 10^{-4}$. A direct analytic last-step argument gives strict failure at every $T \geq 2$; a separate certificate at $T = 3$ gives a quantitative bound. The same query method also beats the pair in Kullback–Leibler divergence and in Rényi divergence of every order we tested, from $\frac{1}{4}$ to 5, in both directions.

1. Introduction

DP-SGD is usually analysed as if every batch were drawn by Poisson subsampling, while implementations mostly shuffle the data and cut it into batches of a fixed size. Chua, Ghazi, Kamath, Kumar, Manurangsi, Sinha and Zhang [1] studied the gap between the two through the adaptive batch linear query mechanism ABLQ, which isolates what the gradient steps release. For Poisson subsampling and for the deterministic batch sampler the worst case over adaptive query methods is known exactly. For the shuffle sampler $\mathcal{S}$ they gave the pair

$$P_{\mathcal{S}} = \frac{1}{T}\sum_{t=1}^{T} \mathcal{N}(2e_t, \sigma^2 I_T), \qquad Q_{\mathcal{S}} = \frac{1}{T}\sum_{t=1}^{T} \mathcal{N}(e_t, \sigma^2 I_T), \tag{1}$$

which is realised by a non-adaptive query method and so gives lower bounds on the privacy loss of $\mathrm{ABLQ}_{\mathcal{S}}$, and they conjectured that these lower bounds are tight.

Conjecture 1.1 ([1, Conjecture 3.2])**.** *The pair $(P_{\mathcal{S}}, Q_{\mathcal{S}})$ tightly dominates $\mathrm{ABLQ}_{\mathcal{S}}$ for all adaptive query methods $\mathcal{A}$.*

The conjecture is restated as open by van Dijk and Ertan [4, Appendix J], who show that the smaller pair $(\mathcal{N}(0, I), \frac{1}{M}\sum_j \mathcal{N}(e_j, I))$ does not dominate the shuffled mechanism; their example is non-adaptive and does not bear on Conjecture 1.1. Random allocation, a different sampler that also fixes how many steps each record takes part in, is analysed by Feldman and Shenfeld [2].

We show that Conjecture 1.1 is false.

Date: Revised 22 October 2026.

2020 *Mathematics Subject Classification.* 68P27, 62B10, 94A17, 60E15.

Key words and phrases. differential privacy, DP-SGD, shuffling, dominating pair, hockey-stick divergence, Rényi divergence, adaptive composition, interval arithmetic.

Theorem 1.2. *Let $\sigma = 1$ and $\varepsilon = \frac{1}{2}$. For every $T \geq 2$, every batch size $b \geq 1$ and every dimension $d \geq 1$ there are an adaptive query method $\mathcal{A}$ and zero-out adjacent datasets $x \to_z x'$ with*

$$H_{e^\varepsilon}(\mathrm{ABLQ}_{\mathcal{S}}(x) \,\|\, \mathrm{ABLQ}_{\mathcal{S}}(x')) > H_{e^\varepsilon}(P_{\mathcal{S}} \| Q_{\mathcal{S}}).$$

At $T = 2$ the excess is at least $1.04 \cdot 10^{-4}$ and at $T = 3$ at least $2.25 \cdot 10^{-5}$.

Numerically, at $T = 2$ the left side is 0.223234 and $H_{e^\varepsilon}(P_{\mathcal{S}} \| Q_{\mathcal{S}}) = 0.221269$; the reverse orientation $H_{e^\varepsilon}(Q_{\mathcal{S}} \| P_{\mathcal{S}}) = 0.205107$ is smaller still, so the example also breaks a symmetric reading of domination. On a grid of non-adaptive query methods the best one is the conjectured one, so the gain comes from adaptivity.

The query method is simple. All records but one hold the same value w, the remaining record holds u in x and $\perp$ in x'. At every step the conjectured method asks for $+1$ on u and -1 on w. Our method does the same at steps $1, \ldots, T-1$ and, when the outputs so far point to the record having been in one of the early batches, asks for $+1$ on both u and w at step T. Section 2 sets up the mechanism and reduces the effect of such a switch to one function Δ of two variables, the same for every T. Section 3 proves the case $T = 2$ and Section 4 the cases $T \geq 3$. Section 5 treats Kullback–Leibler and Rényi divergences, and Section 6 records numerical values for larger T, other σ and other ε.

Chua et al. note that their results do not rely on the conjecture, since the pair (1) is used only for lower bounds; those bounds stand. What changes is that the gap between $\mathrm{ABLQ}_{\mathcal{S}}$ and Poisson subsampling is at least slightly larger than the pair shows, and that the exact privacy of one shuffled epoch is again open.

2. Setup and the last-step switch

2.1. **The mechanism.** A dataset is $x = (x_1, \ldots, x_n)$ with $x_i \in \mathcal{X} \cup \{\perp\}$ and $n = bT$. The shuffle sampler draws a uniformly random permutation π of $[n]$ and sets $S_t = \{\pi((t-1)b+1), \ldots, \pi(tb)\}$. An adaptive query method $\mathcal{A}$ maps the earlier outputs to a query $\psi_t = \mathcal{A}(g_1, \ldots, g_{t-1}; \cdot)\colon \mathcal{X} \to \mathbb{B}^d$, the closed unit ball, with $\psi_t(\perp) = 0$, and

$$g_t = \sum_{i \in S_t} \psi_t(x_i) + \xi_t, \qquad \xi_t \sim \mathcal{N}(0, \sigma^2 I_d) \text{ independent.}$$

$\mathrm{ABLQ}_{\mathcal{S}}(x)$ is the law of $(g_1, \ldots, g_T)$. Datasets are zero-out adjacent, $x \to_z x'$, if they agree except at one index i, where $x_i \in \mathcal{X}$ and $x'_i = \perp$.

For $\gamma \geq 0$ the hockey-stick divergence of probability measures with densities p, q is $H_\gamma(P \| Q) = \int (p - \gamma q)_+$. Following [1, Definition 2.3], a pair (P, Q) dominates a pair (A, B) if $H_{e^\varepsilon}(P \| Q) \geq H_{e^\varepsilon}(A \| B)$ for all $\varepsilon \in \mathbb{R}$; it dominates a mechanism M if it dominates $(M(x), M(x'))$ for all $x \to_z x'$, and it does so tightly if in addition it is dominated by $(M(x), M(x'))$ for some such pair. To refute Conjecture 1.1 it suffices to exhibit one $\mathcal{A}$, one adjacent pair and one ε with $H_{e^\varepsilon}(M(x) \| M(x')) > H_{e^\varepsilon}(P_{\mathcal{S}} \| Q_{\mathcal{S}})$.

2.2. **The datasets and the two queries.** We work with $d = 1$; for $d > 1$ put the queries on the first coordinate and the others carry pure noise. Let $\mathcal{X} = \{u, w\}$, let x have $x_1 = u$ and $x_j = w$ for $j \geq 2$, and let x' have $x'_1 = \perp$ and $x'_j = w$ for $j \geq 2$. Two queries are used:

$$\mathrm{A}: \ \psi(u) = +1, \ \psi(w) = -1, \qquad\qquad \mathrm{S}: \ \psi(u) = \psi(w) = +1.$$

Write $z_t = g_t - b\,\psi_t(w)$; the method knows $\psi_t(w)$, so z_t is a function of its own outputs. If the record is not in S_t, then S_t holds b copies of w and $z_t \sim \mathcal{N}(0, \sigma^2)$ under both datasets. If it is, then

$$\begin{aligned} z_t \sim \mathcal{N}(2, \sigma^2) \text{ under } x, \quad z_t \sim \mathcal{N}(1, \sigma^2) \text{ under } x' \qquad \text{for A,} \\ z_t \sim \mathcal{N}(0, \sigma^2) \text{ under } x, \quad z_t \sim \mathcal{N}(-1, \sigma^2) \text{ under } x' \qquad \text{for S.} \end{aligned}$$

The batch size has dropped out. Since π is uniform, the record lies in each batch with probability $1/T$, and given the batch the noise terms are independent.

Lemma 2.1. *If $\mathcal{A}$ asks* A *at every step, the law of $(z_1, \dots, z_T)$ is $P_{\mathcal{S}}$ under x and $Q_{\mathcal{S}}$ under x'.*

Proof. Given that the record lies in S_t, the vector z is $\mathcal{N}(2e_t, \sigma^2 I_T)$ under x and $\mathcal{N}(e_t, \sigma^2 I_T)$ under x'. Average over t. □

The map $g \mapsto z$ is a common triangular Borel bijection under both datasets: the query at step t is determined by earlier outputs, so its translation can be inverted recursively. For the last-step switch below, the prefix translation is fixed and the final translation is a Borel function of the prefix. Fubini's theorem shows that this map preserves Lebesgue measure, and the hockey-stick divergence is unchanged. For the all-A method it is a fixed translation.

2.3. **The last-step switch.** Let $m = T - 1$, write ϕ for the $\mathcal{N}(0, \sigma^2)$ density and, for $y \in \mathbb{R}^m$,

$$\alpha(y) = \frac{1}{T}\sum_{t=1}^{m} \phi(y_t - 2) \prod_{s \neq t} \phi(y_s), \quad \gamma(y) = \frac{1}{T}\sum_{t=1}^{m} \phi(y_t - 1) \prod_{s \neq t} \phi(y_s), \quad \beta(y) = \frac{1}{T}\prod_{t=1}^{m} \phi(y_t). \tag{2}$$

For $a, c \in \mathbb{R}$ and $\alpha, \beta, \gamma \geq 0$ set

$$I_{a,c}(\alpha, \beta, \gamma) = \int_{\mathbb{R}} \Big(\alpha\phi(s) + \beta\phi(s - a) - e^{\varepsilon}\big(\gamma\phi(s) + \beta\phi(s - c)\big)\Big)_+ \, ds,$$

and write $I_{\mathrm{A}} = I_{2,1}$, $I_{\mathrm{S}} = I_{0,-1}$.

Lemma 2.2. *Let $\mathcal{A}_S$ ask* A *at steps $1, \dots, m$ and, at step T, ask* S *if $(z_1, \dots, z_m) \in S$ and* A *otherwise, where $S \subseteq \mathbb{R}^m$ is Borel. Then*

$$H_{e^\varepsilon}(M(x) \| M(x')) = H_{e^\varepsilon}(P_{\mathcal{S}} \| Q_{\mathcal{S}}) + \int_S (I_{\mathrm{S}} - I_{\mathrm{A}})(\alpha(y), \beta(y), \gamma(y)) \, dy.$$

Proof. Condition on the batch of the record. If it is S_t with $t \leq m$, the prefix $y = (z_1, \dots, z_m)$ has density $\phi(y_t - 2)\prod_{s \neq t}\phi(y_s)$ under x and $\phi(y_t - 1)\prod_{s \neq t}\phi(y_s)$ under x', and $z_T \sim \mathcal{N}(0, \sigma^2)$ whatever is asked at step T. If it is S_T, the prefix has density $\prod \phi(y_s)$ under both datasets and z_T has density $\phi(\cdot - a)$ under x and $\phi(\cdot - c)$ under x', with $(a, c) = (2, 1)$ for A and $(0, -1)$ for S. So the joint densities are

$$p(y, s) = \alpha(y)\phi(s) + \beta(y)\phi(s - a(y)), \qquad q(y, s) = \gamma(y)\phi(s) + \beta(y)\phi(s - c(y)),$$

with $(a(y), c(y))$ fixed by whether $y \in S$. Integrating $(p - e^\varepsilon q)_+$ first in s gives

$$H_{e^\varepsilon}(M(x) \| M(x')) = \int I_{a(y),c(y)}(\alpha(y), \beta(y), \gamma(y)) \, dy.$$

With $S = \emptyset$ this is $H_{e^\varepsilon}(P_{\mathcal{S}} \| Q_{\mathcal{S}})$ by Lemma 2.1, and the difference is the integral over S. □

The best S is therefore $\{I_{\mathrm{S}} > I_{\mathrm{A}}\}$, and the conjecture fails as soon as this set has positive measure.

2.4. **Two variables.** $I_{a,c}$ is positively homogeneous of degree one in (α, β, γ). At $\sigma = 1$ put $x_t = e^{y_t - 1/2}$; then $\phi(y_t - 2)/\phi(y_t) = e^{-1}x_t^2$ and $\phi(y_t - 1)/\phi(y_t) = x_t$, so

$$\frac{\alpha}{\beta} = A := e^{-1}\sum_{t=1}^{m} x_t^2, \qquad \frac{\gamma}{\beta} = G := \sum_{t=1}^{m} x_t, \tag{3}$$

and

$$(I_{\mathrm{S}} - I_{\mathrm{A}})(\alpha, \beta, \gamma) = \beta\,\Delta(A, G), \qquad \Delta(A, G) := I_{\mathrm{S}}(A, 1, G) - I_{\mathrm{A}}(A, 1, G). \tag{4}$$

The function Δ does not depend on T; only the set of (A, G) that the prefix can reach does. For $m = 1$ it is the curve $eA = G^2$. For $m \geq 2$, since $x_t > 0$,

$$\frac{G^2}{m} \leq eA < G^2, \tag{5}$$

with equality on the left only when all x_t are equal. Conversely every point of $\{G^2/m < eA < G^2\}$ is reached: on the open simplex $\{x \in (0,\infty)^m : \sum x_t = G\}$ the function $\sum x_t^2$ is continuous, equals G^2/m at the centre and tends to G^2 at a vertex, and the simplex is connected.

2.5. **A direct analytic proof.** The qualitative assertion of Theorem 1.2 does not require numerical certification. At $\sigma = 1$ and $\varepsilon = \frac{1}{2}$, write

$$C(y) = \frac{\alpha(y) - \sqrt{e}\,\gamma(y)}{\beta(y)} = \sum_{i=1}^{T-1} (e^{2y_i - 2} - e^{y_i}).$$

Choose $S = [3,4]^{T-1}$. For every $y \in S$, $C(y) \geq (T-1)e^3(e-1) > e^2/4$. Let h_{A} and h_{S} be the signed joint densities $p - \sqrt{e}\,q$ at this prefix. With $v = e^{s-1}$,

$$\frac{h_{\mathrm{A}}(y,s)}{\beta(y)\phi(s)} = C(y) + v^2 - ev \geq C(y) - e^2/4 > 0.$$

On the other hand,

$$\frac{h_{\mathrm{S}}(y,s)}{\beta(y)\phi(s)} = C(y) + 1 - e^{-s} < 0 \quad \text{for } s < -\log(C(y)+1).$$

Both signed densities have integral $\beta(y)(C(y) + 1 - \sqrt{e})$ in s. Hence

$$I_{\mathrm{S}}(y) - I_{\mathrm{A}}(y) = \int_{\mathbb{R}} (-h_{\mathrm{S}}(y,s))_+ \, ds > 0.$$

The last inequality follows because the negative region contains a nonempty interval and $\beta(y)\phi(s) > 0$. The difference is measurable and integrable: both positive-part integrals are bounded by the same prefix marginal of p. Since S has positive measure, Lemma 2.2 proves Theorem 1.2 for every $T \geq 2$. The construction uses only unit-norm scalar queries and works at every batch size; embedding the scalar queries in one coordinate adds identical independent Gaussian noise in all other coordinates, proving the assertion for every $d \geq 1$.

This argument also works at every $y \geq 3$ when $T = 2$, so the improvement region has no finite upper endpoint. It establishes strict failure of the proposed pair but does not identify the tight shuffled privacy accountant. The certificates below supply explicit quantitative gains on different switch sets.

3. Two steps

At $T = 2$ the prefix is $y = z_1$, and with $\sigma = 1$, $\varepsilon = \frac{1}{2}$, $\alpha = \frac{1}{2}\phi(z_1 - 2)$, $\gamma = \frac{1}{2}\phi(z_1 - 1)$, $\beta = \frac{1}{2}\phi(z_1)$. Both inner integrals have closed forms. Dividing the integrand of I_{A} by $\phi(s)$ and writing $\tau = e^s$, the set where it is positive is

$$\{\beta e^{-2}\tau^2 - \beta\tau + (\alpha - \sqrt{e}\,\gamma) > 0\},$$

an interval condition on the roots of a quadratic in τ; for I_{S} the same steps give $\{(\alpha - \sqrt{e}\,\gamma + \beta)\,\tau > \beta\}$, which is linear. On each interval the integral of $p - eq$ is a combination of values of the normal distribution function, so I_{A} and I_{S} are explicit functions of z_1. At $z_1 = 2.5$,

$$I_{\mathrm{A}} = 0.0635779652453, \qquad I_{\mathrm{S}} = 0.0641560879254,$$

and the difference $5.78 \cdot 10^{-4}$ is positive. The numerical grid detects a transition near $z_1 = 2.005$. There is no finite upper cutoff: the analytic argument in Section 2.5 proves $I_{\mathrm{S}} > I_{\mathrm{A}}$ at every

$z_1 \geq 3$. Thus a finite grid endpoint must not be interpreted as the boundary of the improvement set.

Theorem 3.1. *At $T = 2$, $\sigma = 1$, $\varepsilon = \frac{1}{2}$ and every batch size, the method $\mathcal{A}_S$ with $S = [2.1, 2.6]$ satisfies*

$$H_{e^\varepsilon}(M(x)\|M(x')) \geq H_{e^\varepsilon}(P_{\mathcal{S}}\|Q_{\mathcal{S}}) + 1.04 \cdot 10^{-4}.$$

Proof. By Lemma 2.2 it is enough to show $\int_{2.1}^{2.6}(I_{\mathrm{S}} - I_{\mathrm{A}})\,dz_1 \geq 1.04 \cdot 10^{-4}$. We evaluate the closed forms in ball arithmetic (Arb [3] through python-flint, 200 bits) on 5000 balls covering $[2.1, 2.6]$. For each ball the roots of the quadratic, the interval endpoints and the normal distribution values are enclosed, and the resulting enclosure of $I_{\mathrm{S}} - I_{\mathrm{A}}$ has lower end at least $2.0828 \cdot 10^{-4}$. Multiplying by the length 0.5 gives the bound. As controls, the same code refuses the interval $[1.5, 1.9]$, where S loses (at $z_1 = 1.7$ the difference is about $-9.96 \cdot 10^{-3}$), and returns no verdict on balls near $z_1 = 2$, where $\alpha - \sqrt{e}\,\gamma$ vanishes. The coefficient $\alpha - \sqrt{e}\,\gamma + \beta$ is instead equal to $\beta > 0$ at that point. □

The theorem proves Theorem 1.2 at $T = 2$. Taking at each z_1 the better of A and S gives 0.223234, against 0.221269 for the pair; allowing any of the 81 answers $(\psi_2(u), \psi_2(w)) \in \{-1, -\frac{3}{4}, \ldots, 1\}^2$ at step 2 moves this only in the fifth digit (0.22324 on the finer of two grids); on a 9^4 grid of non-adaptive pairs of queries $(\psi_1(u), \psi_1(w), \psi_2(u), \psi_2(w)) \in \{-1, -\frac{3}{4}, \ldots, 1\}^4$ the largest divergence is the pair's, 0.221273 on that grid.

4. Every number of steps

Proposition 4.1. *Let $B = [-0.3, 0.1] \times [2.2, 2.3]$. At $\sigma = 1$, $\varepsilon = \frac{1}{2}$ and $T = 3$, $(I_{\mathrm{S}} - I_{\mathrm{A}})(\alpha(y), \beta(y), \gamma(y)) > 0$ for every $y \in B$, and*

$$\int_B (I_{\mathrm{S}} - I_{\mathrm{A}})(\alpha, \beta, \gamma)\,dy \geq 2.25 \cdot 10^{-5}.$$

Proof. As at $T = 2$, the set where the integrand of $I_{a,c}$ is positive is read off a quadratic in $\tau = e^s$, and $I_{a,c}$ is a sum of differences of normal distribution values. The closed form is evaluated in Arb at 128 bits on 21,539 boxes covering B, obtained by splitting a box in four while its enclosure contains zero. Every box gives a positive lower end, the smallest $9.91 \cdot 10^{-8}$, and summing lower end times area gives the bound. At $y = (-0.1, 2.25)$ the difference is $1.535260930 \cdot 10^{-3}$, which an independent quadrature reproduces to twelve digits. A further replay on 22 October 2026, starting from a 200×50 grid and using depth limit 14, produced 33,367 certified leaf boxes and a summed lower endpoint greater than $2.8408 \cdot 10^{-5}$; this independently verifies the stated bound. The original partition count is not needed for that replay. □

Theorem 4.2. *At $\sigma = 1$ and $\varepsilon = \frac{1}{2}$, for every $T \geq 3$ and every batch size there is a Borel set $S_T \subseteq \mathbb{R}^{T-1}$ of positive measure such that*

$$H_{e^\varepsilon}(M(x)\|M(x')) > H_{e^\varepsilon}(P_{\mathcal{S}}\|Q_{\mathcal{S}}) \qquad \text{for } \mathcal{A}_{S_T}.$$

Proof. Let $F_m \colon \mathbb{R}^m \to \mathbb{R}^2$ send y to (A, G) of (3). For $m = 2$ its Jacobian determinant in the variables x_1, x_2 is $2e^{-1}(x_1 - x_2)$, and on B we have $x_2/x_1 = e^{y_2 - y_1} \geq e^{2.1}$, so F_2 is a local diffeomorphism on the interior of B and $U = F_2(\operatorname{int} B)$ is a nonempty open set. By Proposition 4.1 and (4), $\Delta > 0$ on U. On B the ratio $eA/G^2 = (1 + r^2)/(1 + r)^2$ with $r = x_2/x_1 \in [e^{2.1}, e^{2.6}]$ satisfies $1/2 < eA/G^2 < 1$: the first strict inequality follows from $(r - 1)^2 > 0$, and the second from $2r > 0$. Thus $U \subseteq \{G^2/2 < eA < G^2\} \subseteq \{G^2/m < eA < G^2\}$ for every $m \geq 2$. By the remark after (5), $S_T = F_m^{-1}(U)$ with $m = T - 1$ is nonempty; it is open since F_m is continuous, so it has positive measure. On S_T the integrand of Lemma 2.2 is $\beta\Delta > 0$, and the integral is positive. □

Theorems 3.1 and 4.2 together give Theorem 1.2. The proof of Theorem 4.2 gives no explicit constant for $T \geq 4$; Section 6 lists the numerical size of the gain.

5. Kullback–Leibler and Rényi divergences

For $\alpha \in (0,1) \cup (1,\infty)$ write $D_\alpha(P\|Q) = \frac{1}{\alpha-1} \log \int p^\alpha q^{1-\alpha}$ and $D_{\mathrm{KL}}(P\|Q) = \int p \log(p/q)$. At $T = 2$ the step-2 answer at z_1 changes only the slice $p(z_1,\cdot), q(z_1,\cdot)$, so as in Lemma 2.2

$$\int p^\alpha q^{1-\alpha} = \int J_\alpha(z_1)\, dz_1, \qquad D_{\mathrm{KL}} = \int K(z_1)\, dz_1,$$

with J_α and K the inner integrals over z_2 for the answer chosen at z_1. For $\alpha > 1$ a larger $\int J_\alpha$ is a larger D_α, for $\alpha < 1$ a smaller one. So a switch to S on an interval where $J_\alpha^{\mathrm{S}} - J_\alpha^{\mathrm{A}}$ has the right sign, or where $K^{\mathrm{S}} > K^{\mathrm{A}}$, beats the pair, and an Arb enclosure of the difference on balls covering the interval proves it. The inner integrals are computed with Arb's rigorous integration on a bounded interval, about thirty noise scales on each side, and the tails are bounded by explicit Gaussian tails; for non-integer α the power is taken in Arb's analytic mode, so a ball that meets the branch cut gives no verdict and is subdivided. Table 1 lists the certified excesses at $\sigma = 1$ for the orientation $(M(x)\|M(x'))$ against $(P_{\mathcal{S}}\|Q_{\mathcal{S}})$, which we call $P\|Q$, and for $(M(x')\|M(x))$ against $(Q_{\mathcal{S}}\|P_{\mathcal{S}})$, called $Q\|P$.

divergence	certified excess, $P\|Q$	certified excess, $Q\|P$
KL	$9.97 \cdot 10^{-6}$	$6.72 \cdot 10^{-6}$
order 1/4	$9.42 \cdot 10^{-6}$	$9.57 \cdot 10^{-6}$
order 1/2	$1.52 \cdot 10^{-5}$	$1.52 \cdot 10^{-5}$
order 3/4	$9.57 \cdot 10^{-6}$	$9.42 \cdot 10^{-6}$
order 3/2	$9.21 \cdot 10^{-6}$	$2.30 \cdot 10^{-5}$
order 2	$3.22 \cdot 10^{-5}$	$1.58 \cdot 10^{-3}$
order 5/2	$7.14 \cdot 10^{-5}$	$1.80 \cdot 10^{-3}$
order 3	$2.21 \cdot 10^{-4}$	0.150
order 4	$1.40 \cdot 10^{-3}$	
order 5	$3.79 \cdot 10^{-3}$	

Table 1. Certified lower bounds at $T = 2$, $\sigma = 1$ for the excess of the adaptive switch over the pair: in D_{KL}, and in $\pm \int p^\alpha q^{1-\alpha}$ for Rényi order α (the sign that increases D_α). The switch interval is chosen per row and is recorded with each certificate.

Three remarks on the table. First, for $\alpha < 1$ the integral $\int p^\alpha q^{1-\alpha}$ is symmetric under $(P, Q, \alpha) \mapsto (Q, P, 1-\alpha)$, and this holds slice by slice, so the $P\|Q$ entry at order $\frac{1}{4}$ is the $Q\|P$ entry at order $\frac{3}{4}$. If one compares instead with the larger of the pair's two orientations, the switch wins at order $\frac{1}{2}$ in both orientations, at order $\frac{1}{4}$ only in $Q\|P$ and at order $\frac{3}{4}$ only in $P\|Q$. Second, the large $Q\|P$ entries at high order are not all adaptivity: at orders 4 and 5 the fixed query S at step 2, which uses no information from step 1, already beats the pair in $Q\|P$. The order-3 entry is adaptive, since there the pair beats the fixed query S; in $P\|Q$ the pair beats the fixed S at every order and the whole gain is adaptive. Third, van Dijk and Ertan [4, Appendix J] compute $\chi^2(P_{\mathcal{S}}\|Q_{\mathcal{S}}) = 1.5895$ at $T = 2$, $\sigma = 1$, and our grid gives 1.589482 for the pair and 1.591212 for the best switch. They also use the identity $\chi^2(P\|Q) + 1 = 2\int_0^\infty H_\lambda(P\|Q)\, d\lambda$; through it the order-2 certificate is a second proof that the pair fails to dominate in hockey-stick divergence for some ε.

The same certificates were run at $\sigma = \frac{1}{2}$ and $\sigma = 2$ for KL and the orders $\frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 2, 3$ in both orientations, and every one is positive. At $\sigma = 2$ the bounds are of order 10^{-5} to 10^{-4}. At $\sigma = \frac{1}{2}$ the gains in $P\|Q$ are small (the grid gives $2.8 \cdot 10^{-8}$ in KL) and the certified bounds smaller still, $1.9 \cdot 10^{-14}$ in KL, because the enclosure loses roughly in proportion to the ball width and only very short intervals certify.

6. Numerical values

All values in this section are numerical. At $T = 3$, $\sigma = 1$, $\varepsilon = \frac{1}{2}$ the optimum over methods that may switch at both step 2 and step 3, with answers in $\{-1, 1\}^2$, computed as a nested pointwise maximum on a grid, is 0.210464, against 0.207611 for the pair (Monte Carlo 0.207652 ± 0.000143). The reverse orientation of the pair gives 0.18142.

For larger T we computed the gain of the last-step switch alone, $\frac{1}{T}\mathbb{E}_{y\sim\mathcal{N}(0,I_m)}\Delta(A(y), G(y))_+$ by (4), by Monte Carlo with $2 \cdot 10^6$ prefixes (Table 2). Switching at earlier steps as well can only increase the divergence.

T	$H_{e^\varepsilon}(P_{\mathcal{S}}\|Q_{\mathcal{S}})$	gain of the last-step switch
2	0.22127 (quadrature)	$1.966 \cdot 10^{-3}$ (quadrature)
3	0.20761	$1.910 \cdot 10^{-3}$
4	0.1937	$1.646 \cdot 10^{-3}$
8	0.1631	$8.84 \cdot 10^{-4}$
16	0.1290	$3.71 \cdot 10^{-4}$
32	0.0959	$1.27 \cdot 10^{-4}$

Table 2. $\sigma = 1$, $\varepsilon = \frac{1}{2}$. Monte Carlo standard errors are below 1% of the gain; for $T \geq 4$ the pair's value carries an error between 0.0008 and 0.0022.

At $T = 2$ the gain is positive in every cell of $\sigma \in \{\frac{1}{2}, 1, 2\} \times \varepsilon \in \{0.1, 1, 2, 3\}$, the smallest being $5.2 \cdot 10^{-13}$ at $\sigma = 2$, $\varepsilon = 3$, where $H_{e^\varepsilon}(P_{\mathcal{S}}\|Q_{\mathcal{S}})$ itself is $2.3 \cdot 10^{-10}$. At $\varepsilon = 0.1$ and $\sigma = \frac{1}{2}$ the last-step gain stays near 10^{-4} up to $T = 16$ and then falls roughly like $1/T$ ($5.3 \cdot 10^{-6}$ at $T = 512$). A heuristic reason: with $\sigma = \frac{1}{2}$ the variables $x_t = e^{(y_t - 1/2)/\sigma^2}$ have $\mathbb{E}x_t^2 = e^{1/\sigma^2} \approx 55$, so while m is well below that one term dominates G, the prefix lands in the same part of the region (5) whatever T is, and the factor $1/T$ in β is offset.

7. Computations

The repository [5] contains the scripts and their outputs. `cert_t2_arb.py` and `cert_t3_arb.py` carry out the certificates of Theorem 3.1 and Proposition 4.1; `cert_chi2_arb.py`, `cert_kl_arb_sigma.py`, `cert_renyi_arb_sigma.py`, `cert_renyi_real_arb_sigma.py` and `cert_renyi_sub1_arb_sigma.py` those of Section 5, each subdividing an interval into balls, with every sub-ball built from exact endpoints so that the balls cover the interval. `exact_closed.py` evaluates the closed forms of Section 3 to 30 digits and `t2b1_adaptive.py`, `nonadaptive.py`, `t3_dp.py`, `tgen_mc.py` and `renyi_alpha_grid.py` produce the numerical values. `independent_check.py` recomputes the $T = 2$ values 0.221269, 0.223234 and 0.205107 and the point values at $z_1 = 2.5$ and $y = (-0.1, 2.25)$ with plain quadrature, sharing no code with the others. The scripts need Python 3 with numpy, scipy, mpmath and python-flint. The independent 22 October replay used Python 3.12.14, python-flint 0.9.0, NumPy 2.3.5, SciPy 1.17.0 and mpmath 1.3.0. Its commands and interval-coverage checks

are in `reviews/2026-10-22/`; in particular, the main certificates use arguments `-1.6 -1.1 5000` for $T = 2$ and `-0.1 0.3 2.2 2.3 200 50 14` for $T = 3$.

Acknowledgements and funding

The research presented in this paper was supported by the European Research Council (ERC) under the European Union's Horizon Europe research and innovation programme (grant agreement No. 101041711), by the Simons Foundation as part of the Collaboration on the Mathematical and Scientific Foundations of Deep Learning, by Heights Labs, by Convex Nexus Capital, by the Israel Science Foundation (grant number 2258/19), and by the Israel Science Foundation (ISF Grant 4101/25).

Declaration of competing interest

The author holds equity in Heights Labs and in Convex Nexus Capital and has a management role at Convex Nexus Capital. These entities are included in the funding declaration above.

Declaration of generative AI and AI-assisted technologies

The paper comes out of Math Brain, an ongoing project on the use of AI systems for research mathematics. On 19 October 2026 a Claude agent (Anthropic), working in Claude Code, found the last-step switch, the reduction of Section 2, the certificates of Theorem 3.1 and Proposition 4.1 and those of Section 5; two further Claude agents checked each certificate with code of their own. On 20 October 2026 Claude Opus 5.5, working in Claude Code, recomputed the main values independently and wrote the paper. On 22 October 2026 an independent Codex session checked the main hockey-stick theorem against the source conjecture, reran the two principal certificates, checked their interval coverage and point values, supplied the analytic proof in Section 2.5, and corrected several incidental statements. That review did not cover the KL and Rényi certificates or the full numerical optimization claims, and it is not external human peer review. The author directed the work and is responsible for the content of the paper and for decisions concerning its dissemination.

Data availability

No data were used. The code is available at
`https://github.com/yspennstate/shuffled-ablq-dominating-pair`.

References

[1] L. Chua, B. Ghazi, P. Kamath, R. Kumar, P. Manurangsi, A. Sinha and C. Zhang, *How private are DP-SGD implementations?*, Proceedings of the 41st International Conference on Machine Learning (ICML 2024); arXiv:2403.17673.

[2] V. Feldman and M. Shenfeld, *Privacy amplification by random allocation*, arXiv:2502.08202 (2025).

[3] F. Johansson, *Arb: efficient arbitrary-precision midpoint-radius interval arithmetic*, IEEE Trans. Comput. **66** (2017), 1281–1292.

[4] M. van Dijk and M. B. Ertan, *Trade-off functions for DP-SGD with subsampling based on random allocation: tight upper and lower bounds*, arXiv:2605.06259v3 (2026).

[5] Y. Shmalo, *Code for this paper*, `https://github.com/yspennstate/shuffled-ablq-dominating-pair`.

Einstein Institute of Mathematics, The Hebrew University of Jerusalem, Jerusalem, Israel
Email address: `yitzchak.shmalo@gmail.com`