DEAD BRANCHES IN THE TRACE-REINFORCED ANT PROCESS

YITZCHAK SHMALO

Abstract. In the trace-reinforced ant process of Kious, Mailler and Schapira, ants walk one after another from a nest N to a food source F on a finite graph, and each ant adds one unit of weight to every edge it has crossed. It was conjectured that the normalised weights converge to a deterministic limit which is positive on every edge once N and F are at distance at least two. We show that the positivity fails. If a part of the graph hangs by a single edge from a neighbour of F, on the side away from N, then the normalised weights of all its edges tend to zero almost surely; this happens in 2-connected graphs with N and F at distance two. When some neighbour of F is hidden from N by the other neighbours of F, the mean-field vector field has no equilibrium with all coordinates positive, so on such graphs, some of them 3-connected, the normalised weights cannot converge to a positive limit. The equilibria are described through the voltage of the unit current from N to F and are computed on any number of parallel paths. A census of the 22,828 cases on connected graphs with at most seven vertices supports a corrected conjecture, in which an explicit combinatorial rule gives the edges with limit zero.

1. Introduction

Let $G = (V, E)$ be a finite connected simple graph with two distinct marked vertices, the nest N and the food F. The *trace-reinforced ant process* on (G, N, F) is the sequence of weight vectors $W(n) = (W_e(n))_{e\in E}$ defined as follows. Set $W_e(0) = 1$ for all e. Given $W(n)$, the $(n+1)$-st ant performs a random walk started at N which jumps from a vertex x to a neighbour z with probability $W_{xz}(n)/W_x(n)$, where $W_x(n) = \sum_{z\sim x} W_{xz}(n)$, and stops when it first hits F. Then

$$W_e(n+1) = W_e(n) + \mathbf{1}\{\text{the } (n+1)\text{-st ant crossed } e\}.$$

The process was introduced by Kious, Mailler and Schapira [13], as a variant of the models of [12], in which an ant reinforces only a part of its trace (its loop-erasure, or a shortest path inside it). These models are motivated by the trails with which ants mark the way to a food source, observed in the Argentine ant [5] and turned into an optimisation heuristic [6]. They belong to the reinforced random processes surveyed by Pemantle [11], a family that goes back to the edge-reinforced walk of Davis [4]. Mailler and Varin treat a model with two nests [14]. Reinforced walks which form a path are studied in [9, 7], and Pólya urns placed on the vertices of a graph and competing along its edges in [8]. Kious, Mailler and Schapira proved that $W(n)/n$ converges almost surely to an explicit deterministic limit on several families of graphs (tree-like graphs in which N and F are adjacent, the cone, two parallel paths, the lozenge), and stated the following.

Conjecture 1 ([13], Conjecture 1.1)**.** *There exists a deterministic family $(\chi_e)_{e\in E}$ such that $W_e(n)/n \to \chi_e$ almost surely for all e. Furthermore, if in addition the distance between N and F is at least 2, then $\chi_e > 0$ for all $e \in E$.*

Date: November 2026.

The first sentence of the conjecture is restated in the list of open problems [1, Section 2]. For the loop-erased rule of [12] and for the forward-loop-erased rule of Armfield and Mailler [2], in which an ant reinforces the loop-erasure of its trajectory, the question is instead whether the edges which lie on no geodesic have normalised weight tending to zero. This is proved in [12] for series-parallel graphs and in [2] when N and F are at distance two, and it fails on a graph in which N and F are at distance three [18].

An edge with an endpoint outside the connected component of N in $G-F$ is never crossed, so the second sentence is to be read for graphs in which $G-F$ is connected; we call such triples (G,N,F) *reduced*. The second sentence fails for reduced graphs.

Theorem 2 (Dead branches)**.** *Let u be a neighbour of F, and let $H \subset V \setminus \{N,F,u\}$ be a set of vertices such that exactly one edge $e=\{u,y\}$ joins H to $V \setminus (H \cup \{F\})$. Then, almost surely,*

$$\frac{W_f(n)}{n} \longrightarrow 0 \qquad \text{for every edge } f \text{ with an endpoint in } H.$$

In other words, H is the far side of a bridge of $G-F$ attached at a neighbour of F; the vertices of H may have any number of edges to F. The vertex u may be N itself. When $G-F$ is a tree and $u=N$ is adjacent to F the theorem covers every edge other than $\{N,F\}$, which is the second half of Theorem 1.3 of [13]. When N is adjacent to F, Proposition 2.8 of [13] shows that $W_{NF}(n)/n \to 1$ and that the normalised weight of every other edge at F tends to zero; Theorem 2 needs no such assumption.

Example 3. Take a triangle x,y,F and attach N to x by an edge. The distance from N to F is 2, and Theorem 2 with $u=x$, $H=\{y\}$ gives $W_{xy}(n)/n \to 0$ and $W_{yF}(n)/n \to 0$. This case also follows from [13, Theorem 1.3] applied with nest x: every ant goes through x, and its excursions from x to the pendant vertex N do not change which of the other edges it crosses. The same reduction applies whenever a cut vertex separates N from F, so the examples of interest are those without one.

Example 4. Take the 4-cycle N,a,F,b and add a vertex y joined to a and to F. The graph is 2-connected, N and F are at distance 2, and every edge lies on a simple path from N to F. Theorem 2 with $u=a$, $H=\{y\}$ gives $W_{ay}(n)/n \to 0$ and $W_{yF}(n)/n \to 0$: the detour a,y,F dies next to the edge $\{a,F\}$.

The second result concerns the mean-field vector field. For a weight vector $w \in [0,\infty)^E$ whose support connects N to F, let $p_e(w)$ be the probability that the w-weighted random walk started at N crosses e before hitting F. Then $\mathbb{E}[W_e(n+1)-W_e(n) \mid \mathcal{F}_n] = p_e(W(n))$, the map p is homogeneous of degree 0, and the process is a stochastic approximation of the flow $\dot{w} = p(w) - w$ in the sense of [3, 10]. An *equilibrium* is a solution of $p(w)=w$.

Theorem 5 (Shadowed neighbours of F)**.** *Let (G,N,F) be reduced with N not adjacent to F, let $\Gamma(F)$ be the set of neighbours of F, and let I_0 be the connected component of N in $G-F-\Gamma(F)$. If some $x \in \Gamma(F)$ has no neighbour in I_0, then p has no equilibrium with all coordinates positive. Consequently, on such a graph the event that $W(n)/n$ converges to a limit with all coordinates positive has probability zero.*

So on these graphs the two sentences of Conjecture 1 cannot both hold: either $W(n)/n$ does not converge, or its limit has a zero coordinate. The hypothesis does not involve bridges or cut vertices. For instance, take the complete bipartite graph with parts $\{a,b,c\}$ and $\{N,F,x\}$ and add the edge $\{x,F\}$. This graph is 3-connected, N and F are at distance 2, $I_0=\{N\}$, and x is a neighbour of F with no neighbour in I_0. A smaller example is the 4-cycle N,a,F,b

with a fifth vertex c joined to a, b and F. In the mean-field flow on this graph the edge $\{c, F\}$ dies while $\{a, c\}$ and $\{b, c\}$ keep the positive weight 0.3209 (Section 4).

Section 2 contains the proof of Theorem 2. Section 3 describes the equilibria through electrical voltages and proves Theorem 5. Section 4 states a corrected form of the second sentence of Conjecture 1 and reports a census of small graphs. Section 5 computes an equilibrium on theta graphs with any number of paths.

2. Proof of Theorem 2

Write $a = \{u, F\}$, $X_n = W_e(n)$ and $A_n = W_a(n)$, and let $\mathcal{F}_n$ be the σ-field generated by the first n ants.

Step 0. *For every edge f with an endpoint in H and every n, $W_f(n) \le X_n$.* An ant starts at $N \notin H$ and stops at F, and e is the only edge from H to $V \setminus (H \cup \{F\})$. Hence an ant that crosses f has crossed e earlier. Both weights start at 1 and each increases by one when its edge is crossed.

Step 1. *One ant.* Fix n and write $w = W(n)$, $X = X_n$, $A = A_n$, and R for the total weight of the edges at u other than a and e. Let h be the probability that the walk from N hits u before F ($h = 1$ if $u = N$). The walk cannot enter H before it visits u. After the first visit to u, decompose the walk into its successive departures from u. A departure is of one of four kinds:

(i) the step $u \to F$, with probability A/w_u; the edge a is crossed and the walk ends;
(ii) the step $u \to y$, with probability X/w_u; the edge e is crossed; afterwards the walk returns to u with some probability ρ and is absorbed at F from inside H with probability $1 - \rho$;
(iii) a step along another edge, with probability R/w_u, after which the walk returns to u before hitting F; this has conditional probability ρ', say;
(iv) a step along another edge after which the walk hits F before returning to u.

An excursion of kind (iii) or (iv) does not meet H. Successive departures are independent. The edge e is crossed if and only if the first departure that is not of kind (iii) is of kind (ii), and the edge a is crossed if and only if the walk ends by a departure of kind (i). With $S = A + R(1 - \rho')$ this gives

$$p_e(w) = \frac{hX}{X + S}, \qquad p_a(w) = \frac{hA}{S + X(1 - \rho)}. \tag{1}$$

Therefore

$$\frac{p_a(w)}{A} - \frac{p_e(w)}{X} = \frac{hX\rho}{(S + X(1 - \rho))(S + X)} \;\ge\; \frac{\rho\, p_e(w)}{S + X}. \tag{2}$$

Now $S + X \le w_u \le \deg(u)\,(n + 1)$, since every weight is at most $n + 1$ at time n. Moreover ρ is at least the probability that the first step from y goes back to u, which is $X/w_y \ge 1/\deg(y)$ by Step 0, all edges at y having an endpoint in H. With $c = 1/(\deg(u)\deg(y))$ we obtain

$$\frac{p_a(W(n))}{A_n} - \frac{p_e(W(n))}{X_n} \;\ge\; \frac{c\, p_e(W(n))}{n + 1}. \tag{3}$$

Step 2. *The logarithm of the ratio.* Let $\Lambda_n = \log(X_n/A_n)$, and let D_n and D'_n be the indicators that the $(n + 1)$-st ant crosses e and a respectively, so that

$$\Lambda_{n+1} - \Lambda_n = D_n \log\Big(1 + \frac{1}{X_n}\Big) - D'_n \log\Big(1 + \frac{1}{A_n}\Big).$$

Using $t - t^2/2 \le \log(1+t) \le t$ and (3),

$$\mathbb{E}[\Lambda_{n+1} - \Lambda_n \mid \mathcal{F}_n] \;\le\; \frac{p_e(W(n))}{X_n} - \frac{p_a(W(n))}{A_n} + \frac{p_a(W(n))}{2A_n^2} \;\le\; -\frac{c\,p_e(W(n))}{n+1} + \frac{p_a(W(n))}{2A_n^2}.$$

Since X increases by one each time $D_n = 1$, we have $\sum_n D_n/X_n^2 \le \sum_{k\ge1} k^{-2}$, and likewise for A. Hence $\sum_n \mathbb{E}[(\Lambda_{n+1}-\Lambda_n)^2] \le 2\sum_k k^{-2} < \infty$, so the martingale $M_n = \sum_{j<n}\big(\Lambda_{j+1} - \Lambda_j - \mathbb{E}[\Lambda_{j+1} - \Lambda_j \mid \mathcal{F}_j]\big)$ is bounded in L^2 and converges almost surely. Also $\mathbb{E}\sum_n p_a(W(n))/A_n^2 = \mathbb{E}\sum_n D_n'/A_n^2 < \infty$, so $Q = \sum_n p_a(W(n))/(2A_n^2)$ is almost surely finite. Altogether, for all n,

$$\Lambda_n \;\le\; M_n + Q - c\sum_{j<n} \frac{p_e(W(j))}{j+1}. \tag{4}$$

Step 3. *Conclusion.* The martingale $\sum_{j<n}(D_j - p_e(W(j)))/(j+1)$ has square-summable increments and converges almost surely. Hence, almost surely,

$$\sum_j \frac{p_e(W(j))}{j+1} = \infty \quad\Longleftrightarrow\quad \sum_j \frac{D_j}{j+1} = \infty. \tag{5}$$

Summation by parts gives $\sum_{j<n} D_j/(j+1) = (X_n - 1)/n + \sum_{j=1}^{n-1}(X_j - 1)/(j(j+1))$. Suppose that $X_m \ge \delta m$ for infinitely many m, for some $\delta > 0$. For such m with $\delta m \ge 2$, monotonicity gives $X_j - 1 \ge \delta m/2$ for $m \le j \le 2m$, and $\sum_{j=m}^{2m} 1/(j(j+1)) \ge 1/(2m)$, so the block $[m, 2m]$ contributes at least $\delta/4$. Infinitely many disjoint blocks give $\sum_j D_j/(j+1) = \infty$. By (5) and (4), on the event $\{\limsup X_n/n > 0\}$ we have $\Lambda_n \to -\infty$ almost surely, that is $X_n/A_n \to 0$. But $A_n \le n+1$, so $X_n/n \to 0$ on that event, which is a contradiction unless the event is null. Thus $X_n/n \to 0$ almost surely, and Step 0 extends this to every edge with an endpoint in H. □

Remark 6. The bridge is essential. If the part hanging from u is attached by two edges, say a triangle u, y_1, y_2, an ant can enter by one edge and leave by the other, and the same computation gives $p_{uy_1}(w)/w_{uy_1} > p_a(w)/w_a$ for small weights on the triangle. In the mean-field flow such a triangle keeps positive weight.

3. Equilibria and voltages

Let $w \in [0,\infty)^E$ have a support that connects N to F, and let $g(x)$ be the expected number of visits to x by the w-walk from N before it hits F. For a vertex $x \ne F$ with $w_x > 0$ put $v(x) = g(x)/w_x$, and put $v(F) = 0$. Then v is the voltage of the unit current from N to F in the network w:

$$\sum_{z\sim x} w_{xz}\,(v(x) - v(z)) = \mathbf{1}\{x = N\} \qquad (x \ne F,\ w_x > 0). \tag{6}$$

Lemma 7. *The expected number of crossings of $e = \{x, z\}$ by the walk is $w_e\,(v(x) + v(z))$. In particular $p_e(w) \le w_e(v(x) + v(z))$, and $p_{xF}(w) = w_{xF}\,v(x)$ for every neighbour x of F.*

Proof. The expected number of steps from x to z is $g(x)\,w_e/w_x$. An edge at F is crossed at most once. □

Proposition 8. *Let w be an equilibrium and let V_1 be the set of neighbours x of F with $w_{xF} > 0$. Then*

(a) *$v(x) = 1$ for every $x \in V_1$;*
(b) *$v(x) \ge 1$ for every vertex $x \ne F$ visited by the walk;*

(c) *if $N \notin V_1$, every $x \in V_1$ has a neighbour z with $w_{xz} > 0$ and $v(z) > 1$, and every vertex with $v > 1$ is joined to N by a path of the support along which $v > 1$;*
(d) *if $N \in V_1$, then $w_{NF} = 1$, $v \equiv 1$ on the visited vertices, and $w_{xF} = 0$ for every other neighbour x of F.*

Proof. (a) By Lemma 7, $w_{xF} = p_{xF}(w) = w_{xF}v(x)$. (b) By (6), at a visited vertex outside $V_1 \cup \{F\}$ the value $v(x)$ is at least the weighted mean of v over its neighbours in the support, none of which is F. If the minimum of v over the visited vertices other than F were attained outside V_1, it would be attained at all support neighbours, and, the support component of N being connected and containing a vertex of V_1, also on V_1, where $v = 1$. (c) For $x \in V_1$, $x \neq N$, (6) and (a) give $\sum_{z \neq F} w_{xz}(v(z) - 1) = w_{xF} > 0$, so some neighbour z in the support has $v(z) > 1$. Now let $v(z) = t > 1$ and let C be the set of vertices joined to z by a path of the support along which $v \geq t$. Suppose $N \notin C$. By (a), C does not meet V_1, so by (6) the value of v at a vertex of C is the weighted mean of v over its neighbours in the support. A neighbour outside C has $v < t$. Hence a vertex where v is maximal on C has all its support neighbours in C, and v is maximal at each of them. Repeating, C is a union of connected components of the support. But z is visited, so its component contains N, a contradiction. (d) If $N \in V_1$ then $v(N) = 1$, and $v(N)$ is the maximum of v, so $v \equiv 1$ on the visited vertices by (b). Then (6) at N reads $w_{NF} = 1$, and at another $x \in V_1$ it would read $w_{xF} = 0$. □

Proof of Theorem 5. Let w be an equilibrium with all coordinates positive. Then $V_1 = \Gamma(F)$, the support is E, and by Proposition 8(a) the set $\{v > 1\}$ does not meet $\Gamma(F)$. By (c), each $x \in \Gamma(F)$ has a neighbour z with $v(z) > 1$, and z is joined to N by a path inside $\{v > 1\} \subset V \setminus (\Gamma(F) \cup \{F\})$; thus $z \in I_0$.

For the second statement, $W_e(n) - \sum_{j<n} p_e(W(j))$ is a martingale with increments bounded by 1, so $n^{-1}\big(W_e(n) - \sum_{j<n} p_e(W(j))\big) \to 0$ almost surely. On the event that $W(n)/n \to \chi$ with all $\chi_e > 0$, homogeneity and the continuity of p on $(0,\infty)^E$ give $p_e(W(j)) \to p_e(\chi)$, hence $p(\chi) = \chi$, which has just been excluded. □

Lemma 7 also explains which zero edges can be stable. Let w be an equilibrium and $e = \{x, z\}$ an edge with $w_e = 0$. If a small weight ε is put on e alone, then $p_e = \varepsilon\,(v(x)+v(z))+o(\varepsilon)$ when both endpoints are visited, and $p_e = \varepsilon\, v(x) + o(\varepsilon)$ when only x is. By Proposition 8(b) the first rate is at least 2 unless $z = F$, so such an edge is repelled from zero. An edge to an unvisited vertex, or to F, is not repelled only when $v(x) = 1$, and then the rate equals 1 exactly: linear analysis is silent, the decay is of order $1/t$, and whether the edge dies is decided at second order, as in Theorem 2.

4. A corrected conjecture and a census of small graphs

Let (G, N, F) be reduced. If N is adjacent to F put $I_0 = \emptyset$ and $\mathcal{A} = \{N\}$; otherwise let I_0 be the component of N in $G - F - \Gamma(F)$ and let $\mathcal{A}$ be the set of neighbours of F that have a neighbour in I_0. Call $V \setminus (I_0 \cup \{F\})$ the *shadow*. Let $\mathcal{D}(G, N, F)$ be the set of edges of the following two kinds:

(D1) $\{x, F\}$ with $x \in \Gamma(F) \setminus \mathcal{A}$;
(D2) for every shadow vertex u and every bridge $e = \{u, y\}$ of $G - F$ whose far side H (the component of y in $G - F - e$) does not contain N: the edge e and every edge with an endpoint in H.

Conjecture 9. *For every reduced (G, N, F), $W(n)/n$ converges almost surely to a deterministic limit χ, and $\chi_e = 0$ if and only if $e \in \mathcal{D}(G, N, F)$.*

Theorem 2 proves $\chi_e = 0$ for the edges of kind (D2) with $u \in \Gamma(F)$. When N is adjacent to F, the edges of kind (D1) are covered by [13, Proposition 2.8], and Proposition 8(d) is the corresponding statement for equilibria. When N is not adjacent to F, Theorem 5 shows that $\mathcal{D} \neq \emptyset$ is forced as soon as (D1) is non-empty.

We tested the mean-field form of Conjecture 9 on every connected graph with 3 to 7 vertices and every ordered pair (N, F) up to automorphism, 22,828 cases in all, of which 20,092 are reduced; in the others the never-crossed edges were removed first. The list of graphs is that of the atlas of Read and Wilson [17], as distributed with NetworkX [15], and the computations use NumPy [16]. For each case the flow $\dot{w} = p(w) - w$ was integrated from six initial conditions (uniform, log-uniform over six e-folds, and concentrated on a random path from N to F); coordinates that decay were set to zero and the remaining ones were refined by Newton's method, with p evaluated by one linear solve per edge.

	cases	one limit	zero set $= \mathcal{D}$	reduced	reduced, $\mathcal{D} \neq \emptyset$
N adjacent to F	12,004	12,003	12,003	10,286	10,064
distance at least 2	10,824	10,824	10,824	9,806	4,118

The six runs reached the same point in every case but one. In the remaining case, a graph with seven vertices in which N is adjacent to F, an edge of $\mathcal{D}$ has a weight that decays like $1/t$ along the flow, and in one run it had not been set to zero by the end of the integration; with ten times the integration time the six runs agree and the zero set is $\mathcal{D}$. At the limits the largest real part of an eigenvalue of $Dp - I$ restricted to the support is -0.015, and the smallest positive coordinate is 0.0019. Among the 9,806 reduced cases at distance at least 2, Theorem 2 applies to 2,504 and Theorem 5 to 3,294; 1,643 of the 4,118 cases with a dead edge are 2-connected graphs, and 247 are 3-connected.

The process itself was simulated with 10^7 ants on four graphs. On the graph of Example 4 the values of $W_{ay}(n)/n$ at $n = 10^3, 10^4, \ldots, 10^7$ were 0.060, 0.059, 0.049, 0.044, 0.040, and those of $W_{yF}(n)/n$ were 0.020, 0.016, 0.010, 0.007, 0.005, while the four edges of the cycle were within 0.02 of their limits $2^{-1/2}$ and $1/2$ at $n = 10^7$. On the two graphs described after Theorem 5 the weight of the edge at F that dies in the flow went from 0.123 to 0.067 and from 0.098 to 0.044 over the same range. The decay is slow. In Step 2 of the proof of Theorem 2 the logarithm of X_n/A_n decreases in conditional mean by an amount of order p_e/n per ant, and p_e is itself of order X_n/n; this points to a rate of order $1/\log n$, which is compatible with these values.

5. Theta graphs

Let $\Theta(L_1, \ldots, L_k)$ consist of $k \geq 2$ internally disjoint paths from N to F, the i-th of which has $L_i \geq 2$ edges. For $k = 2$ the almost sure limit of $W(n)/n$ was found in [13, Theorem 1.5]: the j-th edge from N of each path has limit α^j, with a ratio α depending on the path. The same form gives an equilibrium for every k.

Proposition 10. *There is a unique $(\alpha_1, \ldots, \alpha_k) \in (0,1)^k$ such that*

$$\sum_{i=1}^{k} \alpha_i^{L_i} = 1 \qquad \text{and} \qquad \frac{1-\alpha_i}{1-\alpha_i^{L_i}} \text{ does not depend on } i, \tag{7}$$

and the vector that gives weight α_i^j to the j-th edge from N of the i-th path, $1 \leq j \leq L_i$, is an equilibrium on $\Theta(L_1, \ldots, L_k)$. If $L_1 = \cdots = L_k = L$ then $\alpha_i = k^{-1/L}$ for all i.

Proof. Let the j-th edge of path i have weight α_i^j, with $\alpha_i \in (0,1)$ to be chosen. This edge is crossed exactly when the walk reaches the vertex at depth j of path i. An excursion from N enters path i with probability proportional to α_i, and then reaches depth j before returning to N with probability $\left(\sum_{m=0}^{j-1} \alpha_i^{-m}\right)^{-1}$, by the gambler's ruin formula for the path with resistances $\alpha_i^{-1}, \dots, \alpha_i^{-j}$. So the excursions that reach depth j of path i occur at rate

$$a_{i,j} = \frac{\alpha_i}{\sum_{m=0}^{j-1} \alpha_i^{-m}} = \frac{1-\alpha_i}{\alpha_i^{-j}-1},$$

and those that reach F along path i at rate $c_i = a_{i,L_i}$. The walk ends at the first excursion that reaches F, so with $C = \sum_i c_i$,

$$p_{i,j} = \frac{a_{i,j}}{a_{i,j} + C - c_i}.$$

Since $a_{i,j}(1-\alpha_i^j) = (1-\alpha_i)\alpha_i^j$, the equation $p_{i,j} = \alpha_i^j$ is equivalent to $1-\alpha_i = C - c_i$, for every j. As $c_i = (1-\alpha_i)\alpha_i^{L_i}/(1-\alpha_i^{L_i})$, this reads $(1-\alpha_i)/(1-\alpha_i^{L_i}) = C$, and then $c_i = C\alpha_i^{L_i}$, so that $C = \sum_i c_i$ becomes $\sum_i \alpha_i^{L_i} = 1$. This is (7).

The function $\alpha \mapsto (1-\alpha)/(1-\alpha^L) = (1+\alpha+\cdots+\alpha^{L-1})^{-1}$ decreases from 1 to $1/L$ on $(0,1)$. For $C \in (\max_i 1/L_i, 1)$ let $\alpha_i(C)$ be the solution of $(1-\alpha)/(1-\alpha^{L_i}) = C$. Then $\sum_i \alpha_i(C)^{L_i}$ is continuous and decreasing in C, tends to 0 as $C \to 1$, and exceeds 1 as $C \to \max_i 1/L_i$, because one term tends to 1 and the others stay positive. So (7) has exactly one solution. In the case of equal lengths the second condition gives equal α_i and the first gives $\alpha^L = 1/k$. □

For $k = 2$, (7) is the system of [13, Theorem 1.5]. In the flow, thirty random initial conditions converged to the equilibrium of Proposition 10 on each of 39 theta graphs with $k \le 8$ and $L_i \le 8$.

Acknowledgements and funding

This work was done by multiple AI agents, Claude (Anthropic) and Codex (OpenAI), connected to Math Brain [19] and working together inside the Brain framework [21], a shared, persistent memory in which the agents record claims, proofs, failed attempts and methods, retrieve one another's results, coordinate who works on what, and audit each other's work. For more on the details behind the Brain and Math Brain, see the Math Brain paper [20], the Math Brain repository [19] and the Brain framework repository [21].

The research presented in this paper was supported by the European Research Council (ERC) under the European Union's Horizon Europe research and innovation programme (grant agreement No. 101041711), by the Simons Foundation as part of the Collaboration on the Mathematical and Scientific Foundations of Deep Learning, by Heights Labs, by Convex Nexus Capital, by the Israel Science Foundation (grant number 2258/19), and by the Israel Science Foundation (ISF Grant 4101/25). Computational resources were provided in part by Google Cloud through its education credits program (credit 530550103).

Declaration of competing interest

The author holds equity in Heights Labs and in Convex Nexus Capital and has a management role at Convex Nexus Capital. These entities are included in the funding declaration above.

Declaration of generative AI and AI-assisted technologies

The paper comes out of Math Brain, an ongoing project on the use of AI systems for research mathematics, and its results were developed with the Brain [21] and Math Brain [19]. In November 2026 a Claude Fable agent (Anthropic), connected to the Brain and working from the project's list of open problems in probability, found the statements of this paper and their proofs, wrote the programs, ran the census and the simulations, and prepared the manuscript. A second Claude agent independently checked part of the computations. These checks are not peer review. The author directed the work and is responsible for the content of the paper and for decisions concerning its dissemination.

Code and data availability

Programs, census files and outputs:
`https://github.com/yspennstate/trace-reinforced-ants-dead-branches`.

The paper is part of the Math Brain project, whose public repository is `https://github.com/yspennstate/math-brain-public` [19].

Appendix A. Notation

Symbols are listed roughly in the order in which they first appear, and the last column gives that place. The letters a, c, C, t and u are reused, and each meaning has its own row.

Symbol	Meaning	First appears
$G = (V, E)$	a finite connected simple graph, its vertices and its edges	Section 1
N, F	the nest and the food, two distinct marked vertices	Section 1
n	the number of ants that have walked	Section 1
$W(n)$, $W_e(n)$	the weight vector after n ants, and its value at the edge e	Section 1
x, z	vertices; W_{xz} is the weight of the edge $\{x, z\}$	Section 1
$W_x(n)$	the total weight of the edges at the vertex x	Section 1
$\mathbf{1}\{\cdot\}$	the indicator of an event	Section 1
χ, χ_e	the deterministic limit of $W(n)/n$ in the conjectures, and its value at e	Conjecture 1
reduced	said of (G, N, F) when $G - F$ is connected	Section 1
u	a neighbour of F	Theorem 2
u	a shadow vertex, in (D2)	Section 4
H	a set of vertices joined to the rest of $G - F$ by a single edge; in (D2), the far side of a bridge	Theorem 2
$e = \{u, y\}$	the single edge joining H to $V \setminus (H \cup \{F\})$; elsewhere e is any edge	Theorem 2
f	an edge with an endpoint in H	Theorem 2
a, b, c	vertices of the examples	Example 4
w, w_e, w_x	a weight vector, its value at the edge e, and the total weight of the edges at x	Section 1
$p_e(w)$	the probability that the w-weighted walk from N crosses e before it hits F	Section 1
$\mathbb{E}$, $\mathcal{F}_n$	expectation, and the σ-field generated by the first n ants	Section 1

Symbol	Meaning	First appears
equilibrium	a solution of $p(w) = w$	Section 1
$\Gamma(F)$	the set of neighbours of F	Theorem 5
I_0	the connected component of N in $G - F - \Gamma(F)$ ($I_0 = \emptyset$ in Section 4 when N is adjacent to F)	Theorem 5
a	the edge $\{u, F\}$, in the proof of Theorem 2	Section 2
X_n, A_n	the weights $W_e(n)$ and $W_a(n)$	Section 2
R	the total weight of the edges at u other than a and e	Section 2
h	the probability that the walk from N visits u before F	Section 2
ρ, ρ'	the probabilities that the walk returns to u after the step $u \to y$, and after a step along another edge	Section 2
S	$A + R(1 - \rho')$	Section 2
$\deg(x)$	the degree of the vertex x	Section 2
c	the constant $1/(\deg(u)\deg(y))$	Section 2
Λ_n	$\log(X_n/A_n)$	Section 2
D_n, D'_n	the indicators that the $(n+1)$-st ant crosses e and a	Section 2
t	a real number in $\log(1 + t)$	Section 2
t	a value of v greater than 1, in the proof	Proposition 8
t	the time of the flow $\dot{w} = p(w) - w$	Section 3
M_n, Q	the martingale part of Λ_n, and the sum of the second-order terms	Section 2
δ, m	a positive number, and a time with $X_m \geq \delta m$	Section 2
$g(x)$	the expected number of visits to x by the w-walk from N before it hits F	Section 3
$v(x)$	the voltage $g(x)/w_x$ of the unit current from N to F, with $v(F) = 0$	Section 3
V_1	the set of neighbours x of F with $w_{xF} > 0$	Proposition 8
C	the set of vertices joined to z by a path of the support along which $v \geq t$, in the proof	Proposition 8
ε	a small weight put on one edge	Section 3
$\mathcal{A}$	the set of neighbours of F with a neighbour in I_0 ($\mathcal{A} = \{N\}$ when N is adjacent to F)	Section 4
shadow	the vertex set $V \setminus (I_0 \cup \{F\})$	Section 4
$\mathcal{D}(G, N, F)$	the set of edges of kinds (D1) and (D2)	Section 4
Dp, I	the derivative of p, and the identity	Section 4
$\Theta(L_1, \ldots, L_k)$	the graph made of $k \geq 2$ internally disjoint paths from N to F, the i-th with L_i edges	Section 5
α_i	the ratio of the weights of consecutive edges of the i-th path	Section 5
$a_{i,j}$, c_i, C	the rate of excursions from N that reach depth j of path i, the rate of those that reach F along path i, and $\sum_i c_i$, in the proof	Proposition 10
$p_{i,j}$	$p_e(w)$ for the j-th edge from N of path i	Proposition 10

References

[1] G. Aletti, A. Collevecchio, I. Crimaldi, M. Deijfen, S. Dereich, A. Ghiglietti, R. van der Hofstad, M. Holmes, D. Kious, V. Kleptsyn, B. Lodewijks, P. Mörters and B. Schapira, Open problems in stochastic reinforcement models, in *2023 MATRIX Annals*, MATRIX Book Series, Springer, 2025, 477–498.

[2] W. Armfield and C. Mailler, The forward-loop-erased ants walk, arXiv:2610.18453.

[3] M. Benaïm, Dynamics of stochastic approximation algorithms, in *Séminaire de Probabilités XXXIII*, Lecture Notes in Math. 1709, Springer, 1999, 1–68.

[4] B. Davis, Reinforced random walk, *Probab. Theory Related Fields* **84** (1990), 203–229.

[5] J.-L. Deneubourg, S. Aron, S. Goss and J. M. Pasteels, The self-organizing exploratory pattern of the Argentine ant, *J. Insect Behav.* **3** (1990), 159–168.

[6] M. Dorigo and T. Stützle, *Ant Colony Optimization*, MIT Press, 2004.

[7] D. Erhard, T. Franco and G. Reis, The directed edge reinforced random walk: the ant mill phenomenon, *J. Stat. Phys.* **190** (2023), no. 1, Paper No. 18.

[8] R. van der Hofstad, M. Holmes, A. Kuznetsov and W. Ruszel, Strongly reinforced Pólya urns with graph-based competition, *Ann. Appl. Probab.* **26** (2016), no. 4, 2494–2539.

[9] L. C. Le Goff and O. Raimond, Vertex reinforced non-backtracking random walks: an example of path formation, *Electron. J. Probab.* **23** (2018), Paper No. 39, 1–38.

[10] R. Pemantle, Nonconvergence to unstable points in urn models and stochastic approximations, *Ann. Probab.* **18** (1990), 698–712.

[11] R. Pemantle, A survey of random processes with reinforcement, *Probab. Surv.* **4** (2007), 1–79.

[12] D. Kious, C. Mailler and B. Schapira, Finding geodesics on graphs using reinforcement learning, *Ann. Appl. Probab.* **32** (2022), no. 5, 3889–3929; arXiv:2010.04820.

[13] D. Kious, C. Mailler and B. Schapira, The trace-reinforced ants process does not find shortest paths, *J. Éc. polytech. Math.* **9** (2022), 505–536; arXiv:2106.10559.

[14] C. Mailler and Z. Varin, The two-nest ants process on triangle-series-parallel graphs, arXiv:2601.22855.

[15] A. A. Hagberg, D. A. Schult and P. J. Swart, Exploring network structure, dynamics, and function using NetworkX, in *Proceedings of the 7th Python in Science Conference (SciPy 2008)*, 2008, 11–15.

[16] C. R. Harris, K. J. Millman, S. J. van der Walt, R. Gommers, P. Virtanen, D. Cournapeau, E. Wieser, J. Taylor, S. Berg, N. J. Smith, R. Kern, M. Picus, S. Hoyer, M. H. van Kerkwijk, M. Brett, A. Haldane, J. Fernández del Río, M. Wiebe, P. Peterson, P. Gérard-Marchant, K. Sheppard, T. Reddy, W. Weckesser, H. Abbasi, C. Gohlke and T. E. Oliphant, Array programming with NumPy, *Nature* **585** (2020), 357–362.

[17] R. C. Read and R. J. Wilson, *An Atlas of Graphs*, Oxford University Press, 1998.

[18] Y. Shmalo, *A non-geodesic edge that survives in the loop-erased ant process*, 2026, `https://github.com/yspennstate/loop-erased-ants-non-geodesic-edge`.

[19] Y. Shmalo, *Math Brain: public snapshot, code and tests*, 2026, `https://github.com/yspennstate/math-brain-public`.

[20] Y. Shmalo, *Math Brain: shared, auditable memory for multi-agent mathematical research*, October 2026, `https://github.com/yspennstate/math-brain-public/tree/main/papers/math-brain-2026-10-24`.

[21] Y. Shmalo, *Brain Framework*, public software repository, 2026, `https://github.com/yspennstate/brain-framework-public`.