

## OXIDATIVE STRESS

**Effects of Ozone on Leaf Senescence, Photochemical Efficiency and Grain Yield in Two Winter Wheat Cultivars**S. Burkart<sup>1</sup>, J. Bender<sup>2</sup>, B. Tarkotta<sup>2</sup>, S. Faust<sup>2</sup>, A. Castagna<sup>3</sup>, A. Ranieri<sup>3</sup> & H.-J. Weigel<sup>2</sup><sup>1</sup> Thünen-Institute of Climate-Smart Agriculture Braunschweig, Germany<sup>2</sup> Thünen-Institute of Biodiversity Braunschweig, Germany<sup>3</sup> Department of Crop Biology, University of Pisa Pisa, Italy**Keywords**

chlorophyll fluorescence; ozone; photosynthesis; senescence; wheat; yield

**Correspondence**

S. Burkart

Thünen-Institute of Climate-Smart Agriculture  
Bundesallee 50  
D-38116 Braunschweig  
Germany

Tel.: +49 531 596 2546

Fax: +49 531 596 2599

Email: stefan.burkart@ti.bund.de

Accepted October 30, 2012

doi:10.1111/jac.12013

**Abstract**

The adverse effects of tropospheric ozone ( $O_3$ ) on crop photosynthesis, growth and yield have been documented in numerous studies over the last decades, but little information from field experimentation exists on how modern European winter wheat cultivars respond to  $O_3$ . Two winter wheat cultivars (Astron and Pegassos) differing in development characteristics were exposed to non-filtered ambient air or non-filtered air plus 30 ppb and non-filtered air plus 60 ppb  $O_3$  ( $8 \text{ h day}^{-1}$ ) in open-top field chambers. At several dates during growth, green leaf area was determined by destructive harvests. Leaf gas exchange, pigment content and xanthophyll cycle activity, and photochemical efficiency by chlorophyll a fluorescence were measured.  $O_3$  exposure induced accelerated senescence with no difference between cultivars. Photosynthesis declined especially in Pegassos; however, stomatal conductance was hardly affected by  $O_3$ . Pigment contents were reduced by  $O_3$  exposure, and de-epoxidation index increased. Photochemical efficiency ( $F_v/F_m$ ) declined, whereas actual quantum yield ( $\Phi_{PSII}$ ) did not respond to  $O_3$ .  $O_3$  exposure reduced grain yield in both cultivars. However, yield of Pegassos was more affected by  $O_3$  exposure than yield of Astron, suggesting a higher  $O_3$  sensitivity of Pegassos. The data presented in this manuscript indicate a need to test whether high-yield varieties such as Pegassos are particularly sensitive to  $O_3$  exposure.

**Introduction**

Tropospheric ozone ( $O_3$ ) is a widespread secondary air pollutant found in all industrialized countries worldwide, and it is considered the most significant phytotoxic pollutant in the atmosphere affecting vegetation. Predictive models indicate that  $O_3$  concentrations will continue to increase at a rate of 0.5–2 % per year in the Northern hemisphere during the next several decades (IPCC 2007). To date, the background  $O_3$  concentration in the Northern hemisphere is within the range of 23–34 ppb (Vingarzan 2004). Current levels of  $O_3$  pollution are known to be high enough to decrease carbon assimilation and to suppress growth and biomass production in many crop and wild plant species (Bender and Weigel 2003, 2011, Ashmore 2005, Mills et al. 2010, Pleijel 2011). Moreover,  $O_3$  has been shown to mitigate possible benefits due to expected climate change in Europe (Clausen et al. 2011). The

observed decline in the yield of major crops may imply a growing threat from ozone pollution to the security of food supplies (Weigel and Bender 2012, Wilkinson et al. 2012).

Ozone enters leaves through stomata, and it is thought to rapidly react with the aqueous phase of the mesophyll cell wall matrix to form highly reactive oxygen species (ROS). As a consequence of this oxidative stress, accelerated senescence and leaf damages were often observed as one of the harmful effects on plants caused by  $O_3$  (Wohlge-muth et al. 2002, Fuhrer and Booker 2003). Photosynthetic membranes are seen as main targets of ROS (Fiscus et al. 2005). Impairments of light reactions of photosynthesis can be detected by studying chlorophyll fluorescence, which is an acknowledged non-invasive tool, even before visible symptoms emerge (Sayed 2003).  $O_3$  stress is known to reduce both the maximum ( $F_v/F_m$ ) and effective quantum yield ( $\Phi_{PSII}$ ) of photosystem II photochemistry (Carrasco-Rodriguez and del Valle Tascon 2001, Calatayud 2007, Feng

et al. 2008). O<sub>3</sub>-related damages of photosynthetic processes will increase the need to dissipate excessive energy due to overreduction of the photosystems. The de-epoxidation of violaxanthin (V) via antheraxanthin (A) to zeaxanthin (Z) is an important process in the dissipation of excess energy and protection of the photosynthetic membranes from injury (Niyogi 1999). The de-epoxidation state of the system (DEPS), that is, the proportion of de-epoxidized epoxide groups in total epoxide groups, indicates the degree of activation of light protection processes. A higher de-epoxidation state is related to lower light harvesting capacity and more energy dissipation in the pigment (Demmig-Adams and Adams 1992). Moreover, carotenoids such as  $\beta$ -carotene and Z are known to act as protectors of photosynthetic membranes against ROS (Young et al. 1997, Jahns et al. 2009). A stimulation of the xanthophyll cycle activity was observed especially in O<sub>3</sub>-sensitive species (Castagna and Ranieri 2009), indicating enhanced non-photochemical dissipation processes.

Winter wheat is the predominant cereal crop across Europe. Wheat is known to be one of the most ozone-sensitive crops (Mills et al. 2007), but the vast majority of the research on wheat has been performed on spring wheat cultivars (Feng et al. 2008, Grünhage et al. 2012). Exposure or dose–response functions gained from these studies are used for quantifying effects of ozone on wheat yield and for the derivation of critical levels for O<sub>3</sub> on crops in Europe (Mills et al. 2007, 2011, Grünhage et al. 2012). However, there is a substantial variability in O<sub>3</sub> responses between cultivars (Bender et al. 1999, Feng et al. 2008, Gonzalez-Fernandez et al. 2010), and knowledge about O<sub>3</sub> effects on modern winter wheat cultivars is particularly scarce. For example, it is unclear whether winter wheat cultivars differing in ontogenesis and performance also differ in their sensitivity to O<sub>3</sub>.

The objective of this study was to test how photosynthesis and senescence of winter wheat cultivars respond to chronic O<sub>3</sub> exposure and whether the responses can be related to yield responses and to differences in growth characteristics between the cultivars.

## Material & Methods

### Plant material, growth conditions and experimental design

An open-top field chamber (OTC) experiment was conducted in 2006 in an agro-ecosystem at the Johann Heinrich von Thünen-Institute (vTI) in Braunschweig, Lower Saxony, Germany (10°26'E, 52°18'N, 79 m a.s.l.), to examine the effects of O<sub>3</sub> on plant growth, yield and physiology in two field-grown winter wheat cultivars (CV): *Triticum aestivum* cv. Astron (*Ast*) and Pegassos (*Peg*). Astron is a medium late ripening blending wheat cultivar, with medium self-shading

capacity (Saaten-Union 2012a). Pegassos is a medium early ripening quality wheat cultivar, with high self-shading capacity and drought resistance (Saaten-Union 2012b).

For the last 30 years, the local climate at the field site is characterized by a mean annual air temperature of 8.8 °C and a total precipitation rate of 618 mm year<sup>-1</sup> (Germany's National Meteorological Service). The soil at the site is a luvisol of a loamy sand texture, with a pH of 6.3–6.5. For at least 30 years, the field has been used for agriculture. After ploughing and preparation of the field with a cultivator, plants were sown with 0.12-m row spacing in October 2005 (230 kg seeds ha<sup>-1</sup>). Agricultural management measures were carried out according to local farming practices (Table 1). Total nitrogen added to the respective experimental area was 188 kg N ha<sup>-1</sup>. Pesticide treatments were performed on demand to control insects and fungal infection. None of the pesticides used (Table 1) have been reported to affect crop responses to O<sub>3</sub>. From begin of May until mid-June, experimental plots were regularly irrigated as needed to prevent drought stress. Total irrigation during this period was 68 mm. Plants were exposed to O<sub>3</sub> in OTCs, 3 m in diameter by 2.3 m height (for details refer to Weigel and Jäger 1988) beginning on 9 May 2006 (stem elongation). The centres of the OTCs were separated from each other by at least 12 m. Plants were fumigated with O<sub>3</sub> for 54 days (Table 1). Three O<sub>3</sub> treatments were employed in four replicate open-top chambers (n = 4): non-filtered (NF) ambient air, NF with additional 30 ppb (NF+) or NF with additional 60 ppb O<sub>3</sub> (NF++) for 8 h day<sup>-1</sup> (10:00–18:00). Starting from May 22nd, O<sub>3</sub>-dosage in NF++ was reduced to 40 ppb due to the occurrence of visible symptoms in the highest ozone treatment.

**Table 1** Timetable of events during the 2006 open-top chamber experiment GS = plant growth stages according to Totman and Broad (1987)

Event/growth stage	Date
Sowing	07 Oct 2005
Emergence	16 Oct 2005
GS 31 (first node)	04 May 2006
GS 65 (mid-anthesis)	14 June
GS 92 (grain maturity)	22 July
Installation of OTCs	April 2006
Start of ozone fumigation	09 May
End of ozone fumigation	02 July
Days of exposure	54
Fertilizer application (kg N ha <sup>-1</sup> )	10 Mar (48)
	21 April (90)
	22 May (50)
Insecticides application ( <i>Karate Zeon</i> <sup>TM</sup> )	31 May
Fungicides application ( <i>Opus Top</i> <sup>TM</sup> )	08 May
	31 May
	21 June

### Assessment of senescence and yield

Five to seven plants per cultivar and OTC were harvested at four harvest dates (04.05.2006 tillering, growth stage (GS) 31; 17.05.2006 booting, GS 39; 07.06.2006, pre-anthesis; 20.06.2006 grain filling, GS 71) and separated in leaves, stems and ears. The final harvest was done at grain maturity. Leaves were separated in green and yellow/dead leaves, and the projected area of green leaves (GLA) was measured with a LAI 3100 instrument (Li-COR, Lincoln, NE, USA). At maturity, grain yield was determined after drying to constant weight and threshing.

### Chlorophyll fluorescence

Chlorophyll fluorescence measurements were taken with a pulse-modulated chlorophyll fluorometer (FMS 2; Hansatech Instruments Ltd, Norfolk UK) *in situ* on flag leaves (Schreiber et al. 1986, Krause and Weis 1991). There is a discrepancy between the need to select the most representative part of the leaf and to avoid using yellow/dead leaves for fluorescence measurements. Anyway, for the measurements, only leaf parts that were still green were chosen, putting up with a possible bias in the results.

Maximum photochemical efficiency of photosystem II ( $F_v/F_m$ , Bolh ar-Nordenkamp et al. 1989) was measured daily from May 9th to July 2nd. Leaf clamps were fixed on the mid of flag leaf of two of the marked plants to pre-darken the leaves for 30 min. After opening the gate of the leaf clamp, a weak pulsed measuring light was given to obtain ground fluorescence  $F_0$ . A short intense light pulse was then given to close all reaction centres, and maximum fluorescence  $F_m$  was recorded.  $F_v/F_m$  was calculated as

$$F_v/F_m = (F_m - F_0)/F_m. \quad (1)$$

The actual quantum yield of photosystem II reaction centres ( $\Phi_{PSII}$ , Genty et al. 1989) was measured twice per week on mid of unshaded flag leaves of all marked plants. The leaf was fixed in the leaf holder and kept under natural light conditions. A weak pulsed measuring light was given to obtain steady-state fluorescence  $F_s$ . A short intense light pulse was then given to close all reaction centres, and then the maximum fluorescence  $F'_m$  was recorded. The actual quantum yield of photosystem II was calculated as

$$\Phi_{PSII} = (F'_m - F_s)/F'_m. \quad (2)$$

### Pigment analysis

Total chlorophyll content was estimated with a SPAD Chlorophyll meter (Minolta SPAD-502 (Uddling et al. 2007)) two times a week on several spots of flag leaves (except leaf base and apex) of the marked plants.

Flag leaves of at least 10 plants were harvested at anthesis (growth stage 61). Pigments were analysed by means of HPLC, according to the method of Castagna et al. (2001). Leaves were homogenized in the dark in 100 % HPLC-grade acetone with sodium ascorbate and filtered through 0.2- $\mu$ m filters. The separation was performed using a Zorbax ODS column (Chrompack SA, 5 mm particle size, 250  $\times$  4.6 mm diameter). Pigments were eluted using 100 % solvent A (acetonitrile/methanol, 75/25, v/v) for the first 15 min, followed by a 2.5-min linear gradient to 100 % solvent B (methanol/ethylacetate, 68/32, v/v) which continued isocratically until the end of the 32-min separation. The flow rate was 1 ml min<sup>-1</sup>. The pigments were detected by their absorbance at 445 nm. To quantify pigment content, known amounts of a pure standard were injected into the HPLC system and an equation, correlating peak area to pigment concentration was formulated. Pigment concentration was expressed as nmol g<sup>-1</sup> fresh weight. The de-epoxidation index (DEPS, %) was calculated as

$$DEPS = (A/2 + Z)/(V + A + Z) \times 100, \quad (3)$$

where A, Z and V represent the content in antheraxanthin, zeaxanthin and violaxanthin, respectively.

### Leaf gas exchange

Leaf gas exchange was measured on flag leaves using an open gas-exchange system (LI-6400; LI-COR). Gas-exchange measurements were taken from May 5th to June 25th, from 08:30–19:00 h. The sequence of OTCs measured was randomized each day to exclude any bias caused by diurnal variation. Measurements were taken under natural light conditions using the transparent cuvette. Photosynthetic photon flux density (PPFD,  $\mu$ mol m<sup>-2</sup> s<sup>-1</sup>) and air temperature (T,  $^{\circ}$ C) were recorded at the leaf level. The water vapour concentration of air entering the chamber was not controlled and therefore tracked ambient conditions. The chamber [CO<sub>2</sub>] was set to 380  $\mu$ mol mol<sup>-1</sup>. Readings were taken once the coefficient of variation (stability variables: sample cell (CO<sub>2</sub>), sample cell (H<sub>2</sub>O) and flow) of the infrared gas analyzer output was below 0.7 %. Each reading took at least eight minutes to ensure that  $g_s$  had enough time to equilibrate to the vapour pressure deficit (VPD, kPa) in the leaf chamber. Net photosynthesis ( $P_N$ ,  $\mu$ mol m<sup>-2</sup> s<sup>-1</sup>), stomatal conductance ( $g_s$ , mol m<sup>-2</sup> s<sup>-1</sup>) and the ratio of leaf internal/ambient CO<sub>2</sub> concentration ( $C_i/C_a$ ) were calculated using the equations of von Von Caemmerer and Farquhar (1981).

### Data processing

The effects of O<sub>3</sub> on leaf growth and senescence, gas exchange and fluorescence as dependent variables were

tested by three-factorial analysis of variance (date  $\times$  O<sub>3</sub>  $\times$  CV) for  $n = 4$  replicated chambers. Average values of the variables calculated for each chamber half were used as statistical unit. Data were analysed with the R Software Package V. 2.12.0 (<http://www.r-project.org/>) as a completely randomized design with NF, NF+ and NF++ treatments ( $n = 4$  each) each split for cultivar (total  $n = 24$ ). As pigments and yield were only analysed for one date (anthesis and final harvest, resp.), two-way ANOVA was applied in this case. P-values lower than 0.05 were considered as significant.

## Results

### Ozone concentration

O<sub>3</sub> concentration in the NF-chambers superseded 40 ppb in the first half of May and at the end of the measuring period, and 60 ppb in mid-June (Fig. 1). In the OTCs with O<sub>3</sub> enrichment, maximum values of O<sub>3</sub>-concentration reached


60 ppb in NF+ and 80 ppb in NF++ in these periods. From the end of May, O<sub>3</sub>-dosage was reduced to +25 ppb (NF+) and +40 ppb (NF++) due to the occurrence of visible injuries. The 8-h seasonal means for the whole exposure period were 36 ppb in NF, 51 ppb in NF+ and 58 ppb in NF++. The resulting average O<sub>3</sub> exposure (accumulated exposure over threshold 40 ppb; AOT40) was 1.70 ppm.h in NF, 6.46 ppm.h in NF+ and 10.04 ppm.h in NF++.

### Visual symptoms of senescence


Ozone exposure caused leaf damages in *Ast* and *Peg*, visible as spots, necrosis and yellowing. In the course of the season, different patterns of injury between the cultivars were observed: a rather homogenous yellowing occurred in *Peg*, whereas in *Ast*, increasing proportions of the leaf blade completely died off with a corresponding loss of green leaf area. The green parts, however, remained nearly completely green, with no substantial change in colour.

### Green leaf area

Green leaf area (GLA) increased to ca. 60 cm<sup>2</sup> per plant at GS 39 and remained constant until anthesis in NF plants (Fig. 2). GLA of *Ast* tended to be lower than of *Peg*, but this was not significant ( $P = 0.23$  at GS 31). In both cultivars, O<sub>3</sub> exposure induced a significant decline of GLA starting from the 2nd harvest in both O<sub>3</sub>-treatments ( $P < 0.001$ ). The reduction in GLA due to O<sub>3</sub>-fumigation reached 75 % at GS 56, with no differences between NF+ and NF++ or between cultivars.



**Fig. 1** Seasonal courses of O<sub>3</sub>-concentrations (8 h-mean, 10–18 h), AOT40 and weather parameters. NF: Non-filtered ambient air, NF+: NF + 30 ppb O<sub>3</sub>, NF++: NF + 60 ppb O<sub>3</sub>. VPD vapour pressure deficit (hPa), T temperature (°C), Rg global radiation (W m<sup>-2</sup>). Arrows indicate the start of the reduction in the amount of O<sub>3</sub> added in NF++ to 40 ppb.




**Fig. 2** Green leaf area as determined from destructive harvest at different growth stages. NF: Non-filtered ambient air, NF+: NF + 30 ppb O<sub>3</sub>, NF++: NF + 60 ppb O<sub>3</sub>. Error bars represent standard error of mean ( $n = 4$ ).



### Gas exchange

Due to high variation of the data, three-way ANOVA indicated that there was neither any difference between cultivars in  $g_s$ , nor any  $O_3$  effect on  $g_s$  (Fig. 3a), although  $g_s$  of *Peg* seemed to be slightly reduced by  $O_3$  after anthesis. In both cultivars,  $O_3$  exposure reduced  $P_N$  after booting stage ( $P = 0.049$ , Fig. 3b). At grain filling stage,  $O_3$  exposure reduced  $P_N$  by 30 % in *Ast* and 43 % in *Peg*.  $P_N$  of *Peg* tended to be lower than that of *Ast* especially at the end of the season ( $P = 0.095$ ). The ratio internal/ambient  $CO_2$  concentration ( $C_i/C_a$ ) did not differ between cultivars, but was increased by  $O_3$  at the 2nd and 3rd measuring date




**Fig. 3** A Stomatal conductance ( $g_s$ ), B net photosynthesis ( $P_N$ ) and C the ratio internal/ambient  $CO_2$  concentration ( $C_i/C_a$ ) averaged across different measuring periods (booting 17.05.–24.05., anthesis 07.06.–14.06., ripening 25.06.). NF: Non-filtered ambient air, NF+: NF + 30 ppb  $O_3$ , NF++: NF + 60 ppb  $O_3$ . Error bars represent standard error of mean ( $n = 4$ ).

( $P = 0.026$ , Fig. 3c). There was no  $O_3 \times CV$  interaction on leaf gas-exchange parameters at any growth stage.

### Pigments

In both cultivars, SPAD values (estimate of total chlorophyll) of flag leaves declined in the course of the growth period ( $P < 0.001$ , Fig. 4). At all growth stages, the SPAD value of *Ast* was higher than that of *Peg* ( $P < 0.001$ ). The  $O_3$ -exposure reduced SPAD at all growth stages ( $P < 0.001$ ). There were no interactions between cultivar and  $O_3$  at any growth stage, although the SPAD values tended to reflect the differences in visual symptoms described above.

As found for SPAD values, the concentrations of all pigments measured were significantly lower in *Peg* than in *Ast* ( $P = 0.031$  for total chlorophylls,  $P = 0.030$  for total carotenoids,  $P = 0.060$  for  $\beta$ -carotene,  $P = 0.032$  for the  $V + A + Z$  sum,  $P = 0.017$  for total xanthophylls) when averaged across all  $O_3$ -treatments, while the de-epoxidation state of the xanthophyll cycle was similar between the two cultivars (Fig. 5).  $O_3$  exposure resulted in a clear decline in pigment concentrations ( $P < 0.001$ ), with no  $O_3 \times CV$  interaction for any pigment. The small differences in pigment concentrations between NF+ and NF++ were not significant. The percentage of decrease was higher for chlorophylls (–41 % and –54 % in *Ast*, and –55 % and –64 % in *Peg*, in NF+ and NF++ samples, respectively) than for carotenoids (–34 % and –44 % in *Ast*, and –41 % and –50 % in *Peg*, in NF+ and NF++ samples, respectively). Statistical analysis showed a significant increase in DEPS due to  $O_3$  exposure ( $P = 0.013$ ) with no



**Fig. 4** Relative chlorophyll content measured with SPAD chlorophyll meter at different growth stages. NF: non-filtered ambient air, NF+: NF + 30 ppb  $O_3$ , NF++: NF + 60 ppb  $O_3$ . Error bars represent standard error of mean ( $n = 4$ ).

difference between NF+ and NF++, whereas there were no differences in DEPS between cultivars.


### Chlorophyll a fluorescence

Throughout the measuring period,  $F_v/F_m$  (potential quantum yield, Fig. 6a) was higher in *Ast* than in *Peg* ( $P = 0.016$ ). At GS 39,  $O_3$ -fumigation did not affect  $F_v/F_m$ . However, starting with GS 56,  $F_v/F_m$  responded to  $O_3$ -fumigation in both cultivars ( $P < 0.001$ ). Moreover, especially in NF++,  $F_v/F_m$  declined much more in *Peg* than in *Ast* at GS 56 ( $P = 0.068$ ). These changes in  $F_v/F_m$  at GS56 were mainly caused by a decrease in  $F_v$  ( $P = 0.024$ ), whereas  $F_0$  remained unaffected by  $O_3$  (Fig. 6c,d). Towards the end of the measuring period,  $F_v/F_m$  recovered in both cultivars. After recovery, there were no significant responses of  $F_v/F_m$  to  $O_3$  nor any interactions between  $O_3$  and cultivars. However, the recovery of  $F_v/F_m$  at GS 71 was mainly due to a decrease in  $F_0$  ( $P = 0.086$ ) which appeared to be more pronounced in *Peg*, but this difference between cultivars was not significant.


With respect to climatic conditions and growth stages, the measuring period was divided into three sub-periods: 1: May 23rd to June 6th, 2: June 7th to June 19th, 3: June 28th.  $\Phi_{PSII}$  (actual quantum yield) varied with PPFD (Fig. 6b), but was not affected by  $O_3$ -fumigation at any period ( $P = 0.27$ ). As was found for  $F_v/F_m$ ,  $\Phi_{PSII}$  of *Peg* was lower than of *Ast* at all growth stages ( $P = 0.004$ ). There was no  $O_3 \times CV$  interaction on  $\Phi_{PSII}$  at any growth stage.

### Grain yield and biomass

Two-factorial ANOVA showed that grain yield of *Peg* was significantly higher than yield of *Ast* (+29.1 % in NF, +19.6 % in NF+ and +10.5 % in NF++,  $P = 0.004$ , Table 2). Similar results were found for straw mass ( $1031.5 \pm 31.8 \text{ gm}^{-2}$  in *Peg*,  $832.0 \pm 34.2 \text{ g m}^{-2}$  in *Ast*, data not shown).  $O_3$ -fumigation reduced grain yield of both cultivars ( $P < 0.001$ ). In *Ast*, the reduction of grain yield amounted -13.1 % in NF+ and -19.4 % in NF++, whereas in *Peg*  $O_3$ -fumigation reduced grain yield by -19.5 % in NF+ and -31.1 % in NF++. However, a



**Fig. 5** Content of A chlorophyll a + b (Chl), B total carotenoids (C), C total xanthophylls (X), D  $\beta$ -carotene, E the sum of violaxanthin (V), antheraxanthin (A) and zeaxanthin (Z), and (F) de-epoxidation index (DEPS) at anthesis. NF: Non-filtered ambient air, NF+: NF + 30 ppb  $O_3$ , NF++: NF + 60 ppb  $O_3$ . Error bars represent standard error of mean (n = 4).



**Fig. 6** (a) Maximum quantum yield ( $F_v/F_m$ ) of photosystem II, (b) effective quantum yield ( $\Phi_{PSII}$ ) of photosystem II, (c) ground fluorescence ( $F_0$ ) and (d) variable fluorescence ( $F_v$ ) at different growth stages. NF: Non-filtered ambient air, NF+: NF + 30 ppb  $O_3$ , NF++: NF + 60 ppb  $O_3$ . Error bars represent standard error of mean ( $n = 4$ ). Please note for panel B: Values of  $\Phi_{PSII}$  were averaged across three different measuring periods. Hence, growth stage numbers in panel B cover 23.05.–06.06. (GS 39), 07.06.–19.06. (GS 56), 28.06. (GS 71). Numbers above columns indicate mean photon flux densities ( $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) during measurements.

**Table 2** Grain yield ( $\text{g m}^{-2}$ ) as determined at maturity (GS 92). NF: non-filtered ambient air, NF+: NF + 30 ppb  $O_3$ , NF++: NF + 60 ppb  $O_3$ . Values represent means  $\pm$  standard error of means (S.E.,  $n = 4$ )

	Astron	Pegassos
NF		
Mean	688.1	888.4
$\pm$ S.E.	30.1	30.4
NF+		
Mean	598.1	715.3
$\pm$ S.E.	50.6	49.8
NF++		
Mean	554.3	612.5
$\pm$ S.E.	54.1	52.8

significant  $O_3 \times \text{CV}$  interaction on grain yield could not be proven ( $P = 0.32$ ).

## Discussion

In this study, chronic  $O_3$  exposure decreased photosynthesis and caused accelerated senescence in two winter wheat cultivars. Both cultivars responded to the chronic  $O_3$  exposure by reducing green leaf area, via an accelerated

senescence and resulting in a loss in grain yield. This was also found in many but not all studies with wheat (Reichenauer et al. 1998, Ewert and Pleijel 1999, Sild et al. 2002).

Our data showed only a slight decrease in stomatal conductance in *Peg* caused by  $O_3$  fumigation, especially after anthesis. However, the overall reduction in  $g_s$  of wheat due to  $O_3$  as reported by Feng et al. (2008) could not be confirmed in the present study, which may be due to small effects and the high variability of the data. Moreover, both the differences in the means of  $g_s$  between the cultivars and the effects of  $O_3$  on  $g_s$  were smaller than the  $O_3$ -effects on  $P_N$ . Ozone exposure led to a decline in net  $\text{CO}_2$  fixation in both cultivars, which is in line with many other experiments with wheat (Feng et al. 2008). However, in several studies, photosynthetic and growth responses to  $O_3$  could not be related to variations in  $g_s$  (Reichenauer et al. 1998, Herbiner et al. 2002, Akhtar et al. 2010, Feng et al. 2011). Although  $g_s$  likewise tended to decrease in this study in response to  $O_3$ , the reduction in photosynthesis was mainly due to decreases in  $\text{CO}_2$ -fixation processes (Singh et al. 2009) because  $C_i/C_a$  was increased under  $O_3$  fumigation, whereas  $\Phi_{PSII}$  did not respond to  $O_3$ .

The ozone-induced decrease in both chlorophyll and carotenoid concentration suggests a generalized negative impact of the pollutant on photosynthetic pigments, although chlorophylls tended to be more affected than carotenoids. Carotenoids, apart from their role as secondary light-absorbing pigments, are able to reduce the chlorophyll triplet state and to prevent the formation of harmful singlet oxygen or to scavenge it after its production by the interaction of triplet chlorophyll with  $O_2$  (Young et al. 1997). It is well known that when absorbed light exceeds plant photochemical requirement as may occur also under normal light irradiation, but in the presence of environmental constraints, this excess energy may be transferred to the ever-present oxygen, leading to the formation of ROS and, ultimately, to damage at the photosystem level (Castagna et al. 2001, Ort 2001, Demmig-Adams and Adams 2006). In this respect,  $O_3$  stress resembles plant responses to many other abiotic stresses such as drought (Singh et al. 2012), UV-B radiation (Lidon et al. 2012) or heat stress (Dias et al. 2011).  $CO_2$  assimilation is the major sink for the reducing equivalents and energy produced by the primary photochemical reactions. Therefore, a diminished  $CO_2$  fixation will induce a decreased demand for ATP and NADPH and, consequently, may lead to an over-reduction in PSII. Ozone-treated samples, which exhibited a decline in net  $CO_2$  fixation in comparison with their respective controls, might be expected to activate the xanthophyll cycle to dissipate the large amount of excitation energy absorbed by the antenna. The increase in DEPS observed in both wheat cultivars under the thylakoid capacity to react and acclimate to  $O_3$ , allowing a decrease in the PSII photochemical rate and subsequent alleviation of excitation pressure on the reaction centres.

In this study, chronic  $O_3$  exposure reduced the performance of photosynthetic electron transport chain. The exposure to  $O_3$  induced a decrease in  $F_v/F_m$  more in *Peg* than in *Ast*. Similar to our results,  $F_v/F_m$  was found to decline under  $O_3$  exposure in soybean (Chen et al. 2009), in *Quercus mongolica* (Wang et al. 2009b) and in wheat at grain filling (Wang et al. 2009a). However, after reducing the amount of  $O_3$  added after 22nd May,  $F_v/F_m$  recovered to nearly initial values especially in *Peg*. These changes in  $F_v/F_m$  may indicate a plasticity of the photosynthetic system (Bussotti et al. 2011). Reichenauer et al. (1998) showed that a recovery of  $F_v/F_m$  may occur even overnight but not in all wheat cultivars of their study. The decrease in  $F_0$  as observed in this study, however, indicates an increased thermal dissipation and not an inactivation of photosystem II (Guidi et al. 2000, 2002, Leipner et al. 2001, Bussotti et al. 2011), which is in line with the observed increase in DEPS. The decrease in  $F_v$  under  $O_3$  fumigation may be attributed to the pigment losses.

In contrast to other experiments with wheat (Feng et al. 2008),  $O_3$  fumigation impaired the performance of photosynthetic electron transport ( $\Phi_{PSII}$ ) only marginally. Adams and Demmig-Adams (2004) showed that  $\Phi_{PSII}$  cannot be seen as a measure of Pn. Together with the sensitivity of  $F_v/F_m$  to  $O_3$ , this is a further evidence that it is not the electron transport chain *per se* that is impaired by  $O_3$  but the light harvesting capacity due to pigment loss and the use of reductive equivalents. This may reflect a strategy described by Bussotti et al. (2011) for poplar showing that leaves try to maintain the efficiency of the remaining green leaf area around necrotic regions at an optimal level. Literature, however, is unequivocal although most studies reported a decrease in  $\Phi_{PSII}$  under  $O_3$  fumigation depending on species, cultivars and exposure duration (Ciompi et al. 1997, Endo et al. 2005, Calatayud et al. 2006, Degl'Innocenti et al. 2007, Wang et al. 2009a). Moreover, in this study, senescence behaviour differed between cultivars, thus preferring leaf parts that are still alive for fluorescence measurements may cause a bias in the data. Only a few studies addressed the spatial heterogeneity of visible damages by likewise selecting green leaves or areas for measurement (Guidi et al. 2000, Carrasco-Rodriguez and del Valle Tascon 2001). Moreover, visible damages do not necessarily entail changes in fluorescence parameters and *vice versa* (Carrasco-Rodriguez and del Valle Tascon 2001, Konishi et al. 2005).

The observed  $O_3$ -induced yield reductions in both cultivars (yield reduction in  $NF^{++}$  of 19 % and 31 % in *Ast* and *Peg*, respectively) suggest that modern winter wheat cultivars exhibited a similar high  $O_3$  sensitivity as spring-sown cultivars do. The meta-analysis performed by Feng et al. (2008) revealed that  $O_3$  exposure to an average of 72 ppb (7-h seasonal mean) reduces wheat grain yield by 29 % (95 % confidence interval: 24–34 %). The majority of studies considered in this meta-analysis were performed with spring wheat. Nevertheless, a direct comparison between spring wheat and winter wheat in the responses to elevated  $O_3$  is difficult to draw due to the lack of comparative experiments with the two wheat types in which they are simultaneously exposed to  $O_3$ . However, in the present study, grain yield of *Peg* responded more sensitively to  $O_3$  exposure as compared to *Ast* which is correlated with the overall higher growth and yield performance of *Peg*. Photosynthesis was also more affected by  $O_3$  in *Peg* than in *Ast*, as indicated by a larger reduction in  $P_n$  and the decrease in  $F_v/F_m$ . The physiological or genetic basis for the variation in  $O_3$  sensitivity between genotypes is poorly understood (Fiscus et al. 2005); however, the differences in  $O_3$  sensitivity between the two cultivars cannot be attributed to differences in  $g_s$  and thus to differences in  $O_3$  uptake or flux. Moreover, there are examples where the  $O_3$  response of grain yield did not necessarily correlate to the extent of leaf



damage (Picchi et al. 2010). The data presented in this manuscript indicate a need to test whether high-yield varieties such as *Peg* are particularly sensitive to O<sub>3</sub> exposure.

## Acknowledgements

We thank Anke Mundt, Carina Trenkler, Peter Braunisch and Ralf Staudte for technical assistance in the field and in the laboratory, and Annette Freibauer for helpful discussions and comments on the manuscript.

## References

- Adams, W. W., and B. Demmig-Adams, 2004: Chlorophyll fluorescence as a tool to monitor plant response to the environment. In chlorophyll a fluorescence. A signature of photosynthesis. In: G. C. Papageorgiou, and G. Govindjee, eds. *Advances in Photosynthesis and Respiration Series*, pp. 583–604. Springer, Dordrecht, the Netherlands.
- Akhtar, N., M. Yamaguchi, H. Inada, D. Hoshino, T. Kondo, and T. Izuta, 2010: Effects of ozone on growth, yield and leaf gas exchange rates of two Bangladeshi cultivars of wheat (*Triticum aestivum* L.). *Environ. Pollut.* 158, 1763–1767.
- Ashmore, M. R., 2005: Assessing the future global impacts of ozone on vegetation. *Plant Cell Environ.* 28, 949–964.
- Bender, J., and H. J. Weigel, 2003: Ozone stress impacts on plant life. In: R. S. Ambasht, and N. K. Ambasht, eds. *Modern Trends in Applied Terrestrial Ecology*, pp. 165–182. Kluwer Academic/Plenum Publishers, New York, NY, USA.
- Bender, J., and H.-J. Weigel, 2011: Changes in atmospheric chemistry and crop health: a review. *Agron. Sustain. Dev.* 31, 81–89.
- Bender, J., U. G. Hertstein, and C. R. Black, 1999: Growth and yield responses of spring wheat to increasing carbon dioxide, ozone and physiological stresses: a statistical analysis of 'ESPACE-wheat' results. *Eur. J. Agron.* 10, 185–195.
- Bolh r-Nordenkamp, H. R., S. P. Long, N. R. Baker, G. Oquist, U. Schreiber, and E. G. Lechner, 1989: Chlorophyll fluorescence as a probe of the photosynthetic competence of leaves in the field: a review of current instrumentation. *Funct. Ecol.* 3, 497–514.
- Bussotti, F., R. Desotgiu, C. Cascio, M. Pollastrini, E. Gravano, G. Gerosa, R. Marzuoli, C. Nali, G. Lorenzini, E. Salvatori, F. Manes, M. Schaub, and R. J. Strasser, 2011: Ozone stress in woody plants assessed with chlorophyll a fluorescence. A critical reassessment of existing data. *Environ. Exp. Bot.* 73, 19–30.
- Calatayud, A., 2007: Chlorophyll a fluorescence as indicator of atmospheric pollutant effects. *Toxicol. Environ. Chem.* 89, 627–639.
- Calatayud, A., F. Pomares, and E. Barreno, 2006: Interactions between nitrogen fertilization and ozone in watermelon cultivar Reina de Corazones in open-top chambers. Effects on chlorophyll a fluorescence, lipid per oxidation, and yield. *Photosynthetica* 44, 93–101.
- Carrasco-Rodriguez, J. L., and S. del Valle Tascon, 2001: Impact of elevated ozone on chlorophyll a fluorescence in field-grown oat (*Avena sativa*). *Environ. Exp. Bot.* 45, 133–142.
- Castagna, A., and A. Ranieri, 2009: Detoxification and repair process of ozone injury: from O<sub>3</sub> uptake to gene expression adjustment. *Environ. Pollut.* 157, 1461–1469.
- Castagna, A., C. Nali, S. Ciompi, G. Lorenzini, G. F. Soldatini, and A. Ranieri, 2001: Ozone exposure affects photosynthesis of pumpkin (*Cucurbita pepo*) plants. *New Phytol.* 152, 223–229.
- Chen, C. P., T. D. Frank, and S. P. Long, 2009: Is a short, sharp shock equivalent to long-term punishment? Contrasting the spatial pattern of acute and chronic ozone damage to soybean leaves via chlorophyll fluorescence imaging. *Plant Cell Environ.* 32, 327–335.
- Ciompi, S., A. Castagna, A. Ranieri, C. Nali, G. Lorenzini, and G. F. Soldatini, 1997: CO<sub>2</sub> assimilation, xanthophyll cycle pigments and PSII efficiency in pumpkin plants as affected by ozone fumigation. *Physiol. Plant.* 101, 881–889.
- Clausen, S. K., G. Frenck, L. G. Linden, T. N. Mikkelsen, C. Lunde, and R. B. J rgensen, 2011: Effects of Single and Multifactor Treatments with Elevated Temperature, CO<sub>2</sub> and Ozone on Oilseed Rape and Barley. *J. Agron. Crop Sci.* 197, 442–453.
- Degl'Innocenti, E., L. Guidi, and G. F. Soldatini, 2007: Effects of elevated ozone on chlorophyll a fluorescence in symptomatic and asymptomatic leaves of two tomato genotypes. *Biol. Plant.* 51, 313–321.
- Demmig-Adams, B., and W. W. Adams, 1992: Photoprotection and other responses of plants to high light stress. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 43, 599–626.
- Demmig-Adams, B., and W. W. Adams, 2006: Photoprotection in an ecological context: the remarkable complexity of thermal energy dissipation. *New Phytol.* 172, 11–21.
- Dias, A. S., J. Semedo, J. C. Ramalho, and F. C. Lidon, 2011: Bread and durum wheat under heat stress: a comparative study on the photosynthetic performance. *J. Agron. Crop Sci.* 197, 50–56.
- Endo, R., A. Konishi, and K. Omasa, 2005: Effect of ozone exposure on the photosynthetic activity at cellular level: chlorophyll fluorescence microscopic imaging analysis. *Phyton-Annales Rei Botanicae* 45, 493–496.
- Ewert, F., and H. Pleijel, 1999: Phenological development, leaf emergence, tillering and leaf area index, and duration of spring wheat across Europe in response to CO<sub>2</sub> and ozone. *Eur. J. Agron.* 10, 171–184.
- Feng, Z., K. Kobayashi, and E. A. Ainsworth, 2008: Impact of elevated ozone concentration on growth, physiology, and yield of wheat (*Triticum aestivum* L.): a meta-analysis. *Glob. Change Biol.* 14, 2696–2708.
- Feng, Z., J. Pang, K. Kobayashi, J. Zhu, and D. R. Ort, 2011: Differential responses in two varieties of winter wheat to elevated ozone concentration under fully open-air field conditions. *Glob. Change Biol.* 17, 580–591.
- Fiscus, E. L., F. L. Booker, and K. O. Burkey, 2005: Crop responses to ozone: uptake, modes of action, carbon assimilation and partitioning. *Plant Cell Environ.* 28, 997–1011.

- Fuhrer, J., and F. Booker, 2003: Ecological issues related to ozone: agricultural issues. *Environ. Int.* 29, 141–154.
- Genty, B., J. M. Briantais, and N. R. Baker, 1989: The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochim. Biophys. Acta* 990, 87–92.
- Gonzalez-Fernandez, I., A. Kaminska, M. Dodmani, E. Goumenaki, S. Quarrie, and J. D. Barnes, 2010: Establishing ozone flux-response relationships for winter wheat: analysis of uncertainties based on data for UK and Polish genotypes. *Atmos. Environ.* 44, 621–630.
- Grünhage, L., H. Pleijel, G. Mills, J. Bender, H. Danielsson, Y. Lehmann, J.-F. Castell, and O. Bethenod, 2012: Updated stomatal flux and flux-effect models for wheat for quantifying effects of ozone on grain yield, grain mass and protein yield. *Environ. Pollut.* 165, 147–157.
- Guidi, L., R. Di Cagno, and G. F. Soldatini, 2000: Screening of bean cultivars for their response to ozone as evaluated by visible symptoms and leaf chlorophyll fluorescence. *Environ. Pollut.* 107, 349–355.
- Guidi, L., E. Degl'Innocenti, and G. F. Soldatini, 2002: Assimilation of CO<sub>2</sub>, enzyme activation and photosynthetic electron transport in bean leaves, as affected by high light and ozone. *New Phytol.* 156, 377–388.
- Herbinger, K., M. Tausz, A. Wonisch, G. Soja, A. Sorger, and D. Grill, 2002: Complex interactive effects of drought and ozone stress on the antioxidant defence systems of two wheat cultivars. *Plant Physiol. Biochem.* 40, 691–696.
- IPCC, 2007: Climate change 2007: the physical science basis. In: Solomon S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor, H.L. Miller, et al., (eds). Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, p. 996. Cambridge University Press, UK and USA.
- Jahns, P., D. Latowski, and K. Strzalka, 2009: Mechanism and regulation of the violaxanthin cycle: the role of antenna proteins and membrane lipids. *Biochim. Biophys. Acta* 1787, 3–14.
- Konishi, A., R. Endo, and K. Omasa, 2005: Imaging of the red and far-red chlorophyll fluorescence of the ozone injured plant leaf. *Phyton – Annales Rei Botanicae* 45, 487–491.
- Krause, G. H., and E. Weis, 1991: Chlorophyll fluorescence and photosynthesis: the basics. *Annu. Rev. Plant Physiol.* 42, 313–349.
- Leipner, J., K. Oxborough, and N. R. Baker, 2001: Primary sites of ozone-induced perturbations of photosynthesis in leaves: identification and characterization in *Phaseolus vulgaris* using high resolution chlorophyll fluorescence imaging. *J. Exp. Bot.* 52, 1689–1696.
- Lidon, F. J. C., M. Teixeira, and J. C. Ramalho, 2012: Decay of the Chloroplast Pool of Ascorbate Switches on the Oxidative Burst in UV-B-Irradiated Rice. *J. Agron. Crop Sci.* 198, 130–144.
- Mills, G., A. Buse, B. Gimeno, V. Bermejo, M. Holland, L. Emberson, and H. Pleijel, 2007: A synthesis of AOT40-based response functions and critical levels of ozone for agricultural and horticultural crops. *Atmos. Environ.* 41, 2630–2643.
- Mills, G., F. Hayes, D. Simpson, L. Emberson, D. Norris, H. Harmens, and P. Büker, 2010: Evidence of widespread effects of ozone on crops and (semi-)natural vegetation in Europe (1990–2006) in relation to AOT40- and flux-based risk maps. *Glob. Change Biol.* 17, 592–613.
- Mills, G., H. Pleijel, S. Braun, P. Büker, V. Bermejo, E. Calvo, H. Danielsson, L. Emberson, I. G. Fernández, L. Grünhage, H. Harmens, F. Hayes, P.-E. Karlsson, and D. Simpson, 2011: New stomatal flux-based critical levels for ozone effects on vegetation. *Atmos. Environ.* 45, 5064–5068.
- Niyogi, K. K., 1999: Photoprotection revisited: genetic and molecular approaches. *Annu. Rev. Plant Physiol. Plant. Mol. Biol.* 50, 333–359.
- Ort, D. R., 2001: When there is too much light. *Plant Physiol.* 125, 29–32.
- Picchi, V., M. Iriti, S. Quaroni, M. Saracchi, P. Viola, and F. Faoro, 2010: Climate variations and phenological stages modulate ozone damages in field-grown wheat. A three-year study with eight modern cultivars in Po Valley (Northern Italy). *Agric. Ecosyst. Environ.* 135, 310–317.
- Pleijel, H., 2011: Reduced ozone by air filtration consistently improved grain yield in wheat. *Environ. Pollut.* 159, 897–902.
- Reichenauer, T. G., B. A. Goodman, P. Kostecki, and G. Soja, 1998: Ozone sensitivity in *Triticum durum* and *T. aestivum* with respect to leaf injury, photosynthetic activity and free radical content. *Physiol. Plant.* 104, 681–686.
- Saaten-Union, 2012a: Astron a winter wheat. <http://www.saaten-union.de/index.cfm/product/1,605,html.html> [last accessed 8 August 2012].
- Saaten-Union, 2012b: Pegasso a winter wheat. <http://www.saaten-union.de/index.cfm/product/1,609,html.html>, [last accessed 8 August 2012].
- Sayed, O. H., 2003: Chlorophyll fluorescence as a tool in cereal crop research. *Photosynthetica* 41, 321–330.
- Schreiber, U., U. Schliwa, and W. Bilger, 1986: Continuous recording of photochemical and nonphotochemical chlorophyll fluorescence quenching with a new type of modulation fluorometer. *Photosynth. Res.* 10, 51–62.
- Sild, E., H. Pleijel, and G. Sellden, 2002: Elevated ozone (O<sub>3</sub>) alters carbohydrate metabolism during grain filling in wheat (*Triticum aestivum* L.). *Agric. Ecosyst. Environ.* 92, 71–81.
- Singh, E., S. Tiwari, and M. Agrawal, 2009: Effects of elevated ozone on photosynthesis and stomatal conductance of two soybean varieties: a case study to assess impacts of one component of predicted global climate change. *Plant Biol.* 11, 101–108.
- Singh, S., A. K. Gupta, and N. Kaur, 2012: Differential Responses of Antioxidative Defence System to Long-Term Field Drought in Wheat (*Triticum aestivum* L.) Genotypes Differing in Drought Tolerance. *J. Agron. Crop Sci.* 198, 185–195.
- Tottman, D. R., and H. Broad, 1987: The decimal code for the growth stages of cereals, with illustrations. *Ann. Appl. Biol.* 110, 441–454.
- Uddling, J., J. Gelang-Alfredsson, K. Piikki, and H. Pleijel, 2007: Evaluating the relationship between leaf chlorophyll concentration and SPAD-502 chlorophyll meter readings. *Photosynth. Res.* 91, 37–46.

- Vingarzan, R., 2004: A review of surface ozone background levels and trends. *Atmos. Environ.* 38, 3431–3442.
- Von Caemmerer, S., and G. D. Farquhar, 1981: Some relationships between the biochemistry of photosynthesis and the gas exchange of leaves. *Planta* 153, 376–387.
- Wang, L., Q. Zeng, Z. Feng, J. Zhu, H. Tang, X. Chen, Z. Xie, G. Liu, and K. Kobayashi, 2009a: Photosynthetic damage induced by elevated  $O_3$  in two varieties of winter wheat with free air controlled enrichment approach. *Huanjing Kexue* 30, 527–534.
- Wang, L. L., X. Y. He, and W. Chen, 2009b: Effects of elevated ozone on photosynthetic  $CO_2$  exchange and chlorophyll a fluorescence in leaves of *Quercus mongolica* grown in urban area. *Bull Environ Contam Toxicol* 82, 478–481.
- Weigel, H.-J., and J. Bender, 2012: Bodennahes Ozon – ein Problem für Kulturpflanzen und Ernährungssicherheit? *Gesunde Pflanzen* 64, 79–87.
- Weigel, H. J., and H.-J. Jäger, 1988: Zur Ökotoxikologie von Luftschadstoffen. II. Aufbau und Funktionsweise einer Expositionsanlage aus open-top Kammern zur Untersuchung von Immissionswirkungen auf Pflanzen. *Landbauforschung Voelkenrode* 38, 182–195.
- Wilkinson, S., G. Mills, R. Illidge, and W. J. Davies, 2012: How is ozone pollution reducing our food supply? *J. Exp. Bot.* 63, 527–536.
- Wohlgemuth, H., K. Mittelstrass, S. Kschieschan, J. Bender, H. J. Weigel, K. Overmyer, J. Kangasjarvi, H. Sandermann, and C. Langebartels, 2002: Activation of an oxidative burst is a general feature of sensitive plants exposed to the air pollutant ozone. *Plant Cell Environ.* 25, 717–726.
- Young, A. J., D. Phillip, and J. Savill, 1997: Carotenoids in higher plant photosynthesis. In: M. Pessarakli, ed. *Handbook of Photosynthesis*, pp. 575–596. Marcel Dekker, New York, NY, USA.