

# Regular Positive Representations

Ivan Ip

Hong Kong University of Science and Technology

March 22, 2023

Advances in Cluster Algebra 2023

# Overview

Algebraic set up:

- Quantum group  $\mathcal{U}_q(\mathfrak{g})$ , semisimple Lie type
- Quantum cluster algebra  $\mathcal{O}_q(\mathcal{X}^{\mathbf{Q}})$  associated to initial quiver  $\mathbf{Q}$
- Regularity:  $\mathcal{U}_q(\mathfrak{g}) \longrightarrow \mathcal{O}_q(\mathcal{X}^{\mathbf{Q}})$

Self-adjoint conditions:  $X_i^* = X_i, |q| = 1$ :

- Polarization of cluster variables  $\implies$  representations of  $\mathcal{O}_q(\mathcal{X}^{\mathbf{Q}})$  on some Hilbert space  $\mathcal{H}$  by Heisenberg operators  $x_i, p_i$ .
- Induces positive representations of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$  on  $\mathcal{H}$  by positive self-adjoint operators.

**Goal:** Classify the regular positive representations obtained this way.

# Overview

Algebraic set up:

- Quantum group  $\mathcal{U}_q(\mathfrak{g})$ , semisimple Lie type
- Quantum cluster algebra  $\mathcal{O}_q(\mathcal{X}^{\mathbf{Q}})$  associated to initial quiver  $\mathbf{Q}$
- Regularity:  $\mathcal{U}_q(\mathfrak{g}) \longrightarrow \mathcal{O}_q(\mathcal{X}^{\mathbf{Q}})$

Self-adjoint conditions:  $X_i^* = X_i$ ,  $|q| = 1$ :

- Polarization of cluster variables  $\implies$  representations of  $\mathcal{O}_q(\mathcal{X}^{\mathbf{Q}})$  on some Hilbert space  $\mathcal{H}$  by Heisenberg operators  $x_i, p_i$ .
- Induces positive representations of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$  on  $\mathcal{H}$  by positive self-adjoint operators.

Goal: Classify the regular positive representations obtained this way.

# Overview

Algebraic set up:

- Quantum group  $\mathcal{U}_q(\mathfrak{g})$ , semisimple Lie type
- Quantum cluster algebra  $\mathcal{O}_q(\mathcal{X}^{\mathbf{Q}})$  associated to initial quiver  $\mathbf{Q}$
- Regularity:  $\mathcal{U}_q(\mathfrak{g}) \longrightarrow \mathcal{O}_q(\mathcal{X}^{\mathbf{Q}})$

Self-adjoint conditions:  $X_i^* = X_i$ ,  $|q| = 1$ :

- Polarization of cluster variables  $\implies$  representations of  $\mathcal{O}_q(\mathcal{X}^{\mathbf{Q}})$  on some Hilbert space  $\mathcal{H}$  by Heisenberg operators  $x_i, p_i$ .
- Induces positive representations of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$  on  $\mathcal{H}$  by positive self-adjoint operators.

**Goal:** Classify the regular positive representations obtained this way.

# Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

# Definition of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$

## Definition

$\mathcal{U}_q(\mathfrak{sl}_2)$  = Hopf algebra  $\langle \mathbf{E}, \mathbf{F}, \mathbf{K}^{\pm 1} \rangle$  over  $\mathbb{C}(q)$  such that

$$\mathbf{K}\mathbf{E} = q^2\mathbf{E}\mathbf{K}, \quad \mathbf{K}\mathbf{F} = q^{-2}\mathbf{F}\mathbf{K}, \quad [\mathbf{E}, \mathbf{F}] = \frac{\mathbf{K} - \mathbf{K}^{-1}}{q - q^{-1}}.$$

Coproduct:

$$\begin{aligned} \Delta(\mathbf{E}) &= 1 \otimes \mathbf{E} + \mathbf{E} \otimes \mathbf{K}, & \Delta(\mathbf{F}) &= \mathbf{F} \otimes 1 + \mathbf{K}^{-1} \otimes \mathbf{F}, \\ \Delta(\mathbf{K}) &= \mathbf{K} \otimes \mathbf{K}. \end{aligned}$$

(Also counit  $\varepsilon$ , antipode  $S$ .)

$\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ :  $(|q| = 1, e^{\pi i b^2}, b^2 \in (0, 1) \setminus \mathbb{Q})$

$$\mathbf{E}^* = \mathbf{E}, \quad \mathbf{F}^* = \mathbf{F}, \quad \mathbf{K}^* = \mathbf{K}, \quad q^* = q^{-1}.$$

# Definition of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$

## Definition

$\mathcal{U}_q(\mathfrak{sl}_2)$  = Hopf algebra  $\langle \mathbf{E}, \mathbf{F}, \mathbf{K}^{\pm 1} \rangle$  over  $\mathbb{C}(q)$  such that

$$\mathbf{K}\mathbf{E} = q^2\mathbf{E}\mathbf{K}, \quad \mathbf{K}\mathbf{F} = q^{-2}\mathbf{F}\mathbf{K}, \quad [\mathbf{E}, \mathbf{F}] = \frac{\mathbf{K} - \mathbf{K}^{-1}}{q - q^{-1}}.$$

Coproduct:

$$\begin{aligned} \Delta(\mathbf{E}) &= 1 \otimes \mathbf{E} + \mathbf{E} \otimes \mathbf{K}, & \Delta(\mathbf{F}) &= \mathbf{F} \otimes 1 + \mathbf{K}^{-1} \otimes \mathbf{F}, \\ \Delta(\mathbf{K}) &= \mathbf{K} \otimes \mathbf{K}. \end{aligned}$$

(Also counit  $\varepsilon$ , antipode  $S$ .)

$\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ :  $(|q| = 1, e^{\pi i b^2}, b^2 \in (0, 1) \setminus \mathbb{Q})$

$$\underline{\mathbf{E}^* = \mathbf{E}}, \quad \underline{\mathbf{F}^* = \mathbf{F}}, \quad \underline{\mathbf{K}^* = \mathbf{K}}, \quad q^* = q^{-1}.$$

# Definition of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

## Definition

$\mathcal{U}_q(\mathfrak{g}) =$  Hopf-algebra  $\langle \mathbf{E}_i, \mathbf{F}_i, \mathbf{K}_i^{\pm 1} \rangle_{i \in I}$  over  $\mathbb{C}(q)$  such that

$$\mathbf{K}_i \mathbf{E}_j = q_i^{a_{ij}} \mathbf{E}_j \mathbf{K}_i, \quad \mathbf{K}_i \mathbf{F}_j = q_i^{-a_{ij}} \mathbf{F}_j \mathbf{K}_i, \quad [\mathbf{E}_i, \mathbf{F}_j] = \delta_{ij} \frac{\mathbf{K}_i - \mathbf{K}_i^{-1}}{q_i - q_i^{-1}}$$

+ Serre's relations.  $(q_i = q^{d_i})$

Coproduct:

$$\begin{aligned} \Delta(\mathbf{E}_i) &= 1 \otimes \mathbf{E}_i + \mathbf{E}_i \otimes \mathbf{K}_i, & \Delta(\mathbf{F}_i) &= \mathbf{F}_i \otimes 1 + \mathbf{K}_i^{-1} \otimes \mathbf{F}_i, \\ \Delta(\mathbf{K}_i) &= \mathbf{K}_i \otimes \mathbf{K}_i. \end{aligned}$$

(Also counit  $\varepsilon$ , antipode  $S$ ).

$$\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}}): \quad (|q| = 1, e^{\pi i b^2}, b^2 \in (0, 1) \setminus \mathbb{Q})$$

$$\mathbf{E}_i^* = \mathbf{E}_i, \quad \mathbf{F}_i^* = \mathbf{F}_i, \quad \mathbf{K}_i^* = \mathbf{K}_i, \quad q^* = q^{-1}.$$



# Definition of $\mathcal{D}_q(\mathfrak{g}_{\mathbb{R}})$

## Definition

$\mathcal{D}_q(\mathfrak{g}) =$  *Drinfeld's Double*:  $\langle \mathbf{E}_i, \mathbf{F}_i, \mathbf{K}_i^{\pm 1}, \mathbf{K}'_i{}^{\pm 1} \rangle_{i \in I}$

of  $\mathcal{U}_q(\mathfrak{g})$

$$\mathbf{K}_i \mathbf{E}_j = q_i^{a_{ij}} \mathbf{E}_j \mathbf{K}_i, \quad \mathbf{K}_i \mathbf{F}_j = q_i^{-a_{ij}} \mathbf{F}_j \mathbf{K}_i, \quad [\mathbf{E}_i, \mathbf{F}_j] = \delta_{ij} \frac{\mathbf{K}_i - \mathbf{K}'_i}{q_i - q_i^{-1}}$$

+ Serre relations + similarly for  $\mathbf{K}'_i$ .

Coproduct:

$$\begin{aligned} \Delta(\mathbf{E}_i) &= 1 \otimes \mathbf{E}_i + \mathbf{E}_i \otimes \mathbf{K}_i, & \Delta(\mathbf{F}_i) &= \mathbf{F}_i \otimes 1 + \mathbf{K}'_i \otimes \mathbf{F}_i, \\ \Delta(\mathbf{K}_i) &= \mathbf{K}_i \otimes \mathbf{K}_i, & \Delta(\mathbf{K}'_i) &= \mathbf{K}'_i \otimes \mathbf{K}'_i. \end{aligned}$$

(Also counit  $\varepsilon$ , antipode  $S$ ).

$$\mathcal{U}_q(\mathfrak{g}) = \mathcal{D}_q(\mathfrak{g}) / \langle \mathbf{K}_i \mathbf{K}'_i = 1 \rangle_{i \in I}.$$

$$\mathbf{e}_i := -\mathbf{i}(q_i - q_i^{-1})\mathbf{E}_i, \quad \mathbf{f}_i := -\mathbf{i}(q_i - q_i^{-1})\mathbf{F}_i, \quad i \in I.$$

# Definition of $\mathcal{D}_q(\mathfrak{g}_{\mathbb{R}})$

## Definition

$\mathcal{D}_q(\mathfrak{g}) =$  *Drinfeld's Double*:  $\langle \mathbf{E}_i, \mathbf{F}_i, \mathbf{K}_i^{\pm 1}, \mathbf{K}'_i{}^{\pm 1} \rangle_{i \in I}$

$$\mathbf{K}_i \mathbf{E}_j = q_i^{a_{ij}} \mathbf{E}_j \mathbf{K}_i, \quad \mathbf{K}_i \mathbf{F}_j = q_i^{-a_{ij}} \mathbf{F}_j \mathbf{K}_i, \quad [\mathbf{E}_i, \mathbf{F}_j] = \delta_{ij} \frac{\mathbf{K}_i - \mathbf{K}'_i}{q_i - q_i^{-1}}$$

+ Serre relations + similarly for  $\mathbf{K}'_i$ .

Coproduct:

$$\begin{aligned} \Delta(\mathbf{E}_i) &= 1 \otimes \mathbf{E}_i + \mathbf{E}_i \otimes \mathbf{K}_i, & \Delta(\mathbf{F}_i) &= \mathbf{F}_i \otimes 1 + \mathbf{K}'_i \otimes \mathbf{F}_i, \\ \Delta(\mathbf{K}_i) &= \mathbf{K}_i \otimes \mathbf{K}_i, & \Delta(\mathbf{K}'_i) &= \mathbf{K}'_i \otimes \mathbf{K}'_i. \end{aligned}$$

(Also counit  $\varepsilon$ , antipode  $S$ ).

$$\mathcal{U}_q(\mathfrak{g}) = \mathcal{D}_q(\mathfrak{g}) / \langle \mathbf{K}_i \mathbf{K}'_i = 1 \rangle_{i \in I}.$$

$$\mathbf{e}_i := -\mathbf{i}(q_i - q_i^{-1})\mathbf{E}_i, \quad \mathbf{f}_i := -\mathbf{i}(q_i - q_i^{-1})\mathbf{F}_i, \quad i \in I.$$

# Definition of $\mathcal{D}_q(\mathfrak{g}_{\mathbb{R}})$

## Definition

$\mathcal{D}_q(\mathfrak{g}) =$  *Drinfeld's Double*:  $\langle \mathbf{E}_i, \mathbf{F}_i, \mathbf{K}_i^{\pm 1}, \mathbf{K}'_i{}^{\pm 1} \rangle_{i \in I}$

$$\mathbf{K}_i \mathbf{E}_j = q_i^{a_{ij}} \mathbf{E}_j \mathbf{K}_i, \quad \mathbf{K}_i \mathbf{F}_j = q_i^{-a_{ij}} \mathbf{F}_j \mathbf{K}_i, \quad [\mathbf{E}_i, \mathbf{F}_j] = \delta_{ij} \frac{\mathbf{K}_i - \mathbf{K}'_i}{q_i - q_i^{-1}}$$

+ Serre relations + similarly for  $\mathbf{K}'_i$ .

Coproduct:

$$\begin{aligned} \Delta(\mathbf{E}_i) &= 1 \otimes \mathbf{E}_i + \mathbf{E}_i \otimes \mathbf{K}_i, & \Delta(\mathbf{F}_i) &= \mathbf{F}_i \otimes 1 + \mathbf{K}'_i \otimes \mathbf{F}_i, \\ \Delta(\mathbf{K}_i) &= \mathbf{K}_i \otimes \mathbf{K}_i, & \Delta(\mathbf{K}'_i) &= \mathbf{K}'_i \otimes \mathbf{K}'_i. \end{aligned}$$

(Also counit  $\varepsilon$ , antipode  $S$ ).

$$\mathcal{U}_q(\mathfrak{g}) = \mathcal{D}_q(\mathfrak{g}) / \langle \mathbf{K}_i \mathbf{K}'_i = 1 \rangle_{i \in I}.$$

$$2 \sin \pi b^2 > 0$$

$$\mathbf{e}_i := -\mathbf{i}(q_i - q_i^{-1})\mathbf{E}_i, \quad \mathbf{f}_i := -\mathbf{i}(q_i - q_i^{-1})\mathbf{F}_i, \quad i \in I.$$

Example:  $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ 

$$e^{2\pi b p} f(x) = f(x - ib)$$

$$e^{\pi b(u-2p)} := q e^{\pi b u} e^{-2\pi b p}$$

Theorem (Faddeev, Teschner (1999))

Irreducible representations  $\mathcal{P}_\lambda$  of  $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$  parametrized by  $\lambda \geq 0$ : ( $p = \frac{1}{2\pi i} \frac{d}{du}$ )

$$\pi_\lambda(\mathbf{e}) := e^{\pi b(u+2p+2\lambda)} + e^{\pi b(-u+2p-2\lambda)} \quad u \quad p$$

$$\pi_\lambda(\mathbf{f}) := e^{\pi b(u-2p)} + e^{\pi b(-u-2p)}$$

$$\pi_\lambda(\mathbf{K}) := e^{\pi b(2u+2\lambda)}$$

 $C \in L^2(\mathbb{R})$ acting on  $L^2(\mathbb{R})$  as (unbounded essentially) positive self-adjoint operators.(Irreducible: the only operators *strongly commute* with them are the scalars.)

Corollary

The Casimir element

$$\mathbf{C} := \mathbf{f}\mathbf{e} - q\mathbf{K} - q^{-1}\mathbf{K}$$

acts on  $\mathcal{P}_\lambda$  by the scalar

$$\pi_\lambda(\mathbf{C}) = e^{2\pi b\lambda} + e^{-2\pi b\lambda} \geq 2.$$

# Example: $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$

Theorem (*Faddeev, Teschner (1999)*)

Irreducible representations  $\mathcal{P}_\lambda$  of  $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$  parametrized by  $\lambda \geq 0$ : ( $p = \frac{1}{2\pi i} \frac{d}{du}$ )

$$\pi_\lambda(\mathbf{e}) := e^{\pi b(u+2p+2\lambda)} + e^{\pi b(-u+2p-2\lambda)}$$

$$\pi_\lambda(\mathbf{f}) := e^{\pi b(u-2p)} + e^{\pi b(-u-2p)}$$

$$\pi_\lambda(\mathbf{K}) := e^{\pi b(2u+2\lambda)}$$

acting on  $L^2(\mathbb{R})$  as (unbounded essentially) positive self-adjoint operators.

(Irreducible: the only operators *strongly commute* with them are the scalars.)

Corollary

The Casimir element

$$\mathbf{C} := \mathbf{f}\mathbf{e} - q\mathbf{K} - q^{-1}\mathbf{K}$$

acts on  $\mathcal{P}_\lambda$  by the scalar

$$\pi_\lambda(\mathbf{C}) = e^{2\pi b\lambda} + e^{-2\pi b\lambda} \geq 2.$$

# Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations  $\mathcal{P}_{\lambda}$  by positive operators on Hilbert space.
- = “Quantization of principal series representations”.
- Generalization of Teschner’s representations of  $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ .

## Theorem (I. (2012))

*There exists a family of irreducible representations  $\mathcal{P}_{\lambda}$  of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ :*

- *Parametrized by  $\lambda \in \mathbb{R}_{\geq 0} P^+ \simeq \mathbb{R}_{\geq 0}^{n=\text{rank } \mathfrak{g}}$ .*
- *Positivity:  $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i\}$  are represented by positive, essentially self-adjoint (unbounded) operators on  $L^2(\mathbb{R}^N)$ ,  $N := \ell(w_0)$ .*
- *$\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i$  are expressed in terms of Laurent polynomials of  $\{e^{\pi b u_k}, e^{2\pi b p_k}\}_{k=1}^N$  where  $[p_j, u_k] = \frac{\delta_{jk}}{2\pi i}$ .*
- *Transcendental relations (compatibility with modular double  $b \longleftrightarrow b^{-1}$ ) (Langland’s duality).*

Can be understood in terms of quantum cluster algebra!

# Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations  $\mathcal{P}_{\lambda}$  by positive operators on Hilbert space.
- = “Quantization of principal series representations”.
- Generalization of Teschner’s representations of  $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ .

## Theorem (I. (2012))

*There exists a family of irreducible representations  $\mathcal{P}_{\lambda}$  of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ :*

- *Parametrized by  $\lambda \in \mathbb{R}_{\geq 0} P^+ \simeq \mathbb{R}_{\geq 0}^{n=\text{rank } \mathfrak{g}}$ .*
- *Positivity:  $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i\}$  are represented by positive, essentially self-adjoint (unbounded) operators on  $L^2(\mathbb{R}^N)$ ,  $N := \ell(w_0)$ .*
- *$\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i$  are expressed in terms of Laurent polynomials of  $\{e^{\pi b u_k}, e^{2\pi b p_k}\}_{k=1}^N$  where  $[p_j, u_k] = \frac{\delta_{jk}}{2\pi i}$ .*
- *Transcendental relations (compatibility with modular double  $b \longleftrightarrow b^{-1}$ ) (Langland’s duality).*

Can be understood in terms of quantum cluster algebra!

# Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations  $\mathcal{P}_{\lambda}$  by positive operators on Hilbert space.
- = “Quantization of principal series representations”.
- Generalization of Teschner’s representations of  $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ .

## Theorem (I. (2012))

*There exists a family of irreducible representations  $\mathcal{P}_{\lambda}$  of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ :*

- *Parametrized by  $\lambda \in \mathbb{R}_{\geq 0} P^+ \simeq \mathbb{R}_{\geq 0}^{n=\text{rank } \mathfrak{g}}$ .*
- *Positivity:  $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i\}$  are represented by positive, essentially self-adjoint (unbounded) operators on  $L^2(\mathbb{R}^N)$ ,  $N := \ell(w_0)$ .*
- *$\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i$  are expressed in terms of Laurent polynomials of  $\{e^{\pi b u_k}, e^{2\pi b p_k}\}_{k=1}^N$  where  $[p_j, u_k] = \frac{\delta_{jk}}{2\pi i}$ .*
- *Transcendental relations (compatibility with modular double  $b \longleftrightarrow b^{-1}$ ) (Langland’s duality).*

Can be understood in terms of quantum cluster algebra!



# Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations  $\mathcal{P}_{\lambda}$  by positive operators on Hilbert space.
- = “Quantization of principal series representations”.
- Generalization of Teschner’s representations of  $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ .

## Theorem (I. (2012))

*There exists a family of irreducible representations  $\mathcal{P}_{\lambda}$  of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ :*

- *Parametrized by  $\lambda \in \mathbb{R}_{\geq 0} P^+ \simeq \mathbb{R}_{\geq 0}^{n=\text{rank } \mathfrak{g}}$ .*
- *Positivity:  $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i\}$  are represented by positive, essentially self-adjoint (unbounded) operators on  $L^2(\mathbb{R}^N)$ ,  $N := \ell(w_0)$ .*
- *$\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i$  are expressed in terms of Laurent polynomials of  $\{e^{\pi b u_k}, e^{2\pi b p_k}\}_{k=1}^N$  where  $[p_j, u_k] = \frac{\delta_{jk}}{2\pi i}$ .*
- *Transcendental relations (compatibility with modular double  $b \longleftrightarrow b^{-1}$ ) (Langland’s duality).*

Can be understood in terms of quantum cluster algebra!

# Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations  $\mathcal{P}_{\lambda}$  by positive operators on Hilbert space.
- = “Quantization of principal series representations”.
- Generalization of Teschner’s representations of  $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ .

## Theorem (I. (2012))

*There exists a family of irreducible representations  $\mathcal{P}_{\lambda}$  of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ :*

- *Parametrized by  $\lambda \in \mathbb{R}_{\geq 0} P^+ \simeq \mathbb{R}_{\geq 0}^{n=\text{rank } \mathfrak{g}}$ .*
- **Positivity:**  $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i\}$  are represented by positive, essentially self-adjoint (unbounded) operators on  $L^2(\mathbb{R}^N)$ ,  $N := \ell(w_0)$ .
- $\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i$  are expressed in terms of Laurent polynomials of  $\{e^{\pi b u_k}, e^{2\pi b p_k}\}_{k=1}^N$  where  $[p_j, u_k] = \frac{\delta_{jk}}{2\pi i}$ .
- *Transcendental relations (compatibility with modular double  $b \longleftrightarrow b^{-1}$ ) (Langland’s duality).*

Can be understood in terms of quantum cluster algebra!

# Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations  $\mathcal{P}_{\lambda}$  by positive operators on Hilbert space.
- = “Quantization of principal series representations”.
- Generalization of Teschner’s representations of  $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ .

## Theorem (I. (2012))

*There exists a family of irreducible representations  $\mathcal{P}_{\lambda}$  of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ :*

- *Parametrized by  $\lambda \in \mathbb{R}_{\geq 0} P^+ \simeq \mathbb{R}_{\geq 0}^{n=\text{rank } \mathfrak{g}}$ .*
- ***Positivity:**  $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i\}$  are represented by positive, essentially self-adjoint (unbounded) operators on  $L^2(\mathbb{R}^N)$ ,  $N := \ell(w_0)$ .*
- *$\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i$  are expressed in terms of Laurent polynomials of  $\{e^{\pi b u_k}, e^{2\pi b p_k}\}_{k=1}^N$  where  $[p_j, u_k] = \frac{\delta_{jk}}{2\pi i}$ .*
- *Transcendental relations (compatibility with modular double  $b \longleftrightarrow b^{-1}$ ) (Langland’s duality).*

Can be understood in terms of quantum cluster algebra!

# Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations  $\mathcal{P}_{\lambda}$  by positive operators on Hilbert space.
- = “Quantization of principal series representations”.
- Generalization of Teschner’s representations of  $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ .

## Theorem (I. (2012))

There exists a family of irreducible representations  $\mathcal{P}_{\lambda}$  of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ :

- Parametrized by  $\lambda \in \mathbb{R}_{\geq 0} P^+ \simeq \mathbb{R}_{\geq 0}^{n=\text{rank } \mathfrak{g}}$ .
- **Positivity:**  $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i\}$  are represented by positive, essentially self-adjoint (unbounded) operators on  $L^2(\mathbb{R}^N)$ ,  $N := \ell(w_0)$ .
- $\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i$  are expressed in terms of Laurent polynomials of  $\{e^{\pi b u_k}, e^{2\pi b p_k}\}_{k=1}^N$  where  $[p_j, u_k] = \frac{\delta_{jk}}{2\pi i}$ .  $q = e^{\pi i b^2}$   
 $\tilde{q} = e^{\pi i b^{-2}}$
- **Transcendental relations** (compatibility with modular double  $b \longleftrightarrow b^{-1}$ ) (Langland’s duality).

Can be understood in terms of quantum cluster algebra!

# Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations  $\mathcal{P}_{\lambda}$  by positive operators on Hilbert space.
- = “Quantization of principal series representations”.
- Generalization of Teschner’s representations of  $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ .

## Theorem (I. (2012))

*There exists a family of irreducible representations  $\mathcal{P}_{\lambda}$  of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ :*

- Parametrized by  $\lambda \in \mathbb{R}_{\geq 0} P^+ \simeq \mathbb{R}_{\geq 0}^{n=\text{rank } \mathfrak{g}}$ .
- **Positivity:**  $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i\}$  are represented by positive, essentially self-adjoint (unbounded) operators on  $L^2(\mathbb{R}^N)$ ,  $N := \ell(w_0)$ .
- $\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i$  are expressed in terms of Laurent polynomials of  $\{e^{\pi b u_k}, e^{2\pi b p_k}\}_{k=1}^N$  where  $[p_j, u_k] = \frac{\delta_{jk}}{2\pi i}$ .
- **Transcendental relations** (compatibility with modular double  $b \longleftrightarrow b^{-1}$ ) (Langland’s duality).

Can be understood in terms of quantum cluster algebra!

# Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations  $\mathcal{P}_{\lambda}$  by positive operators on Hilbert space.
- = “Quantization of principal series representations”.
- Generalization of Teschner’s representations of  $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ .

## Theorem (I. (2012))

*There exists a family of irreducible representations  $\mathcal{P}_{\lambda}$  of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ :*

- Parametrized by  $\lambda \in \mathbb{R}_{\geq 0} P^+ \simeq \mathbb{R}_{\geq 0}^{n=\text{rank } \mathfrak{g}}$ .
- **Positivity:**  $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i\}$  are represented by positive, essentially self-adjoint (unbounded) operators on  $L^2(\mathbb{R}^N)$ ,  $N := \ell(w_0)$ .
- $\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i$  are expressed in terms of Laurent polynomials of  $\{e^{\pi b u_k}, e^{2\pi b p_k}\}_{k=1}^N$  where  $[p_j, u_k] = \frac{\delta_{jk}}{2\pi i}$ .
- **Transcendental relations** (compatibility with modular double  $b \longleftrightarrow b^{-1}$ ) (Langland’s duality).

Can be understood in terms of quantum cluster algebra!

# Quantum Cluster Variety

# Quantum Torus Algebra

“Quantization of cluster  $\mathcal{X}$  variety” [Fock-Goncharov]

## Definition

Seed  $\mathbf{Q} = (Q, Q_0, B, D)$ :

- $Q$  = nodes (finite set),
- $Q_0 \subset Q$  = frozen nodes,
- $B = (\varepsilon_{ij})$  exchange matrix (skew-symmetrizable,  $\frac{1}{2}\mathbb{Z}$ -valued),
- $D = (d_i)$  multipliers (in  $\mathbb{Q}$ ).  $d := \min d_i$ .

Let  $W := DB$  such that  $w_{ij} = d_i \varepsilon_{ij} = -w_{ji}$ .

The quantum torus algebra  $\mathcal{X}_q^{\mathbf{Q}}$  is generated by  $\langle X_i \rangle_{i \in Q}$  over  $\mathbb{C}[q^{\pm d}]$  with

$$X_i X_j = q^{-2w_{ij}} X_j X_i.$$

$X_i$  = quantum cluster variables.



# Quantum Torus Algebra

“Quantization of cluster  $\mathcal{X}$  variety” [Fock-Goncharov]

## Definition

Seed  $\mathbf{Q} = (Q, Q_0, B, D)$ :

- $Q$  = nodes (finite set),
- $Q_0 \subset Q$  = frozen nodes,
- $B = (\varepsilon_{ij})$  exchange matrix (skew-symmetrizable,  $\frac{1}{2}\mathbb{Z}$ -valued),
- $D = (d_i)$  multipliers (in  $\mathbb{Q}$ ).  $d := \min d_i$ .

Let  $W := DB$  such that  $w_{ij} = d_i \varepsilon_{ij} = -w_{ji}$ .

The quantum torus algebra  $\mathcal{X}_q^{\mathbf{Q}}$  is generated by  $\langle X_i \rangle_{i \in Q}$  over  $\mathbb{C}[q^{\pm d}]$  with

$$X_i X_j = q^{-2w_{ij}} X_j X_i.$$

$X_i$  = quantum cluster variables.

# Quantum Torus Algebra

“Quantization of cluster  $\mathcal{X}$  variety” [Fock-Goncharov]

## Definition

Seed  $\mathbf{Q} = (Q, Q_0, B, D)$ :

- $Q$  = nodes (finite set),
- $Q_0 \subset Q$  = frozen nodes,
- $B = (\varepsilon_{ij})$  exchange matrix (skew-symmetrizable,  $\frac{1}{2}\mathbb{Z}$ -valued),
- $D = (d_i)$  multipliers (in  $\mathbb{Q}$ ).  $d := \min d_i$ .

Let  $W := DB$  such that  $w_{ij} = d_i \varepsilon_{ij} = -w_{ji}$ .

The quantum torus algebra  $\mathcal{X}_q^{\mathbf{Q}}$  is generated by  $\langle X_i \rangle_{i \in Q}$  over  $\mathbb{C}[q^{\pm d}]$  with

$$X_i X_j = q^{-2w_{ij}} X_j X_i.$$

$X_i$  = quantum cluster variables.

# Quantum Torus Algebra

“Quantization of cluster  $\mathcal{X}$  variety” [Fock-Goncharov]

## Definition

Seed  $\mathbf{Q} = (Q, Q_0, B, D)$ :

- $\Lambda_{\mathbf{Q}} = \mathbb{Z}$ -Lattice with basis  $\{e_i\}_{i \in Q}$ ,
- $(-, -)$  skew-symmetric form,  $(e_i, e_j) := w_{ij}$ .

The quantum torus algebra  $\mathcal{X}_q^{\mathbf{Q}}$  is generated by  $\langle X_{\lambda} \rangle_{\lambda \in \Lambda_{\mathbf{Q}}}$  over  $\mathbb{C}[q^{\pm d}]$  with

$$X_{\lambda+\mu} = q^{(\lambda, \mu)} X_{\lambda} X_{\mu}.$$

We use notation

$$\begin{aligned} X_i &:= X_{e_i} \\ X_{i_1, i_2, \dots, i_k} &:= X_{e_{i_1} + e_{i_2} + \dots + e_{i_k}}. \end{aligned}$$

# Quantum Torus Algebra

“Quantization of cluster  $\mathcal{X}$  variety” [Fock-Goncharov]

## Definition

Seed  $\mathbf{Q} = (Q, Q_0, B, D)$ :

- $\Lambda_{\mathbf{Q}} = \mathbb{Z}$ -Lattice with basis  $\{e_i\}_{i \in Q}$ ,
- $(-, -)$  skew-symmetric form,  $(e_i, e_j) := w_{ij}$ .

The **quantum torus algebra**  $\mathcal{X}_q^{\mathbf{Q}}$  is generated by  $\langle X_{\lambda} \rangle_{\lambda \in \Lambda_{\mathbf{Q}}}$  over  $\mathbb{C}[q^{\pm d}]$  with

$$X_{\lambda+\mu} = q^{(\lambda, \mu)} X_{\lambda} X_{\mu}.$$

We use notation

$$\begin{aligned} X_i &:= X_{e_i} \\ X_{i_1, i_2, \dots, i_k} &:= X_{e_{i_1} + e_{i_2} + \dots + e_{i_k}}. \end{aligned}$$

$$X_1 X_2 = q^{-2} X_2 X_1$$

$$\begin{aligned} \checkmark \quad X_{1,2} &= X_{e_1 + e_2} \\ &:= q X_1 X_2 \end{aligned}$$

# Quiver

Cluster seed  $\mathbf{Q} \leadsto$  quiver with adjacency matrix  $(c_{ij})$  where

$$c_{ij} := \begin{cases} \varepsilon_{ij} d_i d_j^{-1} & \text{if } d_j > d_i, \\ \varepsilon_{ij} & \text{otherwise.} \end{cases}$$

An arrow



represents the algebraic relation

$$X_i X_j = \begin{cases} q_i^{-2} X_j X_i & \text{if } d_i \geq d_j, \\ q_j^{-2} X_j X_i & \text{if } d_j \geq d_i. \end{cases}$$


( $\square \dashrightarrow \square$  between frozen nodes represents half the weight.)

# Quiver

Cluster seed  $\mathbf{Q} \rightsquigarrow$  quiver with adjacency matrix  $(c_{ij})$  where

$$c_{ij} := \begin{cases} \varepsilon_{ij} d_i d_j^{-1} & \text{if } d_j > d_i, \\ \varepsilon_{ij} & \text{otherwise.} \end{cases}$$

An arrow



represents the algebraic relation


$$X_i X_j = \begin{cases} q_i^{-2} X_j X_i & \text{if } d_i \geq d_j, \\ q_j^{-2} X_j X_i & \text{if } d_j \geq d_i. \end{cases}$$

( $\square \dashrightarrow \square$  between **frozen nodes** represents half the weight.)

# Quiver

We also use **thickness** to indicate commutation relations w.r.t.  $q$ , but thickness is **not** part of the data of  $\mathbf{Q}$ .

For example,



represents the relations

$$X_1 X_2 = q^{-2} X_2 X_1$$

$$X_2 X_3 = q_3^{-2} X_3 X_2 = q^{-4} X_3 X_2$$


$$X_1 X_3 = X_3 X_1$$

Note: we can recover  $\mathbf{Q}$  from the quiver and multipliers.

# Quiver

We also use **thickness** to indicate commutation relations w.r.t.  $q$ , but thickness is **not** part of the data of  $\mathbf{Q}$ .

For example,



represents the relations

$$X_1 X_2 = q^{-2} X_2 X_1$$

$$X_2 X_3 = q_3^{-2} X_3 X_2 = q^{-4} X_3 X_2$$

$$X_1 X_3 = X_3 X_1$$


Note: we can recover  $\mathbf{Q}$  from the quiver and multipliers.



# Quiver

We also use **thickness** to indicate commutation relations w.r.t.  $q$ , but thickness is **not** part of the data of  $\mathbf{Q}$ .

For example,



represents the relations

$$X_1 X_2 = q^{-2} X_2 X_1$$

$$X_2 X_3 = \textcolor{blue}{q_3}^{-2} X_3 X_2 = q^{-4} X_3 X_2$$

$$X_1 X_3 = X_3 X_1$$

Note: we can recover  $\mathbf{Q}$  from the quiver and multipliers.

# Polarization of $\mathcal{X}_q^{\mathbf{Q}}$

Recall  $q = e^{\pi i b^2}$  such that  $|q| = 1$ .

## Definition

A **polarization** of  $\mathcal{X}_q^{\mathbf{Q}}$  is a choice of representation of the cluster variables  $X_k \in \mathcal{X}_q^{\mathbf{Q}}$  of the form  $X_k = e^{2\pi b x_k}$  such that

- $x_k$  is a  $\mathbb{Q}$ -linear combination of  $u_i, p_i = \frac{1}{2\pi i} \frac{\partial}{\partial u_i}$  and scalars  $\lambda_i \in \mathbb{R}$ .
- $x_k$  satisfies the Heisenberg algebra relations

$$[x_j, x_k] = \frac{1}{2\pi i} \varepsilon_{jk},$$

$X_k$  acts on some Hilbert space  $\mathcal{H}_{\mathbf{Q}} \simeq L^2(\mathbb{R}^M, du_i)$  as positive operators.

## Remark

Modular double  $\tilde{X}_k := X^{\frac{1}{b^2}}$  acts by  $e^{2\pi b^{-1} x_k}$  on  $\mathcal{H}_{\mathbf{Q}}$ .

# Polarization of $\mathcal{X}_q^{\mathbf{Q}}$

Recall  $q = e^{\pi i b^2}$  such that  $|q| = 1$ .

$$\text{Dom}(e^{2\pi i b u}) = \{f : e^{2\pi i b u} f \in L^2\}$$

## Definition

A **polarization** of  $\mathcal{X}_q^{\mathbf{Q}}$  is a choice of representation of the cluster variables  $X_k \in \mathcal{X}_q^{\mathbf{Q}}$  of the form  $X_k = e^{2\pi i b x_k}$  such that

- $x_k$  is a  $\mathbb{Q}$ -linear combination of  $u_i$ ,  $p_i = \frac{1}{2\pi i} \frac{\partial}{\partial u_i}$  and scalars  $\lambda_i \in \mathbb{R}$ .
- $x_k$  satisfies the Heisenberg algebra relations

$$[x_j, x_k] = \frac{1}{2\pi i} \varepsilon_{jk},$$

$X_k$  acts on some Hilbert space  $\mathcal{H}_{\mathbf{Q}} \simeq L^2(\mathbb{R}^M, du_i)$  as positive operators.

## Remark

**Modular double**  $\tilde{X}_k := X^{\frac{1}{b^2}}$  acts by  $e^{2\pi i b^{-1} x_k}$  on  $\mathcal{H}_{\mathbf{Q}}$ .

# Polarization of $\mathcal{X}_q^Q$

## Example

For  $X_1 X_2 = q^2 X_2 X_1$ , we have

$$X_1 = e^{2\pi bu}, \quad X_2 = e^{2\pi bp}$$

acting on  $L^2(\mathbb{R}, du)$  as positive operators.

## Definition

- The center of  $\mathcal{X}_q^Q$  is generated by  $|Q| - \text{rank}(B)$  central monomials.
- A polarization is irreducible if every central monomial acts by a scalar.

## Proposition

*Any irreducible polarization with the same action of the central monomials are unitarily equivalent.*

# Polarization of $\mathcal{X}_q^Q$

## Example

For  $X_1 X_2 = q^2 X_2 X_1$ , we have

$$X_1 = e^{2\pi bu}, \quad X_2 = e^{2\pi bp}$$

acting on  $L^2(\mathbb{R}, du)$  as positive operators.

## Definition

- The center of  $\mathcal{X}_q^Q$  is generated by  $|Q| - \text{rank}(B)$  central monomials.
- A polarization is irreducible if every central monomial acts by a scalar.

## Proposition

*Any irreducible polarization with the same action of the central monomials are unitarily equivalent.*

# Polarization of $\mathcal{X}_q^Q$

## Example

For  $X_1 X_2 = q^2 X_2 X_1$ , we have

$$X_1 = e^{2\pi b u}, \quad X_2 = e^{2\pi b p}$$

acting on  $L^2(\mathbb{R}, du)$  as positive operators.

## Definition

- The center of  $\mathcal{X}_q^Q$  is generated by  $|Q| - \text{rank}(B)$  **central monomials**.
- A polarization is **irreducible** if every central monomial acts by a scalar.

## Proposition

*Any irreducible polarization with the same action of the central monomials are unitarily equivalent.*

# Quantum Cluster Mutations

$\mathbf{T}_q^{\mathbf{Q}}$  := (non-commutative) field of fractions of  $\mathcal{X}_q^{\mathbf{Q}}$ .

Usual cluster mutation  $\mu_k$  induces  $\mu_k^q : \mathbf{T}_q^{\mathbf{Q}'} \longrightarrow \mathbf{T}_q^{\mathbf{Q}}$ :

$$\mu_k^q(X'_i) := \begin{cases} X_k^{-1} & i = k, \\ X_i \prod_{r=1}^{|\varepsilon_{ki}|} (1 + q_i^{2r-1} X_k) & i \neq k, \varepsilon_{ki} < 0, \\ X_i \prod_{r=1}^{\varepsilon_{ki}} (1 + q_i^{2r-1} X_k^{-1})^{-1} & i \neq k, \varepsilon_{ki} > 0. \end{cases}$$

Can be rewritten as

$$\mu_k^q = \mu_k^{\#} \circ \mu_k^m,$$

$$\mu_k^m(X'_i) := \begin{cases} X_k^{-1} & i = k, \\ X_i & i \neq k, \varepsilon_{ki} < 0, \\ q_i^{\varepsilon_{ik} \varepsilon_{ki}} X_i X_k^{\varepsilon_{ik}} & i \neq k, \varepsilon_{ki} > 0. \end{cases}$$

$$\mu_k^{\#} := \text{Ad}_{g_b^*}(X_k)$$

$g_b(X)$  = quantum dilogarithm

# Quantum Cluster Mutations

$\mathbf{T}_q^{\mathbf{Q}}$  := (non-commutative) field of fractions of  $\mathcal{X}_q^{\mathbf{Q}}$ .

Usual cluster mutation  $\mu_k$  induces  $\mu_k^q : \mathbf{T}_q^{\mathbf{Q}'} \longrightarrow \mathbf{T}_q^{\mathbf{Q}}$ :

$$\mu_k^q(X'_i) := \begin{cases} X_k^{-1} & i = k, \\ X_i \prod_{r=1}^{|\varepsilon_{ki}|} (1 + q_i^{2r-1} X_k) & i \neq k, \varepsilon_{ki} < 0, \\ X_i \prod_{r=1}^{\varepsilon_{ki}} (1 + q_i^{2r-1} X_k^{-1})^{-1} & i \neq k, \varepsilon_{ki} > 0. \end{cases}$$

Can be rewritten as

$$\mu_k^q = \underbrace{\mu_k^{\#}} \circ \underbrace{\mu_k^m},$$

$$\mu_k^m(X'_i) := \begin{cases} X_k^{-1} & i = k, \\ X_i & i \neq k, \varepsilon_{ki} < 0, \\ q_i^{\varepsilon_{ik}\varepsilon_{ki}} X_i X_k^{\varepsilon_{ik}} & i \neq k, \varepsilon_{ki} > 0. \end{cases}$$

$$\mu_k^{\#} := Ad_{g_b^*}(X_k)$$

$g_b(X)$  = quantum dilogarithm



# Quantum Cluster Mutations

## Proposition

- If  $\pi$  is a polarization of  $\mathcal{X}_q^{\mathbf{Q}}$  on  $\mathcal{H}_{\mathbf{Q}}$ , then

$$\pi'(X_i) := \pi(\mu_k^m(X_i))$$

is a polarization of  $\mathcal{X}_q^{\mathbf{Q}'}$ .

- Cluster mutations  $\longleftrightarrow$  unitary transformations  $\mathcal{H}_{\mathbf{Q}} \simeq \mathcal{H}_{\mathbf{Q}'}$ .

## Definition

$\mathcal{O}_q(\mathcal{X})$  denotes the quantum upper cluster algebra of the cluster variety  $\mathcal{X}$ .

The elements of  $\mathcal{O}_q(\mathcal{X})$  consists of all elements  $f \in \mathcal{X}_q^{\mathbf{Q}}$  which remain Laurent polynomials over  $\mathbb{Z}[q^{\pm d}]$  under any quantum cluster mutations.

We call elements  $f \in \mathcal{O}_q(\mathcal{X})$  universally Laurent polynomials.

Polarization of  $\mathcal{O}_q(\mathcal{X})$ =polarization of  $\mathcal{X}_q^{\mathbf{Q}}$  for every  $\mathbf{Q}$  related by above rule.

# Quantum Cluster Mutations

## Proposition

- If  $\pi$  is a polarization of  $\mathcal{X}_q^{\mathbf{Q}}$  on  $\mathcal{H}_{\mathbf{Q}}$ , then

$$\pi'(X_i) := \pi(\mu_k^m(X_i))$$

is a polarization of  $\mathcal{X}_q^{\mathbf{Q}'}$ .

- Cluster mutations  $\longleftrightarrow$  unitary transformations  $\mathcal{H}_{\mathbf{Q}} \simeq \mathcal{H}_{\mathbf{Q}'}$ .

## Definition

$\mathcal{O}_q(\mathcal{X})$  denotes the quantum upper cluster algebra of the cluster variety  $\mathcal{X}$ .

The elements of  $\mathcal{O}_q(\mathcal{X})$  consists of all elements  $f \in \mathcal{X}_q^{\mathbf{Q}}$  which remain Laurent polynomials over  $\mathbb{Z}[q^{\pm d}]$  under any quantum cluster mutations.

We call elements  $f \in \mathcal{O}_q(\mathcal{X})$  **universally Laurent polynomials**.

**Polarization** of  $\mathcal{O}_q(\mathcal{X})$  = polarization of  $\mathcal{X}_q^{\mathbf{Q}}$  for every  $\mathbf{Q}$  related by above rule.

# Quantum Cluster Mutations

## Proposition

- If  $\pi$  is a polarization of  $\mathcal{X}_q^{\mathbf{Q}}$  on  $\mathcal{H}_{\mathbf{Q}}$ , then

$$\pi'(X_i) := \pi(\mu_k^m(X_i))$$

is a polarization of  $\mathcal{X}_q^{\mathbf{Q}'}$ .

- Cluster mutations  $\longleftrightarrow$  unitary transformations  $\mathcal{H}_{\mathbf{Q}} \simeq \mathcal{H}_{\mathbf{Q}'}$ .

## Definition

$\mathcal{O}_q(\mathcal{X})$  denotes the **quantum upper cluster algebra** of the cluster variety  $\mathcal{X}$ .

The elements of  $\mathcal{O}_q(\mathcal{X})$  consists of all elements  $f \in \mathcal{X}_q^{\mathbf{Q}}$  which remain Laurent polynomials over  $\mathbb{Z}[q^{\pm d}]$  under any quantum cluster mutations.

We call elements  $f \in \mathcal{O}_q(\mathcal{X})$  **universally Laurent polynomials**.

**Polarization** of  $\mathcal{O}_q(\mathcal{X})$  = polarization of  $\mathcal{X}_q^{\mathbf{Q}}$  for every  $\mathbf{Q}$  related by above rule.

# Quantum Cluster Mutations

## Proposition

- If  $\pi$  is a polarization of  $\mathcal{X}_q^{\mathbf{Q}}$  on  $\mathcal{H}_{\mathbf{Q}}$ , then

$$\pi'(X_i) := \pi(\mu_k^m(X_i))$$

is a polarization of  $\mathcal{X}_q^{\mathbf{Q}'}$ .

- Cluster mutations  $\longleftrightarrow$  unitary transformations  $\mathcal{H}_{\mathbf{Q}} \simeq \mathcal{H}_{\mathbf{Q}'}$ .

## Definition

$\mathcal{O}_q(\mathcal{X})$  denotes the **quantum upper cluster algebra** of the cluster variety  $\mathcal{X}$ .

The elements of  $\mathcal{O}_q(\mathcal{X})$  consists of all elements  $f \in \mathcal{X}_q^{\mathbf{Q}}$  which remain Laurent polynomials over  $\mathbb{Z}[q^{\pm d}]$  under any quantum cluster mutations.

We call elements  $f \in \mathcal{O}_q(\mathcal{X})$  **universally Laurent polynomials**.


**Polarization** of  $\mathcal{O}_q(\mathcal{X})$ =polarization of  $\mathcal{X}_q^{\mathbf{Q}}$  for every  $\mathbf{Q}$  related by above rule.

# Quantum Cluster Variety

$S$ =Riemann surface with marked points on  $\partial S$  and punctures.

Fock-Goncharov's  $\mathcal{X}_{G,S}$ -space= “(framed) local  $G$ -system”

- $\mathcal{X}_{G,S}$  has Poisson cluster  $\mathcal{X}$  variety structure  $\leadsto$  quantization  $\mathcal{X}_{G,S}^q$
- To each triangle of **ideal triangulation** of  $S$ , assign a **basic quiver**.
- $G = PGL_{n+1}$ : “ $n$ -triangulation”




(Full generality: [I. (2016)], [Le (2016)], [Goncharov-Shen (2019)])  
 For any longest word  $w_0 \in W \leadsto \mathbf{Q} \leadsto \mathcal{X}_q^{\mathbf{Q}}$ .

# Quantum Cluster Variety

$S$ =Riemann surface with marked points on  $\partial S$  and punctures.

Fock-Goncharov's  $\mathcal{X}_{G,S}$ -space= “(framed) local  $G$ -system”

- $\mathcal{X}_{G,S}$  has Poisson cluster  $\mathcal{X}$  variety structure  $\leadsto$  quantization  $\mathcal{X}_{G,S}^q$
- To each triangle of ideal triangulation of  $S$ , assign a basic quiver.
- $G = PGL_{n+1}$ : “ $n$ -triangulation”




(Full generality: [I. (2016)], [Le (2016)], [Goncharov-Shen (2019)])  
 For any longest word  $w_0 \in W \leadsto \mathbf{Q} \leadsto \mathcal{X}_q^{\mathbf{Q}}$ .

# Quantum Cluster Variety

$S$ =Riemann surface with marked points on  $\partial S$  and punctures.

Fock-Goncharov's  $\mathcal{X}_{G,S}$ -space= “(framed) local  $G$ -system”

- $\mathcal{X}_{G,S}$  has Poisson cluster  $\mathcal{X}$  variety structure  $\leadsto$  quantization  $\mathcal{X}_{G,S}^q$
- To each triangle of **ideal triangulation** of  $S$ , assign a **basic quiver**.
- $G = PGL_{n+1}$ : “ $n$ -triangulation”




(Full generality: [I. (2016)], [Le (2016)], [Goncharov-Shen (2019)])  
 For any longest word  $w_0 \in W \leadsto \mathbf{Q} \leadsto \mathcal{X}_q^{\mathbf{Q}}$ .

# Quantum Cluster Variety

$S$ =Riemann surface with marked points on  $\partial S$  and punctures.

Fock-Goncharov's  $\mathcal{X}_{G,S}$ -space= “(framed) local  $G$ -system”

- $\mathcal{X}_{G,S}$  has Poisson cluster  $\mathcal{X}$  variety structure  $\leadsto$  quantization  $\mathcal{X}_{G,S}^q$
- To each triangle of **ideal triangulation** of  $S$ , assign a **basic quiver**.
- $G = PGL_{n+1}$ : “ $n$ -triangulation”



(Full generality: [I. (2016)], [Le (2016)], [Goncharov-Shen (2019)])  
 For any longest word  $w_0 \in W \leadsto \mathbf{Q} \leadsto \mathcal{X}_q^{\mathbf{Q}}$ .




# Quantum Cluster Variety

$S$ =Riemann surface with marked points on  $\partial S$  and punctures.

Fock-Goncharov's  $\mathcal{X}_{G,S}$ -space= “(framed) local  $G$ -system”

- $\mathcal{X}_{G,S}$  has Poisson cluster  $\mathcal{X}$  variety structure  $\leadsto$  quantization  $\mathcal{X}_{G,S}^q$
- To each triangle of **ideal triangulation** of  $S$ , assign a **basic quiver**.
- $G = PGL_{n+1}$ : “ $n$ -triangulation”




(Full generality: [I. (2016)], [Le (2016)], [Goncharov-Shen (2019)])  
 For any longest word  $w_0 \in W \leadsto \mathbf{Q} \leadsto \mathcal{X}_q^{\mathbf{Q}}$ .


# Quantum Cluster Variety

We can flip the triangulation with  $\binom{n+2}{3}$  quiver mutations.

[Fock-Goncharov]




# Example: Type $A_n$ Case



Amalgamation of 2 quivers

# Example: Type $A_n$ Case




$$\mathbf{D}_{\mathfrak{sl}_{n+1}}\text{-quiver} \leadsto \mathcal{X}_q^{std} \quad [\text{Schrader-Shapiro}]$$

$$\iota : \mathcal{D}_q(\mathfrak{sl}_{n+1}) \hookrightarrow \mathcal{X}_q^{std}$$

$$\mathcal{U}_q(\mathfrak{sl}_{n+1}) \hookrightarrow \mathcal{X}_q^{std} / \langle \iota(\mathbf{K}_i \mathbf{K}'_i) = 1 \rangle$$

### Example: Type $A_n$ Case



Embedding of  $F_i \in \mathfrak{D}_{\mathfrak{sl}_4} \hookrightarrow \mathcal{X}_q^{std}$

$\swarrow q X_1 X_2 \quad \swarrow q^2 X_1 X_2 X_3$


$$\mathbf{f}_1 = X_1 + X_{1,2} + X_{1,2,3} + X_{1,2,3,4} + X_{1,2,3,4,5} + X_{1,2,3,4,5,6}$$

$$\mathbf{f}_2 = X_8 + X_{8,9} + X_{8,9,10} + X_{8,9,10,11}$$

$$\mathbf{f}_3 = X_{13} + X_{13,14}$$

$$\mathbf{K}'_1 = X_{1,2,3,4,5,6,7} \quad \mathbf{K}'_2 = X_{8,9,10,11,12} \quad \mathbf{K}'_3 = X_{13,14,15}$$

# Example: Type $A_n$ Case



Embedding of  $E_i \in \mathcal{D}_{\mathfrak{sl}_4} \hookrightarrow \mathcal{X}_q^{std}$

$$\mathbf{e}_1 = X_7 + X_{7,16}$$

$$\mathbf{e}_2 = X_{12} + X_{12,6} + X_{12,6,17} + X_{12,6,17,2}$$

$$\mathbf{e}_3 = X_{15} + X_{15,11} + X_{15,11,5} + X_{15,11,5,18} + X_{15,11,5,18,3} + X_{15,11,5,18,3,9}$$

$$\mathbf{K}_1 = X_{7,16,1} \quad \mathbf{K}_2 = X_{12,6,17,2,8} \quad \mathbf{K}_3 = X_{15,11,5,18,3,9,13}$$

# Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

*✓ type  $\mathfrak{g}$ .*

Theorem (Schrader-Shapiro, I. (2016))

Given a longest reduced word  $\mathbf{i}_0$

- There exists a quiver  $\mathbf{D}(\mathbf{i}_0)$  associated to  and an embedding

$$\mathcal{D}_q(\mathfrak{g}) \hookrightarrow \mathcal{X}_q^{std}.$$

- We recover the positive representations  $\mathcal{P}_\lambda$  through a polarization of  $\mathcal{X}_q^{std}$ .

Theorem (I. (2016), Goncharov-Shen (2019))

- $\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i, \mathbf{K}'_i$  are universally Laurent polynomials in  $\mathcal{O}_q(\mathcal{X}_q^{std})$ .
- $\mathcal{D}_q(\mathfrak{g}) \hookrightarrow \mathcal{X}_q^{std}$  can be described by paths on the quiver  $\mathbf{D}(\mathbf{i}_0)$ .
- $\mathbf{f}_i$  paths are simple  $\leadsto$  Feigin's homomorphism  $\mathcal{U}_q^-(\mathfrak{g}) \hookrightarrow \mathcal{X}_q$ .

# Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Theorem (*Schrader-Shapiro, I. (2016)*)

Given a longest reduced word  $\mathbf{i}_0$

- There exists a quiver  $\mathbf{D}(\mathbf{i}_0)$  associated to  and an embedding

$$\mathcal{D}_q(\mathfrak{g}) \hookrightarrow \mathcal{X}_q^{std}.$$

- We recover the positive representations  $\mathcal{P}_\lambda$  through a polarization of  $\mathcal{X}_q^{std}$ .

Theorem (*I. (2016), Goncharov-Shen (2019)*)

- $\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i, \mathbf{K}'_i$  are universally Laurent polynomials in  $\mathcal{O}_q(\mathcal{X}^{std})$ .
- $\mathcal{D}_q(\mathfrak{g}) \hookrightarrow \mathcal{X}_q^{std}$  can be described by paths on the quiver  $\mathbf{D}(\mathbf{i}_0)$ .
- $\mathbf{f}_i$  paths are simple  $\leadsto$  Feigin's homomorphism  $\mathcal{U}_q^-(\mathfrak{g}) \hookrightarrow \mathcal{X}_q$ .



# Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

## Theorem (Schrader-Shapiro, I. (2016))

Given a longest reduced word  $\mathbf{i}_0$

- There exists a quiver  $\mathbf{D}(\mathbf{i}_0)$  associated to  and an embedding

$$\mathcal{D}_q(\mathfrak{g}) \hookrightarrow \mathcal{X}_q^{std}.$$

- We recover the positive representations  $\mathcal{P}_\lambda$  through a polarization of  $\mathcal{X}_q^{std}$ .

## Theorem (I. (2016), Goncharov-Shen (2019))

- $\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i, \mathbf{K}'_i$  are universally Laurent polynomials in  $\mathcal{O}_q(\mathcal{X}^{std})$ .
- $\mathcal{D}_q(\mathfrak{g}) \hookrightarrow \mathcal{X}_q^{std}$  can be described by paths on the quiver  $\mathbf{D}(\mathbf{i}_0)$ .
- $\mathbf{f}_i$  paths are simple  $\leadsto$  Feigin's homomorphism  $\mathcal{U}_q^-(\mathfrak{g}) \hookrightarrow \mathcal{X}_q$ .

# Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

## Theorem (Schrader-Shapiro, I. (2016))

Given a longest reduced word  $\mathbf{i}_0$

- There exists a quiver  $\mathbf{D}(\mathbf{i}_0)$  associated to  and an embedding

$$\mathcal{D}_q(\mathfrak{g}) \hookrightarrow \mathcal{X}_q^{\text{std}}.$$

- We recover the positive representations  $\mathcal{P}_\lambda$  through a polarization of  $\mathcal{X}_q^{\text{std}}$ .

## Theorem (I. (2016), Goncharov-Shen (2019))

- $\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i, \mathbf{K}'_i$  are universally Laurent polynomials in  $\mathcal{O}_q(\mathcal{X}^{\text{std}})$ .
- $\mathcal{D}_q(\mathfrak{g}) \hookrightarrow \mathcal{X}_q^{\text{std}}$  can be described by paths on the quiver  $\mathbf{D}(\mathbf{i}_0)$ .
- $\mathbf{f}_i$  paths are simple  $\leadsto$  Feigin's homomorphism  $\mathcal{U}_q(\mathfrak{g}) \hookrightarrow \mathcal{X}_q$ .

$\mathcal{U}_q(\mathfrak{b}^-)$

# Regular positive representations

## Definition

A representation  $\mathcal{P}$  of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$  is called a *regular positive representation* if  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$  acts by a polarization of some quantum upper cluster algebra  $\mathcal{O}_q(\mathcal{X})$ .

In other words,

- There exists a homomorphism  $\iota : \mathcal{D}_q(\mathfrak{g}) \longrightarrow \mathcal{O}_q(\mathcal{X})$ .
- i.e. The image of  $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i, \mathbf{K}'_i\}$  are universally Laurent.  
( $\iff$ ?) The image of  $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i, \mathbf{K}'_i\}$  are universally **polynomial**. [I.-Ye 2023]
- The representation  $\mathcal{P}$  is obtained via an irreducible polarization of  $\mathcal{O}_q(\mathcal{X})$  (where the central monomials  $\iota(\mathbf{K}_i \mathbf{K}'_i)$  act as 1.)

**Main Question:** Classify all irreducible regular positive representations!

- If  $\mathcal{P}$  is irreducible, the Casimir elements act as multiplications by real scalars.
- Which eigenvalues of the Casimirs arise from the regular positive representations?
- Do the actions of the Casimirs uniquely determine a regular positive representation?

# Regular positive representations

## Definition

A representation  $\mathcal{P}$  of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$  is called a *regular positive representation* if  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$  acts by a polarization of some quantum upper cluster algebra  $\mathcal{O}_q(\mathcal{X})$ .

In other words,

- There exists a homomorphism  $\iota : \mathcal{D}_q(\mathfrak{g}) \longrightarrow \mathcal{O}_q(\mathcal{X})$ .
- i.e. The image of  $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i, \mathbf{K}'_i\}$  are universally Laurent.  
( $\iff$ ?) The image of  $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i, \mathbf{K}'_i\}$  are universally **polynomial**. [I.-Ye 2023]
- The representation  $\mathcal{P}$  is obtained via an irreducible polarization of  $\mathcal{O}_q(\mathcal{X})$  (where the central monomials  $\iota(\mathbf{K}_i \mathbf{K}'_i)$  act as 1.)

**Main Question:** Classify all irreducible regular positive representations!

- If  $\mathcal{P}$  is irreducible, the Casimir elements act as multiplications by real scalars.
- Which eigenvalues of the Casimirs arise from the regular positive representations?
- Do the actions of the Casimirs uniquely determine a regular positive representation?

# Regular positive representations

## Definition

A representation  $\mathcal{P}$  of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$  is called a *regular positive representation* if  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$  acts by a polarization of some quantum upper cluster algebra  $\mathcal{O}_q(\mathcal{X})$ .

In other words,

- There exists a homomorphism  $\iota : \mathcal{D}_q(\mathfrak{g}) \longrightarrow \mathcal{O}_q(\mathcal{X})$ .
- i.e. The image of  $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i, \mathbf{K}'_i\}$  are universally Laurent.  
( $\iff$ ?) The image of  $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i, \mathbf{K}'_i\}$  are universally **polynomial**. [I.-Ye 2023]
- The representation  $\mathcal{P}$  is obtained via an irreducible polarization of  $\mathcal{O}_q(\mathcal{X})$  (where the central monomials  $\iota(\mathbf{K}_i \mathbf{K}'_i)$  act as 1.)

**Main Question:** Classify all irreducible regular positive representations!

- If  $\mathcal{P}$  is irreducible, the Casimir elements act as multiplications by real scalars.
- Which eigenvalues of the Casimirs arise from the regular positive representations?
- Do the actions of the Casimirs uniquely determine a regular positive representation?

# Parabolic Positive Representations

# Parabolic Positive Representations

The standard positive representation  $\mathcal{P}_\lambda$  is constructed by quantizing the principal series representation on

$$L^2((B_- \setminus G)_{>0}) \quad (\simeq \text{Ind}_{B_-}^G(\mathbb{C}_\lambda))$$

## Definition

*Parabolic Positive Representations*  $\mathcal{P}_\lambda^J$  is a family of positive representations of  $\mathcal{U}_q(\mathfrak{g}_\mathbb{R})$  based on quantizing the parabolic induction representations on  $L^2((P_J \setminus G)_{>0})$ , where  $P_J \subset G$  is a parabolic subgroup indexed by  $J \subset I$ .

The standard representation  $\mathcal{P}_\lambda$  corresponds to  $J = \emptyset$ , i.e.  $P_J = B_-$  the lower Borel subgroup.

# Parabolic Positive Representations

The standard positive representation  $\mathcal{P}_\lambda$  is constructed by quantizing the principal series representation on

$$L^2((B_- \setminus G)_{>0}) \quad (\simeq \text{Ind}_{B_-}^G(\mathbb{C}_\lambda))$$

## Definition

*Parabolic Positive Representations*  $\mathcal{P}_\lambda^J$  is a family of positive representations of  $\mathcal{U}_q(\mathfrak{g}_\mathbb{R})$  based on quantizing the parabolic induction representations on  $L^2((P_J \setminus G)_{>0})$ , where  $P_J \subset G$  is a parabolic subgroup indexed by  $J \subset I$ .

The standard representation  $\mathcal{P}_\lambda$  corresponds to  $J = \emptyset$ , i.e.  $P_J = B_-$  the lower Borel subgroup.



# Parabolic Positive Representations

Parabolic induction  $\longleftrightarrow$  truncating  $\mathbf{i}_J \subset \mathbf{i}_0$  where  $\mathbf{i}_J, \mathbf{i}_0$  are the longest word of the Weyl groups  $W_J \subset W$ .

$$w_0 = w_J \bar{w}$$

$$\bar{w} \longleftrightarrow \bar{\mathbf{i}}$$

## Example

$$W_{\mathfrak{sl}_3} \subset W_{\mathfrak{sl}_4}$$

$$\mathbf{i}_0 = (1, 2, 1, 3, 2, 1)$$

Observe that

$$Q(\mathbf{i}_0) = Q(\mathbf{i}_J) * Q(\bar{\mathbf{i}}).$$

# Parabolic Positive Representations

Parabolic induction  $\longleftrightarrow$  truncating  $\mathbf{i}_J \subset \mathbf{i}_0$  where  $\mathbf{i}_J, \mathbf{i}_0$  are the longest word of the Weyl groups  $W_J \subset W$ .

$$w_0 = w_J \bar{w}$$

$$\bar{w} \longleftrightarrow \bar{\mathbf{i}}$$

## Example

$$W_{\mathfrak{sl}_3} \subset W_{\mathfrak{sl}_4}$$

$$\mathbf{i}_0 = (1, 2, 1, 3, 2, 1)$$

Observe that

$$Q(\mathbf{i}_0) = Q(\mathbf{i}_J) * Q(\bar{\mathbf{i}}).$$

# Parabolic Positive Representations

Parabolic induction  $\longleftrightarrow$  truncating  $\mathbf{i}_J \subset \mathbf{i}_0$  where  $\mathbf{i}_J, \mathbf{i}_0$  are the longest word of the Weyl groups  $W_J \subset W$ .

$$w_0 = w_J \bar{w}$$

$$\bar{w} \longleftrightarrow \bar{\mathbf{i}}$$

## Example


$$W_{\mathfrak{sl}_3} \subset W_{\mathfrak{sl}_4}$$

$$\mathbf{i}_0 = (1, 2, 1, 3, 2, 1)$$

Observe that


$$Q(\mathbf{i}_0) = Q(\mathbf{i}_J) * Q(\bar{\mathbf{i}}).$$

# Example: Type $A_3$ Case



Embedding of  $\mathfrak{D}_{\mathfrak{sl}_4} \hookrightarrow \mathcal{X}_q^{std}$

# Parabolic Positive Representations



$$D(\bar{\mathbf{i}}) := Q(\bar{\mathbf{i}}^{op}) * Q(\bar{\mathbf{i}}), \quad \bar{\mathbf{i}} = (3, 2, 1)$$

$$\mathbf{e}_1 = X_3 + X_{3,0}$$

$$\mathbf{e}_2 = X_6 + X_{6,2}$$

$$\mathbf{e}_3 = X_9 + X_{9,5}$$

$$\mathbf{f}_1 = X_1 + X_{1,2}$$

$$\mathbf{f}_2 = X_4 + X_{4,5}$$

$$\mathbf{f}_3 = X_7 + X_{7,8}$$

$$\mathbf{K}_1 = X_{3,0,1}$$

$$\mathbf{K}_2 = X_{6,2,4}$$

$$\mathbf{K}_3 = X_{9,5,7}$$

$$\mathbf{K}'_1 = X_{1,2,3}$$

$$\mathbf{K}'_2 = X_{4,5,6}$$

$$\mathbf{K}'_3 = X_{7,8,9}$$

$D_i(g) \rightarrow Q(K)$

# Parabolic Positive Representation

Parabolic positive representation is **regular**.

Theorem (I. (2020), I.-Ye (2023))

- There is a homomorphism

$$\mathcal{D}_q(\mathfrak{g}) \longrightarrow \mathcal{O}_q(\mathcal{X}^{\mathbf{D}(\bar{\mathbf{i}})})$$

such that the image of  $\langle \mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i, \mathbf{K}'_i \rangle$  are universally polynomials.

- A polarization of  $\mathcal{O}_q(\mathcal{X}^{\mathbf{D}(\bar{\mathbf{i}})})$  induces a family of irreducible representations  $\mathcal{P}_\lambda^J$  of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$  parametrized by  $\lambda \in \mathbb{R}^{|I \setminus J|}$  as positive self-adjoint operators on  $L^2(\mathbb{R}^{l(\bar{w})})$ .

# Parabolic Positive Representation

Parabolic positive representation is **regular**.


Theorem (I. (2020), I.-Ye (2023))

- *There is a homomorphism*

$$\mathcal{D}_q(\mathfrak{g}) \longrightarrow \mathcal{O}_q(\mathcal{X}^{\mathbf{D}(\bar{\mathbf{i}})})$$


*such that the image of  $\langle \mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i, \mathbf{K}'_i \rangle$  are universally polynomials.*

- *A polarization of  $\mathcal{O}_q(\mathcal{X}^{\mathbf{D}(\bar{\mathbf{i}})})$  induces a family of irreducible representations  $\mathcal{P}_\lambda^J$  of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$  parametrized by  $\lambda \in \mathbb{R}^{|I \setminus J|}$  as positive self-adjoint operators on  $L^2(\mathbb{R}^{l(\bar{w})})$ .*


Example:  $E_6$ 

$$A_1 \subset A_2 \subset A_3 \subset D_4 \subset D_5 \subset E_6$$




Example:  $E_6$ 

$$D_5 \subset E_6$$

Example:  $B_5$ 


$$A_4 \subset B_5$$

Example:  $B_5$ 

$$A_4 \subset B_5$$

Back to  $\mathcal{U}_q(\mathfrak{sl}_2)$

# Cluster Realization of $\mathcal{U}_q(\mathfrak{sl}_2)$



We have  $\iota : \mathcal{D}_q(\mathfrak{sl}_2) \hookrightarrow \mathcal{X}_q^{std}$ :

$$\mathbf{e} \mapsto X_3 + X_{3,4}$$

$$\mathbf{f} \mapsto X_1 + X_{1,2}$$


$$\mathbf{K} \mapsto X_{3,4,1}$$

$$\mathbf{K}' \mapsto X_{1,2,3}$$

*exactly Teschner's rep*

- The Casimir is given by  $\mathbf{C} \mapsto X_{1,3} + X_{1,2,3,4}$ .
- The center is generated by  $\iota(\mathbf{K}\mathbf{K}') = X_{1^2,2,3^2,4}$  and  $Q = X_{2,4}$ .
- We recover the previous  $\mathcal{P}_\lambda$  with a polarization of  $\mathcal{X}_q^{std}$ .

# Cluster Realization of $\mathcal{U}_q(\mathfrak{sl}_2)$



We have  $\iota : \mathcal{D}_q(\mathfrak{sl}_2) \hookrightarrow \mathcal{X}_q^{std}$ :

$$\mathbf{e} \mapsto X_3 + X_{3,4}$$

$$\mathbf{f} \mapsto X_1 + X_{1,2}$$


$$\mathbf{K} \mapsto X_{3,4,1}$$

$$\mathbf{K}' \mapsto X_{1,2,3}$$

- The Casimir is given by  $\mathbf{C} \mapsto X_{1,3} + X_{1,2,3,4}$ .
- The center is generated by  $\iota(\mathbf{K}\mathbf{K}') = X_{1^2,2,3^2,4}$  and  $Q = X_{2,4}$ .
- We recover the previous  $\mathcal{P}_\lambda$  with a polarization of  $\mathcal{X}_q^{std}$ .

# Cluster Realization of $\mathcal{U}_q(\mathfrak{sl}_2)$

Do a mutation at vertex 2:



The vertex 2 and 4 become *symmetric* in  $\mathcal{X}_q^{sym}$ .

$$\mathbf{e} \mapsto X_3 + X_{3,2} + X_{3,4} + X_{3,2,4}$$

$$\mathbf{f} \mapsto X_1$$

$$\mathbf{K} \mapsto X_{3,2,4,1}$$


$$\mathbf{K}' \mapsto X_{1,3}$$

and the Casimir element is given by

$$\begin{aligned} \mathbf{C} &\mapsto X_{1,2,3} + X_{1,3,4} \\ &= X_{1,3}(X_2 + X_4). \end{aligned}$$

# Cluster Realization of $\mathcal{U}_q(\mathfrak{sl}_2)$

Do a mutation at vertex 2:



The vertex 2 and 4 become *symmetric* in  $\mathcal{X}_q^{sym}$ .

$$\mathbf{e} \mapsto X_3 + X_1^{-1}\mathbf{C} + X_{3,2,4}$$

$$\mathbf{f} \mapsto X_1$$

$$\mathbf{K} \mapsto X_{3,2,4,1}$$

$$\mathbf{K}' \mapsto X_{1,3}$$

and the Casimir element is given by

$$\begin{aligned}\mathbf{C} &\mapsto X_{1,2,3} + X_{1,3,4} \\ &= X_{1,3}(X_2 + X_4).\end{aligned}$$




# Cluster Realization of $\mathcal{U}_q(\mathfrak{sl}_2)$

Setting  $\mathbf{C} = 0 \rightsquigarrow$  **folding**, where we identify the vertex 2 and 4:

$$X_{\star} := X_{2,4}$$

and obtain a new *skew-symmetrizable*  $\mathcal{X}_q^0$  with  $d_{\star} := 2$ .



and a homomorphism  $\mathcal{D}_q(\mathfrak{sl}_2) \longrightarrow \mathcal{X}_q^0$ :

$$\mathbf{e} \mapsto X_3 + X_{3,\star}$$

$$\mathbf{f} \mapsto X_1$$

$$\mathbf{K} \mapsto X_{3,\star,1}$$

$$\mathbf{K}' \mapsto X_{1,3}$$


We obtain the **degenerate representation**  $\mathcal{P}^0$  of  $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$  by polarization!

# Cluster Realization of $\mathcal{U}_q(\mathfrak{sl}_2)$

Setting  $\mathbf{C} = 0 \rightsquigarrow$  **folding**, where we identify the vertex 2 and 4:

$$X_{\star} := X_{2,4}$$

and obtain a new *skew-symmetrizable*  $\mathcal{X}_q^0$  with  $d_{\star} := 2$ .



and a homomorphism  $\mathcal{D}_q(\mathfrak{sl}_2) \longrightarrow \mathcal{X}_q^0$ :

$$\mathbf{e} \mapsto X_3 + X_{3,\star}$$

$$\mathbf{f} \mapsto X_1$$

$$\mathbf{K} \mapsto X_{3,\star,1}$$

$$\mathbf{K}' \mapsto X_{1,3}$$

We obtain the **degenerate representation**  $\mathcal{P}^0$  of  $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$  by polarization!

# Positive Representations at Zero Casimir

# Casimir Operators

## Theorem

The center of  $\dot{\mathcal{U}}_q(\mathfrak{g}) := \mathcal{U}_q(\mathfrak{g})[\mathbf{K}_i^{\frac{1}{h}}]$  is spanned by rank  $\mathfrak{g} = n$  *generalized Casimirs*

$$\mathbf{C}_k := (1 \otimes \mathrm{Tr}|_{V_k}^q)(RR_{21}).$$

- $R \in \dot{\mathcal{U}}_q(\mathfrak{g}) \hat{\otimes} \dot{\mathcal{U}}_q(\mathfrak{g}) = \text{universal } R \text{ matrix.}$
- $\{V_k\}_{k=1}^n = \text{fundamental representations of } \mathcal{U}_q(\mathfrak{g}).$
- The *quantum trace*  $\mathrm{Tr}|_V^q$  is given by

$$\mathrm{Tr}|_V^q(x) := \mathrm{Tr}|_V(xu^{-1}), \quad x \in \dot{\mathcal{U}}_q(\mathfrak{g})$$

where  $u \in \dot{\mathcal{U}}_q(\mathfrak{g})$  invertible such that  $\mathrm{Ad}(u) = S^2$ .

To ensure positivity of  $\mathbf{C}_k$  in  $\mathcal{P}_\lambda$ , we choose [L. (2016)]

$$u := \mathbf{K}_{2\rho} \tilde{\mathbf{K}}_{2\rho} \in \dot{\mathcal{U}}_{q\tilde{q}}(\mathfrak{g}_{\mathbb{R}})$$

where  $\rho = \frac{1}{2} \sum_{\alpha \in \Delta_+} \alpha$ .

# Casimir Operators

## Theorem

The center of  $\dot{\mathcal{U}}_q(\mathfrak{g}) := \mathcal{U}_q(\mathfrak{g})[\mathbf{K}_i^{\frac{1}{h}}]$  is spanned by rank  $\mathfrak{g} = n$  *generalized Casimirs*

$$\mathbf{C}_k := (1 \otimes \mathrm{Tr}|_{V_k}^q)(RR_{21}).$$

- $R \in \dot{\mathcal{U}}_q(\mathfrak{g}) \hat{\otimes} \dot{\mathcal{U}}_q(\mathfrak{g}) = \text{universal } R \text{ matrix.}$
- $\{V_k\}_{k=1}^n = \text{fundamental representations of } \mathcal{U}_q(\mathfrak{g}).$
- The *quantum trace*  $\mathrm{Tr}|_V^q$  is given by

$$\mathrm{Tr}|_V^q(x) := \mathrm{Tr}|_V(xu^{-1}), \quad x \in \dot{\mathcal{U}}_q(\mathfrak{g})$$

where  $u \in \dot{\mathcal{U}}_q(\mathfrak{g})$  invertible such that  $\mathrm{Ad}(u) = S^2$ .

To ensure positivity of  $\mathbf{C}_k$  in  $\mathcal{P}_\lambda$ , we choose [I. (2016)]

$$u := \mathbf{K}_{2\rho} \tilde{\mathbf{K}}_{2\rho} \in \dot{\mathcal{U}}_{q\tilde{q}}(\mathfrak{g}_{\mathbb{R}})$$

where  $\rho = \frac{1}{2} \sum_{\alpha \in \Delta_+} \alpha$ .

# Casimir Operators

## Example

In  $\mathcal{U}_q(\mathfrak{sl}_2)$ , the Casimir is

$$C = fe - qK - q^{-1}K^{-1}.$$

## Example

In  $\mathcal{U}_q(\mathfrak{sl}_3)$ , the two generalized Casimirs are

$$C_1 = K(q^{-2}K_1K_2 + K_1^{-1}K_2 + q^2K_1^{-1}K_2^{-1} - q^{-1}K_2e_1f_1 - qK_1^{-1}e_2f_2 + e_{21}f_{12}),$$

$$C_2 = K^{-1}(q^2K_1^{-1}K_2^{-1} + K_1K_2^{-1} + q^{-2}K_1K_2 - qK_2^{-1}e_1f_1 - q^{-1}K_1e_2f_2 + e_{12}f_{21}),$$

where  $K = K_1^{\frac{1}{3}}K_2^{-\frac{1}{3}}$ , and

$$e_{ij} := \frac{q^{\frac{1}{2}}e_j e_i - q^{-\frac{1}{2}}e_i e_j}{q - q^{-1}}, \quad f_{ij} := \frac{q^{\frac{1}{2}}f_j f_i - q^{-\frac{1}{2}}f_i f_j}{q - q^{-1}}$$

are obtained by the *symmetrized Lusztig's transformations*.

# Casimir Operators

## Example

In  $\mathcal{U}_q(\mathfrak{sl}_2)$ , the Casimir is

$$C = \mathbf{f}\mathbf{e} - q\mathbf{K} - q^{-1}\mathbf{K}^{-1}.$$

## Example

In  $\mathcal{U}_q(\mathfrak{sl}_3)$ , the two generalized Casimirs are

$$C_1 = \mathbf{K}(q^{-2}\mathbf{K}_1\mathbf{K}_2 + \mathbf{K}_1^{-1}\mathbf{K}_2 + q^2\mathbf{K}_1^{-1}\mathbf{K}_2^{-1} - q^{-1}\mathbf{K}_2\mathbf{e}_1\mathbf{f}_1 - q\mathbf{K}_1^{-1}\mathbf{e}_2\mathbf{f}_2 + \mathbf{e}_{21}\mathbf{f}_{12}),$$

$$C_2 = \mathbf{K}^{-1}(q^2\mathbf{K}_1^{-1}\mathbf{K}_2^{-1} + \mathbf{K}_1\mathbf{K}_2^{-1} + q^{-2}\mathbf{K}_1\mathbf{K}_2 - q\mathbf{K}_2^{-1}\mathbf{e}_1\mathbf{f}_1 - q^{-1}\mathbf{K}_1\mathbf{e}_2\mathbf{f}_2 + \mathbf{e}_{12}\mathbf{f}_{21}),$$

where  $\mathbf{K} = \mathbf{K}_1^{\frac{1}{3}}\mathbf{K}_2^{-\frac{1}{3}}$ , and

$$\mathbf{e}_{ij} := \frac{q^{\frac{1}{2}}\mathbf{e}_j\mathbf{e}_i - q^{-\frac{1}{2}}\mathbf{e}_i\mathbf{e}_j}{q - q^{-1}}, \quad \mathbf{f}_{ij} := \frac{q^{\frac{1}{2}}\mathbf{f}_j\mathbf{f}_i - q^{-\frac{1}{2}}\mathbf{f}_i\mathbf{f}_j}{q - q^{-1}}$$

are obtained by the *symmetrized Lusztig's transformations*.

# Casimir Operators

## Theorem (I. (2016))

The generalized Casimirs  $\mathbf{C}_k$  act on  $\mathcal{P}_\lambda$  by the scalars

$$\pi_\lambda(\mathbf{C}_k) = \sum_{\mathcal{V} \subset V_k} \exp\left(-4\pi\mu_{\mathcal{V}}(\vec{\lambda}_{\mathfrak{h}})\right),$$

over all weight subspaces  $\mathcal{V} \subset V_k$  with weight  $\mu_{\mathcal{V}} \in \mathfrak{h}_{\mathbb{R}}^*$ , and

$$\vec{\lambda}_{\mathfrak{h}} = \sum_{i \in I} \lambda_i b_i W_i \in \mathfrak{h}_{\mathbb{R}}$$

where  $W_i =$  fundamental coweights dual to  $\alpha_i$ .

## Remark

$\pi_\lambda(\mathbf{C}_k)$  can also be computed by the [Weyl character formula](#).



# Casimir Operators

## Theorem (I. (2016))

The generalized Casimirs  $\mathbf{C}_k$  act on  $\mathcal{P}_\lambda$  by the scalars

$$\pi_\lambda(\mathbf{C}_k) = \sum_{\mathcal{V} \subset V_k} \exp\left(-4\pi\mu_{\mathcal{V}}(\vec{\lambda}_{\mathfrak{h}})\right),$$

over all weight subspaces  $\mathcal{V} \subset V_k$  with weight  $\mu_{\mathcal{V}} \in \mathfrak{h}_{\mathbb{R}}^*$ , and

$$\vec{\lambda}_{\mathfrak{h}} = \sum_{i \in I} \lambda_i b_i W_i \in \mathfrak{h}_{\mathbb{R}}$$

where  $W_i =$  fundamental coweights dual to  $\alpha_i$ .

## Remark

$\pi_\lambda(\mathbf{C}_k)$  can also be computed by the [Weyl character formula](#).

# Casimir Operators

## Proposition

In type  $A_n$ ,  $V_k = \Lambda^k V_1$ , and the actions of  $\mathbf{C}_k$  on  $\mathcal{P}_\lambda$  are given by

$$\pi_\lambda(\mathbf{C}_k) = \mathcal{E}_k(e^{4\pi b \varpi_0}, \dots, e^{4\pi b \varpi_n}).$$

where  $\mathcal{E}_k$  are the *elementary symmetric polynomials* in  $n+1$  variables, and

$$\varpi_i := \frac{1}{n+1} \sum_{k=1}^n k \lambda_k - \sum_{j=n+1-i}^n \lambda_j, \quad i = 0, \dots, n.$$

## Example

In  $\mathcal{U}_q(\mathfrak{sl}_3)$ , (type  $A_2$ ) we have

$$\begin{aligned} \pi_\lambda(\mathbf{C}_1) &= e^{\frac{4}{3}\pi b \lambda_1 + \frac{8}{3}\pi \lambda_2} + e^{\frac{4}{3}\pi b \lambda_1 - \frac{4}{3}\pi \lambda_2} + e^{-\frac{8}{3}\pi b \lambda_1 - \frac{4}{3}\pi \lambda_2}, \\ \pi_\lambda(\mathbf{C}_2) &= e^{\frac{8}{3}\pi b \lambda_1 + \frac{4}{3}\pi \lambda_2} + e^{-\frac{4}{3}\pi b \lambda_1 + \frac{4}{3}\pi \lambda_2} + e^{-\frac{4}{3}\pi b \lambda_1 - \frac{8}{3}\pi \lambda_2}. \end{aligned}$$

# Casimir Operators

## Proposition

In type  $A_n$ ,  $V_k = \Lambda^k V_1$ , and the actions of  $\mathbf{C}_k$  on  $\mathcal{P}_\lambda$  are given by

$$\pi_\lambda(\mathbf{C}_k) = \mathcal{E}_k(e^{4\pi b\varpi_0}, \dots, e^{4\pi b\varpi_n}).$$

where  $\mathcal{E}_k$  are the *elementary symmetric polynomials* in  $n+1$  variables, and

$$\varpi_i := \frac{1}{n+1} \sum_{k=1}^n k\lambda_k - \sum_{j=n+1-i}^n \lambda_j, \quad i = 0, \dots, n.$$


## Example

In  $\mathcal{U}_q(\mathfrak{sl}_3)$ , (type  $A_2$ ) we have

$$\begin{aligned} \pi_\lambda(\mathbf{C}_1) &= e^{\frac{4}{3}\pi b\lambda_1 + \frac{8}{3}\pi\lambda_2} + e^{\frac{4}{3}\pi b\lambda_1 - \frac{4}{3}\pi\lambda_2} + e^{-\frac{8}{3}\pi b\lambda_1 - \frac{4}{3}\pi\lambda_2}, \\ \pi_\lambda(\mathbf{C}_2) &= e^{\frac{8}{3}\pi b\lambda_1 + \frac{4}{3}\pi\lambda_2} + e^{-\frac{4}{3}\pi b\lambda_1 + \frac{4}{3}\pi\lambda_2} + e^{-\frac{4}{3}\pi b\lambda_1 - \frac{8}{3}\pi\lambda_2}. \end{aligned}$$

# Symmetric Folding


We start with the standard quiver  $\mathbf{D}(\mathbf{i}_0)$  for  $\mathbf{i}_0 = (1 \ 21 \ 321 \ 4321 \cdots n \ \cdots \ 1)$ .



- The center of  $\mathcal{X}_q^{std}$  is generated by  $\iota(\mathbf{K}_i \mathbf{K}'_i)$  and the **monodromies**  $Q_i$  around the puncture.
- The polarization is given by  $\pi_\lambda(\mathbf{K}_i \mathbf{K}'_i) = 1$  and  $\pi_\lambda(Q_i) = e^{4\pi b \lambda_i}$ .

# Symmetric Folding


We start with the standard quiver  $\mathbf{D}(\mathbf{i}_0)$  for  $\mathbf{i}_0 = (1 \ 21 \ 321 \ 4321 \cdots n \ \cdots \ 1)$ .



- The center of  $\mathcal{X}_q^{std}$  is generated by  $\iota(\mathbf{K}_i \mathbf{K}'_i)$  and the **monodromies**  $Q_i$  around the puncture.
- The polarization is given by  $\pi_\lambda(\mathbf{K}_i \mathbf{K}'_i) = 1$  and  $\pi_\lambda(Q_i) = e^{4\pi b \lambda_i}$ .

# Symmetric Folding

We perform the flip to obtain a  
self-folded triangulation




and obtain the following quiver.


There is a mutation sequence by  
[Schrader-Shapiro] which transform the  
top half of the quiver into  $\mathcal{X}_q^{sym}$ .

# Symmetric Folding

We perform the flip to obtain a **self-folded triangulation**




and obtain the following quiver.



There is a mutation sequence by [Schrader-Shapiro] which transform the top half of the quiver into  $\mathcal{X}_q^{sym}$ .


# Symmetric Folding

We perform the flip to obtain a **self-folded triangulation**



and obtain the following quiver.

There is a mutation sequence by [Schrader-Shapiro] which transform the top half of the quiver into  $\mathcal{X}_q^{sym}$ .





# Symmetric Folding

The central monodromies become


$$Q_i := X_{c_i} X_{c_{i-1}}^{-1}, \quad \pi_\lambda(Q_i) = e^{4\pi b \lambda_i}$$

The embedding of the generators  $\mathbf{f}_1, \dots, \mathbf{f}_n$  as well as  $\mathbf{e}_1, \dots, \mathbf{e}_{n-1}$  does not involve the indices  $c_i$ .

However,  $\mathbf{e}_n$  is of the form

$$\iota(\mathbf{e}_n) = \sum A_k B_k$$

where  $B_k = \mathcal{E}_k(X_{c_0}, \dots, X_{c_n})$  which is proportional to  $\iota(\mathbf{C}_k)$ !



# Symmetric Folding

The central monodromies become


$$Q_i := X_{c_i} X_{c_{i-1}}^{-1}, \quad \pi_\lambda(Q_i) = e^{4\pi b \lambda_i}$$

The embedding of the generators  $\mathbf{f}_1, \dots, \mathbf{f}_n$  as well as  $\mathbf{e}_1, \dots, \mathbf{e}_{n-1}$  does not involve the indices  $c_i$ .

However,  $\mathbf{e}_n$  is of the form

$$\iota(\mathbf{e}_n) = \sum A_k B_k$$

where  $B_k = \mathcal{E}_k(X_{c_0}, \dots, X_{c_n})$  which is proportional to  $\iota(\mathbf{C}_k)$ !



# Symmetric Folding

The central monodromies become


$$Q_i := X_{c_i} X_{c_{i-1}}^{-1}, \quad \pi_\lambda(Q_i) = e^{4\pi b \lambda_i}$$

The embedding of the generators  $\mathbf{f}_1, \dots, \mathbf{f}_n$  as well as  $\mathbf{e}_1, \dots, \mathbf{e}_{n-1}$  does not involve the indices  $c_i$ .

However,  $\mathbf{e}_n$  is of the form

$$\iota(\mathbf{e}_n) = \sum A_k B_k$$

where  $B_k = \mathcal{E}_k(X_{c_0}, \dots, X_{c_n})$  which is proportional to  $\iota(\mathbf{C}_k)$ !



# Symmetric Folding

The central monodromies become


$$Q_i := X_{c_i} X_{c_{i-1}}^{-1}, \quad \pi_\lambda(Q_i) = e^{4\pi b \lambda_i}$$

The embedding of the generators  $\mathbf{f}_1, \dots, \mathbf{f}_n$  as well as  $\mathbf{e}_1, \dots, \mathbf{e}_{n-1}$  does not involve the indices  $c_i$ .

However,  $\mathbf{e}_n$  is of the form

$$\iota(\mathbf{e}_n) = \sum A_k B_k$$

where  $B_k = \mathcal{E}_k(X_{c_0}, \dots, X_{c_n})$  which is proportional to  $\iota(\mathbf{C}_k)$ !



# Symmetric Folding

We can now perform the **folding** of the symmetric variables to get  $\mathcal{X}_q^0$ , where

$$X_\star := \prod_{k=0}^n X_{c_k}$$

and assign  $d_\star := n + 1$ .

- The embeddings of  $\mathbf{f}_1, \dots, \mathbf{f}_n$  and  $\mathbf{e}_1, \dots, \mathbf{e}_{n-1}$  are unaffected.
- $\iota(\mathbf{e}_n)$  is modified such that

$$\begin{cases} B_k \mapsto 0 & k = 1, \dots, n, \\ B_{n+1} \mapsto X_\star. \end{cases}$$

By construction  $\mathbf{C}_k = 0$  in  $\mathcal{X}_q^0$ .

# Symmetric Folding

We can now perform the **folding** of the symmetric variables to get  $\mathcal{X}_q^0$ , where


$$X_\star := \prod_{k=0}^n X_{c_k}$$

and assign  $d_\star := n + 1$ .

- The embeddings of  $\mathbf{f}_1, \dots, \mathbf{f}_n$  and  $\mathbf{e}_1, \dots, \mathbf{e}_{n-1}$  are unaffected.
- $\iota(\mathbf{e}_n)$  is modified such that

$$\begin{cases} B_k \mapsto 0 & k = 1, \dots, n, \\ B_{n+1} \mapsto X_\star. \end{cases}$$

By construction  $\mathbf{C}_k = 0$  in  $\mathcal{X}_q^0$ .



# Symmetric Folding

We can now perform the **folding** of the symmetric variables to get  $\mathcal{X}_q^0$ , where


$$X_\star := \prod_{k=0}^n X_{c_k}$$

and assign  $d_\star := n + 1$ .

- The embeddings of  $\mathbf{f}_1, \dots, \mathbf{f}_n$  and  $\mathbf{e}_1, \dots, \mathbf{e}_{n-1}$  are unaffected.
- $\iota(\mathbf{e}_n)$  is modified such that

$$\begin{cases} B_k \mapsto 0 & k = 1, \dots, n, \\ B_{n+1} \mapsto X_\star. \end{cases}$$

By construction  $\mathbf{C}_k = 0$  in  $\mathcal{X}_q^0$ .



# Main Theorem: Type $A_n$

Theorem (I.-Man (2022), I.-Ye (2023))

*There is an embedding*

$$\mathcal{D}_q(\mathfrak{sl}_{n+1})/\langle \mathbf{C}_k = 0 \rangle \hookrightarrow \mathcal{O}_q(X^0)$$

*to a skew-symmetrizable quantum cluster algebra such that*

- *The image of  $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i, \mathbf{K}'_i\}$  are **universally polynomials**.*
- *We have an irreducible representation  $\mathcal{P}^0$  of  $\mathcal{U}_q(\mathfrak{sl}(n+1, \mathbb{R}))$  acting on  $L^2(\mathbb{R}^N)$  as positive operators, such that  $\pi(\mathbf{C}_k) = 0$ .*



# Main Theorem: Type $A_n$

Theorem (I.-Man (2022), I.-Ye (2023))

*There is an embedding*


$$\mathcal{D}_q(\mathfrak{sl}_{n+1}) / \langle \mathbf{C}_k = 0 \rangle \hookrightarrow \mathcal{O}_q(X^0)$$

*to a skew-symmetrizable quantum cluster algebra such that*

- *The image of  $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i, \mathbf{K}'_i\}$  are **universally polynomials**.*
- *We have an irreducible representation  $\mathcal{P}^0$  of  $\mathcal{U}_q(\mathfrak{sl}(n+1, \mathbb{R}))$  acting on  $L^2(\mathbb{R}^N)$  as positive operators, such that  $\pi(\mathbf{C}_k) = 0$ .*

# Degenerate Modular Double

Recall



The vertex 2 and 4 are *symmetric* in  $\mathcal{X}_q^{sym}$ .

$$\mathbf{e} \mapsto X_3 + X_1^{-1} \mathbf{C} + X_{3,2,4}$$


$$\mathbf{f} \mapsto X_1$$

$$\mathbf{K} \mapsto X_{3,2,4,1}$$

$$\mathbf{K}' \mapsto X_{1,3}$$

# Degenerate Modular Double

By reversing the folding, we obtain a new *skew-symmetrizable*  $\tilde{\mathcal{X}}_q^0$



and a homomorphism  $\mathcal{D}_q(\mathfrak{sl}_2) \longrightarrow \tilde{\mathcal{X}}_q^0$ :

$$\mathbf{e} \mapsto X_3 + (q^{\frac{1}{2}} + q^{-\frac{1}{2}})X_{3,*} + X_{3,*^2}$$

$$\mathbf{f} \mapsto X_1$$

$$\mathbf{K} \mapsto X_{3,*^2,1}$$

$$\mathbf{K}' \mapsto X_{1,3}$$

We obtain the [modular double counterpart](#) of the degenerate representation  $\tilde{\mathcal{P}}^0$  of  $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$  by polarization!

# Degenerate representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

# Degenerate Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

- Recall a parabolic subgroup  $\longleftrightarrow J \subset I$  provides a decomposition of  $\mathcal{X}_q^{std}$ .

$$\mathbf{i}_0 = \mathbf{i}_J \mathbf{i}'.$$

- We can fold the  $\mathbf{i}_J$  part if it is of type  $A_{k_1} \times \cdots \times A_{k_m}$ .
- The number of central parameters are reduced by  $|J| = k_1 + \cdots + k_m$ .

# Degenerate Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

- Recall a parabolic subgroup  $\longleftrightarrow J \subset I$  provides a decomposition of  $\mathcal{X}_q^{std}$ .

$$\mathbf{i}_0 = \mathbf{i}_J \mathbf{i}'.$$

- We can **fold** the  $\mathbf{i}_J$  part if it is of type  $A_{k_1} \times \cdots \times A_{k_m}$ .
- The number of central parameters are reduced by  $|J| = k_1 + \cdots + k_m$ .

# Degenerate Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

- Recall a parabolic subgroup  $\longleftrightarrow J \subset I$  provides a decomposition of  $\mathcal{X}_q^{std}$ .

$$\mathbf{i}_0 = \mathbf{i}_J \mathbf{i}'.$$

- We can **fold** the  $\mathbf{i}_J$  part if it is of type  $A_{k_1} \times \cdots \times A_{k_m}$ .
- The number of central parameters are reduced by  $|J| = k_1 + \cdots + k_m$ .

# Degenerate Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

## Theorem (I.-Man (2022))

For each parabolic subgroup  $W_J \subset W$  of type  $A_{k_1} \times \cdots \times A_{k_m}$ :

- We have a homomorphism

$$\mathcal{D}_q(\mathfrak{g}) \longrightarrow \mathcal{O}_q(\mathcal{X}^0)$$

to some skew-symmetrizable quantum cluster algebra.

- $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i, \mathbf{K}'_i\}$  are universally polynomials.
- We have a family of irreducible regular positive representations  $\mathcal{P}_{\lambda}^{0,J}$  of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$  parametrized by  $\lambda \in \mathbb{R}^{n-|J|}$ .

- The same results hold for the modular double counterpart with  $\mathcal{X}_q^0$ , where the multipliers  $d_*$  of any folded vertices are inverted.



# Degenerate Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

## Theorem (I.-Man (2022))

For each parabolic subgroup  $W_J \subset W$  of type  $A_{k_1} \times \cdots \times A_{k_m}$ :

- We have a homomorphism


$$\mathcal{D}_q(\mathfrak{g}) \longrightarrow \mathcal{O}_q(\mathcal{X}^0)$$

to some skew-symmetrizable quantum cluster algebra.


- $\{\mathbf{e}_i, \mathbf{f}_i, \mathbf{K}_i, \mathbf{K}'_i\}$  are universally polynomials.
- We have a family of irreducible regular positive representations  $\mathcal{P}_{\lambda}^{0,J}$  of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$  parametrized by  $\lambda \in \mathbb{R}^{n-|J|}$ .
- The same results hold for the **modular double counterpart** with  $\tilde{\mathcal{X}}_q^0$ , where the multipliers  $d_{\star}$  of any folded vertices are inverted.

# Maximal Degenerate Representations


$$i_0 = i_{A_4} i' = (1213214321)(545345234512345)$$



# Maximal Degenerate Representations



# Maximal Degenerate Representations



Polarization gives the degenerate representation  $P_\lambda^{0,J}$  of  $\mathcal{U}_q(\mathfrak{g}_{B_5})$ .

# Classification of Regular Positive Representations

# Classification of Regular Positive Representations

We have the following list of irreducible regular positive representations of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ :

- (1) The standard positive representations  $\mathcal{P}_{\lambda}$ , parametrized by  $n$  positive scalars.
- (2) The parabolic positive representations  $\mathcal{P}_{\lambda}^J$ , with respect to parabolic subgroup  $W_J \subset W$  parametrized by  $n - |J|$  positive scalars.
- (3) The degenerate representations  $\mathcal{P}_{\lambda}^{0,J}$  with respect to parabolic subgroup  $W_J \subset W$  of type  $A_{k_1} \times \cdots \times A_{k_m}$ , parametrized by  $n - |J|$  positive scalars.
- (4) The modular double counterpart of the degenerate representations  $\mathcal{P}_{\lambda}^{0,J}$ , also parametrized by  $n - |J|$  positive scalars.
- (5) A mixture of type (2)–(4) for disconnected subsets of Dynkin index.

## Conjecture

*The classes of all irreducible regular positive representations of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$  are classified by the list (1)–(5) above.*

# Classification of Regular Positive Representations

We have the following list of irreducible regular positive representations of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ :

- (1) The standard positive representations  $\mathcal{P}_{\lambda}$ , parametrized by  $n$  positive scalars.
- (2) The parabolic positive representations  $\mathcal{P}_{\lambda}^J$ , with respect to parabolic subgroup  $W_J \subset W$  parametrized by  $n - |J|$  positive scalars.
- (3) The degenerate representations  $\mathcal{P}_{\lambda}^{0,J}$  with respect to parabolic subgroup  $W_J \subset W$  of type  $A_{k_1} \times \cdots \times A_{k_m}$ , parametrized by  $n - |J|$  positive scalars.
- (4) The modular double counterpart of the degenerate representations  $\mathcal{P}_{\lambda}^{0,J}$ , also parametrized by  $n - |J|$  positive scalars.
- (5) A mixture of type (2)–(4) for disconnected subsets of Dynkin index.

## Conjecture

*The classes of all irreducible regular positive representations of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$  are classified by the list (1)–(5) above.*

# Classification of Regular Positive Representations

We have the following list of irreducible regular positive representations of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ :

- (1) The standard positive representations  $\mathcal{P}_{\lambda}$ , parametrized by  $n$  positive scalars.
- (2) The parabolic positive representations  $\mathcal{P}_{\lambda}^J$ , with respect to parabolic subgroup  $W_J \subset W$  parametrized by  $n - |J|$  positive scalars.
- (3) The degenerate representations  $\mathcal{P}_{\lambda}^{0,J}$  with respect to parabolic subgroup  $W_J \subset W$  of type  $A_{k_1} \times \cdots \times A_{k_m}$ , parametrized by  $n - |J|$  positive scalars.
- (4) The modular double counterpart of the degenerate representations  $\mathcal{P}_{\lambda}^{0,J}$ , also parametrized by  $n - |J|$  positive scalars.
- (5) A mixture of type (2)–(4) for disconnected subsets of Dynkin index.

## Conjecture

*The classes of all irreducible regular positive representations of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$  are classified by the list (1)–(5) above.*



# Classification of Regular Positive Representations

We have the following list of irreducible regular positive representations of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ :

- (1) The standard positive representations  $\mathcal{P}_{\lambda}$ , parametrized by  $n$  positive scalars.
- (2) The parabolic positive representations  $\mathcal{P}_{\lambda}^J$ , with respect to parabolic subgroup  $W_J \subset W$  parametrized by  $n - |J|$  positive scalars.
- (3) The degenerate representations  $\mathcal{P}_{\lambda}^{0,J}$  with respect to parabolic subgroup  $W_J \subset W$  of type  $A_{k_1} \times \cdots \times A_{k_m}$ , parametrized by  $n - |J|$  positive scalars.
- (4) The modular double counterpart of the degenerate representations  $\mathcal{P}_{\lambda}^{0,J}$ , also parametrized by  $n - |J|$  positive scalars.
- (5) A mixture of type (2)–(4) for disconnected subsets of Dynkin index.

## Conjecture

*The classes of all irreducible regular positive representations of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$  are classified by the list (1)–(5) above.*

# Classification of Regular Positive Representations

We have the following list of irreducible regular positive representations of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ :

- (1) The standard positive representations  $\mathcal{P}_{\lambda}$ , parametrized by  $n$  positive scalars.
- (2) The parabolic positive representations  $\mathcal{P}_{\lambda}^J$ , with respect to parabolic subgroup  $W_J \subset W$  parametrized by  $n - |J|$  positive scalars.
- (3) The degenerate representations  $\mathcal{P}_{\lambda}^{0,J}$  with respect to parabolic subgroup  $W_J \subset W$  of type  $A_{k_1} \times \cdots \times A_{k_m}$ , parametrized by  $n - |J|$  positive scalars.
- (4) The modular double counterpart of the degenerate representations  $\mathcal{P}_{\lambda}^{0,J}$ , also parametrized by  $n - |J|$  positive scalars.
- (5) A mixture of type (2)–(4) for disconnected subsets of Dynkin index.

## Conjecture

*The classes of all irreducible regular positive representations of  $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$  are classified by the list (1)–(5) above.*

# Example: Type $A_2$

We illustrate the joint spectrum  $(\pi_\lambda(\mathbf{C}_1), \pi_\lambda(\mathbf{C}_2))$  of the Casimirs of the irreducible regular positive representations of  $\mathcal{U}_q(\mathfrak{sl}_3)$ . We assume  $\lambda \geq 0$  and  $q$  not a root of unity.

Representations	$J$	Action of Casimirs
$\mathcal{P}_\lambda$		$\pi_\lambda(\mathbf{C}_1) = e^{\frac{4}{3}\pi b\lambda_1 + \frac{8}{3}\pi b\lambda_2} + e^{\frac{4}{3}\pi b\lambda_1 - \frac{4}{3}\pi b\lambda_2} + e^{-\frac{8}{3}\pi b\lambda_1 - \frac{4}{3}\pi b\lambda_2}$ $\pi_\lambda(\mathbf{C}_2) = e^{\frac{8}{3}\pi b\lambda_1 + \frac{4}{3}\pi b\lambda_2} + e^{-\frac{4}{3}\pi b\lambda_1 + \frac{4}{3}\pi b\lambda_2} + e^{-\frac{4}{3}\pi b\lambda_1 - \frac{8}{3}\pi b\lambda_2}$
$\mathcal{P}_\lambda^J$	$\{1\}$	$\pi_\lambda(\mathbf{C}_1) = e^{-\frac{8}{3}\pi b\lambda} - (q + q^{-1})e^{\frac{4}{3}\pi b\lambda}$ $\pi_\lambda(\mathbf{C}_2) = e^{\frac{8}{3}\pi b\lambda} - (q + q^{-1})e^{-\frac{4}{3}\pi b\lambda}$
	$\{2\}$	$\pi_\lambda(\mathbf{C}_1) = e^{\frac{8}{3}\pi b\lambda} - (q + q^{-1})e^{-\frac{4}{3}\pi b\lambda}$ $\pi_\lambda(\mathbf{C}_2) = e^{-\frac{8}{3}\pi b\lambda} - (q + q^{-1})e^{\frac{4}{3}\pi b\lambda}$
$\mathcal{P}_\lambda^{0,J}$	$\{1, 2\}$	$\pi_\lambda(\mathbf{C}_1) = 0$ $\pi_\lambda(\mathbf{C}_2) = 0$
	$\{1\}$	$\pi_\lambda(\mathbf{C}_1) = e^{\frac{8}{3}\pi b\lambda}$ $\pi_\lambda(\mathbf{C}_2) = e^{-\frac{8}{3}\pi b\lambda}$
	$\{2\}$	$\pi_\lambda(\mathbf{C}_1) = e^{-\frac{8}{3}\pi b\lambda}$ $\pi_\lambda(\mathbf{C}_2) = e^{\frac{8}{3}\pi b\lambda}$
$\mathcal{P}_{\bar{\lambda}}^{0,J}$	$\{1, 2\}$	$\pi_\lambda(\mathbf{C}_1) = q^{\frac{2}{3}} + 1 + q^{-\frac{2}{3}}$ $\pi_\lambda(\mathbf{C}_2) = q^{\frac{2}{3}} + 1 + q^{-\frac{2}{3}}$
	$\{1\}$	$\pi_\lambda(\mathbf{C}_1) = e^{\frac{4}{3}\pi b\lambda} + (q^{\frac{1}{2}} + q^{-\frac{1}{2}})e^{-\frac{4}{3}\pi b\lambda}$ $\pi_\lambda(\mathbf{C}_2) = e^{-\frac{4}{3}\pi b\lambda} + (q^{\frac{1}{2}} + q^{-\frac{1}{2}})e^{\frac{4}{3}\pi b\lambda}$
	$\{2\}$	$\pi_\lambda(\mathbf{C}_1) = e^{-\frac{4}{3}\pi b\lambda} + (q^{\frac{1}{2}} + q^{-\frac{1}{2}})e^{\frac{4}{3}\pi b\lambda}$ $\pi_\lambda(\mathbf{C}_2) = e^{\frac{4}{3}\pi b\lambda} + (q^{\frac{1}{2}} + q^{-\frac{1}{2}})e^{-\frac{4}{3}\pi b\lambda}$


# Example: Type $A_2$

We illustrate the joint spectrum  $(\pi_\lambda(\mathbf{C}_1), \pi_\lambda(\mathbf{C}_2))$  of the Casimirs of the irreducible regular positive representations of  $\mathcal{U}_q(\mathfrak{sl}_3)$ . We assume  $\lambda \geq 0$  and  $q$  not a root of unity.

Representations	$J$	Action of Casimirs
$\mathcal{P}_\lambda$		$\pi_\lambda(\mathbf{C}_1) = e^{\frac{4}{3}\pi b\lambda_1 + \frac{8}{3}\pi b\lambda_2} + e^{\frac{4}{3}\pi b\lambda_1 - \frac{4}{3}\pi b\lambda_2} + e^{-\frac{8}{3}\pi b\lambda_1 - \frac{4}{3}\pi b\lambda_2}$ $\pi_\lambda(\mathbf{C}_2) = e^{\frac{8}{3}\pi b\lambda_1 + \frac{4}{3}\pi b\lambda_2} + e^{-\frac{4}{3}\pi b\lambda_1 + \frac{4}{3}\pi b\lambda_2} + e^{-\frac{4}{3}\pi b\lambda_1 - \frac{8}{3}\pi b\lambda_2}$
$\mathcal{P}_\lambda^J$	$\{1\}$	$\pi_\lambda(\mathbf{C}_1) = e^{-\frac{8}{3}\pi b\lambda} - (q + q^{-1})e^{\frac{4}{3}\pi b\lambda}$ $\pi_\lambda(\mathbf{C}_2) = e^{\frac{8}{3}\pi b\lambda} - (q + q^{-1})e^{-\frac{4}{3}\pi b\lambda}$
	$\{2\}$	$\pi_\lambda(\mathbf{C}_1) = e^{\frac{8}{3}\pi b\lambda} - (q + q^{-1})e^{-\frac{4}{3}\pi b\lambda}$ $\pi_\lambda(\mathbf{C}_2) = e^{-\frac{8}{3}\pi b\lambda} - (q + q^{-1})e^{\frac{4}{3}\pi b\lambda}$
$\mathcal{P}_\lambda^{0,J}$	$\{1, 2\}$	$\pi_\lambda(\mathbf{C}_1) = 0$ $\pi_\lambda(\mathbf{C}_2) = 0$
	$\{1\}$	$\pi_\lambda(\mathbf{C}_1) = e^{\frac{8}{3}\pi b\lambda}$ $\pi_\lambda(\mathbf{C}_2) = e^{-\frac{8}{3}\pi b\lambda}$
	$\{2\}$	$\pi_\lambda(\mathbf{C}_1) = e^{-\frac{8}{3}\pi b\lambda}$ $\pi_\lambda(\mathbf{C}_2) = e^{\frac{8}{3}\pi b\lambda}$
$\mathcal{P}_{\tilde{\lambda}}^{0,J}$	$\{1, 2\}$	$\pi_\lambda(\mathbf{C}_1) = q^{\frac{2}{3}} + 1 + q^{-\frac{2}{3}}$ $\pi_\lambda(\mathbf{C}_2) = q^{\frac{2}{3}} + 1 + q^{-\frac{2}{3}}$
	$\{1\}$	$\pi_\lambda(\mathbf{C}_1) = e^{\frac{8}{3}\pi b\lambda} + (q^{\frac{1}{2}} + q^{-\frac{1}{2}})e^{-\frac{4}{3}\pi b\lambda}$ $\pi_\lambda(\mathbf{C}_2) = e^{-\frac{8}{3}\pi b\lambda} + (q^{\frac{1}{2}} + q^{-\frac{1}{2}})e^{\frac{4}{3}\pi b\lambda}$
	$\{2\}$	$\pi_\lambda(\mathbf{C}_1) = e^{-\frac{8}{3}\pi b\lambda} + (q^{\frac{1}{2}} + q^{-\frac{1}{2}})e^{\frac{4}{3}\pi b\lambda}$ $\pi_\lambda(\mathbf{C}_2) = e^{\frac{8}{3}\pi b\lambda} + (q^{\frac{1}{2}} + q^{-\frac{1}{2}})e^{-\frac{4}{3}\pi b\lambda}$

# Example: Type $A_2$


A generic plot of  $(x, y) = (\pi_\lambda(\mathbf{C}_1), \pi_\lambda(\mathbf{C}_2))$  with  $b \sim 0.5$ .



If  $q$  is a root of unity, these family of Casimir values can overlap!

## Example: Type $A_2$

A generic plot of  $(x, y) = (\pi_\lambda(\mathbf{C}_1), \pi_\lambda(\mathbf{C}_2))$  with  $b \sim 0.5$ .



If  $q$  is a root of unity, these family of Casimir values can overlap!

Thank you for your attention!