

2016/8/26

1. Semisimple Lie algebras

Def of Lie alg

A Lie algebra is a \mathbb{C} -vector space \mathfrak{g} equipped with a bilinear map

$[\cdot, \cdot]: \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ satisfying

- $[x, y] = -[y, x] \quad (\Leftrightarrow [x, x] = 0)$
- $[x, [y, z]] = [[x, y], z] + [y, [x, z]]$

Jacobi identity //

Notion of homomorphism, subalg, ideal are obviously defined.

A Lie alg \mathfrak{g} is abelian

\Leftrightarrow def $\forall x, y \in \mathfrak{g} \quad [x, y] = 0$

Ex

(1) A : associative \mathbb{C} -alg

$$[x, y] := xy - yx$$

$\rightarrow (A, [,]) \text{ Lie alg}$

In particular, $\text{End}_{\mathbb{C}}(V)$ for a \mathbb{C} -vec
sp. V is a Lie alg.

Denote the Lie alg $\text{End}(\mathbb{C}^n)$ by $\mathfrak{gl}_n(\mathbb{C})$
(general linear Lie alg)

(2)

$$\mathfrak{sl}_n(\mathbb{C}) := \{x \in \mathfrak{gl}_n(\mathbb{C}) \mid \text{tr } x = 0\}$$

is a Lie subalg of $\mathfrak{gl}_n(\mathbb{C})$

(special linear Lie alg)

$$x, y \in \mathfrak{sl}_n(\mathbb{C}) \Rightarrow \text{tr}[x, y] = \text{tr } xy - \text{tr } yx \\ = 0 \quad \square$$

$$\mathfrak{sl}_2 = \mathfrak{sl}_2(\mathbb{C})$$

3

$$= \mathbb{C} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \oplus \mathbb{C} \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \oplus \mathbb{C} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$\begin{matrix} \Downarrow \\ e \end{matrix}$
 $\begin{matrix} \Downarrow \\ f \end{matrix}$
 $\begin{matrix} \Downarrow \\ h \end{matrix}$

$$\begin{cases} [e, f] = ef - fe = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} - \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} = h \\ [h, e] = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} - \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = 2e \\ [h, f] = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} - \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} = -2f \end{cases}$$

Universal enveloping alg

$$\begin{array}{ccc} \text{Category of} & \xrightarrow{\text{Forget}} & \text{Category of} \\ \text{ass } \mathbb{C}\text{-alg} & & \text{Lie alg } / \mathbb{C} \\ & \nwarrow & \\ & \exists \cup \text{ left adjoint} & \end{array}$$

$$\text{Lie. Hom}_{\text{Lie alg}}(\mathfrak{g}, \text{For}(A))$$

$$\cong \text{Hom}_{\mathbb{C}\text{-alg}}(\cup(\mathfrak{g}), A)$$

Construction of $\cup(\mathfrak{g})$:

$$T(\mathfrak{g}) := \bigoplus_{n=0}^{\infty} \mathfrak{g}^{\otimes n} \quad \text{tensor alg}$$

$$\cup(\mathfrak{g}) := T(\mathfrak{g}) / \langle x \otimes y - y \otimes x - [x, y] \mid x, y \in \mathfrak{g} \rangle$$

two-sided

Denote the image of $x_1 \otimes \dots \otimes x_n \in \mathfrak{g}^{\otimes n} \subset$
by $x_1 \dots x_n \in U(\mathfrak{g})$.

$$U(\mathfrak{g}) = \text{span}_{\mathbb{C}} \{ x_1 \dots x_n \mid \substack{n \geq 0 \\ x_i \in \mathfrak{g}} \}$$

Ex

$$U(\mathfrak{sl}_2) = \text{span}_{\mathbb{C}} \{ x_1 \dots x_n \mid \substack{n \geq 0 \\ x_i = e, f, h} \}$$

We can rewrite a product w.r.t.
the order f, h, e .

e.g. $efhe$

$$= ([ef] + fe)he$$

$$[ef] = h$$

$$= h^2e + f \underline{e} h e$$

$$\begin{aligned} eh &= [e, h] + he \\ &= -2e + he \end{aligned}$$

$$= h^2e - 2fe^2 + fhe^2 //$$

$$\therefore U(\mathfrak{sl}_2) = \text{span}_{\mathbb{C}} \{ f^k h^l e^m \mid k, l, m \geq 0 \}$$

In fact $f^k h^l e^m$ ($k, l, m \in \mathbb{N}_{\geq 0}$) are
(linearly) independent.

\therefore It gives a \mathbb{C} -basis of $U(\mathfrak{sl}_2)$.

$\mathfrak{g} = \text{Lie alg}$

5

Define a filtration $F^\bullet U(\mathfrak{g})$ of $U(\mathfrak{g})$ by

$$F^k U(\mathfrak{g}) := \text{Im} \left(\bigoplus_{h=0}^k \mathfrak{g}^{\otimes h} \rightarrow U(\mathfrak{g}) \right)$$

Thm (Poincaré - Birkhoff - Witt)

$$\mathfrak{g} \cap F^\bullet U(\mathfrak{g}) \xleftarrow{\cong} \mathfrak{S}(\mathfrak{g}) \quad \text{symmetric alg}$$

Cor 1

(x_1, x_2, \dots) = an ordered basis of \mathfrak{g}

$\Rightarrow \{ x_1^{k_1} \dots x_n^{k_n} \mid \substack{k_i \geq 0 \\ k_i \in \mathbb{Z}} \}$ gives a \mathbb{C} -basis of $U(\mathfrak{g})$.

Cor 2

$\mathfrak{g} = \text{fin. dim'l } \mathbb{C}$

$\Rightarrow U(\mathfrak{g})$ is left and right Noetherian.

Representation

6

A representation of \mathcal{G} is a pair of a \mathbb{C} -vec sp M and a Lie alg hom

$$\varphi: \mathcal{G} \longrightarrow \text{End}_{\mathbb{C}}(M).$$

We simply say that M is a rep of \mathcal{G} or M is a \mathcal{G} -module.

Rem

The category of \mathcal{G} -modules and the category of left $U(\mathcal{G})$ -modules are isomorphic.

$$\therefore \text{Hom}_{\text{Lie alg}}(\mathcal{G}, \text{End}(M)) \cong \text{Hom}_{\text{alg}}(U(\mathcal{G}), \text{End}(M))$$

We will not distinguish \mathcal{G} -modules and $U(\mathcal{G})$ -modules.

Ex

7

(1) $M = \{0\}$

(2) $M = \mathbb{C} \quad \forall x \in \mathfrak{g} \text{ acts by } 0$

trivial rep

(3) $M = \mathfrak{g}$

$\text{ad}: \mathfrak{g} \longrightarrow \text{End}_{\mathbb{C}}(\mathfrak{g})$ adjoint rep

$x \longmapsto [x, -]$

$\star \text{ ad is a Lie alg hom} \Leftrightarrow \text{Jacobi Identity}$

$\therefore \text{ ad } [x, y] \stackrel{?}{=} [\text{ad } x, \text{ad } y]$

"

$\text{ad } x \text{ ad } y - \text{ad } y \text{ ad } x$

LHS: $\text{ad}[x, y](z) = [[x, y], z]$

RHS: $[\text{ad } x, \text{ad } y](z)$

$= [x, [y, z]] - [y, [x, z]] \quad \square$

Rep of \mathfrak{sl}_2

• $L(0) = \mathbb{C}$

trivial rep

• $L(1) = \mathbb{C}^2$

$\mathfrak{sl}_2 \subset \mathfrak{gl}_2 = \text{End } \mathbb{C}^2 \simeq \mathbb{C}^3$
acts as matrices

vector rep

• $L(2) = \mathfrak{sl}_2 \cong \mathbb{C}^3$ adjoint rep δ

What does $L(\underline{n})$ mean?

• $L(1) = \mathbb{C}(h) \oplus \mathbb{C}(e)$

$$A \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \begin{array}{l} \text{eigenvalue} \\ \underline{1} \end{array}$$

$$A \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = - \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad -1$$

eigenvalue of A is called weight.

★ $e \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = 0$

~~e~~ $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ is called highest weight vector

1 is called highest weight

• $L(2) = \mathbb{C}e \oplus \mathbb{C}A \oplus \mathbb{C}f$
 $\quad \quad \quad \text{wt} \quad \quad \quad 2 \quad \quad \quad 0 \quad \quad \quad -2$

($[A, e] = 2e$, $[A, f] = -2f$)

$[e, e] = 0$

$\Rightarrow e$ is a h.w. vector

with h.w. 2

Then

$\{ \text{fin. dim. simple } \mathfrak{sl}_2\text{-modules} \} / \cong$

$$= \{ L(n) \mid \begin{array}{l} (n+1)\text{-dim'l} \\ \text{h.w. } n \in \mathbb{Z}_{\geq 0} \end{array} \} //$$

NOTICE:

$M: \mathfrak{sl}_2\text{-module}$

Suppose $v \in M$ has wt $m \in \mathbb{C}$ i.e.

$$Av = mv.$$

Then $e v$ has wt $m+2$ if $e v \neq 0$

$f v$ has wt $m-2$ if $f v \neq 0$

$$\therefore A e v = ([A, e] + e A) v \quad [A, e] = 2e$$

$$= 2e v + m e v$$

$$= (m+2) e v \quad \square$$

Important

$$\bullet \mathfrak{sl}_2 = \mathbb{C} f \oplus \mathbb{C} h \oplus \mathbb{C} e$$

\bullet eigen space decomposition w.r.t.

h -action (eigenvalue = weight)

\bullet e raises wt by 2

f lowers wt by 2.

Semisimple Lie alg

10

A Lie alg \mathfrak{g} is simple

\Leftrightarrow def \bullet \mathfrak{g} is not abelian

\bullet \mathfrak{g} does not have nontrivial ideals //

"Def"

semisimple Lie algebra

= direct product of simple Lie algebras

ordinary def:

$\left(\begin{array}{l} \text{s.s.} \xLeftrightarrow{\text{def}} \text{radical} = 0 \quad (\text{maximal solvable ideal}) \\ \Leftrightarrow \text{direct prod of simples} \end{array} \right)$

Then
analog of Wedderburn then

Ex

$$\mathfrak{sl}_n(\mathbb{C}) = \{ x \in \mathfrak{gl}_n(\mathbb{C}) = \text{End}(\mathbb{C}^n) \mid \text{tr } x = 0 \}$$

is simple.

$\mathfrak{gl}_n(\mathbb{C})$ is not simple

but reductive = semisimple \times center //

Assume \mathfrak{g} = fin. dim. simple Lie alg
in the sequel.

Root space decomposition

11

$\mathfrak{g} \subset \mathfrak{g}$ Lie subalg $\rightarrow \mathfrak{g} \curvearrowright \mathfrak{g}$ adjoint action

\mathfrak{g} is a Cartan subalg

- \Leftrightarrow def
- (1) The action of any element of \mathfrak{g} on \mathfrak{g} is semisimple i.e. diagonalizable
 - (2) \mathfrak{g} is maximal among subalgebras satisfying (1). //

Prop

$\mathfrak{g} = \text{CSA} \Rightarrow \mathfrak{g}$ is maximal abelian

Ex

$\mathfrak{sl}_n(\mathbb{C}) \supset \{\text{diagonal matrices}\}$

is a Cartan subalg.

For $\alpha \in \mathfrak{g}^*$ define

$$\mathfrak{g}_\alpha := \{x \in \mathfrak{g} \mid \forall h \in \mathfrak{g} \quad [h, x] = \alpha(h)x\}$$

Then

$$\mathfrak{g} = \bigoplus_{\alpha \in \mathfrak{g}^*} \mathfrak{g}_\alpha$$

$$\text{and } \mathfrak{g} = \mathfrak{g}_0$$

(\because max. abelian)

$$\Delta := \{ \alpha \in \mathfrak{g}^* \mid \mathfrak{g}_\alpha \neq 0 \} \setminus \{0\} \quad 12$$

We call an element of Δ a root.

$$\mathfrak{g} = \mathfrak{g} \oplus \bigoplus_{\alpha \in \Delta} \mathfrak{g}_\alpha \quad \text{root sp decomposition}$$

Ex $\mathfrak{g} = \mathfrak{sl}_3(\mathbb{C})$

basis: $\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ upper triangular

$\mathfrak{h}_1 = \begin{pmatrix} 1 & & \\ & -1 & \\ & & 0 \end{pmatrix} \quad \mathfrak{h}_2 = \begin{pmatrix} 0 & & \\ & 1 & \\ & & -1 \end{pmatrix}$ diag

$\begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}$ lower tri

$\mathfrak{g} = \mathbb{C}\mathfrak{h}_1 \oplus \mathbb{C}\mathfrak{h}_2 \oplus \dots$ is a CSA.

$$[\mathfrak{h}_1, E_{\alpha_j}] = [E_{11}, E_{\alpha_j}] - [E_{22}, E_{\alpha_j}]$$

$$= (\delta_{1\alpha_j} E_{1j} - \delta_{1j} E_{\alpha_1}) - (\delta_{2\alpha_j} E_{2j} - \delta_{2j} E_{\alpha_2})$$

$$= \begin{cases} (\alpha_j) = (1,2) & (2,3) & (1,3) \\ 2E_{12} & -E_{23} & E_{13} \\ (2,1) & (3,2) & (3,1) \\ -2E_{21} & E_{32} & -E_{31} \end{cases}$$

$$[A_2, E_{\bar{j}}]$$

$$= \begin{pmatrix} -E_{12} & 2E_{23} & E_{13} \\ E_{21} & -2E_{32} & -E_{31} \end{pmatrix}$$

$$\left([E_{22}, E_{\bar{j}}] - [E_{33}, E_{\bar{j}}] \right) = (\delta_{2j} E_{2j} - \delta_{3j} E_{3j}) - (\delta_{3j} E_{3j} - \delta_{2j} E_{2j})$$

Define $E_1, E_2, E_3 \in \mathfrak{g}^*$ by

$$E_{\bar{n}} : h \in \mathfrak{g} \mapsto (\bar{n}, \bar{n})\text{-entry of } h.$$

$$(\longrightarrow E_1 + E_2 + E_3 = 0)$$

$$\longrightarrow [A, E_{\bar{j}}] = (E_{\bar{n}} - E_{\bar{j}})(A) E_{\bar{j}}$$

$$\therefore \Delta = \{ E_{\bar{n}} - E_{\bar{j}} \mid (\bar{n} \neq \bar{j}) \}$$

Similar for $\mathfrak{sl}_n(\mathbb{C})$

$$h_1 = \begin{pmatrix} 1 & & \\ & -1 & \\ & & \ddots \end{pmatrix}, \dots, h_{n-1} = \begin{pmatrix} & & \\ & 1 & \\ & & \ddots \end{pmatrix}, E_1, \dots, E_n \in \mathfrak{g}^*$$

In this example

$$\bullet \alpha = E_{\bar{n}} - E_{\bar{j}} \in \Delta \Rightarrow \mathfrak{g}_{\alpha} = \mathbb{C} E_{\bar{j}} \text{ 1-dim'l}$$

• We can divide Δ into

$$\Delta^+ = \{ E_{\bar{n}} - E_{\bar{j}} \mid (\bar{n} > \bar{j}) \} \text{ and } \Delta^- = \{ E_{\bar{n}} - E_{\bar{j}} \mid (\bar{n} < \bar{j}) \}.$$

\longleftrightarrow upper tri.

\longleftrightarrow lower tri

These hold for general s.s. Lie alg 14

Prop $\dim \mathfrak{g}_\alpha = 1$ for $\forall \alpha \in \Delta$

Thm $n := \dim \mathfrak{g}$

$\exists \Pi = \{\alpha_1, \dots, \alpha_n\} \subset \Delta \subset \mathfrak{g}^*$ s.t.

- Π is a \mathbb{Q} -basis of \mathfrak{g}^*
- $\forall \alpha \in \Delta$ can be written as

$$\alpha = \sum_{i=1}^n m_i \alpha_i \quad \text{with} \quad \begin{cases} m_i \in \mathbb{Z}_{\geq 0} & \alpha_i \text{ — (+)} \\ \text{or} \\ m_i \in \mathbb{Z}_{\leq 0} & \alpha_i \text{ — (-)} \end{cases}$$

//

We call an element of

$\begin{cases} \Pi & \text{a simple root} \\ (+) & \text{a positive root} \\ (-) & \text{a negative root} \end{cases}$

$$\Delta = \Delta^+ \sqcup \Delta^-$$

the set of the set of
pos. roots neg. roots

We have $\Delta^- = -\Delta^+$
(property of roots: $\alpha \in \Delta \Rightarrow -\alpha \in \Delta$)

$$\mathfrak{g} = \mathfrak{h} \oplus \bigoplus_{\alpha \in \Delta} \mathfrak{g}_{\alpha}$$

15

$$= \left(\bigoplus_{\alpha \in \Delta^-} \mathfrak{g}_{\alpha} \right) \oplus \mathfrak{h} \oplus \left(\bigoplus_{\alpha \in \Delta^+} \mathfrak{g}_{\alpha} \right)$$

\Downarrow \mathfrak{n}^- \Downarrow \mathfrak{n}^+

triangular decomp.

(cf. $\mathfrak{sl}_2 = \mathbb{C}f \oplus \mathbb{C}h \oplus \mathbb{C}e$)

$\mathfrak{b} := \mathfrak{h} \oplus \mathfrak{n}^+$ is called a Borel subalg
(Borel $\stackrel{\text{def}}{\Leftrightarrow}$ maximal solvable subalg)

Ex $\mathfrak{sl}_n(\mathbb{C})$

We can take

$$\Pi = \{ \varepsilon_1 - \varepsilon_2, \varepsilon_2 - \varepsilon_3, \dots, \varepsilon_{n-1} - \varepsilon_n \}$$

$$\Rightarrow \Delta^+ = \{ \varepsilon_{\tilde{n}} - \varepsilon_{\tilde{r}} \mid \tilde{n} > \tilde{r} \}$$

We can also take

$$\Pi = \{ \varepsilon_2 - \varepsilon_1, \varepsilon_3 - \varepsilon_2, \dots, \varepsilon_n - \varepsilon_{n-1} \} \text{ for example.}$$

\Rightarrow pos. roots and neg. roots are interchanged.

Rem

16

Notion of roots depends on the choice of CDA.

Notion of pos. roots depends on the neg.

choice of simple roots.

Killing form, Weyl group, Cartan matrix

$$\text{ad} : \mathfrak{g} \rightarrow \text{End}(\mathfrak{g})$$

Define $(,) : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{C}$ by

$$(x, y) := \text{tr ad}_x \text{ad}_y \quad \text{Killing form}$$

$(,)$ is a nondegenerate sym bil. form.

(For $\mathfrak{g} = \text{fin. dim. Lie alg}$

$\mathfrak{g} = \text{s.s.} \iff \text{Killing form is nondeg}$)

$(,)|_{\mathfrak{g}}$ is nondeg.

$$\leadsto \quad \tau : \mathfrak{g} \xrightarrow{\sim} \mathfrak{g}^*$$

$(,)$ on \mathfrak{g}^* is defined via τ .

For $\alpha \in \Delta$, define the coroot $\alpha^\vee \in \mathfrak{g}$ by

$$\alpha^\vee := \frac{2}{(\alpha, \alpha)} \bar{z}^1(\alpha)$$

$$(\rightarrow \langle \alpha^\vee, \beta \rangle = \frac{2(\alpha, \beta)}{(\alpha, \alpha)} \quad \alpha, \beta \in \Delta)$$

For each $\alpha \in \Delta$, we can take

$e_\alpha \in \mathfrak{g}_\alpha, f_\alpha \in \mathfrak{g}_{-\alpha}$ satisfying \mathfrak{sl}_2 -rel:

$$[e_\alpha, f_\alpha] = \alpha^\vee, \quad [\alpha^\vee, e_\alpha] = 2e_\alpha, \quad [\alpha^\vee, f_\alpha] = -2f_\alpha$$

For $\alpha \in \Delta$, define $S_\alpha \in \text{End}(\mathfrak{g}^*)$ by

$$S_\alpha(\beta) := \beta - \langle \alpha^\vee, \beta \rangle \alpha$$

The Weyl group W is a ~~finite~~ subgroup of $\text{GL}(\mathfrak{g}^*)$ gen by $\{S_\alpha \mid \alpha \in \Delta\}$.

Ex $\mathfrak{g} = \mathfrak{sl}_n(\mathbb{C}) \Rightarrow W \cong \mathfrak{S}_n$ //

W is gen by $\Delta = \{S_{\alpha_i} \mid \alpha_i \in \Pi\}$.

(W, Δ) is a finite Coxeter group.

Classification of s.s. Lie algebras? 18

$(\mathfrak{g}, \mathfrak{g}) \rightsquigarrow \Delta$ & satisfies the axioms of "root system"

→ reduced to classification of root systems.

$\Pi = \{\alpha_i \mid i \in I\} \subset \Delta$ simple roots

$$a_{ij} := \langle \alpha_i^\vee, \alpha_j \rangle = \frac{2(\alpha_i, \alpha_j)}{(\alpha_i, \alpha_i)}$$


$(a_{ij})_{i,j \in I}$: Cartan matrix of \mathfrak{g} .

Semistable Lie algebras are determined by their Cartan matrices.

Ex $\mathfrak{sl}_n(\mathbb{C})$, $\Pi = \{\alpha_i = \epsilon_i - \epsilon_{i+1} \mid i = 1, \dots, n-1\}$

$$(a_{ij}) = \begin{pmatrix} 2 & -1 & & 0 \\ -1 & 2 & -1 & \\ & \ddots & \ddots & \\ 0 & & -1 & 2 \end{pmatrix}$$

Dynkin diagram



type A_{n-1}

Simple Lie algebras? / \cong

$\xleftrightarrow{(\cong)}$ $A_n, B_n, C_n, D_n, E_6, E_7, E_8, F_4, G_2$

name of root system

//

2. Representation theory

Fix a triangular decomp $\mathfrak{g} = \mathfrak{n} \oplus \mathfrak{g} \oplus \mathfrak{n}^+$

$$\Rightarrow U(\mathfrak{g}) \cong U(\mathfrak{n}) \underset{\otimes}{\otimes} U(\mathfrak{g}) \underset{\otimes}{\otimes} U(\mathfrak{n}^+)$$

PBW

as \mathbb{C} -vec sp

For a \mathfrak{g} -module M and $\lambda \in \mathfrak{g}^*$ define

$$M_\lambda := \{v \in M \mid \forall h \in \mathfrak{g} \quad hv = \lambda(h)v\}$$

λ is called a weight of M if $M_\lambda \neq 0$.

Prop

$M = \text{fin. dim.} \Rightarrow \mathfrak{g} \curvearrowright M$ semisimple

$$\text{true.} \quad M = \bigoplus_{\lambda \in \mathfrak{g}^*} M_\lambda$$

//

Rem

$M = \bigoplus_{\lambda \in \mathfrak{g}^*} M_\lambda$ does not hold in general.

Notice:

$$v \in M_\lambda, \quad x \in \mathfrak{g}_\alpha \quad (\alpha \in \Delta)$$

$$\Rightarrow xv \in M_{\lambda+\alpha} \quad (\text{similar as sl}_2\text{-case})$$

$$\therefore [h, x] = \alpha(h)x$$

$$h(xv) = [h, x]v + xhv$$

$$= \alpha(h)xv + \lambda(h)xv$$

$$= (\lambda + \alpha)(h)xv \quad \square$$

Prop

20

M : fin. dim. simple \mathfrak{g} -module

$\Rightarrow \exists \lambda \in \mathfrak{g}^*$ s.t.

- $\dim M_\lambda = 1$

- $n^+ M_\lambda = 0$.

Proof

existence:

M = fin. dim. \Rightarrow the set of weights is finite

$\Rightarrow \exists \lambda$ s.t. $M_\lambda \neq 0$ and $\forall \alpha \in \Delta^+, M_{\lambda+\alpha} = 0$.

$\Rightarrow n^+ M_\lambda = 0$.

Take $v_{\neq 0} \in M_\lambda \Rightarrow M = U(\mathfrak{g})v$

since M is simple

$$M = U(n^-) \underbrace{U(\mathfrak{g}) U(n^+)}_{\text{acts by scalar}} v$$

$$= U(n^-)v$$

Recall $n^- = \bigoplus_{\alpha \in \Delta^-} \mathfrak{g}_\alpha$ $\dim \mathfrak{g}_\alpha = 1$

Let $\Delta^- = \{\beta_1, \dots, \beta_r\}$ and take

nonzero $f_{\beta_i} \in \mathfrak{g}_{\beta_i}$ for each β_i .

$\rightarrow \{ f_{\beta_1}^{m_1} \dots f_{\beta_r}^{m_r} \mid m_i \geq 0 \}$ is a \mathbb{Q} -basis of $\mathcal{U}(n)$

$$\therefore \mathcal{U}(n)v = \text{span}_{\mathbb{Q}} \{ \underbrace{f_{\beta_1}^{m_1} \dots f_{\beta_r}^{m_r}}_{M_{\lambda + \sum_{j=1}^r m_j \beta_j}} v \mid m_j \geq 0 \}$$

$$\therefore \dim M_{\lambda} = 1 //$$

uniqueness:

Suppose λ_1 and λ_2 satisfy the condition.

$$\Rightarrow \lambda_2 = \lambda_1 + \sum_{i=1}^r m_i \beta_i \quad \forall m_i \in \mathbb{Z}_{\geq 0}$$

$$\beta_i \in \sum_{j=1}^n \mathbb{Z}_{\leq 0} \alpha_j$$

$$\lambda_2 - \lambda_1 \in \sum_{j=1}^n \mathbb{Z}_{\leq 0} \alpha_j$$

$$\text{We also have } \lambda_1 - \lambda_2 \in \sum_{j=1}^n \mathbb{Z}_{\leq 0} \alpha_j$$

$$\therefore \lambda_1 = \lambda_2 \quad \square$$

Highest weight module

Looking at the proof of Prop.

"the set of weights is finite"

\leadsto "bounded from above" is enough.

Def

22

Define a partial order \geq on \mathfrak{g}^* by

$$\lambda \geq \mu \stackrel{\text{def}}{\iff} \lambda - \mu \in \sum_{i=1}^n \mathbb{Z}_{\geq 0} \alpha_i \quad //$$

M : \mathfrak{g} -module

$v_{\neq 0} \in M$ is maximal vector of wt λ

$$\stackrel{\text{def}}{\iff} v \in M_{\lambda} \text{ and } \mathfrak{n}^+ v = 0.$$

M is a highest weight module with highest weight λ

$$\stackrel{\text{def}}{\iff} M = U(\mathfrak{g})v \text{ for a maximal vector } v \text{ of wt } \lambda$$

v is called a highest weight vector //

Prop \rightarrow fin. dim. simple \mathfrak{g} -module
is a h.w. module.

Assume $M = \text{h.w. module}$ with h.w. λ

$$(1) \quad M = \bigoplus_{\mu \leq \lambda} M_{\mu}$$

$$\dim M_{\mu} < \infty \quad \forall \mu \quad \text{and} \quad \dim M_{\lambda} = 1$$

(2) M has a unique simple quotient.
(i.e. simple top)

In particular, M is indecomposable.

Proof

(1) Start as for dual case

(2) Let $M' \subsetneq M$ be a proper submodule

$$\Rightarrow M'_{\lambda} = 0$$

$$\left(\begin{array}{l} \because \dim M_{\lambda} = 1, \\ M'_{\lambda} = M_{\lambda} \cap M' \neq 0 \\ \Rightarrow M_{\lambda} \subset M' \\ \Rightarrow M = \bigcup_{\mu} M_{\mu} \subset M' \\ \Rightarrow M = M' \end{array} \right)$$

$\therefore \sum_{\text{all proper sub}} M'$ is also a proper submodule.

This gives a unique maximal submodule.



Q. Are there h.w. modules other than fin. dim'l ones?

24

→ Yes.

Verma module

$\lambda \in \mathfrak{g}^* \leadsto \mathbb{C}_\lambda := 1\text{-dim'l rep of } \mathfrak{g}$

$$\mathbb{C}_\lambda \cong \mathbb{C} \ni 1 \quad \hbar 1 = \lambda(\hbar)1$$

Extend \mathbb{C}_λ to a \mathfrak{b} -module by

$$\mathfrak{b} = \mathfrak{g} \oplus \mathfrak{n}^+ \longrightarrow \mathfrak{g} \curvearrowright \mathbb{C}_\lambda$$

$$\text{i.e. } \mathfrak{n}^+ 1 = 0.$$

$$M(\lambda) := \underbrace{\mathcal{U}(\mathfrak{g})}_{\mathcal{U}(\mathfrak{b})} \otimes \mathbb{C}_\lambda \cong 1 \otimes 1 = \mathbb{C}_\lambda$$

Verma module

$M(\lambda)$ is a h.w. module with h.w. vector v_λ

$$\text{PBW: } \mathcal{U}(\mathfrak{g}) \cong \mathcal{U}(\mathfrak{n}^-) \underset{\mathbb{C}}{\otimes} \mathcal{U}(\mathfrak{b})$$

→ $M(\lambda) \cong \mathcal{U}(\mathfrak{n}^-)$ as \mathbb{C} -vec sp.

$M(\lambda)$ is universal among h.w. modules ²⁵

T.Q. If M : h.w. module with
h.w. vector v of wt λ .

$\Rightarrow \exists$ surjective \mathfrak{g} -module hom

$$M(\lambda) \longrightarrow M$$

$$v_\lambda \longmapsto v$$

$$\therefore \text{Hom}_{U(\mathfrak{g})}(M(\lambda), M)$$

$$\cong \text{Hom}_{U(\mathfrak{g})}(\mathbb{C}_\lambda, M)$$

$$\cong \{m \in M_\lambda \mid m^+ m = 0\} \quad \square$$

Denote the unique simple quotient
of $M(\lambda)$ by $L(\lambda)$.

Thm

(1) $L(\lambda)$ is fin. dim'l.

$$\Leftrightarrow \forall \alpha \in \Delta^+ \quad \lambda(\alpha^\vee) \in \mathbb{Z}_{\geq 0}$$

\uparrow

This λ is said to be dominant
integral

(2) { fin. dim. simple \mathfrak{g} -module } / \cong

$$\stackrel{1:1}{\longleftrightarrow} \{ \lambda : \text{dominant integral?} \}$$

$$L(\lambda) \longleftarrow \lambda$$

(3) The category of fin. dim'l \mathfrak{g} -modules is semisimple.

26

Ex $\mathfrak{sl}_2 = \mathbb{C}f \oplus \mathbb{C}h \oplus \mathbb{C}e$

$\alpha := e_1 - e_2 \quad \Delta = \{ \pm \alpha \} \supset \Pi = \{ \alpha \}$

$h = \alpha^\vee$

$\mathfrak{g} = \mathbb{C}h \longleftrightarrow \mathfrak{g}^* = \mathbb{C}\omega \cong \mathbb{C}$
dual basis $\omega \longleftrightarrow 1$

$\alpha = 2\omega \longleftrightarrow 2$

order: $\lambda, \mu \in \mathbb{C} \quad \lambda \geq \mu \Leftrightarrow \lambda - \mu \in 2\mathbb{Z}_{\geq 0}$

$U(\mathfrak{n}^-) = \mathbb{C}[f] = \bigoplus_{k \geq 0} \mathbb{C}f^k$

$M(\lambda) = \bigoplus_{k \geq 0} \mathbb{C} \underbrace{f^k v_\lambda}_{\text{wt } \lambda - 2k}$

* $k \geq 1$, condition for $f^k v_\lambda = 0$?

$f^k v_\lambda = [e, f^k] v_\lambda + \underbrace{f^k e v_\lambda}_{=0}$

$[e, f^k] = k f^{k-1} h - k(k-1) f^{k-1}$

$[e, f^k] v_\lambda = k(\lambda - (k-1)) f^{k-1} v_\lambda$

$\stackrel{?}{=} 0 \Leftrightarrow \lambda = k-1$

∴

$$\lambda \notin \mathbb{Z}_{\geq 0} \Rightarrow M(\lambda) \text{ is simple}$$

$$\lambda \in \mathbb{Z}_{\geq 0} \Rightarrow M(\lambda) \neq U(\mathfrak{g}) f^{\lambda+1} v_{\lambda}$$

max proper submodule

$$L(\lambda) = M(\lambda) / \underbrace{U(\mathfrak{g}) f^{\lambda+1} v_{\lambda}}_{\cong L(-\lambda-2)} \quad \lambda+1 - \text{dim } L$$

$$f^{\lambda+1} v_{\lambda}$$

$$\text{wt } \lambda - 2(\lambda+1) = -\lambda-2$$

$$0 \rightarrow L(-\lambda-2) \rightarrow M(\lambda) \rightarrow L(\lambda) \rightarrow 0$$

$$\text{if } \lambda \in \mathbb{Z}_{\geq 0} //$$

In general

$$[M(\lambda) : L(\lambda)] = 1$$

$$[M(\lambda) : L(\mu)] \neq 0 \Rightarrow M(\lambda)_{\mu} \neq 0$$

$$D_{\lambda\mu}$$

$$\Rightarrow \mu \leq \lambda$$

(M(λ) : finite length)

D = (D_{λμ}) upper triangular matrix

In Grothendieck grp

$$[M(\lambda)] = \sum_{\mu \leq \lambda} D_{\lambda\mu} [L(\mu)]$$

$$\leadsto [L(\lambda)] = \sum_{\mu \leq \lambda} (D^{-1})_{\lambda\mu} [M(\mu)]$$

Kazhdan-Lusztig conjecture:

28

(for integral weights)

$$[M(\lambda) : L(\mu)] = P_{y,w}(1)$$

where $P_{y,w}(q) \in \mathbb{Z}[q]$ is KL polynomial attached to some elements $y, w \in W$.

y, w are determined by λ, μ .

KL conj: 1979 (introduce KL poly via Hecke alg)

Solved by Berenstein-Bernstein (1981)
Brylinski-Kashiwara

Use D-modules
pervershe sheaves) on flag variety

(2012-) 2014 algebraic proof by
Elias-Wilkinson

(inspired by geometric method)

Category \mathcal{O}

The category of all \mathfrak{g} -modules is too big.
Hopeless to classify simples etc.

→ Study a good full subcategory.

Def (Bernstein-Gelfand-Gelfand '76)

$\mathcal{O} \subset \mathfrak{g}$ -modules full subcat.

$M \in \mathcal{O} \stackrel{\text{def}}{\iff} \bullet$ finitely generated over $U(\mathfrak{g})$

\bullet \mathfrak{g} -semisimple

$\bullet \forall v \in M$

$$\dim U(\mathfrak{n}^+)v < \infty$$

(locally \mathfrak{n}^+ -finite) //

\mathcal{O} is closed by taking sub, quot,

fin. direct sum.

$U(\mathfrak{g})$: noetherian

→ abelian category

Note: \mathcal{O} is not closed by extension.

↗ failure of \mathfrak{g} -semisimplicity.

Prop(1) \forall h.w. module $\in \mathcal{O}$ In particular $M(\lambda), L(\lambda) \in \mathcal{O}$ (2) $M \in \mathcal{O}$ $\Rightarrow \exists \lambda_1, \dots, \lambda_r \in \mathfrak{g}^*$ weights of $M \subset \bigcup_{i=1}^r \{ \mu \in \mathfrak{g}^* \mid \mu \leq \lambda_i \}$ (3) $\{ \text{Simple } \tau \in \mathcal{O} \} / \cong \xleftrightarrow{|\cdot|} \mathfrak{g}^*$ $L(\lambda) \longleftrightarrow \lambda$ Proof(1) • fin. gen \leftarrow gen by h.w. vec• \mathfrak{g} -s.s. \leftarrow done• loc n^+ -fin.

$$M = \bigoplus_{\mu \leq \lambda} M_\mu \quad \lambda = \text{h.w.}$$

 $v \in M_\mu \Rightarrow$ wt of $\tau(n^+)v$ satisfy $\mu \leq \tau \leq \lambda$

of such wts is finite

 $\& \forall$ wt sp are fin. dim'l \rightarrow OK //

(2) Take a finite set of generators 31

$$S \subset M$$

$$\text{loc. fin.} \rightarrow \dim U(n^+)S < \infty$$

We can take maximal weights

$$\lambda_1, \dots, \lambda_r \text{ of } U(n^+)S \text{ w.r.t. } \geq.$$

$$\rightarrow M = U(\mathfrak{g})S$$

$$= U(n^-)U(\mathfrak{g})U(n^+)S$$

$$\text{wt } \subset \bigcup_{|\vec{\alpha}| \geq 1} \{ \mu \leq \lambda_i \} //$$

(3) Let M be a simple \mathfrak{u} in \mathcal{O} .

(2) \Rightarrow M has a maximal vector v
of wt λ

$$\text{simple} \Rightarrow M = U(\mathfrak{g})v$$

$\therefore M$ is a h.w. module with h.w. λ

$$\therefore M(\lambda) \rightarrow M \\ \cong L(\lambda) \quad \square$$

Prop

- (1) \mathcal{O} is finite length
 (2) \mathcal{O} has enough projectives
 (Use central character etc. for a proof)

Rem

\mathcal{O} has finitely many simples.

But each block has finitely many simples.

Hence each block of \mathcal{O} is Morita equiv
 to some fin. dim. alg.

$P(\lambda) :=$ projective cover of $L(\lambda)$.

Thm $\forall \lambda \in \mathfrak{g}^*$

$\exists \mathcal{O}$ -module filtration

$$P(\lambda) = M_0 \supsetneq M_1 \supsetneq \dots \supsetneq M_d = 0$$

s.t. $M_0/M_1 \cong M(\lambda)$

$$M_{\bar{n}}/M_{\bar{n}+1} \cong M(\lambda_{\bar{n}}) \quad \exists \lambda_{\bar{n}} > \lambda$$

$$\bar{n} = 1, \dots, d-1 //$$

Thm. \mathcal{O} is a h.w. cat. whose standard modules are Verma modules.

Its poset is (\mathcal{G}^*, \geq) .

By CPS thm, the fin. dim alg which is Morita equiv to a block of \mathcal{O} is quasi-hereditary.

BGG reciprocity

duality:

\exists exact contravariant functor $v: \mathcal{O} \rightarrow \mathcal{O}$ s.t.

- $M^{vv} \cong M$
- $L(\lambda)^v \cong L(\lambda)$

$\Rightarrow M(\lambda)^v$ is a costandard module of \mathcal{O} .

$[M(\lambda)^v : L(\mu)] = [M(\lambda) : L(\mu)]$ holds.

Cor of Thm

$$(P(\lambda) : M(\mu)) = [M(\mu) : L(\lambda)]$$

BGG reciprocity

The reciprocity for a general h.w. at
TS $(P(\lambda) : M(\mu)) = [M(\mu)^V : L(\lambda)]$. //

$$C_{\lambda\mu} := [P(\lambda) : L(\mu)]$$

$$D_{\lambda\mu} := [M(\lambda) : L(\mu)]$$

$$C = (C_{\lambda\mu}) \quad \text{Cartan matrix}$$

$$D = (D_{\lambda\mu}) \quad \text{decomposition matrix} \\ \& \text{ upper tri.}$$

$$C_{\lambda\mu} = \sum_{\nu} (P(\lambda) : M(\nu)) \underset{\text{"BGG"}}{[M(\nu) : L(\mu)]} \\ [M(\nu) : L(\lambda)]$$

$$\therefore C = {}^t D \cdot D$$

In particular
symmetric matrix //

Ex sl_2

35

$\mathcal{O}_0 :=$ the block containing the trivial rep $L(0)$

stamples $L(0)$ $L(-2)$ $\subset \mathcal{O}$
 $0 > -2$

$M(0):$ basis v_0 $f v_0$ $f^2 v_0 \dots$

 $\text{sub} \cong L(-2)$
 $\cong M(-2)$

$$[M(0) : L(0)] = 1$$

$$[M(0) : L(-2)] = 1$$

$$[M(-2) : L(0)] = 0$$

$$[M(-2) : L(-2)] = 1$$

projectives?

$0 \cong$ maximal in $so, -2?$

$$\rightarrow P(0) \cong M(0)$$

$$P(-2): (P(-2) : M(-2)) = 1$$

$$\text{By BGG} \quad (P(-2) : M(0)) = [M(0) : L(-2)] = 1$$


$$\therefore 0 \rightarrow M(0) \rightarrow P(-2) \rightarrow M(-2) \rightarrow 0$$

$P := P(0) \oplus P(-2)$ proj generator 36

$\text{End}_{\mathcal{O}} P = ?$

- $\text{Hom}_{\text{M}(0)}(P(0), P(-2)) = \mathbb{Q}a$ $M(0) \hookrightarrow P(-2)$
- $\text{Hom}(P(-2), P(0)) = \mathbb{Q}b$ $P(-2) \rightarrow M(-2) \cong L(-2) \hookrightarrow M(0)$
 $\rightarrow b \circ a = 0$
- $\text{Hom}_{\text{M}(0)}(P(0), P(0)) = \mathbb{Q}e_0$ corresp. to idempotent
- $\text{Hom}(P(-2), P(-2)) = \mathbb{Q}e_{-2} \oplus \mathbb{Q}a \circ b$

$\dim \text{End}_{\mathcal{O}} P = 5$



rel: $ab=0$
 $(\Leftrightarrow b \circ a = 0)$

$\mathcal{O}_0 \cong \text{mod} - \text{End}_{\mathcal{O}} P$

cat. of fin. dim. right modules

(*)

37

Apply $\text{Hom}(P(-2), -)$ to

$$0 \rightarrow M(0) = P(0) \rightarrow P(-2) \rightarrow M(-2) \rightarrow 0$$

$$0 \rightarrow (\text{Hom}(P(-2), P(0))) = \mathbb{Q}b$$

$$\rightarrow \text{Hom}(P(-2), P(-2)) \quad \downarrow$$

$$a \circ b$$

$$\rightarrow \text{Hom}(P(-2), M(-2)) = \mathbb{Q}[P(-2) \rightarrow M(-2)]$$

$$\rightarrow 0$$

$$\therefore \dim \text{Hom}(P(-2), P(-2)) = 2 \quad \square$$

\mathcal{G} : fin. dim. ss.

38

$$\rho := \frac{1}{2} \sum_{\alpha \in \Delta^+} \alpha$$

$\mathcal{G}_0 :=$ the block of \mathcal{G} containing $L(0)$

\rightarrow simples of \mathcal{G}_0 are $L(-w\rho - \rho)$
 $w \in W$

We often use "Bruhat order" on W
instead of \geq .

KL conj:

$$[M(-w\rho - \rho) : L(-w\rho - \rho)] \\ = P_{w_0 w, w_0} (1)$$

w_0 : longest element of W

Roigel: 90

$$P := \bigoplus_{w \in W} P(-w\rho - \rho), \quad L := \bigoplus_{w \in W} L(-w\rho - \rho)$$

$$\text{End}_{\mathcal{G}} P \cong \text{Ext}_{\mathcal{G}}^{\bullet}(L, L)$$

and it is Koszul.

given & rel

Vertex set = W

$$\dim \operatorname{Ext}_{\mathcal{O}}^1(L(-yp-p), L(-wp-p)) \\ = \text{coeff of } q^{\frac{1}{2}(\ell(w)-\ell(y)-1)} \text{ in } P_{y,w}(q)$$

if $y < w$ ($\ell(w)$ = length of w)

(\because radical layer of Verma modules)
is captured by KL poly

$$\text{By duality, } \operatorname{Ext}_{\mathcal{O}}^1(L(\lambda), L(\mu)) \\ \cong \operatorname{Ext}_{\mathcal{O}}^1(L(\mu), L(\lambda))$$

$\therefore y \Leftrightarrow w$ in the given.

rel: \exists algorithm to compute relations

by Stroppel 03

Vybornov 07

using Soergel's work.