```
{"2":"1","3":"L15E-1677898349435","4":{"fv":"0"},"5":[{"3":"9582-1677840728986","4":
{"I":"title"},"5":[{"2":"2","3":"80tG-1677898349435","7":[{"8":"地图结构体"}]}],"6":"h"},{"3":"5040-
1677857973283","5":[{"2":"2","3":"llyj-1677898349435"}]},{"3":"8356-1677857973411","5":
[{"2":"2","3":"WMQt-1677898349435"}]},{"3":"6193-1677857973570","5":[{"2":"2","3":"JCEf-
1677898349435"}]},{"3":"8957-1677857973938","4":
{"u":"DFF765DD155843A485C92451DF2C41C2","w":620,"h":555},"6":"im"},{"3":"9943-
1677857973938","4":{"I":"h2"},"5":[{"2":"2","3":"LA3L-1677898349435","7":[{"8":"思
想: "}]}],"6":"h"},{"3":"7730-1677859215117","4":{"l":"h2"},"5":[{"2":"2","3":"Fo4t-
1677898349435","7":[{"8":"x轴走到10步,就会往下走2步,在往回走10步,步长从0开始算,也就是x
初始会走10步,在往下走2步,然后又往回走10步,又往下走2步,。。。。。"}]}],"6":"h"},{"3":"1040-
1677859378860","4":{"l":"h2"},"5":[{"2":"2","3":"8TTI-1677898349435"}],"6":"h"},{"3":"2699-
1677859382021","4":{"l":"h2"},"5":[{"2":"2","3":"ydPX-1677898349435","7":[{"8":"所以得出结论,每
走10步,下2步,在走10步,下2步,走10步要设置^运算之类的操作,因为之后2个选择:前、后,而且
是一次前下一次必然是相反的结果: "}]}],"6":"h"},{"3":"2622-1677859840105","4":
{"u":"C5DE092F863B46DFAF88220B2E96A692","w":620,"h":775},"6":"im"},{"3":"4758-
1677860635351", "5":[{"2": "2", "3": "phg1-1677898349435"}]}, {"3": "8165-1677860635351", "4":
{"u":"16A09B96A8D74D8E9186EF3BC03C2270","w":348,"h":103},"6":"im"},{"3":"3436-
1677860634991","4":{"l":"h2"},"5":[{"2":"2","3":"NFsO-1677898349435","7":
[{"8":"output:"}]}],"6":"h"},{"3":"6279-1677859846235","4":
{"u":"38EE9D094CE3489AAF1E94838A9D1B22","w":433,"h":536},"6":"im"},{"3":"1310-
1677859846235", "5":[{"2": "2", "3": "9xvq-1677898349435"}]}], compress ":true}
```