Mycology media

Huang MEGA Lab (NCKU)
Lab plain-PDF edition, rebuilt 2026-10-21

Source of record: Huang MEGA Lab (NCKU). Originally authored on protocols.io (dx.doi.org/10.17504/protocols.io.e6nvw53 created 2022-05-16, version 1. This PDF is the lab's self-hosted copy and supersedes the protocols.io export.

Abstract

Mycology media

Procedure

PROCEDURE

1. WA+C agar

Water agar + chloramphenicol (氯黴素 kill bacteria): As an environmental isolating media

Component	Amount
Agar	20g
Chloramphenicol	0.05g
DI water	**1000 ml**

chloramphenicol is autoclavable

2. PDA+C agar

potato dextrose agar + chloramphenicol (氯黴素 kill bacteria): Use when WA+C agar is not working

Component	Amount
PDA	39g (or per instruction on label)
Agar	5g
Chloramphenicol	0.05g
DI water	**1000 ml**

5. Emerson YpSs Agar: For chytrids

Component	Amount
Yeast extract	4g
Soluble Starch	15g
K2HPO4	1g
MgSO4 x 7H2O	0.5g
Agar	20g
DI water	**1000ml**

Adujust pH to 7.0 +/- 0.2. Autoclave at 121ºC.

6. 10% Unclarified V8 Agar

1. Combine 100 ml V8 juice and 900 ml of distilled water.
2. Add 1g $CaCO_3$and 0.05 g b-sitosterol and mix well. Leave stir bar in the flask for later mixing.
3. Add 15g of agar.
4. Autoclave at 15 psi for 20 minutes.
5. Stir medium while dispensing to insure good mixing of $CaCO_3$. 20% : 200 ml V8, 800ml d. water, 2g $CaCO_3$, and 0.05g b-sitosterol.

15% : 150 ml V8, 850 ml d. water, 1.5g $CaCO_3$, and 0.05g b-sitosterol.

Reference: Miller, P. M. 1955. V-8 juice agar as a general purpose medium for fungi and bacteria. Phytopathology 45:461-462.

---

7. SceSel+ medium (Kaltseis et al.,2009)

For isolation of Scedosporium

Component	Amount
Malt extract	6.25 g
maltose	6.25 g
KH_2PO_4	1.25 g
yeast extract	1 g
Magnesium Sulfate Heptahydrate	0.625 g
soy peptone	0.625 g
chloramphenicol	0.1 g
ciprofloxacin	0.1 g
streptomycin sulfate	0.1 g
dichloran	**2 mg**
benomyl	**6 mg**
DI water	**1000ml**

---

3. quarter-strength PDA

1. Less nutrient than regular PDA (1/4)
2. For beetle-associated fungi isolation

Component	Amount
PDA	9.75g (or per instruction on label)
Agar	5g
DI water	**1000 ml**

---

4. MEA (malt extraction agar)

For isolate subculture

Component	Amount
MEA	33.6g (or per instruction on label)
Agar	5g
DI water	**1000 ml**