# From specimen to sequence: Huang MEGA Lab Nanopore workflow

amplicon (two-step PCR or one-pot PCR) or whole genome · scan or click a QR code to open the protocol · LIMS stages 1 → 2 → 3a → 3b → 4a

**Stage 1** Specimen# / Culture#

**Specimen**
soil / environmental / insect | fungal strain in culture

**LIMS register** `/lims-register` | Stage 1 → Specimen# (soil, environmental) or Culture# (strain)

**isolate fungi (strains, Route B / C)**

soil / environmental sample, or an existing pure culture: extract directly (skip isolation)

**Stage 2** isolate + culture · Culture#

**Microbe isolation**
environmental sample → vial

**Culture media**
MEA, PDA and other recipes

**Isolation and culturing**
primary plate on MEA (Stage 2a) → subculture one colony / hyphal tip on PDA
one Stage-2b row per isolate · preserve in vials
sequence from a pure culture only

**LIMS register** `/lims-register` | Stage 2a plate → Stage 2b isolate → Culture# per strain

**DNA extraction**

**Stage 3a** extraction + Qubit · MolWork#

**Soil DNA extraction**
SuperSoil · Zixpress 32

**Fungal DNA extraction**
modified BOMB v9.0

**DNA extraction**
soil: SuperSoil, Zixpress 32 | fungal / HMW gDNA: BOMB v9
genome route: keep the DNA high-molecular-weight

**quantify**

**Qubit Flex**
concentration of the extract (dsDNA HS)

**LIMS register** `/lims-register` | Stage 3a → MolWork# (minted only after quantification)

choose a route

**Stage 3b** PCR + gel + Qubit · pcr_reaction

**Route A · amplicon, two-step PCR**
environmental samples (soil, water) and mixed templates

**1st PCR (gene regions)**

**2-step PCR protocol**
mixtures + cycling

1st PCR products (per sample)
🟦 head seq — gene region —

**2nd PCR (barcoding)**
gel check #1: good amplicons only

2nd PCR products: one tube per sample, one barcode per tube

**Tube 1** ▶ barcode 1
**Tube 2** ▶ barcode 2
…
**Tube n** ▶ barcode n

▶ barcode 🟦 head seq — gene — 🟦 head seq ◀ barcode

**Route B · amplicon, one-pot PCR**
strains: single- or multi-locus

**Set up one-pot PCR**

One reaction tube (per strain × locus)
**gene-region primers** (head-tailed, 0.1 µM each) + **barcode primers** (barcode + head seq, 0.3 µM each)
50 µL: KAPA HiFi 2X + both primer sets + 3 µL template

**One-pot PCR protocol**
50 µL mix + 25/15 cycles

**One-pot PCR**
phase 1: 25 cycles, 60–66 °C
phase 2: 15 cycles, 65 °C

One-pot PCR products: one tube per sample, one barcode per tube

**Tube 1** ▶ barcode 1
**Tube 2** ▶ barcode 2
…
**Tube n** ▶ barcode n

▶ barcode 🟦 head seq — gene — 🟦 head seq ◀ barcode

**Route C ↓**

**Route C · genome, no PCR (Stage 3a → 4a, no 3b)**

same barcoded molecule from either route

**Gel check → Qubit Flex (dsDNA BR)**
confirm the barcoded product

**LIMS register** `/lims-register` | Stage 3b → pcr_reaction: method, conc., 2nd_pcr_ok, barcode

**Pooling (equal molar)**

**Stage 4a** Nanopore experiment · Nanopore#

**Pool all barcoded samples**
equal molar of every sample

**Size selection 1X**
bead purification protocol

**Purification (bead clean-up)**
PCR sample : bead = 1 : 1

**Library construction**

**EvoPrep library prep**
KAPA EvoPrep NBD114.96

**KAPA EvoPrep library**
PCR-barcoded pool, ~200 fmol
end-prep → adapter ligation → clean-up
PromethION

**QC the gDNA**

**High-molecular-weight gDNA**
integrity: gel / TapeStation
input per ONT table
(10 kb avg → 200 fmol ≈ 1.3 µg)

**Library construction**

**KAPA EvoPrep + NBD114.96**
end-prep → native barcode ligation (NB01–96) → pool
→ adapter ligation
~140 min

**EvoPrep library prep**
KAPA EvoPrep NBD114.96

**LIMS register** `/lims-register` | Stage 4a → Nanopore#: experiment, samples (PCR barcode or NB), flow cell, NanoAmp

**Load the flow cell and sequence**
PromethION R10.4.1 (FLO-PRO114M)

run data (POD5)

**/nanoamp**
**amplicon runs (Routes A and B)**
basecall → demux → trim → cluster → BLAST
run it on the run folder with the LIMS sample sheet

**/nanogenome**
**genome runs (Route C)**
basecall → assemble → annotate → compare
run it on the project directory

**after the run**
(bench)

**Flow cell wash**
EXP-WSH004 PromethION

**Flow cell wash and reuse**
EXP-WSH004

**LIMS register** `/lims-register` | Stage 4a → flow cell: pores after wash, date of use

**Legend**

- 🟦 Head seq: universal tail on the gene-region primers
- ▶ Barcode: sample-specific, on the barcode primers
- — Gene region: target amplicon (colour = sample)

QR code: scan, or click on the website, to open that step's protocol PDF
Left bands = LIMS stages; /lims-register ends every stage; dashed = QC / info
Route C uses native barcodes (NBD114.96) in the library: no PCR, no Stage 3b
/nanoamp, /nanogenome: Claude Code analysis skills for the run data

Huang MEGA Lab (NCKU) · chart built 2026-10-21 from doc/protocols/registry.csv · live version with clickable links: ythuangmyco.github.io/protocol