# Mycology media

Huang MEGA Lab (NCKU)

(Lab plain-PDF edition, rebuilt 2026-10-20)

**Source of record:** Huang MEGA Lab (NCKU). Originally authored on protocols.io (dx.doi.org/10.17504/protocols. created 2022-05-16, version 1. This PDF is the lab's self-hosted copy and supersedes the protocols.io export.

**Abstract**

Mycology media

**Procedure**

**Procedure**

---

**1. WA+C agar**

Water agar + chloramphenicol (氯黴素 kill bacteria): As an environmental isolating media

| Component | Amount |
|---|---|
| Agar | 20g |
| Chloramphenicol | 0.05g |
| DI water | **1000 ml** |

*chloramphenicol is autoclavable*

---

**2. PDA+C agar**

potato dextrose agar + chloramphenicol (氯黴素 kill bacteria): Use when WA+C agar is not working

| Component | Amount |
|---|---|
| PDA | 39g (or per instruction on label) |
| Agar | 5g |
| Chloramphenicol | 0.05g |
| DI water | **1000 ml** |

---

**5. Emerson YpSs Agar**: For chytrids

| Component | Amount |
|---|---|
| Yeast extract | 4g |
| Soluble Starch | 15g |
| K2HPO4 | 1g |
| MgSO4 x 7H2O | 0.5g |
| Agar | 20g |
| DI water | **1000ml** |

*Adujust pH to 7.0 +/- 0.2. Autoclave at 121℃.*

---

**6. 10% Unclarified V8 Agar**

1. Combine 100 ml V8 juice and 900 ml of distilled water.
2. Add 1g CaCO3and 0.05 g b-sitosterol and mix well. Leave stir bar in the flask for later mixing.
3. Add 15g of agar.
4. Autoclave at 15 psi for 20 minutes.
5. Stir medium while dispensing to insure good mixing of CaCO3. 20% : 200 ml V8, 800ml d. water, 2g CaCO3, and 0.05g b-sitosterol.

15% : 150 ml V8, 850 ml d. water, 1.5g CaCO3, and 0.05g b-sitosterol.

Reference: Miller, P. M. 1955. V-8 juice agar as a general purpose medium for fungi and bacteria. Phytopathology 45:461-462.

---

**7. SceSel+ medium** (Kaltseis et al.,2009)

For isolation of Scedosporium

| **Component** | **Amount** |
|---|---|
| Malt extract | 6.25 g |
| maltose | 6.25 g |
| KH2PO4 | 1.25 g |
| yeast extract | 1 g |
| Magnesium Sulfate Heptahydrate | 0.625 g |
| soy peptone | 0.625 g |
| chloramphenicol | 0.1 g |
| ciprofloxacin | 0.1 g |
| streptomycin sulfate | 0.1 g |
| dichloran | **2 mg** |
| benomyl | **6 mg** |
| DI water | **1000ml** |

---

**3. quarter-strength PDA**

1. Less nutrient than regular PDA (1/4)
2. For beetle-associated fungi isolation

| **Component** | **Amount** |
|---|---|
| PDA | 9.75g (or per instruction on label) |
| Agar | 5g |
| DI water | **1000 ml** |

---

**4. MEA** (malt extraction agar)

For isolate subculture

| **Component** | **Amount** |
|---|---|
| MEA | 33.6g (or per instruction on label) |
| Agar | 5g |
| DI water | **1000 ml** |