

# Introduction to Statistical Machine Learning

## Homework 5

Yota Toyama

December 10, 2016

1.

$$\begin{aligned}
 p(y=1|\mathbf{x}) &= \frac{p(\mathbf{x}|y=1)p(y=1)}{p(\mathbf{x})} \\
 &= \frac{p(y=1)}{p(\mathbf{x})} N(\mu_1, \sum_1) \\
 &= \frac{p(y=1)}{p(\mathbf{x})} \prod_{j=1}^d N(\mu_{1,j}, \sigma_j^2) \\
 &= \frac{p(y=1)}{p(\mathbf{x})} \prod_{j=1}^d \frac{1}{\sqrt{2\pi\sigma_j^2}} \exp\left(-\frac{(x_j - \mu_{1,j})^2}{2\sigma_j^2}\right) \\
 &= \frac{p(y=1) \prod_{j=1}^d \frac{1}{\sqrt{2\pi\sigma_j^2}} \exp\left(-\frac{(x_j - \mu_{1,j})^2}{2\sigma_j^2}\right)}{p(y=0) \prod_{j=1}^d \frac{1}{\sqrt{2\pi\sigma_j^2}} \exp\left(-\frac{(x_j - \mu_{0,j})^2}{2\sigma_j^2}\right) + p(y=1) \prod_{j=1}^d \frac{1}{\sqrt{2\pi\sigma_j^2}} \exp\left(-\frac{(x_j - \mu_{1,j})^2}{2\sigma_j^2}\right)} \\
 &= \frac{1}{1 + \frac{p(y=0)}{p(y=1)} \frac{\prod_{j=1}^d \frac{1}{\sqrt{2\pi\sigma_j^2}} \exp\left(-\frac{(x_j - \mu_{0,j})^2}{2\sigma_j^2}\right)}{\prod_{j=1}^d \frac{1}{\sqrt{2\pi\sigma_j^2}} \exp\left(-\frac{(x_j - \mu_{1,j})^2}{2\sigma_j^2}\right)}} \\
 &= \frac{1}{1 + \frac{p(y=0)}{p(y=1)} \prod_{j=1}^d \exp((x_j - \mu_{1,j})^2 - (x_j - \mu_{0,j})^2)} \\
 &= \frac{1}{1 + \exp(\log \frac{p(y=0)}{p(y=1)} + \sum_{j=1}^d (\mu_{1,j}^2 - \mu_{0,j}^2) + 2 \sum_{j=1}^d (\mu_{0,j} - \mu_{1,j}) x_j)} \tag{1}
 \end{aligned}$$

$$\therefore \begin{cases} w_0 = \sum_{j=1}^d (\mu_{1,j}^2 - \mu_{0,j}^2) \\ \mathbf{w} = 2(\mu_0 - \mu_1) \end{cases} \tag{2}$$

2. Yes, they produce the same classifier because it's a convex optimization problem.

3. Please, see a Jupyter notebook file submitted together.

## References

- [1] Christopher M. Bishop, Pattern Recognition and Machine Learning
- [2] Discussion with Tomoki Tsujimura and Bowen Shi