

# Homework 1

Zhaoxia Yu

2023-04-11

Due on **April 17 2023**

## Problem 1

- Suppose  $X_1, X_2, Y_1, Y_2$  are mutually independent.
  - $X_1$  and  $X_2$  are iid from  $N(\mu = 0, \sigma_x^2 = 2^2)$
  - $Y_1$  and  $Y_2$  are iid from  $N(\mu = 0, \sigma_y^2 = 1^2)$  Consider the two pairs  $(X_1, X_2)$  and  $(Y_1, Y_2)$ . Which pair tends to have a larger difference? To answer the question, please calculate and estimate the following two probabilities:

$$P(|X_1 - X_2| > 4), P(|Y_1 - Y_2| > 4)$$

- The hints for calculating/estimating  $P(|X_1 - X_2| > 4)$  can be found in the two slides. Using similar strategies, you can calculate/estimate  $P(|Y_1 - Y_2| > 4)$

**Calculate**  $P(|X_1 - X_2| > 4)$

- Hints for calculating  $P(|X_1 - X_2| > 4)$ .
  - First find the distribution of  $X_1 - X_2$ . Then standard it to have mean 0 and SD 1.
  - Second, express the probability to  $P(|Z| > z)$ , where  $Z \sim N(0, 1)$ .
  - Next, expression the probability in terms of  $\Phi(\cdot)$ , the CDF of the standard normal distribution.
  - Last, use the “pnorm” function in R to find the numerical value.

**Estimate**  $P(|X_1 - X_2| > 4)$

- The probability can be estimated by doing simulations/sampling.
- If you sample many (say 10,000) pairs of  $X_1$  and  $X_2$ , count how many pairs satisfying  $|X_1 - X_2| > 4$ . The probability can be used to estimate  $P(|X_1 - X_2| > 4)$

## Problem 2

- Find a matrix  $A$  such that  $AY$  gives the difference of mean vectors between iris setosa and iris versicolor
- Find a matrix  $B$  such that  $YB$  is column-standardized, i.e., the standard deviation of each column/feature is 1.
- Check the following
  - Let  $C = \mathbf{I}_{150} - \frac{1}{150}J$ , where  $J_{150 \times 150}$  is an all-ones matrices . Use R to verify that  $CY$  centers each column/feature. The R code for  $C$  is ”

`C=diag(1,150) - \frac{1}{150}matrix(1, 150, 150)`

- Let  $S$  be the sample covariance matrix. Use R to verify that each column of  $CYS^{-1/2}$  has been centered and standardized (in fact, the columns have also been de-correlated). Hints:

`S=cov(Y)`

To compute  $S^{-1/2}$ , you may need an R package, such as “qtl2pleio”.

### **Problem 3**

- Choose a picture you like and conduct approximations using singular value decomposition (SVD).