

IE 522 HW12

Q1

5 Points

Consider the time series model given by the following:

$$\text{Equation 1 : } Y_t - \mu = \phi_1(Y_{t-1} - \mu) + \phi_2(Y_{t-2} - \mu) + \epsilon_t,$$

where $\phi_1 = -0.3$, $\phi_2 = 0.1$, $\mu = 1$, $\epsilon_t \sim N(0, \sigma_\epsilon^2)$, $\sigma_\epsilon^2 = 1$.

Q1.1

1 Point

 No file chosen

What are the roots of $\phi(z) = 1 - \phi_1 z - \phi_2 z^2$?

$Z_1 = 5$, $Z_2 = -2$.

CURRENTLY UPLOADED FILES

▼ A12Q1-1.png

 Download  Remove

Q1

5 Points

Consider the time series model given by the following:

$$\text{Equation 1 : } Y_t - \mu = \phi_1(Y_{t-1} - \mu) + \phi_2(Y_{t-2} - \mu) + \epsilon_t,$$

where $\phi_1 = -0.3$, $\phi_2 = 0.1$, $\mu = 1$, $\epsilon_t \sim N(0, \sigma_\epsilon^2)$, $\sigma_\epsilon^2 = 1$.

Q1.1

1 Point

What are the roots of $\phi(z) = 1 - \phi_1 z - \phi_2 z^2$?

$$\phi(z) = 1 - \phi_1 z - \phi_2 z^2 \rightarrow \left\{ \begin{array}{l} \phi_1 = -0.3 \\ \phi_2 = 0.1 \end{array} \right.$$

$$\begin{aligned} \Rightarrow \phi(z) &= 1 + 0.3z - 0.1z^2 \\ &\quad - 0.1(z^2 - 3z - 10) = 0 \\ &\quad -0.1(z-5)(z+2) = 0 \\ \Rightarrow z_1 &= 5, z_2 = -2 \end{aligned}$$

Choose Files

No file chosen

$$\phi(z) = (1 - \frac{1}{3}z)(1 + \frac{1}{2}z)$$

 Please select file(s)

Select file(s)

Save Answer

Last saved on **Nov 15 at 1:19 AM**

Q1.2

1 Point

Is $\{Y_t\}$ stationary? Why?

Yes. Since $|1/2|$ and $| - 1/5 | < 1$, they are *absolutely summable*, which make Y_t stationary.

Choose Files

UPLOADED FILES

No file chosen

▼ A12Q1-2.png

 Download

 Remove

Q1.2

1 Point

Is $\{Y_t\}$ stationary? Why?

$$\phi(z) = (1 - \frac{1}{5}z)(1 + \frac{1}{5}z)$$

↓

$$Y_t - M = -0.3(Y_{t-1} - M) + 0.1(Y_{t-2} - M) + \varepsilon_t$$

↓

$$(1 + 0.3L - 0.1L^2)(Y_t - M) = \varepsilon_t$$

↓

$$(1 + \frac{1}{5}L)(1 - \frac{1}{5}L)(Y_t - M) = \varepsilon_t$$

$$\begin{aligned} Y_t &= M + C(1 + \frac{1}{5}L)^{-1}(1 - \frac{1}{5}L)^{-1}\varepsilon_t \\ &= M + (1 + \frac{1}{5}L)^{-1}(1 - \frac{1}{5}L)^{-1}\varepsilon_t \end{aligned}$$

 No file chosen

 Please select file(s)

Select file(s)

Save Answer

Last saved on **Nov 15 at 1:19 AM**

Q1.3

1 Point

Using the Yule-Walker equation, derive an equation relating $\gamma(0)$ and $\gamma(1)$.

$$-3\gamma(1) = \gamma(0).$$

Choose Files

LOADED FILES

No file chosen

▼ A12Q1-3.png

 Download  Remove

Q1.3

1 Point

Using the Yule-Walker equation, derive an equation relating $\gamma(0)$ and $\gamma(1)$.

$$\begin{aligned}Y_t - M &= -0.3(Y_{t-1} - M) + 0.1(Y_{t-2} - M) + \varepsilon_t \\ \rightarrow \gamma(h) &= -0.3\gamma(h-1) + 0.1\gamma(h-2) \\ \rightarrow \rho(h) &= -0.3\rho(h-1) + 0.1\rho(h-2) \\ \rho(1) &= -0.3\rho(0) + 0.1\rho(-1) \quad \leftarrow \rho(-1) = \rho(1) \\ 0.9\rho(1) &= -0.3\rho(0) = -0.3 \\ \downarrow \\ \rho(1) &= -\frac{1}{3} = \frac{\gamma(1)}{\gamma(0)} \\ \Rightarrow \gamma(1) &= \gamma(0) \times\end{aligned}$$

Choose Files No file chosen

 Please select file(s)

Select file(s)

Save Answer

Last saved on **Nov 15 at 1:19 AM**

Q1.4

1 Point

Compute the variance of the two sides of Equation 1 to get another relationship for $\gamma(0)$ and $\gamma(1)$. Using this and the relationship in question 1.3, compute $\gamma(0)$ and $\gamma(1)$.

$$(1) 0.9\gamma(0) = -0.06\gamma(1) + 1$$

$$(2) \gamma(0) = 3/2.64, \gamma(1) = 1/ - 2.64.$$

No file chosen

CURRENTLY UPLOADED FILES

▼ A12Q1-4.png

 Download  Remove

Q1.4

1 Point

Compute the variance of the two sides of Equation 1 to get another relationship for $\gamma(0)$ and $\gamma(1)$. Using this and the relationship in question 1.3, compute $\gamma(0)$ and $\gamma(1)$.

$$\textcircled{1} \quad Y_t - M = \phi_1(Y_{t-1} - M) + \phi_2(Y_{t-2} - M) + \varepsilon_t$$

$$\begin{aligned} V(0) &= \text{Var}(\underbrace{\phi_1(Y_{t-1} - M)}_{\phi_1^2 V(0)} + \underbrace{\phi_2(Y_{t-2} - M)}_{\phi_2^2 V(0)} + \underbrace{\varepsilon_t}_{\sigma^2}) \\ &= \phi_1^2 V(0) + \phi_2^2 V(0) + \sigma^2 \\ &\quad + 2\phi_1\phi_2 r(1) \\ &\quad + 2\phi_1 \text{cov}(Y_{t-1} - M, \varepsilon_t) \\ &\quad + 2\phi_2 \text{cov}(Y_{t-2} - M, \varepsilon_t) \end{aligned}$$

$$\begin{aligned} V(0) &= \phi_1^2 V(0) + \phi_2^2 V(0) + 2\phi_1\phi_2 r(1) + \sigma^2 \\ \Rightarrow V(0) &= 0.87 V(0) + 0.01 V(0) - 0.06 r(1) + 1 \\ &\quad \left(\text{var}(aX - bY) = \text{var}[aX + (-b)Y] = a^2 \text{var}(X) + (-b)^2 \text{var}(Y) + 2a(-b)\text{cov}(X, Y) = a^2 \text{var}(X) + b^2 \text{var}(Y) - 2ab\text{cov}(X, Y) \right) \end{aligned}$$

Choose Files No file chosen $= -0.06 r(1) + 1$ *

$$\begin{aligned} \textcircled{2} \quad \begin{cases} - \Rightarrow r(1) = V(0) \\ 0.9 V(0) = -0.06 r(1) + 1 \end{cases} \\ \Rightarrow -2.7 V(0) = -0.06 r(1) + 1 \\ \Rightarrow -2.64 V(0) = 1, \quad r(1) = \frac{1}{-2.64} \\ V(0) = \frac{3}{2.64} \end{aligned}$$

 Please select file(s)

Select file(s)

Save Answer

Last saved on **Nov 16 at 12:38 PM**

Q1.5

1 Point

Suppose the model in Equation 1 is expanded to an ARMA(2,1) model:

$$Y_t - \mu = \phi_1(Y_{t-1} - \mu) + \phi_2(Y_{t-2} - \mu) + \epsilon_t + \theta_1\epsilon_{t-1},$$

where $\theta_1 = 2$, $\phi_1, \phi_2, \sigma_\epsilon^2$ stay the same. Is $\{Y_t\}$ stationary? Why?

Choose Files No file chosen
 $|1/2|, |1/5| < 1$, it is *stationary*.

CURRENTLY UPLOADED FILES

▼ A12Q1-5.png

 Download  Remove

Q1.5

1 Point

Suppose the model in Equation 1 is expanded to an ARMA(2,1) model:

$$Y_t - \mu = \phi_1(Y_{t-1} - \mu) + \phi_2(Y_{t-2} - \mu) + \epsilon_t + \theta_1 \epsilon_{t-1},$$

where $\theta_1 = 2$, $\phi_1, \phi_2, \sigma_\epsilon^2$ stay the same. Is $\{Y_t\}$ stationary? Why?

$$\begin{aligned} Y_t - \mu &= \phi_1(Y_{t-1} - \mu) + \phi_2(Y_{t-2} - \mu) + \epsilon_t + \theta_1 \epsilon_{t-1} \\ \Rightarrow Y_t - \mu &= -0.3(Y_{t-1} - \mu) + 0.1(Y_{t-2} - \mu) + \epsilon_t + 2\epsilon_{t-1} \\ &\quad \downarrow \\ (1 + 0.3L - 0.1L^2)(Y_t - \mu) &= \epsilon_t + 2\epsilon_{t-1} \\ &\quad \downarrow \\ (1 + \frac{1}{2}L)(1 - \frac{1}{5}L)(Y_t - \mu) &= \epsilon_t + 2\epsilon_{t-1} \\ \Rightarrow (1 + \frac{1}{2}L)(1 - \frac{1}{5}L)(Y_t - \mu) &= (1 + 2L)\epsilon_t \\ \phi(L)(Y_t - \mu) &= \theta(L)\epsilon_t \end{aligned}$$

Choose Files No file chosen

 Please select file(s)

Select file(s)

Save Answer

Last saved on **Nov 15 at 1:19 AM**

Q2

5 Points

From R library Ecdat, get Monthly risk free rate from Jan 1960 to Dec 2002. Compute monthly changes of the risk free rate.

```
library(Ecdat)
data(Capm)
Choose Files No file chosen
ar=arir(r)
```

Q2.1

1 Point

Suppose Conditional MLE is used to estimate the parameters of an AR(2) model for $\{Y_t = dr_t, 1 \leq t \leq m\}$:

$$Y_t = a + \phi_1 Y_{t-1} + \phi_2 Y_{t-2} + \epsilon_t, \quad \epsilon_t \sim N(0, \sigma_\epsilon^2).$$

Conditional on the first two observations y_1, y_2 , write down the expression of the log-likelihood of observing $Y_3 = y_3, \dots, Y_m = y_m$.

(shown below)

CURRENTLY UPLOADED FILES

▼ Page1.png

 Download  Remove

No file chosen

Q2.1

1 Point

Suppose Conditional MLE is used to estimate the parameters of an AR(2) model for $\{Y_t = dr_t, 1 \leq t \leq m\}$:

$$Y_t = a + \phi_1 Y_{t-1} + \phi_2 Y_{t-2} + \epsilon_t, \quad \epsilon_t \sim N(0, \sigma_\epsilon^2).$$

Conditional on the first two observations y_1, y_2 , write down the expression of the log-likelihood of observing $Y_3 = y_3, \dots, Y_m = y_m$.

$$Y_3 \sim N(a + \phi_1 y_1 + \phi_2 y_2, \sigma_\epsilon^2)$$

$$f_{Y_3}(y_3) = \frac{1}{\sqrt{2\pi\sigma_\epsilon^2}} e^{-\frac{(y_3 - a - \phi_1 y_1 - \phi_2 y_2)^2}{2\sigma_\epsilon^2}}$$

$$\text{Log-likelihood func}$$

$$= \log \left(\frac{1}{\sqrt{2\pi\sigma_\epsilon^2}} e^{-\frac{(y_3 - a - \phi_1 y_1 - \phi_2 y_2)^2}{2\sigma_\epsilon^2}} \right)$$

Choose Files No file chosen

$$\log \left(\frac{1}{\sqrt{2\pi\sigma_\epsilon^2}} e^{-\frac{(y_3 - a - \phi_1 y_1 - \phi_2 y_2)^2}{2\sigma_\epsilon^2}} \right)$$

$$= -\frac{1}{2} \log 2\pi\sigma_\epsilon^2 - \frac{(y_3 - a - \phi_1 y_1 - \phi_2 y_2)^2}{2\sigma_\epsilon^2}$$

$$f_{Y_4|Y_3}(y_4|y_3) = \frac{1}{\sqrt{2\pi\sigma_\epsilon^2}} e^{-\frac{(y_4 - a - \phi_1 y_2 - \phi_2 y_3)^2}{2\sigma_\epsilon^2}}$$

log-likelihood func:

$$\log [f_{Y_4|Y_3}(y_4|y_3) \times f_{Y_5|Y_3}(y_5|y_3) \dots f_{Y_m|Y_3}(y_m|y_3)]$$

$$= \log \left[\left(\frac{1}{\sqrt{2 \pi \zeta}} \right) \right]$$

▼ Page2.png

 Download  Remove

Choose Files No file chosen

$$\times e^{\left[\frac{(y_4 - a - \phi_1 y_2 - \phi_2 y_3)}{2\sigma^2} - \frac{(y_3 - a - \phi_1 y_1 - \phi_2 y_2)}{2\sigma^2} \right]}$$

$$= - \log 2\pi\sigma^2 - \sum_{i=3}^4 \frac{(y_i - a - \phi_1 y_{i-2} - \phi_2 y_{i-1})^2}{2\sigma^2}$$



log-likelihood func for y_m

$$\Rightarrow -\frac{(n-2)}{2} \log 2\pi\sigma^2 - \sum_{i=3}^n \frac{(y_i - a - \phi_1 y_{i-2} - \phi_2 y_{i-1})^2}{2\sigma^2}$$

Choose Files No file chosen

 Please select file(s) Select file(s)

Save Answer

Last saved on **Nov 15 at 1:19 AM**

Q2.2

1 Point

Suppose Exact MLE is used to estimate the parameters of a MA(2) model for $\{Y_t = dr_t, 1 \leq t \leq m\}$:

$$Y_t = \mu + \epsilon_t + \theta_1 \epsilon_{t-1} + \theta_2 \epsilon_{t-2}, \quad \epsilon_t \sim N(0, \sigma_\epsilon^2).$$

What's the mean vector and covariance matrix of the multivariate normal distribution for

Choose Files No file chosen $(\epsilon_1, \epsilon_2, \dots, \epsilon_m)$ in terms of $\mu, \theta_1, \theta_2, \sigma_\epsilon^2$?

(shown below)

CURRENTLY UPLOADED FILES

▼ Page1.png

 Download  Remove

Q2.2

1 Point

Suppose Exact MLE is used to estimate the parameters of a MA(2) model for $\{Y_t = dr_t, 1 \leq t \leq m\}$:

$$Y_t = \mu + \epsilon_t + \theta_1 \epsilon_{t-1} + \theta_2 \epsilon_{t-2}, \quad \epsilon_t \sim N(0, \sigma_\epsilon^2).$$

What's the mean vector and covariance matrix of the multivariate normal distribution for $(Y_1, \dots, Y_m)^\top$ in terms of $\mu, \theta_1, \theta_2, \sigma_\epsilon^2$?

$$\textcircled{1} \text{ mean vector} = (\underbrace{\mu, \mu, \dots, \mu}_m)^\top$$

$$\begin{aligned}\textcircled{2} \quad \gamma(0) &= \text{Var}(Y_t) = \text{Var}(\mu + \epsilon_t + \theta_1 \epsilon_{t-1} + \theta_2 \epsilon_{t-2}) \\ &= \sigma_\epsilon^2 + \theta_1^2 \sigma_\epsilon^2 + \theta_2^2 \sigma_\epsilon^2 = (1 + \theta_1^2 + \theta_2^2) \sigma_\epsilon^2\end{aligned}$$

$$\gamma(1) = \text{Cov}(Y_t, Y_{t+1})$$

Choose Files

No file chosen

$$\begin{aligned}&= E[(\mu + \theta_1 \epsilon_{t-1} + \theta_2 \epsilon_{t-2}) \times (\mu + \theta_1 \epsilon_t + \theta_2 \epsilon_{t-1})] \\ &= E(\theta_1 \epsilon_t) + E(\theta_1 \theta_2 \epsilon_{t-1}^2) \\ &= \theta_1 \sigma_\epsilon^2 + \theta_1 \theta_2 \sigma_\epsilon^2 = (\theta_1 + \theta_1 \theta_2) \sigma_\epsilon^2\end{aligned}$$

$$\gamma(2) = \text{Cov}(Y_t, Y_{t+2})$$

$$\begin{aligned}&= E[(\mu + \theta_1 \epsilon_{t-1} + \theta_2 \epsilon_{t-2}) \times (\mu + \theta_1 \epsilon_{t+1} + \theta_2 \epsilon_t)] \\ &= \theta_2 \sigma_\epsilon^2\end{aligned}$$

$$\gamma(h) \text{ for } h > 3 = 0$$

▼ Page2.png

 Download  Remove

No file chosen

$\Rightarrow \sum =$

$$\begin{bmatrix} (1 + \theta_1^2 + \theta_2^2) \zeta_2^2 & (\theta_1 + \theta_1 \theta_2) \zeta_2^2 & \theta_2 \zeta_2^2 & \dots & \dots \\ (\theta_1 + \theta_1 \theta_2) \zeta_2^2 & (1 + \theta_1^2 + \theta_2^2) \zeta_2^2 & \ddots & & \vdots \\ \theta_2 \zeta_2^2 & \ddots & \ddots & \ddots & \theta_2 \zeta_2^2 \\ \vdots & \vdots & \vdots & \ddots & (\theta_1 + \theta_1 \theta_2) \zeta_2^2 \\ \vdots & \ddots & \theta_2 \zeta_2^2 & (\theta_1 + \theta_1 \theta_2) \zeta_2^2 & (1 + \theta_1^2 + \theta_2^2) \zeta_2^2 \end{bmatrix}$$

Choose Files No file chosen

 Please select file(s)

Select file(s)

Save Answer

Last saved on **Nov 15 at 1:19 AM**

Q2.3

1 Point

Use auto.arima to fit dr. Using the fit you obtain, forecast the monthly change from Dec 2002 to Jan 2003. What's the corresponding forecast for the interest rate in Jan 2003?

Series: dr

ARIMA(1,0,1) with zero mean

Choose Files **No file chosen**

Coefficients:

ar1 ma1

0.4551 -0.6761

s.e. 0.1104 0.0903

$\sigma^2 = 0.004751$: log likelihood = 647.65

AIC=-1289.29 AICc=-1289.25 BIC=-1276.56

$$Y_t - \mu = \phi_1(Y_{t-1} - \mu) + \epsilon_t + \theta_1 \epsilon_{t-1}$$

$$= Y_t - \mu = 0.4551(Y_{t-1} - \mu) + \epsilon_t - 0.6761\epsilon_{t-1}$$

$$\text{Forecast model} = \hat{Y}_{t+1} = \mu + 0.4551(Y_t - \mu) - 0.6761\epsilon_t$$

Monthly Change = Y_hat = -0.01154477

The IRR of January = J2003 = 0.09845523

CURRENTLY UPLOADED FILES

▼ Q2-3.R

 Download  Remove

```
1 #Q2-3
2 #Data Preparation
3 library(Ecdat)
4 data(Capm)
5 r = Capm$rf
6 dr = diff(r)
7
8 #Fit ARIMA model
9 library(forecast)
10 auto.arima(dr)
11
12 #Get December 2002 to January 2003 monthly change
```

Choose Files No files chosen

```
14 Y_t = dr[length(dr)]
15 Y_t1 = dr[length(dr)-1]
16 Y_hat = mu + 0.4551*(Y_t - mu) - 0.6761*(Y_t - Y_t1)
17 Y_hat
18
19 #Get January 2003
20 D2002 = r[length(r)]
21 J2003 = D2002 + Y_hat
22 J2003
23
```

Please select file(s)

Select file(s)

Save Answer

Last saved on **Nov 15 at 1:19 AM**

Q2.4

1 Point

What's the 90% prediction interval for the monthly change from Dec 2002 to Jan 2003? What's the corresponding 90% prediction interval for the interest rate in Jan 2003?

The 90% prediction interval of montely change from December 2022 to January 2023 =
[-0.1079146, 0.1188449]

The 90% prediction interval of January 2023 = [0.00208540143421547,
0.228844940056894]

CURRENTLY UPLOADED FILES

▼ Q2-4.R

 Download  Remove

Choose Files **No file chosen**

```
2 #Data Preparation
3 library(Ecdat)
4 data(Capm)
5 r = Capm$rf
6 dr = diff(r)
7
8 #Fit ARIMA model
9 library(forecast)
10 fit = auto.arima(dr)
11 fit
12
```

```

13 #Get December 2002 to January 2003 monthly change
14 mu = mean(dr)
15 Y_t = dr[length(dr)]
16 Y_t1 = dr[length(dr)-1]
17 Y_hat = mu + 0.4551*(Y_t - mu) - 0.6761*(Y_t - Y_t1)
18 Y_hat
19
20 #Get Prediction Interval for monthly change
21 alpha = 0.1
22 f = forecast(fit, 1, level = 90)
23 f
24 #Get Prediction Interval for January 2003
25 D2002 = r[length(r)]
26 J2003ub = D2002 + f$upper
27 J2003lb = D2002 + f$lower
28 print(paste(J2003lb, J2003ub))
29
30

```

 Please select file(s)

Select file(s)

Save Answer

Last saved on **Nov 30 at 11:59 PM**

Choose Files **No file chosen**

Q2.5

1 Point

If you try to produce a k -step-ahead forecast, when $k \rightarrow +\infty$, what'll be your forecast for the monthly change? What'll be the variance of the forecasting error?

- (1) multi-step-ahead forecast converges to $\mu = -0.0004271845$.
- (2) The variance converges to 0.005044025 .

CURRENTLY UPLOADED FILES

▼ Q2-5.R

 [Download](#) [Remove](#)

```
1 #Q2-5
2 #Data Preparation
3 library(Ecdat)
4 data(Capm)
5 r = Capm$rf
6 dr = diff(r)
7
8 #Fit ARIMA model
9 library(forecast)
10 fit = auto.arima(dr)
11 fit
12
13 #Get December 2002 to January 2003 monthly change
14 mu = mean(dr)
15 Y_t = dr[length(dr)]
16 Y_t1 = dr[length(dr)-1]
17 Y_hat = mu + 0.4551*(Y_t - mu) - 0.6761*(Y_t - Y_t1)
18 mu
19
20 #Get the gamma(0) after infinity periods
21 gamma = fit$sigma2 * (1 + (-0.6761)^2 + 2*0.4551*(-0.6761)) / (1 - 0.4551^2)
22 gamma
23
```

 Please select file(s)Select file(s)Save AnswerLast saved on **Nov 15 at 1:19 AM**Save All AnswersSubmit & View Submission ➔

