

Physics 7501 (Autumn 2026)
Quantum Mechanics 1 (graduate core course)

Yuan-Ming Lu

August 25, 2026

Abstract. In this elementary course of quantum mechanics, we will learn the mathematical basis of quantum mechanics (i.e. linear algebra), the postulates of quantum theory, and how to use quantum mechanics to understand microscopic quantum phenomena.

Contents

0	Course Information	1
1	Postulates and Mathematics of Quantum Mechanics	5
1.1	Key experiments that established quantum mechanics (8/26)	6
1.1.1	Energy quantization of light in terms of photons	6
1.1.2	Quantization of atomic energy levels	8
1.1.3	Quantization of angular momentum	8
1.1.4	Davisson-Germer experiment demonstrated the wave nature of electrons	10
1.2	Hilbert space and state vectors (8/28)	13
1.3	Linear operators and observables (9/2)	16
1.4	Projective measurements (9/4)	19
1.5	Unitary evolution of a closed quantum system (9/9)	21
1.5.1	Appendix: Functions of operators/matrices	22
1.6	Composite quantum systems and tensor product (9/11)	24
A	Homework 1 (due 9/4)	26
A.1	Franck-Hertz experiment	26
A.2	Davisson-Germer experiment	26
A.3	Hilbert space $\mathbb{L}_{\mathbb{V}}$ of linear operators on $\mathbb{V} = \mathbb{C}^n$	28
B	Homework 2 (due 9/11)	30
B.0.1	Pauli matrices	30
B.0.2	Linear algebra	30
B.0.3	Consecutive measurements of incompatible operators	31
B.0.4	Quantum Zeno effect	32
	Bibliography	33

0 Course Information

Instructor

Professor **Yuan-Ming Lu**

Contact Information

- **Email:** lu.1435@osu.edu
- **Office Phone:** 614-292-3084
- **Office:** 2012 Physics Research Building (PRB)

Office Hours

Friday 2:00–3:00 pm in PRB 2015

Lecture Logistics

- **Location:** Scott Lab N050
- **Time:** Wed. Fri. 11:10 am – 12:30 pm
- **Schedule:** See Google Spreadsheet

Teaching Assistant & Guided Group Work

- **Graduate Teaching Assistant (GTA):** Mr. Hyungjun Jeon (jeon.258@osu.edu)
- **GTA Office Hours:** Thursday 3:00–4:00 pm in PRB 2015
- **Guided Group Work (GGW) Sessions:** Thursdays 9:30 am – 11:00 am in PRB 0118. For questions regarding GGW sessions, please contact Dr. Beatriz E Burrola Gabilondo (burrolagabilondo.1@osu.edu).

Prerequisites

The course will assume physics and mathematics knowledge at the advanced undergraduate level (e.g., basic quantum mechanics, linear algebra, and differential equations), which we will review as needed.

Lecture Notes

Lecture notes will be uploaded to Carmen as PDF files generated from LaTeX sources at least 4 days before each lecture.

Main Textbooks (Recommended but Not Required)

- *Modern Quantum Mechanics* by J.J. Sakurai and Jim Napolitano (3rd Edition, Cambridge University Press, 2020)
- *Quantum Mechanics* (Vol. 3, Lectures on Theoretical Physics) by David Tong (Cambridge University Press, 2025). Problem sets available at <https://www.damtp.cam.ac.uk/user/tong/books>
- *Principles of Quantum Mechanics* by Ramamurti Shankar (2nd Edition, Plenum Press, 2011)
- *Quantum Computation and Quantum Information* by Michael Nielsen and Isaac Chuang (Cambridge University Press, 2011)

Other References

- *Lectures on Quantum Mechanics and on Topics in Quantum Mechanics*, David Tong, University of Cambridge
- *Quantum Mechanics: Non-Relativistic Theory* by Lev Landau and L. Lifshitz (2nd Edition, Pergamon Press)
- *Lectures on Quantum Mechanics* by Steven Weinberg (Cambridge University Press, 2013)
- *Quantum theory: a foundational approach* by Charis Anastopoulos (Cambridge University Press, 2024)
- *Quantum Mechanics* by Arjun Berera and Luigi Del Debbio (Cambridge University Press, 2021)

Reading Assignments

For each lecture, you will be expected to complete reading assignments before class. These will be announced on Carmen. On days when reading assignments are discussed in class, you must answer questions about the reading using Carmen "Quizzes" by 10:00 am the day of class. Your answers will not be graded for correctness; full credit is given for submitting on time. Late submissions will not be counted.

Homework

Homework will be assigned weekly on Carmen (except during holidays or exams). Submissions are due on Carmen by 11:59 pm on Fridays (or occasionally Wednesdays).

HW Quizzes

To verify individual effort in solving homework problems, a 10-minute quiz will take place during the last 10 minutes (12:20–12:30 pm) of lecture on homework due dates. Handwritten answers will be scanned and uploaded to Carmen/Gradescope.

Grading Scheme

All homework, quizzes, and exams will be graded by the GTA. Refer to Table 1 for the grading allocation.

Table 1: Grading basis for the course.

Item	Time	Assignments	Percentage
Reading Quizzes	10:00 am on lecture day	23	6%
Homework	Friday at 11:59 pm	12	18%
HW Quizzes	Last 10 mins of Friday lecture	12	18%
Mid Term Exam	Oct. 9 (Fri.), 11:30 am–1:30 pm	1	29%
Final Exam	Dec. 17 (Thu.), 12:00–2:00 pm	1	29%

Exams

- **Mid Term Exam:** In-person exam scheduled for 11:30 am–1:30 pm on Friday, Oct. 9 in PRB 1080 (Smith Seminar Room).
- **Final Exam:** In-person exam scheduled for 12:00–2:00 pm on Thursday, Dec. 17 in PRB 1080 (Smith Seminar Room).

Course Topics

- Linear algebra: mathematics of quantum mechanics
- Postulates of quantum mechanics

- Quantum entanglement
- Schrödinger equation and quantum dynamics
- Heisenberg's uncertainty principle
- One-dimensional potentials
- The harmonic oscillator and coherent states
- Squeezed states and quantum optics
- Symmetry and conservation laws
- Rotational symmetry and angular momentum
- Central potentials and the hydrogen atom
- Variational methods and WKB approximation
- Perturbation theory: time-independent and time-dependent
- Scattering theory

Shared Values

Please note The Ohio State University Shared Values.

Academic Misconduct

Academic integrity is essential to maintaining an environment that fosters excellence in teaching, research, and other educational activities. The Ohio State University and the Committee on Academic Misconduct (COAM, <https://oaa.osu.edu/resources/policies-and-procedures/committee-academic-misconduct>) expect that all students understand the University's Code of Student Conduct (<https://trustees.osu.edu/bylaws-and-rules/code>). Any activity that compromises academic integrity or subverts the educational process will be reported to COAM.

Artificial Intelligence and Academic Integrity

Generative AI tools are permitted as learning aids to support your understanding (*e.g.*, asking for alternative explanations or math reviews). However, every calculation, derivation, and explanation submitted must reflect your own understanding. You may not copy or lightly modify AI-generated solutions. AI tools are strictly prohibited during timed assessments, exams, and quizzes. When AI is permitted and used for homework, include a brief acknowledgment detailing the tool and how you verified its output.

Religious Accommodations

Ohio State accommodates students' religious beliefs and practices. Requests for accommodations must be communicated in writing to the instructor within the first 14 days of the semester. See the official policy at OAA Religious Holidays Policy.

Disability Statement & Illness

Students requesting disability accommodations should register with Student Life Disability Services (<https://slds.osu.edu/>) and notify the instructor as early as possible. If you are ill and need to miss class, notify the instructor immediately.

Intellectual Diversity

Ohio State is committed to open inquiry and intellectual diversity. Classroom discussions aim to foster critical thinking on course material without promoting any single point of view.

Grievances and Problem Solving

To resolve grievances concerning grades or academic practices, contact the instructor first, followed by the department chairperson, college dean, and provost (Faculty Rule 3335-8-23).

Creating an Environment Free from Harassment and Discrimination

The Ohio State University does not tolerate harassment, discrimination, or sexual misconduct. Reports can be submitted to the Civil Rights Compliance Office (CRCO) at <http://civilrights.osu.edu/>, by phone at 614-247-5838, or via email at civilrights@osu.edu.

1 Postulates and Mathematics of Quantum Mechanics

1.1 Key experiments that established quantum mechanics (8/26)

- Refer to Weinberg Chap 1. For further reading, try Anastopoulos Chap 2, Shankar 3.

Quantum mechanics was first motivated by a series of experiments that cannot be explained by classical mechanics and statistical physics. These exceptions to classical physics happen in the microscopic world, when the scale of physical quantities, such as energy, angular momentum and electric charge becomes comparable to certain characteristic scales.

The quantization of energy was observed in both light and atomic spectra, which motivated Planck, Einstein and others to propose a discretized (“quantized”) energy levels for photons and atoms. Bohr was able to explain the Rydberg formula[1] of hydrogen atom spectral lines under the assumption of a quantized angular momentum for electrons, which was later observed in the Stern-Gerlach experiment[2]. After de Broglie first proposed the wave nature of microscopic particles such as electrons, Davisson and Germer[3] were able to directly observe it in the diffraction pattern of electrons scattered by a nickel crystal. These experiments call for a new theory framework that defeats the intuition of classical physics.

1.1.1 Energy quantization of light in terms of photons

Young’s double-slit experiment clearly demonstrated the wave nature of light, whose energy should change continuously in classical electromagnetism. However, this point of view is seriously challenged by black-body radiation.

1.1.1.1 Black-body radiation and Planck’s formula

Consider light in a cubic box (a simplified model of a black body) of volume L^3 . A standing wave in the box satisfies the following condition for its wavevector $\vec{k} = (k_x, k_y, k_z)$:

$$\lambda_i = \frac{2\pi}{k_i} = \frac{2L}{n_i}, \quad n_i = 1, 2, 3, \dots, \quad i = x, y, z \quad (1.1)$$

Counting the number of standing wave modes in the box in terms of frequency $\nu = ck/2\pi$

$$\int dk_x dk_y dk_z = \frac{1}{8} 4\pi \int k^2 dk = \left(\frac{\pi}{L}\right)^3 \sum_{\{n_i\}} = \left(\frac{\pi}{L}\right)^3 N_{\text{modes}} \quad (1.2)$$

we have the density of states for light-wave modes per unit volume:

$$\frac{1}{L^3} \frac{dN}{d\nu} = \left(\frac{1}{\pi}\right)^3 \frac{\pi}{2} k^2 \frac{dk}{d\nu} = \frac{8\pi}{c^3} \nu^2 \quad (1.3)$$

Assuming the equipartition theorem in classical physics, i.e. each mode carries an averaged energy of $k_B T$, the black-body radiation energy per unit volume is

$$\frac{dE}{L^3 d\nu} = \frac{k_B T}{L^3} \frac{dN}{d\nu} = \frac{8\pi k_B T}{c^3} \nu^2 \quad (1.4)$$

This is known as the Rayleigh-Jeans formula and it correctly captures the long-wavelength (small ν) behavior of universal black-body radiation, first established by Kirchhoff around 1860. However, it diverges in the short wavelength limit, known as the ultraviolet catastrophe. Experimentally the Wien’s Distribution Law was observed at short wavelengths:

$$\frac{dE}{L^3 d\nu} \sim \nu^3 e^{-\alpha(T)\nu} \quad (1.5)$$




Figure 1.1: (Left) Schematic setup of Compton's experiment. (Right) The light-electron scattering process in the Compton experiment.

In order to reconcile long-wavelength Rayleigh-Jeans formula to short-wavelength Wien's law, Planck came up with his famous formula:

$$\frac{dE}{L^3 d\nu} = \frac{8\pi\nu^2}{c^3} \frac{h\nu}{e^{\alpha\nu} - 1}, \quad \alpha = \frac{h}{k_B T} \quad (1.6)$$

where the Planck constant is

$$h = 6.63 \times 10^{-34} \text{ J} \cdot \text{s} \quad (1.7)$$

Planck's formula (1.6) indicates that the statistical-averaged energy of light at frequency ν is given by

$$U(\nu) = \frac{h\nu}{e^{h\nu/K_B T} - 1} = \frac{\sum_{n \in \mathbb{Z}_+} nh\nu e^{-nh\nu/K_B T}}{\sum_{n \in \mathbb{Z}_+} e^{-nh\nu/K_B T}} \quad (1.8)$$

under Planck's quantization hypothesis that the microscopic states of light have energy $nh\nu$ where $n \in \mathbb{Z}_+$ is a positive integer.

1.1.1.2 Einstein's "light quantum" and photoelectric effect

Motivated by Planck's quantization hypothesis, based on the assumption that light gives out energy in multiples of energy quantum $h\nu$, Einstein predicted the photoelectric effect

$$K_{\max} = eV = h\nu - \Phi \quad (1.9)$$

where $K_{\max}$ is the maximal kinetic energy of emitted photoelectrons when light of frequency ν shines on a metal, and Φ is the work function of the metal. V is minimal voltage that is required to stop photoelectron emission. This experiment was carried out by Millikan[4], who obtained Planck's constant

$$h = e \frac{dV}{d\nu} \quad (1.10)$$

1.1.1.3 Compton effect demonstrated quantization of light in terms of photons

Discovered by Arthur Compton in 1923[5], when the energy of the X ray photon (~ 17 keV) was significantly larger than the binding energy of the atomic electron, the electrons could be treated as being free after scattering. In this case, a *Compton shift* between the wavelengths

of injected light (λ) and the scattered light (λ') was observed to satisfy the following relation with the scattering angle θ (see Fig. 1.1):

$$\lambda' - \lambda = \frac{h}{m_e c} (1 - \cos \theta) \quad (1.11)$$

where h is the Planck's constant.

The Compton shift can be explained by energy and momentum conservation, by assuming the photon energy of $E = h\nu = hc/\lambda$.

1.1.2 Quantization of atomic energy levels

1.1.2.1 Rydberg formula for the spectral lines in a hydrogen atom

After the discovery of dark lines in the spectrum of the sun at the beginning of 18th century, it was later realized that hot atomic gases absorb and emit light only at certain definite frequencies. Building on works of Balmer and others, Rydberg discovered the following rule[1] for the frequency of emitted light from hydrogen gases:

$$\nu = \frac{c}{\lambda} = cR_H \left(\frac{1}{n^2} - \frac{1}{m^2} \right), \quad m, n \in \mathbb{Z}_+. \quad (1.12)$$

This clearly demonstrated the discreteness of atomic energy levels $E_n \propto \frac{1}{n^2}$, further motivating Bohr's theory of the hydrogen atom.

1.1.2.2 Franck-Hertz experiment

In the Franck-Hertz experiment (1914) illustrated in Fig. 1.2, in a vacuum tube fitted with three electrodes (an electron-emitting, hot cathode; a metal mesh grid; and an anode), when a grid voltage is applied between the cathode and the anode, the electric current through the tube filled with thin mercury vapours, due to electrons that pass through the grid and reach the anode, is measured in the experiment. As shown in the middle figure, the current increases rapidly at low voltages, and then sharply drops almost back to zero at 4.9 V, with similar effects observed as 9.8 V and other multiples of 4.9 V. This can be understood by the quantized energy levels of mercury atoms. inside the single mercury droplet in the vacuum tube, as shown in Fig. 1.2.

1.1.3 Quantization of angular momentum

1.1.3.1 Bohr's explanation of hydrogen atom spectral lines

Rydberg's rule[1] indicates that the photon energy emitted by the hydrogen atom is given by

$$h\nu = E_m - E_n, \quad E_n = -\frac{hcR_H}{n^2}, \quad n \in \mathbb{Z}_+. \quad (1.13)$$

How to explain the $1/n^2$ quantization of hydrogen atom energy levels? According to classical mechanics, for an electron in the Coulomb force field of hydrogen nucleus, its speed v and radius r satisfies

$$\frac{mv^2}{r} = \frac{ke^2}{r^2} \implies E = \frac{1}{2}mv^2 - \frac{ke^2}{r} = -\frac{ke^2}{2r} \quad (1.14)$$




Figure 1.2: (Left) Photograph of a vacuum tube used for the Franck-Hertz experiment. (Middle) Anode current (arbitrary units) versus the grid voltage relative to the cathode. (Right) Wavelengths of light emitted by a mercury vapor discharge and by a Franck-Hertz tube in operation at 10 V. (Picture from Wikipedia)

where k is the Coulomb constant. To fit the n^2 dependence of the orbit radius r , Niels Bohr came up with the following quantization condition for angular momentum:

$$mvr = n\hbar \quad (1.15)$$

where $\hbar$ turns out to be $h/2\pi$. This quantization condition leads to the famous energy levels of electrons in a hydrogen atom:

$$E_n = -\frac{mk^2e^4}{2\hbar^2} \frac{1}{n^2} = -\frac{13.6 \text{ eV}}{n^2} \quad (1.16)$$

1.1.3.2 Stern-Gerlach experiment demonstrated the quantization of angular momentum

In 1922 Stern and Gerlach performed an experiment[2] measuring the deflection of silver atoms by an inhomogeneous magnetic field, as illustrated in Fig. 1.3. Due to the interaction energy $U = -\vec{\mu} \cdot \vec{B}$ of a magnetic moment $\vec{\mu}$ under a magnetic field $\vec{B} = B_z \hat{z}$, a non-uniform magnetic field exerts a force $\vec{F} = -\vec{\nabla}U = \mu_z \vec{\nabla}B_z$ on the moment-carrying silver atoms, therefore deflecting the silver atom beams in the inhomogeneous magnetic field. As shown in Fig. 1.3, in contrast a continuous range of values of the magnetic moment as expected in classical mechanics, the experimental observations pointed to quantization of magnetic moment $\mu_z = g\mu_B S_z/\hbar$, where we defined μ_B as the Bohr magneton:

$$\mu_B = \frac{\hbar e}{2m_e} \quad (1.17)$$

$g \approx 2$ as the electron g -factor and

$$S_z = \pm \frac{\hbar}{2} \quad (1.18)$$

is the quantized spin angular momentum of a silver atom observed in Stern-Gerlach experiment.

This experiment demonstrated that in addition to energy level quantization, other physical observable such as spin angular momentum are also quantized in microscopic scales such as in




Figure 1.3: Schematic setup of Stern-Gerlach experiment: Silver atoms travelling through an inhomogeneous magnetic field, being deflected up or down depending on their spin. (1) furnace, (2) beam of silver atoms, (3) inhomogeneous magnetic field, (4) the classically expected result, (5) the observed result.

silver atoms. Moreover, we can consider sequential Stern-Gerlach experiments shown in Fig. 1.4, where silver atoms are deflected by multiple inhomogeneous magnetic fields in a sequence. Remarkably, unlike the classical case of light passing through polarizers, which filter out everything after two polarizers with perpendicular polarization axes, two quantum mechanical “beam splitters” with perpendicular magnetic field directions still allows two splitted beams to pass through, as illustrated in the middle and bottom parts of Fig. 1.4.

1.1.4 Davisson-Germer experiment demonstrated the wave nature of electrons

In 1924 de Broglie formally proposed the wave nature of electrons and other particles, in terms of their de Broglie wavelengths:

$$\lambda = \frac{h}{p} \quad (1.19)$$

where p is the momentum of the particle. If this proposal were true, it means that microscopic particles such as electrons should exhibit interference phenomena in the same way as acoustic and light waves.

In 1927 Davisson and Germer[3] performed an experiment to observe the diffraction pattern of electrons scattered by the surface of a crystal of nickel metal, as shown in Fig. 1.5. The highest intensity was observed at an angle $\theta = 50^\circ$ with a voltage of 54 V, giving the electrons a kinetic energy of 54 eV. To be compatible with the Bragg’s law

$$n\lambda = 2d \sin(90^\circ - \theta/2), \quad n \in \mathbb{Z}_+. \quad (1.20)$$

with the lattice spacing of nickel $d = 0.091$ nm, the $n = 1$ Bragg peak in (1.20) exactly corresponds to the de Broglie wavelength (1.19) of electrons. We will examine the experimental




Figure 1.4: Sequential Stern-Gerlach experiments. (Picture from S&N)




Figure 1.5: (Left) Schematic setup of Davisson-Germer experiment. (Right) Intensity of electron scattering vs. azimuth angle at 54 volts, co-latitude $\theta = 50^\circ$.

results in Fig. 1.5 more carefully in Homework 1.

1.2 Hilbert space and state vectors (8/28)

- Refer to N&C 2.1, Tong 3.1, Shankar 1.1-1.4

How to explain the above experimental observations? 4 postulates, derived (or more precisely, guessed) after a long process of trial and error, form the foundation for a theory framework that explains all observed experimental phenomena, known as quantum mechanics. Below we will introduce these postulates one by one, along the way of introducing linear algebra as the mathematics of quantum mechanics.

Postulate 1.1. *Associated to any isolated physical system is a complex vector space with inner product (i.e. a Hilbert space) known as the state space of the system. The system is completely described by its state vector, which is a unit vector in the system's state space.*

The key mathematical framework for quantum mechanics is linear algebra. The basic objects of linear algebra are vector spaces:

A *vector space* over a *scalar field* $\mathbb{F}$ is a set $\mathbb{V}$ of objects (called *vectors*) equipped two binary operations: a *vector addition* (“+”)

$$|v\rangle + |w\rangle \in \mathbb{V}, \quad \forall |v\rangle, |w\rangle \in \mathbb{V}. \quad (1.21)$$

and a *scalar multiplication*:

$$a|v\rangle \in \mathbb{V}, \quad \forall |v\rangle \in \mathbb{V}, a \in \mathbb{F}. \quad (1.22)$$

Mathematically, the set $\mathbb{V}$ forms an abelian group under vector addition operation and the scalar multiplication defines a ring homomorphism from the field $\mathbb{F}$ into the endomorphism ring of this abelian group. We will break down these terminology below.

It is worthwhile to define a group since we will study symmetry in quantum mechanics later. A group is a set $V = \{u, v, \dots\}$ equipped with a *closed* binary operation $+$: $V \times V \rightarrow V$

$$u + v = w \in V, \quad \forall u, v \in V. \quad (1.23)$$

where 3 group axioms are satisfied:

(1) *Identity element*:

$$\exists 0 \in V, \text{ s.t. } \forall u \in V, u + 0 = 0 + u = u. \quad (1.24)$$

(2) *Inverse*:

$$\forall u \in V, \exists (-u) \in V, \text{ s.t. } u + (-u) = 0. \quad (1.25)$$

(3) *Associativity*:

$$(u + v) + w = u + (v + w), \quad \forall u, v, w \in V. \quad (1.26)$$

An Abelian group is a group whose addition operation is commutative:

$$u + v = v + u, \quad \forall u, v \in V. \quad (1.27)$$

The scalar multiplication is another map $\mathbb{F} \times \mathbb{V} \rightarrow \mathbb{V}$ that is *associative*

$$a(b|v\rangle) = (ab)|v\rangle, \quad \forall a, b \in \mathbb{F}, |v\rangle \in \mathbb{V} \quad (1.28)$$

and *distributive in both the vectors and the scalars*:

$$a(|u\rangle + |v\rangle) = a|u\rangle + a|v\rangle, \quad (1.29)$$

$$(a + b)|v\rangle = a|v\rangle + b|v\rangle, \quad \forall a, b \in \mathbb{F}, |u\rangle, |v\rangle \in \mathbb{F}. \quad (1.30)$$

The above conditions satisfied by vector addition and scalar multiplication are summarized well in Chapter 1.1 of Shankar.

Of particular interests to quantum mechanics is a *Hilbert space*, which is a (n -dimensional) vector space $\mathbb{C}^n$ over the field $\mathbb{C}$ of complex numbers, equipped with a map $\mathbb{V} \times \mathbb{V} \rightarrow \mathbb{C}$ known as an *inner product*:

$$\langle u|v\rangle \in \mathbb{C}, \quad \forall |u\rangle, |v\rangle \in \mathbb{V}. \quad (1.31)$$

In the Dirac notation, $\langle u|$ is known as a *bra*, and $|v\rangle$ is known as a *ket*. The inner product is *skew-symmetric*

$$\langle u|v\rangle = \langle v|u\rangle^* \quad (1.32)$$

positive semidefinite

$$\langle v|v\rangle \geq 0, \quad \langle v|v\rangle = 0 \iff |v\rangle = 0. \quad (1.33)$$

and *linear in ket*

$$\langle u|(\sum_i \lambda_i |v_i\rangle) = \sum_i \lambda_i \langle u|v_i\rangle \quad (1.34)$$

These properties are summarized in Chapter 1.2 of Shankar and Chapter 2.1.4 of N&C.

Note that an inner product also defines a norm for each vector in vector space $\mathbb{V}$:

$$||v\rangle| \equiv \sqrt{\langle v|v\rangle} \quad (1.35)$$

Two important properties of an inner product are

(1) **Cauchy-Schwarz inequality**:

$$\langle v|v\rangle \langle u|u\rangle \geq |\langle u|v\rangle|^2 \quad (1.36)$$

The equality holds if and only if $|v\rangle$ and $|w\rangle$ are linearly dependent.

(2) **Triangle inequality** for the norms of vectors:

$$||v\rangle + |w\rangle| \leq ||v\rangle| + ||w\rangle| \quad (1.37)$$

Triangle inequality can be derived from the Cauchy-Schwarz inequality, which is a part of Homework 2.

Strictly speaking, in addition to being a vector space equipped with an inner product (known as an inner product space), a Hilbert space must also be *complete* in the sense that every Cauchy sequence $\{v^{(k)}\}$ of vectors in $\mathbb{V}$ has a limit that also lies in $\mathbb{V}$:

$$\lim_{k \rightarrow \infty} v^{(k)} \in \mathbb{V}, \quad \text{if } \forall \epsilon > 0, \quad \exists N \in \mathbb{Z}_+, \text{ s.t. } |v^{(m)} - v^{(n)}| < \epsilon, \quad \forall m, n > N. \quad (1.38)$$

For an inner product space of finite dimensions on any complete field $\mathbb{F}$, such as real and complex numbers $\mathbb{F} = \mathbb{R}, \mathbb{C}$, it is automatically a Hilbert space. Below are a few examples of Hilbert spaces, and inner product spaces that are not Hilbert spaces:

- n -dimensional Hilbert space on complex numbers: $\mathbb{V} \simeq \mathbb{C}^n$
- $n \times n$ complex matrices equipped with the Hilbert-Schmidt inner product forms a Hilbert space $L_{\mathbb{C}^n}$ (see Homework 1)
- n -dimensional Hilbert space on real numbers: $\mathbb{V} \simeq \mathbb{R}^n$
- n -dimensional vector space $\mathbb{V} \simeq \mathbb{Q}^n$ on rational numbers is not a Hilbert space because it is not complete, since a Cauchy sequence of rational numbers can converge to an irrational number.
- The inner product space of continuous square-integrable complex functions on interval $[a, b]$ with the following L^2 inner product:

$$\langle f_1 | f_2 \rangle \equiv \int_a^b f_1^*(x) f_2(x) dx \quad (1.39)$$

is not a Hilbert space due to its incompleteness: a Cauchy sequence of continuous functions can be discontinuous. Its completion is the Hilbert space $L^2[a, b]$.

- For the same reason of incompleteness, the inner product space of polynomials on $[a, b]$ with the same L^2 inner product is not a Hilbert space, even though it's a subspace of $L^2[a, b]$.

A Hilbert space forms the *state space* described in Postulate 1.1. With the help of the inner product structure, we can always choose a complete set of orthonormal basis $\{|i\rangle\}$ that expands the full state space $\mathbb{V}$, such that

$$|v\rangle = \sum_i v_i |i\rangle, \quad \forall |v\rangle \in \mathbb{V}; \quad \langle i | j \rangle = \delta_{i,j} \quad (1.40)$$

where $\delta_{i,j}$ is the Kronecker delta function. Given a n -dimensional vector space V , one can always find a set of n linearly independent vectors $\{e_j | 1 \leq j \leq n\}$, known as a basis of vector space V , such that any vector $v \in V$ can be uniquely expressed as a linear superposition of the basis vectors: $v = \sum_j v_j e_j$ with $v_j \in \mathbb{F}$. Given the inner product structure in a Hilbert space, a set of complete orthonormal basis can be constructed following the Gram-Schmidt procedure, discussed in details in Chapter 1.2 of Shankar and Chapter 2.1.4 of N&C. In such a complete orthonormal basis of Hilbert space $\mathbb{V} = \mathbb{C}^n$, a Dirac ket $|v\rangle$ can be represented by a column vector $\vec{v} = (v_1, \dots, v_n)^T$, a Dirac bra represented by a row vector, and the inner product reduces to the familiar form in $\mathbb{C}^n$:

$$\langle u | v \rangle = \sum_i u_i^* v_i = (u_1^*, \dots, u_n^*) \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix} \quad (1.41)$$

1.3 Linear operators and observables (9/2)

- Refer to N&C 2.1; Tong 3.2, 4.1; S&N 1.3-1.4; Shankar 1.5-1.6

Given two vector spaces $\mathbb{V}$ and $\mathbb{W}$, we can also define a linear operator $\hat{O}$ as any function that maps the vector space $\mathbb{V}$ to $\mathbb{W}$, which is linear in its inputs:

$$\hat{O}(\sum_i a_i |v_i\rangle) = \sum_i a_i \hat{O}(|v_i\rangle) \quad (1.42)$$

In the case of $\mathbb{V} = \mathbb{W}$ we call it a linear operator $\hat{O} \in \mathbb{L}_{\mathbb{V}}$ on vector space $\mathbb{V}$, which is of particular interest to quantum mechanics.

Using a complete set of orthonormal basis $\{|j\rangle | 1 \leq j \leq \dim \mathbb{V}\}$ for Hilbert space $\mathbb{V}$, each linear operator has a matrix representation:

$$\hat{O} = \sum_{a,b} O_{a,b} |a\rangle \langle b|, \quad O_{a,b} \equiv \langle a | \hat{O} | b \rangle \quad (1.43)$$

In particular, for the identity matrix which maps every vector to itself, we have the **completeness relation** for basis $\{|i\rangle\}$, also known as the **resolution of identity**:

$$\sum_i |i\rangle \langle i| = \hat{1}. \quad (1.44)$$

We can also define the product of two linear operators $\hat{M}, \hat{N} \in \mathbb{L}_{\mathbb{V}}$:

$$\hat{M}\hat{N} = \sum_{a,b,c} M_{a,b} N_{b,c} |a\rangle \langle c| \quad (1.45)$$

their commutator

$$[\hat{M}, \hat{N}] = \hat{M}\hat{N} - \hat{N}\hat{M} \quad (1.46)$$

and anti-commutator

$$\{\hat{M}, \hat{N}\} = \hat{M}\hat{N} + \hat{N}\hat{M} \quad (1.47)$$

and the *Hermitian adjoint* of an operator

$$\hat{O}^\dagger = \sum_{a,b} O_{b,a}^* |a\rangle \langle b| = (O^T)^* = (O^*)^T \quad (1.48)$$

Specifically in Postulate 1.2 about quantum (projective) measurements, each physical observable corresponds to a Hermitian linear operator, represented as a Hermitian matrix in the orthonormal basis:

$$\hat{M} = \hat{M}^\dagger = \sum_{a,b} M_{a,b} |a\rangle \langle b|, \quad (M^\dagger)_{a,b} = M_{b,a}^* = M_{a,b} \equiv \langle a | \hat{M} | b \rangle. \quad (1.49)$$

The diagonal form

$$\hat{O} = \sum_a O_a |a\rangle \langle a| \quad (1.50)$$

where $\{|a\rangle\}$ is a complete set of orthonormal basis of the Hilbert space, is also known as the *spectral decomposition* of a matrix $\hat{O}$. If there is a complete set of orthonormal basis $\{|a\rangle\}$ such

that the linear operator $\hat{O}$ can be rewritten as the diagonal form (1.50), the matrix $\hat{O}$ is called “diagonalizable”.

Not all matrices are diagonalizable (i.e. have a diagonal form), two non-diagonalizable examples are

$$\hat{M}_1 = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}, \quad \hat{M}_2 = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}, \quad (1.51)$$

They either have less eigenvectors than the dimension of the Hilbert space, or have at least two non-orthogonal eigenvectors with different eigenvalues.

Below are a few useful theorems on the diagonalization of matrices:

The spectral decomposition theorem states: A linear operator $\hat{O}$ on vector space $\mathbb{V}$ is diagonal w.r.t. some orthonormal basis of $\mathbb{V}$ if and only if operator $\hat{O}$ is *normal* i.e. $\hat{O}^\dagger \hat{O} = \hat{O} \hat{O}^\dagger$. (Refer to N&C Box 2.2).

Theorem: A normal matrix is Hermitian if and only if it has real eigenvalues.

Theorem: The eigenkets corresponding to different eigenvalues of a Hermitian matrix are orthogonal.

Theorem: A positive linear operator must be Hermitian.

Simultaneous diagonalization theorem: Two Hermitian linear operators $\hat{A}$ and $\hat{B}$ can be diagonalized simultaneously if and only if $[\hat{A}, \hat{B}] = 0$. In other words, there is a complete set of simultaneous eigenket $\{|a, b\rangle\}$ such that

$$\hat{A}|a, b\rangle = a|a, b\rangle, \quad \hat{B}|a, b\rangle = b|a, b\rangle. \quad (1.52)$$

if and only if $[\hat{A}, \hat{B}] = 0$.

An Hermitian operator can be diagonalized into a special orthonormal basis $\{|i\rangle\}$, known as its eigenbasis, in which the linear operator is represented by a diagonal matrix

$$\hat{M} = \sum_i \lambda_i |i\rangle \langle i| \quad (1.53)$$

If we use the set $\{m\}$ to label all different (real) eigenvalues of matrix $\hat{M}$, we can rewrite the above diagonal form (1.53) as

$$\hat{M} = \sum_m m \hat{P}_m, \quad \hat{P}_m = \sum_j \delta_{\lambda_j, m} |j\rangle \langle j| \quad (1.54)$$

where operators $\hat{P}_m$ are known as *projection operators* (or *projectors*) onto the eigenspace of $\hat{M}$ with eigenvalue m . Mathematically, a projector $\hat{P}$ is defined as a linear operator satisfying:

$$\hat{P}^2 = \hat{P} \quad (1.55)$$

You will learn more about projectors in HW2.

Given a state $|\psi\rangle$ and a physical observable $\hat{M}$, the (probability-weighted) *expectation value*

$\langle \hat{M} \rangle$ of operator $\hat{M}$ w.r.t. state vector $|\psi\rangle$ is

$$\langle \hat{M} \rangle \equiv \frac{\langle \psi | \hat{M} | \psi \rangle}{\langle \psi | \psi \rangle} \quad (1.56)$$

Naturally, the expectation value of any physical observable is a real number, since any physical observable corresponds to a Hermitian operator.

We note that the expectation value (1.56) of any observable is the same for two linearly dependent states $|\psi\rangle$ and $\alpha|\psi\rangle$ for any complex scalar $\alpha \in \mathbb{C}$. This means $|\psi\rangle$ and $\alpha|\psi\rangle$ describe the same physical state. In other words, in a N -dimensional Hilbert space $\mathbb{V} = \mathbb{C}^N$, for a generic state

$$|\psi\rangle \equiv \sum_{j=1}^N \psi_j |j\rangle \quad (1.57)$$

where $\{|j\rangle | 1 \leq j \leq N\}$ is a complete set of orthonormal basis of $\mathbb{V}$, although $\{\psi_j \in \mathbb{C} | 1 \leq j \leq N\}$ correspond to $2N$ independent real variables, only $2N - 2$ of them are physically measurable, because the global phase factor and normalization leads to 2 redundancies that cannot be physically observed.

In the simplest example of a single qubit, we have $\mathbb{V} = \mathbb{C}^2$ for the $N = 2$ -dimensional Hilbert space. There are 4 linearly independent Hermitian operators in $\mathbb{C}^2$, whose matrix representations are

$$\hat{1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \hat{\sigma}_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \hat{\sigma}_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \hat{\sigma}_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (1.58)$$

As you will learn in HW1, they (up to normalization) form a complete set of orthonormal basis for the Hilbert space $L_{\mathbb{C}^2}$ of linear operators. In the matrix representation, a generic state vector can be written as

$$|\psi\rangle = \alpha \begin{pmatrix} \cos \theta e^{i\phi} \\ \sin \theta \end{pmatrix}, \quad \alpha \in \mathbb{C} \quad (1.59)$$

θ and ϕ are the two variables in the state vector that can be physically measured. One can carry out measurements for two observables to extract θ and ϕ :

$$\langle \hat{\sigma}_z \rangle = \cos(2\theta), \quad \langle \hat{\sigma}_x \rangle = \sin(2\theta) \cos \phi \quad (1.60)$$

1.4 Projective measurements (9/4)

- Refer to N&C 2.2.5, Tong 3.4, 4.2, 16.4.1

Postulate 1.2. A *projective measurements* is described by a physical observable, $\hat{M} = \hat{M}^\dagger$ on the Hilbert space of the system being observed. The observable has a spectral decomposition

$$\hat{M} = \sum_m m \hat{P}_m \quad (1.61)$$

where $\hat{P}_m$ is the projector onto the eigenspace of M with eigenvalue m . The possible outcomes of the measurement correspond to the eigenvalues m of the observable. Upon measuring state $|\psi\rangle$, the probability of getting result m is given by

$$p(m) = \langle \psi | \hat{P}_m | \psi \rangle \quad (1.62)$$

If the outcome m occurred, the state of the quantum system collapsed into a new state

$$\hat{P}_m |\psi\rangle / \sqrt{p(m)} \quad (1.63)$$

immediately after the measurement.

Postulate 1.2 is in fact only a special form of quantum measurements, which will be discussed in detail in later parts of this course. Most generally, a generalized quantum measurement is described as follows:

Postulate 2b: A **generalized quantum measurement** is described by a collection $\{M_m\}$ of measurement operators. These are operators acting on the state space of the system being measured. The index m refers to the measurement outcomes that may occur in the experiment. If the (normalized) state of the quantum system is $|\psi\rangle$ immediately before the measurements, then the probability that result m occurs is given by

$$p(m) = \langle \psi | M_m^\dagger M_m | \psi \rangle \quad (1.64)$$

and the state of the system after the measurement is $\frac{M_m |\psi\rangle}{\sqrt{\langle \psi | M_m^\dagger M_m | \psi \rangle}}$. The measurement operators satisfy the completeness equation

$$\sum_m M_m^\dagger M_m = \hat{1}. \quad (1.65)$$

When we choose $\hat{M}_m$ to be projector $\hat{P}_m$ into the eigenspace of observable $\hat{M}$ with eigenvalue m , the above generalized measurement reduces to the a projective measurement. Now it also becomes clear that the expectation value of observable $\hat{M}$ w.r.t. to (normalized) state vector $|\psi\rangle$ is nothing but the probability-weighted average of measurement outcomes:

$$\langle \hat{M} \rangle \equiv \langle \psi | \hat{M} | \psi \rangle = \sum_m m \times p(m) \quad (1.66)$$

One can also find out the *standard deviation* of measuring observable $\hat{M}$ in state $|\psi\rangle$:

$$\Delta(M) \equiv \sqrt{\langle (\hat{M} - \langle \hat{M} \rangle)^2 \rangle} = \sqrt{\langle \hat{M}^2 \rangle - \langle \hat{M} \rangle^2} \quad (1.67)$$

A significant consequence of the collapse part in Postulate 1.2 is the following: starting from any state vector $|\psi\rangle$, if we measure the same observable $\hat{M}$ multiple times, the measurement

outcome does not change any more after the 1st measurement. This is because the state already collapse into an eigenstate of $\hat{M}$ with some eigenvalue m after the 1st measurement.

Another interesting aspect comes from *incompatible measurements* of physical observables that do not commute with each other. If we perform two measurements $\hat{A}$ and $\hat{B}$ that satisfy $[\hat{A}, \hat{B}] \neq 0$, for a generic quantum state, the measurement outcome depend on the order of the two measurements: i.e. measuring $\hat{A}$ after $\hat{B}$ versus $\hat{B}$ after $\hat{A}$ will yield different results. For example, starting from the eigenstate $|\uparrow\rangle$ of Pauli matrix σ_z , if we measure σ_z and then σ_x , there is a probability of 1/2 for us to end up with +1 outcomes for both measurements. However, if σ_x is measured before the σ_z measurement, there is only a probability of 1/4 of +1 outcomes for both measurements.

1.5 Unitary evolution of a closed quantum system (9/9)

- Refer to Tong 4.3; S&N 2.1; N&C 2.1.8, 2.2.2, 4.2; Shankar 4.3

In addition to Postulates 1.1 and 1.2 introduced earlier, we have two more postulates, one regarding how a quantum system evolves with time:

Postulate 1.3. *The time evolution of the state of a closed quantum system is described by the Schrodinger equation:*

$$i\hbar \frac{d|\psi\rangle}{dt} = \hat{H}|\psi\rangle \quad (1.68)$$

where $\hbar = h/2\pi \approx 1.05 \times 10^{-34}$ J.s is a constant, and $\hat{H}$ is a fixed Hermitian operator known as the Hamiltonian of the closed system.

Postulate 1.3 states that the state vector associated with a closed quantum system evolves following the Schrodinger equation (1.68). This postulate can be states in an equivalent manner:

Postulate 3b. *The evolution of a closed quantum system is described by a unitary transformation. That is, the state $|\psi(t_2)\rangle$ of the system at time t_2 is related to the state $|\psi(t_1)\rangle$ at time t_1 by a unitary operator U which only depends on the times t_1 and t_2 :*

$$|\psi(t_2)\rangle = U(t_1, t_2)|\psi(t_1)\rangle, \quad U(t_1, t_2) = e^{-i(t_2-t_1)\hat{H}/\hbar} \quad (1.69)$$

One can further generalize this to a time-dependent Hamiltonian $\hat{H}(t)$ where we have

$$U(t_1, t_2) = \mathcal{T} e^{-i \int_{t_1}^{t_2} \hat{H}(t) dt / \hbar} \quad (1.70)$$

where $\mathcal{T}$ denotes time ordering in the integral, as explained in S&N 2.1.

In linear algebra, a unitary operator $\hat{U}$ is defined as any linear operator satisfying

$$\hat{U}^\dagger \hat{U} = \hat{1}. \quad (1.71)$$

which also implies $\hat{U}\hat{U}^\dagger = \hat{1}$. Due to the spectrum decomposition theorem, a unitary operator can be diagonalized, and its eigenvalues all have modulus 1:

$$\hat{U}|\psi\rangle = e^{i\theta}|\psi\rangle \quad (1.72)$$

Three examples of unitary operators in quantum mechanics are the following: (1) unitary time evolution operator $U(t_1, t_2) = e^{-i(t_2-t_1)\hat{H}/\hbar}$ for a closed quantum system discussed above; (2) a symmetry operator in a quantum system as we will discuss later in this course; (3) a quantum (logic) gate in quantum computing. For instance, in a two-dimensional closed quantum system, also known as a qubit, the common quantum gates are summarized in Table 1.1.

In the case of unitary time evolution $U(t_1, t_2)$, the Hamiltonian $\hat{H}$ of a closed quantum system plays a crucial role. In its spectral decomposition:

$$\hat{H} = \sum_a E_a |a\rangle\langle a| \quad (1.73)$$

the eigenkets $|a\rangle$ are called the *energy eigenstates*, or *stationary states*, and the eigenvalue E_a is called the energy of state $|a\rangle$. In particular, the eigenstate with the lowest energy is known as the *ground state*, while other states with higher energy are known as *excited states*. According

Single-qubit gate	Symbol	Unitary matrix
Pauli-X (bit flip)	X	$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$
Pauli-Y	Y	$\begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$
Pauli-Z (phase flip)	Z	$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$
Hadamard	H	$\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$
Phase	S	$\begin{pmatrix} 1 & 0 \\ 0 & i \end{pmatrix}$
$\pi/8$	T	$\begin{pmatrix} 1 & 0 \\ 0 & e^{i\pi/4} \end{pmatrix}$

Table 1.1: Common single-qubit gates and associated unitary matrices.

to Schrodinger equation (1.68), a generic state vector

$$|\psi\rangle = \sum_a c_a |a\rangle \quad (1.74)$$

evolves as

$$|\psi(t)\rangle = e^{-i\hat{H}t/\hbar} |\psi\rangle = \sum_a c_a e^{-iE_a t/\hbar} |a\rangle \quad (1.75)$$

and the different oscillation frequency $\omega_a = E_a/\hbar$ in different components of state $|\psi(t)\rangle$ can result in oscillations of physical observables. Two such examples, i.e. (1) spin precession; and (2) neutrino oscillation are discussed in detail in S&N 2.1, and will be a part of HW3. In particular, a physical observable you will encounter a lot in the future is the *transition probability*:

$$P(i \rightarrow f) \equiv |\langle f | e^{-i\hat{H}t/\hbar} | i \rangle|^2 \quad (1.76)$$

which measures the probability that an initial state $|i\rangle$ evolves into a final state $|f\rangle$ after time t . This quantity is related to the outgoing beam intensity in scattering experiments.

1.5.1 Appendix: Functions of operators/matrices

A function $f(\hat{M})$ of a square matrix $\hat{M}$ can be defined by the following Taylor expansion

$$f(\hat{M}) = \sum_{n=0}^{\infty} \frac{\hat{M}^n}{n!} \frac{d^n f(x)}{dx^n} \Big|_{x=0} \quad (1.77)$$

if the Taylor expansion converges at $x = 0$. In quantum mechanics we will mostly encounter square matrices that can be diagonalized, such as Hermitian and unitary matrices. For a diagonalizable matrix $\hat{M}$ with eigenkets $\{|j\rangle\}$ and eigenvalues $\{\lambda_j\}$, the matrix can be written as

$$\hat{M} = \sum_j \lambda_j |j\rangle \langle j| \quad (1.78)$$

and a function of this diagonalizable matrix can be defined as

$$f(\hat{M}) = \sum_j f(\lambda_j) |j\rangle \langle j| \quad (1.79)$$

Name	Controlled-Z (CPHASE)	CNOT	SWAP
Unitary matrix	$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$	$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$	$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$

Table 1.2: Common 2-qubit gates and associated unitary matrices.

Name	Toffoli (Controlled-CNOT)	Fredkin (Controlled-SWAP)
Unitary matrix	$\frac{1+Z_1}{2} \otimes \text{CNOT}_{23} + \frac{1-Z_1}{2} \otimes \hat{1}_{23}$	$\frac{1+Z_1}{2} \otimes \text{SWAP}_{23} + \frac{1-Z_1}{2} \otimes \hat{1}_{23}$

Table 1.3: Common 3-qubit gates and associated unitary matrices.

1.6 Composite quantum systems and tensor product (9/11)

- Refer to N&C 2.1.7, 2.2.8, 4.3; Tong 3.6

Previously we have focused on a single closed quantum system. When two closed quantum systems are combined together e.g. through some interactions b/w the two systems, how to describe such as composite quantum system? The last postulate addresses this issue, under the mathematical frameworks of *tensor product*:

Postulate 1.4. *The state space of a composite physical system is the tensor product of the state spaces of the component physical systems. Moreover, if we have systems numbered 1 through n , and the system number i is prepared in the state $|\psi_i\rangle$, then the joint state of the total system is $|\psi_1\rangle \otimes |\psi_2\rangle \otimes \cdots \otimes |\psi_n\rangle$.*

Specifically, consider two quantum systems with Hilbert space V of dimension m and W of dimension n , their composite system is described by the tensor product of the two Hilbert spaces:

$$V \otimes W \equiv \{|v\rangle \otimes |w\rangle | |v\rangle \in V, |w\rangle \in W\} \quad (1.80)$$

which is a Hilbert space of dimension mn . Linear operators on $V \otimes W$ are given by $A \otimes B$, where A is a linear operator on V and B is a linear operator on W , satisfying

$$(A \otimes B)|v\rangle \otimes |w\rangle = A|v\rangle \otimes B|w\rangle \quad (1.81)$$

An inner product on $V \otimes W$ is naturally defined as

$$\left(\sum_i a_i |v_i\rangle \otimes |w_i\rangle, \sum_j b_j |v'_j\rangle \otimes |w'_j\rangle \right) = \sum_{i,j} a_i^* b_j \langle v_i | v'_j \rangle \langle w_i | w'_j \rangle \quad (1.82)$$

One can therefore define a complete set of orthonormal basis $\{|v_i\rangle \otimes |w_j\rangle\}$ for $V \otimes W$, and the matrix representation of linear operators in $V \otimes W$ is given by

$$(A \otimes B)_{i,j} = A_{i_1,j_1} B_{i_2,j_2}, \quad A \in L_V, B \in L_W. \quad (1.83)$$

Since $A \otimes B$ is a matrix of dimension mn , where m and n are dimensions of matrices A and B respectively, the row and column index of matrix $A \otimes B$ can be written as

$$i = (i_1 - 1)n + i_2, \quad j = (j_1 - 1)n + j_2. \quad (1.84)$$

More properties of tensor product can be found in N&C section 2.1.7.

With the help of tensor product, we can define Hamiltonian and unitary operators for a composite quantum system. We will study the transverse field Ising model for a 2-qubit system

$$H_{\text{TFIM}} = -JX_1 \otimes X_2 - h(Z_1 \otimes \hat{1}_2 + \hat{1}_2 \otimes Z_2) \quad (1.85)$$

as a demonstration. Due to a conserved quantity $Z_1 \otimes Z_2$ that commutes with the Hamiltonian, it can be diagonalized in the $Z_1 \otimes Z_2 = \pm 1$ subspaces respectively.

A good set of examples of unitary operators in a composite system are 2-qubit and 3-qubit quantum gates, shown in Table 1.2 and 1.3 respectively, such as a controlled- U gate in the 2-qubit system.

A Homework 1 (due 9/4)

A.1 Franck-Hertz experiment

In the Franck-Hertz experiment (see Fig. 1.2 in lecture notes), the sharp drop of electric current through the vacuum tube at a voltage of 4.9 V corresponds to the absorption of electron energy by the mercury atoms in the vacuum tube, where the atomic energy level of mercury atoms go through a transition. Compute the wavelength of light associated with this atomic energy level transition, show that it is approximately 254 nm, as observed in optical emission of the same vacuum tube.

A.2 Davisson-Germer experiment

(1) *Bragg's law* Consider the diffraction of a wave by a lattice of atoms with a lattice constant of d between layers. Derive the Bragg's law

$$n\lambda = 2d \cos \frac{\theta}{2}, \quad n = 0, 1, 2, 3, \dots \quad (\text{A.1})$$

for the diffraction pattern.

(2) Consider the Davisson-Germer experiment illustrated in Fig. 1.5 of overleaf lecture notes. For $\theta = 50^\circ$, $d = 0.091$ nm for nickel crystals, and an electron beam accelerated by a $V = 54$ volts voltage, show that it satisfies the Bragg's law with $n = 1$.

In part (1) and (2), we have dealt with a simplified version of the actual Davisson-Germer experiment[3], assuming the electron beam was scattered by a three-dimensional simple cubic lattice. Below we will seriously consider the actual situation of Davisson-Germer experiment in history, where an electron beam was scattered by the (111) surface of a nickel crystal in which Ni atoms form a face-centered cubic (FCC) lattice. In this case, we need to first derive the general Bragg condition for two-dimensional diffraction peaks and apply it to the setup of Davisson-Germer experiment. Below we carry out this calculation step by step.

(3) For electrons of energy 54 eV in the Davisson-Germer experiment, their penetration depth into the nickel crystal is less than 1 nanometer, comparable to the FCC lattice constant $a = 3.52$ Å. This means as a zeroth order approximation, we can model the Davisson-Germer experiment as electrons scattered by the top layer of Ni atoms parallel to the crystal surface. In Davisson-Germer experiment the incident beam of electrons is perpendicular to the cleaved (111) surface of the nickel crystal. Can you draw the two-dimensional lattice formed on the (111) surface of a nickel crystal? Given the FCC lattice constant $a = 3.52$ Å in nickel crystals, what is the distance $\tilde{a}$ between two nearest Ni atoms on its (111) surface?

(4) For the two-dimensional (2d) plane parallel to the (111) surface, we can choose the following two orthonormal vectors as two basis vectors for the 2d plane:

$$\hat{e}_1 = (-1, 1, 0)/\sqrt{2}, \quad \hat{e}_2 = (-1, -1, 2)/\sqrt{6}, \quad \hat{e}_1 \times \hat{e}_2 = (1, 1, 1)/\sqrt{3}. \quad (\text{A.2})$$

In this two-dimensional lattice on (111) surface, the periodic pattern of Ni atoms are described




Figure A.1: Scattering of a wave by a lattice of atoms with a lattice constant d .

by two primitive vectors

$$\vec{a}_1 = \tilde{a}\hat{e}_1, \quad \vec{a}_2 = \tilde{a}\frac{-\hat{e}_1 + \sqrt{3}\hat{e}_2}{2}. \quad (\text{A.3})$$

where $\tilde{a}$ is the distance $\tilde{a}$ between two nearest Ni atoms on its (111) surface. They are called primitive vectors in the sense that any lattice site position $\vec{R}$ in the 2d lattice is a summation of integer multiples of $\vec{a}_i$

$$\vec{R} = \sum_{j=1}^2 m_j \vec{a}_j, \quad m_j \in \mathbb{Z}. \quad (\text{A.4})$$

Most generally, the Bragg condition for diffraction peaks is given by

$$\sum_{\vec{R}} e^{i\vec{q}\cdot\vec{R}} \neq 0 \iff \vec{q} \cdot \vec{a}_j = 0 \pmod{2\pi}, \quad \forall j = 1, 2. \quad (\text{A.5})$$

where $\vec{q}$ is the in-plane component of electron momentum transfer before and after scattered by the nickel crystal:

$$\vec{q} \equiv (\vec{k}_{\text{out}} - \vec{k}_{\text{in}}) - \frac{(1, 1, 1)}{3} (\vec{k}_{\text{out}} - \vec{k}_{\text{in}}) \cdot (1, 1, 1). \quad (\text{A.6})$$

In later lectures we will come back to (A.5) when discussing Fourier transform as a change of basis (i.e. a unitary transformation). From (A.5), show that

$$\vec{q} = \sum_{j=1}^2 n_j \vec{b}_j, \quad n_j \in \mathbb{Z} \quad (\text{A.7})$$

A Homework 1 (due 9/4)

and where the *reciprocal lattice primitive vectors* $\vec{b}_{1,2}$ are defined by the following conditions

$$\vec{a}_i \cdot \vec{b}_j = 2\pi\delta_{i,j} \quad (\text{A.8})$$

Note that $\vec{b}_{1,2}$ are both perpendicular to (1,1,1) as well. Express $\vec{b}_{1,2}$ using $\tilde{a}$ and $\hat{e}_{1,2}$.

(5) In the Davisson-Germer experiment, the incident momentum $\vec{k}_{\text{in}}$ is perpendicular to the (111) surface i.e. $\vec{k}_i = \frac{2\pi}{\lambda}(1,1,1)/\sqrt{3}$ where λ is the wavelength of electrons with a kinetic energy of 54 eV. For elastic scatterings responsible for the diffraction pattern, we have

$$|\vec{k}_i| = |\vec{k}_f| = \frac{2\pi}{\lambda} \quad (\text{A.9})$$

In the experimental setup of Fig. 1.5 (left), using results (A.6)-(A.7) of the previous part, derive the following Bragg's law for Davisson-Germer experiment:

$$\lambda\sqrt{n_1^2 + n_1n_2 + n_2^2} = \sqrt{\frac{3}{8}}a \sin \theta, \quad n_{1,2} \in \mathbb{Z} \quad (\text{A.10})$$

Comment: specifically, if we choose $(n_1, n_2) = n(1, 0)$ or $n(0, 1)$ or $n(1, -1)$, Eq. (A.10) reduces to the more familiar one

$$n\lambda = d_{(111)} \sin \theta, \quad n \in \mathbb{Z}, \quad d_{(111)} = a\sqrt{3/8} \approx 2.16 \text{ \AA}. \quad (\text{A.11})$$

(6) Note that the in-plane momentum transfer $\vec{q}$ can be written as

$$\vec{q} = |q|(\cos \phi \hat{e}_1 + \sin \phi \hat{e}_2) \quad (\text{A.12})$$

where ϕ is the azimuth angle in Fig. 1.5 of the lecture notes. Use relations (A.6)-(A.9) to explain the ϕ -dependent peaks in the right panel of Fig. 1.5.

(7) **Bonus** Can you qualitatively explain in Fig. 1.5 right panel, why are the peak heights at $\phi = \frac{\pi}{2} \bmod 2\pi/3$ different from those at $\phi = \frac{\pi}{6} \bmod 2\pi/3$?

A.3 Hilbert space $\mathbb{L}_{\mathbb{V}}$ of linear operators on $\mathbb{V} = \mathbb{C}^n$

(1) Consider linear operators on complex vector space $\mathbb{V} = \mathbb{C}^n$, i.e. $n \times n$ complex matrices. Prove that they form a vector space, which we label as $\mathbb{L}_{\mathbb{V}}$. For the definition of a vector space, refer to Chapter 1.1 of Shankar.

(2) Prove that the following binary operation $(\cdot, \cdot)$ on $\mathbb{L}_{\mathbb{V}} \times \mathbb{L}_{\mathbb{V}}$ defined by

$$(A, B) \equiv \text{tr}(A^\dagger B), \quad A, B \in \mathbb{L}_{\mathbb{V}} \quad (\text{A.13})$$

is an inner product. For the definition of an inner product, refer to Shankar 1.2 or N&C 2.1.4. This is known as the Hilbert-Schmidt inner product on operators.

(3) Using the above Hilbert-Schmidt inner product, show that 3 Pauli matrices $\sigma_x, \sigma_y, \sigma_z$ defined below

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (\text{A.14})$$

and the 2×2 identity matrix $\hat{1}$ together can form a complete orthonormal basis for the vector space $\mathbb{L}_{\mathbb{C}^2}$ of all 2×2 complex matrices.

(4) **Bonus** Prove the following *completeness relation of Pauli matrices*:

$$\sum_{i=x,y,z} (\sigma_i)_{a,b} (\sigma_i)_{c,d} = \vec{\sigma}_{a,b} \cdot \vec{\sigma}_{c,d} = 2\delta_{a,d}\delta_{b,c} - \delta_{a,b}\delta_{c,d} \quad (\text{A.15})$$

Hint: use the conclusion of part (3), i.e. any 2×2 Hermitian matrix $\hat{M}$ can be written as

$$\hat{M} = \frac{\text{Tr}(\hat{M})}{2} \hat{1}_{2 \times 2} + \sum_{a=x,y,z} \frac{\text{Tr}(\hat{M}\sigma_a)}{2} \sigma_a \quad (\text{A.16})$$

B Homework 2 (due 9/11)

B.0.1 Pauli matrices

(1) Diagonalize the 3 Pauli matrices and find out their eigenvalues and eigenkets:

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (\text{B.1})$$

(2) For an arbitrary function $f(\cdot)$ from complex numbers to complex numbers, if $\vec{n}$ is a normalized vector in three dimensions and θ is a real number, prove the following relation

$$f(\theta \vec{n} \cdot \vec{\sigma}) = \frac{f(\theta) + f(-\theta)}{2} \hat{1} + \frac{f(\theta) - f(-\theta)}{2} \vec{n} \cdot \vec{\sigma} \quad (\text{B.2})$$

where $\vec{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$ are three Pauli matrices.

Hint: a function $f(\hat{M})$ of a matrix (operator) $\hat{M}$ can be defined by the Taylor expansion:

$$f(\hat{M}) = \sum_{n=0}^{\infty} \frac{\hat{M}^n}{n!} \left. \frac{d^n f}{dx^n} \right|_{x=0} \quad (\text{B.3})$$

B.0.2 Linear algebra

(1) Use the Cauchy-Schwarz inequality to prove the triangle inequality

$$||u\rangle + |v\rangle| \leq ||u\rangle| + ||v\rangle|, \quad \forall |u\rangle, |v\rangle \in V \quad (\text{B.4})$$

(2) A linear operator $\hat{P}$ is a projection operator (or a projector) if $\hat{P}^2 = \hat{P}$. Prove that the eigenvalues of a projector are all either 0 or 1.

(3) **Bonus** A projector has the following form:

$$\hat{P} = \alpha |1\rangle\langle 1| + \beta |2\rangle\langle 2| \quad (\text{B.5})$$

where $|1\rangle$ and $|2\rangle$ are two normalized vectors in the Hilbert space that are linearly independent. What conditions should coefficients α and β satisfy in order for $\hat{P}$ defined above to be a projector (i.e. to satisfy $\hat{P}^2 = \hat{P}$)?

(4) Assume Hamiltonian H commutes with two linear operators, A_1 and A_2 , which do not commute with each other:

$$[H, A_1] = [H, A_2] = 0, \quad [A_1, A_2] \neq 0 \quad (\text{B.6})$$

Show that the eigenstates of H are in general degenerate.

(5) Assume two observables A_1 and A_2 anticommute with each other, i.e.

$$\{A_1, A_2\} = A_1 A_2 + A_2 A_1 = 0 \quad (\text{B.7})$$

Show that for any eigenstate $|\psi_1\rangle$ of observable A_1 with a nonzero eigenvalue, its expectation

value w.r.t. A_2 must vanish

$$\langle \psi_1 | A_2 | \psi_1 \rangle = 0 \quad (\text{B.8})$$

Comment: one such example is Pauli matrices $A_1 = \sigma_x$ and $A_2 = \sigma_y$.

(6) Consider two observables $A_{1,2}$ that anticommute with each other. Is it possible to have simultaneous eigenket of A_1 and A_2 ? Explain.

(7) **Bonus** For two linear operators satisfying $[\hat{A}, \hat{B}] = \alpha \hat{1}$ where $\hat{1}$ is the identity operator, prove the following identity:

$$[\hat{A}, e^{\hat{B}}] = \alpha e^{\hat{B}} \quad (\text{B.9})$$

Hint: first show that

$$\hat{f}(x) = e^{-x\hat{B}} \hat{A} e^{x\hat{B}} = \hat{A} + x\alpha \hat{1} \quad (\text{B.10})$$

(8) For a quantum-mechanical particle in one dimension, its position $\hat{x}$ and momentum $\hat{p}$ operators satisfy the canonical commutation relation:

$$[\hat{x}, \hat{p}] = i\hbar = ih/2\pi \quad (\text{B.11})$$

Use the results of part (7) to show that for every $\hat{x}$ eigenstate $|x_0\rangle$ with eigenvalue x_0

$$\hat{x}|x_0\rangle = x_0|x_0\rangle \quad (\text{B.12})$$

the following state

$$|x_0 + a\rangle \equiv e^{-i\hat{p}a/\hbar}|x_0\rangle \quad (\text{B.13})$$

is also an eigenstate of $\hat{x}$ with eigenvalue $x_0 + a$.

Comment: physically this means the linear operator $\hat{T}(a) \equiv e^{-i\hat{p}a/\hbar}$ translate the position by a , from x_0 to $x_0 + a$.

B.0.3 Consecutive measurements of incompatible operators

Two consecutive projective measurements, for Pauli matrices $\hat{\sigma}_x$ and $\hat{\sigma}_z$ respectively, are carried out one after another on a quantum state of a single qubit (i.e. a Hilbert space $\mathbb{V} = \mathbb{C}^2$). In the first procedure, we first measured $\hat{\sigma}_x$ with an outcome $+1$ and probability $p(X = +1)$, and then measured $\hat{\sigma}_z$ with an outcome $+1$ with probability $p(Z = +1|X = +1)$. In the second procedure, we switched the order of the two measurements, i.e. we first measured $\hat{\sigma}_z$ with an outcome $+1$ and probability $p(Z = +1)$, and then measured $\hat{\sigma}_x$ with an outcome $+1$ with probability $p(X = +1|Z = +1)$. It is found that the probabilities do not depend on the order of the two incompatible measurements, i.e.

$$p(X = +1) = p(X = +1|Z = +1), \quad p(Z = +1|X = +1) = p(Z = +1) \quad (\text{B.14})$$

(1) Find a single-qubit pure state $|\psi\rangle$ that satisfies the condition (B.14).

(2) **Bonus** If the quantum state is a single-qubit mixed state, it is generally described by a density matrix of the following form:

$$\hat{\rho} = \frac{\hat{1} + \vec{r} \cdot \vec{\sigma}}{2}, \quad \vec{r} = (x, y, z) \in \mathbb{R}^3, \quad |\vec{r}| \leq 1. \quad (\text{B.15})$$

Find those mixed states that satisfy the same condition (B.14).

Comment: you will need to learn about measurements on mixed states in N&C 2.4.1.

B.0.4 Quantum Zeno effect

Consider a single qubit evolving under the following Hamiltonian:

$$\hat{H} = -\mu_B B \hat{\sigma}_z \quad (\text{B.16})$$

where $\mu_B = \hbar e / 2m_e$ is the Bohr magneton. We define time T as follows:

$$T = \frac{\pi m_e}{eB} \quad (\text{B.17})$$

Starting from the initial state

$$|\psi_0\rangle = |+\rangle = \frac{|\uparrow\rangle + |\downarrow\rangle}{\sqrt{2}} \quad (\text{B.18})$$

at time $t = 0$, we evolve the system by time T/n , followed by a measurement of $\hat{\sigma}_x$, and then repeat the same 2-step (evolution by $\frac{T}{n} + \hat{\sigma}_x$ measurement) procedure n times. What is the probability at time T (after n such 2-step procedures) to produce the measurement outcome of $\sigma_x = +1$?

Bibliography

- [1] J. R. Rydberg, The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science **29**, 331 (1890), <https://doi.org/10.1080/14786449008619945> .
- [2] W. Gerlach and O. Stern, Zeitschrift für Physik **9**, 349 (1922).
- [3] C. Davisson and L. H. Germer, Nature **119**, 558 (1927).
- [4] R. A. Millikan, Phys. Rev. **7**, 355 (1916).
- [5] A. H. Compton, Phys. Rev. **21**, 483 (1923).