

# Holomorphic Fields of Hilbert Spaces in Finite von Neumann algebra

WEI YUAN AND WENMING WU

ABSTRACT. Enter abstract here

## 1. INTRODUCTION

Let  $\mathcal{H}$  be a Hilbert space,  $Proj(\mathcal{H})$  be the set of self-adjoint projections and  $Uint(\mathcal{H})$  be the set of unitary operators. We will routinely identify a closed subspace with its associated orthogonal projection in  $B(\mathcal{H})$ . For a set  $\mathcal{L}$  of orthogonal projections in  $Proj(\mathcal{H})$ , we denote by  $Alg\mathcal{L}$  the set of all bounded linear operators on  $\mathcal{H}$  leaving each element in  $\mathcal{L}$  invariant. Then  $Alg\mathcal{L}$  is an unital weak-operator closed subalgebra of  $B(\mathcal{H})$ . Similarly, for a subset  $\mathcal{S}$  of  $B(\mathcal{H})$ , let  $Lat\mathcal{S}$  be the set of invariant projections for every operators in  $\mathcal{S}$ . Then  $Lat\mathcal{S}$  is a strong-operator closed lattice of projections. A subalgebra  $\mathcal{A}$  of  $B(\mathcal{H})$  is said to be reflexive if  $\mathcal{A} = AlgLat\mathcal{A}$ , similarly a lattice  $\mathcal{L}$  of projections is reflexive if  $\mathcal{L} = LatAlg\mathcal{L}$ .

## 2. HILBERT FIELDS OVER SPHERE

Let

$$\mathfrak{J}_1 = \begin{pmatrix} 0 & T_1 \\ 0 & I \end{pmatrix} \text{ and } \mathfrak{J}_2 = \begin{pmatrix} 0 & T_2 \\ 0 & I \end{pmatrix}$$

be two (unbounded) idempotents acting on  $\mathcal{H} \oplus \mathcal{H}$ , where  $T_1$  and  $T_2$  are two closed operators affiliated with a finite von Neumann subalgebra of  $\mathcal{B}(\mathcal{H})$ . The ranges of  $\mathfrak{J}_1$  and  $\mathfrak{J}_2$  are closed subspaces of  $\mathcal{H} \oplus \mathcal{H}$ .

Recall that the set of closed, densely defined operators affiliated with a finite von Neumann algebra  $\mathfrak{A}$  admits algebraic operations of addition and multiplication. In another word, the set is an unital  $*$  algebra (cf. [23]), and the elements in it can be manipulated as if they were bounded operators. In the rest of the paper, we will repeatedly make use this fact without mentioning it explicitly. For a elegant treatment of this subject, we refer readers to [24].

Let  $Q(\infty) = Ker(\mathfrak{J}_1)$ ,  $Q(0) = Ran(\mathfrak{J}_1)$  and  $Q(-1) = Ran(\mathfrak{J}_2)$  be there trace half projections in  $\mathfrak{A} \otimes M_2(\mathbb{C})$ . Note that

$$Alg(Q(\infty), Q(0), Q(-1)) = \{A | A\mathfrak{J}_i \subset \mathfrak{J}_i A, i = 1, 2\}.$$

By conjugating a unitary, we could assume that

$$T_1 - T_2 = \begin{pmatrix} K & 0 \\ 0 & 0 \end{pmatrix}$$

where  $K \geq 0$  and  $K$  has a closed inverse.

We also write  $T_1$  as  $\begin{pmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{pmatrix}$ . It is not hard to check that  $\mathcal{Alg}(Q(\infty), Q(0), Q(-1))$  contains the following elements

$$\begin{pmatrix} A_1 & 0 & T_{11}K^{-1}A_1K - A_1T_{11} & -A_1T_{12} \\ 0 & 0 & T_{21}K^{-1}A_1K & 0 \\ 0 & 0 & K^{-1}A_1K & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & A_2 & -A_2T_{21} & -A_2T_{22} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\begin{pmatrix} 0 & A_3 & -A_3T_{21} & -A_3T_{22} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & T_{12}D_1 & 0 \\ 0 & 0 & T_{22}D_1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & D_1 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & 0 & T_{12}D_2 \\ 0 & 0 & 0 & T_{22}D_2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & D_2 \end{pmatrix}$$

If  $Q \in \mathcal{Lat}(\mathcal{Alg}(Q(\infty), Q(0), Q(-1)))$  such that  $Q \wedge Q(\infty) = 0$  and  $Q \vee Q(\infty) = I$ , then there is an idempotent

$$\mathfrak{J} = \begin{pmatrix} 0 & T_1 + S_1 \\ 0 & I \end{pmatrix}$$

such that  $\mathfrak{J}A = A\mathfrak{J}$  for any  $A \in \mathcal{Alg}(Q(\infty), Q(0), Q(-1))$ . This implies that

$$\begin{pmatrix} 0 & 0 \\ 0 & I \end{pmatrix} \begin{pmatrix} S_{11} & S_{12} \\ S_{21} & S_{22} \end{pmatrix} = 0 \text{ and } \begin{pmatrix} S_{11} & S_{12} \\ S_{21} & S_{22} \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & I \end{pmatrix} = 0.$$

Therefore  $S_{12}$ ,  $S_{21}$  and  $S_{22}$  are all equals 0. Since

$$\begin{pmatrix} A_1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} S_{11} & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} S_{11} & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} K^{-1}A_1K & 0 \\ 0 & 0 \end{pmatrix} \quad \text{for all } A_1 \in \text{Ran}(K) \mathfrak{A} \text{Ran}(K).$$

This implies that  $S = zK$  for some  $z \in \mathbb{C}$ . We will denote this idempotent by  $\mathfrak{J}(z)$  and its range projection by  $Q(z)$ . Note that  $\mathfrak{J}_1 = \mathfrak{J}(0)$  and  $\mathfrak{J}_2 = \mathfrak{J}(-1)$ .

If  $Q \wedge Q(\infty) = 0$  and  $Q \vee Q(\infty) \neq I$ , then

$$E \equiv Q \vee Q(\infty) = \begin{pmatrix} I & 0 & 0 & 0 \\ 0 & I & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & I \end{pmatrix}.$$

Therefore, there must exist a  $\beta = (\xi_1, \xi_2, \xi_3, \xi_4)^T \in Q\mathcal{H}$  such that  $\xi_4 \neq 0$ . This implies that

$$\{(T_{12}\xi, T_{22}\xi, 0, \xi)^t | \xi\} \subset Q\mathcal{H}.$$

Consider the trace of  $Q$ , we know that

$$\{(T_{12}\xi, T_{22}\xi, 0, \xi)^t | \xi\} = Q\mathcal{H}, \text{ and } Q = \text{Ran}\left(\begin{pmatrix} 0 & T_1(I - \text{Ran}(T_1 - T_2)) \\ 0 & (I - \text{Ran}(T_1 - T_2)) \end{pmatrix}\right).$$

It is not hard to check that

$$Q(z_1) \wedge Q(z_2) = \{(T_{12}\xi, T_{22}\xi, 0, \xi)^t | \xi\}$$

for any  $z_1 \neq z_2 (\in \mathbb{C})$ .

If  $Q \wedge Q(\infty) \neq 0$ , then it is easy to see that

$$F \equiv Q \wedge Q(\infty) = \begin{pmatrix} I & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

If  $Q \neq F$ , we have  $E \leq Q \vee Q(\infty)$  and  $Q(z_1) \wedge Q(z_2) \leq Q$ . If  $Q \vee Q(\infty) = E$ , then  $\tau(Q) = \frac{1}{2}$ . Hence

$$Q\mathcal{H} = \{(\xi_1, T_{22}\xi, 0, \xi)^t | \xi_1, \xi\} = F \vee (Q(z_1) \wedge Q(z_2)).$$

The last possibility is  $Q \vee Q(\infty) = I$ . Then there must exists a vector  $(\xi_1, \xi_2, \xi_3, \xi_4)^T \in Q$  such that  $\xi_3 \neq 0$ . Note that  $\tau(Q) = \frac{1}{2} + \tau(F)$  and

$$\{(0, T_{21}\xi, \xi, 0)^t | \xi\} \subset Q\mathcal{H}.$$

Then it is not hard to see that

$$Q = \{(\xi_1, T_{21}\xi_2 + T_{22}\xi_3, \xi_2, \xi_3)^t | \xi_1, \xi_2, \xi_3\} = F \vee Q(z) = Q(z_1) \vee Q(z_2)$$

for any  $z_1$  and  $z_2 \in \mathbb{C}$ .

**Remark 2.1.**  $E = Q(\infty) \vee (Q(0) \wedge Q(-1))$  and  $F = (Q(0) \vee Q(-1)) \wedge Q(\infty)$ .

**Example 2.1** (Tautological line bundle over  $\mathbb{CP}^1$ ). Let

$$Q(\infty) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, Q(0) = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, Q(-1) = \begin{pmatrix} \frac{1}{2} & -\frac{1}{2} \\ -\frac{1}{2} & \frac{1}{2} \end{pmatrix},$$

then

$$Q(z) = \frac{1}{1+|z|^2} \begin{pmatrix} |z|^2 & z \\ \bar{z} & 1 \end{pmatrix}.$$

Note that the map  $\phi : \mathbb{CP}^1 \rightarrow S^2$  given by  $\phi([z_1, z_2]) = z_1/z_2$  is a homeomorphism. We have

$$\phi^*(Q)([z_1, z_2]) = \frac{1}{|z_1|^2 + |z_2|^2} \begin{pmatrix} |z_1|^2 & z_1 \bar{z}_2 \\ \bar{z}_1 z_2 & |z_2|^2 \end{pmatrix}$$

is a line bundle over  $\mathbb{CP}^1$ . It is the tautological line bundle (or Hopf bundle) over  $\mathbb{CP}^1$ .

If  $T_1$  and  $T_2$  are not bounded, then the connected component of  $\mathcal{Lat}(\mathcal{Alg}(Q(\infty), Q(0), Q(-1)))$  is not a Hilbert bundle.

**Definition 2.1** (Definition 2.2.1 and Definition 2.2.2 in [27]). Let  $X$  be a smooth manifold. A field of Hilbert spaces in a map  $p : \mathcal{H} \rightarrow X$ , with each fiber  $\mathcal{H}_x = p^{-1}(x)$  endowed with the structure of a Hilbert space. A smooth structure on a field  $\mathcal{H} \rightarrow X$  of Hilbert spaces is given by specifying a set  $\Gamma^\infty$  of sections of  $\mathcal{H}$ , closed under addition and under multiplication by elements of  $C^\infty(X)$ , and linear operators  $\nabla_\xi : \Gamma^\infty \rightarrow \Gamma^\infty$  for each  $\xi \in \text{Vect}(X)$ , such that for  $\xi, \eta \in \text{Vect}(X)$ ,  $f \in C^\infty(X)$  and  $\varphi, \psi \in \Gamma^\infty$

- (1)  $\nabla_{\xi+\eta} = \nabla_\xi + \nabla_\eta, \nabla_{f\xi} = f\nabla_\xi, \nabla_\xi(f\varphi) = (\xi f)\varphi + f\nabla_\xi\varphi;$
- (2)  $\langle \varphi, \psi \rangle \in C^\infty(X)$  and  $\xi \langle \varphi, \psi \rangle = \langle \nabla_\xi \varphi, \psi \rangle + \langle \varphi, \nabla_\xi \psi \rangle;$
- (3)  $\{\varphi(x) : \varphi \in \Gamma^\infty\} \subset \mathcal{H}_x$  is dense, for all  $x \in X$ .

Let  $\text{Proj}(1, \mathcal{H})$  denote the set of projections in  $\mathcal{B}(\mathcal{H})$  such that  $\dim P\mathcal{H} = 1$  for any  $P \in \text{Proj}(1, \mathcal{H})$ . Although  $\text{Proj}(1, \mathcal{H})$  is only a real analytic manifold, there exists a canonical complex structure on  $\text{Proj}(1, \mathcal{H})$ .

In general, an almost complex structure on a Banach manifold  $X$  is a smooth tensor field  $J$  on  $X$  which associates to each  $x \in X$  a linear operator  $J$  on the tangent space  $T_x(X)$  to  $X$  at  $x$  which satisfies  $J_x(J\xi) = -\xi$  for all  $\xi \in T_x(X)$ . An almost complex structure  $J$  on  $X$  is said to be a complex structure on  $X$  if and only if, given any point  $x \in X$ , there exists an open neighbourhood  $\Omega$  of  $x$  in  $X$  and a smooth map  $\Phi : \Omega \rightarrow B$ , where  $B$  is some complex Banach space which satisfies the following conditions:

- (1)  $\Phi$  maps  $\Omega$  diffeomorphically onto some open set in  $B$ ;
- (2) for each  $x \in \Omega$  the derivative  $(d\Phi)_x : T_m(\Omega) \rightarrow B$  of the map  $\Omega$  at  $x$  satisfies  $(d\Phi)_m(J_m\zeta) = i(d\Phi)_m\zeta$  for all  $\zeta \in T_m(X)$ .

We will describe the complex structure of  $\text{Proj}(1, \mathcal{H})$ . Fixed a unit vector  $\beta$  in  $P\mathcal{H}$ , it easy to see that the tangent space  $T_P\text{Proj}(1, \mathcal{H})$  at  $P$  is

$$\begin{aligned} T_P\text{Proj}(1, \mathcal{H}) &= \{H | H^* = H, PHP = 0, (I - P)H(I - P) = 0, H \in \mathcal{B}(\mathcal{H})\} \\ &= \{|\zeta\rangle\langle\beta| + (|\zeta\rangle\langle\beta|)^* | \zeta \in (I - P)\mathcal{H}\} \cong (I - P)\mathcal{H}. \end{aligned}$$

Let

$$J_P : \{|\zeta\rangle\langle\beta| + (|\zeta\rangle\langle\beta|)^* \rightarrow i|\zeta\rangle\langle\beta| + (i|\zeta\rangle\langle\beta|)^*$$

It is easy to see that  $J_P^2 = -I$ . Actually the operator is just the multiplication of  $i$  on the Hilbert space  $(I - P)\mathcal{H}$ . Let  $\Omega = \{Q | \langle Q\beta, \beta \rangle > 0\}$  and

$$\Phi : Q \rightarrow \frac{(I - P)Q\beta}{\langle Q\beta, \beta \rangle}.$$

Then

$$d\Phi : H \rightarrow H\beta, \quad \forall H \in T_P\text{Proj}(1, \mathcal{H}).$$

Now it obvious that  $(d\Phi)_P(J_P H) = iH\beta = i(d\Phi)_P H$  (Much general result is proved for the Flag manifolds in a  $C^*$ -algebra).

**Remark 2.2.** If we identify the projection with its range, then the complex structure described above is the same as the complex structure of the 1-dimensional Grassmann manifold  $\text{Gr}(1, \mathcal{H})$ , the set of all 1-dimensional subspaces of  $\mathcal{H}$ , as the complex projective space.

**Definition 2.2.** Suppose that  $\Phi : X \rightarrow \text{Proj}(1, \mathcal{H})$  is a holomorphic embedding, where  $X$  is a complex manifold. We say the embedding is locally constant with respect to a von Neumann algebra  $\mathfrak{A}$  if for any  $x \in X$

**Definition 2.3.** Let  $\mathfrak{A}$  be a von Neumann algebra acting on a separable Hilbert space  $\mathcal{H}$ . Suppose that  $\Phi : X \rightarrow \text{Proj}(1, \mathcal{H})$  is a holomorphic embedding, where  $X$  is a complex manifold. Then the field of Hilbert spaces induce by the holomorphic embedding and  $\mathfrak{A}$  is

$$\mathfrak{F}(\Phi, \mathfrak{A}) \equiv \coprod_{x \in X} \bigvee_{U \in U(\mathfrak{A})} U^* \Phi(x) U.$$

The following fact is immediate from the Definition 2.3.

**Proposition 2.1.** Let  $\mathfrak{A}$  be a von Neumann algebra acting on a separable Hilbert space  $\mathcal{H}$  and  $\Phi : X \rightarrow \text{Proj}(1, \mathcal{H})$  is a holomorphic embedding of a complex manifold. Then for any  $x \in X$ ,  $\mathfrak{F}(\Phi, \mathfrak{A})(x) \in \mathfrak{A}'$ .

### 3. CONCLUSION

#### REFERENCES

- [1] R. Kadison and John R. Ringrose, *Fundamentals of the theory of Operator Algebras*, **Volume II** (1997)
- [2] B. Fuglede and R. Kadison, *Determinant Theory in Finite Factors*, The Annals of Mathematics, Second Series, Vol. 55, No. 3 (1952), 520-530
- [3] M. Takesaki, *Theory of Operator Algebras*, **Volume I**, Encyclopedia of Mathematical Sciences, vol 124, Springer-Verlag, Berlin (2002)
- [4] R. H. Levene, S. C. Power *Manifold of Hilbert space projection*, Proc. London Math. Soc. 1-25 (2009)
- [5] Mohan Ravichandran, University of New Hampshire Ph.D. Thesis (2009)
- [6] L. Ge and W. Yuan, *Kadison-Singer algebras, II—General case*, to appear, 2009

- [7] Uffe Haagerup and Flemming Larsen, *Brown's Spectral Distribution Measure for R-diagonal Elements in Finite von Neumann Algebras*, 1999
- [8] Uffe Haagerup and Hanne Schultz, *Brown Measures of Unbounded Operators Affiliated with Finite von Neumann Algebras*, 1999
- [9] Hari Bercovici and Dan Voiculescu, *Free Convolution of Measures with Unbounded Support*,
- [10] K.S.KÖLBIG, *On The Value Of A Logarithmic-Trigonometric Integral*, BIT 11(1971), 21-28
- [11] Mehmet Koca, Ramazan Koc and Muataz Al-Barwani, *Breaking  $SO(3)$  into its closed subgroups by Higgs mechanism*, J.Phys.A: Math.Gen. 30(1997) 2109-2125
- [12] Beardon, Alan F, *The Geometry of Discrete Groups*, New York: Springer-Verlag GTM 91.
- [13] Alexandru Nica and Roland Speicher, *Lectures on the Combinatorics of Free Probability*, London Mathematical Society Lecture Note Series:335
- [14] Elias M. Stein and Rami Shakarchi, *complex analysis*, Princeton lectures in analysis II.
- [15] Chengjun Hou and Wei Yuan, *Minimal Generating Reflexive Lattices of Projections in Finite von Neumann Algebras*
- [16] W.Wu and W. Yuan, *On generators of abelian Kadison–Singer algebras in matrix algebras* Linear Algebra and its Applications, Vol 440, 2014, Pages 197–205.
- [17] R. C. Lyndon and J. L. Ullman *Groups of Elliptic Linear Fractional Transformations*, Proceedings of the American Mathematical Society, Vol. 18, No. 6, (1967) 1119-1124
- [18] S. Sakai, *On automorphism groups of  $II_1$ -factors*, Tôkoku Math. J. (2) 26 (1974), 423-430.
- [19] William L. Green and Anthony To-Ming Lau *Strong finite von Neumann algebras* , Math. Scand. 40 (1977), 105- 112.
- [20] M. Koca, R. Koc and M. Al-Barwani *Breaking  $SO(3)$  into its closed subgroups by Higgs mechanism*, J.Phys. A: Math. Gen. 30 (1997) 2109-2125
- [21] D. Voiculescu, K. Dykema and A. Nica, *Free Random Variables*, CMR Monograph Series 1, American Mathematical Society (1992)
- [22] A. Nica and R. Speicher, *Lectures on the Combinatorics of Free Probability*, London Mathematical Society Lecture Note Series: 335 (2006)
- [23] F. J. Murray and J. von Neumann, *On rings of operators, II*, Trans. Amer. Math. Soc. 41 (1937), 208 - 248
- [24] Z. Liu, *On Some Mathematical Aspects of The Heisenberg Relation* , Science China series Mathematics: Kadison's proceedings
- [25] Masoud Khalkhaili, *Basic Noncommutative Geometry*, EMS series of Lectures in Mathematics (2009)
- [26] N. Salinas *The Grassmann manifold of a  $C^*$ -algebra, and Hermitian holomorphic bundles*, Operator Theory: Advances and Applications 28, Birkhäuser Verlag Basel, 1998, 267-289.
- [27] L. Lempert and R. Szoke, *Direct Images, Fields of Hilbert spaces, and Geometric Quantization*
- [28] D. R. Wilkings, *The Grassmann manifold of a  $C^*$ -algebra*, Proc. of the Royal Irish Acad. 90A, 1990, 99-116.
- [29] M. Martin and N. Salinas *The Canonical Complex Structure of Flag Manifolds in a  $C^*$ -algebra*, Operator Theory: Advances and Applications, Vol.104, 1998, 173-187.

AMSS