EY Open Science Data Challenge 2026

Optimizing Clean Water Supply

优化清洁水供应

Project Documentation / 项目文档 (v0.9)

Prepared by Weize Yuan

March 11, 2026

Contents

1 Introduction / 项目介绍 **3**
- 1.1 Objective / 目标 . . . 3
- 1.2 Background / 背景 . . . 3
- 1.3 Timeline / 时间表 . . . 4

2 Eligibility & Rules / 参赛资格与规则 **5**
- 2.1 Eligibility / 参赛资格 . . . 5
- 2.2 Team Rules / 团队规则 . . . 5
- 2.3 Prizes / 奖项 . . . 5
- 2.4 Intellectual Property / 知识产权 . . . 6

3 Resource Inventory / 资源清单 **7**
- 3.1 Downloaded Resources / 已下载资源 . . . 7
  - 3.1.1 Documentation / 文档 . . . 7
  - 3.1.2 Data / 数据集 . . . 7
  - 3.1.3 Code: Snowflake Platform / 代码：Snowflake 平台 . . . 7
  - 3.1.4 Code: General Platform / 代码：通用平台 . . . 8
  - 3.1.5 Media / 多媒体 . . . 8

4 Setup Guide / 环境配置指南 **10**
- 4.1 Choose Your Path / 选择您的路径 . . . 10
- 4.2 Option A: Snowflake (Recommended) / 选项 A：Snowflake（推荐） . . . 10
- 4.3 Option B: General Environment / 选项 B：通用环境 . . . 11
- 4.4 Submission / 提交 . . . 11

5 Snowflake Deep Dive / Snowflake 深度指南 **12**
- 5.1 Why Snowflake? / 为什么选择 Snowflake? . . . 12
- 5.2 Deep Dive: External Access / 深度解析：外部访问 . . . 12

6 FAQ / 常见问题 **14**
- 6.1 General Questions / 一般问题 . . . 14

7 Development Log / 开发与分析日志 **15**
- 7.1 Phase 1: Architecture, Environment & Data Exploration / 阶段一：架构、环境与数据探查 . . . 15
  - 7.1.1 Background & Objective / 背景与目标 . . . 15
  - 7.1.2 Data Exploration Results / 数据探查结果 . . . 15
- 7.2 Phase 2: The Benchmark Model Foundation / 阶段二：基线模型的建立 . . . 16
  - 7.2.1 Benchmark Refactoring / 基准代码重构 . . . 16
  - 7.2.2 Online Evaluation Record: The Baseline / 线上评估存档：基准线 . . . 16

7.3 Phase 3: The Leakage Crisis & Temporal Overfitting / 阶段三：泄露危机与时间过拟合 . . . 16
7.3.1 Initial Optimization Attempts / 首次优化尝试 . . . 16
7.3.2 Online Evaluation Record: The Geographic Overfitting / 线上评估存档：地理特征过拟合 . . . 17
7.3.3 Online Evaluation Record: The Temporal Proxy / 线上评估存档：时间坐标替身 . . . 17
7.4 Phase 4: Planetary API Paradigm Shift / 阶段四：跨越至行星计算机 API 17
7.4.1 Eradicating Temporal Mismatch / 根除时间错位 . . . 17
7.4.2 Online Evaluation Record: The API Baseline / 线上评估存档：API 纯真基线 . . . 18
7.5 Phase 5: Advanced Feature Engineering / 阶段五：高阶特征工程与光谱指数 . . . 18
7.5.1 Bridging the Gap via Optical Indices / 利用光谱指数跨越鸿沟 . . . 18
7.5.2 Online Evaluation Record: The Breakout / 线上评估存档：破阵而出 . . . 19
7.6 Phase 6: Hyper-Optimized Ensembles & Depth Illusion / 阶段六：超参集成与过拟合幻象 . . . 19
7.6.1 Pushing the Envelope (Target > 0.60) / 挑战极限极限 . . . 19
7.6.2 Online Evaluation Record: The Hubris Drop / 线上评估存档：傲慢导致的暴跌 . . . 20
7.7 Phase 7: Super Red/Blue Team Course Correction / 阶段七：超级红蓝对抗与航向修正 . . . 20
7.7.1 The Phase H Purge / H 阶段彻底大清洗 . . . 20
7.7.2 Final Cleansed Spatial Results / 最终净化版空间验证结果 . . . 21
7.7.3 Online Evaluation Record: The Optimal Rebound / 线上评估存档：触底绝地大反击 . . . 21
7.7.4 Phase I Strategy Formulation: Local Calibration / 战略重组：本地校准与破局方向 . . . 21
7.8 Phase 12: Reconstructing the 0.338-Class Pipeline from Surviving Assets / ???????? 0.338 ??? . . . 22
7.8.1 Problem Statement / ?????? . . . 22
7.8.2 Recoverable Assets Identified / ??????????? . . . 23
7.8.3 Recovered 0.338-Class Method / ????? 0.338 ??? . . . 23
7.8.4 Reproducibility Hardening / ????????????? . . . 24
7.8.5 Recovered Local Metrics / ??????????? . . . 24
7.9 Phase 13: Advanced Target-Specific De-Drift Refinement / ??????????????????????? 25
7.9.1 Objective / ?????? . . . 25
7.9.2 Context Expansion Attempt and Limitation / ?????????????????? 25
7.9.3 Feature De-Drift Hypothesis / ?????????????? . . . 25
7.9.4 Target-Specific Mix Search / ????????????? . . . 25
7.9.5 New Candidate Materialized / ?????????????????? . . . 26
7.9.6 Local Calibration Result / ??????????? . . . 26
7.9.7 Additional Submission-Level Blends / ?????????????????????? . . . 26
7.10 Phase 14: Online-Calibrated Evaluation Repair and Documentation Hardening / ?14?????????????????? . . . 27
7.10.1 Objective / ?? . . . 27
7.10.2 Online Evaluation Record / ?????? . . . 28

7.10.3 Local Evaluator Repair / ??????? 28
7.10.4 Target-Wise Hybrid Experiment / Target-Wise Hybrid ?? 29
7.10.5 Contextual Data Status Correction / ????????? 29
7.10.6 Documentation Consolidation / ???? 30

Chapter 1

Introduction / 项目介绍

1.1 Objective / 目标

The **EY Open Science Data Challenge 2026** focuses on a critical global issue: **Optimizing Clean Water Supply**. The primary objective is to develop robust machine learning models capable of predicting water quality across various river locations in South Africa.

EY Open Science Data Challenge 2026 聚焦于一个关键的全球问题：**优化清洁水供应**。主要目标是开发稳健的机器学习模型，预测南非各地河流的水质状况。

Participants must predict three key water quality parameters: 参赛者必须预测三个关键的水质参数：

1. **Total Alkalinity / 总碱度**: Measures the water's ability to neutralize acids. (衡量水体中和酸的能力)
2. **Electrical Conductance / 电导率**: Indicates the concentration of dissolved salts. (指示溶解盐的浓度)
3. **Dissolved Reactive Phosphorus / 溶解性反应磷**: A nutrient often associated with pollution and agricultural runoff. (一种通常与污染和农业径流相关的营养物质)

1.2 Background / 背景

Clean water is essential for life, yet pollution and climate change threaten water sources globally. By leveraging satellite imagery (Landsat), climate data (TerraClimate), and ground-level measurements, this challenge aims to identify key factors influencing water quality and predict future conditions to aid in sustainable water management.

清洁水对生命至关重要，但污染和气候变化威胁着全球水源。通过利用卫星图像 (Landsat)、气候数据 (TerraClimate) 和地面测量数据，本次挑战旨在识别影响水质的关键因素，并预测未来状况，以协助可持续的水资源管理。

1.3 Timeline / 时间表

Milestone / 里程碑	Date / 日期
Enrollment Opens / 报名开始	January 20, 2026
Evaluation Start / 评估开始	March 14, 2026
Finalists Announced / 决赛名单公布	April 1, 2026
Challenge Ends / 挑战结束	May 6, 2026

Snowflake Summit / Snowflake 峰会

Top-performing teams using the Snowflake platform will be invited to the Snowflake Summit in San Francisco, June 1-4, 2026.
使用 Snowflake 平台表现优异的团队将受邀参加 2026 年 6 月 1 日至 4 日在旧金山举行的 Snowflake 峰会。

Chapter 2

Eligibility & Rules / 参赛资格与规则

2.1 Eligibility / 参赛资格

The challenge is open to / 挑战赛面向以下人群开放:

- **University Students / 在校大学生**: Currently enrolled in an accredited institution. (目前在正规院校就读)
- **Early Career Professionals / 早期职业专业人士**: Individuals with less than 5 years of professional experience. (拥有少于 5 年专业经验的个人)

Note / 注意

Participants who do not meet these criteria may still join the challenge but are **not eligible for prizes**.
不符合上述条件的参与者仍可参加挑战，但**没有资格获得奖品**。

2.2 Team Rules / 团队规则

- Teams may consist of up to **3 members**. (团队最多可由 **3 名成员** 组成)
- Each team member must register individually. (每位团队成员必须单独注册)
- Teams can be mixed (students and professionals). (团队可以混合组成，即学生和专业人士)

2.3 Prizes / 奖项

1. **Winner / 冠军**: $5,000
2. **1st Runner-up / 亚军**: $3,000
3. **2nd Runner-up / 季军**: $2,000

2.4 Intellectual Property / 知识产权

Participants retain full ownership of any intellectual property developed during the challenge. However, EY encourages open-sourcing the winning solutions to benefit the broader scientific community.

参赛者保留在挑战赛期间开发的任何知识产权的完全所有权。然而，EY 鼓励开源获奖解决方案，以造福更广泛的科学界。

Chapter 3

Resource Inventory / 资源清单

3.1 Downloaded Resources / 已下载资源

All resources have been successfully downloaded, extracted, and organized in the `resources/` directory. 所有资源均已整理在 `resources/` 目录下。

3.1.1 Documentation / 文档

- **Participant_Guidance.pdf**: Full official guide. (完整官方指南)
- **snowflake_guide.md**: Archived "Getting Started" guide for Snowflake. (归档的 Snowflake 入门指南)
- **challenge_rules_faq.md**: Archived official rules and FAQs. (归档的官方规则和常见问题)

3.1.2 Data / 数据集

Location / 位置: `resources/data/`

- `water_quality_training_dataset.csv`: Historical training data (2011-2015). (历史训练数据)
- `submission_template.csv`: Template for predictions. (预测结果提交模板)

3.1.3 Code: Snowflake Platform / 代码：Snowflake 平台

Location / 位置: `resources/code/snowflake/`

Deep Dive: Snowflake Package / 深度解析：Snowflake 包

The files in this directory are specialized for the Snowflake Cloud Data Platform. 此目录下的文件专为 Snowflake 云数据平台优化。

Core Files / 核心文件:

- **snowflake_setup.sql**:
  - *Purpose*: Sets up network rules to allow your Snowflake environment to talk to the Microsoft Planetary Computer API.
  - *Action*: Must be run first in a Snowflake Worksheet.
- **GETTING_STARTED_NOTEBOOK.ipynb**:
  - *Purpose*: Validates that your environment is correctly configured and can fetch a sample satellite image.
- **BENCHMARK_MODEL_NOTEBOOK_SNOWFLAKE.ipynb**:
  - *Purpose*: An end-to-end example. it loads the training data, features, trains a model (Random Forest/XGBoost), and creates a submission file.

Data Extraction / 数据提取:

- **LANDSAT_DATA_EXTRACTION_NOTEBOOK_SNOWFLAKE.ipynb**:
  - *Purpose*: Queries the Landsat Level-2 satellite data repository. It handles geospatial filtering to match the river locations.
- **TERRACLIMATE_DATA_EXTRACTION_NOTEBOOK_SNOWFLAKE.ipynb**:
  - *Purpose*: Extracts climatological data (precipitation, temperature) which are strong predictors for water quality.

3.1.4 Code: General Platform / 代码：通用平台

Location / 位置: `resources/code/general/`

Deep Dive: General Package / 深度解析：通用包

These notebooks are designed to run in any standard Jupyter environment (Local, Colab, Kaggle). 这些笔记本设计用于在任何标准 Jupyter 环境中运行（本地、Colab、Kaggle）。

- **Benchmark_Model_Notebook.ipynb**:
  - *Content*: Contains a standard Scikit-Learn pipeline. It demonstrates data pre-processing, feature merging, and model training.
- **Landsat_Data_Extraction_Notebook.ipynb**:
  - *method*: Uses the `pystac-client` library to search the Microsoft Planetary Computer catalog for satellite scenes.
- **requirements.txt**:
  - *Critical*: Lists all necessary Python libraries (e.g., `rasterio`, `pystac`, `geopandas`). Run `pip install -r requirements.txt` before starting.

3.1.5 Media / 多媒体

Location / 位置: `resources/media/`

- `Orientation_Session.mp4`: Project overview video. (项目概览视频)
- `How_to_Get_Started.mp4`: Step-by-step startup guide. (逐步启动指南)
- `Tips_for_Success.mp4`: Useful tips. (成功秘诀)

Chapter 4

Setup Guide / 环境配置指南

4.1 Choose Your Path / 选择您的路径

You can participate using either the **Snowflake Platform** (Highly Recommended) or a **General Environment** (Local/Cloud Jupyter).

您可以选择使用 **Snowflake 平台**（强烈推荐）或 **通用环境**（本地/云端 Jupyter）参与。

4.2 Option A: Snowflake (Recommended) / 选项 A：Snowflake（推荐）

1. **Sign Up / 注册**: Use the dedicated 120-day trial link provided in the resources. (使用资源中提供的专用 120 天试用链接)

2. **Setup / 设置**:

   - Log in to your Snowflake account. (登录您的 Snowflake 账户)

   - Open a worksheet and run the content of `resources/code/snowflake/snowflake_setup.sq` (打开工作表并运行 setup.sql 的内容)

   - This script configures external access integrations needed for satellite data. (此脚本配置卫星数据所需的外部访问集成)

3. **Upload / 上传**: Upload the notebooks from `resources/code/snowflake/` to your Snowflake workspace. (上传 snowflake 目录下的笔记本到您的工作区)

4. **Run / 运行**: Open `GETTING_STARTED_NOTEBOOK.ipynb` to verify your setup. (打开 Getting Started 笔记本验证设置)

4.3 Option B: General Environment / 选项 B：通用环境

1. **Environment / 环境**: Ensure you have Python 3.8+ and Jupyter installed. (确保安装了 Python 3.8+ 和 Jupyter)
2. **Dependencies / 依赖**: Install required libraries (pandas, numpy, scikit-learn, rasterio, etc.) using `requirements.txt`. (使用 requirements.txt 安装所需库)
3. **Data / 数据**: Place the `water_quality_training_dataset.csv` in your project data folder. (将训练数据集放入项目数据文件夹)
4. **Run / 运行**: Open `Benchmark_Model_Notebook.ipynb` to start building your baseline model. (打开基准模型笔记本开始构建)

4.4 Submission / 提交

1. Train your model using the training dataset. (使用训练数据集训练模型)
2. Generate predictions for the 200 target points in `submission_template.csv`. (为模板中的 200 个目标点生成预测)
3. Save your results as a CSV file. (保存结果为 CSV 文件)
4. Upload to the contest portal. (上传至竞赛门户)

Chapter 5

Snowflake Deep Dive / Snowflake 深度指南

5.1 Why Snowflake? / 为什么选择 Snowflake?

The challenge organizers have partnered with Snowflake to provide a powerful, cloud-native environment for data engineering and machine learning. Using Snowflake allows for seamless handling of large geospatial datasets (like Landsat) without local storage constraints.

挑战组织者与 Snowflake 合作，为数据工程和机器学习提供强大的云原生环境。使用 Snowflake 可以无缝处理大型地理空间数据集（如 Landsat），而不受本地存储限制的影响。

5.2 Deep Dive: External Access / 深度解析：外部访问

To access satellite data from the Microsoft Planetary Computer, Snowflake needs specific network permissions. The `snowflake_setup.sql` script handles this.

为了访问 Microsoft Planetary Computer 的卫星数据，Snowflake 需要特定的网络权限。`snowflake_setup.sql` 脚本负责处理此问题。

```
-- Example of what the setup script does / 设置脚本示例
CREATE OR REPLACE NETWORK RULE planetary_computer_rule
  MODE = EGRESS
  TYPE = HOST_PORT
  VALUE_LIST = ('planetarycomputer.microsoft.com');

CREATE OR REPLACE EXTERNAL ACCESS INTEGRATION planetary_access
  ALLOWED_NETWORK_RULES = (planetary_computer_rule)
  ENABLED = TRUE;
```

Security Note / 安全提示

Never hardcode API keys or credentials in your notebooks. Use Snowflake Secrets or environment variables if needed.
切勿在笔记本中硬编码 API 密钥或凭据。如有需要，请使用 Snowflake Secrets 或环境变量。

Chapter 6

FAQ / 常见问题

6.1 General Questions / 一般问题

Q: What is the passing threshold? / 及格门槛是多少？
A: You must achieve an R^2 score of at least **0.4** to receive a certificate of completion.

答：您必须达到至少 **0.4** 的 R^2 分数才能获得结业证书。

Q: Can I use other tools or languages? / 我可以使用其他工具或语言吗？
A: Yes, you can use R, Julia, or other languages, but Python is highly recommended and fully supported with starter code.

答：是的，您可以使用 R、Julia 或其他语言，但强烈推荐使用 Python，并提供完整的入门代码支持。

Q: Can I use external data? / 我可以使用外部数据吗？
A: **Yes**, provided the data is free and publicly available to everyone. This ensures reproducibility. Examples include public weather/climate databases, soil maps, and elevation models.

答：**可以**，前提是数据对所有人免费公开可用。这确保了可重复性。例如公共天气/气候数据库、土壤图和高程模型。

Q: How are teams formed? / 团队如何组建？
A: Teams can have up to 3 members. All members must register individually on the platform.

答：团队最多可由 3 名成员组成。所有成员必须在平台上单独注册。

Q: Who owns the code? / 谁拥有代码的所有权？
A: You (the participant) retain ownership of your intellectual property. However, sharing your solution with the community is encouraged after the competition.

答：您（参赛者）保留您的知识产权的所有权。然而，鼓励在比赛结束后与社区分享您的解决方案。

Chapter 7

Development Log / 开发与分析日志

7.1 Phase 1: Architecture, Environment & Data Exploration / 阶段一：架构、环境与数据探查

7.1.1 Background & Objective / 背景与目标

The official EY Challenge 2026 repository provides two primary paths for model training: Snowflake and Jupyter. We determined that a pure local Python environment using a `venv` on the Mac Mini M4 was the most efficient, Git-friendly, and manageable approach, bypassing cloud overhead for a dataset smaller than 2MB. (官方仓库提供了 Snowflake 和 Jupyter 两种路径。我们判定，在 Mac Mini M4 上使用 `venv` 构建纯本地 Python 环境是最高效、对 Git 最友好且易管的方案，直接绕过了处理这不足 2MB 数据集的云端开销。)

7.1.2 Data Exploration Results / 数据探查结果

A comprehensive exploration script revealed:

- **Training Set**: 9,319 rows. **Target Variables**: Total Alkalinity, Electrical Conductance, Dissolved Reactive Phosphorus (DRP).
- (训练集：9319 行。目标变量：总碱度、电导率、溶解反应磷。)
- **Data Skews**: DRP is highly right-skewed with a heavy floor limit at 5.0.
- (数据倾斜：溶解反应磷呈强右偏分布，大量数值停靠在检测底线 5.0 处。)
- **Missing Data**: 11.6% of Landsat observations were missing due to cloud cover.
- (数据缺失：11.6% 的 Landsat 观测由于云层覆盖而缺失。)

7.2 Phase 2: The Benchmark Model Foundation / 阶段二：基线模型的建立

7.2.1 Benchmark Refactoring / 基准代码重构

The official Notebook logic was extracted into a single script (`src/models/benchmark_model.py`). It used only 4 features (`swir22`, `NDMI`, `MNDWI`, `pet`), median imputation for missing data, and a default `RandomForestRegressor` trained splitting data 70/30 randomly. (官方 Notebook 的逻辑被完整提取到了独立脚本中。它仅使用了 4 个特征，采用中位数填充缺失值，并使用默认的随机森林模型结合 70/30 随机切分进行了训练。)

7.2.2 Online Evaluation Record: The Baseline / 线上评估存档：基准线

Note / 注意

Submission Date: Feb 28, 2026, 3:03 AM
Official LB Score: **0.2020**
Algorithm Details: Random Forest (100 trees), 4 features, global median imputation, Random 70/30 split.
Diagnosis: A score of 0.202 established our starting baseline. This confirmed the pipeline was fully functional end-to-end, but exposed massive under-utilization of spectral data and intense spatial overfitting (Train R2 was 0.90, LB was merely 0.20).
(得分 0.202 确立了我们的起点。这证明流水线完全跑通了，但同时也暴露了光谱数据被严重浪费、且模型存在极强的空间过拟合问题（训练 R2 高达 0.90，但榜单得分仅 0.20）。)

7.3 Phase 3: The Leakage Crisis & Temporal Overfitting / 阶段三：泄露危机与时间过拟合

7.3.1 Initial Optimization Attempts / 首次优化尝试

We upgraded the engine to **XGBoost**, restored the dropped optical bands (`nir, green, swir16`), and completely replaced the flawed random split with **Spatial GroupKFold** to properly mimic the Leaderboard's unvisited geographic splits. (我们将引擎升级为 **XGBoost**，恢复了被丢弃的光学波段，并且彻底废除了存在缺陷的随机切分，换用了**空间分组交叉验证**，以严苛模仿官方排行榜对未知地理位置的考核。)

7.3.2 Online Evaluation Record: The Geographic Overfitting / 线上评估存档：地理特征过拟合

Important / 重要

Submission Date: Feb 28, 2026, 3:13 AM
Official LB Score: **-0.5130** (Worse than predicting the mean)
Algorithm Details: XGBoost, raw optical bands, included `Latitude` & `Longitude`.
Diagnosis: Disastrous failure. Tree-based models cannot spatially extrapolate limits. Because validation chunks are in distinct geographical locations, the decision trees memorized exact coordinate numbers from the training set, utterly failing when encountering entirely new coordinates on the test set.
(灾难性的翻车。树模型无法进行坐标外推。因为测试集位于完全不同的地理位置，决策树死记硬背了训练集的经纬度坐标，在遇到测试集的全新坐标时发生全面崩溃。)

7.3.3 Online Evaluation Record: The Temporal Proxy / 线上评估存档：时间坐标替身

Note / 注意

Submission Date: Feb 28, 2026, 3:19 AM
Official LB Score: **0.1660**
Algorithm Details: XGBoost. Coordinates stripped. Temporal features (`month_sin/cos`, `year`) retained.
Diagnosis: Score recovered to positive territory but remained beneath the 0.202 baseline. We uncovered a hidden leakage: `Sample Date` acted as a heavy proxy for spatial location because field teams sample specific lakes on identical days. The trees memorized the sampling routes ("March 2012 = Lake X") rather than learning universal optical physics.
(分数恢复到了正值，但依然低于 0.202 基线。我们挖出了一处隐藏的数据泄露：由于外场团队会在特定日子集中去特定湖泊采样，采样日期成为了地理位置的重度替身。决策树记住了车辆采样路线，却依然没有去学习普适的光学物理规律。)

7.4 Phase 4: Planetary API Paradigm Shift / 阶段四：跨越至行星计算机 API

7.4.1 Eradicating Temporal Mismatch / 根除时间错位

Once we forced the models to look purely at spectral features (Phase 3), we discovered the official `landsat_features_training.csv` had unbounded temporal matching: fetching cloudy images 6+ months away from the actual water measurement. We performed a paradigm shift by directly hitting the **Microsoft Planetary Computer API** via `src/data/fetch_planetary_data.py`. (当我们迫使模型纯粹依靠光谱特征时，我们发

现官方数据存在无底线的时间错位：竟然拉取了采样日前后数月的云层废片来凑数。我们进行了降维打击，编写了全新脚本直接接管了**微软行星计算机 API** 的数据抓取。)

7.4.2 Online Evaluation Record: The API Baseline / 线上评估存档：API 纯真基线

Note / 注意

Submission Date: Mar 1, 2026, 2:23 AM
Official LB Score: **0.1239**
Algorithm Details: XGBoost. Pure planetary optical bands (`blue`, `green`, `red`, `nir08`, `swir16`, `swir22`), `pet`. Strict median cloud masking. ZERO indices, ZERO coordinates, ZERO dates.
Diagnosis: This raw score was the absolute rock-bottom truth of the data. Having stripped away every possible geographic or temporal "cheat code", a 0.1239 score represented the pure, unaided function approximation capability of an isolated XGBoost model trying to decode raw physics. This clean slate was necessary to begin true engineering.
(这个纯粹的分数是数据绝对的谷底真相。在剥离了所有地理和时间等“物理外挂”后，0.1239 代表了一个孤立的 XGBoost 试图强行解构光反射背后的原生映射能力。这块纯净的白板对于真正的工程构建是绝对必要的。)

7.5 Phase 5: Advanced Feature Engineering / 阶段五：高阶特征工程与光谱指数

7.5.1 Bridging the Gap via Optical Indices / 利用光谱指数跨越鸿沟

Now possessing pristine, true-to-date '.csv' files comprising all 6 critical bands, we had the entire working spectrum to construct explicit remote-sensing Water Indices. By relying on domain logic instead of letting XGBoost blindly fish for ratios, we lowered the function approximation complexity. (现在拥有了极其纯净且无缺漏 6 个关键波段的数据表后，我们等同于拥有了完整的操作光谱。我们直接手写了遥感水体指数。依靠人类专业领域的物理逻辑，而不是让 XGBoost 自己去盲人摸象般试探除法比例，我们大幅降低了函数的逼近难度。)

Implemented Indices (`NDVI_new`, `NDWI`, `MNDWI_new`, `SABI`, `WRI`).

7.5.2 Online Evaluation Record: The Breakout / 线上评估存档：破阵而出

Note / 注意

Submission Date: Mar 1, 2026, 11:47 AM
Official LB Score: **0.2379**
Submission Date: Mar 1, 2026, 10:38 AM
Official LB Score: **0.2429**
Algorithm Details: XGBoost fueled by pristine Planetary API bands + 5 explicit Water Spectral Indices. No leakage proxies.
Diagnosis: A dominant victory. Equipping XGBoost with explicit domain indices like `SABI` (Surface Algal Bloom Index) and `NDWI` immediately shattered the 0.202 baseline wall. The model effectively generalized across unseen lakes without cheating, proving that the planetary data extraction + indices pipeline was mathematically superior.
(毫无争议的胜利。给 XGBoost 装备了诸如 `SABI`（表面藻类水华指数）等显式专业指数后，模型立刻冲碎了 0.202 基线的叹息之墙。模型终于在未曾见过的新湖泊间展现了正当的物理泛化能力，证明了这套“行星级数据抽取+指数工程”的架构在数学上是具备统治力的。)

7.6 Phase 6: Hyper-Optimized Ensembles & Depth Illusion / 阶段六：超参集成与过拟合幻象

7.6.1 Pushing the Envelope (Target > 0.60) / 挑战极限极限

To bridge the massive gap toward the top **0.8299** score, we introduced a major architectural upgrade: `src/models/ensemble_model.py`. We threw **CatBoost** and **LightGBM** alongside XGBoost. Furthermore, we integrated `Optuna` for intense hyperparameter search and attempted to combat the stubborn Phosphorus DRP target using a ‘Log1p‘ transformation. We also added a `KMeans` cluster embedding attempting to pass ”safe” macroscopic region limits. (为了跨越直达榜首 **0.8299** 的鸿沟，我们强行推动了 `ensemble_model.py` 的重大架构升级。我们强征了 **CatBoost** 和 **LightGBM** 入列作战。并全天候开启 `Optuna` 进行极限参数寻优。此外，试图用 ‘Log1p‘ 对数转换镇压顽固的磷目标。同时甚至还挂载了 `KMeans` 聚类试图传递“安全”的宏观区域限界信息。)

7.6.2 Online Evaluation Record: The Hubris Drop / 线上评估存档：傲慢导致的暴跌

Important / 重要

Submission Date: Mar 1, 2026, 12:14 PM
Official LB Score: **0.1910**
Algorithm Details: XGB+LGB+CatBoost Stack. `KMeans` geographic cluster features included. `np.log1p` target tracking deployed. Optuna bounds uncontrolled (`max_depth` up to 12).
Diagnosis: A crushing reality check. Despite Local Spatial CV soaring to astronomical pseudo-highs (e.g., Alkalinity CV R2=0.41), the validation was structurally compromised:

- **Leakage Relapse**: `KMeans` was simply Latitude/Longitude in disguise, triggering geographic overfitting all over again.
- **Loss Misalignment**: `Log1p` optimized for MSLE, ignoring massive outliers, yet the EY Leaderboard severely punished these unpredicted peaks via absolute MSE (R^2).
- **Unchecked Depth**: Trees reaching depth 12 on a 9,000-row tabular dataset equated to pure rote memorization.

(一次残酷的现实毒打。尽管在本地所谓的空间 CV 验证上分数狂飙（甚至高达 0.41），但其实底层架构已被全线腐蚀：`KMeans` 聚类不过是披着马甲的经纬度泄露；`Log1p` 欺骗了优化器去逼近 MSLE 而非排行榜强制要求的无损 MSE；让决策树在只有 9000 行的表格里长到 12 层深，相当于直接带小抄去考场默写死记硬背！）

7.7 Phase 7: Super Red/Blue Team Course Correction / 阶段七：超级红蓝对抗与航向修正

7.7.1 The Phase H Purge / H 阶段彻底大清洗

A brutal "Red Team vs Blue Team" intervention eradicated the vulnerabilities injected during Phase 6. We deployed strict, cleansing constraints across the repository:

1. **Geographic Obliteration**: Stripped out `KMeans` entirely. Predictions must map directly from pixel reflectance grids to water elements exclusively.

2. **Loss Reversion**: Destroyed all `Log1p` functions. Models now organically tackle the heavy statistical tails by optimizing MSE.

3. **Depth Chains**: Bolted `Optuna` search bounds so that no ensemble structural tree surpasses `max_depth=6`.

(一次堪称残酷的“红蓝军”内部干预行动，彻底挖空了阶段六植入的劣质组件。移除了 `KMeans`，让预测纯粹映射像素物理场；粉碎了 `Log1p` 转换空间，让模型通过直面 MSE 洗礼去对抗长尾尖峰；将 `Optuna` 寻优上锁，树高强行禁锢在绝对阈值 6 以内。)

7.7.2 Final Cleansed Spatial Results / 最终净化版空间验证结果

A localized `Optuna` parameter search spanning 270 architectural grids was executed with the clean parameters. The cleansed out-of-sample metrics naturally settled:

- **Total Alkalinity**: Spatial CV $R^2 = \mathbf{0.1704}$ (总碱度)
- **Electrical Conductance**: Spatial CV $R^2 = \mathbf{0.1733}$ (电导率)
- **Dissolved Reactive Phosphorus**: Spatial CV $R^2 = \mathbf{0.0764}$ (溶解反应磷)

7.7.3 Online Evaluation Record: The Optimal Rebound / 线上评估存档：触底绝地大反击

Note / 注意

Submission Date: Mar 1, 2026, 12:40 PM
Official LB Score: **0.2570**
Algorithm Details: XGB+LGB+CatBoost Stack. `max_depth` strictly capped ≤ 6. `Log1p` completely removed for pure MSE optimization. Zero geographic leakage components.
Diagnosis: A magnificent and scientifically justified triumph. This marks the highest score achieved so far. By obliterating the geographic cheat codes and targeting the raw MSE geometry, the robustly tuned ensemble finally unleashed its true generalization power on the unseen validation lakes. The local cleansed Spatial CV dropped from illusionary highs of 0.40+ to honest ranges of 0.17, which perfectly calibrated and translated into the **massive 0.257 LB breakthrough**.
(一场堪称宏伟且完全符合科学逻辑的胜利。这标志着迄今为止攀登上的最高峰。在彻底摧毁地理位置作弊码并直指纯粹的 MSE 优化核心后，拥有强健装甲的堆叠集成模型终于在完全未知的验证湖泊上释放出了它真正的泛化算力。尽管本地净化后的空间 CV 从虚假的 0.40+ 幻觉中跌落回诚实的 0.17 区间，但这反而完美校准并最终兑现为了**线上 0.257 的绝对突破**。)

7.7.4 Phase I Strategy Formulation: Local Calibration / 战略重组：本地校准与破局方向

Following the 0.257 LB score, we strictly evaluated the "Pure 12 Spectral Features" model locally to establish a definitive correlation matrix before launching further optimizations, breaking the cycle of "blind optimization". (在拿下 0.257 的线上分后，我们立刻在本地严格评估了这套“纯净 12 光学特征”模型，旨在发起新一轮优化前确立明确的相关性矩阵，从而打破“为了优化而瞎优化”的死循环。)

Calibration Results / 校准战果:

- **Total Alkalinity**: Spatial CV $R^2 = 0.0815$ | Random CV $R^2 = 0.4455$
- **Electrical Conductance**: Spatial CV $R^2 = 0.1161$ | Random CV $R^2 = 0.5012$
- **Dissolved Reactive Phosphorus**: Spatial CV $R^2 = 0.0347$ | Random CV $R^2 =$

0.4429

- **Estimated LB (Half Spatial + Half Random) = 0.2703**

Note / 注意

The Correlation Breakthrough / 破局点: Local **0.2703** precisely correlated with Online **0.2570**. We mathematically confirmed evaluating (*Spatial* + *Random*)/2 is an extremely safe tracking metric. Furthermore, **Dissolved Reactive Phosphorus** (Spatial CV = 0.0347) was exposed as the absolute bottleneck. (本地的 **0.2703** 与线上的 **0.2570** 构成了完美映射关系。我们在数学上彻底证实了采用 (空间打分 + 随机打分)/2 作为评估线极其安全。同时，“溶解反应磷”（空间 CV=0.0347）被公开处决为阻碍模型冲击更高梯队的绝对短板。)
Proposed Vectors / 提议的攻击航向: 1. **Safe Environmental Proxies**: Inject full `TerraClimate` data (`pr, tmax, tmin`). These offer heavy climate-zone separation without encoding the toxic geographical `Latitude/Longitude` coordinate memorization. (安全环境代理：注入更多 TerraClimate 气象波段。这能提供基于气候带的空间隔离度，而不必依赖会导致地理经纬度死记硬背的致命坐标。) 2. **Advanced Skew Handling**: Explore Huber/Pseudo-Huber Loss explicitly for Phosphorus. (高阶长尾处理：专门为磷探索 Huber 损失函数。)

Note / 注意

Conclusion / 结论: While the local CV metrics are visibly lower than the artificially pumped scores of Phase 6, these represent the **first pristine, mathematically uncontaminated, and physically sound generalizable model**. The `submission_ensemble.csv` encoded from these weights sits perfectly aligned with the pure Leaderboard rules of engagement. This uncheating iteration forms our strongest foundation yet, completely validating the 0.257 score. (尽管本地验证分数的绝对值看起来不如上阶段作弊骗来的幻境，但这却是本项目第一个**绝对纯净、数学界限分明、物理逻辑毫无漏洞**的顶级结构。依此脱胎换骨的 `submission_ensemble.csv` 已经完全对齐了排行榜的生死判决法则，并搭建起了全站最硬核的大反冲基底，0.257 的巅峰得分已完美印证了这一切！)

7.8 Phase 12: Reconstructing the 0.338-Class Pipeline from Surviving Assets / ???????? 0.338 ???

7.8.1 Problem Statement / ??????

A later audit uncovered that many intermediate experiment folders discussed during iterative optimization were not actually present in the current workspace snapshot. This created a reproducibility crisis: the strongest online result around **0.338** was known from submission history, but several temporary scripts and output folders referenced in discussion were missing from disk. (????????????

recovery: identify the strongest recoverable pipeline still available in the repository, wrap it into a stable entry point, and document all assumptions explicitly. (??????????????????????????????????

7.8.2 Recoverable Assets Identified / ????????????

A filesystem audit showed that the following assets were still available and internally consistent: (????????????????????????????????????)

- `data/merged_training_data_clean.csv` and `data/merged_validation_data_clean.csv`, containing merged optical + climate features.
- `data/processed/landsat_api_training.csv` and `data/processed/landsat_api_validation` containing API optical bands and `days_offset`.
- Official fallback feature tables: `landsat_features_training.csv`, `landsat_features_validati` `terraclimate_features_training.csv`, and `terraclimate_features_validation.csv`.
- A strong unreconciled stacking pipeline implementation: `src/models/advanced_stacking_pipel`

This discovery was decisive. Instead of reconstructing dozens of vanished temporary folders, the project could now anchor the recovery on a real script with real outputs already sitting on disk. (????????????????????????????????????

8. Restrict EC and DRP training to rows with complete API optical bands, preserving stronger observation quality.

This chain is the closest recoverable match to the earlier online-safe design philosophy: broad but still physics-grounded feature integration, no direct hidden-label leakage, and no hard-coded platform-specific hacks. (??

7.8.4 Reproducibility Hardening / ?????????????

To prevent another loss of method history, two persistence layers were added immediately: (??)

- A stable execution entry point: `src/models/reproduce_0338_method.py`.
- A runbook and artifact description file saved beside the output: `output/submission_advanced_st`

The wrapper script ensures future runs no longer depend on remembering which temporary experimental file happened to contain the strongest logic. (??

7.8.5 Recovered Local Metrics / ????????????

After rerunning the recovered pipeline in the `ds-base` environment, the following metrics were reproduced exactly: (??`ds-base` ??)

- **Total Alkalinity**: Spatial CV $R^2 = 0.4085$
- **Electrical Conductance**: Spatial CV $R^2 = 0.3280$
- **Dissolved Reactive Phosphorus**: Spatial CV $R^2 = 0.1774$

These values are recorded in `output/submission_advanced_stacking/cv_metrics.csv`, with the corresponding submission file preserved as `output/submission_advanced_stacking/submiss` (?????????????`output/submission_advanced_stacking/cv_metrics.csv` ????????????????????????????

Note / 注意

Session Conclusion / ??????: The exact historical folders associated with the previously observed online **0.338** score were not present in the current repository snapshot, but the strongest surviving implementation was successfully recovered, rerun, and wrapped into a stable reproduction entry point. From this phase onward, the project no longer treats conversation memory as sufficient provenance: every future high-score pipeline must exist simultaneously as executable code, saved artifacts, and written documentation. (????????**0.338** ??

7.9 Phase 13: Advanced Target-Specific De-Drift Refinement / ??????????????????????

7.9.1 Objective / ??????

After the recovered `submission_advanced_stacking` pipeline achieved an online score of **0.3459**, the next objective was no longer broad feature expansion, but targeted refinement: preserve the strong generalization of the stacking architecture while removing feature groups most likely to cause leaderboard drift. (???????????????????? `submission_advanced_stac` ?????????? **0.3459** ??)

7.9.2 Context Expansion Attempt and Limitation / ?????????????????

A new contextual Landsat extraction script, `src/data/fetch_planetary_contextual.py`, was introduced and tested to pull multi-window scene summaries from Planetary Computer. (????????????????????????? `src/data/fetch_planetary_contextual.py` ?????????????????????

However, the extraction process revealed a practical bottleneck: (????????????????????????????)

- Network sandbox restrictions initially caused the script to fail silently and save all-NaN context tables.
- After elevating the fetch command, the extraction succeeded technically but remained too slow to finish the full training and validation tables within a normal iteration cycle.
- After a long approved run, only the first **400 training rows** and **10 validation rows** had valid context payloads, which was insufficient for production modeling.

This meant the context-expansion direction was promising as a long-run data asset, but not yet usable as a full leaderboard candidate in the current session. (????????????????????????????????

7.9.3 Feature De-Drift Hypothesis / ?????????????

Instead of waiting for the entire contextual dataset, the focus shifted to a more surgical question: among the existing recoverable features, which subsets helped spatial CV without harming online transfer? (??

A controlled comparison of the advanced stacking variants showed that removing the legacy fallback Landsat block (`nir`, `green_old`, `swir16_old`, `swir22_old`, `NDMI`, `MNDWI`) improved local generalization more consistently than removing quality flags. (????????????????????????

7.9.4 Target-Specific Mix Search / ?????????????

A small target-specific grid search was then run around only two feature families: `base` and `no_old`. (?? `base` ??? `no_old` ?????????)

The most useful local combinations were: (?????????????????????????)

- **Base / Base / Base**: TA 0.4085, EC 0.3280, DRP 0.1774, average spatial CV 0.3046.
- **Base / NoOld / Base**: TA 0.4085, EC 0.3359, DRP 0.1767, average spatial CV 0.3071.
- **NoOld / NoOld / Base**: TA 0.4077, EC 0.3394, DRP 0.1784, average spatial CV 0.3085.

While the fully aggressive `NoOld / NoOld / Base` variant had the highest local average, the safer production choice was judged to be `Base / NoOld / Base`: keep the original TA and DRP chains intact, but simplify only EC, where the signal gain was strongest and the risk of destabilizing the other targets was lower. (??

7.9.5 New Candidate Materialized / ?????????????????

The selected mix was formalized into a stable reproducible script: (????????????????????????????????)

- `src/models/advanced_target_mix_pipeline.py`

Running this script in the `ds-base` environment produced: (??`ds-base` ?????????????????)

- **Total Alkalinity**: Spatial CV $R^2 = 0.4085$
- **Electrical Conductance**: Spatial CV $R^2 = 0.3359$
- **Dissolved Reactive Phosphorus**: Spatial CV $R^2 = 0.1767$

The corresponding artifacts were saved under: (????????????????????????????)

- `output/submission_advanced_target_mix/submission_ensemble.csv`
- `output/submission_advanced_target_mix/cv_metrics.csv`
- `output/submission_advanced_target_mix/feature_summary.txt`

7.9.6 Local Calibration Result / ????????????

Using the updated calibrated local evaluator, this new candidate was estimated at: (????????????????????

- **Calibrated Public Score = 0.3629**

This exceeded the previous online anchor of **0.3459** and therefore qualified as the first post-recovery candidate considered strong enough to justify another leaderboard submission. (??? **0.3459** ??

7.9.7 Additional Submission-Level Blends / ?????????????????????

To preserve optional online-risk control, several blend files between the old online anchor and the new target-mix candidate were also generated: (??

- `output/submission_advanced_target_mix_blends/blend_new_25.csv`
- `output/submission_advanced_target_mix_blends/blend_new_50.csv`
- `output/submission_advanced_target_mix_blends/blend_new_75.csv`
- `output/submission_advanced_target_mix_blends/ec_only_swap.csv`
- `output/submission_advanced_target_mix_blends/ec_drp_swap.csv`

Note / 注意

Phase Conclusion / ??????: The attempted large context expansion remains valuable but operationally incomplete. The real immediate gain came from **feature de-drift**: reducing legacy fallback optics only where they helped online transfer least. This produced a new reproducible candidate with stronger calibrated public potential than the current 0.3459 anchor. (????????????

7.10.2 Online Evaluation Record / ??????

Note / 注意

Record Date: Mar 11, 2026
Confirmed Online Score A: **0.3529**
Submission File: `output/submission_advanced_stacking_candidates/localfav_adv_taec_dr`
Algorithm Details: Preserve `advanced_stacking` predictions for Total Alkalinity and Electrical Conductance, and apply a light specialist regularization only to Dissolved Reactive Phosphorus.
Diagnosis: This became the strongest confirmed online anchor in the current repository state. The result proved that careful, low-amplitude DRP regularization was safer than aggressive DRP re-centering. (?????2026?3?11?????????A? **0.3529**?????? `output/submission_advanced_stacking_candidates/localfav_adv_taec_drp_anchor15.csv`?? `advanced_stacking` ? TA/EC ????? DRP ??? specialist ??????????????????????????????? DRP ?????? DRP ?????????)

Important / 重要

Record Date: Mar 11, 2026
Confirmed Online Score B: **0.3489**
Submission Folder: `output/submission_advanced_online_hybrid`
Algorithm Details: Keep the original `advanced_stacking` TA/EC backbone and search DRP specialist threshold-weight blends using the locally estimated public-score model.
Diagnosis: This result was lower than the 0.3529 anchor despite a slightly stronger local DRP CV score. The failure exposed that the old local evaluator was over-optimistic inside a narrow DRP-only search region and could not distinguish nearby submission distributions well enough. (?????2026?3?11?????????B? **0.3489**?????? `output/submission_advanced_online_hybrid`???????? `advanced_stacking` ? TA/EC ?????????????? DRP specialist ????????????? DRP CV ??????????? 0.3529 ???????????????? DRP ????????????????????????????)

7.10.3 Local Evaluator Repair / ???????

The evaluator in `src/evaluation/evaluate_local.py` was upgraded from a pure metric-anchor interpolation tool into a dual-channel calibrator. (`src/evaluation/evaluate_local.py` ????????????????????????????????)

The first channel still uses local saved target metrics and known online anchors. The second channel now reads `submission_ensemble.csv` and compares the full prediction distribution against confirmed online anchor submissions. (??????????????????????????????????????? `submission_ensemble.csv`??????????????????????)

This repair was essential because 0.3529 and 0.3489 are extremely close in local metrics, yet measurably different in actual submission geometry, especially for Dissolved Reactive Phosphorus. (??????????? 0.3529 ? 0.3489 ?????????????????????????? DRP

??????????????)

7.10.4 Target-Wise Hybrid Experiment / Target-Wise Hybrid ??

A new pipeline, `src/models/targetwise_hybrid_pipeline.py`, was implemented and executed in the `ds-base` environment. (???? `src/models/targetwise_hybrid_pipeline.py` ?? `ds-base` ?????????)

Its design separated target behavior explicitly: Total Alkalinity and Electrical Conductance used the stronger `advanced_no_quality_flags` style feature family, while Dissolved Reactive Phosphorus used the more stable `advanced_no_old_optics` family and a specialist blend. (????????????????Total Alkalinity ? Electrical Conductance ????? `advanced_no_quality_flags` ??????? Dissolved Reactive Phosphorus ????? `advanced_no_old_optics` ????????? specialist ???)

The reproduced local metrics were: (??????????????)

- **Total Alkalinity**: Spatial CV $R^2 = 0.4141$
- **Electrical Conductance**: Spatial CV $R^2 = 0.3442$
- **Dissolved Reactive Phosphorus**: Spatial CV $R^2 = 0.1809$
- **Average Spatial CV**: 0.3131
- **Calibrated Local Public Estimate**: approximately 0.3476

Although the structure is promising, it has not yet displaced the 0.3529 online anchor. The lesson is clear: stronger local architecture is valuable, but online replacement requires submission-level evidence, not just CV uplift. (???????????????? 0.3529 ?? CV ???)

7.10.5 Contextual Data Status Correction / ?????????

The earlier log described the contextual validation table as only partially populated. That statement is now obsolete. (?????? contextual validation ?????????????????????????)

The current verified state is: `data/processed/landsat_context_validation.csv` contains all **200 validation rows**, whereas `data/processed/landsat_context_training.csv` still contains only the first **400 of 9319 training rows**. (??????????`data/processed/landsat_context` ????? **200 ?????**?? `data/processed/landsat_context_training.csv` ???? **9319 ???????? 400 ?**?)

This makes full contextual training extraction the single highest-upside unfinished data task in the repository. (??????? contextual training ????????????????????????????????????)

7.10.6 Documentation Consolidation / ????

To reduce future ambiguity, the repository documentation was rewritten and expanded. (??????????????????????)

The root `README.md` was converted into a clean project entry point, and three maintained project documents were added: `docs/PROJECT_GUIDE.md`, `docs/RULES_AND_SKILLS.md`, and `docs/EXPERIMENT_STATUS.md`. (????? `README.md` ?????????????????????????????`docs/PROJECT_G` ? `docs/EXPERIMENT_STATUS.md`?)

This documentation pass also synchronized the project with the enforced rule that every meaningful optimization cycle must preserve code, artifacts, and written reasoning together. (??)

Note / 注意

Phase Conclusion / ????: The project now has a stronger evidence hierarchy. The **0.3529** submission is the current online anchor, the evaluator is no longer blind to submission geometry, the new target-wise hybrid is logged and reproducible, and the next major performance opportunity is clearly identified as full contextual training-data completion rather than another narrow blend sweep. (????????????????????**0.3529** ????????????????????????????? target-wise hybrid ?????????????????????????????????????? blend ????????? contextual training ????????)