

MATH-UA 263 Partial Differential Equations

Recitation Summary

Yuanxun (Bill) Bao

Office Hour: Wednesday 2-4pm, WWH 1003

Email: yxb201@nyu.edu

1 February 2, 2018

Topics: verifying solution to a PDE, dispersion relations, well-posedness, general solution via integration.

1. (EPDE, Exercise 1.6) Verify the general solution of the heat equation on the real line:

$$\begin{cases} u_t = ku_{xx}, & -\infty < x < \infty, t > 0 \\ u(x, 0) = g(x) \end{cases}$$

is given by

$$u(x, t) = \frac{1}{\sqrt{4\pi kt}} \int_{-\infty}^{\infty} e^{-(x-s)^2/4kt} g(s) ds. \quad (1)$$

2. (PDE p.51-52) Solve the heat equation with the initial condition $u(x, 0) = g(x) = e^{-x}$. To do so, use (1) and the integral identity $\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-x^2/2} dx = 1$ to show that the solution is

$$u(x, t) = e^{kt-x}. \quad (2)$$

3. Find the dispersion relation of the linear PDE: $u_t = -u - \delta u_{xx} - u_{xxxx}$, $\delta > 0$.
4. (Well-posed and ill-posed problems) Consider the following PDEs for u and v :

$$(\square) \begin{cases} u_{tt} - u_{xx} = 0 \\ u(x, 0) = 0 \\ u_t(x, 0) = 0 \end{cases} \quad \text{and} \quad (\triangle) \begin{cases} v_{tt} - v_{xx} = 0 \\ v(x, 0) = 0 \\ v_t(x, 0) = \epsilon \sin\left(\frac{x}{\epsilon}\right) \end{cases} \quad (3)$$

It is straightforward to check that $u(x, t) = 0$ and $v(x, t) = \epsilon^2 \sin\left(\frac{x}{\epsilon}\right) \sin\left(\frac{t}{\epsilon}\right)$. For small $\epsilon > 0$, note that (\triangle) is a small perturbation to (\square) in the initial derivative data. Use the notion of “stability with respect to initial data” to argue that (\triangle) is well-posed. (*Hint:* if $\|v_t(x, 0) - u_t(x, 0)\| \leq \epsilon$, then $\|v(x, t) - u(x, t)\| \leq \epsilon^2$, for all $t > 0$.)

Similarly, consider

$$(\square) \begin{cases} u_{tt} + u_{xx} = 0 \\ u(x, 0) = 0 \\ u_t(x, 0) = 0 \end{cases} \quad \text{and} \quad (\triangle) \begin{cases} v_{tt} + v_{xx} = 0 \\ v(x, 0) = 0 \\ v_t(x, 0) = \epsilon \sin(\frac{x}{\epsilon}) \end{cases} \quad (4)$$

Note that $u(x, t) = 0$ and $v(x, t) = \epsilon^2 \sin(\frac{x}{\epsilon}) \sinh(\frac{t}{\epsilon})$. Argue that (\triangle) is an ill-posed problem.

5. Find the general solution of the PDE: $u_{xt} + 3u_x = 1$.

2 February 9, 2018

Topics: conservations laws, differential operators in polar/spherical coordinates, BVP and eigenvalues/eigenfunctions, method of characteristics.

1. (*1D gas dynamics*) Consider the one dimensional, time-dependent flow of gas under the assumption of constant pressure p . If u denotes the gas velocity, ρ the density and e the energy per unit volume, the basic equations of gas dynamics are

$$u_t + uu_x = 0 \quad (5)$$

$$\rho_t + u\rho_x + \rho u_x = 0 \quad (6)$$

$$e_t + ue_x + eu_x + pu_x = 0. \quad (7)$$

Rewrite the above equations in terms of a conservation of law. In other words, find the quantity (vector) \mathbf{U} and flux $\mathbf{F}(\mathbf{U})$ so that the above equations can be written in the form:

$$\mathbf{U}_t + [\mathbf{F}(\mathbf{U})]_x = \mathbf{0}. \quad (8)$$

2. Suppose that $u(x, y)$ satisfies the two-dimensional Laplace equation $\nabla^2 u = 0$ and u only depends on the distance from the origin $r = \sqrt{x^2 + y^2}$, ie, $u(x, y) \equiv v(r, t)$. Show that

$$v_{rr} + \frac{1}{r}v_r = 0. \quad (9)$$

3. For what value(s) of λ does the following linear system $(\mathbf{A} - \lambda\mathbf{I})\mathbf{x} = \mathbf{0}$ have a unique solution or infinitely many solutions, where

$$\mathbf{A} = \begin{bmatrix} 3 & 1 \\ 2 & 2 \end{bmatrix}.$$

Now consider the following two-point *Boundary Value Problem* (BVP) for $0 < x < 2\pi$:

$$u''(x) + \lambda u = 0, \quad (10)$$

$$u(0) = u'(2\pi) = 0. \quad (11)$$

For what value(s) of λ does the BVP has a unique solution? Infinitely many solutions?

4. (APDE, §1.2, Exercise 8, p.26) Solve the initial value problem

$$u_t + u_x - 3u = t, \quad x \in \mathbb{R}, t > 0, \quad (12)$$

$$u(x, 0) = x^2, \quad x \in \mathbb{R}. \quad (13)$$

5. (APDE, §1.2, Example 1.10, p.18) Find the general solution to $u_t + 2tu_x = 0$.

3 February 16, 2018

Topics: first-order PDEs and method of characteristics.

1. Revisit APDE §1.2, Exercise 8, solve by two approaches: (1) using a suitable choice of coordinate transformation, and, (2) using the method of characteristics.
2. Solve the following PDE by method of characteristics. Sketch some characteristic curves.

$$\begin{cases} y^2 u_x + u_y = 0, \\ u(x, 0) = x^2. \end{cases} \quad (14)$$

3. APDE, §1.2, Example 1.11.
4. Classify and solve the following PDE

$$\begin{cases} 2u_{xx} + 5u_{xt} + 3u_{tt} = 0, \\ u(x, 0) = 0, \quad u_t(x, 0) = xe^{-x^2}. \end{cases} \quad (15)$$

4 February 23, 2018

Topics: Wave equation, d'Alembert's formula, domain of dependence, Heat equation and Cauchy problem.

1. Review the solution of #5 in HW3.
2. (*The hammer blow*, PDE 2.1 Exercise #5,6) Consider the wave equation $u_{tt} = u_{xx}$ on the entire real line $-\infty < x < \infty$ with zero initial position $u(x, 0) = 0$ and initial velocity $u_t(x, 0) = g(x)$, where $g(x) = 1$ for $|x| < 1$ and $g(x) = 0$ for $|x| \geq 1$. Sketch the solution at time instants $t = \frac{1}{2}, 1, \frac{3}{2}, 2$ and $t = \frac{5}{2}$. What is the maximum displacement $\max_x u(x, t)$?
3. Solve the heat equation on the whole line with given initial data $u(x, 0) = \phi(x)$:
 - (a) $\phi(x) = 1$ for $|x| < l$, and $\phi(x) = 0$ for $|x| > l$.
 - (b) $\phi(x) = x^2$.
 - (c) $\phi(x) = e^{3x}$.

5 March 2, 2018

Topics: Duhamel's principle for the wave, heat, and linear advection equations.

1. Review the solution of #4 in HW3.
2. Principle of superposition for breaking down the solution of the inhomogeneous equation (with sources) with nonzero initial data into two subproblems: one homogeneous problem with nonzero initial data, and one inhomogeneous problem with zero initial data (Duhamel's principle).
3. Explain Duhamel's principle for the wave, heat, and linear advection equations.
4. Exercise 1, §3.4, PDE.
5. Part of #6 in HW4.
6. APDE, §2.5, Exercise #3 and #4.

6 March 9, 2018

Topics: Review midterm solutions.

7 March 16, 2018

Spring break.

8 March 23, 2018

Topics: Review #3, #4 on the midterm, linear PDEs in bounded domains, separation of variables.

1. Use separation of variables to find the solution to the heat equation with homogeneous Neumann BCs:

$$\begin{aligned}u_t &= ku_{xx}, \quad \text{for } 0 < x < L, t > 0, \\u(x, 0) &= \phi(x), \\u_x(0, t) &= u_x(L, t) = 0.\end{aligned}$$

2. Use separation of variables to find the solution to the wave equation with homogeneous Dirichlet BCs:

$$\begin{aligned}u_{tt} &= c^2 u_{xx}, \quad \text{for } 0 < x < L, t > 0, \\u(x, 0) &= f(x), \\u_t(x, 0) &= g(x), \\u(0, t) &= u(L, t) = 0.\end{aligned}$$

9 March 30, 2018

Topics: separation of variables and Fourier series.

1. Consider

$$X''(x) + \lambda X(x) = 0, \quad 0 < x < L. \quad (16)$$

For each of the following BCs, find its eigenvalues and eigenfunctions $\{\lambda_n, \varphi_n\}$ to the above 2-point BVP.

| BCs | λ_n | φ_n |
|--|-------------|-------------|
| Dirichlet: $X(0) = X(L) = 0$ | | |
| Neumann: $X'(0) = X'(L) = 0$ | | |
| Mixed: $X'(0) = X(L) = 0$ | | |
| Mixed: $X(0) = X'(L) = 0$ | | |
| Periodic: $X(0) = X(L), X'(0) = X'(L)$ | | |

2. Solve the following heat equation with initial condition and boundary conditions:

$$\begin{aligned} u_t &= k u_{xx}, \quad \text{for } 0 < x < L, t > 0, \\ u(x, 0) &= 1 + \frac{1}{2018} \cos\left(\frac{2018\pi x}{L}\right), \\ u_x(0, t) &= u_x(L, t) = 0. \end{aligned}$$

3. Solve the wave equation with Dirichlet BCs:

$$\begin{aligned} u_{tt} &= c^2 u_{xx}, \quad \text{for } 0 < x < L, t > 0, \\ u(x, 0) &= f(x), \quad u_t(x, 0) = g(x), \\ u(0, t) &= u(L, t) = 0. \end{aligned}$$

Use Fourier series to express the solution.

10 April 6, 2018

1. Use separation of variables to solve the wave equation with Neumann BCs and initial conditions:

$$\begin{aligned} u_{tt} &= c^2 u_{xx}, \quad \text{for } 0 < x < 1, t > 0, \\ u(x, 0) &= f(x), \quad u_t(x, 0) = 0, \\ u_x(0, t) &= u_x(1, t) = 0, \end{aligned}$$

where $f(x)$ is a “bump” function

$$f(x) = \begin{cases} \cos(4\pi x) & \frac{3}{8} < x < \frac{5}{8}, \\ 0, & \text{otherwise.} \end{cases} \quad (17)$$

Plot the solution by using MATLAB, Python, Maple or Mathematica, etc.

2. Pointwise, uniform convergence of Fourier series. Convergence in L^2 -norm, Parseval's identity, and completeness of an orthogonal set (notion of orthonormal basis).

11 April 13, 2018

Topics: inhomogeneous BCs and sources in bounded domains, Sturm-Liouville theory.