

# 1 Hamiltonians and EOM

## 1.1 Toy Problem

Consider simplest spin Hamiltonian  $H = -\vec{B} \cdot \vec{s}$ . It's clear that if we set up initial conditions  $\vec{s}$  misaligned from  $\vec{B}$ , it will simply spin around  $\vec{B}$ , which is fixed. Thus, let  $\hat{B} \cdot \hat{s} = \cos \theta$  the angle between the two, and let  $\phi$  measure the azimuthal angle.

We claim that  $\cos \theta, \phi$  are canonical variables. Since  $\phi$  is ignorable, immediately  $\frac{d\theta}{dt} = \frac{d \cos \theta}{dt} = -\frac{\partial H}{\partial \phi} = 0$ , while  $\frac{d\phi}{dt} = \frac{\partial H}{\partial (\cos \theta)} = Bs$  tells us the rate at which the spin precesses around  $\vec{B}$ .

## 1.2 Cassini State Hamiltonian

This Hamiltonian is Kassandra's Eq. 13, in the co-rotating frame with the perturber's angular momentum:

$$\mathcal{H} = \frac{1}{2}(\hat{s} \cdot \hat{l})^2 - \eta(\hat{s} \cdot \hat{l}_p). \quad (1)$$

In this frame, we can choose  $\hat{l} \equiv \hat{z}$  fixed, and  $\hat{l}_p = \cos I \hat{z} + \sin I \hat{x}$  fixed as well. Then

$$\hat{s} = \cos \theta \hat{z} - \sin \theta (\sin \phi \hat{y} + \cos \phi \hat{x}).$$

We can choose the convention for  $\phi = \phi$  azimuthal angle requiring  $\phi = 0, \pi$  mean coplanarity between  $\hat{s}, \hat{l}, \hat{l}_p$  in the  $\hat{x}, \hat{z}$  plane such that  $\hat{l}_p, \hat{s}$  lie on the same side of  $\hat{l}$ . Then we can evaluate in coordinates

$$\begin{aligned} \hat{s} \cdot \hat{l} &= \cos \theta, \\ \hat{s} \cdot \hat{l}_p &= \cos \theta \cos I - \sin I \sin \theta \cos \phi, \\ \mathcal{H} &= \frac{1}{2} \cos^2 \theta - \eta(\cos \theta \cos I - \sin I \sin \theta \cos \phi). \end{aligned}$$

Note that if we take  $\cos \theta$  to be our canonical variable,  $\sin \theta = \sqrt{1 - \cos^2 \theta}$  can be used.

## 1.3 Equation of Motion

The correct EOM comes from Kassandra's Eq. 12:

$$\begin{aligned} \frac{d\hat{s}}{dt} &= (\hat{s} \cdot \hat{l})(\hat{s} \times \hat{l}) - \eta(\hat{s} \times \hat{l}_p), \\ &= (s_y s_z - \eta s_y \cos I) \hat{x} - (s_x s_z + \eta(s_x \cos I - s_z \sin I)) \hat{y} + \eta s_y \sin I \hat{z}. \end{aligned}$$

Alternatively, consider Hamilton's equations applied to the Hamiltonian:

$$\frac{\partial \phi}{\partial t} = \frac{\partial \mathcal{H}}{\partial (\cos \theta)} = \cos \theta - \eta(\cos I + \sin I \cot \theta \cos \phi), \quad (2)$$

$$\frac{\partial (\cos \theta)}{\partial t} = -\frac{\partial \mathcal{H}}{\partial \phi} = +\eta \sin I \sin \theta \sin \phi. \quad (3)$$

This produces the same trajectories as the Cartesian EOM, so this is correct. However, since  $\frac{\partial \phi}{\partial t} \propto 1/\sin \theta$ , this is not a desirable system of equations to use, as they are very stiff near  $\theta \approx 0$ .

## 1.4 Tidal Dissipation

We can add a tidal dissipation term; we write it in form  $\left(\frac{d\hat{s}}{dt}\right)_{tide} = \epsilon \hat{s} \times (\hat{l} \times \hat{s})$ . Expanding,

$$\begin{aligned} \left(\frac{d\hat{s}}{dt}\right)_{tide} &= \epsilon(\hat{z} - s_z \hat{s}), \\ &= \epsilon(-s_z s_x \hat{x} - s_z s_y \hat{y} + (1 - s_z^2) \hat{z}). \end{aligned} \quad (4)$$

We run numerical simulations for weaker  $\epsilon \ll \eta \ll 1$  and stronger  $\epsilon \lesssim \eta \ll 1$ .

We can seek equilibria of the the system including tides, which requires

$$\begin{aligned} 0 &= s_y s_z - \eta s_y \cos I - \epsilon s_z s_x, \\ 0 &= -s_x s_z - \eta(s_x \cos I - s_z \sin I) - \epsilon s_z s_y, \\ 0 &= \eta s_y \sin(I) + \epsilon(1 - s_z^2). \end{aligned}$$

We expect at least two equilibria, based on the simulations: one near  $s_z \approx 1$  and one  $s_z \approx 0$ .

For near alignment/near Cassini state 1,  $1 - s_z \sim 1 - s_\perp^2$ , so we can set  $s_z = 1$  to first order:  $s_y - \epsilon s_x - \eta s_y \cos I = -s_x - \eta(s_x \cos I - \sin I) - \epsilon s_y = \eta s_y \sin I = 0$ . This can be satisfied if we set  $s_x = \tan(I) \ll 1, s_y = \mathcal{O}(\epsilon s_x)$ ; this coarsely corresponds to Cassini state 1.

The other solution should be near Cassini state 2, where  $s_x \approx 1$ ; dropping second order terms forces  $\eta s_y + \epsilon s_z = -s_z - \eta(\cos I - s_z \sin I) = \eta s_y \sin(I) + \epsilon = 0$ . This can thus be satisfied for  $s_y \approx -\frac{\epsilon}{\eta \sin(I)}$ . Thus, this explains why as  $\epsilon$  is increased, we first start to get points that don't converge to Cassini state 2 in the absence of tides, before starting to see points that fail to converge to Cassini state 1.