

Perturbation-Stable Maximum Cut

Yuchong Pan

UBC Beyond Worst-Case Analysis Seminars

June 30, 2020

MAXIMUM CUT

Problem (MAXIMUM CUT)

Input: An undirected graph $G = (V, E)$ with edge weights $w_e > 0$ for each $e \in E$.

Goal: A cut (A, B) that maximizes the weight of the *crossing* edges.

MAXIMUM CUT

Problem (MAXIMUM CUT)

Input: An undirected graph $G = (V, E)$ with edge weights $w_e > 0$ for each $e \in E$.

Goal: A cut (A, B) that maximizes the weight of the *crossing* edges.

- ▶ MAXIMUM CUT is a type of 2-clustering problem (e.g. weights measure dissimilarities).

MAXIMUM CUT Is *NP*-Hard

Problem (MAXIMUM CUT, Decision Version)

Input: An undirected graph $G = (V, E)$ with edge weights $w_e > 0$ for each $e \in E$, and a positive integer W .

Output: Yes iff. there is a set $S \subseteq V$ such that the weight of the *crossing* edges is at least W .

MAXIMUM CUT Is NP-Hard

Problem (MAXIMUM CUT, Decision Version)

Input: An undirected graph $G = (V, E)$ with edge weights $w_e > 0$ for each $e \in E$, and a positive integer W .

Output: Yes iff. there is a set $S \subseteq V$ such that the weight of the *crossing* edges is at least W .

Problem (PARTITION, Decision Version)

Input: $(c_1, \dots, c_n) \in \mathbb{Z}^n$.

Output: Yes iff. there is $I \subseteq [n]$ such that $\sum_{i \in I} c_i = \sum_{i \notin I} c_i$.

MAXIMUM CUT Is *NP*-Hard

Problem (MAXIMUM CUT, Decision Version)

Input: An undirected graph $G = (V, E)$ with edge weights $w_e > 0$ for each $e \in E$, and a positive integer W .

Output: Yes iff. there is a set $S \subseteq V$ such that the weight of the *crossing* edges is at least W .

Problem (PARTITION, Decision Version)

Input: $(c_1, \dots, c_n) \in \mathbb{Z}^n$.

Output: Yes iff. there is $I \subseteq [n]$ such that $\sum_{i \in I} c_i = \sum_{i \notin I} c_i$.

Proof Sketch ($\text{PARTITION} \leq_P \text{MAXIMUM CUT}$)

- ▶ $G = K_n$.
- ▶ $w_{ij} = c_i c_j$ for all $i, j \in V, i \neq j$.
- ▶ $W = \lceil \frac{1}{4} \sum c_i^2 \rceil$.

MAXIMUM CUT Is *NP*-Hard

- ▶ MINIMUM CUT is *not NP*-hard and can be solved by the Maximum-Flow Minimum-Cut Theorem.

MAXIMUM CUT Is *NP*-Hard

- ▶ MINIMUM CUT is *not NP*-hard and can be solved by the Maximum-Flow Minimum-Cut Theorem.
- ▶ **Question:** Can't we negate the edge weights, yielding a MINIMUM CUT instance?

MAXIMUM CUT Is *NP*-Hard

- ▶ MINIMUM CUT is *not NP*-hard and can be solved by the Maximum-Flow Minimum-Cut Theorem.
- ▶ **Question:** Can't we negate the edge weights, yielding a MINIMUM CUT instance?
- ▶ No! Polynomial-time algorithms solving MINIMUM CUT require nonnegative edge weights.

Beyond Worst-Case: Exact Recovery

- **Theme:** To recover the optimal solution in polynomial time in γ -*perturbation-stable* instances, where γ is as small as possible.

Beyond Worst-Case: Exact Recovery

- **Theme:** To recover the optimal solution in polynomial time in γ -*perturbation-stable* instances, where γ is as small as possible.

Definition (γ -Perturbation-Stability)

For $\gamma \geq 1$, an instance of MAXIMUM CUT is γ -*perturbation-stable* if a cut (A, B) is the *unique* optimal solution to all γ -*perturbations*, where each original edge weight w_e is replaced with an edge weight $w'_e \in [\frac{1}{\gamma} w_e, w_e]$.

LP Relaxation, Take 1

- **Question:** Can we use an LP relaxation similar to the one for MINIMUM CUT, i.e.

$$\begin{array}{ll} \max & \sum_{e \in E} w_e x_e \\ \text{s.t.} & x_e \geq |d_u - d_v|, \quad \forall e = uv \in E. \\ & x_e \in [0, 1], \quad \forall e \in E. \\ & d_v \in [0, 1], \quad \forall v \in V. \end{array}$$

LP Relaxation, Take 1

- **Question:** Can we use an LP relaxation similar to the one for MINIMUM CUT, i.e.

$$\begin{array}{ll}\max & \sum_{e \in E} w_e x_e \\ \text{s.t.} & x_e \geq |d_u - d_v|, \quad \forall e = uv \in E. \\ & x_e \in [0, 1], \quad \forall e \in E. \\ & d_v \in [0, 1], \quad \forall v \in V.\end{array}$$

- No! $x_e = 1$ for each $e \in E$ is a feasible solution and maximizes the objective value.

LP Relaxation, Take 1

- **Question:** Can we use an LP relaxation similar to the one for MINIMUM CUT, i.e.

$$\begin{array}{ll}\max & \sum_{e \in E} w_e x_e \\ \text{s.t.} & x_e \geq |d_u - d_v|, \quad \forall e = uv \in E. \\ & x_e \in [0, 1], \quad \forall e \in E. \\ & d_v \in [0, 1], \quad \forall v \in V.\end{array}$$

- No! $x_e = 1$ for each $e \in E$ is a feasible solution and maximizes the objective value.
- **Question:** What about $x_e \leq d_u - d_v$ and $x_e \leq d_v - d_u$?

LP Relaxation, Take 1

- **Question:** Can we use an LP relaxation similar to the one for MINIMUM CUT, i.e.

$$\begin{array}{ll}\max & \sum_{e \in E} w_e x_e \\ \text{s.t.} & x_e \geq |d_u - d_v|, \quad \forall e = uv \in E. \\ & x_e \in [0, 1], \quad \forall e \in E. \\ & d_v \in [0, 1], \quad \forall v \in V.\end{array}$$

- No! $x_e = 1$ for each $e \in E$ is a feasible solution and maximizes the objective value.
- **Question:** What about $x_e \leq d_u - d_v$ and $x_e \leq d_v - d_u$?
- This forces $x_e = 0$, instead of $x_e \leq |d_u - d_v|$.

LP Relaxation, Take 2

- ▶ Let $x_{ij} \in \{0, 1\}$ denote whether or not i, j are on different sides of the cut, for all distinct $i, j \in V$. We denote by x_{ij} and x_{ji} the same variable.

LP Relaxation, Take 2

- ▶ Let $x_{ij} \in \{0, 1\}$ denote whether or not i, j are on different sides of the cut, for all distinct $i, j \in V$. We denote by x_{ij} and x_{ji} the same variable.
- ▶ **Intuition:** If i, j are on different sides, and i, k are also on different sides, then j, k must be on the same sides.

LP Relaxation, Take 2

- ▶ Let $x_{ij} \in \{0, 1\}$ denote whether or not i, j are on different sides of the cut, for all distinct $i, j \in V$. We denote by x_{ij} and x_{ji} the same variable.
- ▶ **Intuition:** If i, j are on different sides, and i, k are also on different sides, then j, k must be on the same sides.
- ▶ For any distinct $i, j, k \in V$, at most two of x_{ij}, x_{ik}, x_{jk} are 1.

LP Relaxation, Take 2

- ▶ Let $x_{ij} \in \{0, 1\}$ denote whether or not i, j are on different sides of the cut, for all distinct $i, j \in V$. We denote by x_{ij} and x_{ji} the same variable.
- ▶ **Intuition:** If i, j are on different sides, and i, k are also on different sides, then j, k must be on the same sides.
- ▶ For any distinct $i, j, k \in V$, at most two of x_{ij}, x_{ik}, x_{jk} are 1.

$$x_{ij} + x_{ik} + x_{jk} \leq 2, \quad \forall i, j, k \in V \text{ distinct.}$$

LP Relaxation, Take 2

- ▶ Let $x_{ij} \in \{0, 1\}$ denote whether or not i, j are on different sides of the cut, for all distinct $i, j \in V$. We denote by x_{ij} and x_{ji} the same variable.
- ▶ **Intuition:** If i, j are on different sides, and i, k are also on different sides, then j, k must be on the same sides.
- ▶ For any distinct $i, j, k \in V$, at most two of x_{ij}, x_{ik}, x_{jk} are 1.

$$x_{ij} + x_{ik} + x_{jk} \leq 2, \quad \forall i, j, k \in V \text{ distinct.}$$

- ▶ **Intuition:** If i, j are on the same side, and i, k are on the same side, then j, k are on the same side.

LP Relaxation, Take 2

- ▶ Let $x_{ij} \in \{0, 1\}$ denote whether or not i, j are on different sides of the cut, for all distinct $i, j \in V$. We denote by x_{ij} and x_{ji} the same variable.
- ▶ **Intuition:** If i, j are on different sides, and i, k are also on different sides, then j, k must be on the same sides.
- ▶ For any distinct $i, j, k \in V$, at most two of x_{ij}, x_{ik}, x_{jk} are 1.

$$x_{ij} + x_{ik} + x_{jk} \leq 2, \quad \forall i, j, k \in V \text{ distinct.}$$

- ▶ **Intuition:** If i, j are on the same side, and i, k are on the same side, then j, k are on the same side.
- ▶ For any distinct $i, j, k \in V$, $x_{ij} = x_{ik} = 0$ implies $x_{jk} = 0$.

LP Relaxation, Take 2

- ▶ Let $x_{ij} \in \{0, 1\}$ denote whether or not i, j are on different sides of the cut, for all distinct $i, j \in V$. We denote by x_{ij} and x_{ji} the same variable.
- ▶ **Intuition:** If i, j are on different sides, and i, k are also on different sides, then j, k must be on the same sides.
- ▶ For any distinct $i, j, k \in V$, at most two of x_{ij}, x_{ik}, x_{jk} are 1.

$$x_{ij} + x_{ik} + x_{jk} \leq 2, \quad \forall i, j, k \in V \text{ distinct.}$$

- ▶ **Intuition:** If i, j are on the same side, and i, k are on the same side, then j, k are on the same side.
- ▶ For any distinct $i, j, k \in V$, $x_{ij} = x_{ik} = 0$ implies $x_{jk} = 0$.

$$x_{jk} \leq x_{ij} + x_{ik}, \quad \forall i, j, k \in V \text{ distinct.}$$

LP Relaxation, Take 2

- Hence we obtain the LP relaxation (LP-MAXCUT):

$$\begin{array}{ll}\max & \sum_{(i,j) \in E} w_{ij} x_{ij} \\ \text{s.t.} & x_e \geq |d_u - d_v|, \quad \forall e = uv \in E. \\ & x_{ij} + x_{ik} + x_{jk} \leq 2, \quad \forall i, j, k \in V \text{ distinct.} \\ & x_{jk} \leq x_{ij} + x_{ik}, \quad \forall i, j, k \in V \text{ distinct.} \\ & x_{ij} \in [0, 1], \quad \forall i, j \in V \text{ distinct.}\end{array}$$

Main Theorem

Theorem

There is a constant $c > 0$ such that in every $(c \log n)$ -perturbation-stable instance of MAXIMUM CUT with n vertices, (LP-MAXCUT) solves to integers.