

# The Tales of Planar Graphs

The Extremal Problems, the Criterion, and the Conjectures

Alan Bu<sup>1</sup>   **Yuchong Pan**<sup>2</sup>

<sup>1</sup>Harvard University   <sup>2</sup>MIT

February 19, 2025  
Meet-and-Think  
CMU Tepper School of Business

**Disclaimer:** I am **not** a combinatorialist.

an extremal problem

an extremal problem (Kenyon 1996)

How large is the maximum number  $T_m$  of spanning trees over  $m$ -edge planar graphs?

## an extremal problem (Kenyon 1996)

How large is the **maximum number  $T_m$  of spanning trees** over  $m$ -edge planar graphs?

some StackExchange answer (user JimT):

This is a very, very tough question. It is obvious that for any graph  $G$  with  $n$  edges the number of spanning trees  $t(G)$  does not exceed  $2^n$  (each edge is either included in or excluded from a subtree; this is also the upper bound of the number of connected subgraphs of  $G$ ). We can somewhat improve this - if  $G$  has  $m$  vertices ( $m \leq n + 1$ ) then we have

$$t(G) \leq \binom{n}{m-1} < 2^n,$$

since we need to choose  $m - 1$  edges out of  $n$  (not arbitrarily, of course).

## an extremal problem (Kenyon 1996)

How large is the **maximum number  $T_m$  of spanning trees** over  $m$ -edge planar graphs?

some StackExchange answer (user JimT):

This is a very, very tough question. It is obvious that for any graph  $G$  with  $n$  edges the number of spanning trees  $t(G)$  does not exceed  $2^n$  (each edge is either included in or excluded from a subtree; this is also the upper bound of the number of connected subgraphs of  $G$ ). We can somewhat improve this - if  $G$  has  $m$  vertices ( $m \leq n + 1$ ) then we have

$$t(G) \leq \binom{n}{m-1} < 2^n,$$

since we need to choose  $m - 1$  edges out of  $n$  (not arbitrarily, of course).

a **lower bound**:  $T_m \geq 1.7916^m$  achieved by **square grid graphs**

some **upper bounds** ( $T_m \leq \tau^m$ ):

---

2 As a matter of fact, the conjecture above has a relatively elementary proof (found by yours truly) for  $\tau \approx 1.96$ . Of course, that is not very close to the hypothetical upper bound of  $e^{2Cl\pi}$ , but at least it is something... – [JimT](#) Jan 8, 2021 at 4:21 ✍

---

2 1.9328... is the new  $\tau$  - with an elementary proof. :) – [JimT](#) Jan 15, 2021 at 3:32

---

There was a recent improvement of  $1.96^n$  down to  $1.913^n$ . I could not find the proof, but the link is here: [sspcdn.blob.core.windows.net/files/Documents/SEP/STS/2024/...](https://sspcdn.blob.core.windows.net/files/Documents/SEP/STS/2024/...) – [Igor Pak](#) Aug 2, 2024 at 1:18

---

Actually, it is now at  $1.8637^n$ . See [arxiv.org/abs/2103.10523](https://arxiv.org/abs/2103.10523) – [JimT](#) Aug 3, 2024 at 3:34

---

Quick update: Here is a result by Stoimenow in knot theory, which, in their abstract, mentions giving an upper bound of  $1.8393^n$  on this number. Here: [projecteuclid.org/journals/tokyo-journal-of-mathematics/...](https://projecteuclid.org/journals/tokyo-journal-of-mathematics/...) – [Alien](#) Sep 28, 2024 at 0:17

---

@Alien Nice! One comment though - I quickly looked thru the article; it is very non-elementary, as it uses a bunch of other results in links and knot theory, as well as related polynomial algebra. But still, it proves the even lower upper bound, so the gap between this and  $1.7916...$  is getting smaller with every post here :) . – [JimT](#) Sep 29, 2024 at 2:32 ✍

---

some **upper bounds** ( $T_m \leq \tau^m$ ):

---

2 As a matter of fact, the conjecture above has a relatively elementary proof (found by yours truly) for  $\tau \approx 1.96$ . Of course, that is not very close to the hypothetical upper bound of  $e^{2Cl\pi}$ , but at least it is something... – [JimT](#) Jan 8, 2021 at 4:21 ✍

---

2 1.9328... is the new  $\tau$  - with an elementary proof. :) – [JimT](#) Jan 15, 2021 at 3:32

---

There was a recent improvement of  $1.96^n$  down to  $1.913^n$ . I could not find the proof, but the link is here: [spcndn.blob.core.windows.net/files/Documents/SEP/STS/2024/...](https://spcndn.blob.core.windows.net/files/Documents/SEP/STS/2024/...) – [Igor Pak](#) Aug 2, 2024 at 1:18

---

Actually, it is now at  $1.8637^n$ . See [arxiv.org/abs/2103.10523](https://arxiv.org/abs/2103.10523) – [JimT](#) Aug 3, 2024 at 3:34

---

Quick update: Here is a result by Stoimenow in knot theory, which, in their abstract, mentions giving an upper bound of  $1.8393^n$  on this number. Here: [projecteuclid.org/journals/tokyo-journal-of-mathematics/...](https://projecteuclid.org/journals/tokyo-journal-of-mathematics/...) – [Alien](#) Sep 28, 2024 at 0:17

---

@Alien Nice! One comment though - I quickly looked thru the article; it is very non-elementary, as it uses a bunch of other results in links and knot theory, as well as related polynomial algebra. But still, it proves the even lower upper bound, so the gap between this and  $1.7916...$  is getting smaller with every post here :) .  
– [JimT](#) Sep 29, 2024 at 2:32 ✍

---

**theorem (Stoimenow 2007)**

$T_m \leq \delta^m$ , where  $\delta \simeq 1.8393$  is the unique real root of  $x^3 - x^2 - x - 1 = 0$ .



In this talk, we provide a **simple, linear-algebraic** proof of Stoimenow's upper bound (in contrast to his “non-elementary”, knot-theoretic, polynomial-algebraic proof)

In this talk, we provide a **simple, linear-algebraic** proof of Stoimenow's upper bound (in contrast to his “non-elementary”, knot-theoretic, polynomial-algebraic proof)

... and also some nice by-products on **planarity criteria** from our techniques.

another extremal problem

### another extremal problem

How large is the **maximum determinant**  $\Delta_m$  over  $m \times m$  matrices that are concatenations  $[M|N]$  of two submatrices of (possibly different) incidence matrices?

### another extremal problem

How large is the **maximum determinant**  $\Delta_m$  over  $m \times m$  matrices that are concatenations  $[M|N]$  of two submatrices of (possibly different) incidence matrices?

### theorem (Bu, P. 2025)

$T_m \leq \Delta_m \leq \delta^m$ , where  $\delta \simeq 1.8393$  is the unique real root of  $x^3 - x^2 - x - 1 = 0$ .

$$\Delta_m \leq \delta^m$$

## definitions

An **incidence submatrix** is a matrix where each row has at most one 1, at most one  $-1$ , and all other entries zero.

## definitions

An **incidence submatrix** is a matrix where each row has at most one 1, at most one  $-1$ , and all other entries zero.

A **bi-incidence matrix** is a concatenation  $[M|N]$  of two incidence submatrices  $M$  and  $N$ .



## definitions

An **incidence submatrix** is a matrix where each row has at most one 1, at most one  $-1$ , and all other entries zero.

A **bi-incidence matrix** is a concatenation  $[M|N]$  of two incidence submatrices  $M$  and  $N$ .

**Recall:**  $\Delta_m$  is the maximum determinant over  $m \times m$  bi-incidence matrices.

## definitions

An **incidence submatrix** is a matrix where each row has at most one 1, at most one  $-1$ , and all other entries zero.

A **bi-incidence matrix** is a concatenation  $[M|N]$  of two incidence submatrices  $M$  and  $N$ .

**Recall:**  $\Delta_m$  is the maximum determinant over  $m \times m$  bi-incidence matrices.

proof of  $\Delta_m \leq \delta^m$

Let  $[M|N]$  achieve  $\Delta_m$ .

## definitions

An **incidence submatrix** is a matrix where each row has at most one 1, at most one  $-1$ , and all other entries zero.

A **bi-incidence matrix** is a concatenation  $[M|N]$  of two incidence submatrices  $M$  and  $N$ .

**Recall:**  $\Delta_m$  is the maximum determinant over  $m \times m$  bi-incidence matrices.

## proof of $\Delta_m \leq \delta^m$

Let  $[M|N]$  achieve  $\Delta_m$ .

Extend  $[M|N]$  by two columns to  $[M'|N']$  so that each row of  $M'$  (resp. each row of  $N'$ ) has **exactly** one 1 and **exactly** one  $-1$ .

## definitions

An **incidence submatrix** is a matrix where each row has at most one 1, at most one  $-1$ , and all other entries zero.

A **bi-incidence matrix** is a concatenation  $[M|N]$  of two incidence submatrices  $M$  and  $N$ .

**Recall:**  $\Delta_m$  is the maximum determinant over  $m \times m$  bi-incidence matrices.

## proof of $\Delta_m \leq \delta^m$

Let  $[M|N]$  achieve  $\Delta_m$ .

Extend  $[M|N]$  by two columns to  $[M'|N']$  so that each row of  $M'$  (resp. each row of  $N'$ ) has **exactly** one 1 and **exactly** one  $-1$ .

# nonzero entries in  $[M'|N']$ :  $4m$

## definitions

An **incidence submatrix** is a matrix where each row has at most one 1, at most one  $-1$ , and all other entries zero.

A **bi-incidence matrix** is a concatenation  $[M|N]$  of two incidence submatrices  $M$  and  $N$ .

**Recall:**  $\Delta_m$  is the maximum determinant over  $m \times m$  bi-incidence matrices.

## proof of $\Delta_m \leq \delta^m$

Let  $[M|N]$  achieve  $\Delta_m$ .

Extend  $[M|N]$  by two columns to  $[M'|N']$  so that each row of  $M'$  (resp. each row of  $N'$ ) has **exactly** one 1 and **exactly** one  $-1$ .

# nonzero entries in  $[M'|N']$ :  $4m$

# columns in  $[M'|N']$ :  $m + 2$

## definitions

An **incidence submatrix** is a matrix where each row has at most one 1, at most one  $-1$ , and all other entries zero.

A **bi-incidence matrix** is a concatenation  $[M|N]$  of two incidence submatrices  $M$  and  $N$ .

**Recall:**  $\Delta_m$  is the maximum determinant over  $m \times m$  bi-incidence matrices.

## proof of $\Delta_m \leq \delta^m$

Let  $[M|N]$  achieve  $\Delta_m$ .

Extend  $[M|N]$  by two columns to  $[M'|N']$  so that each row of  $M'$  (resp. each row of  $N'$ ) has **exactly** one 1 and **exactly** one  $-1$ .

# nonzero entries in  $[M'|N']$ :  $4m$

# columns in  $[M'|N']$ :  $m+2$

$\implies [M'|N']$  has a column with at most  $\lfloor 4m/(m+2) \rfloor = 3$  nonzero entries.

## definitions

An **incidence submatrix** is a matrix where each row has at most one 1, at most one  $-1$ , and all other entries zero.

A **bi-incidence matrix** is a concatenation  $[M|N]$  of two incidence submatrices  $M$  and  $N$ .

**Recall:**  $\Delta_m$  is the maximum determinant over  $m \times m$  bi-incidence matrices.

## proof of $\Delta_m \leq \delta^m$

Let  $[M|N]$  achieve  $\Delta_m$ .

Extend  $[M|N]$  by two columns to  $[M'|N']$  so that each row of  $M'$  (resp. each row of  $N'$ ) has **exactly** one 1 and **exactly** one  $-1$ .

# nonzero entries in  $[M'|N']$ :  $4m$

# columns in  $[M'|N']$ :  $m+2$

$\implies [M'|N']$  has a column with at most  $\lfloor 4m/(m+2) \rfloor = 3$  nonzero entries.

W.l.o.g. (by column operations), this column is in  $[M|N]$ .

proof of  $\Delta_m \leq \delta^m$  (cont'd)

$$\begin{matrix} r_1 \\ r_2 \\ r_3 \\ r_4 \\ \vdots \\ r_m \end{matrix} \left[ \begin{array}{c|c} \begin{matrix} a_{11} \\ a_{21} \\ a_{31} \end{matrix} & \begin{matrix} \\ \\ \end{matrix} \\ \hline \begin{matrix} 0 \\ \vdots \\ 0 \end{matrix} & \begin{matrix} \\ \\ \end{matrix} \end{array} \right]$$



proof of  $\Delta_m \leq \delta^m$  (cont'd)

$$\begin{array}{c} r_1 \\ r_2 \\ r_3 \\ r_4 \\ \vdots \\ r_m \end{array} \left[ \begin{array}{c|c} \begin{array}{c} a_{11} \\ a_{21} \\ a_{31} \end{array} & \begin{array}{c} \\ \\ \end{array} \\ \hline \begin{array}{c} 0 \\ \vdots \\ 0 \end{array} & \begin{array}{c} \\ \\ \end{array} \end{array} \right]$$

$$\left[ \begin{array}{c|c} \text{green} & \text{orange} \end{array} \right] = \left[ \begin{array}{c|c} \text{green} & 0 \dots 0 \end{array} \right] + \left[ \begin{array}{c|c} 0 \dots 0 & \text{orange} \end{array} \right]$$

proof of  $\Delta_m \leq \delta^m$  (cont'd)

$$\begin{array}{c|c} \begin{matrix} r_1 \\ r_2 \\ r_3 \\ r_4 \\ \vdots \\ r_m \end{matrix} & \begin{bmatrix} a_{11} \\ a_{21} \\ a_{31} \\ 0 \\ \vdots \\ 0 \end{bmatrix} \end{array}$$

$$\left[ \begin{array}{c|c} \text{green} & \text{orange} \end{array} \right] = \left[ \begin{array}{c|c} \text{green} & 0 \dots 0 \end{array} \right] + \left[ \begin{array}{c|c} 0 \dots 0 & \text{orange} \end{array} \right]$$

$$\det [M \mid N] = \sum_{\alpha, \beta, \gamma \in \{0,1\}} \det \begin{bmatrix} r_1^\alpha & & \\ r_2^\beta & & \\ r_3^\gamma & & \\ r_4 & 0 & \\ \vdots & \vdots & \\ r_m & 0 & \end{bmatrix}$$

proof of  $\Delta_m \leq \delta^m$  (cont'd)

$$\begin{matrix} r_1^1 \\ r_2^1 \\ r_3^1 \\ r_4 \\ \vdots \\ r_m \end{matrix} \left[ \begin{array}{cccc|c} 0 & \cdots & 0 & & \\ 0 & \cdots & 0 & & \\ 0 & \cdots & 0 & & \\ 0 & & & & \\ \vdots & & & & \\ 0 & & & & \end{array} \right]$$

**case 1:**  $\alpha = \beta = \gamma = 1$

proof of  $\Delta_m \leq \delta^m$  (cont'd)

$$\det \begin{matrix} r_1^1 \\ r_2^1 \\ r_3^1 \\ r_4 \\ \vdots \\ r_m \end{matrix} \left[ \begin{array}{cccc|c} 0 & \cdots & 0 & & \\ 0 & \cdots & 0 & & \\ 0 & \cdots & 0 & & \\ 0 & & & & \\ \vdots & & & & \\ 0 & & & & \end{array} \right] = 0.$$

**case 1:**  $\alpha = \beta = \gamma = 1$

proof of  $\Delta_m \leq \delta^m$  (cont'd)

$$\sum_{\beta, \gamma \in \{0,1\}} \det \begin{bmatrix} r_1^0 & 0 \dots 0 \\ r_2^\beta & \\ r_3^\gamma & \\ r_4 & \\ \vdots & \\ r_m & \end{bmatrix} = \det \begin{bmatrix} r_1^0 & 0 \dots 0 \\ r_2 & \\ r_3 & \\ r_4 & \\ \vdots & \\ r_m & \end{bmatrix}$$

case 2:  $\alpha = 0$

proof of  $\Delta_m \leq \delta^m$  (cont'd)

$$\sum_{\beta, \gamma \in \{0,1\}} \det \begin{bmatrix} r_1^0 & 0 \dots 0 \\ r_2^\beta & \\ r_3^\gamma & \\ r_4 & \\ \vdots & \\ r_m & \end{bmatrix} = \det \begin{bmatrix} r_1^0 & 0 \dots 0 \\ r_2 & \\ r_3 & \\ r_4 & \\ \vdots & \\ r_m & \end{bmatrix}$$

case 2:  $\alpha = 0$

removing-zeros lemma (Bu, P. 2025; informal)

If an  $m \times m$  bi-incidence matrix has a row whose restriction to the left/right side has zeros only,

proof of  $\Delta_m \leq \delta^m$  (cont'd)

$$\sum_{\beta, \gamma \in \{0,1\}} \det \begin{array}{c} r_1^0 \\ r_2^\beta \\ r_3^\gamma \\ r_4 \\ \vdots \\ r_m \end{array} \left[ \begin{array}{c|c} \text{green} & 0 \dots 0 \\ \text{purple} & \text{purple} \\ \hline & \end{array} \right] = \det \begin{array}{c} r_1^0 \\ r_2 \\ r_3 \\ r_4 \\ \vdots \\ r_m \end{array} \left[ \begin{array}{c|c} \text{green} & 0 \dots 0 \\ & \text{orange} \\ \hline & \end{array} \right]$$

case 2:  $\alpha = 0$

removing-zeros lemma (Bu, P. 2025; informal)

If an  $m \times m$  bi-incidence matrix has a row whose restriction to the left/right side has zeros only, then one can remove this row and perform column operations to obtain an  $(m-1) \times (m-1)$  square bi-incidence matrix, while preserving the determinant.

proof of  $\Delta_m \leq \delta^m$  (cont'd)

$$\sum_{\beta, \gamma \in \{0,1\}} \det \begin{bmatrix} r_1^0 & \text{green} & 0 \dots 0 \\ r_2^\beta & \text{purple} & \text{purple} \\ r_3^\gamma & & \\ r_4 & & \\ \vdots & & \\ r_m & & \end{bmatrix} = \det \begin{bmatrix} r_1^0 & \text{green} & 0 \dots 0 \\ r_2 & \text{green} & \text{orange} \\ r_3 & & \\ r_4 & & \\ \vdots & & \\ r_m & & \end{bmatrix} \leq \Delta_{m-1}.$$

case 2:  $\alpha = 0$

removing-zeros lemma (Bu, P. 2025; informal)

If an  $m \times m$  bi-incidence matrix has a row whose restriction to the left/right side has zeros only, then one can remove this row and perform column operations to obtain an  $(m-1) \times (m-1)$  square bi-incidence matrix, while preserving the determinant.



proof of  $\Delta_m \leq \delta^m$  (cont'd)

$$\sum_{\gamma \in \{0,1\}} \det \begin{array}{c} r_1^1 \\ r_1^0 \\ r_2^0 \\ r_3^\gamma \\ r_4 \\ \vdots \\ r_m \end{array} \left[ \begin{array}{c|c} 0 \dots 0 & \text{orange} \\ \hline \text{green} & 0 \dots 0 \\ \hline \text{purple} & \text{purple} \\ \hline & \\ \hline & \\ \hline & \end{array} \right] = \det \begin{array}{c} r_1^1 \\ r_1^0 \\ r_2^0 \\ r_3 \\ r_4 \\ \vdots \\ r_m \end{array} \left[ \begin{array}{c|c} 0 \dots 0 & \text{orange} \\ \hline \text{green} & 0 \dots 0 \\ \hline & \text{orange} \\ \hline & \\ \hline & \\ \hline & \end{array} \right] \leq \Delta_{m-2}$$

case 3:  $\alpha = 1$  and  $\beta = 0$

proof of  $\Delta_m \leq \delta^m$  (cont'd)

$$\sum_{\gamma \in \{0,1\}} \det \begin{bmatrix} r_1^1 & 0 & \dots & 0 & \text{orange} \\ r_2^0 & \text{green} & & 0 & \dots & 0 \\ r_3^\gamma & \text{purple} & & \text{purple} \\ r_4 & & & \\ \vdots & & & \\ r_m & & & \end{bmatrix} = \det \begin{bmatrix} r_1^1 & 0 & \dots & 0 & \text{orange} \\ r_2^0 & \text{green} & & 0 & \dots & 0 \\ r_3 & \text{green} & & \text{orange} \\ r_4 & & & \\ \vdots & & & \\ r_m & & & \end{bmatrix} \leq \Delta_{m-2}$$

**case 3:**  $\alpha = 1$  and  $\beta = 0$

$$\det \begin{bmatrix} r_1^1 & 0 & \dots & 0 & \text{orange} \\ r_2^1 & 0 & \dots & 0 & \text{orange} \\ r_3^0 & \text{green} & & 0 & \dots & 0 \\ r_4 & & & \\ \vdots & & & \\ r_m & & & \end{bmatrix} \leq \Delta_{m-3}$$

**case 4:**  $\alpha = \beta = 1$  and  $\gamma = 0$

$$\Delta_m \leq \Delta_{m-1} + \Delta_{m-2} + \Delta_{m-3}$$

$$T_m \leq \Delta_m$$

proof of  $T_m \leq \Delta_m$

**Recall:**  $\Delta_m$  is the maximum determinant over  $m \times m$  bi-incidence matrices;  
 $T_m$  is the maximum number of spanning trees over  $m$ -edge planar graphs.

proof of  $T_m \leq \Delta_m$

**Recall:**  $\Delta_m$  is the maximum determinant over  $m \times m$  bi-incidence matrices;  
 $T_m$  is the maximum number of spanning trees over  $m$ -edge planar graphs.

**Goal:** Given an  $m$ -edge planar graph  $G$  (achieving  $T_m$ ), construct an  $m \times m$  bi-incidence matrix with determinant at least the number of spanning trees in  $G$ .

## proof of $T_m \leq \Delta_m$

**Recall:**  $\Delta_m$  is the maximum determinant over  $m \times m$  bi-incidence matrices;  
 $T_m$  is the maximum number of spanning trees over  $m$ -edge planar graphs.

**Goal:** Given an  $m$ -edge planar graph  $G$  (achieving  $T_m$ ), construct an  $m \times m$  bi-incidence matrix with determinant at least the number of spanning trees in  $G$ .

### definitions

A **truncated incidence matrix** of a digraph is the incidence matrix of the digraph with an arbitrary column removed.

## proof of $T_m \leq \Delta_m$

**Recall:**  $\Delta_m$  is the maximum determinant over  $m \times m$  bi-incidence matrices;  
 $T_m$  is the maximum number of spanning trees over  $m$ -edge planar graphs.

**Goal:** Given an  $m$ -edge planar graph  $G$  (achieving  $T_m$ ), construct an  $m \times m$  bi-incidence matrix with determinant at least the number of spanning trees in  $G$ .

### definitions

A **truncated incidence matrix** of a digraph is the incidence matrix of the digraph with an arbitrary column removed.

### key lemma (Bu, P. 2025)

Let  $G$  be a connected planar graph with orientation  $D$ . Let  $D^*$  be the dual of  $D$ .

Let  $M$  and  $M^*$  be truncated incidence matrices of  $D$  and  $D^*$ , of which the  $i^{\text{th}}$  rows correspond to the same arc and its dual arc for each  $i$ .


Then  $|\det[M|M^*]|$  is equal to the number of spanning trees  $\tau(G)$  in  $G$ .



proof of  $T_m \leq \Delta_m$  (cont'd)

key lemma (Bu, P. 2025)


Let  $G$  be a connected planar graph with orientation  $D$ .



proof of  $T_m \leq \Delta_m$  (cont'd)

key lemma (Bu, P. 2025)

Let  $G$  be a connected planar graph with orientation  $D$ . Let  $D^*$  be the dual of  $D$ .




## proof of $T_m \leq \Delta_m$ (cont'd)

### key lemma (Bu, P. 2025)

Let  $G$  be a connected planar graph with orientation  $D$ . Let  $D^*$  be the dual of  $D$ .

Let  $M$  and  $M^*$  be truncated incidence matrices of  $D$  and  $D^*$ , of which the  $i^{\text{th}}$  rows correspond to the same arc and its dual arc for each  $i$ .



$v_1$	$v_2$	$v_3$	$f_1$	$f_2$
-1			-1	
1	-1		-1	
	1		-1	1
		-1		1
	-1	1		1


## proof of $T_m \leq \Delta_m$ (cont'd)

### key lemma (Bu, P. 2025)

Let  $G$  be a connected planar graph with orientation  $D$ . Let  $D^*$  be the dual of  $D$ .

Let  $M$  and  $M^*$  be truncated incidence matrices of  $D$  and  $D^*$ , of which the  $i^{\text{th}}$  rows correspond to the same arc and its dual arc for each  $i$ .

Then  $|\det[M|M^*]|$  is equal to the number of spanning trees  $\tau(G)$  in  $G$ .



$v_1$	$v_2$	$v_3$	$f_1$	$f_2$
-1			-1	
1	-1		-1	
	1		-1	1
		-1		1
	-1	1		1


## proof of $T_m \leq \Delta_m$ (cont'd)

### key lemma (Bu, P. 2025)

Let  $G$  be a connected planar graph with orientation  $D$ . Let  $D^*$  be the dual of  $D$ .

Let  $M$  and  $M^*$  be truncated incidence matrices of  $D$  and  $D^*$ , of which the  $i^{\text{th}}$  rows correspond to the same arc and its dual arc for each  $i$ .

Then  $|\det[M|M^*]|$  is equal to the number of spanning trees  $\tau(G)$  in  $G$ .



$v_1$	$v_2$	$v_3$	$f_1$	$f_2$
-1			-1	
1	-1		-1	
	1		-1	1
		-1		1
	-1	1		1

### corollary

For all  $m \in \mathbb{N}$ , we have  $T_m \leq \Delta_m$ .

## proof of key lemma

### key lemma (Bu, P. 2025)

Let  $G$  be a connected planar graph with orientation  $D$ . Let  $D^*$  be the dual of  $D$ .

Let  $M$  and  $M^*$  be truncated incidence matrices of  $D$  and  $D^*$ , of which the  $i^{\text{th}}$  rows correspond to the same arc and its dual arc for each  $i$ .

Then  $|\det[M|M^*]|$  is equal to the number of spanning trees  $\tau(G)$  in  $G$ .

## proof of key lemma

### key lemma (Bu, P. 2025)

Let  $G$  be a connected planar graph with orientation  $D$ . Let  $D^*$  be the dual of  $D$ .

Let  $M$  and  $M^*$  be truncated incidence matrices of  $D$  and  $D^*$ , of which the  $i^{\text{th}}$  rows correspond to the same arc and its dual arc for each  $i$ .

Then  $|\det[M|M^*]|$  is equal to the number of spanning trees  $\tau(G)$  in  $G$ .

$$(\det [M \mid M^*])^2 = \det [M \mid M^*]^T [M \mid M^*]$$

## proof of key lemma

### key lemma (Bu, P. 2025)

Let  $G$  be a connected planar graph with orientation  $D$ . Let  $D^*$  be the dual of  $D$ .

Let  $M$  and  $M^*$  be truncated incidence matrices of  $D$  and  $D^*$ , of which the  $i^{\text{th}}$  rows correspond to the same arc and its dual arc for each  $i$ .

Then  $|\det[M|M^*]|$  is equal to the number of spanning trees  $\tau(G)$  in  $G$ .

$$\begin{aligned} (\det [M \mid M^*])^2 &= \det [M \mid M^*]^T [M \mid M^*] \\ &= \det \begin{bmatrix} M^T M & M^T M^* \\ (M^*)^T M & (M^*)^T M^* \end{bmatrix} \end{aligned}$$



## proof of key lemma

### key lemma (Bu, P. 2025)

Let  $G$  be a connected planar graph with orientation  $D$ . Let  $D^*$  be the dual of  $D$ .

Let  $M$  and  $M^*$  be truncated incidence matrices of  $D$  and  $D^*$ , of which the  $i^{\text{th}}$  rows correspond to the same arc and its dual arc for each  $i$ .

Then  $|\det[M|M^*]|$  is equal to the number of spanning trees  $\tau(G)$  in  $G$ .

$$\begin{aligned} (\det [M \mid M^*])^2 &= \det [M \mid M^*]^T [M \mid M^*] \\ &= \det \begin{bmatrix} M^T M & M^T M^* \\ (M^*)^T M & (M^*)^T M^* \end{bmatrix} \\ &= \det \begin{bmatrix} M^T M & 0 \\ 0 & (M^*)^T M^* \end{bmatrix} \quad (\text{if } M^T M^* = 0) \end{aligned}$$

## proof of key lemma

### key lemma (Bu, P. 2025)

Let  $G$  be a connected planar graph with orientation  $D$ . Let  $D^*$  be the dual of  $D$ .

Let  $M$  and  $M^*$  be truncated incidence matrices of  $D$  and  $D^*$ , of which the  $i^{\text{th}}$  rows correspond to the same arc and its dual arc for each  $i$ .

Then  $|\det[M|M^*]|$  is equal to the number of spanning trees  $\tau(G)$  in  $G$ .

$$\begin{aligned}(\det [M \mid M^*])^2 &= \det [M \mid M^*]^T [M \mid M^*] \\&= \det \begin{bmatrix} M^T M & M^T M^* \\ (M^*)^T M & (M^*)^T M^* \end{bmatrix} \\&= \det \begin{bmatrix} M^T M & 0 \\ 0 & (M^*)^T M^* \end{bmatrix} \quad (\text{if } M^T M^* = 0) \\&= \det M^T M \cdot \det (M^*)^T M^*\end{aligned}$$

## proof of key lemma

### key lemma (Bu, P. 2025)

Let  $G$  be a connected planar graph with orientation  $D$ . Let  $D^*$  be the dual of  $D$ .

Let  $M$  and  $M^*$  be truncated incidence matrices of  $D$  and  $D^*$ , of which the  $i^{\text{th}}$  rows correspond to the same arc and its dual arc for each  $i$ .

Then  $|\det[M|M^*]|$  is equal to the number of spanning trees  $\tau(G)$  in  $G$ .

$$\begin{aligned}(\det [M \mid M^*])^2 &= \det [M \mid M^*]^T [M \mid M^*] \\&= \det \begin{bmatrix} M^T M & M^T M^* \\ (M^*)^T M & (M^*)^T M^* \end{bmatrix} \\&= \det \begin{bmatrix} M^T M & 0 \\ 0 & (M^*)^T M^* \end{bmatrix} \quad (\text{if } M^T M^* = 0) \\&= \det M^T M \cdot \det (M^*)^T M^* \\&= \tau(G) \cdot \tau(G^*)\end{aligned}$$

## proof of key lemma

### key lemma (Bu, P. 2025)

Let  $G$  be a connected planar graph with orientation  $D$ . Let  $D^*$  be the dual of  $D$ .

Let  $M$  and  $M^*$  be truncated incidence matrices of  $D$  and  $D^*$ , of which the  $i^{\text{th}}$  rows correspond to the same arc and its dual arc for each  $i$ .

Then  $|\det[M|M^*]|$  is equal to the number of spanning trees  $\tau(G)$  in  $G$ .

$$\begin{aligned}(\det [M \mid M^*])^2 &= \det [M \mid M^*]^T [M \mid M^*] \\&= \det \begin{bmatrix} M^T M & M^T M^* \\ (M^*)^T M & (M^*)^T M^* \end{bmatrix} \\&= \det \begin{bmatrix} M^T M & 0 \\ 0 & (M^*)^T M^* \end{bmatrix} \quad (\text{if } M^T M^* = 0) \\&= \det M^T M \cdot \det (M^*)^T M^* \\&= \tau(G) \cdot \tau(G^*) \\&= \tau(G)^2.\end{aligned}$$

proof of  $M^T M^* = 0$

proof of  $M^T M^* = 0$


W.l.o.g.,  $G$  is loopless.

proof of  $M^T M^* = 0$

W.l.o.g.,  $G$  is **loopless**. (Why?)

proof of  $M^T M^* = 0$



W.l.o.g.,  $G$  is **loopless**. (Why?)







proof of  $M^T M^* = 0$

W.l.o.g.,  $G$  is **loopless**. (Why?)





proof of  $M^T M^* = 0$

W.l.o.g.,  $G$  is **loopless**. (Why?)



proof of  $M^T M^* = 0$

W.l.o.g.,  $G$  is **loopless**. (Why?)



Hence, column  $v$  in  $M$  and column  $f$  in  $M^*$  are **orthogonal**.

a planarity criterion

### yet another extremal problem

Given a connected graph  $G$  with  $m$  edges, how large is the **maximum determinant**  $\Delta(G)$  over  $m \times m$  matrices  $[M|N]$ , where  $M$  is a truncated incidence matrix of  $G$  and  $N$  is an incidence submatrix of appropriate size?

### yet another extremal problem

Given a connected graph  $G$  with  $m$  edges, how large is the **maximum determinant**  $\Delta(G)$  over  $m \times m$  matrices  $[M|N]$ , where  $M$  is a truncated incidence matrix of  $G$  and  $N$  is an incidence submatrix of appropriate size?

In other words, can we achieve better determinants than the previous construction, where  $N$  is a truncated incidence matrix of the dual?

### yet another extremal problem

Given a connected graph  $G$  with  $m$  edges, how large is the **maximum determinant**  $\Delta(G)$  over  $m \times m$  matrices  $[M|N]$ , where  $M$  is a truncated incidence matrix of  $G$  and  $N$  is an incidence submatrix of appropriate size?

In other words, can we achieve better determinants than the previous construction, where  $N$  is a truncated incidence matrix of the dual?

**proposition** (Bu, P. 2025)

For any connected graph  $G$ , we have  $\Delta(G) \leq \tau(G)$ .

### yet another extremal problem

Given a connected graph  $G$  with  $m$  edges, how large is the **maximum determinant**  $\Delta(G)$  over  $m \times m$  matrices  $[M|N]$ , where  $M$  is a truncated incidence matrix of  $G$  and  $N$  is an incidence submatrix of appropriate size?

In other words, can we achieve better determinants than the previous construction, where  $N$  is a truncated incidence matrix of the dual?

### proposition (Bu, P. 2025)

For any connected graph  $G$ , we have  $\Delta(G) \leq \tau(G)$ .

In other words, the previous construction is the **best possible**.



## definitions

Given a **connected** graph  $G$ , define its **excess** to be  $\epsilon(G) := \tau(G) - \Delta(G)$ .

## definitions

Given a **connected** graph  $G$ , define its **excess** to be  $\varepsilon(G) := \tau(G) - \Delta(G)$ .

Given a **disconnected** graph  $G$ , define its **excess**  $\varepsilon(G)$  to be the sum of the excesses of its connected components.

## definitions

Given a **connected** graph  $G$ , define its **excess** to be  $\varepsilon(G) := \tau(G) - \Delta(G)$ .

Given a **disconnected** graph  $G$ , define its **excess**  $\varepsilon(G)$  to be the sum of the excesses of its connected components.

For **planar** graphs, we have seen that the previous construction is the **best possible**.

## corollary of key lemma

For any **planar** graph  $G$ , we have  $\varepsilon(G) = 0$ .

## definitions

Given a **connected** graph  $G$ , define its **excess** to be  $\varepsilon(G) := \tau(G) - \Delta(G)$ .

Given a **disconnected** graph  $G$ , define its **excess**  $\varepsilon(G)$  to be the sum of the excesses of its connected components.

For **planar** graphs, we have seen that the previous construction is the **best possible**.

## corollary of key lemma

For any **planar** graph  $G$ , we have  $\varepsilon(G) = 0$ .

For **non-planar** graphs, the excesses are at least 18.

## lemma (Bu, P. 2025)

For any **non-planar** graph  $G$ , we have  $\varepsilon(G) \geq 18$ .

## definitions

Given a **connected** graph  $G$ , define its **excess** to be  $\varepsilon(G) := \tau(G) - \Delta(G)$ .

Given a **disconnected** graph  $G$ , define its **excess**  $\varepsilon(G)$  to be the sum of the excesses of its connected components.

For **planar** graphs, we have seen that the previous construction is the **best possible**.

## corollary of key lemma

For any **planar** graph  $G$ , we have  $\varepsilon(G) = 0$ .

For **non-planar** graphs, the excesses are at least 18.

## lemma (Bu, P. 2025)

For any **non-planar** graph  $G$ , we have  $\varepsilon(G) \geq 18$ .


The lower bound **18** is tight and achieved by  $K_{3,3}$ .

### theorem (Wagner 1937)

A graph is non-planar if and only if it can produce either  $K_5$  or  $K_{3,3}$  by a sequence of **edge contractions** and **edge deletions**.


### theorem (Wagner 1937)

A graph is non-planar if and only if it can produce either  $K_5$  or  $K_{3,3}$  by a sequence of **edge contractions** and **edge deletions**.



### theorem (Wagner 1937)


A graph is non-planar if and only if it can produce either  $K_5$  or  $K_{3,3}$  by a sequence of **edge contractions** and **edge deletions**.







### theorem (Wagner 1937)

A graph is non-planar if and only if it can produce either  $K_5$  or  $K_{3,3}$  by a sequence of **edge contractions** and **edge deletions**.



### theorem (Wagner 1937)

A graph is non-planar if and only if it can produce either  $K_5$  or  $K_{3,3}$  by a sequence of **edge contractions** and **edge deletions**.



### merge-cut lemma (Bu, P. 2025)

For connected  $G = (V, E)$  and non-loop non-bridge  $e \in E$ ,  $\varepsilon(G) \geq \varepsilon(G/e) + \varepsilon(G \setminus e)$ .

### merge-cut lemma (Bu, P. 2025)

For connected  $G = (V, E)$  and non-loop non-bridge  $e \in E$ ,  $\varepsilon(G) \geq \varepsilon(G/e) + \varepsilon(G \setminus e)$ .

proof of  $\varepsilon(G) \geq 18$  for non-planar  $G$

W.l.o.g., assume that  $G$  is connected.

## merge-cut lemma (Bu, P. 2025)

For connected  $G = (V, E)$  and non-loop non-bridge  $e \in E$ ,  $\varepsilon(G) \geq \varepsilon(G/e) + \varepsilon(G \setminus e)$ .

## proof of $\varepsilon(G) \geq 18$ for non-planar $G$

W.l.o.g., assume that  $G$  is connected.

merge-cut lemma & non-negative of  $\varepsilon(\cdot) \implies$  for any **non-loop non-bridge** edge  $e$ ,

$$\varepsilon(G) \geq \varepsilon(G/e) + \varepsilon(G \setminus e) \geq \max\{\varepsilon(G/e), \varepsilon(G \setminus e)\}.$$

## merge-cut lemma (Bu, P. 2025)

For connected  $G = (V, E)$  and non-loop non-bridge  $e \in E$ ,  $\varepsilon(G) \geq \varepsilon(G/e) + \varepsilon(G \setminus e)$ .

## proof of $\varepsilon(G) \geq 18$ for non-planar $G$

W.l.o.g., assume that  $G$  is connected.

merge-cut lemma & non-negative of  $\varepsilon(\cdot) \implies$  for any **non-loop non-bridge** edge  $e$ ,

$$\varepsilon(G) \geq \varepsilon(G/e) + \varepsilon(G \setminus e) \geq \max\{\varepsilon(G/e), \varepsilon(G \setminus e)\}.$$

removing-zeros lemma  $\implies$  for any **loop** edge  $e$ ,  $\varepsilon(G) \geq \varepsilon(G \setminus e)$ .

## merge-cut lemma (Bu, P. 2025)

For connected  $G = (V, E)$  and non-loop non-bridge  $e \in E$ ,  $\varepsilon(G) \geq \varepsilon(G/e) + \varepsilon(G \setminus e)$ .

## proof of $\varepsilon(G) \geq 18$ for non-planar $G$

W.l.o.g., assume that  $G$  is connected.

merge-cut lemma & non-negative of  $\varepsilon(\cdot) \implies$  for any **non-loop non-bridge** edge  $e$ ,

$$\varepsilon(G) \geq \varepsilon(G/e) + \varepsilon(G \setminus e) \geq \max\{\varepsilon(G/e), \varepsilon(G \setminus e)\}.$$

removing-zeros lemma  $\implies$  for any **loop** edge  $e$ ,  $\varepsilon(G) \geq \varepsilon(G \setminus e)$ .

Wagner's theorem  $\implies$  a sequence of **edge contractions** and **edge deletions**

## merge-cut lemma (Bu, P. 2025)

For connected  $G = (V, E)$  and non-loop non-bridge  $e \in E$ ,  $\varepsilon(G) \geq \varepsilon(G/e) + \varepsilon(G \setminus e)$ .

## proof of $\varepsilon(G) \geq 18$ for non-planar $G$

W.l.o.g., assume that  $G$  is connected.

merge-cut lemma & non-negative of  $\varepsilon(\cdot) \implies$  for any **non-loop non-bridge** edge  $e$ ,

$$\varepsilon(G) \geq \varepsilon(G/e) + \varepsilon(G \setminus e) \geq \max\{\varepsilon(G/e), \varepsilon(G \setminus e)\}.$$

removing-zeros lemma  $\implies$  for any **loop** edge  $e$ ,  $\varepsilon(G) \geq \varepsilon(G \setminus e)$ .

Wagner's theorem  $\implies$  a sequence of **edge contractions** and **edge deletions** such that each intermediate graph is **connected**



## merge-cut lemma (Bu, P. 2025)

For connected  $G = (V, E)$  and non-loop non-bridge  $e \in E$ ,  $\varepsilon(G) \geq \varepsilon(G/e) + \varepsilon(G \setminus e)$ .

## proof of $\varepsilon(G) \geq 18$ for non-planar $G$

W.l.o.g., assume that  $G$  is connected.

merge-cut lemma & non-negative of  $\varepsilon(\cdot)$   $\implies$  for any **non-loop non-bridge** edge  $e$ ,

$$\varepsilon(G) \geq \varepsilon(G/e) + \varepsilon(G \setminus e) \geq \max\{\varepsilon(G/e), \varepsilon(G \setminus e)\}.$$

removing-zeros lemma  $\implies$  for any **loop** edge  $e$ ,  $\varepsilon(G) \geq \varepsilon(G \setminus e)$ .

Wagner's theorem  $\implies$  a sequence of **edge contractions** and **edge deletions** such that each intermediate graph is **connected** (i.e., if we delete an edge  $e$ , then  $e$  is **non-bridge**).

### merge-cut lemma (Bu, P. 2025)

For connected  $G = (V, E)$  and non-loop non-bridge  $e \in E$ ,  $\varepsilon(G) \geq \varepsilon(G/e) + \varepsilon(G \setminus e)$ .

### proof of $\varepsilon(G) \geq 18$ for non-planar $G$

W.l.o.g., assume that  $G$  is connected.

merge-cut lemma & non-negative of  $\varepsilon(\cdot) \implies$  for any **non-loop non-bridge** edge  $e$ ,

$$\varepsilon(G) \geq \varepsilon(G/e) + \varepsilon(G \setminus e) \geq \max\{\varepsilon(G/e), \varepsilon(G \setminus e)\}.$$

removing-zeros lemma  $\implies$  for any **loop** edge  $e$ ,  $\varepsilon(G) \geq \varepsilon(G \setminus e)$ .

Wagner's theorem  $\implies$  a sequence of **edge contractions** and **edge deletions** such that each intermediate graph is **connected** (i.e., if we delete an edge  $e$ , then  $e$  is **non-bridge**).

Hence,  $\varepsilon(G) \geq \min\{\varepsilon(K_{3,3}), \varepsilon(K_5)\}$ .

proposition (Bu, **P.** 2025)

We have  $\varepsilon(K_{3,3}) = 18$  and  $\varepsilon(K_5) = 25$ .

proposition (Bu, P. 2025)

We have  $\varepsilon(K_{3,3}) = 18$  and  $\varepsilon(K_5) = 25$ .

These two quantities can be computed in a computer-assisted manner with brute-force and some clever pruning.

proposition (Bu, P. 2025)

We have  $\varepsilon(K_{3,3}) = 18$  and  $\varepsilon(K_5) = 25$ .

These two quantities can be computed in a **computer-assisted** manner with **brute-force** and some clever **pruning**.

In the paper, we provide a purely **combinatorial** proof of these two quantities. It is quite involved and has many cases, but might provide intuition on the combinatorial structure. In particular, the tight case for  $K_5$  is quite elegant.

proposition (Bu, P. 2025)

We have  $\varepsilon(K_{3,3}) = 18$  and  $\varepsilon(K_5) = 25$ .

These two quantities can be computed in a **computer-assisted** manner with **brute-force** and some clever **pruning**.

In the paper, we provide a purely **combinatorial** proof of these two quantities. It is quite involved and has many cases, but might provide intuition on the combinatorial structure. In particular, the tight case for  $K_5$  is quite elegant.

**Open question:** Are there simpler proofs?

## definitions

Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two graphs with  $|E_1| = |E_2| = |V_1| + |V_2| - 2$ .

## definitions

Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two graphs with  $|E_1| = |E_2| = |V_1| + |V_2| - 2$ .

Let  $f : E_1 \rightarrow E_2$  be a bijection.



## definitions

Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two graphs with  $|E_1| = |E_2| = |V_1| + |V_2| - 2$ .

Let  $f : E_1 \rightarrow E_2$  be a bijection.

A spanning tree  $T$  of  $G_1$  is **annihilated** by  $f$  if  $f(E_1 \setminus T)$  is not a spanning tree of  $G_2$ .

## definitions

Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two graphs with  $|E_1| = |E_2| = |V_1| + |V_2| - 2$ .

Let  $f : E_1 \rightarrow E_2$  be a bijection.

A spanning tree  $T$  of  $G_1$  is **annihilated** by  $f$  if  $f(E_1 \setminus T)$  is not a spanning tree of  $G_2$ .

A spanning tree of  $G_1$  not annihilated by  $f$  is a **cospanning tree** of  $G_1$  and  $G_2$  under  $f$ .

## definitions

Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two graphs with  $|E_1| = |E_2| = |V_1| + |V_2| - 2$ .

Let  $f : E_1 \rightarrow E_2$  be a bijection.

A spanning tree  $T$  of  $G_1$  is **annihilated** by  $f$  if  $f(E_1 \setminus T)$  is not a spanning tree of  $G_2$ .

A spanning tree of  $G_1$  not annihilated by  $f$  is a **cospanning tree** of  $G_1$  and  $G_2$  under  $f$ .

Let  **$\text{cospans}(G_1, G_2)$**  be the maximum number of cospanning trees of  $G_1$  and  $G_2$  over  $f$ .

## definitions

Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two graphs with  $|E_1| = |E_2| = |V_1| + |V_2| - 2$ .

Let  $f : E_1 \rightarrow E_2$  be a bijection.

A spanning tree  $T$  of  $G_1$  is **annihilated** by  $f$  if  $f(E_1 \setminus T)$  is not a spanning tree of  $G_2$ .

A spanning tree of  $G_1$  not annihilated by  $f$  is a **cospanning tree** of  $G_1$  and  $G_2$  under  $f$ .

Let  **$\text{cspan}(G_1, G_2)$**  be the maximum number of cospanning trees of  $G_1$  and  $G_2$  over  $f$ .

## proposition (Bu, P. 2025)

Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two graphs with  $|E_1| = |E_2| = |V_1| + |V_2| - 2$ .

## definitions

Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two graphs with  $|E_1| = |E_2| = |V_1| + |V_2| - 2$ .

Let  $f : E_1 \rightarrow E_2$  be a bijection.

A spanning tree  $T$  of  $G_1$  is **annihilated** by  $f$  if  $f(E_1 \setminus T)$  is not a spanning tree of  $G_2$ .

A spanning tree of  $G_1$  not annihilated by  $f$  is a **cospanning tree** of  $G_1$  and  $G_2$  under  $f$ .

Let  **$\text{cspan}(G_1, G_2)$**  be the maximum number of cospanning trees of  $G_1$  and  $G_2$  over  $f$ .

## proposition (Bu, P. 2025)

Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two graphs with  $|E_1| = |E_2| = |V_1| + |V_2| - 2$ .

Let  $M$  and  $N$  be truncated incidence matrices of  $G_1$  and  $G_2$ , respectively.

## definitions

Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two graphs with  $|E_1| = |E_2| = |V_1| + |V_2| - 2$ .

Let  $f : E_1 \rightarrow E_2$  be a bijection.

A spanning tree  $T$  of  $G_1$  is **annihilated** by  $f$  if  $f(E_1 \setminus T)$  is not a spanning tree of  $G_2$ .

A spanning tree of  $G_1$  not annihilated by  $f$  is a **cospanning tree** of  $G_1$  and  $G_2$  under  $f$ .

Let  **$\text{cospan}(G_1, G_2)$**  be the maximum number of cospanning trees of  $G_1$  and  $G_2$  over  $f$ .

## proposition (Bu, P. 2025)

Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two graphs with  $|E_1| = |E_2| = |V_1| + |V_2| - 2$ .

Let  $M$  and  $N$  be truncated incidence matrices of  $G_1$  and  $G_2$ , respectively.

Then  **$\det[M|N] \leq \text{cospan}(G_1, G_2)$** .

## definitions

Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two graphs with  $|E_1| = |E_2| = |V_1| + |V_2| - 2$ .

Let  $f : E_1 \rightarrow E_2$  be a bijection.

A spanning tree  $T$  of  $G_1$  is **annihilated** by  $f$  if  $f(E_1 \setminus T)$  is not a spanning tree of  $G_2$ .

A spanning tree of  $G_1$  not annihilated by  $f$  is a **cospanning tree** of  $G_1$  and  $G_2$  under  $f$ .

Let  **$\text{cospan}(G_1, G_2)$**  be the maximum number of cospanning trees of  $G_1$  and  $G_2$  over  $f$ .

## proposition (Bu, P. 2025)

Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two graphs with  $|E_1| = |E_2| = |V_1| + |V_2| - 2$ .

Let  $M$  and  $N$  be truncated incidence matrices of  $G_1$  and  $G_2$ , respectively.

Then  **$\det[M|N] \leq \text{cospan}(G_1, G_2)$** .

In the proof for  $\varepsilon(K_{3,3})$  and  $\varepsilon(K_5)$ , we repeatedly **restrict the space of feasible graphs** corresponding to the right incidence submatrix through various bounds.

## definitions

Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two graphs with  $|E_1| = |E_2| = |V_1| + |V_2| - 2$ .

Let  $f : E_1 \rightarrow E_2$  be a bijection.

A spanning tree  $T$  of  $G_1$  is **annihilated** by  $f$  if  $f(E_1 \setminus T)$  is not a spanning tree of  $G_2$ .

A spanning tree of  $G_1$  not annihilated by  $f$  is a **cospanning tree** of  $G_1$  and  $G_2$  under  $f$ .

Let  **$\text{cospan}(G_1, G_2)$**  be the maximum number of cospanning trees of  $G_1$  and  $G_2$  over  $f$ .

## proposition (Bu, P. 2025)

Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two graphs with  $|E_1| = |E_2| = |V_1| + |V_2| - 2$ .

Let  $M$  and  $N$  be truncated incidence matrices of  $G_1$  and  $G_2$ , respectively.

Then  **$\det[M|N] \leq \text{cospan}(G_1, G_2)$** .

In the proof for  $\varepsilon(K_{3,3})$  and  $\varepsilon(K_5)$ , we repeatedly **restrict the space of feasible graphs** corresponding to the right incidence submatrix through various bounds.

These bounds give us **increasingly restrictive information** about both the graph and properties of the bijection, which we then use to prove the desired results.



a conjecture

### corollary

For all  $m \in \mathbb{N}$ , we have  $T_m \leq \Delta_m$ .

### corollary

For all  $m \in \mathbb{N}$ , we have  $T_m \leq \Delta_m$ .

$T_m$  and  $\Delta_m$  for small values of  $m$

$m$	1	2	3	4	5	6	7	8	9	10
$T_m$	1	2	3	5	8	16	24	45	75	130
$\Delta_m$	1	2	3	5	8	16	24	45	75	130

### corollary

For all  $m \in \mathbb{N}$ , we have  $T_m \leq \Delta_m$ .

### $T_m$ and $\Delta_m$ for small values of $m$

$m$	1	2	3	4	5	6	7	8	9	10
$T_m$	1	2	3	5	8	16	24	45	75	130
$\Delta_m$	1	2	3	5	8	16	24	45	75	130

### conjecture

For all  $m \in \mathbb{N}$ , we have  $T_m = \Delta_m$ .

towards the conjecture

conjecture

For all  $m \in \mathbb{N}$ , we have  $T_m = \Delta_m$ .

### conjecture

For all  $m \in \mathbb{N}$ , we have  $T_m = \Delta_m$ .

### another conjecture

If  $G$  is a connected non-planar graph with  $m$  edges, then  $\Delta(G) \leq T_m$ .

### conjecture

For all  $m \in \mathbb{N}$ , we have  $T_m = \Delta_m$ .

### another conjecture

If  $G$  is a connected non-planar graph with  $m$  edges, then  $\Delta(G) \leq T_m$ .

**In other words:** The maximum  $\Delta(\cdot)$  value over graphs with the same number of edges is always achieved by **planar** graphs.



### conjecture

For all  $m \in \mathbb{N}$ , we have  $T_m = \Delta_m$ .

### another conjecture

If  $G$  is a connected non-planar graph with  $m$  edges, then  $\Delta(G) \leq T_m$ .

**In other words:** The maximum  $\Delta(\cdot)$  value over graphs with the same number of edges is always achieved by **planar** graphs.

**In other words:** Although a non-planar graph might have more spanning trees than the best planar graph with the same number of edges, we conjecture that it is “**dominated**” by the best planar graph with the same number of edges in some **linear-algebraic** sense.

### conjecture

For all  $m \in \mathbb{N}$ , we have  $T_m = \Delta_m$ .

### another conjecture

If  $G$  is a connected non-planar graph with  $m$  edges, then  $\Delta(G) \leq T_m$ .

**In other words:** The maximum  $\Delta(\cdot)$  value over graphs with the same number of edges is always achieved by **planar** graphs.

**In other words:** Although a non-planar graph might have more spanning trees than the best planar graph with the same number of edges, we conjecture that it is “**dominated**” by the best planar graph with the same number of edges in some **linear-algebraic** sense.

These two conjectures are equivalent.

### conjecture

For all  $m \in \mathbb{N}$ , we have  $T_m = \Delta_m$ .

### another conjecture

If  $G$  is a connected non-planar graph with  $m$  edges, then  $\Delta(G) \leq T_m$ .

**In other words:** The maximum  $\Delta(\cdot)$  value over graphs with the same number of edges is always achieved by **planar** graphs.

**In other words:** Although a non-planar graph might have more spanning trees than the best planar graph with the same number of edges, we conjecture that it is “**dominated**” by the best planar graph with the same number of edges in some **linear-algebraic** sense.

These two conjectures are equivalent. (Why?)

theorem (Bu, P. 2025)

If  $G$  is a subdivision of  $K_{3,3}$  or  $K_5$  with  $m$  edges, then  $\Delta(G) \leq T_m$ .

theorem (Bu, P. 2025)

If  $G$  is a subdivision of  $K_{3,3}$  or  $K_5$  with  $m$  edges, then  $\Delta(G) \leq T_m$ .

Idea for  $K_{3,3}$ : The **right** side of the matrix  $P$  attaining  $\Delta(G)$  is “**planar.**”

theorem (Bu, P. 2025)

If  $G$  is a subdivision of  $K_{3,3}$  or  $K_5$  with  $m$  edges, then  $\Delta(G) \leq T_m$ .

Idea for  $K_{3,3}$ : The **right** side of the matrix  $P$  attaining  $\Delta(G)$  is “**planar.**”

In particular, we use **multilinearity of determinants** to argue that, w.l.o.g., the right sides of rows corresponding to edges from subdividing the same edge in  $K_{3,3}$  are the same.

theorem (Bu, P. 2025)

If  $G$  is a subdivision of  $K_{3,3}$  or  $K_5$  with  $m$  edges, then  $\Delta(G) \leq T_m$ .

Idea for  $K_{3,3}$ : The **right** side of the matrix  $P$  attaining  $\Delta(G)$  is “**planar.**”

In particular, we use **multilinearity of determinants** to argue that, w.l.o.g., the right sides of rows corresponding to edges from subdividing the same edge in  $K_{3,3}$  are the same.

The right side of  $P$  has 5 columns and at most  $m$  rows that are not all-zero, among which there are at most 9 **distinct** rows.

theorem (Bu, P. 2025)

If  $G$  is a subdivision of  $K_{3,3}$  or  $K_5$  with  $m$  edges, then  $\Delta(G) \leq T_m$ .

Idea for  $K_{3,3}$ : The **right** side of the matrix  $P$  attaining  $\Delta(G)$  is “**planar.**”

In particular, we use **multilinearity of determinants** to argue that, w.l.o.g., the right sides of rows corresponding to edges from subdividing the same edge in  $K_{3,3}$  are the same.

The right side of  $P$  is a truncated incidence matrix of a graph  $H$  with 5 vertices and at most  $m$  edges, among which there are at most 9 **distinct** edges.



theorem (Bu, P. 2025)

If  $G$  is a subdivision of  $K_{3,3}$  or  $K_5$  with  $m$  edges, then  $\Delta(G) \leq T_m$ .

Idea for  $K_{3,3}$ : The **right** side of the matrix  $P$  attaining  $\Delta(G)$  is “**planar.**”

In particular, we use **multilinearity of determinants** to argue that, w.l.o.g., the right sides of rows corresponding to edges from subdividing the same edge in  $K_{3,3}$  are the same.

The right side of  $P$  is a truncated incidence matrix of a graph  $H$  with 5 vertices and at most  $m$  edges, among which there are at most 9 **distinct** edges.

Wagner says  $H$  is **planar**!

theorem (Bu, P. 2025)

If  $G$  is a subdivision of  $K_{3,3}$  or  $K_5$  with  $m$  edges, then  $\Delta(G) \leq T_m$ .

Idea for  $K_{3,3}$ : The **right** side of the matrix  $P$  attaining  $\Delta(G)$  is “**planar.**”

In particular, we use **multilinearity of determinants** to argue that, w.l.o.g., the right sides of rows corresponding to edges from subdividing the same edge in  $K_{3,3}$  are the same.

The right side of  $P$  is a truncated incidence matrix of a graph  $H$  with 5 vertices and at most  $m$  edges, among which there are at most 9 **distinct** edges.

Wagner says  $H$  is **planar**!  $\implies \Delta(G) = \det(P) = \Delta(H) \leq \tau(H) \leq T_m$

theorem (Bu, P. 2025)

If  $G$  is a subdivision of  $K_{3,3}$  or  $K_5$  with  $m$  edges, then  $\Delta(G) \leq T_m$ .

Idea for  $K_5$ : The **left** side of the matrix attaining  $\Delta(G)$  can be made “**planar**” by careful matrix operations

theorem (Bu, P. 2025)


If  $G$  is a subdivision of  $K_{3,3}$  or  $K_5$  with  $m$  edges, then  $\Delta(G) \leq T_m$ .

Idea for  $K_5$ : The **left** side of the matrix attaining  $\Delta(G)$  can be made “**planar**” by careful matrix operations (“**edge relocation method**”).

theorem (Bu, P. 2025)

If  $G$  is a subdivision of  $K_{3,3}$  or  $K_5$  with  $m$  edges, then  $\Delta(G) \leq T_m$ .

Idea for  $K_5$ : The **left** side of the matrix attaining  $\Delta(G)$  can be made “**planar**” by careful matrix operations (“**edge relocation method**”).



theorem (Bu, P. 2025)

If  $G$  is a subdivision of  $K_{3,3}$  or  $K_5$  with  $m$  edges, then  $\Delta(G) \leq T_m$ .

Idea for  $K_5$ : The **left** side of the matrix attaining  $\Delta(G)$  can be made “**planar**” by careful matrix operations (“**edge relocation method**”).





graphs of bounded genus

(ideas due to **Jack Spalding-Jamieson**)


Intuitively, the **genus** of a surface is the number of its “holes.”



Intuitively, the **genus** of a surface is the number of its “holes.”



Intuitively, the **genus** of a surface is the number of its “holes.”



### definition

The **genus** of a graph  $G$  is the smallest genus of a surface in which  $G$  can be embedded without crossings.

theorem (Bu, P. 2025)

If a graph  $G$  is **planar**, then  $\Delta(G) = \tau(G)$ .

If a graph  $G$  is **non-planar**, then  $\Delta(G) \leq \tau(G) - 18$ .

theorem (Bu, P. 2025)

If a graph  $G$  is **planar**, then  $\Delta(G) = \tau(G)$ .

If a graph  $G$  is **non-planar**, then  $\Delta(G) \leq \tau(G) - 18$ .

**Question:** For certain subclasses of **non-planar** graphs, can we obtain non-trivial lower bounds on  $\Delta(G)$ ?

### theorem (Bu, P. 2025)

If a graph  $G$  is **planar**, then  $\Delta(G) = \tau(G)$ .

If a graph  $G$  is **non-planar**, then  $\Delta(G) \leq \tau(G) - 18$ .

**Question:** For certain subclasses of **non-planar** graphs, can we obtain non-trivial lower bounds on  $\Delta(G)$ ?

### idea (Spalding-Jamieson)

If a graph has genus  $g$  and maximum degree  $D$ , then maybe  $\Delta(G) \geq f(g, D) \cdot \tau(G)$  for some function  $f(g, D) \in (0, 1)$ .

### theorem (Bu, P. 2025)

If a graph  $G$  is **planar**, then  $\Delta(G) = \tau(G)$ .

If a graph  $G$  is **non-planar**, then  $\Delta(G) \leq \tau(G) - 18$ .

**Question:** For certain subclasses of **non-planar** graphs, can we obtain non-trivial lower bounds on  $\Delta(G)$ ?

### idea (Spalding-Jamieson)

If a graph has genus  $g$  and maximum degree  $D$ , then maybe  $\Delta(G) \geq f(g, D) \cdot \tau(G)$  for some function  $f(g, D) \in (0, 1)$ .

planar graphs

graphs of genus  $g$

### theorem (Bu, P. 2025)

If a graph  $G$  is **planar**, then  $\Delta(G) = \tau(G)$ .

If a graph  $G$  is **non-planar**, then  $\Delta(G) \leq \tau(G) - 18$ .

**Question:** For certain subclasses of **non-planar** graphs, can we obtain non-trivial lower bounds on  $\Delta(G)$ ?

### idea (Spalding-Jamieson)

If a graph has genus  $g$  and maximum degree  $D$ , then maybe  $\Delta(G) \geq f(g, D) \cdot \tau(G)$  for some function  $f(g, D) \in (0, 1)$ .

planar graphs

graphs of genus  $g$

dual

dual, **but need more columns**

### theorem (Bu, P. 2025)

If a graph  $G$  is **planar**, then  $\Delta(G) = \tau(G)$ .

If a graph  $G$  is **non-planar**, then  $\Delta(G) \leq \tau(G) - 18$ .

**Question:** For certain subclasses of **non-planar** graphs, can we obtain non-trivial lower bounds on  $\Delta(G)$ ?

### idea (Spalding-Jamieson)

If a graph has genus  $g$  and maximum degree  $D$ , then maybe  $\Delta(G) \geq f(g, D) \cdot \tau(G)$  for some function  $f(g, D) \in (0, 1)$ .

planar graphs	graphs of genus $g$
dual	dual, <b>but need more columns</b>
$n - m + f = 2$	$n - m + f = 2 - 2g$



### theorem (Bu, P. 2025)

If a graph  $G$  is **planar**, then  $\Delta(G) = \tau(G)$ .

If a graph  $G$  is **non-planar**, then  $\Delta(G) \leq \tau(G) - 18$ .

**Question:** For certain subclasses of **non-planar** graphs, can we obtain non-trivial lower bounds on  $\Delta(G)$ ?

### idea (Spalding-Jamieson)

If a graph has genus  $g$  and maximum degree  $D$ , then maybe  $\Delta(G) \geq f(g, D) \cdot \tau(G)$  for some function  $f(g, D) \in (0, 1)$ .

planar graphs	graphs of genus $g$
dual	dual, <b>but need more columns</b>
$n - m + f = 2$	$n - m + f = 2 - 2g$
$M^T M^* = 0$	Schur complement

### theorem (Bu, P. 2025)

If a graph  $G$  is **planar**, then  $\Delta(G) = \tau(G)$ .

If a graph  $G$  is **non-planar**, then  $\Delta(G) \leq \tau(G) - 18$ .

**Question:** For certain subclasses of **non-planar** graphs, can we obtain non-trivial lower bounds on  $\Delta(G)$ ?

### idea (Spalding-Jamieson)

If a graph has genus  $g$  and maximum degree  $D$ , then maybe  $\Delta(G) \geq f(g, D) \cdot \tau(G)$  for some function  $f(g, D) \in (0, 1)$ .

planar graphs	graphs of genus $g$
dual	dual, <b>but need more columns</b>
$n - m + f = 2$	$n - m + f = 2 - 2g$
$M^T M^* = 0$	Schur complement
$\tau(G^*) = \tau(G)$	$\tau(G^*) = 2^{2g} \tau(G)$

future

Does  $T_m = \Delta_m$  hold?

Does  $T_m = \Delta_m$  hold?

Can we find better **lower bounds** than the square grid graphs?

Does  $T_m = \Delta_m$  hold?

Can we find better **lower bounds** than the square grid graphs?

Can we extend the **planar dominance theorem** beyond subdivisions of  $K_{3,3}$  and  $K_5$ ?

Does  $T_m = \Delta_m$  hold?

Can we find better **lower bounds** than the square grid graphs?

Can we extend the **planar dominance theorem** beyond subdivisions of  $K_{3,3}$  and  $K_5$ ?

Consequences on **algorithms**?

Does  $T_m = \Delta_m$  hold?

Can we find better **lower bounds** than the square grid graphs?

Can we extend the **planar dominance theorem** beyond subdivisions of  $K_{3,3}$  and  $K_5$ ?

Consequences on **algorithms**?

Consequences on **mathematical programming**?



Thanks (for listening and for the ice cream)!