

# *Roundtrip Spanners and Roundtrip Routing in Directed Graphs*

by Roditty, Thorup, and Zwick (2008)

Yuchong Pan

MIT 6.890

December 9, 2021

# Spanners in **Undirected** Graphs

## Definition (Spanner)

Let  $G = (V, E)$  be a weighted **undirected** graph. A subgraph  $H$  of  $G$  is a  $t$ -**spanner** if  $d_H(u, v) \leq t \cdot d_G(u, v)$  for all  $u, v \in V$ .

# Spanners in **Undirected** Graphs

## Definition (Spanner)

Let  $G = (V, E)$  be a weighted **undirected** graph. A subgraph  $H$  of  $G$  is a  $t$ -**spanner** if  $d_H(u, v) \leq t \cdot d_G(u, v)$  for all  $u, v \in V$ .

## Theorem (Althöfer et al., 1993)

*For any  $k \in \mathbb{N}$ , any weighted **undirected** graph on  $n$  vertices has a  $(2k - 1)$ -spanner with  $O(n^{1+1/k})$  edges.*

# Spanners in **Undirected** Graphs

## Definition (Spanner)

Let  $G = (V, E)$  be a weighted **undirected** graph. A subgraph  $H$  of  $G$  is a  $t$ -**spanner** if  $d_H(u, v) \leq t \cdot d_G(u, v)$  for all  $u, v \in V$ .

## Theorem (Althöfer et al., 1993)

*For any  $k \in \mathbb{N}$ , any weighted **undirected** graph on  $n$  vertices has a  $(2k - 1)$ -spanner with  $O(n^{1+1/k})$  edges.*

**Directed graphs?**

# Spanners in **Undirected** Graphs


## Definition (Spanner)

Let  $G = (V, E)$  be a weighted **undirected** graph. A subgraph  $H$  of  $G$  is a  $t$ -**spanner** if  $d_H(u, v) \leq t \cdot d_G(u, v)$  for all  $u, v \in V$ .

## Theorem (Althöfer et al., 1993)

*For any  $k \in \mathbb{N}$ , any weighted **undirected** graph on  $n$  vertices has a  $(2k - 1)$ -spanner with  $O(n^{1+1/k})$  edges.*

**Directed graphs?**



# Spanners in **Undirected** Graphs


## Definition (Spanner)

Let  $G = (V, E)$  be a weighted **undirected** graph. A subgraph  $H$  of  $G$  is a  $t$ -**spanner** if  $d_H(u, v) \leq t \cdot d_G(u, v)$  for all  $u, v \in V$ .

## Theorem (Althöfer et al., 1993)

For any  $k \in \mathbb{N}$ , any weighted **undirected** graph on  $n$  vertices has a  $(2k - 1)$ -spanner with  $O(n^{1+1/k})$  edges.

**Directed graphs?**



# Roundtrip Spanners

## Definition (Roundtrip spanner)

Let  $G$  be a weighted **directed** graph. A subgraph  $H$  of  $G$  is a  **$t$ -roundtrip-spanner** if  $d_H(u \rightrightarrows v) \leq t \cdot d_G(u \rightrightarrows v)$  for  $u, v \in V$ .

# Roundtrip Spanners

## Definition (Roundtrip spanner)

Let  $G$  be a weighted **directed** graph. A subgraph  $H$  of  $G$  is a  **$t$ -roundtrip-spanner** if  $d_H(u \rightrightarrows v) \leq t \cdot d_G(u \rightrightarrows v)$  for  $u, v \in V$ .

## Theorem (Cowen and Wagner, 2000)

*For any  $k \in \mathbb{N}$ , any weighted **directed** graph on  $n$  vertices has a  $(2^k - 1)$ -roundtrip-spanner with  $\tilde{O}(n^{1+1/k})$  edges.*



# Roundtrip Spanners

## Definition (Roundtrip spanner)

Let  $G$  be a weighted **directed** graph. A subgraph  $H$  of  $G$  is a  **$t$ -roundtrip-spanner** if  $d_H(u \rightleftarrows v) \leq t \cdot d_G(u \rightleftarrows v)$  for  $u, v \in V$ .

## Theorem (Cowen and Wagner, 2000)

*For any  $k \in \mathbb{N}$ , any weighted **directed** graph on  $n$  vertices has a  $(2^k - 1)$ -roundtrip-spanner with  $\tilde{O}(n^{1+1/k})$  edges.*

## Theorem (Roditty, Thorup, and Zwick, 2008)

*For any  $k \in \mathbb{N}, \varepsilon > 0$ , any weighted **directed** graph on  $n$  vertices with edge weights from  $[1, W]$  has a  $(2k + \varepsilon)$ -roundtrip-spanner with  $O((k^2/\varepsilon)n^{1+1/k} \log(nW))$  edges.*

# A General Technique of Cohen (1998)

# A General Technique of Cohen (1998)

## Procedure PARTIALSPANNER( $R$ )

**Input:**  $R \in [1, 2nW]$ .

**Output:** A subgraph  $H$  of  $G$  with  $O(kn^{1+1/k})$  edges such that  $d_H(u \rightleftharpoons v) \leq 2kR$  for all  $u, v \in V$  with  $d_G(u \rightleftharpoons v) \leq R$ .

# A General Technique of Cohen (1998)

## Procedure PARTIALSPANNER( $R$ )

**Input:**  $R \in [1, 2nW]$ .

**Output:** A subgraph  $H$  of  $G$  with  $O(kn^{1+1/k})$  edges such that  $d_H(u \rightleftharpoons v) \leq 2kR$  for all  $u, v \in V$  with  $d_G(u \rightleftharpoons v) \leq R$ .

## The Approach of Cohen (1998)

```
1 foreach  $i \leftarrow 1, \dots, \log_{1+\varepsilon}(2nW)$  do  
2    $R_i \leftarrow (1 + \varepsilon)^i$   
3    $H_i \leftarrow \text{PARTIALSPANNER}(R_i)$   
4  $H \leftarrow \bigcup \{H_i : i \in [\log_{1+\varepsilon}(2nW)]\}$   
5 return  $H$ 
```

# A General Technique of Cohen (1998)

## Procedure PARTIALSPANNER( $R$ )

**Input:**  $R \in [1, 2nW]$ .

**Output:** A subgraph  $H$  of  $G$  with  $O(kn^{1+1/k})$  edges such that  $d_H(u \rightleftharpoons v) \leq 2kR$  for all  $u, v \in V$  with  $d_G(u \rightleftharpoons v) \leq R$ .

## The Approach of Cohen (1998)

```
1 foreach  $i \leftarrow 1, \dots, \log_{1+\varepsilon}(2nW)$  do  
2    $R_i \leftarrow (1 + \varepsilon)^i$   
3    $H_i \leftarrow \text{PARTIALSPANNER}(R_i)$   
4  $H \leftarrow \bigcup \{H_i : i \in [\log_{1+\varepsilon}(2nW)]\}$   
5 return  $H$ 
```

**Claim:**  $H$  is a  $2k(1 + \varepsilon)$ -roundtrip-spanner of  $G$  with  $O(kn^{1+1/k} \log_{1+\varepsilon}(nW))$  edges.

# Constructing Partial Spanners

## Definition (Ball)

For every  $V' \subset V$ ,  $v \in V$  and  $r \geq 0$ , let

$$\text{ball}_{V'}(v, r) = \{u \in V' : d_{G[V']}(v \rightleftharpoons u) \leq r\},$$

called a **ball** centered at  $v$  of radius  $r$ .

# Constructing Partial Spanners

## Definition (Ball)

For every  $V' \subset V$ ,  $v \in V$  and  $r \geq 0$ , let

$$\text{ball}_{V'}(v, r) = \{u \in V' : d_{G[V']}(v \rightleftarrows u) \leq r\},$$

called a **ball** centered at  $v$  of radius  $r$ .

## Definition ( $(k, R)$ -cover)

A set  $\mathcal{C}$  of balls is a  $(k, R)$ -**cover** of a directed graph  $G = (V, E)$  if each ball in  $\mathcal{C}$  is of radius at most  $kR$ , and for all  $u, v \in V$  with  $d_G(u \rightleftarrows v) \leq R$ , there is a ball  $B \in \mathcal{C}$  such that  $u, v \in B$ .

# Constructing Partial Spanners

## Procedure COVER( $G, k, R$ )

```
1  $\mathcal{C} \leftarrow \emptyset$ 
2  $V_{k-1} \leftarrow V(G)$ 
3 foreach  $i \leftarrow k-1, \dots, 0$  do
4    $S_i \leftarrow \text{SAMPLE}(V_i, n^{-i/k})$ 
5    $\mathcal{C} \leftarrow \mathcal{C} \cup \{\text{ball}_{V_i}(v, (i+1)R) : v \in S_i\}$ 
6    $V_{i-1} \leftarrow V_i \setminus \bigcup_{v \in S_i} \text{ball}_{V_i}(v, iR)$ 
7 return  $\mathcal{C}$ 
```



# Constructing Partial Spanners

## Procedure COVER( $G, k, R$ )

```
1  $\mathcal{C} \leftarrow \emptyset$ 
2  $V_{k-1} \leftarrow V(G)$ 
3 foreach  $i \leftarrow k-1, \dots, 0$  do
4    $S_i \leftarrow \text{SAMPLE}(V_i, n^{-i/k})$ 
5    $\mathcal{C} \leftarrow \mathcal{C} \cup \{\text{ball}_{V_i}(v, (i+1)R) : v \in S_i\}$ 
6    $V_{i-1} \leftarrow V_i \setminus \bigcup_{v \in S_i} \text{ball}_{V_i}(v, iR)$ 
7 return  $\mathcal{C}$ 
```

## Theorem

*The collection  $\mathcal{C} = \text{COVER}(G, k, R)$  is a  $(k, R)$ -cover of a directed graph  $G = (V, E)$ . For each  $v \in V$ , the **expected** number of balls in  $\mathcal{C}$  containing  $v$  is at most  $kn^{1/k}$ .*

COVER( $G, k, R$ ) Is a  $(k, R)$ -Cover

## COVER( $G, k, R$ ) Is a $(k, R)$ -Cover

- Let  $u, v \in V$  be such that  $d_G(u \rightleftharpoons v) \leq R$ .

## COVER( $G, k, R$ ) Is a $(k, R)$ -Cover

- Let  $u, v \in V$  be such that  $d_G(u \rightleftharpoons v) \leq R$ .
- Let  $c$  be a closed tour in  $G$  of length  $\leq R$  containing  $u$  and  $v$ .

## COVER( $G, k, R$ ) Is a $(k, R)$ -Cover

- Let  $u, v \in V$  be such that  $d_G(u \rightleftharpoons v) \leq R$ .
- Let  $c$  be a closed tour in  $G$  of length  $\leq R$  containing  $u$  and  $v$ .
- Let  $i$  be the largest index so that a vertex  $u'$  on  $c$  is contained in the **core**  $\text{ball}_{V_i}(w, iR)$  of  $\text{ball}_{V_i}(w, (i+1)R)$  added to  $\mathcal{C}$ .

## COVER( $G, k, R$ ) Is a $(k, R)$ -Cover

- Let  $u, v \in V$  be such that  $d_G(u \rightleftharpoons v) \leq R$ .
- Let  $c$  be a closed tour in  $G$  of length  $\leq R$  containing  $u$  and  $v$ .
- Let  $i$  be the largest index so that a vertex  $u'$  on  $c$  is contained in the **core**  $\text{ball}_{V_i}(w, iR)$  of  $\text{ball}_{V_i}(w, (i+1)R)$  added to  $\mathcal{C}$ .
- By maximality of  $i$ , all the vertices on  $c$  are contained in  $V_i$ .

## COVER( $G, k, R$ ) Is a $(k, R)$ -Cover

- Let  $u, v \in V$  be such that  $d_G(u \rightleftharpoons v) \leq R$ .
- Let  $c$  be a closed tour in  $G$  of length  $\leq R$  containing  $u$  and  $v$ .
- Let  $i$  be the largest index so that a vertex  $u'$  on  $c$  is contained in the **core**  $\text{ball}_{V_i}(w, iR)$  of  $\text{ball}_{V_i}(w, (i+1)R)$  added to  $\mathcal{C}$ .
- By maximality of  $i$ , all the vertices on  $c$  are contained in  $V_i$ .
- By the triangle inequality,

$$\begin{aligned} d_{G[V_i]}(w \rightleftharpoons v) &\leq d_{G[V_i]}(w \rightleftharpoons u) + d_{G[V_i]}(u \rightleftharpoons v) \\ &\leq iR + R = (i+1)R. \end{aligned}$$

## COVER( $G, k, R$ ) Is a $(k, R)$ -Cover

- Let  $u, v \in V$  be such that  $d_G(u \rightleftharpoons v) \leq R$ .
- Let  $c$  be a closed tour in  $G$  of length  $\leq R$  containing  $u$  and  $v$ .
- Let  $i$  be the largest index so that a vertex  $u'$  on  $c$  is contained in the **core**  $\text{ball}_{V_i}(w, iR)$  of  $\text{ball}_{V_i}(w, (i+1)R)$  added to  $\mathcal{C}$ .
- By maximality of  $i$ , all the vertices on  $c$  are contained in  $V_i$ .
- By the triangle inequality,

$$\begin{aligned} d_{G[V_i]}(w \rightleftharpoons v) &\leq d_{G[V_i]}(w \rightleftharpoons u) + d_{G[V_i]}(u \rightleftharpoons v) \\ &\leq iR + R = (i+1)R. \end{aligned}$$

- Therefore,  $v \in \text{ball}_{V_i}(w, (i+1)R)$ .



## COVER( $G, k, R$ ) Is a $(k, R)$ -Cover

- Let  $u, v \in V$  be such that  $d_G(u \rightleftharpoons v) \leq R$ .
- Let  $c$  be a closed tour in  $G$  of length  $\leq R$  containing  $u$  and  $v$ .
- Let  $i$  be the largest index so that a vertex  $u'$  on  $c$  is contained in the **core**  $\text{ball}_{V_i}(w, iR)$  of  $\text{ball}_{V_i}(w, (i+1)R)$  added to  $\mathcal{C}$ .
- By maximality of  $i$ , all the vertices on  $c$  are contained in  $V_i$ .
- By the triangle inequality,

$$\begin{aligned} d_{G[V_i]}(w \rightleftharpoons v) &\leq d_{G[V_i]}(w \rightleftharpoons u) + d_{G[V_i]}(u \rightleftharpoons v) \\ &\leq iR + R = (i+1)R. \end{aligned}$$

- Therefore,  $v \in \text{ball}_{V_i}(w, (i+1)R)$ .
- Similarly,  $u \in \text{ball}_{V_i}(w, (i+1)R)$ , completing the proof.

# Expected Number of Balls Containing a Vertex

# Expected Number of Balls Containing a Vertex

## Lemma

*In each iteration, the expected number of balls containing  $v \in V$  that are added to  $\mathcal{C}$  is at most  $n^{1/k}$ .*

# Expected Number of Balls Containing a Vertex

## Lemma

*In each iteration, the expected number of balls containing  $v \in V$  that are added to  $\mathcal{C}$  is at most  $n^{1/k}$ .*

- Let  $B = \text{ball}_{V_{i+1}}(v, (i+1)R)$ .

# Expected Number of Balls Containing a Vertex

## Lemma

*In each iteration, the expected number of balls containing  $v \in V$  that are added to  $\mathcal{C}$  is at most  $n^{1/k}$ .*

- Let  $B = \text{ball}_{V_{i+1}}(v, (i+1)R)$ .
- If  $u \in B$  is chosen to  $S_{i+1}$  (with probability  $n^{-(i+1)/k}$ ), then  $v \in \text{ball}_{V_{i+1}}(u, (i+1)R)$  and thus removed in iteration  $i+1$ .

# Expected Number of Balls Containing a Vertex

## Lemma

*In each iteration, the expected number of balls containing  $v \in V$  that are added to  $\mathcal{C}$  is at most  $n^{1/k}$ .*

- Let  $B = \text{ball}_{V_{i+1}}(v, (i+1)R)$ .
- If  $u \in B$  is chosen to  $S_{i+1}$  (with probability  $n^{-(i+1)/k}$ ), then  $v \in \text{ball}_{V_{i+1}}(u, (i+1)R)$  and thus removed in iteration  $i+1$ .
- If  $v \in \text{ball}_{V_i}(u, (i+1)R) \subset \text{ball}_{V_{i+1}}(u, (i+1)R)$ , then  $u \in B$ .

# Expected Number of Balls Containing a Vertex

## Lemma

*In each iteration, the expected number of balls containing  $v \in V$  that are added to  $\mathcal{C}$  is at most  $n^{1/k}$ .*

- Let  $B = \text{ball}_{V_{i+1}}(v, (i+1)R)$ .
- If  $u \in B$  is chosen to  $S_{i+1}$  (with probability  $n^{-(i+1)/k}$ ), then  $v \in \text{ball}_{V_{i+1}}(u, (i+1)R)$  and thus removed in iteration  $i+1$ .
- If  $v \in \text{ball}_{V_i}(u, (i+1)R) \subset \text{ball}_{V_{i+1}}(u, (i+1)R)$ , then  $u \in B$ .
- $\mathbb{P}[u \in S_i \wedge v \in \text{ball}_{V_i}(u, (i+1)R)] = (1 - n^{-(i+1)/k})^{|B|} n^{-i/k}$ .

# Expected Number of Balls Containing a Vertex


## Lemma

*In each iteration, the expected number of balls containing  $v \in V$  that are added to  $\mathcal{C}$  is at most  $n^{1/k}$ .*


- Let  $B = \text{ball}_{V_{i+1}}(v, (i+1)R)$ .
- If  $u \in B$  is chosen to  $S_{i+1}$  (with probability  $n^{-(i+1)/k}$ ), then  $v \in \text{ball}_{V_{i+1}}(u, (i+1)R)$  and thus removed in iteration  $i+1$ .
- If  $v \in \text{ball}_{V_i}(u, (i+1)R) \subset \text{ball}_{V_{i+1}}(u, (i+1)R)$ , then  $u \in B$ .
- $\mathbb{P}[u \in S_i \wedge v \in \text{ball}_{V_i}(u, (i+1)R)] = (1 - n^{-(i+1)/k})^{|B|} n^{-i/k}$ .
- The expected number of balls containing  $v$  in iteration  $i$  is  $|B| (1 - n^{-(i+1)/k})^{|B|} n^{-i/k}$ .



# Expected Number of Balls Containing a Vertex



# Expected Number of Balls Containing a Vertex



$\mathbb{E}[\# \text{ balls containing } v \text{ in iteration } i]$

$$= |B| \left(1 - n^{-(i+1)/k}\right)^{|B|} n^{-i/k} \leq \frac{n^{-i/k}}{en^{-(i+1)/k}} = \frac{n^{1/k}}{e}.$$

# Double-Trees

## Definition (Double-tree)

Let  $G = (V, E)$  be a weighted directed graph. Let  $V' \subset V$ ,  $v \in V$  and  $r \geq 0$ . Let  $B = \text{ball}_{V'}(v, r)$ .

Let  $\text{OutTree}(B, v)$  be a tree containing directed shortest paths in  $G[V']$  from  $v$  to all the vertices in  $B$ .

Let  $\text{InTree}(B, v)$  be a tree containing directed shortest paths in  $G[V']$  from all the vertices in  $B$  to  $v$ .

Let  $\text{InOutTrees}(B, v) = \text{InTree}(B, v) \cup \text{OutTree}(B, v)$ , referred to as a **double-tree**.

# Double-Trees

## Definition (Double-tree)

Let  $G = (V, E)$  be a weighted directed graph. Let  $V' \subset V$ ,  $v \in V$  and  $r \geq 0$ . Let  $B = \text{ball}_{V'}(v, r)$ .

Let  $\text{OutTree}(B, v)$  be a tree containing directed shortest paths in  $G[V']$  from  $v$  to all the vertices in  $B$ .

Let  $\text{InTree}(B, v)$  be a tree containing directed shortest paths in  $G[V']$  from all the vertices in  $B$  to  $v$ .

Let  $\text{InOutTrees}(B, v) = \text{InTree}(B, v) \cup \text{OutTree}(B, v)$ , referred to as a **double-tree**.

## Lemma

*Let  $V' \subset V$  and  $v \in V$ . If  $u \in B := \text{ball}_{V'}(v, r)$  and  $w$  is on a shortest path in  $G[V']$  from  $u$  to  $v$ , or from  $v$  to  $u$ , then  $w \in B$ .*

# Double-Trees

## Definition (Double-tree)

Let  $G = (V, E)$  be a weighted directed graph. Let  $V' \subset V$ ,  $v \in V$  and  $r \geq 0$ . Let  $B = \text{ball}_{V'}(v, r)$ .

Let  $\text{OutTree}(B, v)$  be a tree containing directed shortest paths in  $G[V']$  from  $v$  to all the vertices in  $B$ .

Let  $\text{InTree}(B, v)$  be a tree containing directed shortest paths in  $G[V']$  from all the vertices in  $B$  to  $v$ .

Let  $\text{InOutTrees}(B, v) = \text{InTree}(B, v) \cup \text{OutTree}(B, v)$ , referred to as a **double-tree**.

## Lemma

*Let  $V' \subset V$  and  $v \in V$ . If  $u \in B := \text{ball}_{V'}(v, r)$  and  $w$  is on a shortest path in  $G[V']$  from  $u$  to  $v$ , or from  $v$  to  $u$ , then  $w \in B$ .*

## Corollary

$V(\text{InOutTrees}(B, v)) \subset B$ , so  $e(\text{InOutTrees}(B, v)) \leq 2(|B| - 1)$ .

# Double-Trees

## Lemma

*Let  $V' \subset V, v \in V$ . If  $u_1, u_2 \in B := \text{ball}_{V'}(v, r)$ ,  $\text{InOutTrees}(B, v)$  contains a closed directed tour containing  $u_1, u_2$  of length  $\leq 2r$ .*

# Double-Trees

## Lemma

*Let  $V' \subset V, v \in V$ . If  $u_1, u_2 \in B := \text{ball}_{V'}(v, r)$ ,  $\text{InOutTrees}(B, v)$  contains a closed directed tour containing  $u_1, u_2$  of length  $\leq 2r$ .*

**Obtaining a  $(2k + \varepsilon)$ -roundtrip-spanner of  $G$ :**

# Double-Trees

## Lemma

*Let  $V' \subset V, v \in V$ . If  $u_1, u_2 \in B := \text{ball}_{V'}(v, r)$ ,  $\text{InOutTrees}(B, v)$  contains a closed directed tour containing  $u_1, u_2$  of length  $\leq 2r$ .*

**Obtaining a  $(2k + \varepsilon)$ -roundtrip-spanner of  $G$ :**

- Let  $\varepsilon' = \varepsilon/(2k)$ .



# Double-Trees

## Lemma

*Let  $V' \subset V, v \in V$ . If  $u_1, u_2 \in B := \text{ball}_{V'}(v, r)$ ,  $\text{InOutTrees}(B, v)$  contains a closed directed tour containing  $u_1, u_2$  of length  $\leq 2r$ .*

## Obtaining a $(2k + \varepsilon)$ -roundtrip-spanner of $G$ :

- Let  $\varepsilon' = \varepsilon/(2k)$ .
- For  $i \in [\log_{1+\varepsilon'}(2nW)]$ , let  $\mathcal{C}_i = \text{PARTIALSPANNER}(G, k, R_i)$  be a  $(k, R_i)$ -cover, where  $R_i = (1 + \varepsilon')^i$ .

# Double-Trees

## Lemma

*Let  $V' \subset V, v \in V$ . If  $u_1, u_2 \in B := \text{ball}_{V'}(v, r)$ ,  $\text{InOutTrees}(B, v)$  contains a closed directed tour containing  $u_1, u_2$  of length  $\leq 2r$ .*

## Obtaining a $(2k + \varepsilon)$ -roundtrip-spanner of $G$ :

- Let  $\varepsilon' = \varepsilon/(2k)$ .
- For  $i \in [\log_{1+\varepsilon'}(2nW)]$ , let  $\mathcal{C}_i = \text{PARTIALSPANNER}(G, k, R_i)$  be a  $(k, R_i)$ -cover, where  $R_i = (1 + \varepsilon')^i$ .
- Let  $H = \bigcup \{\text{InOutTrees}(B) : B \in \bigcup_i \mathcal{C}_i\}$ .

# Double-Trees

## Lemma

*Let  $V' \subset V, v \in V$ . If  $u_1, u_2 \in B := \text{ball}_{V'}(v, r)$ ,  $\text{InOutTrees}(B, v)$  contains a closed directed tour containing  $u_1, u_2$  of length  $\leq 2r$ .*

## Obtaining a $(2k + \varepsilon)$ -roundtrip-spanner of $G$ :

- Let  $\varepsilon' = \varepsilon/(2k)$ .
- For  $i \in [\log_{1+\varepsilon'}(2nW)]$ , let  $\mathcal{C}_i = \text{PARTIALSPANNER}(G, k, R_i)$  be a  $(k, R_i)$ -cover, where  $R_i = (1 + \varepsilon')^i$ .
- Let  $H = \bigcup \{\text{InOutTrees}(B) : B \in \bigcup_i \mathcal{C}_i\}$ .
- Therefore,  $H$  is a  $(2k + \varepsilon)$ -roundtrip-spanner of  $G$  with  $O((k^2/\varepsilon)n^{1+1/k} \log(nW))$  edges **in expectation**.

# Double-Trees

## Lemma

*Let  $V' \subset V, v \in V$ . If  $u_1, u_2 \in B := \text{ball}_{V'}(v, r)$ ,  $\text{InOutTrees}(B, v)$  contains a closed directed tour containing  $u_1, u_2$  of length  $\leq 2r$ .*

## Obtaining a $(2k + \varepsilon)$ -roundtrip-spanner of $G$ :

- Let  $\varepsilon' = \varepsilon/(2k)$ .
- For  $i \in [\log_{1+\varepsilon'}(2nW)]$ , let  $\mathcal{C}_i = \text{PARTIALSPANNER}(G, k, R_i)$  be a  $(k, R_i)$ -cover, where  $R_i = (1 + \varepsilon')^i$ .
- Let  $H = \bigcup \{\text{InOutTrees}(B) : B \in \bigcup_i \mathcal{C}_i\}$ .
- Therefore,  $H$  is a  $(2k + \varepsilon)$ -roundtrip-spanner of  $G$  with  $O((k^2/\varepsilon)n^{1+1/k} \log(nW))$  edges **in expectation**.

**Application:** Compact roundtrip routing schemes.

# State-of-the-Art

## Theorem (Cen, Duan, and Gu, 2019)

*For any  $k \in \mathbb{N}$ , any weighted directed graph on  $n$  vertices has a  $(2k - 1)$ -roundtrip-spanner with  $O(kn^{1+1/k} \log n)$  edges.*

# State-of-the-Art

## Theorem (Cen, Duan, and Gu, 2019)

*For any  $k \in \mathbb{N}$ , any weighted directed graph on  $n$  vertices has a  $(2k - 1)$ -roundtrip-spanner with  $O(kn^{1+1/k} \log n)$  edges.*

## Proposition

*If the Erdős girth conjecture is true, there is an **undirected** graph on  $n$  vertices such that any  $(2k - 1)$ -spanner has  $\Omega(n^{1+1/k})$  edges.*

Thank you and happy holidays!