# A projective plane of order 18 admits no collineation of order 17 (computer-assisted, conditional on assumptions A1–A5)

Yudai Kobayashi

**Abstract**

We prove that a projective plane of order 18 has no collineation $\sigma$ of order 17. Counting arguments and Bruck's bound on subplanes (only $m^2 \le n$ is used) show that $\sigma$ has either exactly three fixed points, forming a triangle, or exactly twenty, namely a line plus an external point. The second case is excluded by an orbit-matrix argument over $\mathbb{Z}[\zeta_{17}]$: since 2 has order 8 modulo 17, the primes above 2 are self-conjugate, which forces the number of non-fixed orbits to be even, but it is 19. For the first case we derive from the axioms a reduction to 58,161 equivalence classes of orthomorphisms of $\mathbb{Z}_{17}$ with pairwise compatibility conditions on "rows"; an exhaustive computer search shows that no class has a solution. The triangle case is therefore computer-assisted, and the theorem holds under assumptions A1–A5 (Section 7.3). Baer's theorem on involutions is not used.

## 1 Introduction

A projective plane of order $n$ has $n^2 + n + 1$ points, $n + 1$ points on every line, and a unique line through any two points. Planes of prime-power order exist; none is known otherwise, and the Bruck–Ryser theorem [4] excludes infinitely many orders. Order 10 was excluded by a large computation [7] building on coding theory [8]. Order $18 = 3^2 + 3^2$ is not excluded by Bruck–Ryser.

**Theorem 1.1.** *A projective plane $\Pi$ of order 18 has no collineation of order 17. This is computer-assisted: the case of three fixed points is proved by an exhaustive search, and the theorem holds under the assumptions A1–A5 of Section 7.3.*

**Context.** Baer [2, Thm. 5, Cor. 6] shows that in a plane of non-square order every involution is a perspectivity; we do not use this. MacWilliams–Sloane–Thompson [8] (text not obtained; we rely on the report in [7, Sec. 2]), the survey [6], and [1, 9] concern order 10 and codes of designs. Janko–Tran Van Trung [5] (text not obtained, not relied on) restrict collineation groups for small orders. Orbit matrices over group rings are a standard tool; we found no primary reference stating the norm-parity lemma of Section 4 in the form used here and make no claim that it is new. We were *unable to verify* whether Theorem 1.1 or the exclusion of the homology configuration already appears in the literature (e.g. for orders $12, 15, 18$ or orbit matrices of $(343, 19, 1)$ designs); the elementary parts may be routine to experts. We do not claim that Theorem 1.1 is new; we were unable to find it in the literature we could access. The reduction of the triangle case to orthomorphisms of $\mathbb{Z}_{17}$ and its complete enumeration may be of interest.

## 2 Preliminaries

Throughout, $\Pi$ has order $n = 18$, $v = n^2 + n + 1 = 343$, and $\sigma$ is a collineation of prime order $p = 17$; orbit indices live in $\mathbb{Z}_{17} = \mathbb{Z}/17$. Quoted facts (nothing else from the literature is used):

**Q1.** (Counting.) The incidence matrix $N$ satisfies $NN^T = N^T N = nI + J$.

**Q2.** (Bruck [3, Lemma 3.1].) If a plane of order $n$ has a proper subplane of order $m$, then $n = m^2$ or $n \ge m^2 + m$; we use only $m^2 \le n$.

**Q3.** (Cyclotomic fields.) For $\zeta = e^{2\pi i/p}$, the ring of integers of $K = \mathbb{Q}(\zeta)$ is $\mathbb{Z}[\zeta]$, $\mathrm{Gal}(K/\mathbb{Q}) \cong (\mathbb{Z}/p)^*$ via $\gamma_a : \zeta \mapsto \zeta^a$, and $\gamma_{-1}$ is complex conjugation. A prime $q \neq p$ is unramified, and the decomposition group of each prime above $q$ is generated by $\gamma_q$.

**Lemma 2.1.** *Let $\sigma$ have prime order $p$ in a plane of order $n$, with $f$ fixed points.*

*(a) Orbits on points and on lines have length 1 or $p$; $f \equiv n^2 + n + 1 \pmod p$, and a fixed line carrying $f_L$ fixed points has $f_L \equiv n + 1 \pmod p$.*

*(b) The numbers of fixed points and of fixed lines are equal; hence so are the numbers of non-fixed point orbits and line orbits.*

*(c) The line through two fixed points is fixed; the intersection of two fixed lines is fixed.*

*Proof.* (a) Orbit lengths divide $p$. Non-fixed points form orbits of length $p$; so do the non-fixed points of a fixed line, which has $n + 1$ points. (b) Let $A, B$ be the permutation matrices of $\sigma$ on points and lines; $AN = NB$. By Q1, $\det N \neq 0$ ($\det(nI + J) = n^{v-1}(n + v)$), so $B = N^{-1}AN$ and $\operatorname{tr} A = \operatorname{tr} B$, these being the numbers of fixed points and lines. (c) $\sigma$ maps the line $xy$ ($x, y$ fixed) to a line through $x, y$; dually. □

Here $343 \equiv 3$ and $19 \equiv 2 \pmod{17}$.

## 3 The fixed structure

Let $F = \mathrm{Fix}(\sigma)$, $f = |F|$. A *quadrangle* is a set of four points, no three collinear.

**Lemma 3.1.** *If $F$ contains a quadrangle, then $F$ is the point set of a proper subplane of order $m \geq 2$, and $f = m^2 + m + 1$.*

*Proof.* Let $\mathcal{L}_F$ be the fixed lines. By Lemma 2.1(c), two points of $F$ lie on a unique line, which is in $\mathcal{L}_F$, and two lines of $\mathcal{L}_F$ meet in a point of $F$. The quadrangle makes the structure non-degenerate, so $(F, \mathcal{L}_F)$ is a projective plane of some order $m \geq 2$. If $m = n$ then $F$ is everything and $\sigma = \mathrm{id}$; so $m < n$. □

**Lemma 3.2.** *If $F$ contains no quadrangle, there are a line $\ell$ and a point $P$ with $F \subseteq \ell \cup \{P\}$.*

*Proof.* If $F$ is collinear, take $\ell \supseteq F$. Otherwise take non-collinear $A, B, C \in F$ with *sides* $AB, AC, BC$, and $X = F \setminus \{A, B, C\}$. *Step 1: each $D \in X$ lies on a side.* Otherwise no three of $A, B, C, D$ are collinear, so $\{A, B, C, D\}$ is a quadrangle. *Step 2.* Two sides meet in a vertex, which is not in $X$; so by Step 1 the parts of $X$ on the three sides partition $X$. *Step 3: $X$ meets at most one side.* Let two sides be $UV, UW$ (common vertex $U$), and suppose $D \in X \cap UV$, $E \in X \cap UW$; then $D, E \neq U$. We claim $\{V, W, D, E\}$ is a quadrangle: $V, W, D$ collinear would put $W$ on $VD = UV$ (but $W \notin UV$); $V, W, E$ collinear would put $V$ on $WE = UW$ (but $V \notin UW$); $V, D, E$ collinear would put $E$ on $VD = UV$, hence $E \in UV \cap UW = \{U\}$; $W, D, E$ collinear would put $D = U$. All are impossible, a contradiction. Hence $X$ lies on one side, say $AB$, and $F \subseteq AB \cup \{C\}$. □

**Theorem 3.3** (Fixed structure)**.** *Let $\sigma$ have order 17 in a plane of order 18. Then $f \in \{3, 20\}$, and*

*(i) if $f = 3$, the fixed points $P_0, P_1, P_2$ form a triangle, the fixed lines are exactly its three sides, and each side carries exactly two fixed points;*

*(ii) if $f = 20$, $F = \ell \cup \{P\}$ with $P \notin \ell$, and there are exactly 20 fixed lines.*

*The number of non-fixed point orbits (equal to that of line orbits) is $r = (343 - f)/17$, i.e. $r = 20$ for $f = 3$ and $r = 19$ for $f = 20$.*

*Proof.* By Lemma 2.1(a), $f \equiv 3 \pmod{17}$. If $F$ contains a quadrangle, Lemma 3.1 and Q2 give $m^2 \leq 18$, $m \in \{2, 3, 4\}$, $f \in \{7, 13, 21\}$, none $\equiv 3$. So Lemma 3.2 applies and $f \leq 19 + 1 = 20$; thus $f \in \{3, 20\}$. If $f = 20$, then $F = \ell \cup \{P\}$ with 20 distinct points, so $P \notin \ell$, and there are 20

fixed lines by Lemma 2.1(b). If $f = 3$ and the fixed points lay on a line $L$, $L$ would be fixed with $f_L = 3 \not\equiv 2$, contradicting Lemma 2.1(a). So they form a triangle; its sides are fixed, and there are exactly three fixed lines by (b). □

## 4 The homology configuration is impossible

**Theorem 4.1** (Norm-parity lemma)**.** *Let $\sigma$ have odd prime order $p$ in a plane of order $n$, with $r$ orbits of length $p$ on points. Let $q \neq p$ be a prime with $-1 \in \langle q \bmod p \rangle \leq (\mathbb{Z}/p)^*$. Then $r\, v_q(n)$ is even.*

*Proof.* Let $\zeta = e^{2\pi i/p}$. Choose representatives $x_1, \dots, x_r$ of the point orbits and $y_1, \dots, y_r$ of the line orbits of length $p$ (equal numbers by Lemma 2.1(b)), and set

$$A_{ab} = \sum_{k \in \mathbb{Z}_p} \zeta^k N[x_a, \sigma^k y_b] \in \mathbb{Z}[\zeta].$$

Put $u_a = \sum_k \zeta^k e_{\sigma^k x_a}$ and $w_b = \sum_k \zeta^k e_{\sigma^k y_b}$ (standard Hermitian form); they have disjoint supports, $\langle u_a, u_{a'} \rangle = p\delta_{aa'}$, $\langle w_b, w_{b'} \rangle = p\delta_{bb'}$. *Claim:* $Nw_b = \sum_a A_{ab} u_a$. At $x = \sigma^m x_a$: $(Nw_b)(x) = \sum_k \zeta^k N[x_a, \sigma^{k-m} y_b] = \zeta^m A_{ab} = A_{ab} u_a(x)$. At a fixed point $x$: $(Nw_b)(x) = N[x, y_b] \sum_k \zeta^k = 0$. Since $\sum_k \zeta^k = 0$, $Jw_{b'} = 0$, so by Q1 $\langle Nw_b, Nw_{b'} \rangle = \langle w_b, N^T N w_{b'} \rangle = n \langle w_b, w_{b'} \rangle = np\, \delta_{bb'}$. By the claim the left side is $p \sum_a A_{ab} \overline{A_{ab'}}$. Hence $A^* A = nI_r$ and $\det A \cdot \overline{\det A} = n^r$. Let $\mathfrak{p} \mid q$ be a prime of $\mathbb{Z}[\zeta]$. By Q3 its decomposition group $\langle \gamma_q \rangle$ contains $\gamma_{-1}$, so complex conjugation fixes $\mathfrak{p}$ and $v_\mathfrak{p}(\overline{\alpha}) = v_\mathfrak{p}(\alpha)$. As $q$ is unramified, $v_\mathfrak{p}(n) = v_q(n)$, and applying $v_\mathfrak{p}$ gives $2 v_\mathfrak{p}(\det A) = r\, v_q(n)$. □

**Corollary 4.2.** *There is no collineation of order 17 with $f = 20$ in a plane of order 18.*

*Proof.* Take $p = 17$, $q = 2$: $2^4 = 16 \equiv -1$, so 2 has order 8 and $-1 \in \langle 2 \rangle$; $v_2(18) = 1$. So $r$ is even, but $r = 19$ by Theorem 3.3. □

**Remark 4.3.** For $f = 3$, $r = 20$ is even and nothing follows; for $q = 3$, $v_3(18) = 2$ and the condition is vacuous. $\mathbb{Q}(\zeta_{17})$ has $16/8 = 2$ primes above 2, each fixed by complex conjugation.

## 5 The triangle case: reduction to a finite problem

Now $\sigma$ has exactly three fixed points $P_0, P_1, P_2$. Write $\{a, b, k\} = \{0, 1, 2\}$ and $L_k = P_a P_b$. Only $p = 17$ prime and $n = p + 1$ are used, so the derivation can be tested with other $p$ (Section 7). The letters $r_i, c_j, \tau_k$ below are gauge parameters.

**Lemma 5.1** (Orbit structure)**.** *A point is* general *if on no side; a line is* general *if not a side and containing no $P_k$.*

*(a) $S_k := L_k \setminus \{P_a, P_b\}$ is one orbit of length 17.*

*(b) The set $V_k$ of the 17 lines through $P_k$ other than the sides $L_a, L_b$ is one orbit. A line of $V_k$ meets $L_k$ in one point (of $S_k$), meets $L_a, L_b$ only in $P_k$, and its other 17 points are general.*

*(c) There are 289 general points, forming orbits $O_1, \dots, O_{17}$, and 289 general lines, forming orbits $H_1, \dots, H_{17}$.*

*(d) A general line meets each $L_k$ in one point of $S_k$, these three points are distinct, and it has 16 general points. A general point lies on exactly one line of each $V_0, V_1, V_2$ and on 16 general lines.*

*(e) Every line of $V_k$ contains exactly one point of each $O_i$.*

*Proof.* (a) $L_k$ has 19 points, two fixed; the other 17 are non-fixed, hence one orbit. (b) Of the 19 lines through $P_k$, two are sides; the other 17 are not fixed (the fixed lines are the sides), so form one orbit. Such a line meets $L_k$ outside $P_a, P_b$ (else it is a side $P_k P_a$ or $P_k P_b$), and meets $L_a, L_b$ only

in $P_k$; of its 19 points one is $P_k$, one is on $L_k$, and 17 are general. (c) $343-3-3\cdot 17=289$ general points, non-fixed, so 17 orbits. Lines containing a vertex or being a side number $3+3\cdot 17=54$ (a non-side contains at most one vertex), leaving 289 non-fixed lines. (d) A general line $\Lambda$ meets each side in a non-vertex point, so in $S_k$; two of these points coincide only at a vertex, so they are distinct and $\Lambda$ has $19-3=16$ general points. For a general point $Q$, the lines $QP_k$ are distinct non-sides, so $QP_k\in V_k$ (the three lines $QP_0, QP_1, QP_2$ are distinct, else $Q$ lies on a side), and the other 16 lines through $Q$ are general. (e) By (d), $\sum_{v\in V_k}|v\cap O_i|=17$; as $V_k$ is one orbit and $O_i$ is invariant, $|v\cap O_i|=c_i$ is constant, so $c_i=1$. □

### 5.1 Coordinates and incidence identities

Choose $v_k\in V_k$, $s_k:=v_k\cap L_k$, and put $V_{k,w}=\sigma^w v_k$, $(k,z)=\sigma^z s_k$ $(w,z\in\mathbb{Z}_{17})$, so $V_{k,w}\cap L_k=\{(k,w)\}$. Choose $Q_i\in O_i$, $\Lambda_j\in H_j$ $(i,j\in[17]:=\{1,\dots,17\})$, and put $(i,X)=\sigma^X Q_i$, $H_{j,y}=\sigma^y\Lambda_j$. Define

$$f_{ij}=\{t: Q_i\in\sigma^t\Lambda_j\},\quad s_j^k=\text{the } t \text{ with } s_k\in\sigma^t\Lambda_j,\quad u_i^k=\text{the } t \text{ with } Q_i\in\sigma^t v_k.$$

(Well defined: $\Lambda_j\cap L_k$ is one point of $S_k$ and $Q_i$ is on one line of $V_k$, by Lemma 5.1(d), and $\sigma$ acts regularly on $S_k$ and $V_k$.) Equivariance gives, for all $i,j,k$ and $X,y,z,w\in\mathbb{Z}_{17}$:

$$(i,X)\in H_{j,y}\iff y-X\in f_{ij}, \tag{I1}$$
$$(k,z)\in H_{j,y}\iff y-z=s_j^k, \tag{I2}$$
$$(i,X)\in V_{k,w}\iff w-X=u_i^k, \tag{I3}$$
$$(k,z)\in V_{k,w}\iff z=w. \tag{I4}$$

By Lemma 5.1(b), $V_{k,w}$ contains no point of $S_{k'}$ $(k'\neq k)$; general lines contain no vertex, and sides no general point.

**Proposition 5.2** (Incidence identities)**.** *(I) For each $i$, the multiset $\{t-t': j\in[17],\ t,t'\in f_{ij},\ t\neq t'\}$ equals $\mathbb{Z}_{17}^*$, each element once.*

*(II) For each $i,k$, the sets $\{t-s_j^k: t\in f_{ij}\}$ $(j\in[17])$ are pairwise disjoint with union $\mathbb{Z}_{17}\setminus\{u_i^k\}$.*

*(III) For each $j,k$, the sets $\{u_i^k-t: t\in f_{ij}\}$ $(i\in[17])$ are pairwise disjoint with union $\mathbb{Z}_{17}\setminus\{-s_j^k\}$.*

*(IV) For $a\neq b$, $j\mapsto s_j^a-s_j^b$ is a bijection $[17]\to\mathbb{Z}_{17}$.*

*(V) For $i\neq i'$ put $\delta_k=u_i^k-u_{i'}^k$. Then $\delta_0,\delta_1,\delta_2$ are distinct, and $\{y-y': j,\ y\in f_{ij},\ y'\in f_{i'j}\}$ consists of the 14 elements of $\mathbb{Z}_{17}\setminus\{\delta_0,\delta_1,\delta_2\}$, each once.*

*(VI) For $j\neq j'$, the elements $s_{j'}^k-s_j^k$ $(k=0,1,2)$ are distinct, and $\{t'-t: i,\ t\in f_{ij},\ t'\in f_{ij'}\}$ consists of the 14 elements outside them, each once. For $j=j'$, $\{t'-t: i,\ t\neq t'\in f_{ij}\}=\mathbb{Z}_{17}^*$, each once.*

*Proof.* Each item counts the unique line through two points (or the unique common point of two lines), and decides to which family that line belongs.

(I) For $d\neq 0$ the points $(i,0),(i,d)$ lie on one line $\lambda$, not a side, and not in $V_k$ (Lemma 5.1(e)); so $\lambda=H_{j,y}$ for a unique $(j,y)$, and by (I1) $y,\ y-d\in f_{ij}$. Conversely every such triple gives a line through both points.

(II) For $z\in\mathbb{Z}_{17}$, the line through $(i,0)$ and $(k,z)$ is no side and not in $V_{k'}$, $k'\neq k$. It lies in $V_k$ iff it is $V_{k,w}$ with $w=z$ (I4) and $w=u_i^k$ (I3), i.e. iff $z=u_i^k$. Otherwise it is $H_{j,y}$ with $y\in f_{ij}$ (I1) and $y-z=s_j^k$ (I2), i.e. $z=t-s_j^k$, $t=y\in f_{ij}$, for a unique $(j,t)$.

(III) $H_{j,0}$ and $V_{k,w}$ meet in one non-vertex point $Z$. If $Z$ is a side point it is $(k,w)$, and lies on $H_{j,0}$ iff $w=-s_j^k$ (I2). If $Z=(i,X)$ is general, (I3) gives $X=w-u_i^k$ and (I1) gives $-X\in f_{ij}$, i.e. $w=u_i^k-t$, $t=-X\in f_{ij}$.

(IV) $(a,z)$ and $(b,z')$ lie on one line, not a side and not in any $V_c$ (a line of $V_c$ has exactly one point on $S_0 \cup S_1 \cup S_2$). So it is $H_{j,y}$ with $y - z = s_j^a$, $y - z' = s_j^b$, i.e. $z' - z = s_j^a - s_j^b$, for a unique $j$.

(V) $(i,0)$ and $(i',X)$ lie on one line, not a side. It is in $V_k$ iff it is $V_{k,w}$ with $w = u_i^k$ and $w - X = u_{i'}^k$, i.e. $X = \delta_k$. If $\delta_k = \delta_{k'}$ for $k \neq k'$, two distinct lines would contain the same two points; so the $\delta_k$ are distinct. For $X \notin \{\delta_k\}$ the line is $H_{j,y}$ with $y \in f_{ij}$, $y - X \in f_{i'j}$, for a unique $(j,y)$.

(VI) For $(j',y) \neq (j,0)$, $H_{j,0}$ and $H_{j',y}$ meet in one non-vertex point. It is a side point $(k,z)$ iff $-z = s_j^k$ and $y - z = s_{j'}^k$ (I2), i.e. iff $y = s_{j'}^k - s_j^k$. It is general $(i,X)$ iff $t := -X \in f_{ij}$, $t' := y - X \in f_{ij'}$ (I1), i.e. $y = t' - t$. If $j = j'$, $y \neq 0$, a side point would need $y = 0$, so the point is general. If $j \neq j'$, the three $s_{j'}^k - s_j^k$ are distinct (else the lines share two side points), and the count follows. □

**Types of rows.** By (I), a block $f_{ij}$ of size $m$ contributes $\binom{m}{2}$ of the 8 classes $\{\pm d\}$, each exactly once over $j$. So $\sum_j \binom{|f_{ij}|}{2} = 8$, whence $|f_{ij}| \leq 4$, and $\sum_j |f_{ij}| = 16$ by (II). With $\nu_m = \#\{j : |f_{ij}| = m\}$ (not to be confused with the order $n$): $6\nu_4 + 3\nu_3 + \nu_2 = 8$, $4\nu_4 + 3\nu_3 + 2\nu_2 + \nu_1 = 16$, with solutions $(\nu_4,\nu_3,\nu_2,\nu_1) \in \{(0,0,8,0),(0,1,5,3),(0,2,2,6),(1,0,2,8)\}$ ($\nu_4 \leq 1$; $\nu_4 = 1$ forces $\nu_3 = 0, \nu_2 = 2$; if $\nu_4 = 0$, $\nu_3 \leq 2$).

### 5.2 Gauge, normal form, nets

Replacing $Q_i \mapsto \sigma^{r_i} Q_i$, $\Lambda_j \mapsto \sigma^{c_j}\Lambda_j$, $v_k \mapsto \sigma^{\tau_k} v_k$ changes the data to

$$f'_{ij} = f_{ij} + r_i - c_j, \qquad s'^k_j = s^k_j - c_j + \tau_k, \qquad u'^k_i = u^k_i + r_i - \tau_k$$

(e.g. $Q'_i \in \sigma^t \Lambda'_j \iff Q_i \in \sigma^{t+c_j-r_i}\Lambda_j$).

**Proposition 5.3** (Normal form)**.** *The choices can be made with $s_j^0 = 0$ and $u_i^0 = 0$ for all $i,j$. Then (a) for each $i$ the sets $f_{ij}$ ($j \in [17]$) partition $\mathbb{Z}_{17}^*$, and for each $j$ the sets $f_{ij}$ ($i \in [17]$) partition $\mathbb{Z}_{17}^*$; (b) $j \mapsto s_j^1$, $j \mapsto s_j^2$, $j \mapsto s_j^2 - s_j^1$ are bijections $[17] \to \mathbb{Z}_{17}$.*

*Proof.* Take $\tau_k = 0$, $c_j = s_j^0$, $r_i = -u_i^0$. (a) is Proposition 5.2(II),(III) for $k = 0$ (in (III) the sets $-f_{ij}$ partition $\mathbb{Z}_{17} \setminus \{0\}$). (b) is (IV) applied to the three pairs of sides, using $s^0 = 0$. □

The residual freedom is $c_j = r_i = \tau_0$ constant and arbitrary $\tau_1, \tau_2$: all $f_{ij}$ are unchanged and $s_j^k \mapsto s_j^k + \tau_k - \tau_0$ ($k = 1,2$).

**The net.** By (b) name the columns by $\alpha = s_j^1 \in \mathbb{Z}_{17}$, write $f_{i\alpha}$, and define $\theta(\alpha) = s_j^2$. Then $\theta$ and $\theta - \mathrm{id}$ are bijections: $\theta$ is an *orthomorphism* of $\mathbb{Z}_{17}$, and $\Theta = \{(s_j^1, s_j^2)\}$ is a transversal of the cyclic Latin square of order 17 (it meets each of $x = c$, $y = c$, $y - x = c$ once, as $\theta$ and $\theta - \mathrm{id}$ are bijections). Residual translations replace $\Theta$ by $\Theta + (a,b)$, so we may assume $\theta(0) = 0$; let $\mathrm{Orth}_0$ be the set of such $\theta$.

### 5.3 Rows and compatibility

For each row $i$ define $g_i : \mathbb{Z}_{17}^* \to \mathbb{Z}_{17}$ by $g_i(x) = \alpha$ iff $x \in f_{i\alpha}$ (Proposition 5.3(a)); the *cells* are $C_\alpha = g^{-1}(\alpha)$.

**Definition 5.4** (Rows)**.** $\mathrm{Rows}(\theta)$ is the set of $g : \mathbb{Z}_{17}^* \to \mathbb{Z}_{17}$ such that: (R1) the multiset of differences $x - x'$ ($x \neq x'$ in a common cell) is exactly $\mathbb{Z}_{17}^*$; (R2) $x \mapsto x - g(x)$ is injective; $u^1(g)$ denotes the missing element of $\mathbb{Z}_{17}$; (R3) $x \mapsto x - \theta(g(x))$ is injective; $u^2(g)$ denotes the missing element; (R4) for columns $\alpha \neq \beta$ the differences $y - x$ ($x \in C_\alpha, y \in C_\beta$) are pairwise distinct and differ from $\beta - \alpha$ and $\theta(\beta) - \theta(\alpha)$.

**Proposition 5.5** (Rows)**.** *In the normal form each $g_i \in \mathrm{Rows}(\theta)$, with $u^k(g_i) = u_i^k$ ($k = 1,2$).*

For a fixed net the rows are independent objects: $\mathrm{Rows}(\theta)$ depends on $\theta$ only, and the only coupling is pairwise compatibility.

*Proof.* (R1) is (I). (R2), (R3) are (II) for $k = 1, 2$ with $s_j^1 = \alpha_j$, $s_j^2 = \theta(\alpha_j)$. (R4): for columns $\alpha \neq \beta$ ($j \neq j'$), by (VI) the differences $t' - t$ over all rows are distinct and avoid $s_{j'}^k - s_j^k$, i.e. $\beta - \alpha$ and $\theta(\beta) - \theta(\alpha)$; restrict to row $i$. □

**Definition 5.6** (Compatibility)**.** Rows $g, g' \in \mathrm{Rows}(\theta)$ are *compatible* if, with $\delta_0 = 0$, $\delta_k = u^k(g) - u^k(g')$: (C1) $\delta_0, \delta_1, \delta_2$ are distinct and $\{y - y' : \alpha \in \mathbb{Z}_{17},\ y \in C_\alpha,\ y' \in C'_\alpha\}$ consists of the 14 elements of $\mathbb{Z}_{17} \setminus \{\delta_0, \delta_1, \delta_2\}$, each once; (C2) for columns $\alpha \neq \beta$ the sets $\{y - x : x \in C_\alpha, y \in C_\beta\}$ and $\{y - x : x \in C'_\alpha, y \in C'_\beta\}$ are disjoint.

**Proposition 5.7** (Pairs)**.** *In the normal form, $g_i, g_{i'}$ are compatible for $i \neq i'$.*

*Proof.* (C1) is (V) (with $\delta_0 = u_i^0 - u_{i'}^0 = 0$); (C2) is (VI), since all differences over all rows in a column pair are distinct. □

**Remark 5.8.** Compatible rows satisfy $\sum_\alpha |C_\alpha||C'_\alpha| = 14$ and $g(x) \neq g'(x)$ for all $x$ (since $0 \notin$ (C1)), and no two difference classes $\pm d$ occur once in a cell of $g$ and once in a cell of $g'$ of the same column. The program uses these as pre-filters; they add nothing logically.

**Lemma 5.9** (Bucket lemma)**.** *In the normal form, for each $x \in \mathbb{Z}_{17}^*$ the map $i \mapsto g_i(x)$ is a bijection $[17] \to \mathbb{Z}_{17}$. Hence with $B_\theta(x, c) = \{g \in \mathrm{Rows}(\theta) : g(x) = c\}$ (the* buckets*), if some $g_i \in B_\theta(1, 0)$, then for every $c \neq 0$ the bucket $B_\theta(1, c)$ contains a row compatible with $g_i$ (namely the $g_{i'}$ with $g_{i'}(1) = c$).*

*Proof.* By Proposition 5.3(a), for each column $\alpha$ the cells $f_{i\alpha}$ ($i \in [17]$) partition $\mathbb{Z}_{17}^*$, so $x$ lies in exactly one. □

### 5.4 Symmetry and classes

Let $G$ be the group generated by translations of $\mathbb{Z}_{17}^2$, the maps $(x, y) \mapsto (mx, my)$, $m \in \mathbb{Z}_{17}^*$, and $(x, y) \mapsto (y, x)$, $(x, y) \mapsto (-x, y - x)$ (these two generate $S_3$).

**Proposition 5.10** (Symmetry)**.** *If $(\Pi, \sigma)$ of triangle type yields the net $\Theta$ in normal form, then for every $\gamma \in G$ there are coordinates of the same type yielding $\gamma(\Theta)$.*

*Proof.* Translations: the residual freedom. Scalars: replace $\sigma$ by the generator $\sigma^m$; then $s_j^k$ becomes $m^{-1}s_j^k$. Relabel the vertices by a permutation $\pi$: $s_j^k \mapsto s_j^{\pi(k)}$, and renormalise by subtracting $s_j^{\pi(0)}$; the new net is $\{(s_j^{\pi(1)} - s_j^{\pi(0)},\ s_j^{\pi(2)} - s_j^{\pi(0)})\}$. In the coordinates $(x, y) = (s_j^1 - s_j^0, s_j^2 - s_j^0)$, swapping $1, 2$ is $(x, y) \mapsto (y, x)$ and swapping $0, 1$ is $(x, y) \mapsto (-x, y - x)$. □

Orthomorphisms $\theta, \theta' \in \mathrm{Orth}_0$ are *equivalent* if $\gamma(\mathrm{graph}\,\theta) = \mathrm{graph}\,\theta'$ for some $\gamma \in G$. A class consists of the normalised orthomorphisms obtained from $L(\mathrm{graph}\,\theta)$, for $L$ in the group of $16 \cdot 6 = 96$ linear maps, by translating one of its 17 points to the origin; so it has at most $17 \cdot 96 = 1632$ elements. Its *representative* is the lexicographically smallest $(\theta(1), \ldots, \theta(16))$.

**Theorem 5.11** (Reduction)**.** *If $\Pi$ has order 18 and a collineation $\sigma$ of order 17 with exactly 3 fixed points, there are a class representative $\theta$ and pairwise compatible rows $g_1, \ldots, g_{17} \in \mathrm{Rows}(\theta)$ such that $g_1(x), \ldots, g_{17}(x)$ are distinct for every $x$. In particular, for the row $g$ with $g(1) = 0$ and every $c \in \{1, \ldots, 16\}$, $B_\theta(1, c)$ contains a row compatible with $g$.*

*Proof.* Normal form (Proposition 5.3) with $\theta$ a representative (Proposition 5.10); rows by Proposition 5.5, compatibility by Proposition 5.7, distinctness and the last claim by Lemma 5.9. □

All conditions used are necessary; compatibility is weaker than the full set of incidence identities, which suffices since only non-existence is to be shown. The search therefore tests, for each class representative $\theta$,

$$\forall g \in B_\theta(1, 0)\ \ \exists c \in \{1, \ldots, 16\} :\ \text{no row of } B_\theta(1, c) \text{ is compatible with } g. \qquad (\star_\theta)$$

## 6 The computation

The 1,606,008,513 orthomorphisms of $\mathbb{Z}_{17}$, of which 94,471,089 are normalised, fall into 58,161 classes under $G$ (exhaustive enumeration with a canonicity test, `orth.cpp`); Table 1 gives the class sizes. The set $\mathcal{F}$ of partitions of $\mathbb{Z}_{17}^*$ satisfying (R1) ("families") has $490{,}801 = 3857 + 351360 + 119168 + 16416$ members, for the four types $(\nu_4, \nu_3, \nu_2, \nu_1)$ of Section 5, in the order listed there (obtained by enumeration; no closed formula is known; the count is used only to describe the search).

| class size | 3 | 6 | 102 | 204 | 408 | 816 | 1632 | total |
|---|---|---|---|---|---|---|---|---|
| number of classes | 1 | 2 | 3 | 6 | 27 | 486 | 57 636 | 58 161 |
| orthomorphisms | 3 | 12 | 306 | 1 224 | 11 016 | 396 576 | 94 061 952 | 94 471 089 |

Table 1: Classes of normalised orthomorphisms of $\mathbb{Z}_{17}$. The sizes $3, 6, 6$ account for the 15 linear orthomorphisms $\theta(x) = \lambda x$, $\lambda \neq 0, 1$, on which $S_3$ acts anharmonically (a consistency check).

**Algorithm 1** Test of $(\star_\theta)$ for one representative $\theta$

**Require:** $\theta \in \mathrm{Orth}_0$; the list $\mathcal{F}$ (same for all $\theta$)

1: generate lazily $B(c) = B_\theta(1, c)$, $c = 0, \ldots, 16$: place the blocks of each family in distinct columns with 1 in column $c$; prune by (R2),(R3); check (R4); record $u^1, u^2$
2: $\mathit{Cand} \leftarrow B(0)$
3: **for** $c = 1, \ldots, 16$ **while** $\mathit{Cand} \neq \emptyset$ **do**
4: $\quad \mathit{Cand} \leftarrow \{g \in \mathit{Cand} : \exists h \in B(c)$ compatible with $g\}$ ▷ necessary by Lemma 5.9
5: **end for**
6: **return** ELIMINATED if $\mathit{Cand} = \emptyset$; else run an exact 17-clique search with one row from each bucket containing a survivor

Compatibility is tested in the order: (a) $\sum_\alpha |C_\alpha||C'_\alpha| = 14$ (SIMD); (b) $g(x) \neq h(x)$ for all $x$; (c) no common same-column difference class; (d) (C1) by bitmasks; (e) (C2) by bitsets indexed by (column pair, difference). Tests (a)–(c) are implied by (d),(e) (Remark 5.8).

**Theorem 6.1** (Computational)**.** $(\star_\theta)$ *holds for each of the* 58,161 *class representatives. Indeed* $\mathit{Cand} = \emptyset$ *after the step* $c = 3$ *for every* $\theta$*; the exact search is never invoked (zero survivors,* 17*-cliques and nodes).*

Established by computer; data in `run_out.txt` (one line per net), summarised in Table 2.

| quantity | value |
|---|---|
| class representatives processed | 58 161 (ids 0–58160, once each) |
| $\lvert B(0)\rvert$, range (mean) | 23 696–27 776 (26 774) |
| nets with $\mathit{Cand} = \emptyset$ first at $c = 1$ / $c = 2$ / $c = 3$ | 0 / 57 538 / 623 |
| survivors after $c = 1$: total (max per net) | 971 489 (37) |
| survivors after $c = 2$: total (max per net) | 627 (2) |
| survivors after $c = 3$; 17-cliques; exact-search nodes | 0; 0; 0 |
| total CPU time | 179 471 s ($\approx$ 49.9 core-hours) |

Table 2: The run for $p = 17$, recomputed from `run_out.txt` by one `awk` line.

*Proof of Theorem 1.1.* By Theorem 3.3, $f \in \{3, 20\}$. Corollary 4.2 excludes 20. If $f = 3$, Theorem 5.11 gives a representative $\theta$ for which $(\star_\theta)$ fails, contradicting Theorem 6.1. □

# 7 Verification, assumptions and limitations

## 7.1 Controls

The reduction uses only that $\sigma$ has prime order $p$, $n = p+1$, and a fixed triangle. Replacing 17 by $p$ throughout ($(p-1)/2$ difference classes, $p-3$ for 14, etc.), the same program (`-DP=`$p$) applies to $n = p+1$.

| $p$ | $n$ | $\lvert\mathrm{Orth}_0\rvert$ | classes | with sol. | plane of order $n$ | output |
|---|---|---|---|---|---|---|
| 3 | 4 | 1† | 1 | 1 | exists | as expected (1 clique) |
| 5 | 6 | 3 | 1 | 0 | none | as expected |
| 7 | 8 | 19 | 3 | 2 | exists | as expected |
| 11 | 12 | 3 441 | 11 | 0 | open | no information |
| 13 | 14 | 79 259 | 117 | 0 | none | as expected |
| 31 | 32 | — | — | — | PG(2,32) | true plane found (below) |
| 17 | 18 | 94 471 089 | 58 161 | 0 | open | — |

Table 3: Controls. $p = 7$: 4 cliques on the 3 representatives; on all 133 transversal nets (no reduction) 35 nets carry 70 solutions, each verified to be a plane of order 8 (`plane.py`). $p = 5$: the family list has one member and no valid row exists. †`orth.cpp` needs $p \geq 5$; for $p = 3$ the unique $\theta(x) = 2x$ was entered by hand. Entries with output 0 are weak tests (a bug discarding too much also gives 0); the positive controls are $p = 3, 7$, and $p = 31$ only via ported predicates (see below).

**Why positive controls are in characteristic 2.** A genuine control needs a plane of order $n = p + 1$ with a collineation of prime order $p$ with a fixed triangle. All known planes have prime-power order $q^k$ with $q^k - 1$ prime: for odd $q$ this forces $n = 3$, $p = 2$ (excluded, $p$ must be odd); for $q = 2$, $p = 2^k - 1$ is a Mersenne prime, $n = 4, 8, 32, 128, \ldots$. We use the Desarguesian planes $\mathrm{PG}(2, 2^k)$ with $2^k - 1$ prime as controls; they have such a collineation (diag of order $2^k - 1$). Other known planes with such an automorphism may exist (e.g. non-Desarguesian planes of order 32) but were not used. A non-Desarguesian plane of order $q^2$, $q > 2$, has $n - 1 = (q - 1)(q + 1)$ composite. So the controls available are $p = 3, 7, 31, 127$; $p = 127$ was not attempted. The $p = 3, 7$ controls test that true solutions survive the necessary conditions, pruning and the bucket lemma, but not features specific to 17: the family list, blocks of size 4 (which need $p \geq 13$), or the code paths for 17 columns.

**Control $p = 31$ ($\mathrm{PG}(2, 32)$).** Elements $\sigma_c = \mathrm{diag}(1, g, g^c)$, $c = 2, \ldots, 30$ (for $c = 2$ alone all blocks have size $\leq 2$, as for $p = 7$; for $\sigma_c$ blocks are roots of a trinomial, with sizes $1, 2, 3, 5$; size 4 occurs only at $N = 63$, composite). From the explicit geometry of the 29 elements, the quotient data were computed, and the normal form, (R1)–(R3), the net conditions, the bucket argument and the compatibility predicates accepted all: 31/31 true rows, 465/465 true row pairs; about 5000 single-entry perturbations of true rows rejected; the exact search finds exactly one 31-clique, the true plane; the 1057-point structure rebuilt from the quotient data satisfies the axioms. These tests used hand-ported 64-bit predicates and generators (`ctrl_p.cpp`; regression on $\mathrm{PG}(2, 8), \mathrm{PG}(2, 16)$ passes). A parametrised copy of `fast4.cpp` reproduces the first 500 lines of the $P = 17$ run and the $P = 7$ results on all 133 nets, and at $P = 31$ returns exactly the true plane for $c = 3, \ldots, 30$ except 16, and 6 variants, with and without decoy (extra random false) families; at $P = 63$ blocks of size 4 were tested with injected rows (generator bypassed). *Not done:* the original `fast4.cpp` at $p = 31$ (its SIMD tables are fixed at 17); $c = 2$, $c = 16$ and the all-pairs row type (generator memory); the full $\mathrm{Rows}(\theta)$ at $p = 31$; the generator on families with 4-blocks; equality of the parametrised build with the $p = 17$ binary on all 58,161 nets. The control shows the quotient derivation is not contradicted by real planes; it says nothing about non-existence.

### 7.2 Independent checks

- **Independent re-implementations from the same derivation.** `indep.cpp` agrees with the main program byte for byte on the row sets $B(1,0), B(1,1), B(1,2)$ of a test net and on survivor counts. `mine.cpp` ($p = 3, 5, 7, 11, 13$) and `mine17b.cpp` ($p = 17$) agree on all controls and, for $p = 17$, on $|B(1,0)|, a_1, a_2, a_3$ for 40 random nets and 8 nets with $a_2 > 0$ ($a_i$: candidates left after filtering step $i$).
- **Reproduction from this paper alone.** A second implementation, written without the original code, reproduced the 490,801 families, the 94,471,089 orthomorphisms and the 58,161 classes with the same size histogram (its own canonical form), passed the controls $p = 3, 5, 7, 11, 13$ (these controls were run by the independent reproducer; I cannot confirm a re-run of the original implementation when preparing this text), and decided all 58,161 classes with 0 cliques (interrupted once at 8,926 classes, resumed from checkpoint). Verdicts agree; per-class statistics are not comparable.
- **Counting.** The class sizes sum to 94,471,089; `cnt.cpp` (depth-first) gives the same and $\times 17 =$ 1,606,008,513; 200 random orthomorphisms canonicalise into the list; `orbsum.cpp, orb.cpp` enumerate orbits explicitly; the class list regenerated from scratch has the same sorted content. The family count and its split were reproduced by `fam.cpp` and `indep.cpp`.
- **Oracle and SAT.** `oracle_pairs.py` tests compatibility of 257 pairs on a net by building the partial incidence structure from the axioms; it agrees with the C++ function. A direct CP-SAT model of the axioms with only the gauge conditions $s^0 = u^0 = 0$ confirms Proposition 5.3 for $p = 3, 7$ (infeasible when negated) and shows infeasibility for $p = 5$.
- **Re-run.** For this paper 2,760 nets of the $p = 17$ run (ids 0–759, 30000–31999) were re-run with output identical to `run_out.txt` except run time. The full $p = 17$ search was not repeated.

### 7.3 Assumptions

Theorem 1.1 rests on:

**A1.** the quoted facts Q1–Q3 (Bruck's bound is the only non-elementary plane-geometric input);

**A2.** the correctness of the hand derivation of Section 5 (re-derived by a second, non-blind reader and consistent with the CP-SAT model for $p = 3, 7$; no human referee);

**A3.** the correctness of the enumeration of class representatives and of Proposition 5.10;

**A4.** the correctness of the search with respect to Definitions 5.4 and 5.6: its row generator produces all of $B_\theta(1, c)$, and its compatibility test implies (C1),(C2). Only necessary conditions are tested, so a test that is too weak cannot lose a solution, whereas one that is too strong could;

**A5.** correct execution on the hardware used.

### 7.4 Limitations

This is a computer-assisted result with no human referee. The row generator (forward checking, trie of block structures, SIMD) is checked against a naive generator and independent implementations on some nets but is not formally verified; a proof-producing check (an independent simple program recomputing $B(1, c)$, or a certificate for "no compatible row") would be valuable. All implementations, including the reproduction from the paper, share the derivation of Section 5: they are independent in implementation, not in logic, and the logic was re-derived only by a second non-blind reader. Proposition 5.10 was also checked numerically ($p = 7, 11$: row counts and solution counts are constant on classes). At $p = 31$ the production code was ported and run, but the full row set was not enumerated, $c = 2, 16$ and `a28_b25` were not run, and the 4-block generator was not tested; this control is partial, and a derivation-independent check is open. Novelty is not verified. The result concerns only the prime 17; the primes $2, 3, 5, 7, 19$ remain open for order 18, and existence of a plane of order 18 is untouched.

**AI assistance.** Most derivations, programs and verification were produced by AI agents (large language models, Claude by Anthropic) under the author's direction; the author set the problem, directed the research and judged the results. The argument does not rely on their authority: every step of Sections 3–4 and 5 is written out for human checking, and the code and data are held by the author (Section A). The work has not been independently re-checked by a human specialist; a human re-derivation of Section 5 and a run on other hardware are the most useful checks.

# A Reproduction

The programs are in the directory `code/` of the repository (`fast4.cpp`, `orth.cpp`, and the 133 nets of the $p = 7$ control, `nets7_all.txt`; MIT licence). The output of the original run is not distributed: the computation is deterministic, so a new run reproduces it (the time column excepted). Requirements: AVX2, OpenMP; `-march=native` makes the build machine dependent. Compiler: MinGW-w64 GCC 16.1.0 (WinLibs); the compiler of the original run was not recorded and is assumed the same.

```
g++ -O3 -march=native -fopenmp orth.cpp -o orth
./orth classes classes17.txt                          # 58161 class representatives
g++ -O3 -march=native -fopenmp -DP=17 fast4.cpp -o netrun17.exe
OMP_NUM_THREADS=12 ./netrun17.exe classes17.txt 0 58161 run_out.txt run
awk '{a2+=$7;s+=$13;c+=$17} END{print NR,a2,s,c}' run_out.txt   # 58161 627 0 0
```

Interrupted runs resume by re-running the command (ids already in the output are skipped). The class list is written by `./orth classes <file>` (about 10 s; the order of the lines depends on thread scheduling, the set does not, so compare sorted files). Cost: about 3 s CPU per net on average (maximum 6.75 s), about 4 h wall clock on 12 threads; memory a few hundred MB per thread.

**Interrupted run.** The original run was started as `fastrun.exe`; after 727 nets the executable vanished from disk and the process ended (cause unknown; anti-virus suspected). The same source, compiled as `netrun17.exe` (byte-identical to `fast4.cpp`), was restarted by `watchdog.sh` (log: `done=727`) and finished in about 4 h. The output has 58,161 lines, ids $0, \ldots, 58160$ once each; the first 727 come from the first executable, whose exact command line (notes: `-O3 -march=native -DP=17`) was not preserved. These 727 nets were recomputed with identical results.

**Checksums (SHA-256).**

| | |
|---|---|
| `fast4.cpp` | `eb3f3ca800d6cee24255b14f25e543c7a30495051ed2f62e650ac112b37713bb` |
| `classes17.txt` | `aacf1791e82d5ad9b69a17ec32ef0d4c803a9a58e75585554bcbdcc72e8d7bbc` |
| `run_out.txt` | `86cff376cd5a3468aac2403618a6a046b7d260670ee347e85510b3f9f8598f1b` |
| `orth.cpp` | `74665a45165fc4c430778c9ade9b72209c521a073be2e8a9140c4792ab0af7a5` |
| `orbsum.cpp` | `3a96c58b848891994825974e3ca895e893dfb3484784a57164fd472ff24189ab` |

The files `classes17.txt` and `run_out.txt` of the author's run are not distributed; their checksums are listed for the record (the time column of `run_out.txt` changes from run to run). The file `orth.cpp` was edited after the first checksum was taken (output handling only); the hash above is that of the archived file. The sorted regenerated class list has SHA-256 `c3e3937b...e111e`, equal to that of the archived list sorted likewise.

# References


[1] E. F. Assmus, Jr. and J. D. Key. *Designs and Their Codes.* Cambridge University Press, Cambridge, 1992.

[2] Reinhold Baer. Projectivities with fixed points on every line of the plane. *Bull. Amer. Math. Soc.*, 52:273–286, 1946.

[3] R. H. Bruck. Difference sets in a finite group. *Trans. Amer. Math. Soc.*, 78:464–481, 1955.

[4] R. H. Bruck and H. J. Ryser. The nonexistence of certain finite projective planes. *Canad. J. Math.*, 1:88–93, 1949.

[5] Z. Janko and Tran Van Trung. Projective planes of order 10 do not have a collineation of order 3. *J. Reine Angew. Math.*, 325:189–209, 1981.

[6] C. W. H. Lam. The search for a finite projective plane of order 10. *Amer. Math. Monthly*, 98:305–318, 1991.

[7] C. W. H. Lam, L. Thiel, and S. Swiercz. The non-existence of finite projective planes of order 10. *Canad. J. Math.*, 41:1117–1123, 1989.

[8] F. J. MacWilliams, N. J. A. Sloane, and J. G. Thompson. On the existence of a projective plane of order 10. *J. Combinatorial Theory Ser. A*, 14:66–78, 1973.

[9] Xander Perrott. Existence of projective planes, 2016. Student report, arXiv:1603.05333 [math.HO].