

# STAT680\_HW1

Yudi Zhang, Wangqian Ju

9/24/2021

## Q1

a)

We have  $g(X) = X^{\alpha-1}$  and  $X \sim \text{Exp}(1)$

```
n <- 10000
u <- runif(n)

# obtain the Exp(1) random variable
ee <- -log(1-u)

alpha <- 1.89

e.gamma <- ee^(alpha - 1) %>% mean

# check the result using gamma function
c(e.gamma, gamma(alpha))

## [1] 0.9480089 0.9583793
```

b)

$$\begin{aligned} \text{Var}\left(\frac{1}{n} \sum X_i^{\alpha-1}\right) &= \frac{1}{n} \text{Var}(X^{\alpha-1}) \\ &= \frac{1}{n} (E[X^{2(\alpha-1)}] - [E(X^{\alpha-1})]^2) \\ &= \frac{1}{n} (\Gamma(2\alpha - 1) - [\Gamma(\alpha)]^2) \end{aligned}$$

c)

```
estimate_gamma <- function(alpha, n = 1000) {
  u <- runif(n)
  ee <- -log(1-u)

  tt <- ee^(alpha - 1)
  est <- mean(tt)
  sd <- sqrt(var(tt) / n)

  result <- c(est, sd)
  names(result) <- c("estimate", "sd")
}
```

```

    return(result)
}

```

For  $\alpha = 1.5$ ,  $\Gamma(\alpha) = 0.8862269$ . And for different simulation sizes, we have:

```

##      n estimate      sd
## 1    10 0.9443317 0.203278476
## 2   100 0.8065015 0.046315510
## 3  1000 0.8900198 0.014497273
## 4 10000 0.8886879 0.004659166

```

d)

```

estimate_gamma_antithetic <- function(alpha, n = 1000) {
  u <- runif(n)
  u2 <- 1 - u
  u <- c(u, u2)
  ee <- -log(1-u)

  tt <- ee^(alpha - 1)
  est <- mean(tt)
  sd <- sqrt(var(tt) / (2*n))

  result <- c(est, sd)
  names(result) <- c("estimate", "sd")
  return(result)
}

```

We can see that the method of antithetic variates can reduce the standard error.

```
ttd.df
```

```

##      estimate      sd      type
## 1 0.8860105 0.004628369      MC
## 2 0.8873709 0.003308856 Antithetic

```

e)

```

tr_val <- 1 - exp(-1)
n <- 100000

### i)
u <- runif(n)
tt <- as.numeric(u < exp(-(1:n) / n))
mean(tt)

```

```

## [1] 0.63459
sqrt(var(tt) / n)

```

```

## [1] 0.001522786

### ii)
u <- runif(n)
tt <- exp(-u)
mean(tt)

```

```
## [1] 0.633433
sqrt(var(tt) / n)

## [1] 0.00057364
### iii)
u <- runif(n)
u2 <- 1 - u
u <- c(u, u2)
tt <- exp(-u)
mean(tt)

## [1] 0.6321267
sqrt(var(tt) / (2*n))

## [1] 0.0004047294
### iv)
is_beta <- function(a=2, b=2, n=1000) {
  bb <- rbeta(n, a, b)
  tt <- exp(-bb) / dbeta(bb, a, b)
  est <- mean(tt)
  sd <- sqrt(var(tt) / n)
  result <- c(est, sd)
  names(result) <- c("estimate", "sd")
  return(result)
}

grid <- readRDS("grid.rds")

grid

## # A tibble: 625 x 5
##       a      b estimate      sd      diff
##   <dbl> <dbl>   <dbl>   <dbl>   <dbl>
## 1  0.2  0.2    0.635 0.00226 0.00331
## 2  0.4  0.2    0.631 0.00208 0.00112
## 3  0.6  0.2    0.630 0.00231 0.00215
## 4  0.8  0.2    0.629 0.00273 0.00275
## 5  1    0.2    0.633 0.00334 0.000962
## 6  1.2  0.2    0.631 0.00418 0.00106
## 7  1.4  0.2    0.640 0.00563 0.00823
## 8  1.6  0.2    0.643 0.00802 0.0107
## 9  1.8  0.2    0.627 0.00888 0.00545
## 10 2    0.2    0.621 0.0129 0.0111
## # ... with 615 more rows

# grid <- expand.grid(a=seq(0.2,5,by=.2), b=seq(0.2,5,by=.2))
#
# grid %<>% as_tibble %>% mutate(
#   is_beta_result = purrr::map2(a, b, function(aa, bb) {
#     rr <- is_beta(aa, bb, n = 100000) %>% t %>% as_tibble
#     colnames(rr) <- c("estimate", "sd")
#     rr
#   })
# }
```

```

# ) %>% unnest(is_beta_result) %>%
#   mutate(
#     diff = abs(tr_val - estimate)
#   )
#
# grid$sd %>% which.min() %>% {grid[.,]}
# grid$diff %>% which.min() %>% {grid[.,]}
# grid %>% filter(a==1, b==1)
# is_beta(1, 1, n = 100000)
#
# saveRDS(grid, "homework1/grid.rds")

```

## Q2

a)

```

library(boot)
require(fitdistrplus)

## Loading required package: fitdistrplus
## Warning: package 'fitdistrplus' was built under R version 4.0.2
## Loading required package: MASS
##
## Attaching package: 'MASS'
## The following object is masked from 'package:dplyr':
##
##   select
## Loading required package: survival
##
## Attaching package: 'survival'
## The following object is masked from 'package:boot':
##
##   aml
data("aircondit")


fit.gamma <- fitdist(aircondit$hours, distr = "gamma", method = "mle")
summary(fit.gamma)

## Fitting of the distribution ' gamma ' by maximum likelihood
## Parameters :
##           estimate Std. Error
## shape 0.706402862  0.2419358
## rate  0.006535456  0.0030685
## Loglikelihood: -67.64542   AIC:  139.2908   BIC:  140.2607
## Correlation matrix:
##           shape      rate
## shape 1.000000 0.700236
## rate  0.700236 1.000000


```

```
plot(fit.gamma)
```


**Empirical and theoretical dens.**




**Q-Q plot**



**Empirical and theoretical CDFs**



**P-P plot**



```
tb.find <- function(alpha, data) {
  xbar <- mean(data)
  result <- log(alpha) - log(xbar) - digamma(alpha) + mean(log(data))
  return(result)
}

alpha.est.result <- uniroot(tb.find, interval = c(0.001, 4), data = aircondit$hours)
alpha.est.result
```

```
## $root
## [1] 0.7064924
##
## $f.root
## [1] 1.100713e-06
##
## $iter
## [1] 10
##
## $init.it
## [1] NA
##
## $estim.prec
## [1] 6.103516e-05
```

```
alpha.est <- alpha.est.result$root
beta.est <- alpha.est / mean(aircondit$hours)
c(alpha.est, beta.est)
```

```
## [1] 0.706492376 0.006536552
# https://www.math.arizona.edu/~jwatkins/03\_mle.pdf

## b)
jk.est <- sapply(seq_along(aircondit$hours), function(i) {
  data <- aircondit$hours
  data.tmp <- data[-i]

  est <- uniroot(tb.find, interval = c(0.001, 4), data = data.tmp)
  est$root
})
n <- length(aircondit$hours)
eiv <- (n-1) * (alpha.est - jk.est)
bias.alpha <- - mean(eiv)

# var(jk.est) * (n-1)^2 / n

(sum(eiv^2) - n * bias.alpha^2) / (n * (n-1))

## [1] 0.06605922
```

### Q3.

(d) infinitesimal jackknife, regular jackknife and non-parametric bootstrap

i.

```
corr = cor(cd4$baseline, cd4$oneyear)
infjack_var <- function(x, y) {
  mux = mean(x)
  muy = mean(y)
  sx = sd(x)
  sy = sd(y)
  zx = (x - mux)/sx
  zy = (y - muy)/sy

  ifunc = zx * zy - 1/2 * cor(x, y) * (zx^2 + zy^2)
  return(sum(ifunc^2) * 1/(length(ifunc)^2))
}
infjack_var(cd4$baseline, cd4$oneyear)

## [1] 0.005701791
```

ii.

```
jackknife <- function(x, y) {
  est = c()
  for (i in 1:length(x)) {
    est = c(est, cor(x[-i], y[-i]))
  }
  corr = cor(x, y)
  bias = (length(x) - 1) * (mean(est) - corr)
  n = length(x)
```

```

tmp = (n - 1) *(corr - est)
variance = (sum(tmp^2) - n * bias^2)/((n - 1) * n)
return(c(bias, variance))
}
jackknife(cd4$baseline, cd4$oneyear)

```

```
## [1] -0.006784288  0.008187035
```

iii.

```

boot <- function(x, y, b = 50) {
  n = length(x)
  est = c()
  for(i in 1:b) {
    s = sample(c(1:n), n, replace = TRUE)
    est = c(est, cor(x[s], y[s]))
  }
  corr = cor(x, y)
  bias = mean(est) - corr
  variance = var(est - corr)
  return(c(bias, variance))
}
boot(cd4$baseline, cd4$oneyear)

```

```
## [1] -0.011534387  0.007191871
```

(e).

percentile interval

```

bootci <- function(x, y, b = 1000) {
  n = length(x)
  est = c()
  for(i in 1:b) {
    s = sample(c(1:n), n, replace = TRUE)
    est = c(est, cor(x[s], y[s]))
  }
  return(quantile(est, probs = c(0.025, 0.975)))
}
bootci(cd4$baseline, cd4$oneyear)

```

```
##      2.5%      97.5%
## 0.4969143 0.8646504
```

normal approximation

```

bootci_norm <- function(x, y) {
  n = length(x)
  corr = cor(x, y)
  l = (1 + corr - (1 - corr) * exp(2*1.96/sqrt(n-3)))/(1 + corr + (1 - corr)*exp(2*1.96/sqrt(n-3)))
  r = (1 + corr - (1 - corr) * exp(-2*1.96/sqrt(n-3)))/(1 + corr + (1 - corr)*exp(-2*1.96/sqrt(n-3)))
  return(c(l, r))
}

```

```
bootci_norm(cd4$baseline, cd4$oneyear)
```

```
## [1] 0.4127149 0.8830870
```

#### invariance to transformation

```
fishertans <- function(x, y) {  
  n = length(x)  
  corr = cor(x, y)  
  
  z = 0.5 * log((1 + corr)/(1-corr))  
  sd = sqrt(1/(n-3))  
  l = z + qnorm(0.975) * sd  
  r = z - qnorm(0.975) * sd  
  up = (exp(2 * l) - 1)/(exp(2*l) + 1)  
  lo = (exp(2 * r) - 1)/(exp(2*r) + 1)  
  return(c(lo, up))  
}  
fishertans(cd4$baseline, cd4$oneyear)
```

```
## [1] 0.4127222 0.8830851
```

#### Q4. (b).

##### CI Interval example:

bootstrap normal: 0.62 2.78

bootstrap basic: 0.29 1.82

bootstrap studentized: 1.37 3.41

#### Based on 1600 independent data sets:

##### Coverage:

bootstrap normal: 0.99875

bootstrap basic: 0.935625

bootstrap studentized: 0.94375

##### Mean and sd of length:

bootstrap normal: 2.3482 +/- 0.977

bootstrap basic: 2.51 +/- 0.71

bootstrap studentized: 2.61 +/- 1.3

The studentized bootstrap method was closest to 95% and bootstrap normal tends to give far large interval.

#### Below is the implementation

##### 1. normal

```
normalci <- function(n = 15, theta = 2, r = 999, a = 0.025) {  
  obs = rexp(n, theta)
```



```

xbar = rep(0, r)
for(i in 1:r) {
  x=sample(obs, n, replace=T)
  xbar[i] = 1/mean(x)
}
deltastar = xbar - theta
d = quantile(deltastar, c(a,1-a))
ci = mean(xbar) - c(d[2], d[1])
return(ci)
}
normalci()

n = 15
theta = 2
r = 999
a = 0.025
sims <- 1600
results <- as.numeric(sims)
len = rep(0, sims)
for (l in 1:1600) {
  pop <- rexp(100, theta)
  mu <- 1/mean(pop)
  obs = sample(pop, n)
  xbar = rep(0, r)
  for(i in 1:r) {
    x=sample(obs, n, replace=T)
    xbar[i] = 1/mean(x)
  }
  deltaxstar = xbar - theta
  d = quantile(deltastar, c(a,1-a))
  ci = mean(xbar) - c(d[2], d[1])
  len[l] = ci[2] - ci[1]
  results[l] <- ci[1] < mu & ci[2] > mu
}
sum(results)/sims

```

## 2. basic

```

basicci <- function(n = 15, theta = 2, r = 999, a = 0.025) {
  obs = rexp(n, theta)
  est = 1/mean(obs)
  xbar = rep(0, r)
  for(i in 1:r) {
    x = rexp(n, est)
    xbar[i] = 1/mean(x)
  }
  xbar = sort(xbar)
  l = (r + 1) * (1 - a)
  u = (r + 1) * (a)
  return(c(2*est - xbar[l], 2*est - xbar[u]))
}
basicci()

```

```

results <- 0
lens = rep(0, sims)
for (j in 1:1600) {
  pop <- rexp(100, theta)
  mu <- 1/mean(pop)
  obs = sample(pop, n)
  est = 1/mean(obs)
  xbar = rep(0, r)
  for(i in 1:r) {
    x = rexp(n, est)
    xbar[i] = 1/mean(x)
  }
  xbar = sort(xbar)
  l = (r + 1) * (1 - a)
  u = (r + 1) * (a)
  ci = c(2*est - xbar[l], 2*est - xbar[u])
  lens[j] = xbar[l] - xbar[u]
  if(ci[1] < mu & ci[2] > mu)
    results = results + 1
}
results/sims

```

### 3. studentized (found 2 ways to implement this)

```

stuci <- function(n = 15, theta = 2, r = 999, a = 0.025, M = 50) {
  obs = rexp(n, theta)
  est = 1/mean(obs)
  zr = rep(0, r)
  xbar = rep(0, r)
  for(i in 1:r) {
    x=sample(obs, n, replace=T)
    xbar[i] = 1/mean(x)
    thetahat = rep(0, M)
    for (j in 1:M) {
      sx=sample(x, n, replace=T)
      thetahat[j] = 1/mean(sx)
    }
    vr = sd(thetahat)
    zr[i] = (xbar[i] - est)/vr
  }
  v = sd(xbar)
  zr = sort(zr)
  d = quantile(zr, c(a,1-a))
  ci = est - c(d[2], d[1])
  return(ci)
}
stuci()

```

```

M = 10
results <- 0
for (l in 1:1600) {
  pop <- rexp(100, theta)
  mu <- 1/mean(pop)
  obs = sample(pop, n)

```

```

est = 1/mean(obs)
zr = rep(0, r)
xbar = rep(0, r)
for(i in 1:r) {
  x=sample(obs, n, replace=T)
  xbar[i] = 1/mean(x)
  thetahat = rep(0, M)
  for (j in 1:M) {
    sx=sample(x, n, replace=T)
    thetahat[j] = 1/mean(sx)
  }
  vr = sd(thetahat)
  zr[i] = (xbar[i] - est)/vr
}
v = sd(xbar)
zr = sort(zr)
d = quantile(zr, c(a,1-a))
ci = est - c(d[2], d[1])
if(ci[1] < mu & ci[2] > mu)
  results = results + 1
}
sum(results)/sims

```

```

## or use (another implementation)
pop <- rexp(1000, rate = 1/theta)
x <- sample(pop, n)
mu <- mean(pop)
B <- 1000
samples <- matrix(sample(x, size = n * B, replace = T), nrow = n, ncol = B)
est <- (apply(samples, 2, mean) - mean(x))/(apply(samples, 2, sd)/sqrt(n))
mean(x) - quantile(est, probs = c(1-a, a)) * sd(x)/sqrt(n)

```

```

ci <- matrix(FALSE, sims, 2)
results <- as.numeric(sims)
lens = rep(0, sims)
for (i in 1:sims) {
  pop <- rexp(1000, rate = 1/theta)
  x <- sample(pop, n)
  mu <- mean(pop)
  B <- 1000
  samples <- matrix(sample(x, size = n * B, replace = T), nrow = n, ncol = B)
  est <- (apply(samples, 2, mean) - mean(x))/(apply(samples, 2, sd)/sqrt(n))
  ci[i, ] <- mean(x) - quantile(est, probs = c(1-a, a)) * sd(x)/sqrt(n)
  lens[i] = ci[i, 2] - ci[i, 1]
  results[i] <- ci[i, 1] < mu & ci[i, 2] > mu
}
sum(results)/sims

```