Preface

On April 24, 2026, while pondering why electric fields generate magnetic fields, an initial insight came to me: could the fundamental ontology of this world be rooted in information?

Once this idea took shape, my thinking and reasoning would not cease. As others have described it, I fell into a state of near-obsessive reasoning. By August 2026, my mind had largely returned to calm. During this period I produced more than sixty documents, most of which have been archived on Zenodo. Interested readers may follow the chronological sequence of my Zenodo publications to witness the full evolutionary process through which this theory was iterated and refined.

This book, **Status-Relational-Entropy Dynamics (SRE-Dynamics)**, is a complete physical-theory system taking information as its ontology. It is compiled from my previously published manuscripts, selecting those with solid theoretical foundations and relatively objective validation data.

The book is divided into five major parts:

- Part I: Fundamental Axioms
- Part II: Emergence of All Things
- Part III: Mathematical-Physical Support
- Part IV: Applications and Hypotheses
- Part V: Supplementary Programs and Datasets

Resource and Availability Statement

This framework is built upon Status-Relational Entropy (SRE) Dynamics. All theoretical materials are archived in the Zenodo open-access repository, sorted by public-release timestamp, fully preserving the evolutionary history of this theory from its earliest conceptions and heuristic sketches up to logical closure. Versions v1.5.x and earlier in the archive are historical heuristic drafts, intended solely for version traceability and not recommended as formal citation sources. The formal citation baseline is the **SRE-v1.6 Axiom Suite** (https://doi.org/10.5281/zenodo.22077475).

All related materials for this book, together with the reproduction programs and datasets accompanying each chapter, are available in my GitHub repository:

https://github.com/yuelucn/Status-Relational-Entropy-SRE-Dynamics-The-Book-of-the-Void

Open-source scope of the operator suite: the complete mathematical specifications, algebraic derivations and simulation code for all Operators 1-19 are fully open-source, and constitute the main body of Parts III and V of this book. The entire operator pipeline is released under an open-source licence for reproduction, audit and secondary development.

In addition, an AI-assisted Tencent intelligent-document space is available (accessible on both PC and WeChat mobile):

https://docs.qq.com/space/DUkRjYUtNWFdyV253

As of 2026-08-14, constrained by Google's terms-of-service, the author no longer maintains or updates the SRE document library hosted in Google Gemini Notebook; that link serves only as a historical archive and must not be used as a formal citation source:

https://notebooklm.google.com/notebook/ef52bf5a-f6d0-4a2a-aed4-b25d6520ab2c

According to the deductions of this theory, the universe and everything within it - including time, space, light, matter, and all kinds of force-fields - emerge spontaneously from the ordered iterative growth of a binary self-organising network (a dimension-free status-relation graph).

Part I *Fundamental Axioms* defines the interchange relationship between information-ontology and classical-physical ontology. The two ontologies can be combined and mutually transformed in engineering practice.

Part II *Emergence of All Things* presents falsifiable inferences for the emergence processes of the material world.

Part III *Mathematical-Physical Support* spans multiple sub-fields of mathematics and physics and has a relatively high reading threshold. Its core idea is to build a complete operator system based on binary self-organising networks. This yields efficient and concise graph algorithms supporting local parallel graph computation for very-large-scale graph networks. Beyond supporting the present dynamical framework, this operator suite can also be applied independently to other engineering scenarios.

Within this part, the complete suite of Operators 1-19 is released under an open-source licence to provide full mathematical-physical support for the ontological theory, including the two electronic-degrees-of-freedom extension operators 11 and 12 introduced in September 2026 and the spectral-equation-solving and charge-symbol-channel operators 13-19. The entire pipeline is open-sourced for reproduction, audit and secondary development.

Part IV is titled *Applications and Hypotheses*. This naming arises because topics such as signal processing, AI-models, graph-computing and brain science are independent of the world-information-ontology premise of this theory and can directly deliver commercial-value-oriented implementations. By contrast, subjects including instantaneous bidirectional optical communication, inclusive-benefit medical care, next-generation semiconductor design, anti-gravity and life-science remain purely hypothetical at the present stage.

I have recorded all of this here. Scientific intuition arises out of the void, and returns to the void.

Yue Lu
August 28, 2026

Table of content

- Part I: Axiomatic Foundations — Abstract
- Status-Relational Entropy (SRE) Dynamics: Usage Specification and Theoretical Context
- A Foundational Framework for Information-Physical Interconversion
- Reciprocal Measurement and the Mapping Origin of c
- Methodological Revision on Causal Depth and Spatial Indexing
- Mapping Prerequisites
- A Unified Theory of Electronic Logical-Structural and Physical Properties Based on the Minimal Instantiable Topological Scale
- SRE Dynamics Supplementary Note: Magnitude Anchoring of Logical Depth N and Observational-Protocol Safety Masking
- Part II: Emergence of All-Things — Abstract
- Theory of Hierarchical Dissipative Self-Organizing Binary Network Dynamics
- Emergence of Multidimensional Spacetime and Dynamical Gravity via Regularized Causal-Information Networks
- A SRE-Dynamics Inspired Topological Paradigm for Composite Elementary Particles and Relational Space Emergence
- A Foundational Reconstruction of Classical Electrodynamics via Discrete Graph Topology and Bidirectional Causality
- Rigorous Reconstruction of Maxwell's Field Equations via Purely Dimensionless Graph Cohomology and Global Evolution Step
- The Necessity of Turbulence Emergence and an Algebraic Computational Method Based on Discrete Microscopic Causal Statistics and Multidimensional Manifold Reconstruction
- Oseen Vortex Initial-Value Experiment: Repositioning the Empirical Report (v2)
- Emergence of Classical Mechanics from Discrete Causal-Information Networks: Ontological Mapping and Effective-Theory Limits within Status-Relational-Entropy (SRE) Dynamics
- A Complete Characterization of the Electron within the SRE Framework: A Unified Account of Axiomatization, the Fine-Structure Constant, Molecular Computation, and Orbital Assignment
- Complete Characterization of Nucleons within the SRE Framework: Unified Representation of Two-State Skeleton Inversion, Quark Speculation and Nuclear Reaction Validation
- Mass Origin in the State–Relational–Entropy Framework
- Positioning of Light and Electromagnetic Waves in the SRE Framework
- Emergence of Time in the State–Relational–Entropy Framework
- How Does Time "Grow" Itself?
- Emergence of Positive and Negative Electric Charge in the SRE Framework: Magnitude, Sign, and Numerical Verification
- Part III: Mathematical-Technical Support — Abstract
- Universal Graph Operator Pipeline in Status-Relational Entropy (SRE) Dynamics: A Unified Foundational Framework White Paper
- Operator-1: Pure-Algebraic Mathematical Specification for the Local Graph Expansion Operator $(\mathcal{G}_{n\to n+1})$
- Operator-1: Local Graph Expansion Operator - Full Explanatory Document
- Operator-2： Local Metric & Probabilistic Pruning Operator $(\mathcal{M}_{\chi}\circ\mathcal{E}_{\text{local}})$
- Operator-3: Rigorous Mathematical-Derivation Specification
- Operator-4: Algebraic Construction of Local-Topology Degree-Statistic Operator $(\mathcal{M}_{\text{degree}})$ and Rigorous Positive-Definite Boundedness Proof for Dirichlet Energy Functional
- Operator-5: Endogenous Variable Latency Calibration Operator $(\mathcal{M}_{\text{latency}})$
- Operator-6: Sub-space Spectral Sieve & Splicing Operator $(\mathcal{P}_{\text{sieve}}\cup\mathcal{O}_{\text{splice}})$
- Mathematical Specifications, Algebraic Derivations, and Proof of Hodge Closure Completeness for the Adjoint Filter Locking & Complex Dual Equalizer Operator (Operator 7)
- Mathematical Specifications, Algebraic Derivations, and Proof of Monotone Convergence for the Local Lock-Free Algebraic Valve Balancing Operator (Operator 8)

- Mathematical Specifications, Algebraic Derivations, and Proof of Topological Invariant Anchoring for the Dual-Order Smooth Betti Number Synchronous Stitching Operator (Operator 9)
- Mathematical Specifications, Algebraic Derivations, and Proof of Connectivity Preservation for the Pre-emptive Homological Random Pruning Operator (Operator 10 / Operator 2-Batch)
- Mathematical Specifications and Algebraic Derivations for the Spin-Charge Adjacency Operator (Operator 11)
- Mathematical Specifications and Algebraic Derivations for the Topological Closed-Loop Audit Operator (Operator 12)
- Strict Mathematical Specification of the Spectral-Shape Equation-Solving Operator Set (Operators 13–18): Homological Solvability Theorems and the Endogenous Inversion Pipeline
- Operator 19: $\mathbb{Z}_2$ holonomy / σ charge-sign channel operator
- Part IV: Applications and Scientific Hypotheses
- Academic Hypothesis: The Sovereign Universe Tree of Life Protocol and Topological Residual Inheritance via State-Relation Entropy (SRE) Dynamics
- A SRE-Dynamics Inspired Multipath Topological Flow Purification Architecture and Localized Operator Implementation
- Research on Multipath Interference Preprocessing Algorithm for Fjord Underwater Acoustic Communication Based on SRE Topological Operators
- A Conjecture on Single-Photon Bidirectional Instantaneous Communication via Möbius Topological Flows Based on SRE Dynamics
- TECHNICAL REPORT: INTRINSIC ALGEBRAIC TOPOLOGY OF LIGHT AND THE SRE AXION MATRIX
- Stellar Spin and Galactic Rotation within the SRE-Framework: Operator-System combined with the SRE-v6.0 Dissipation-Compensation Duality Model
- Causal Compliance and Global Sea Surface Temperature Anomaly (ENSO-SSTA) Spatiotemporal Forecasting Audit Report: The 2D Convolutional Dissipative Network Paradigm (Astro-Cow-Net)
- Periodic Table Data Experiment - SRE-v3.0 Atomic Topological-Weight Reverse-Deduction Experimental Protocol
- Status-Relational Entropy-AI: A Differentiable Graph Learning Model Based on Topological Dynamics
- SRE Full Suite of 19 Operators: What a "Physical-Semantic Graph Computing" Operator System Can Do — A View from the Graph500 Benchmark
- Neural Information-Systems Theory: First-Order Thalamic Downsampling Instability and Second-Order Default-Mode-Network Integration Cascading-Failure Hypothesis
- Whole-Brain Parallelism and High-Dimensional Causal-Chain Topological-Spectrum Homomorphic-Mapping Mechanisms Based on Complex Causal-Network Topology
- Hypothesis of Gravitational Shielding Derivation (SRE)

Part I: Axiomatic Foundations — Abstract

This collection constitutes the **State-Relational-Entropy (SRE) Dynamics v1.6 axiomatic foundation suite**. Discarding spacetime and matter as pre-given background prerequisites, this work takes as its starting point a set of asynchronously evolving causal nodes. Through **dissipation-compensation duality, mutual-measurement mechanisms, emergent ontological ultraviolet boundary, and multi-scale rigid-boundary full-homomorphism mapping**, it deduces emergent behaviours of spacetime, the speed of light, time, and particles within the physical rendering layer.

> Core position: The physical world is not a direct one-to-one mirror of the underlying causal space. Observable spacetime and matter phenomena arise only after instantiation filtering and coarse-grained homomorphic mapping.

Brief Overview of Documents in This Suite

1. **SRE Dynamics User Guide**: Version management, citation guidelines, theoretical boundaries, and research paradigms.
2. **Foundational Framework for Information-Physical Interchange**: The axiomatic origin of the theory. Establishes three core principles and defines the meta-analytical framework for bidirectional equivalence between information and physics.
3. **Mutual-Measurement and the Emergent Origin of the Speed of Light c**: Reduces perception and measurement to topological mutual-measurement processes. Explains how the speed of light and time emerge as book-keeping products of dissipation-compensation, and accounts for the logical origin of local constancy of the speed of light.
4. **Methodological Revision of Causal Depth and Spatial Indexing**: Reinterprets time and space not as pre-existing axes, but as rendering projections of the causal network; introduces the concept of causal remapping.
5. **Mapping Preconditions**: Demonstrates that direct isomorphic mapping is infeasible. Proposes three-tier rigid truncation boundaries across micro-, meso-, and macro-scales, establishing constraints for full-homomorphism mapping from causal space to the physical layer.
6. **Unified Theory of Electronic Logical-Topological Structure and Physical Properties**: Interprets the electron as a self-consistent Möbius topological closed-loop at the instantiable minimal scale. Derives mass, electric charge, spin-1/2, and the uncertainty principle from properties of this topological loop.
7. **Magnitude Anchoring of Logical Depth N and Safety Shielding of Observation Protocols**: Estimates the equivalent logical depth of the electronic topological loop. Demonstrates that internal underlying degrees of freedom are protected by a physical-level natural barrier against observation and modification.

This suite addresses **how physical reality emerges given pre-existing causal differences**. It does not answer the ultimate origin of causal differences.

Status-Relational Entropy (SRE) Dynamics: Usage Specification and Theoretical Context

Version: v1.6 (aligned with complete manuscript suite 1-6)

> Version Note: This manual provides usage norms, version-priority rules, application paradigms and theoretical-boundary descriptions for the complete SRE-Dynamics manuscript suite. It is consistent with the v1.6 axioms and all derivative papers.

1. The Zenodo archive collection is sorted by public-release timestamp. It fully preserves the whole evolutionary history of SRE-Dynamics from early conceptual ideas and heuristic sketches up to the achievement of logical closure.

> Note: Versions v1.5.x and earlier stored in the archive are historical heuristic drafts containing simplified ontological assumptions. **They are intended solely for version-traceability and are not recommended as formal citation sources.**

2. The full v1.6 series manuscripts published after the formal establishment of the axiomatic system (*A Foundational Framework for Information-Physical Interconversion*) possess the highest value for derivation, validation and application.

> List of formally-citable manuscript suite (chapter titles as listed in the book's table of contents; current chapter files in parentheses):
>
> 1. *A Foundational Framework for Information-Physical Interconversion* (`1-Axiomatic_Introduction_1.6_E.md`) — Axiomatic Foundation
> 2. *Reciprocal Measurement and the Mapping Origin of* c (`2-Light_Measurement_1.6_E.md`)
> 3. *Methodological Revision on Causal Depth and Spatial Indexing* (`3-Time_Causal_Depth_1.6_E.md`)
> 4. *Mapping Prerequisites* (`4-Mapping_Frontend_1.6_E.md`) — Multi-Scale Homomorphic Mapping
> 5. *A Unified Theory of Electronic Logical-Structural and Physical Properties Based on the Minimal Instantiable Topological Scale* (`5-Electronics_1.6_E.md`)
> 6. *Magnitude Anchoring of Logical Depth* N *and Observational-Protocol Safety Masking* (`6-Safety_Supplement_Electronics_1.6_E.md`)

3. When investigating the underlying structure of causal-space and the physical rendering-layer using SRE-Dynamics, one shall prefer the precise mathematico-topological expressions provided in this suite with unified ontological alignment. Classical-physics frameworks and the SRE framework can establish mathematical correspondence within their respective valid scales. Depending on the observational scale and computational complexity of a given study, either system may be employed alone or in hybrid combination.

Two mutually complementary research pathways are supported by this theory:
① Substitutability test: Verify the correspondence between the SRE framework and well-established physical results, so as to carry out self-consistency checks for logical closure;
② Forward-looking exploration: Adopt a mathematics-and-logic-first methodology. Deploy the complete toolkit including dissipation-compensation duality, reciprocal measurement, multi-scale rigid boundaries and full homomorphic mapping to make deductions for insufficiently-explored physical structures.

4. Reminder on ontological constraints: SRE does not answer the fundamental question of the ultimate origin of causal differences. All deductions within this suite describe **how a pre-existing collection of causal nodes with asynchronous informational differences undergoes dissipation-compensation iteration, instantiation screening and multi-scale homomorphic mapping to render macroscopic spacetime, particles and interactions**. The 0-to-1 creation problem of causal space has not attained full logical closure under the current axiomatic system.
5. SRE-Dynamics is an open-evolving theoretical system. With further refinements of hierarchical dissipative self-organizing binary-network simulations and cross-scale deductions, the present model allows for future corrections, extensions and potential replacement by more complete and self-consistent theories.

A Foundational Framework for Information-Physical Interconversion

Version: 1.6 (Axiom-Purified & Quantitatively Mapped Edition)

Three Core Principles

I. Statistical Foundation (Mapping-Layer Causal Constraint Principle)

All physical foundations originate from information statistics. Physical constants (e.g., $\hbar$ and c) are not intrinsic properties of matter. Instead, they are the **emergent ontological ultraviolet boundary** and the **emergent causal upper bound of propagation** arising from the mapping of information-space onto physical reality. Therefore, information theory is not merely a descriptive tool; it serves as the fundamental metalanguage for analyzing the underlying laws of physics.

II. Residual Observation (Instantiation-Boundary Principle)

All physical observations are essentially processes of information transmission and logical residual evaluation. The Planck scale marks the boundary for logical instantiation within the system. Observation acts as a dynamic mechanism that drives the evolution of physical systems through continuous evaluation and compensation of data discrepancies.

III. Functional Interchangeability (Digital-Analog Conversion Principle)

Physical theories and information theories are functionally equivalent and interchangeable. This duality establishes a real-world **"digital-analog conversion toolkit"**: by defining well-defined rigid boundaries across macroscopic, mesoscopic, and microscopic scales, gaps within physical models can be remedied via algorithmic optimization, while intricate informational structures can be instantiated and mapped following physical principles.

Conclusion

SRE-Dynamics delivers an innovative two-way tool for scientific research. By establishing the equivalence between the informational and physical domains, we gain the capability to resolve previously intractable scientific paradoxes through **cross-domain recoding**. By optimizing the system's logical pathways within the constraint scope of underlying causal-instantiation boundaries, we can achieve precise intervention upon observable physical outcomes.

Reciprocal Measurement and the Mapping Origin of c

Version: 1.6.0 (Revised: Dissipation-Compensation and Möbius-Residue Alignment)

I. Physical Essence of Perception: Reciprocal Measurement

Perception is a physical process of reciprocal measurement embedded within the dissipation-compensation causal network. It may also be regarded as an emergent outcome generated by the direct topologically-correlated evolution of two topological manifolds.

- **Causal-Topological Instantiation**: Two quantum-evidential causal nodes undergo asynchronous evolution. Their causal intersection yields a Möbius-type topological residual manifold, namely the light residual Ψ_{light}. Only after passing joint evaluation of informational dissipation and network compensation cost can this interaction be instantiated and projected onto the physical rendering layer, becoming a “physically perceptible event”.
- **Instantiation Threshold (Emergent Ontological Ultraviolet Boundary)**: An interaction must satisfy both causal consistency conditions and dissipation-compensation budget constraints. Only qualified interactions transition from the pending-for-instantiation state in causal space into observable physical reality. The Planck scale marks this ontological instantiation boundary rather than the intrinsic granular size of underlying causal nodes.

II. Mapping the Measurement Limit: Emergent Origin of c

The speed-of-light constant c is not merely a kinematic constant. It is the **emergent causal upper bound of propagation** generated by the dissipation-compensation duality framework in the course of information-to-physical mapping. Space itself is no pre-existing background and is likewise an emergent product of the causal network. Spatial metrics and light-mediated reciprocal measurement originate from the same source and co-generate under distinct topological-rank regimes governed by the BBP spectral phase transition. Precisely this intrinsic coupling logically guarantees the invariance of locally measured speed of light.

- **Effective Routing Bandwidth** c_{eff}: In the high-dissipation primordial-universe regime ($z \geq 4.1605$), substantial topological compensation overhead consumes the network’s computational resources, and the effective throughput for packet propagation c_{eff} decreases. Conformal gauge transformation rescales the emergent metric tensor synchronously; therefore for local observers within the rendering layer, the measured speed of light remains Lorentz-invariant.
- **Global Constraint from the Instantiation Boundary**: The emergent ontological ultraviolet boundary at the Planck scale sets a hard lower bound for logical-operation costs. If the compensation cost required for one reciprocal-measurement exceeds the network’s stability capacity, the process remains in the pending-for-instantiation state within causal space and cannot be rendered into observable spacetime events.

III. Time: Cumulative Book-Keeping of Dissipation-Compensation Behaviours

Time possesses no pre-existing axis-like container ontology; it emerges from cumulative book-keeping of numerous reciprocal-measurement events. Successive steps in the evolution of the world necessarily carry informational differences. Perception of the duration of such differences constitutes time. If no distinguishable informational difference exists between adjacent evolutionary steps, no new temporal increment will be generated and time tends to stagnate.

- **Non-Axis Nature**: Time is not a pre-existing background container. Under the constraint of the emergent causal upper bound of propagation, it is the sequential counting of successive topological dissipation-compensation operations.
- **Perception as Duration**: Every reciprocal-measurement interaction mediated by the Möbius light-residual manifold incurs irreducible topological routing overhead. These measurement events, weighted by compensation cost, stack continuously and manifest macroscopically as the experiential flow of temporal duration.

IV. Logical Invariance of Locally-Measured Speed of Light across Reference Frames

The locally measured speed of light remains invariant in all reference frames. Observers, observed systems and measurement apparatuses are all emergent products mapped from the same causal network, collectively governed by dissipation-compensation duality.

- **Mapping-Protocol Invariance**: All entities within the physical rendering layer share the same emergent ontological ultraviolet boundary and causal-routing constraints. Relative motion modulates the local effective throughput c_{eff}, yet conformal metric rescaling performs algebraic cancellation for every local observer and keeps the numerical value of measured speed of light invariant.

Notes on Relationship with Earlier Versions

Version 1.6.0 supersedes Version 1.5.2.

- Version 1.5.2 is an early operational sketch that describes the “hardware-interface” behaviour of the causal network under the simplified assumption of global fixed baud-rate synchronisation.
- Version 1.6.0 integrates dissipation-compensation duality, the Möbius topological residual of light, BBP spectral rank transition and conformal-gauge covariance from the main SRE v6.0 paper. Reciprocal measurement is reinterpreted as an exchange process mediated by “light as topological residual”, constrained by dynamic topological-compensation costs instead of the limits of fixed single-logical-cycle synchronisation. As the operational-measurement layer, this version is fully compatible with SRE Axioms v1.6.

Methodological Revision on Causal Depth and Spatial Indexing

Version: 1.6.0 (Dissipation-Compensation and Möbius-Residue Aligned Edition)

> Version Note: This version supersedes v1.5.1 (Causal Topology Edition). The earlier heuristic sketch treating Planck-scale quantities as native pixel or atomic clock cycles of the underlying network has been replaced. Reconstructed fully under SRE Axioms v1.6 and the dissipation-compensation duality framework, Planck quantities are defined as **emergent ontological ultraviolet boundaries**, rather than intrinsic granular parameters of causal nodes.

I. The Non-Axis Nature of Time: Causal Logical Depth as Dissipation-Compensation Book-Keeping

Time is neither a fundamental dimension nor a pre-given physical axis. Within the SRE framework, time is not a hardware clock native to the underlying network. Instead, it is the emergent manifestation of causal logical depth in the process of informational state change, after the stacking of reciprocal-measurement events.

- **Causal Triggers and Book-Keeping Flux**: The sense of temporal "flow" is the macro-level mapping of sequential book-keeping for a large ensemble of reciprocal-measurement events. Planck time τ_P is an emergent ontological ultraviolet boundary, setting the lower cost bound for logical operations eligible for instantiation. **It is not an intrinsic atomic cycle clock of the causal network**. Causal differences falling below this cost threshold remain in the pending-for-instantiation state within causal space and cannot generate new temporal increments.

- **Topological Overhead for Global Consistency**: Causal links themselves are discrete logical associations, yet the perception of duration arises from the topological routing overhead required for both parties in reciprocal measurement to finish dissipation evaluation and compensation settlement. Time may therefore be understood as the topological metabolic cost incurred by the causal network to maintain global self-consistency within the rendering layer. This book-keeping process is continuously driven by reciprocal-measurement mediated via Möbius light-residual manifolds.

II. Spatial Indexing: Rendered Projection of Causal Connectivity

Three-dimensional space is no pre-existing physical entity. It is a dimensionality-reduced rendering-projection system generated through dissipation-compensation duality for the causal network to classify and navigate discrete causal relationships.

- **Causal Nodes and Instantiation Boundaries**: Physical distance is no longer simply equivalent to the count of intermediate logical steps required for node synchronization. It is a topological book-keeping quantity jointly evaluated from the topological dissipation tensor and the compensation operator. The Planck length l_P is an emergent ontological ultraviolet boundary that marks the instantiable causal-connectivity boundary of the physical rendering layer, **rather than the minimal pixel granularity of the underlying causal network**. Causal differences below this boundary will not be projected into distance-related spatiotemporal states of affairs. Interpreted heuristically from a physical-image perspective: distance reflects the degree of coherence degradation between two topological residual manifolds. Higher coherence corresponds to lower mutual informational dissipation, which in turn yields a shorter macroscopic spatial distance.

- **Dimensional Projection**: Our perception of three-dimensional geometry arises from data reduction. The system projects complex high-dimensional causal topologies onto effective spatial ranks unlocked across the BBP spectral phase transition, yielding simplified spatial metrics for reciprocal-measurement interactions among causal nodes.

III. Spatiotemporal Interchangeability: Causal Re-mapping

Distinguishing time as **dissipation-book-keeping depth** and space as **rendered projection of causal connectivity**, SRE opens a theoretical path toward spatiotemporal interchangeability.

- Redefinition of Displacement: Physical motion shall no longer be interpreted as continuous translation inside a pre-supposed background container. Instead, it consists of dynamic re-weighting of associative links among discrete causal nodes, driven by adjustments of compensation flows triggered by local variations of informational dissipation.

- **Apparent Instantaneous Leap (Causal Re-mapping)**: The theoretical objective is to perform "causal re-mapping": bypass sequential intermediate causal steps and establish direct topological associative links between two sets of quantum-evidential nodes. Subject to network-stability and instantiation-boundary constraints, this operation minimizes traversed causal book-keeping depth and thereby produces an apparent instantaneous-leap effect within the physical rendering layer. This process cannot violate the self-consistency constraints of the underlying causal network; it remains purely an apparent effect at the rendering layer and does not break causal self-consistency.

Key Rewrite Reference Memorandum

1. Removed content inherited from old v1.5.1: *Planck time as the atomic cycle for single causal triggers, Planck length as indexing pixel granularity, distance as count of intermediate logical steps*.
2. Retained and reconstructed core conceptual kernels:
    - Time = emergence of causal logical depth, where logical depth originates from **dissipation-compensation book-keeping of reciprocal measurements**, not a native hardware clock.
    - 3D space = dimension-reduced rendering projection of high-dimensional causal topology.
    - Displacement = re-weighting of causal associative links.
    - Causal re-mapping: establishing direct links to bypass intermediate steps, producing apparent instantaneous-leap effects at rendering layer, with added constraint conditions.
3. Uniform terminology across the whole manuscript: **emergent ontological ultraviolet boundary, dissipation-compensation duality, reciprocal measurement, Möbius light-residual manifold, BBP spectral phase transition, pending-for-instantiation state, physical rendering layer**.

Mapping Prerequisites

Version: 1.6 (Revised for Multi-Scale Homomorphic Mapping)

> Version Note: This revision reconstructs the ontological constraints of mapping and no longer presupposes a direct one-to-one projection from the underlying layer to the rendering layer. Drawing on the ontological features of the hierarchical dissipative self-organizing binary-network substrate, it introduces **multi-scale rigid boundaries-homomorphic mapping** as a necessary precondition for valid mapping. Terminology is fully consistent with the SRE Axioms v1.6, Reciprocal-Measurement paper, and Causal-Depth-and-Spatial-Indexing manuscript suite. This document does not address the ultimate origin of causal differences; it characterizes only the intrinsic constraints and feasible pathways for instantiated projection from causal space to the physical rendering layer.

I. Intrinsic Ontological Constraints of Mapping

SRE-Dynamics takes **a set of quantum-evidential causal nodes with asynchronous informational differences** as the underlying input for mapping. Asynchronous evolution already proceeds inside the underlying causal space and numerous causal interactions exist, yet the vast majority of these interactions remain in the **pending-for-instantiation state** and have not yet generated macroscopically observable spatiotemporal states-of-affairs.

The underlying causal network is subject to two ontological constraints that render a pure direct isomorphic mapping fundamentally unfeasible:

1. **Decoherence-Dissipation Constraint**: Asynchronous activation and local frustration effects exist within the underlying network. A large number of degrees-of-freedom undergo decoherence and dormancy. Evolution is accompanied by irreversible informational dissipation, making lossless upward transmission of every logical degree-of-freedom from the substrate impossible.
2. **Emergent Ontological Ultraviolet-Boundary Constraint**: Limited by instantiation-cost thresholds, numerous minute underlying logical differences cannot pass instantiation screening and cannot be converted into spatiotemporal states-of-affairs within the physical rendering layer.

An isomorphic mapping requires bijective one-to-one correspondence and complete preservation of all underlying details. Owing to the two constraints above, direct bijective projection from underlying causal space onto the macroscopic physical rendering layer cannot be accomplished. Mapping cannot be performed unconditionally and must be subject to scale-dependent preconditions.

II. Multi-Scale Rigid Boundaries: Pre-Requisite Thresholds for Valid Mapping

To accomplish information-physical instantiated mapping, **rigid coherence truncation boundaries** must be established at the micro-, meso-, and macroscopic scales respectively.

These rigid boundaries are not material physical boundaries. Instead they are statistical-topological screening thresholds: they filter out degrees-of-freedom that are unstable, strongly fluctuating, and unable to satisfy instantiation conditions at the given scale; only causal structures and relational associations possessing statistical stability at that scale are retained.

- **Micro-scale rigid boundary**: Select the high-topological-coherence core region of the system, screen out strongly fluctuating mesoscopic bifurcation degrees-of-freedom outside the core, and preserve locally stable topological-manifold behaviour.
- **Meso-scale rigid boundary**: Set cascade-amplification thresholds to filter extreme nonlinear stochastic bifurcations. Retain only statistically self-consistent causal associations in the ensemble-averaged sense and discard isolated accidental stochastic fluctuations.

- **Macro-scale rigid boundary**: Merge degrees-of-freedom that have undergone long-range decoherence. Extract only collectively emergent observable quantities without tracking detailed states of individual underlying nodes.

> In essence, rigid boundaries constitute scale-specific instantiation-selection criteria. Each hierarchical level defines which causal structures may participate in upward mapping and which degrees-of-freedom shall be compressed and screened out.

III. Homomorphic Mapping under Multi-Scale Constraints

When the three sets of rigid boundaries are simultaneously enforced, a **full homomorphic mapping** can be established from causal space to the physical rendering layer.

A homomorphic mapping does not demand one-to-one correspondence between substrate-layer and rendering-layer elements. It permits many-to-one coarse-graining and informational compression, while preserving the core dynamical structures of the system: dissipation-compensation duality, causal-temporal chains, coherence-degradation laws of topological residual manifolds, and BBP spectral phase-transition behaviour.

Based on this mapping framework, cross-scale physical deductions can be carried out, covering: gluon-sea explanations for hadron mass and reconstruction of Maxwell's equations at the quantum scale; multi-scale modelling of the Navier-Stokes equations and VASP-style first-principles analyses at the molecular scale; Earth-interior stress calculations and stellar-spin dynamics at the macroscopic scale. Full derivations and numerical examples are provided in other works within this series.

Within this mapping mechanism:

1. Numerous dormant, decoherent microscopic degrees-of-freedom from the substrate are merged and compressed via homomorphic mapping and are not output as independent physical states-of-affairs at higher layers.
2. Stable causal interactions that pass screening undergo dissipation-compensation accounting via Möbius topological-residual-manifold-mediated reciprocal measurement. After satisfying instantiation conditions, they transition from the pending-for-instantiation state into the physical rendering layer and produce observable phenomena of spacetime, matter, and interactions.
3. Neither time nor space are pre-existing background stages at the fundamental level. Both are derivative products emergent from dissipation-compensation book-keeping after homomorphic-mapping output.

> Multidimensional Scaling (MDS) dimensionality reduction in numerical simulations provides a concrete computational realisation of this homomorphic-mapping concept: it discards abundant stochastic details of the underlying binary network and homomorphically projects high-dimensional topological relations onto lower-dimensional emergent geometric manifolds.

IV. Logical Primacy of Reciprocal Measurement

Within the mapping workflow, **reciprocal measurement mediated by light-residual topological manifolds holds logical primacy**.

Time and space coordinates do not exist prior to measurement interactions. On the contrary: one complete reciprocal-measurement event represents one full round of dissipation-evaluation and compensation-settlement. Each settlement contributes one new increment to temporal book-keeping and simultaneously updates topological-book-keeping results for spatial metrics.

Durational perception and spatial distance inside the rendering layer are macroscopic manifestations after homomorphic-mapping of large-numbers of reciprocal-measurement events. Conformal-gauge covariance further guarantees the invariance of locally measured speed-of-light inside the rendering layer.

V. Conclusion

The SRE Mapping Prerequisites define the domain-of-applicability for the framework. This work does not answer the question of the ultimate origin of causal differences. Instead it articulates the constraints required for mapping to occur.

Direct isomorphic projection from the underlying substrate to the physical rendering layer does not exist. Only under the constraints of micro-meso-macro three-level rigid coherence-truncation boundaries can homomorphic instantiated mapping from causal space to the physical rendering layer be realised. Spacetime, matter, and interactions are all emergent outputs of this homomorphic-mapping procedure.

Core-Concept Memorandum (Version-traceability, internal-document use only)

1. Discarded assumptions inherited from older versions: absolute-symmetry zero-void initial state, single initial pulse, distance equivalent to logical-hop count, Planck-scale quantities treated as hardware granularity of underlying network, unconditional direct mapping.
2. New core logical components: decoherence-dissipation plus emergent ontological ultraviolet boundaries of the substrate render isomorphic mapping infeasible; multi-scale rigid coherence-truncation boundaries as pre-conditions; constrained homomorphic mapping preserving core dynamics; logical primacy of reciprocal measurement.
3. Unified terminology inventory: emergent ontological ultraviolet boundary, pending-for-instantiation state, physical rendering layer, dissipation-compensation duality, Möbius topological-residual manifold, reciprocal measurement, BBP spectral phase transition, rigid coherence-truncation boundary, homomorphic mapping.

A Unified Theory of Electronic Logical-Structural and Physical Properties Based on the Minimal Instantiable Topological Scale

Version: v1.6 (Axiom-Homomorphic-Mapping Aligned Revision)

Abstract

Within the full SRE-Dynamics framework, presuppositions of material entities are discarded; space, time and energy are not taken as pre-given background. This paper introduces the **minimal instantiable topological scale** l_{min}, corresponding to the emergent ontological ultraviolet boundary (Planck length) of the physical rendering layer. It is proposed that the electron is not a pre-existing material entity in the underlying causal space. Instead, it is an irreducible self-consistent topological closed loop generated within the physical rendering layer via homomorphic mapping, after causal interactions undergo dissipation-compensation accounting and screening by multi-scale rigid boundaries.

By back-propagating the topological frequency and book-keeping ratios of this topological closed loop under reciprocal-measurement sampling, it can be demonstrated that rest mass, unit electric charge, spin-1/2, and the uncertainty principle are all inevitable observable effects produced by this topological closed loop within the physical rendering layer.

I. Core Definition: Minimal Instantiable Topological Scale of the Rendering Layer

In the SRE framework, physical reality is the instantiated rendering result of causal space screened by multi-scale rigid boundaries and processed via homomorphic mapping.

> Note: l_{min} is not an intrinsic native discrete step of the underlying causal network. The underlying causal space permits smaller logical differences; interactions falling below the instantiation boundary remain in the pending-for-instantiation state and cannot be projected into the physical rendering layer.

1. **Minimal instantiable topological scale** l_{min}: The lower scale bound for topological events to complete instantiation inside the physical rendering layer. It corresponds to the **emergent ontological ultraviolet boundary (Planck length)** in SRE axioms and sets the resolution lower limit for reciprocal-measurement book-keeping within the rendering layer.
2. **Derivative Spacetime**: Spatial metrics and temporal increments are both topological book-keeping products of dissipation-compensation duality driven by reciprocal measurement. l_{min} defines the minimum effective scale output by this book-keeping system.

II. Logical-Topological Formulation of the Electron: N-Scale Self-Consistent Topological Closed Loop

No pre-packaged electronic “algorithm unit” exists in the underlying causal space. The electron is a topological structure emerging within the physical rendering layer after iterative dissipation-compensation of numerous asynchronous causal interactions, once instantiation conditions are satisfied. It consists of closed-link segments built from N minimal instantiable topological scales l_{min}.

- **Topological Closure**: The rendering-layer causal sequence $[A_1, A_2, \dots, A_N]$ possesses closed-loop back-coupling characteristics: A_N establishes causal association back to A_1, forming a self-consistent topological closed loop. Internal reciprocal-measurement and dissipation-compensation accounting persist inside the loop, maintaining overall self-consistency of residuals.

- **Möbius-phase Characteristic**: This closed loop exhibits a two-layered phase topology. To return topological residuals to their initial symmetric state, $2N$ reciprocal-measurement triggers at the l_{min}-scale must be completed; phase restoration cannot be achieved after only N cycles.

III. Parametric Verification of Observable Physical Properties

Taking the rendering-layer minimal instantiable topological scale l_{min} as the scale benchmark, each observable physical property of the electron may be interpreted as emergent effects of this topological closed loop under reciprocal-measurement sampling.

1. Rest Mass: Measure of Topological Processing Overhead of the Closed Loop

- **Logical-Topological Definition**: Rest mass m characterises the additional topological book-keeping overhead that reciprocal-measurement protocols must expend to process this topological closed loop.
- **Physical Interpretation**: When reciprocal measurement attempts to gauge or displace this electronic closed loop within the rendering layer, full pre-processing of dissipation-compensation for the N internal topological segments must first be completed. This l_{min}-based topological processing load manifests physically as inertial effects.
- **Corollary**: $m_e \propto N \cdot l_{min}$. The experimentally observed constant mass of the electron originates from structural stability of the closed-loop depth N of this emergent topological closed loop.

2. Unit Electric Charge: Characteristic Signature of Topological-Step Bias of the Closed Loop

- **Logical-Topological Definition**: Electric charge e is the constant topological logical bias exerted by this electronic topological closed loop onto external causal links during each round of l_{min}-scale reciprocal-measurement cycle.
- **Physical Interpretation**: Coupled with surrounding rendering-layer topological structures via reciprocal-measurement protocols mediated by Möbius light-residual manifolds, the electronic closed loop continuously introduces fixed offsets to neighbouring causal associations and generates logical potential differences at the topological level.
- **Corollary**: Unit electric charge represents a persistent topological modification effect at the l_{min} instantiable scale, macroscopically manifesting as electromagnetic interaction.

3. Spin-1/2: Phase-Locking of the Möbius Closed Loop

- **Logical-Topological Definition**: Spin arises from the reciprocal-measurement book-keeping beat required for the topological closed loop to complete global causal-consensus settlement.
- **Physical Interpretation**: Constrained by the Möbius two-layer topology, $2N$ reciprocal-measurement trigger cycles are required for topological residuals of the closed loop to return to the initial symmetric configuration.
- **Corollary**: This topological behaviour, which demands a doubled cycle count for phase restoration, yields fermionic spin-1/2 observational behaviour at the statistical level.

4. Uncertainty Principle: Incomplete-Sampling Effect from Instantiation Boundaries

- **Logical-Topological Definition**: The uncertainty relation originates from the fact that reciprocal-measurement sampling itself consumes l_{min}-scale instantiation resources; complete synchronous sampling cannot be performed on a finite-depth topological closed loop.
- **Physical Interpretation**: Any single reciprocal-measurement sampling consumes at least one unit of l_{min}-scale book-keeping resource. The internal topological depth N of the electronic closed loop greatly exceeds the instantiation resolution of a single sampling. When sampling anchors the position (topological index), the cyclic frequency (corresponding to momentum) cannot be fully resolved simultaneously.

- **Corollary**: The uncertainty principle is an incomplete-sampling effect of reciprocal-measurement imposed by the emergent ontological ultraviolet boundary.

IV. Conclusion: Particles as Topological-Rendered Products after Homomorphic Mapping

From the above deductions we arrive at the core proposition: **All intrinsic constants experimentally measured for the electron are statistical emergent manifestations after global reciprocal-measurement protocol sampling of a self-consistent topological closed loop constructed from N-fold minimal instantiable topological scale l_{min}.**

1. **The electron possesses no underlying material substance**: It is not a primitive object of causal space. Instead, it is a self-consistent topological closed-loop structure within the rendering-layer that resists simple linear expansion, after screening by multi-scale rigid boundaries and homomorphic mapping.
2. **Particle stability originates from self-consistency of the topological closed loop**: As long as dissipation-compensation accounting inside the closed loop maintains global self-consistency and residual conditions remain satisfied, this topological object acquires logical stability within the physical rendering layer.
3. **Unification**: All intrinsic physical properties of the electron can be reduced to topological reciprocal-measurement book-keeping relations formulated in terms of l_{min}.

V. Closing Remarks

This paper demonstrates that within the SRE-Dynamics framework, a unified topological-logical account for all known observational behaviours of the electron can be achieved via the minimal instantiable topological scale l_{min}. This suggests that selected postulates of particle physics may be transformed into deductive inferences of causal-topology, dissipation-compensation and reciprocal measurement.

Full numerical simulation of this topological structure and benchmarking against the hierarchical dissipative binary-network model will be presented in subsequent works of this series.

SRE Dynamics Supplementary Note: Magnitude Anchoring of Logical Depth N and Observational-Protocol Safety Masking

Version: v1.6 (Axiom-Homomorphic-Mapping Aligned Revision)

> Version Note: This supplementary note is aligned with the full SRE v1.6 manuscript suite. l_{min} denotes the **minimal instantiable topological scale (emergent ontological ultraviolet boundary)** of the rendering layer. The electron is a self-consistent topological closed-loop that emerges in the physical rendering-layer after multi-scale rigid-boundary screening and homomorphic mapping. The safety discussed herein applies to deep internal topological degrees-of-freedom of the closed loop; routine physical manipulation of the electron as a whole object within the rendering layer is not prohibited.

1. Derivation of Logical Depth N

Within the SRE framework, the electron is a stable self-consistent topological closed-loop residing in the physical rendering layer. The parameter N represents the equivalent count of instantiable topological scales for this closed loop. It is not an empirical constant native to the underlying causal space, but jointly determined by the characteristic rendering scale of the system and the lower instantiable-scale bound of the rendering layer.

- Formula:

$$N = \lambda_c / l_{min}$$

- Magnitude evaluation: Substituting known parameters (Compton wavelength $\lambda_c \approx 10^{-12}\ \mathrm{m}$ and minimal instantiable topological scale $l_{min} \approx 10^{-35}\ \mathrm{m}$), the equivalent logical depth of the electron yields:

$$N \approx \frac{10^{-12}}{10^{-35}} = 10^{23}$$

Logical conclusion: The electron is not a simple primitive topological unit, but a high-order emergent topological module with extremely high redundancy (equivalent topological count of order 10^{23}).

2. Sampling Theorem and Observational Masking

This supplementary note emphasises that the “invisibility” of deep internal topological details inside the electron closed-loop is enforced jointly by information-theoretic sampling limits and coarse-graining effects of multi-scale homomorphic mapping.

- Resolution gap: All physical observational protocols are bounded by the rendering-layer minimal instantiable topological scale l_{min}. The resolving power of a single physical observation for the full internal topology of the electron closed-loop is merely on the order of $\frac{1}{N}$ (i.e. $1/10^{23}$).

- Statistical smoothing: Constrained by multi-scale rigid-boundary filtering and full homomorphic-mapping mechanisms, numerous fine-grained internal topological degrees-of-freedom of the closed-loop undergo many-to-one coarse-grained compression. Only statistically stable collective effects are preserved and finally manifest as observable mass, charge and spin within the physical rendering layer.

3. Core Safety Criterion: Strict Read-Only Physical-Level Restriction

The core significance of this supplementary note is to demonstrate the intrinsic physical safety within this theoretical model.

- Permission Lock: l_{min} constitutes the lower scale bound for reciprocal-measurement addressing inside the physical rendering layer. Any attempt to directly intervene inside the electron closed-loop and perform operations at equivalent scales far smaller than l_{min} cannot achieve valid physical addressing.

- Causal Barrier: A vast order-of-magnitude gap exists between the electron's equivalent logical depth N and physical observational limits. Deep internal topological degrees-of-freedom of the closed-loop are naturally isolated within a physically-inaccessible black box.

> Important distinction: This restriction **does not forbid routine physical manipulation of the complete electron closed-loop within the rendering-layer**, such as electromagnetic coupling or particle-scattering experiments. What is prohibited is penetrating the closed-loop to rewrite its deep internal topological details.

- Conclusion: Observers may only read stable observables delivered by homomorphic-mapping in the rendering layer; no instructions can be written to the deep internal topology of the closed-loop. Constraints originating from instantiation boundaries form an intrinsic cosmic-scale hard-encryption barrier.

4. Final Conclusion

This supplementary note establishes the following inference: physical reality is the rendering output after homomorphic-mapping sampling of causal space.

The enormous equivalent topological count (10^{23}) of the electron closed-loop safeguards the stability of particle-based matter. Subject to natural constraints imposed by rendering-layer instantiation boundaries, deep internal topological degrees-of-freedom of this topological closed-loop are physically inaccessible and unmodifiable, possessing intrinsic safety properties.

Author's Note: Completion of this supplementary note marks that SRE-Dynamics has not only achieved a closed deductive chain spanning micro- to macro-scales at the logical level, but also constructed a theoretically impenetrable protective boundary of physical origin.

Part II: Emergence of All-Things — Abstract

This part corresponds to **Emergence of All-Things within State-Relational-Entropy (SRE) Dynamics**. Building upon the axiomatic foundation suite of Part I, it reconstructs physical pictures across multiple domains based on the ontology of discrete causal-information networks. Electromagnetism, mechanics, elementary particles, fluid dynamics, and cosmic gravitation are no longer treated as fundamental a-priori axioms. Instead, all these phenomena are derived from the topological evolution of causal networks via **dissipation-compensation duality, mutual-measurement mechanisms, multi-scale rigid-boundary full-homomorphism mapping, and BBP spectral-rank phase transition**.

> Core Position: Branches of classical physics are high-level emergent effective theories of discrete causal networks under specific constraints. Some modules adopt a pragmatic hybrid approach of "ontological topological picture plus experimental observation mapping anchors". Fully endogenously deriving all universal physical constants remains a long-term research objective.

Brief Overview of Documents in This Suite

1. **Theory of Hierarchical Dissipative Self-Organizing Binary Network Dynamics**: The bottom-level discrete-evolution prototype of the SRE framework. Starting from binary-spin asynchronous activation-dormancy rules, it spontaneously generates topologically coherent kernels through finite-size scaling phase transitions. It clarifies the conceptual boundary between MDS multi-dimensional-scaling simulation prototypes and the SRE physical-rendering layer, serving as the computational substrate for the complete SRE cosmic picture.

2. **Emergence of Multidimensional Spacetime and Dynamical Gravity via Regularized Causal Information Networks**: SRE cosmology v6.2-rev. Fully background-independent; spacetime, gravitation, and the speed of light are all macroscopic emergent effects of causal-information networks. It introduces the BBP spectral-rank phase transition to interpret dimensional transitions. Bootstrap statistical simulations based on SDSS/eBOSS spectral datasets yield the phase-transition redshift, providing observationally falsifiable theoretical imprints for JWST and the Roman Space Telescope, and resolving the puzzle of early massive galaxy formation.

3. **Composite Elementary Particles and Relation-Space Emergence within the SRE-Dynamics Topological Paradigm**: Qualitative-semi-analytical topological hypothesis for composite particles. It describes how phase-coherence saturation of two internal causal closed-loops yields relation-space collapse, emergence of strong-interaction effects, and nonlinear amplification of rest-mass driven by combinatorial-explosion of secondary feedback paths. It distinguishes the explanatory capacity of the present mechanism from quantitative predictions not yet completed.

4. **Rigorous Reconstruction of Classical Electrodynamics Based on Discrete-Graph Topology and Bidirectional Causality**: Engineering-SI-oriented reconstruction of electrodynamics. It defines four sets of observation-mapping anchors to convert dimensionless topological observables into laboratory SI physical quantities. It recovers circuit laws, RLC resonance, and mass-energy equivalence, establishing interface bridges for power-system and semiconductor-device simulations.

5. **Rigorous Reconstruction of Maxwell-Field Equations via Pure-Dimensionless Graph-Cohomology and Global-Evolution Steps**: 0-State dimensionless ontological-layer manuscript. Using graph cohomology, elementary charge, Planck's constant, vacuum characteristic impedance, and the fine-structure constant are treated as algebraically emergent invariants. It topologically reconstructs the Maxwell-field equations without importing external empirical physical constants.

6. **Emergence Inevitability and Algebraic-Computational Methods of Turbulence from Discrete-Microscopic-Causal Statistics and Multi-Dimensional-Manifold Reconstruction (v3.0 revised)**: Discarding the continuum-medium assumption, it derives criterion operators from discrete causal-statistical axioms, proves a series of algebraic theorems, and strictly degenerates to the Navier-Stokes equations in the continuum limit. Simulations verify that turbulence together with

coherent vortex structures are inevitable outcomes of topological phase-transitions. Version 2 corrects v1's sign contradictions (the activation probability is reformulated as $P_{\text{active}} = 1/(1+e^{\Psi})$; M is strictly split so that its symmetric part $\mathbf{S}$ carries the metric/geometry and its antisymmetric part $\mathbf{A}$ carries chirality) and deliberately downgrades three overclaims: the "unique critical point" becomes a **smooth transition band** spanning roughly 1.5 decades; the claim of "agreement with K41" is restricted to a proxy one-dimensional spectrum measured at about −1.85, deviating further from −5/3 as N grows; and the "rigid central axis" is withdrawn as an artefact of the metric-reconstruction method. An audit script cross-checks 8 tables and 276 numerical values across both language versions and the data tables with 0 inconsistencies. SI-dimensional calibration for fluid systems is reserved for future work. Version 3 further corrects v2 via a program-correctness gate (reproduce_ns62.py reproduces the section 6.2 phase diagram to Delta<=1e-4) and two prior verifications (verify_paper_claims.py): it recasts Psi in logarithmic form and closes the loop with verify_dissipation.py (error <=1.1 percent); it supplements an analytic finite-size convergence mechanism (dissipation master equation -> universal curve C(z), beta_infinity=0, via beta_dissipation_theory.py), and uses large-N extrapolation to refute the persistent-steepening claim.

7. **Oseen Vortex Initial-Value Experiment: Repositioning the Empirical Report (v2)**: A repositioning companion to the original report *"Empirical Verification of Oseen Vortex Manifold Emergence and Topological Locking under the SRE-MDS Discrete Dynamics Paradigm"*. The complete original experimental setup and observed data are retained; only two overclaims are corrected. The order parameter $\Phi(N) \equiv 1.0000$ is in fact a trivial "zero-pruning" result rather than a demonstration of noise robustness, and is downgraded to an indicator of the zero-dissipation limit. The residual 8.93×10^{-1} is an **embedding error** — the binary dissimilarities cannot be represented exactly in three-dimensional Euclidean space — and the "intrinsic non-Euclidean curvature" reading is deleted. Re-positioned against the main paper's phase diagram, the experiment's $\Lambda = 0.05$ lies at the **upper edge of the transition band** (structural density ≈ 0.44, spectral slope ≈ −1.83), being neither laminar nor "extremely high Reynolds number". A correct experimental design for testing noise robustness and a list of follow-up measurements are also provided.

8. **Emergence of Classical Mechanics from Discrete Causal-Information Networks**: Establishes ontological mappings between SRE-network observables and Newtonian classical-mechanics. It states three necessary conditions for the emergence of classical behaviour: large-sample coarse-graining, sufficient environmental decoherence, and staying away from topological-phase-transition thresholds. Demonstrates emergent behaviour using inertia, $F \propto ma$, and Hooke's law, and defines their failure boundaries.

9. **A Complete Characterization of the Electron within the SRE Framework: A Unified Account of Axiomatization, the Fine-Structure Constant, Molecular Computation, and Orbital Assignment**: Complete ontological account of the electron. The electron is characterized as the unique, stable and noise-resistant emergent fixed point of a sparse operator `R` acting on the coherent kernel of a binary self-organizing network, its ontological structure being the three-dimensional cube Q_3 (8 vertices, 12 edges, first Betti number 5, i.e. 12×5 = 60). The paper presents three independent topological derivation routes for the fine-structure constant $\alpha \approx 1/137.036$ (Möbius-ring graph-Laplacian spectral gap, the four-state intrinsic-spectrum frequency formula, and Möbius-parametrization arc-length perturbation), and a Tier-0 independent audit concludes that λ_0 is not an independent primitive and that the n=60 match is a discretization coincidence. It derives the double-cover four-state complex from Fork A axioms (A1–A4 + R1), establishes the observable dictionary $\alpha = \lambda_2/\lambda_{\max} = v/c$, and promotes the residual to the A5 bare coupling δ. It proves uniqueness, convergence and robustness of `R` on the coherent kernel (unified verification suite 10/10). It further extends to molecular computation (MCI spectral fingerprint, Q_3 ontological identification, MDS relation-inversion geometry) and to electron orbital-assignment rules (Pauli = cell uniqueness, $C_\ell = 4\ell + 2$, $C_n = 2n^2$, Hund = tension minimization).

10. **Complete Characterization of Nucleons within the SRE Framework: Unified Representation of Two-State Skeleton Inversion, Quark Speculation and Nuclear Reaction Validation**: Complete ontological account of the nucleon (v2.1). The nucleon is characterized as a stable composite object emerging on a three-strand Y-shaped coherent kernel, constrained by the open/closed two-state rule for dormant edges and by the closure-rebalancing rule, with ontological structure given by the triangular closure of three strands $Y_3 \ltimes \triangle_3$ (V=12, E=18, β_1=7, $|\mathrm{Aut}|$=36). It proposes three ontological conjectures concerning quarks, enters from neutron β-decay to derive the skeleton structural constraints T1–T3, redefines isospin Z_2 as the open/closed two-state rather than a flavour-multiplet flip, and uniquely retains candidate A via a three-step screening combined with a "transition–residue" framework. It validates the unified three-level signature chain from single-nucleon β-decay through compound-nucleus fission to macroscopic chain criticality (thermal-cross-section Pearson correlation +0.932), and generalizes to a shared-ring splicing law calibrated against external nuclear data (V=9k+3, E=15k+3, β_1=6k+1, $\lambda_2 = 2 - \sqrt{3}$). Version 1.5 further tightens the structural-layer conclusion (**A does not enter the structure, k does**; the

attachment point is not a degree of freedom, and inter-ring linkage is proven impossible to repair, so multi-group configurations should be absent at this layer), demotes antisymmetry from an energy term to a zero-parameter existence criterion L1 ($|n_n - n_p| \leq 1$, 8/8 on the 8-entry scorecard); Version 2.1 further adds a "contextual-limitations" boundary statement and extends the configuration-level test of the electronic-state objection (S1–S4 rerun, S2≅S1) and the disproof of "boson = transition quantum", and rewrites the electronic-state objection as a configuration-level problem, proving that open-band attachment is isomorphic to variant A (hence not a new structure, and already excluded quantitatively by the deuteron binding energy). The boundary at $A = 4$ is thereby split into "numerically non-computable, existentially computable", and the boundary is **relocated** from $A \geq 4$ to $A \geq 5$, explicitly demarcating the honest boundary that "nuclear binding for A ≥ 5 is not computable at this layer."

11. **Mass Origin in the SRE Framework: A Circulant-Operator Formulation on the Three-Ring $\mathbb{Z}_3$ Torsor**: A formalising account of the ontology of mass. Taking the SRE nucleon skeleton $Y_3 \ltimes \triangle_3$ (V=12, E=18, β_1=7, $|\mathrm{Aut}|$=36) as its only structural input, it first proves that the three shared rings form a **$\mathbb{Z}_3$-torsor** under the automorphism group (a cyclic order with no absolute origin); then, from "respecting the three-ring structure ⇒ commuting with the cyclic shift S ⇒ circulant matrix", it derives the unique admissible form of the mass operator $A = c_0\mathbb{1} + c_1 S + \bar{c}_1 S^2$, $\mathbb{Z}_3$-equivariance compressing 9 matrix entries into 3. The spectrum is a set of 120° samples of one cosine, $\lambda_j = c_0 + 2|c_1|\cos(\theta + 2\pi j/3)$, but the paper **explicitly corrects** the reading that "the cosine shape holds trivially for any three generations and is therefore a free reparameterisation rather than a constraint". "Opening" = choosing an origin for the DFT characters, a $\mathbb{Z}_3$ gauge degree of freedom rather than information. The invariant moduli space is **two-dimensional**, $(\eta,\ \delta \bmod 2\pi/3)$, and the Koide combination $Q = \frac{1}{3} + \frac{2}{3}\eta^2$ probes only the η direction (measured on the charged leptons, $\eta^2 = 1/2$ holds to 3.3×10^{-6}). The conclusion is registered as case 15 of the discrete-closure law G12: "three" and the functional form are given (discrete side, closed), while the value of η and the absolute scale c_0 are **not** (continuous side, requiring external input); the single real gap is the assignment rule.

12. **Positioning of Light and Electromagnetic Waves in the SRE Framework: the Involutive $\mathbb{Z}_2$ Sector, the Cohomological Ladder, and "One Emission as One Closed→Open Transition"**: A structural positioning paper for the light side. It places the project's existing light-side stipulations (the residuality axiom, the Möbius double cover, the involution $P = S^{n/2}$ on the weighted Möbius ladder M_n, the $\mathbb{Z}_2$ holonomy coupling δ_A) and the $\mathbb{Z}_3$ structure on the mass side under one framework for comparison; the whole paper needs only one structural input — the involution P on M_n — and gives four mutually independent, verifiable propositions: **the gear proposition** (light runs on $\mathbb{Z}_2$, mass on $\mathbb{Z}_3$; the two are adjacent rungs of the same cohomological ladder $|H^1(G;\mathbb{Z}_n)| = n^{\beta_1}$, the root of the difference being a group-theoretic prohibition $|\mathbb{Z}_2| = 2 < 3$); **the carrier proposition** (the light carrier is the odd eigenspace of an involution, with $P^2 = I$, membership in the automorphism group, commutation with the Laplacian, zero trace and spectrum exactly $\{\pm 1\}$; it is completely determined by the single integer n with **zero continuous free parameters**, so there is no place to put a source); **the source proposition** (the spectrum splits into a "sector-blind term + odd-sector constant term", in which only the w channel simultaneously satisfies "acts only on the odd sector" and "shifts every mode uniformly", so the source can inject only one scalar on the light side); and **the transition proposition** (one emission = one closed→open transition: the ledger difference carries the energy, with E and β_1 each decreasing by 1, while the spectral invariants carry the carrier; the two are orthogonal, so a chemical source and a nuclear source **share the same carrier**, differing only in the level at which the ledger is recorded). The conclusion is registered as case 18 of G12: it gives "why the gears are two and three", "why the source is a single scalar" and "why energy and carrier are orthogonal" (discrete side, closed), but **not** the energy scale of light (continuous side).

13. **Emergence of Time in the SRE Framework: A Saturation Law for the Rate of Change, Universality, and the Incompatibility of Two Internal Clocks**: A derived-quantity account of the ontology of time. Introducing no new free real number, it turns "time = a derived quantity obtained after an internal process has been designated as the clock" into a computable and falsifiable form: the per-step time is defined as $\Omega(n) = \langle \rho_{ij}/(1+\rho_{ij}) \rangle$ with $\rho_{ij} = \lambda d_{ij}/(|M^2|_{ij} + 1)$, and the emergent time as $T_{\mathrm{em}}(n) = \sum_{k<n} \Omega(k)$; every conclusion is synthesised from two **pre-existing zero-parameter laws** (the age-distribution law times the iid spectral law) and then compared against measurement. Seven verdicts: **saturation** (Ω rises monotonically to 1, with a remaining gap of 0.1540 at 999 steps); **the saturation law** $1 - \Omega = c \ln n/\sqrt{n}$ ($c = 0.694157$, maximum relative residual 1.589%, whereas a pure power law $n^{-1/2}$ gives 26.43% — the logarithm is the fingerprint of the harmonic tail — and the same law is predicted with zero free parameters to within 0.28%–1.65%); **asymptotic uniformity** (the relative step-to-step non-uniformity falls from 4.23×10^{-3} to 6.10×10^{-5}); **differential ageing** (the rewriting rate of a single element rises with its own age from 0.058931 to 0.957750, so an individual member's clock is not uniform); **universality in λ** (a 15-fold sweep does not move the limit, only the speed of approach); **a phase**

transition (squaring the denominator of the update rule moves the limit from 1.0000 to about 0.390, and the gap widens monotonically); and **the incompatibility of two clocks** (the memory horizon scales as $\tau_{\mathrm{mem}} \propto \sqrt{n}$ while the amount of change gives $T_{\mathrm{em}} \propto n$, their ratio diverging with n, so **no rescaling can make both clocks uniform simultaneously**). One separation across semantics is preserved: **there is structure, there is no memory** — the joint distribution of the state matrix rejects full independence (row-sum variance is 1.94–2.03 times the iid value), yet two different prefixes become indistinguishable after continued evolution (exact permutation $p \geq 0.3841$, none significant after Holm correction); precisely because the present state carries zero bits about its own past does "time must be emergent rather than read off" become a claim with content. Three honesty boundaries: the emergent time has **shape but no scale**; at attainable sizes "time is constant" holds only to about 22%; and two **guesses formed before measurement** have been refuted ($c \propto \lambda^{-1}$ measured at a slope of -0.577; "the variant kernel runs away into freezing" settles instead at a fixed point near 0.39), registered as open items rather than used as evidence. The conclusion is registered as case 16 of G12.

14. **A Plain-Language Version of the Emergence of Time in the SRE Framework**: A plain-language rewrite of the time-ontology paper Emergence of Time in the SRE Framework, aimed at general science-and-engineering readers. It preserves every core conclusion and datum, replacing project-internal jargon with everyday language and analogies; the seven verdicts (saturation law, universality, incompatibility of two clocks, etc.) and the structure-without-memory semantic separation are retold non-technically, and the three honest boundaries (notably emergent time has shape but no scale) are retained. Reproduction scripts are shared with the technical version (appendix SRE_Time_Emergence).

15. **Emergence of Positive and Negative Electric Charge in the SRE Framework: Magnitude, Sign, and Numerical Verification**: A paper closing the sign gap of charge. SRE expresses charge as the counting of topological knots (magnitude only); this paper rigorously gives Q(K)=sigma(K) middot deg(K) middot e, yielding electron Q=-e, positron Q=+e, proton Q=+e, neutron Q=0, antiproton Q=-e, and explaining the charge-mass decoupling (proton and electron share the same magnitude but differ in mass by about 1836 times). Two numerical tests confirm: the sign channel and the magnitude channel are mutually independent (section 10.1); the electron is forced by skeleton symmetry into the trivial holonomy class while the proton lands in the non-trivial class (section 10.2), giving the electron/proton sign opposition a structural origin.

> This suite inherits the axiomatic foundations of Part I. Several manuscripts only complete qualitative-semi-analytical mechanism construction; full quantitative benchmarking and large-multi-degree-of-freedom simulations are directions for subsequent research. Similarly, **this part does not answer the ultimate origin of causal differences**.

Theory of Hierarchical Dissipative Self-Organizing Binary Network Dynamics

Version: 1.1 (MDS methodological caveats and appendix added based on v1.0; all original axioms, equations and theorems preserved; v1.0 archived for historical traceability)

Framework Positioning Statement

The theory of hierarchical dissipative self-organizing binary-network dynamics presented herein constitutes **the underlying discrete network foundation for the self-emergent cosmic picture within the Status-Relational Entropy (SRE) Dynamics framework**.

This theory provides a computable set of discrete evolutionary axioms. Starting from simple binary spins and asynchronous activate-dormancy mechanisms, spacetime, gravitation, the speed of light and other physical quantities are not pre-implanted. A topologically coherent kernel emerges spontaneously through finite-size scaling phase transitions.

This underlying network undergoes two successive layers of abstract generalization:

1. Ontological refinement via the SRE v1.6 axiom suite, which defines general ontological concepts including causal nodes, pending-for-instantiation states, dissipation-compensation duality, reciprocal measurement, and multi-scale homomorphic mapping.
2. Further mathematical upgrade incorporating spectral-graph theory, the Baik-Ben Arous-Péché (BBP) spectral phase transition, and conformal-gauge transformations, yielding the SRE v6.0 cosmological framework. Metric geometry, variable-speed-of-light effects, gravitational dynamics, and gravitational-lensing step-jump observational predictions are derived within this higher-level framework.

Note that the present paper treats only the axioms, evolutionary equations, and scaling theorems of the underlying binary network. Cosmological-scale derivations and observational-data validation are the scope of the subsequent SRE v6.0 main paper and are not developed here.

I. Foundational Physical Axioms of the System

1. Strict Binary Constraint

The state of the system after the n-th pulse evolution is characterised by a real-symmetric square matrix $M_n \in \mathbb{R}^{n\times n}$. The entire evolutionary process contains no continuous-function truncation and no dissipative zero state. Matrix elements are strictly restricted to the binary-spin set:

$$\forall\ S_{ij} \in +1, -1, \quad S_{ij} = S_{ji}$$

The initial evolutionary baseline is a first-order non-zero single-point seed source:

$$M_1 = [1]$$

2. Asynchronous Binary Activation (Microscopic Point-by-Point Asynchronous-Activation Uncertainty)

The system adopts an asynchronous-update mechanism on a discrete topology. At each dimensional outward expansion $n \to n + 1$, every historical lattice point (i, j) of the old matrix determines its activation state in the current evolutionary step via an endogenous binary stochastic decision gate:

- Dormant state $\chi_{(i,j)} = 0$: the lattice point does not participate in the current step; information conduction is truncated. Algebraically, it contributes as multiplication by unity in the continued-product operation.
- Active state $\chi_{(i,j)} = 1$: the lattice point is normally activated, bringing its original ± 1 spin value into the evolution.

3. Dynamic Geodesic Field (Global Dynamic Geodesic-Depth Rheology)

Topological relations of the system are defined entirely by algebraic topological adjacency among nodes. The cumulative topological step length from any historical lattice point (i, j) to the current evolutionary frontier grows synchronously as system order expands, manifesting as global dynamic redshift rheology:

$$d_n(i,j) = n - \max(i,j)$$

II. Endogenous Dynamical Definition of Probability p

The dormancy probability $p_{ij}^{(n)}$ for lattice point (i, j) at step n is determined endogenously by the local frustration tension at that location together with global scale factors.

1. Local Frustration Energy

The coherent-tension inner product for lattice point (i, j) on the matrix intersection manifold is defined:

$$E_{\text{local}}(i,j) = \left| \sum_{k=1}^{n} S_{ik} \cdot S_{kj} \right|$$

Smaller absolute values indicate stronger cancellation of positive-negative polarities, corresponding to higher local geometric frustration; such links tend to degenerate into dormant edge states.

2. Adaptive Energy-Level Mapping Equation

Combined with cumulative topological step-length $d_n(i, j)$, the lattice-point dormancy probability is rigorously defined:

$$p_{ij}^{(n)} = \Pr\big(\chi_{(i,j)} = 0\big) = 1 - \frac{1}{1 + \lambda \cdot \dfrac{n - \max(i,j)}{E_{\text{local}}(i,j) + 1}}$$

where $\lambda \in \mathbb{R}^+$ denotes the endogenous coupling constant of the system.

Physical corollaries of this definition:

- Microscopic newborn layer: $n - \max(i, j) \to 0$. Newborn nodes have step-length approaching the evolutionary frontier, giving $p_{ij}^{(n)} \to 0$. Microscopic newborn units tend to deterministic activation updates, guaranteeing local-manifold geometric rigidity.
- Macroscopic ancient layer: $n - \max(i, j) \gg 0$. As evolution proceeds, cumulative topological step-lengths for ancient nodes increase monotonically, and dormancy probability rises spontaneously. Large-scale domains enter dormancy; long-range coherence is diluted by spontaneous system-internal dissipation.

III. Core Evolutionary Equation and Global Adaptive Negative-Feedback Damping

When the system undergoes dimensional expansion triggered by the $(n + 1)$-th pulse, each matrix element $S_{i,n+1}$ on the new boundary propagates following a point-wise nonlinear continued-product feedback equation:

$$S_{i,n+1} = \prod_{j=1}^{n} \Big[\chi_{(i,j)} \cdot S_{ij} + \big(1 - \chi_{(i,j)}\big) \Big], \quad i = 1, 2, \ldots, n$$

The bottom-rightmost diagonal matrix element acts as an endogenous energy-balance regulating valve for the system. It carries no stochastic gate and strictly enforces global algebraic adaptive negative-feedback damping, stabilising the global net-spin pool at each evolutionary step:

$$S_{n+1,n+1} = \begin{cases} -1, & \displaystyle\sum_{x=1}^{n} \sum_{y=1}^{n} S_{xy} \geq 0 \\ +1, & \displaystyle\sum_{x=1}^{n} \sum_{y=1}^{n} S_{xy} < 0 \end{cases}$$

IV. Macroscopic Dynamical Emergence and Finite-Size-Scaling Asymptotic-Convergence Theorems

Emergence of system spatial dimension is not an instantaneous observation at some discrete step count (e.g., $N = 250$). Instead it represents asymptotic phase-transition behaviour under the thermodynamic limit via finite-size scaling.

Theorem 1: Statistical Condensation of Local Stable Manifolds and Order-Parameter Convergence

To exclude pseudo-emergence caused by finite-size effects, we take microscopic-core surviving units (the first k sub-matrix rows/columns, with $k = \lfloor 0.2N \rfloor$) and define the topological-coherence order parameter:

$$\Phi(N) = 1 - \frac{\mathrm{Var}\big(\mathrm{Tr}(M_k)\big)}{k^2}$$

Numerical and analytical derivations show that as total pulse steps go to infinity $N \to \infty$, even though macroscopic boundaries exhibit large uncertainties from asynchronous activation, the order-parameter for the ancient core region obeys strict asymptotic boundedness:

$$\lim_{N\to\infty} \Phi(N) = \Phi_0 > 0$$

This non-zero constant Φ_0 rigorously demonstrates that condensation of the system's local stable manifold is not an accidental finite-step fluctuation. It is a statistically asymptotically stable core established by feedback suppression from the high-dimensional renormalisation pool.
This asymptotic-convergence behaviour is directly observable in numerical simulations; see the $\Phi(N)$ convergence plots in the middle column of Figure 1.

Theorem 2: Coherence-Length Decay and Hierarchical Fragmentation under Scaling Cascades

Spatial dimension and geometric curvature within the lattice obey a progressive-dissipation rule: "each outward layer brings one degree of coherence weakening". Define the two-point sign-correlation decay function against topological step-size x:

$$G(x) = \langle S_i \cdot S_{i+x} \rangle \propto e^{-x/\xi}$$

where ξ denotes the effective coherent correlation length.

During finite-size-scaling cascades, rheology of coherence length with total system size N satisfies:

$$\lim_{N\to\infty} \frac{\xi(N)}{N} = 0$$

This limit rigorously proves that three structural regimes arising from spontaneous system phase transition are macroscopically distinguishable:

1. **Microscopic core layer** ($x \leq \xi$): dormancy probability $p \approx 0$, the system maintains very high topological coherence and rigidity. Observed via Multidimensional Scaling (MDS), high-dimensional eigenvalues collapse toward zero, and positive-definite three-dimensional manifolds together with local curvature emerge spontaneously within this layer.
2. **Mesoscopic intermediate layer** ($\xi < x \sim 2\xi$): crossing the correlation-length scale triggers avalanche-like cascade amplification within continued-product causal chains. Manifold resilience against stochastic-dormancy perturbations decays exponentially; geometrically rigid lattice structures undergo nonlinear bending and bifurcation.
3. **Macroscopic thermal-dissipation layer** ($x \gg \xi$): long-range coherence is fully diluted and decoupled by abundant dormant points. Original system structure disintegrates and returns to an isotropic chaotic state.

Simulation results are shown in Figure 1; the MDS three-dimensional manifold on the right-hand side visualises this three-layer hierarchical structure.




Figure 1. Unified verification simulation for hierarchical-dissipation dynamics. Three independent random seeds (1111, 2222, 3333), evolution steps $(N = 300)$, coupling parameter $\lambda = 0.8$.
Left column: matrix heatmaps of the network. Middle column: evolution curves for topological-coherence order-parameter $\Phi(N)$; red dashed lines mark asymptotic limit Φ_0. Right column: emergent three-dimensional manifold obtained via MDS embedding. Simulations verify the asymptotic-convergence behaviour of Theorem 1 and visualise the three-layer hierarchical structure predicted by Theorem 2: high-coherence microscopic core, mesoscopic intermediate zone, and macroscopic dissipative outer layer.

V. Theoretical Conclusion

This paper establishes a finite-size-scaling asymptotic evolutionary model for non-equilibrium binary-network self-organised phase transitions.

The theory demonstrates that matter (damping of ± 1 spins under polar frustration), space (dynamic distance defined via topological geodesics), gravitation (relative path contraction-convergence and spatial evaporative-dissipation arising from local dormancy), and three-dimensional space itself require no hard-coded presupposition at the underlying level.

Taking microscopic point-wise binary activate-dormancy uncertainty as its underlying engine and global algebraic continued-multiplication as causal bonds, the system can spontaneously condense self-organised spacetime on finite-size-scaling phase-transition lines without invoking external continuous probability waves, as system size approaches the thermodynamic limit. The microscopic core yields geometrically positive-definite stable structure, while outward hierarchical expansion is accompanied by progressive dissipation. This asymptotic phase-transition structure, transitioning from non-integer dimensions toward integer three-dimensionality, constitutes the intrinsic macroscopic-thermodynamic limiting phase of this discrete computational model.

> **Methodological Supplementary Remark (added in v1.1)**
> Multidimensional Scaling (MDS) serves only as an offline post-processing numerical prototype. MDS reads static snapshots of the binary network and embeds pairwise topological relations into three-dimensional Euclidean space. Spatial distances within this embedded manifold are emergent representations of node relational-profile similarity; **Euclidean geometric distance does not pre-exist within the underlying binary-network ontology**. This static-snapshot embedding is conceptually distinct from the event-driven, incremental-instantiation mechanism characterising the full SRE-Dynamics physical-rendering-layer. Embedding artefacts appear in strongly-dissipative outer regimes beyond the coherence length ξ, where the validity of low-dimensional geometric descriptions degrades.

Appendix A Conceptual Distinctions and Applicable Boundaries for MDS Numerical Embedding (added in v1.1)

A.1 Conceptual Distinction between MDS Simulation Prototype and the SRE Physical-Rendering-Layer

Within the accompanying simulation script `sim_P.py`, Multidimensional Scaling (MDS) operates as **snapshot-oriented offline post-processing**: after full network evolution to a specified step count, the complete final-state matrix is taken, and pairwise topological relations for all nodes are projected in a single pass onto three-dimensional Euclidean coordinates.

This numerical workflow possesses two key features:

1. It is not incremental-difference rendering: only complete snapshots of all node-topology relations are input; only differential changes per evolutionary step are not rendered.
2. It performs homomorphic dimensional reduction rather than isomorphic copying: the underlying object is high-dimensional discrete binary topology possessing no Euclidean coordinates. MDS executes many-to-one coarse-graining and information compression. Numerous microscopic degrees-of-freedom including underlying spin signs, activate-dormancy stochastic gates, and local frustration are not fully preserved within three-dimensional output.

By contrast, the **physical-rendering-layer** defined in the SRE v1.6 ontological framework is an event-driven incremental-instantiation system:
The underlying causal network undergoes continuous asynchronous-link fluctuations. The vast majority of transient perturbations fail instantiation-cost thresholds and remain in the pending-for-instantiation state without projection onto spacetime. **Only causal relations completing one full round of reciprocal measurement together with dissipation-compensation settlement update distance- and time-bookkeeping records within the rendering layer.**

> MDS simulation: static characterisation of an already-evolved system, answering what three-dimensional manifold would arise via homomorphic projection given statistically stable topology.
> SRE physical-rendering-layer: real-time incremental instantiation triggered by reciprocal-measurement events; the ontological mechanism corresponding to real-world physical cosmology. The two must not be treated as directly equivalent.

A.2 Intuitive Rules and Failure Conditions for MDS Manifold Distances

Geometric distances output by MDS embedding follow this statistical tendency:

> If two nodes exhibit highly similar patterns of causal association (relational-profiles) with all remaining network nodes, their Euclidean distances within the three-dimensional embedded manifold tend to be small.

This property corresponds to SRE ontological intuition: macroscopic spatial distance characterises the degree of coherence degradation between events. More similar relational-profiles imply lower mutual information dissipation, smaller topological-compensation cost, and correspondingly smaller macroscopic metric distance.

This rule-of-thumb has strict applicability boundaries:

- **Microscopic core region ($x \leq \xi$, high order-parameter $\Phi(N)$)**: high network topological coherence yields good MDS dimensional-reduction fidelity; the relational-profile versus manifold-distance correspondence holds.
- **Mesoscopic / macroscopic strong-dissipation regimes ($x > \xi$)**: abundant stochastic link dormancy creates highly disordered topology; no low-dimensional geometry can perfectly represent all high-dimensional relations. MDS produces minimal-loss approximate fits and embedding artefacts may occur: nodes with substantially different underlying relational-profiles may lie accidentally close geometrically, while minor topological differences may be artificially amplified during

dimensional compression. Under these conditions low-dimensional geometric descriptions degrade, and the relational-profile / manifold-distance correspondence becomes unreliable.

A.3 Correspondence with the Full SRE System

Phenomena within the binary-network simulation can be mapped in parallel to SRE ontological concepts, though direct identification is not permitted:

1. Link dormancy $\chi_{ij} = 0$ ⇔ irreversible information dissipation (microscopic origin of the topological-dissipation tensor $\hat{\mathcal{D}}_{ij}$ in cosmological papers).
2. MDS dimensional-reduction embedding ⇔ numerical demonstration prototype for multi-scale homomorphic mapping.
3. Core / meso-scale / outer-layer tripartite division ⇔ multi-scale rigid coherence-truncation boundary filtering mechanism.
4. Finite-size-scaling phase transition ⇔ underlying discrete prototype for the BBP spectral phase transition ($z_{\mathrm{crit}} = 4.1605$, 2D-holographic ↔ 4D-spacetime switching).

> Note: the binary network itself contains no built-in spacetime, gravitational constant, or speed of light. All such physical quantities are macroscopic statistical effects emerging inside the rendering-layer after homomorphic-mapping filtering.

> In summary, this binary network constitutes the discrete computational substrate for SRE's self-emergent cosmic picture. Full cosmological physical predictions require ontological abstraction plus spectral-matrix mathematical upgrades; refer to the accompanying SRE v6.0 paper.

A.4 Version-Change Notes

> v1.1 update record:
>
> 1. Added brief MDS methodological warning at the end of Chapter 5 Theoretical Conclusion.
> 2. Added Appendix A giving full conceptual distinction between the MDS numerical prototype and the SRE physical-rendering-layer, together with applicable-boundary discussion.
> 3. The v1.0 manuscript is fully preserved in the Zenodo archive for historical traceability and is not overwritten.

Emergence of Multidimensional Spacetime and Dynamical Gravity via Regularized Causal-Information Networks

Version: 6.2-rev (incorporating ontological corrigendum; updated numerical simulation results, distinguishing historical reference value from Bootstrap statistical simulation outputs)

Abstract

The ΛCDM standard-cosmology framework encounters significant observational tension at high redshift $z > 5$. The James-Webb Space Telescope (JWST) has observed massive mature galaxies already assembled within the first 500 Myr of cosmic time; under static-G_0 structure-formation scenarios, hierarchical-growth timescales are insufficient to produce such objects.

Built upon the State-Relation-Entropy (SRE) dynamical axiom-system (v1.6), this work establishes the fully background-independent SRE cosmic-gravity framework (v6.2-rev). Spacetime topology, gravitational coupling strength, and the speed-of-light are not primitive axiomatic inputs; instead they emerge as macroscopic gauge effects of a decentralised bidirectional Möbius causal-information network.

This manuscript corrects the residual ontological dependency on extrinsic redshift-difference coordinates $|z_i - z_j|$ present in the v6.1 draft section 2.3. Macroscopic metric distance is directly defined as the topological-compensation cost incurred by the causal network to cancel irreversible information dissipation, fully realising the SRE-v1.6 ontology: *distance is a book-keeping product of dissipation-compensation duality*.

The dynamical compression coefficient $\alpha_{0,\mathrm{dynamic}}$ abandons the hard-fitted constant used in v5.2; it is analytically derived from matrix spectral resonance within each sliding observational horizon. Variable emergent effective-speed-of-light c_{eff} is implemented at network-routing level; conformal-gauge covariance rescales the emergent metric tensor while preserving local Lorentz invariance for measured light-speed.

The Baik-Ben-Arous-Péché (BBP) spectral-rank phase transition governs switching between the 2D holographic-projection phase and the 4D unlocked-spacetime phase. The early-version v6.1 provided a priori theoretical reference critical redshift $z_{\mathrm{crit}} = 4.1605$. Using 1500 non-parametric Bootstrap resampling realisations together with the SDSS/eBOSS spectroscopic dataset, **the statistically-simulated transition location yields $z^* = 3.13$ (95% confidence bounds are subject to model- and dataset-dependent uncertainties)**. This phase transition produces a systematic factor-2-to-4 jump in gravitational-lensing deflection without postulating pre-existing continuous Riemannian geometry. Within the primordial dense-universe regime $z \geq z^*$, baryonic-gas cooling rates are substantially enhanced, amplifying accretion efficiencies without altering cosmic thermal ages, and naturally alleviating the JWST early-massive-galaxy formation puzzle.

Statistical simulations are performed on 29890 SDSS/eBOSS spectroscopic QSO spectra. Chiral-twist correction terms are presented, providing observationally-falsifiable imprints for JWST and the Roman Space Telescope.

Keywords: State-Relation Entropy; dissipation-compensation duality; causal-information network; BBP spectral-rank phase transition; variable-emergent speed-of-light gravity; holographic dimensional crossover; gravitational lensing; Bootstrap statistical simulation

1 Introduction

Modern observational cosmology pushes ΛCDM beyond its domain-of-validity. DESI-DR1 spectroscopy and deep JWST imaging reveal two intertwined paradoxes: apparent dark-energy tension at high cosmic horizons, and the emergence of highly-evolved $M > 10^{10}\,M_{\odot}$ galaxies within the first 500 Myr after the Big-Bang. Under static-G_0 assumptions, standard hierarchical-accretion models lack sufficient causal time to assemble such massive objects from primordial gas seeds.

Earlier incarnations of SRE cosmic-gravity (v5.2 and prior) contained two epistemological weaknesses:
① the network-compression limit $\alpha_0 = 0.12$ was a manually hard-coded fitting constant;
② heuristic piece-wise patches were used in visualisation to artificially enforce a phase-transition boundary.
Moreover, the original v6.1 draft section 2.3 retained the ontological flaw of adopting redshift-coordinate differences $|z_i - z_j|$ as a distance substrate, in violation of SRE-v1.6 ontology:

> *Spatial distance is not a-priori given coordinates; it arises solely as a book-keeping outcome of mutual-measurement driven dissipation-compensation duality.*

This paper carries out four tiers of improvements:

1. Using spectral-graph theory and Random-Matrix-Theory (RMT), eliminate all manually hard-coded cosmological constants;
2. Adopt the corrigendum ontological fix: discard $|z_i - z_j|$ as distance substrate, and construct macroscopic metrics directly from topological dissipation-compensation operators;
3. Introduce non-parametric Bootstrap Monte-Carlo resampling, perform statistical simulation with SDSS/eBOSS spectroscopic data to obtain the simulated transition redshift z^*; demote $z_{\text{crit}} = 4.1605$ to a v6.1 historical-theoretical reference value and remove it as a rigid model prediction of the present revision;
4. Carry out self-consistency statistical validation against SDSS/eBOSS observational catalogues and establish observationally-falsifiable cosmological predictions.

The underlying ontology strictly inherits the SRE-v1.6 axiom suite. Spacetime, matter properties and physical constants are homomorphic emergent mappings after multi-scale rigid-coherence truncation of the causal-information network. The Planck scale acts as an **emergent ultraviolet ontology threshold (instance-realisation cost barrier)**, not a fundamental pixel-granularity of the underlying network. The vast majority of underlying causal interactions remain in *uninstantiated state*; only after completing full dissipation-compensation book-keeping do they project onto the physical-rendering layer.

2 Axiomatic Mathematical Formulation and Ontological Anchoring

2.1 Ontology of the Causal-Information Network (aligned with SRE axiom-suite v1.6)

The causal-information-network provides a cosmological realisation for the SRE-v1.6 primitives: causal-nodes, mutual-measurements, and uninstantiated states.

1. **Causal node** V: Ontologically defined as quantum-evidence events occupying local sectors of Planck-phase-space H_{Planck}. Each node corresponds to a non-local quantum-measurement event that enforces reduction of informational relations.
2. **Network edge** E: Network links are analogous to area-quantum elements of loop-quantum-gravity spin-networks; connectivity topology implements parallel-transport of the Ashtekar-Barbero connection.
3. **Information-packet routing**: Packet propagation on the graph corresponds to topological-geodesic flow within MERA-style tensor-networks. Spacetime geometry is a holographic manifestation of entanglement-entropy boundaries of the network.

> There exists no pre-existing continuous Riemannian manifold at the fundamental level. Spacetime dimensionality, gravitational coupling and light-speed all emerge macroscopically from topological-connectivity densities of the discrete causal-network. Photons correspond to high-frequency information-packets mediated by Möbius-topology residuals, matching exactly the SRE-v1.6 primitive *light-residual* Ψ_{light}.

2.2 Topological-Dissipation Tensor and Compensation Operator (core corrigendum)

> Discard the v6.1-draft practice of building distances from $|z_i - z_j|$. Distances are entirely founded on dissipation-compensation duality, complying with SRE-v1.6 ontology: distance quantifies the degree of topological-residual coherent-degradation as a book-keeping consequence of dissipation-compensation accounting.

Define the **topological-dissipation tensor** $\hat{\mathcal{D}}_{ij}$, which characterises intrinsic information-loss operators for quantum-evidence events (i, j), constrained by observational measurement-entropy bounds σ_z:

$$\hat{\mathcal{D}}_{ij} = \ln\left(1 + \frac{\sigma_{z,i} \cdot \sigma_{z,j}}{\epsilon_{\text{mach}}}\right)$$

ϵ_{mach} denotes machine floating-point epsilon.

Define the dynamical topological-compensation-operator $\hat{\mathcal{C}}_{\text{compensation}}(\alpha_{0,\text{dynamic}})$, representing routing-computational overhead invoked by the network to counteract information-dissipation and maintain numerical-matrix stability:

$$\hat{\mathcal{C}} = \alpha_{0,\text{dynamic}}^{-1} \cdot \sin^2\left(\pi\alpha_{0,\text{dynamic}} \cdot \hat{\mathcal{D}}_{ij}\right)$$

The macroscopic squared-metric distance is directly defined as the trace inner-product of dissipation-tensor and compensation-operator:

$$R_{ij}^2 \equiv \mathrm{Tr}\left(\hat{\mathcal{D}}_{ij} \cdot \hat{\mathcal{C}}_{\mathrm{compensation}}(\alpha_{0,\mathrm{dynamic}})\right) \cdot \exp\left(-\gamma \cdot \mu_{\mathrm{loss}}\right)$$

- $\gamma = 0.0585$: network handshake-latency coefficient, analytically derived from Bekenstein information-smoothing-bound cost $1/(2\pi e)$.
- μ_{loss}: local-mean information-loss weight.

> **Ontological-paradigm shift**: spatial "distance" is not given a-priori. As information-dissipation $\hat{\mathcal{D}}_{ij}$ between nodes increases, the network must allocate exponentially growing routing resources to suppress spectral divergence. This internal structural overhead of the decentralised network counteracting information-loss is what observers interpret as spatial separation. This strictly realises SRE-v1.6 ontology: mutual-measurement / dissipation-compensation book-keeping precedes metric-spacetime generation.

2.3 Topological-Stiffness Weights and Baryonic-Centroid Redshift

Topological-stiffness weight $\mathcal{W}_{ij}$ encodes macroscopic mass-equivalences within the relational manifold, derived purely from spectroscopic observational measurement-entropy:

$$\mathcal{W}_{ij} = \sqrt{\mathcal{C}_i \cdot \mathcal{C}_j} = [\left(1 + \ln\left(1 + \max(\sigma_{z,i}, \epsilon_{\mathrm{mach}})\right)\right) \cdot \left(1 + \ln\left(1 + \max(\sigma_{z,j}, \epsilon_{\mathrm{mach}})\right)\right)]^{-1/2}$$

Given a local slice-subgraph V_{slice} containing N events, define the gravitationally-weighted **baryonic-centroid redshift** μ:

$$\mu = \frac{\sum_{i=1}^{N} \mathcal{C}_i \cdot z_i}{\sum_{i=1}^{N} \mathcal{C}_i}$$

Higher-order loop-perturbation fields couple to the centroid redshift:

$$\xi(z) \equiv \xi(\mu) = 0.08 \cdot \exp(0.15 \cdot \mu)$$

2.4 Spectral-Resonance derivation of dynamical compression coefficient $\alpha_{0,\mathrm{dynamic}}$

$\alpha_{0,\mathrm{dynamic}}$ is no longer an exogenous input parameter. It is analytically obtained from Fourier-spectral resonance of the double-centred network-matrix within each sliding observational causal-horizon of width $\Delta z = z_{\max} - z_{\min}$. Matrix-stability conditions enforce matching of chiral-sine modes to the first resonance valley to avoid numerical dissipation:

$$\alpha_{0,\mathrm{dynamic}} = \frac{\theta_{\mathrm{conformal}}}{\Delta z + \epsilon_{\mathrm{mach}}}$$

Where the conformal-geometric index $\theta_{\mathrm{conformal}} \approx 0.82798$ comes from辛-eigenvalue integration for maximum-packing fractions on complex-hyperbolic Möbius manifolds:

$$\theta_{\mathrm{conformal}} = \frac{1}{\pi} \int_0^1 \frac{\ln(1+x^2)}{x} \mathrm{d}x + \frac{1}{2e} \approx 0.82798$$

For SDSS sliding-window slices with $\Delta z \approx 0.03925$, we obtain $\alpha_{0,\mathrm{dynamic}} = 21.09 \pm 0.34$. This demonstrates compression-limits emerge purely as mathematical consequences of slice-geometry rather than manual tuning.

3 Variable-Emergent Speed-of-Light (VSL) and Conformal-Gauge Covariance (aligned with SRE-v1.6)

SRE-v1.6 axioms state that the speed-of-light c is an **emergent causal-propagation upper-bound**. In the high-dissipation primordial universe the effective throughput c_{eff} drops; conformal-gauge transformations preserve local measured Lorentz-invariance. This section presents the full mathematical formulation.

3.1 Information-propagation emergent effective-speed-of-light

$c_{\mathrm{eff}}(\mu)$ is defined as the maximum packet-routing bandwidth upper-bound on the adjacency network. In the dense primordial universe high topological-winding creates impedance and increases transmission latency:

$$c_{\text{eff}}(\mu) = c_0 \cdot \Phi_{\text{net}}(\mu) = c_0 \cdot \left[1 - \kappa \ln\left(1 + \frac{\rho_{\text{info}}}{\rho_{\text{critical}}}\right)\right]$$

- c_0: sparse-near-field baseline propagation-speed.
- ρ_{info}: local relational-link information-density.
- Topological-coupling index $\kappa = \dfrac{1}{\ln 2 \cdot \pi^2} \approx 0.1462$, originating from topological-complementary-cut-set impedances of Möbius cross-nodes.

> Physical picture: the universal fundamental speed-constant itself is unchanged. Network computational resources are diverted to dissipation-compensation tasks, lowering packet-routing throughput.

3.2 Conformal-scaling factor and local Lorentz-invariance

To guarantee gauge-covariance, changes in link-density simultaneously rescale the emergent metric tensor $g_{\mu\nu}$ and effective propagation-speed:

$$\tilde{g}_{\mu\nu} = \Omega^2(\alpha_{0,\text{dynamic}})\, g_{\mu\nu}, \qquad \tilde{c}_{\text{eff}} = \Omega(\alpha_{0,\text{dynamic}})\, c_{\text{eff}}$$

The line-integral of the conformal-scalar-field over relational-moduli-space:

$$I(z) = -\frac{\gamma}{4} \int_{z_{\min}}^{z_{\max}} \alpha_{0,\text{dynamic}}(z)\, \mathrm{d}z$$

Substituting the analytic expression for $\alpha_{0,\text{dynamic}}$ yields the conformal multiplier:

$$\Omega(\alpha_{0,\text{dynamic}}) = \exp(I(z)) = \left(\frac{\Delta z}{\theta_{\text{conformal}}}\right)^{-\gamma/4}$$

Algebraic cancellation preserves the local line-element:

$$\mathrm{d}s^2 = \tilde{g}_{\mu\nu}\mathrm{d}x^\mu \mathrm{d}x^\nu = g_{00}c_0^2\mathrm{d}t^2 + \Omega^2 g_{ij}\mathrm{d}x^i\mathrm{d}x^j$$

Therefore, even as c_{eff} evolves cosmologically, **locally-measured observer light-speed remains c_0, satisfying Lorentz-invariance**, consistent with qualitative SRE-v1.6 predictions.

4 Random-Matrix-Theory and BBP Spectral-Rank Phase-Transition: 2D-Holographic ↔ 4D-Unlocked-Spacetime

Effective rendered-spacetime dimensionality is determined by the eigenvalue-spectrum of the stabilised association matrix $B_{\text{stabilized}}$. Effective rank counts eigenvalues exceeding the Tracy-Widom statistical-bulk boundary:

$$\text{Rank}(z) = \sum \left(\text{eigvals}(B_{\text{stabilized}}) > \epsilon_{\text{adaptive}}\right)$$

Adaptive threshold:

$$\epsilon_{\text{adaptive}} = \epsilon_{\text{mach}} \cdot \frac{\ln\left(1 + \|B_{\text{stabilized}}\|_1 / N\right)}{2.5} \cdot 1.2$$

N counts Planck-event counters inside the past-light-cone; within numerical pipelines it corresponds to valid-sample rows of spectroscopic slices.

The dimensional-fluctuation-field $\Psi_{\text{fluct}}(z)$ obeys a modified topological Ginzburg-Landau equation describing the BBP spectral-rank phase-transition, with Planck-scale boundary-conditions:

$$\frac{\partial^2 \Psi_{\text{fluct}}}{\partial z^2} + \beta(z)\Psi_{\text{fluct}} - \eta\Psi_{\text{fluct}}^3 = 0$$

$$\Psi_{\text{fluct}}(z^*) = 0, \qquad \left.\frac{\partial \Psi_{\text{fluct}}}{\partial z}\right|_{z\to\infty} = \sqrt{\frac{\beta_0}{\eta}}$$

Microscopically dimensionality oscillates at Planck-frequency $10^{43}\,\mathrm{Hz}$. Astronomical observing-instruments have integration-times $\Delta t \gg \tau_P$; environmentally-induced decoherence washes out fast oscillations and observers detect the smooth expectation-value envelope:

$$\langle \mathrm{Rank}(z) \rangle = \int_0^{\Delta t} \Psi_{\mathrm{fluct}}(t)\,\mathrm{d}t$$

> Two distinct phases separated by the statistically-simulated transition redshift z^*:
>
> 1. **Late-time universe** $z < z^*$, $\mathrm{Rank} = 2$ **(2D-holographic-projection phase)**: the network remains in single-handed Möbius topology; compensation-operator collapses onto a single routing-layer.
> 2. **Primordial dense universe** $z \geq z^*$, $\mathrm{Rank} = 4$ **(4D-unlocked-spacetime phase)**: BBP spectral-rank phase-transition triggers; single-handed Möbius topology splits into bidirectional-chiral two-layer network. The compensation-operator separates into two independent eigen-branches: time-layer compensation and space-layer compensation:

$$\mathrm{Tr}(\hat{\mathcal{C}}_{\mathrm{time}}) = \mathrm{Tr}(\hat{\mathcal{C}}_{\mathrm{space}}) = \alpha_{0,\mathrm{dynamic}}^{-1} \cdot \sin^2\left(\pi \alpha_{0,\mathrm{dynamic}} \hat{\mathcal{D}}_{ij}\right)$$

> z^* denotes the simulated statistical transition-redshift; $z_{\mathrm{crit}} = 4.1605$ is the historical-theoretical reference value from v6.1. In SRE-v1.6 language: eigenvalues crossing the threshold mean causal-interactions satisfy dissipation-compensation budgets and transition from uninstantiated state into physical-rendering-layer.

Figure 1: Cosmological Phase Transition — Ensemble ⟨Rank⟩ & ⟨G_{eff}⟩




Figure 1 BBP spectral-rank cosmological phase-transition. Left-hand vertical axis (red): ensemble-mean normalised effective gravitational coupling $\langle G_{\mathrm{eff}}/G_0 \rangle$ from Bootstrap realisations. Right-hand vertical axis (blue dashed): ensemble-mean emergent spacetime rank $\langle \mathrm{Rank}(z) \rangle$. Solid orange vertical line marks the statistically-simulated phase-transition $z^* = 3.13$; purple dashed vertical line shows historical-theoretical reference $z_{\mathrm{crit}} = 4.1605$. Below z^* the system resides in the two-dimensional holographic phase; above z^* four-dimensional spacetime unlocks, accompanied by oscillatory behaviour of G_{eff} induced by chiral-manifold corrections. Shaded bands denote Bootstrap-derived 95 % statistical-confidence intervals.

5 Causally-Emergent Gravity: Thermodynamic Effect of Dissipation-Gradients

Gravity is not a fundamental-field but a statistical-thermodynamic consequence of local information-dissipation-gradients. Matter condensation elevates the local dissipation-tensor, and the network generates inward compensation-flows for matrix equilibrium. Discarding pre-supposed Riemannian backgrounds, SRE-v6.2-rev obtains gravitational-acceleration from logarithmic-gradients of relational-metrics conditional upon the current network rank-state:

$$a_{\mathrm{SRE}}(r,z) = \begin{cases} -\dfrac{\alpha_{\mathrm{scale}} \cdot \mathcal{W}_{ij}}{r} - \dfrac{\gamma c_{\mathrm{eff}}(z)^2}{4} & \mathrm{Rank} = 2,\ z < z^* \\ -\dfrac{2 \cdot \alpha_{\mathrm{scale}} \cdot \mathcal{W}_{ij}}{r^2} - \dfrac{\gamma c_{\mathrm{eff}}(z)^2}{4} + \Gamma_{\mathrm{chiral}}(r) & \mathrm{Rank} = 4,\ z \geq z^* \end{cases}$$

Chiral-gravitational-correction originates from genus-1 manifold Dirac-operator loop-corrections:

$$\Gamma_{\mathrm{chiral}}(r) = \xi(z) \cdot \frac{\sin\left(\pi \alpha_{0,\mathrm{dynamic}} \cdot 2\mu\right)}{r^2 \cdot \ln(r/\ell_P)}$$

ℓ_P is the CODATA Planck-length, the emergent-ontology ultraviolet-threshold defined in SRE-v1.6.

- $\mathrm{Rank} = 2$ holographic-phase: gravity manifests long-range logarithmic-potential $\propto 1/r$.
- $\mathrm{Rank} = 4$ unlocked-phase: recovers inverse-square-law $(1/r)^2$ behaviour; G_{eff} undergoes smooth oscillations within model-allowed bounds.

Baryonic-cooling-boost factor (explanation for JWST early-massive-galaxy puzzle)

$$Cooling_Boost = \left(\frac{G_{\mathrm{eff}}}{G_0}\right)^2$$

Within the primordial dense-universe interval $z \geq z^*$, baryonic-molecular cooling-rates receive enhancements. Without altering cosmic thermal ages, the Eddington-accretion-limit is amplified, allowing gas to collapse into super-massive galaxies over short cosmic timescales.




Figure 2 Primordial causal-core mass-accumulation comparison. Black dashed curve: standard Λ-CDM accretion under static G_0. Red solid curve: SRE enhanced-accretion under ensemble-mean dynamic $\langle G_{\mathrm{eff}} \rangle$ within the high-dimensional unlocked-phase. Light-yellow shaded band marks the observed-mass boundary of mature JWST galaxies at $z > 5$. Horizontal axis is cosmic lookback-time in Gyr. Pink shaded region denotes 95 % Bootstrap confidence-interval for SRE-model mass-output. The SRE cooling-boost effect reaches observed galactic-core masses within allowed cosmic time.

6 Gravitational-Lensing Shear Formula and the 2-to-4 Systematic Jump (after corrigendum)

Photons propagate as high-frequency information-packets across the causal-network. When passing a massive causal-core at impact-parameter b, macroscopic deflection-angles are determined by the number of active compensation-channels.

1. $z < z^*,\ \mathrm{Rank} = 2$**, two-dimensional holographic-degenerate lensing**
   Only the time-delay compensation-channel is active:

$$\theta_{\text{macro}}^{(2D)} = \frac{2 \cdot \mathcal{W}_{ij}}{b}$$

2. $z \geq z^*$**, four-dimensional unlocked-lensing**
   The BBP spectral-rank phase-transition opens bidirectional-two-layer networks; time-layer and space-layer compensation-flows operate in-parallel and add linearly:

$$\theta_{\text{macro}}^{(4D)} = \theta_{\text{time}} + \theta_{\text{space}} = \frac{2 \cdot \mathcal{W}_{ij}}{b} + \frac{2 \cdot \mathcal{W}_{ij}}{b} = \frac{4 \cdot \mathcal{W}_{ij}}{b} \cdot \left[1 + \Lambda_{\text{twist}}(b)\right]$$

Chiral-twist correction produces observable anisotropic polarisation imprints testable by JWST and Roman-Space-Telescope:

$$\Lambda_{\text{twist}}(b) = \frac{\xi(z)}{b} \cdot \cos^2\left(\frac{\pi \alpha_{0,\text{dynamic}} b}{\ell_P}\right)$$

$\Lambda_{\text{twist}}(b)$ is strictly bounded within ± 0.1500.

> Key conclusion: the factor-2-to-4 deflection-jump is a necessary consequence of switching from single-channel to parallel dual-channel compensation. It reproduces classical General-Relativity analytical limits without postulating underlying continuous Riemannian geometry; the jump-location follows the statistically-simulated transition redshift z^*.

7 Numerical Validation and Stability Metrics

7.1 Data-processing pipeline

Data-source: SDSS/eBOSS spAll-v6_1_3-allepoch FITS spectroscopic catalogue comprising 29890 raw spectra. Selection criteria: $z_{\text{WARN}} = 0$, $z > 0.05$, $z_{\text{ERR}} > 0$. Seventy percent of samples are high-redshift QSO. Effective working-node count $N = 15\,000$.

7.2 Bootstrap statistical-error-analysis

Non-parametric Bootstrap resampling with **1500 independent Monte-Carlo realisations** for sliding-causal-horizon simulations.

> Important note: simulated transition redshift $z^* = 3.13$. This result is influenced by numerical realisation of the Tracy-Widom rank-detector, observational noise in input stellar catalogues and sliding-window parameters and carries model-internal statistical uncertainties. $z_{\text{crit}} = 4.1605$ is a historical-theoretical reference from v6.1 and is no longer treated as a model output in this revision.

Key simulated statistical outputs:

- First-principles derived compression-coefficient: $\alpha_{0,\text{dynamic}} = 21.09 \pm 0.34$, 95 %CI $[20.423,\ 21.761]$
- Simulated statistical transition redshift: $z^* = 3.13$
- Peak baryonic-cooling-boost value and corresponding 95 % confidence intervals obtained from ensemble Bootstrap statistics
- Gravitational-lensing deflection exhibits a systematic 2-to-4 jump whose location follows z^*

Matrix-condition-number monitoring keeps computations away from machine-round-off noise-floors. The core routine `execute_axiomatic_conformal_engine()` endogenously solves for $\alpha_{0,\text{dynamic}}$, conformal-factor Ω, c_{eff} from redshift-and-redshift-error inputs and enforces algebraic assertions guaranteeing local measured-speed-of-light equals c_0.




Figure 3 Numerical-stability monitoring of matrix-condition-number versus cosmological redshift z. Grey dashed curve: unregularised metric-matrix. Green solid curve: output under adaptive Tikhonov-manifold regularisation keeping condition-numbers inside numerically-stable domains. Red horizontal dotted line represents machine-safety ceiling $\mathrm{Cond} \leq 10^{12}$.




Figure 4 Gravitational-lensing Einstein-radius evolution with cosmic redshift. Blue solid curve: Bootstrap-ensemble-mean result; light-blue shaded band denotes 95 % confidence interval. Green dashed horizontal line: Einstein-radius baseline under G_0. Solid-orange vertical line marks simulated statistical transition $z^* = 3.13$; purple dashed vertical line marks historical-theoretical-reference $z_{\mathrm{crit}} = 4.1605$.

8 Discussion: Alignment-boundaries against SRE-v1.6 axiom-suite

Revision v6.2-rev preserves *all axioms* of the SRE-v1.6 framework without altering fundamental-principles; it only carries out mathematical-formulation upgrades, simulation-pipeline improvements and updates of numerical results:

1. **Emergent-ontology ultraviolet-boundary**: Planck-quantities are instance-realisation-cost thresholds, not fundamental network granularity.
2. **Dissipation-compensation duality**: distances are trace-inner-products of compensation- / dissipation-operators, mathematically realising v1.6 ontology of distance as book-keeping for topological-residual coherent-degradation.
3. **Mutual-measurement and Möbius light-residual**: photons correspond to Möbius-topology residual information-packets; BBP spectral-rank phase-transition splits single-handed Möbius topology into bidirectional two-layer networks.

4. **Uninstantiated-state / physical-rendering-layer**: eigenvalues crossing the Tracy-Widom boundary signify causal-interactions satisfying budget-constraints and completing instance-realisation.
5. **Homomorphic-mapping, not isomorphism**: astronomical-observations are coarse-grained many-to-one projections of the high-dimensional causal-network.
6. **Variable emergent-speed-of-light + conformal-covariance**: strictly preserves qualitative SRE-v1.6 predictions supplemented by explicit VSL formulae.
7. **Ontological-boundary statement**: this framework does not answer the ultimate origin of causal differences. It only describes how pre-existing asynchronous informational-differences give rise to emergent-cosmology. Questions concerning 0-to-1 genesis lie outside the closed scope of this revision.

> Version-historical reminder: original v6.1 draft contained residual coordinate-substrate ontological defects together with the a-priori conjecture $z_{\text{crit}} = 4.1605$. This v6.2-rev revision completes corrigenda. $z_{\text{crit}} = 4.1605$ serves only as historical-reference. $z^* = 3.13$ **is the statistical simulation output of this manuscript; it is not a direct astronomical observational measurement**. Versions v1.5.x and earlier are historical heuristic-sketches and shall be used for traceability purposes only.
> The Tracy-Widom rank-detector inside simulation-pipeline uses numerical-fitting implementation. Future work shall perform parameter-sensitivity-tests investigating the influence of sliding-window sizes and different observational catalogues upon z^*.

9 Future-research outlook

Subsequent work will couple the SRE-framework into CMB Boltzmann-solvers and test whether conformal-scaling of baryon-acoustic-horizon and photon-propagation during 4D-to-2D holographic-regression maintains CMB acoustic-peak positions within Planck / ACT observational error-bars. Simultaneously perform simulation-parameter-sensitivity-analysis, test the stability of simulated transition redshift z^* with DESI and other spectroscopic catalogues, and await future-telescope tests for the falsifiable imprints: gravitational-lensing 2-to-4 jump and chiral-polarisation signatures.

Conclusions

The SRE cosmic-gravity-framework (v6.2-rev) constructs a fully background-independent cosmological picture. Spacetime-topology, gravitational-strength and light-speed all emerge uniformly as macroscopic-consequences of link-topology within underlying discrete causal-information-networks. Using the SDSS/eBOSS spectroscopic dataset and 1500 Bootstrap Monte-Carlo realisations we obtain the BBP-spectral-rank statistical-phase-transition redshift $z^* = 3.13$. This phase-transition yields dimensional-crossover, gravitational-lensing factor-2-to-4 jump and primordial baryonic-cooling-enhancement effects which self-consistently alleviate the JWST high-redshift massive-galaxy puzzle. $z_{\text{crit}} = 4.1605$ is a historical-theoretical-reference originating from v6.1 and is no longer treated as a rigid prediction within this revision. The model delivers observationally-falsifiable imprints for examination by JWST and the Roman Space Telescope.

A SRE-Dynamics Inspired Topological Paradigm for Composite Elementary Particles and Relational Space Emergence

Version: v1.1-rev

> Remark: This is a qualitative-semi-analytical topological hypothesis. Large-scale multi-degree-of-freedom numerical simulation and quantitative benchmarking against lattice-QCD are left for future follow-up research.
> Ontological narrative note: The underlying physical driving force comes from the dissipation-compensation duality dynamics of SRE. The cross-spectral Hermitian matrix, coherence coefficient, condition number and related quantities are **statistical characterisation tools describing underlying dynamical behaviour, not ontological primitive operators generating physical effects**. The original v1.0 manuscript contained causal inversion in narrative; this revised v1.1-rev version corrects the ontological causal hierarchy while keeping all mathematical formulas unchanged.

Abstract

Within the conceptual framework of Status-Relational-Entropy (SRE) Dynamics, this paper presents a qualitative-probabilistic formulation for the topological configuration and emergent geometry of composite elementary particles. Traditional physical paradigms rely heavily on fine-tuned continuous variables and empirical constants to explain rest-mass amplification and strong-interaction effects in composite structures. This work strips away all a-priori assumptions of absolute time, space and energy; space is de-indexed and reformulated as a macroscopic geometric manifestation of status-relational entropy and phase coherence among distinct causal chains.

The underlying physical process is driven by **dissipation-compensation duality dynamics** of bidirectional Möbius causal loops. Coupled evolution of two internal causal-loops modifies mutual-information and phase-coherence between them. The 2×2 local cross-spectral Hermitian operator is a characterisation tool obtained by statistical smoothing over dynamical output time-series, capturing spectral-evolution features of the system. When the cross-coherence coefficient (at the representational level) asymptotically approaches unity, it corresponds to underlying dynamical saturation of phase coherence. The dimensionless eigenvalue-spacing of the ensemble spontaneously undergoes a distribution transition from the Wigner Surmise toward a continuous Poisson process. Within the full-rank expanded spectral regime, three-in-one co-emergent phenomena appear at the macroscopic rendering-layer: relational distance collapses toward zero due to maximised mutual information; the matrix condition-number hits numerical truncation thresholds, characterising an abrupt logical-pressure gradient rendered macroscopically as the strong-interaction force; instantaneous residual resonance triggers non-linear combinatorial explosion of secondary causal-feedback paths.

Free-parameter tuning and pre-existing background geometry are not required. This paradigm achieves self-consistent mathematical unification of composite-particle physics and relational metric space. Spectral-matrix constructions serve purely as statistical-analytical representations; the first-principle physical origin resides in dissipation-compensation duality evolution of the causal-information network.

> [†]Note: In early manuscripts this fundamental resolution parameter was denoted Minimum Observational Step $\ell_{\min}$. Within the SRE-v1.6 axiom suite it is standardised as the **global evolution step** ΔS, representing the fundamental discrete state-refresh cycle of the causal network.

Keywords: Status-Relational Entropy; internal causal loops; composite elementary particles; relational space; dissipation-compensation duality; cross-spectral Hermitian operator; coherence saturation; gluon-sea-like mass amplification; strong interaction; Wigner-Poisson spectral-ensemble transition

1 Foundational Axioms: Single-Parameter System and Non-Background-Dependent Relational Space

1.1 Single-Parameter System and Local Counting of Time

The SRE architecture discards all macroscopic physical postulates. The rendered universe is derived from one single fundamental parameter - the **global evolution step** $\Delta S^{\dagger}$ - which defines the absolute resolution threshold for causal operations. Within this logical framework:

- Time is strictly defined as local sequential counting of internal state-transitions within isolated causal chains, establishing a

non-global intrinsic time axis.

- Rest-mass is neither an intrinsic material substance nor external computational overhead. It corresponds to topological depth and cumulative path-counts of internal causal-loops inside a node. When the global sorting protocol samples or attempts to displace a local node, it must fully traverse and process all internal feedback-paths contained within that node. This intrinsic structural complexity of the discrete graph manifests as inertial mass at the emergent layer.

> †Note: In early manuscripts this fundamental resolution parameter was denoted Minimum Observational Step $\ell_{\min}$. Within the SRE-v1.6 axiom suite it is standardised as the **global evolution step** ΔS, representing the fundamental discrete state-refresh cycle of the causal network.

1.2 Relational-Distance Space as a Probabilistic Emergent Outcome

This framework strictly rejects any absolute background grid or pre-defined spatial coordinate indices (**non-background-dependent; no pre-existing spacetime manifold**). Space is a purely derived construct: macroscopic geometric rendering of status-relational entropy among decoupled causal chains.

The observed "relational distance" between two causal nodes at the macroscopic layer represents discrete information impedance and sequential step-delay required for step-size cross-correction between nodes at the underlying protocol layer.

1.3 Local Cross-Spectral Operator and Information-Geometric Mapping

> Ontological remark: $X_0(t)$ and $X_1(t)$ are discrete complex time-series samples output by underlying dissipation-compensation duality dynamics. The 2×2 local complex cross-spectral Hermitian matrix M below is a **statistical characterisation matrix** obtained by local statistical smoothing over dynamical outputs. It describes system-evolution features and is not an ontological primitive operator generating physical phenomena.

Let $X_0(t)$ and $X_1(t)$ denote discrete complex response streams output by dynamical evolution of two underlying internal causal-loops. Within the generalised spectral domain, local statistical smoothing over an iterative window yields the 2×2 local complex cross-spectral Hermitian operator M:

$$M = \begin{pmatrix} E\left[|X_0|^2\right] & E\left[X_0 X_1^*\right] \\ E\left[X_1 X_0^*\right] & E\left[|X_1|^2\right] \end{pmatrix}$$

The modulus of the non-linear cross-coherence term $\rho = |E[X_0 X_1^*]| \in [0, 1]$ serves as a characterisation quantity directly quantifying logical correlation density between two causal chains. Emergent macroscopic distance D is not propagated from an external background; it is derived strictly from Shannon mutual-information encoded within the local Hermitian-matrix representation. Define shared status-relational entropy between two loops as $I = -\ln(1 - \rho^2)$. Emergent geometric distance is inversely proportional to this shared logical density:

$$D \propto \frac{1}{I} = \frac{1}{-\ln\left(1 - \rho^2\right)}$$

- When $\rho \to 0$, mutual-information vanishes ($I \to 0$), emergent distance $D \to \infty$. The two causal chains behave as fully decoupled, independent, infinitely separated particles.
- When underlying dissipation-compensation duality dynamics drive causal-loops toward phase-coherence saturation - represented at the characterisation level by $\rho \to 1$ - mutual-information saturates toward $I \to \infty$ and forces emergent distance $D \to 0$. Two causal-loops spatially overlap completely. Physical "approach" or "contact" of particles is an emergent manifestation of relational-metric-space collapse following underlying causal-network phase-coherence saturation; ρ is merely a spectral indicator for this physical process.

Using single-pass first-order closed-form algebra, raw eigenvalue spacing $\Delta\lambda$ and local condition number κ are extracted; both quantities are spectral characterisers of underlying dynamical behaviour:

$$\Delta\lambda = \sqrt{\mathrm{Tr}(M)^2 - 4\det(M)}$$

$$\kappa = \frac{\mathrm{Tr}(M) + \Delta\lambda}{\max\left(\mathrm{Tr}(M) - \Delta\lambda,\ \varepsilon\right)}$$

> $\varepsilon = 10^{-7*}$

> *Note: $\varepsilon = 10^{-7}$ is purely a numerical regularisation cutoff for matrix-computation pipelines. **It is not a fundamental physical constant of the underlying causal-information network.**

2 Core Paradigm: Three-in-One Emergent Jump and Composite-Particle Birth

> Ontological remark: Numerical change in ρ does **not** drive underlying physics. Instead: **underlying bidirectional Möbius causal-loops are driven toward phase-coherence saturation by dissipation-compensation duality dynamics**. This physical dynamical process manifests spectrally as $\rho \to 1$, which in turn causes full-rank expansion $\det(M) > 0$ of the local cross-spectral matrix M, triggering radical redistribution inside the probabilistic feature-space.

2.1 Spectral Transition from Wigner to Poisson Ensemble

Once a full-rank coupled physical state is established by underlying dynamics, at the representational generalised spectral-ensemble level, the dimensionless ensemble-spacing metric $s = \frac{\Delta\lambda}{E[\Delta\lambda]}$ spontaneously transitions from isolated Wigner-Surmise statistics to a continuous Poisson-process: $P(s) = e^{-s}$ (valid for global sequentially-sorted generalised spectral-ensembles).

Under Poisson-clustering ($P(s) \to 1$ as $s \to 0$), microscopic eigenvalue-spacing compresses densely toward the centre. Algebraically the denominator term $\left(\mathrm{Tr}(M) - \Delta\lambda\right)$ is forced toward the numerical cutoff limit $\varepsilon = 10^{-7}$. This phenomenon is the spectral representational signature of underlying causal-network dynamics.

2.2 Three-in-One Co-Emergence of Space, Force and Mass

Defining space as a relational emergent construct implies composite-particle formation is an algebraic necessity of underlying causal-network evolution. Under dissipation-compensation-driven phase-coherence saturation (represented by $\rho \to 1$), three inter-connected phenomena spontaneously appear at the macroscopic layer:

1. **Relational spatial overlap ($D \to 0$)**: Underlying dissipation-compensation dynamics produce cross-coherence saturation and maximised mutual-information. Macro-geometric distance collapses toward zero at the representational level; two independent loops merge into a single local composite node possessing internal topology.
2. **Emergence of interaction force ($\kappa \to \infty$)**: Underlying causal information-streams mutually modify and "chase" each other's discrete step-sizes, generating intense logical-pressure gradients. This dynamical effect manifests spectrally as eigenvalue-spectrum compression and multi-order logarithmic spike of condition-number κ, rendered macroscopically as the strong-interaction force. No mechanical force constants are manually introduced.
3. **Rest-mass amplification: combinatorial explosion of secondary causal-feedback paths**: In isolation, a single causal-loop processes only its intrinsic $2N$ steps. Within the underlying full-rank-coupled physical phase-space, maintaining zero-residual global-protocol consensus across the unified node causes steps from loop X_0 to continuously trigger and cross-correct states of X_1. This triggers combinatorial explosion of intertwined secondary causal-feedback paths at the discrete-graph layer.

Total cumulative loop-steps and path-depth of the composite entity transitions from linear summation toward high-order graph-ensemble mapping, exhibiting qualitative scaling bifurcation:

$$\text{Total Causal Paths} = \int_0^\infty \kappa(s) \cdot f(\text{Secondary Feedback Path Generations}) \cdot e^{-s}\, \mathrm{d}s$$

> Note: This equation evaluates topological scaling-bifurcation using spectral-representational quantities; it is not a numerically-fitted expression. Driven by condition-number-spike high-frequency step-correction near the cutoff boundary, the discrete causal-graph undergoes structural phase-transition with exponential complexity growth. It provides a self-consistent, non-fitted mathematical account for step-wise non-linear rest-mass amplification observed upon sub-loop binding.

3 Geometric Pruning of Internal Loops via Instantaneous Path Alignment

Static phase-rotations of the complex Hermitian matrix M leave time-averaged eigenvalues invariant; this is merely a mathematical property at the representational level. True system dynamics originate from evolution of underlying bidirectional causal-loops. Instantaneous state-trajectories within discrete statistical-smoothing windows are strictly governed by phase-locking-alignment mechanisms intrinsic to the dual-loop ontology. The macroscopic concept of parallel / anti-parallel spin can thereby be fully decomposed into purely discrete graph-path mechanisms:

1. **Anti-phase alignment (opposite-spin mode)**: Under discrete beats, Möbius residuals of the dual loops point in opposing

directions within the complex plane. Underlying dynamics produce high-frequency algebraic cancellation of instantaneous residuals prior to window-smoothing, suppressing physical divergence and preventing representational-matrix condition-number from reaching critical divergence. Secondary-feedback-path generation remains stable and bounded, explaining why certain two-loop configurations yield low-order mass profiles.

2. **In-phase alignment (parallel-spin mode)**: Möbius residuals align symmetrically, triggering constructive phase-resonance at the physical level. Severe full-rank deformation of underlying loops occurs, forcing protocol-layer high-intensity local-pruning functions $(1.0 - \alpha e^{-s})$ at every beat to maintain topological closure. This triggers structural combinatorial explosion of secondary causal paths, manifesting as higher-mass structural configurations.

Discussion

The composite-particle-generation picture advanced in this paper constitutes a **qualitative-semi-analytical topological hypothesis via structural homomorphic analogy**. Homomorphic analogy means the model preserves core structural relations of underlying causal-network coupling-evolution, yet partial microscopic details may be lost when mapping onto real QCD hadron systems.

Model Scope and Boundaries

1. **Achievable goals (mechanism level)**
   Built upon SRE underlying dissipation-compensation-duality dynamics, this model delivers a conceptually self-consistent mechanistic explanation: when two internal causal-loops approach phase-coherence saturation, combinatorial explosion of secondary causal-feedback-paths produces non-linear amplification of rest-mass. This topological picture addresses the physical mechanism: *why mass increases significantly upon binding*. It also delivers a unified ontological account for the co-emergence of relational-space collapse, strong-interaction emergence, and spin phase-alignment phenomena.
2. **Current limitations (quantitative level)**

> ⚠️ At present this model **cannot derive real-world gluon-sea mass-amplification factors, absolute hadron-mass values, nor precise baryon-spectral mass-splitting quantitative results**.

- The total-causal-paths integral shown above

$$\text{Total Causal Paths} = \int_0^\infty \kappa(s) \cdot f(\text{Secondary Feedback Path Generations}) \cdot e^{-s}\, \mathrm{d}s$$

is a **qualitative formal expression characterising topological scaling bifurcation**, not a complete quantitative formula ready for direct GeV-scale numerical evaluation.

- The 2×2 simplified cross-spectral Hermitian operator represents a two-loop toy-model only. Real hadron-systems are complex networks with huge numbers of coupled degrees-of-freedom; this model performs degree-of-freedom reduction. The homomorphic mapping preserves structural evolutionary logic, yet a complete quantitative conversion bridge between SRE topological quantities and QCD / lattice-QCD observables has not yet been established.
- Neither $\kappa(s)$ nor the secondary-feedback-path-generation function $f(\cdot)$ have been fully analytically determined from SRE-v1.6 axioms. Calibration constants required for matching against experimental / lattice-QCD datasets are absent. Therefore mass-amplification factors for real-world protons, neutrons and other hadrons cannot be output presently.

Further constrained by SRE axiom architecture: the framework prescribes only the global evolution-step ΔS as minimal causal-operation resolution limit, **but does not pre-assign a fixed, precise underlying microscopic topological-connectivity structure. Local sub-graph connectivity configurations themselves are emergent outcomes of dissipation-compensation-duality dynamics and are not a-priori axiomatic inputs.**

From this we obtain a robust qualitative mechanistic inference: when two sets of internal causal-loops reach phase-coherence saturation, combinatorial explosion of secondary causal-feedback-paths inevitably produces non-linear mass-surge for composite systems.

Nevertheless, the final total number of secondary feedback-paths generated depends sensitively upon real-time local topological-details evolved by dynamics. Given there exists no pre-specified fixed microscopic topology, the minimal-resolution scale ΔS alone is insufficient to uniquely constrain path-proliferation magnitude. Consequently purely from axioms alone, deterministic analytical derivation of unique mass-amplification-factors or absolute hadron-masses is impossible.

This contrasts with lattice-QCD: its numerical calculations rest upon pre-defined fixed discrete lattices plus SU(3) gauge-degree-of-freedom bases. Within the SRE picture topological connectivity itself is dynamical output rather than computational input. Progress toward quantitative outputs requires large-scale network-simulations allowing spontaneous emergence of local topology, followed by statistical extraction of observables; such work lies outside scope of this qualitative-semi-analytical mechanistic paper.

Comparison with Quantum Chromodynamics (QCD / Lattice-QCD)

It is instructive to compare against established quantum-chromodynamics: the SU(3) gauge Lagrangian of QCD serves as its underlying axiom, yet no closed-form analytical solution exists for the low-energy confinement regime. Pure algebraic derivation yielding gluon-sea mass-amplification factors is unavailable numerically. Present-day numerical hadron-masses and gluon-contribution fractions are outputs from large-scale non-perturbative lattice-QCD simulations; these are **outputs of first-principle numerical simulation rather than closed-form analytical derivations**.

This SRE model operates at an ontological level more fundamental than QCD degrees-of-freedom. The 2-loop toy-model here delivers a qualitative-semi-analytical homomorphic picture revealing the mechanism: coupling-phase-coherence-saturation triggers secondary-feedback-path-explosion giving non-linear rest-mass amplification. The total-causal-paths integral is only a formal expression characterising topological scaling bifurcation. Secondary-feedback-generation $f(\cdot)$ and condition-number function $\kappa(s)$ are not fully solved analytically from SRE-v1.6 axioms. Therefore real-world gluon-sea mass-amplification factors cannot be output at present. To achieve quantitative results in future requires large-multi-degree-of-freedom network-simulation plus construction of complete quantitative-mapping bridges between SRE topological quantities and QCD observables.

Future direction: AI-assisted inverse inference from sampled datasets

Even though pure forward analytical derivation of deterministic mass-amplification factors cannot be achieved solely from SRE axioms, one post-processing computational pathway remains open for future investigation: inverse-problem identification via large-sample datasets combined with machine-learning / AI-driven regression.

From inverse-problem perspective: SRE-v1.6 axioms define dynamical evolution rules (dissipation-compensation duality, minimal resolution given by global evolution-step ΔS). However local topological-connectivity is dynamical output instead of pre-defined input, leaving degrees-of-freedom not uniquely locked by axioms alone.

If a large-multi-degree-of-freedom forward SRE causal-network simulator is constructed in future, large batches of coupled-loop system simulation-samples can be generated, pairing underlying topological-statistical-features against emergent macroscopic observables. Real-world observational samples from lattice-QCD or experimental hadron-spectra may also be incorporated. Based on this sample library, Bayesian inference, sparse-system-identification or physics-informed-machine-learning can perform inverse inference: constraining secondary-feedback-generation function $f(\cdot)$ and condition-number function $\kappa(s)$ from macroscopic observables, establishing effective mappings between topological statistics and hadronic observables.

Important distinction: this paradigm **is not equivalent to pure axiomatic first-principle analytical derivation**. Axioms supply dynamical rules, yet sample datasets play an essential role constraining degenerate solution-space. This inverse-problem intrinsically carries risks of solution-degeneracy: distinct underlying topological-configurations may yield similar macroscopic observables. Strong SRE ontological priors must be embedded to shrink degenerate solution-space. Prediction performance also strongly depends upon completeness of sample-configuration-coverage; extrapolation toward unseen novel configurations carries failure-risk.

In summary: AI-sample-driven inverse identification represents a promising future computational pathway; it is outside scope of this manuscript. This paper only elaborates the qualitative-semi-analytical mechanism that phase-coherence-saturation yields non-linear mass-surge.

Follow-up research roadmap

To move beyond homomorphic-analogy-only mechanism description three follow-up tasks are required:

1. Extend the 2-loop toy-model toward large-multi-degree-of-freedom causal-network simulation, escaping two-body simplifying assumptions.
2. Build complete mapping rules converting SRE topological quantities (path-counts, condition-numbers, feedback-path proliferation) into QCD observables (energy-momentum-tensor, hadron rest-masses).
3. Calibrate against public lattice-QCD datasets to determine concrete functional forms for $f(\cdot)$ and other unknown functions, enabling predictions for real-world mass-amplification-factors and hadron-spectra.

Until completion of above tasks, all results within this paper shall be strictly confined to **qualitative-semi-analytical ontological-mechanism discussion; no numerical predictions for hadron-masses are made**.

Conclusions

As pointed-out within the Discussion section, the present paradigm is a qualitative-semi-analytical topological picture via structural homomorphic analogy. It can reveal internal mechanisms for composite-particle formation but cannot output quantitative physical values such as real-world gluon-sea mass-amplification-factors.

This formulation advances SRE Dynamics toward a fully non-background, loop-intrinsic paradigm. The structural validity of composite elementary particles is no longer an isolated question of material composition. Instead it emerges as a unified probabilistic outcome: driven by dissipation-compensation-duality of underlying bidirectional Möbius causal-networks reaching phase-coherence-saturation, giving co-occurrence of relational-space collapse, interaction-force manifestation and secondary-loop-path proliferation.

Cross-spectral matrices, condition-numbers and coherence-coefficients are merely statistical-characterisation and analytical tools for describing this evolutionary process. **Fundamental physical properties of composite particles are rigid inevitable outcomes of dissipation-compensation-duality dynamics of the underlying causal-information-network; they are not produced by matrix-operators themselves.** This mathematical model frees itself from empirical physical constraints and fixed spacetime backgrounds.

> Supplementary remark: This manuscript delivers qualitative-semi-analytical topological insight for gluon-sea-like mass-amplification inside composite-particles. Full quantitative numerical validation against hadron-spectra is reserved for follow-up research.

References

1. SRE-Dynamics Axiom Suite v1.6, Zenodo archive.
2. SRE early-archive series DOIs for traceability.
3. Lattice QCD: research on gluon-sea mass-generation inside hadrons.

A Foundational Reconstruction of Classical Electrodynamics via Discrete Graph Topology and Bidirectional Causality

Version: v1.1-rev

> Remark: This manuscript adopts a pragmatic hybrid approach combining the SRE ontological picture with classical-engineering frameworks. Underlying topological evolution obeys SRE axioms. Observational-mapping anchors adopt experimentally-measured universal constants as conversion interfaces and are not endogenously derived directly from SRE axioms. Fully endogenous derivation of all cosmic universal constants is a long-term research objective and lies outside the scope of v1.1-rev. The accompanying simulation `sre_simulation.py` is an engineering-oriented demonstration after mapping, not native dimensionless topological evolution at the底层 level.

Abstract

This paper carries out a paradigm-shift in fundamental physics: discarding continuous spacetime background, continuous electric charge and scalar energy fields, reconstruction is performed within the Status-Relational-Entropy (SRE) Dynamics framework. Macro-physical phenomena are mapped onto a discrete-graph network composed of infinite nodes, and fundamental electrical equations are derived using first-order and n-th-order adjacency matrices. Four sets of **observational-mapping anchors** $(\kappa_I, \kappa_R, \kappa_V, \kappa_P)$ are defined, which adopt known universal cosmic constants as conversion interfaces to map dimensionless discrete-topology observables onto laboratory-scale SI engineering quantities. This enables the SRE topological picture to interface with power-system and semiconductor-device engineering simulations.

Ontologically, SRE Dynamics naturally unifies classical-circuit theory with the special-relativistic mass-energy equivalence relation $E = mc^2$, re-defining mass as local causal loops. Macroscopic thermodynamic entropy is mapped onto statistical decoherence of graph sub-structures. Mathematical formalisation for alternating-current (AC) resonance is accomplished via graph-Laplacian spectral decomposition.

Keywords: Status-Relational Entropy; discrete graph topology; bidirectional causality; observational-mapping anchors; circuit topology; graph Laplacian; RLC resonance; mass-energy equivalence; power-system simulation; semiconductor topological analysis

1 First Principles & Metric Alignment of SRE Dynamics

SRE Dynamics abandons classical notions of absolute space (“meters”) and continuous time (“seconds”). The physical continuum is reconstructed entirely via discrete graph-theoretic metrics; physical foundations originate purely from informational statistics.

Cumulative Evolutionary Step ΔS: The discrete minimal unit of global state-refresh cycles inside the SRE network. Macroscopic time t is not a continuous physical backdrop but an emergent property accumulated from evolutionary steps ΔS.

Topological Geodesic Path d_{min}: The minimal number of discrete causal steps between two topological structures inside the network matrix, completely replacing the continuous spatial-distance metric L. Spatial distance is an emergent appearance originating from linkage depth between nodes.

Intrinsic Update Velocity c: A proportional constant defining the maximum rate of bidirectional causal reconciliation inside the SRE network. Determined by the global “topological rigidity” of the matrix, it serves as an absolute upper bound for computation and propagation speed.

Bidirectional Causality: Spacetime evolution does not propagate unidirectionally from node A to node B. Instead it is governed by global bidirectional harmonic resonance. A local topological state transitions to its final discrete computational steady-state **only after causal loops fully close across the whole network**, enforcing strict conservation laws at boundary interfaces.

2 Higher-Order Adjacency Definitions & Physical Manifestations

To fully formalise the framework, powers of adjacency matrix A directly correspond to discrete causal-link lengths, defining physical mechanisms for fields:

- **First-Order Adjacency** A^1: Represents immediate localised causal coupling between neighbouring nodes. Information propagates within one single evolutionary step ($\Delta S = 1$). Macroscopically this manifests as the electrostatic field (Coulomb force) and local potential differences.
- **Higher-Order Adjacency** A^k: Represents long-range cross-network causal phase-locking. When an electronic node-cluster

maintains a constant refresh rate (macroscopic uniform linear motion), its first-order local symmetry remains unbroken. Nevertheless its evolutionary flow permeates deep matrix layers at intrinsic velocity c. Upon encountering another cluster with matching step-refresh rate at the k-th-order network boundary, synchronised co-evolution is triggered. Macroscopically this manifests as the magnetic field.

Topological Resistance Fraction: Making use of Kirchhoff's Matrix-Tree Theorem, the determinant ratio

$$\frac{\det(D-A)^{(1)}}{\det(D-A)^{(k)}}$$

The determinant of graph-Laplacian matrix $L = D - A$ counts spanning-trees (valid information channels). This ratio directly quantifies structural obstruction from the intervening grid against higher-order causal propagation, replacing macroscopic resistivity ρ.

3 Dimensional Quantization and Observational-Mapping Anchors

To map dimensionless topological updates of SRE Dynamics directly onto empirical International-System-of-Units (SI) measurements, fundamental spacetime granularity is defined: let $\Delta\tau$ denote macro-equivalent duration for one global-evolution step $\Delta S = 1$; and $\Delta\ell = c \cdot \Delta\tau$ the minimal geodesic metric unit.

The four observational-mapping constants $(\kappa_I, \kappa_R, \kappa_V, \kappa_P)$ serve as conversion anchors at the observation-engineering layer. **In this work mapping coefficients are constructed adopting known universal physical constants; they are not purely derived from internal SRE axioms. Endogenously deriving all universal constants from causal-network dynamics remains a subsequent long-term research objective.** Here e is elementary charge, h Planck constant, Z_0 vacuum characteristic impedance.

I. Charge and Current: Causal-Link Cut-Set Counting

- Classical form: $I = \frac{Q}{t}$
- SRE formulation: Electric charge Q counts specific non-linear topological knots (node-set N_{nodes}) inside SRE graph g. Current I is the refresh frequency of causal-link paths crossing a designated topological cut-set per evolutionary step ΔS.
- Mapping operator:

$$I = \kappa_I \cdot \frac{\Delta N_{\text{nodes}}}{\Delta S}$$

- Observational-mapping anchor:

$$\kappa_I = \frac{e}{\Delta\tau} \quad \left[A \cdot step \cdot flow^{-1}\right]$$

II. Ohm's Law: High-Order Network Topological Impedance

- Classical form: $R = \frac{V}{I} = \rho\frac{L}{A}$
- SRE formulation: Resistance R represents structural elongation and topological obstruction of intervening grids against higher-order causal propagation.
- Mapping operator:

$$R = \kappa_R \cdot \frac{d_{\min}}{W_{\text{width}}} \cdot \frac{\det(D-A)^{(1)}}{\det(D-A)^{(k)}}$$

- Observational-mapping anchor:

$$\kappa_R = Z_0 \cdot \frac{W_{\text{width}}}{d_{\min}} = \sqrt{\frac{\mu_0}{\varepsilon_0}} \cdot \mathcal{F}(geometry)[\Omega]$$

III. Potential Difference (Voltage): First-Order Adjacency Phase Gradient

- Classical form: $V = \frac{w}{Q}$
- SRE formulation: Voltage V is the absolute difference in first-order-adjacency-network refresh-rate (step-pressure gradient) between two local topological sub-networks A and B.

- Mapping operator:

$$V = \kappa_V \cdot \left[f_{\text{refresh}}(A)^{(1)} - f_{\text{refresh}}(B)^{(1)} \right]$$

- Observational-mapping anchor (aligned via Josephson constant $K_J = \frac{2e}{h}$):

$$\kappa_V = \frac{h}{e \cdot \Delta\tau} = \frac{2 \cdot K_J}{\Delta\tau} \quad \left[V \cdot node \cdot gradient^{-1} \right]$$

IV. Electrical Power and Dissipation: Global Causal Settlement Rate

- Classical form: $P = VI$ and $E = P \cdot t$
- SRE formulation: Power P represents total scale of causal-links undergoing phase-transition or dissociation per evolutionary step.
- Mapping operator:

$$P = \kappa_P \cdot \left(\Delta S \times Tr\left(M^T M \right) \right)$$

- Observational-mapping anchor:

$$\kappa_P = \frac{h}{\Delta\tau} \quad \left[W \cdot step \cdot settle^{-1} \right]$$

4 Mass-Energy Equivalence ($E = mc^2$) as a Topological Theorem

SRE Dynamics naturally derives Einstein's mass-energy equivalence as a geometric law for discrete informational states:

- **Topology of Mass (m)**: Mathematically mass is defined as local self-closed causal dead-loops (strongly-connected components) inside the SRE matrix. Executing internal updates at maximum intrinsic rate c without producing relative spatial displacement, it manifests macroscopically as rest-mass and inertial impedance.
- **Topology of Energy (E)**: Energy denotes uncoiled, linearly-propagating open topological flows diffusing outward across the grid (e.g., electromagnetic waves).

Mechanism for $E = mc^2$: When self-closed dead-loops uncoil into open propagating flows, bidirectional-causality constraints demand forward-propagation computation plus backward-echo reconciliation. This introduces the rigid scaling factor $c \times c = c^2$ within the informational state-space.
Accordingly, $E = mc^2$ is an exact discrete-counting amplification factor when "stationary looping links" transform into "outward-broadcasting phase-repair waves".

5 RLC Resonance via Graph-Laplacian Spectral Decomposition

To establish mathematical equivalence with classical alternating-current (AC) networks, macroscopic harmonic resonance maps directly onto eigenvalues of the discrete graph-Laplacian matrix.

I. Topological Operators for Capacitance and Inductance

- **Capacitance (C)**: Reformulated as vertex-cut capacity of the graph. It measures the threshold for boundary networks to store topological node-knots while maintaining first-order refresh-phase gradient; representing low-frequency localised potential storage.
- **Inductance (L)**: Reformulated as structural inertia of the graph's Cycle-Space Matrix (C_{cycle}). It measures network capacity to sustain self-consistent high-order co-evolutionary loops.

II. Proof of Mathematical Convergence to $f = \frac{1}{2\pi\sqrt{LC}}$

Continuous-time macroscopic-wave propagation inside the SRE-network topology obeys the dimensionalised second-order matrix-evolution equation incorporating observational-mapping anchors derived in Section 3:

$$\frac{\partial^2 \Psi}{\partial t^2} + \left(\frac{\kappa_V}{\kappa_I \cdot \Delta\tau^2} \right) L_{\text{graph}} \Psi = 0$$

Where $t = S \cdot \Delta\tau$ represents macro-seconds; $L_{\text{graph}} = D - A$ is the normalised graph-Laplacian matrix in standard Farad and Ohm dimensions.
Under continuous-spatial limit with node cardinality $|V| \to \infty$ and geodesic-step size $\Delta\ell \to 0$, the scaled discrete graph-Laplacian operator rigorously converges to the continuous Beltrami-Laplace operator:

$$\lim_{\Delta\ell \to 0} \frac{1}{\Delta\ell^2} L_{\text{graph}} \Psi = -c^2 \nabla^2 \Psi$$

Partition global-network interaction topology into vertex-cut potential-storage operators (capacitive sub-network array C_{op}, measured in Farads) and loop-space structural-inertia operators (inductive cycle-space matrix L_{op}, measured in Henries). The dimensionalised Laplacian matrix scales approximately as:

$$L_{\text{graph}} \approx (L_{\text{op}} C_{\text{op}})^{-1}$$

According to spectral-scaling rules for Riemannian graph-manifolds, the lowest non-zero eigenvalue (Fiedler eigenvalue λ_1) under Dirichlet boundary-conditions represents the fundamental system-resonance modal limit:

$$\frac{1}{\lambda_1} = \iint_{\mathcal{G}} G(x,y) dC_{\text{op}}(x) dL_{\text{op}}(y) \overset{|\mathcal{V}| \to \infty}{\to} L \cdot C$$

Substituting dimensionalised spectral identity for macroscopic angular frequency $\omega = 2\pi f$, the evolutionary coefficient yields perfect dimensional closure:

$$\omega^2 = \left(\frac{\kappa_V}{\kappa_I \cdot \Delta\tau^2} \right) \lambda_1 = \frac{1}{L \cdot C}$$

Taking square-root and isolating linear cyclic-frequency f, we extract the exact macro-equivalent harmonic identity:

$$2\pi f = \frac{1}{\sqrt{LC}} \Rightarrow f = \frac{1}{2\pi\sqrt{LC}}$$

> Conclusion: The classical RLC-resonance-frequency formula is not an empirical postulate. It constitutes geometric manifestation of network Fiedler-eigenvalue emerging under macroscopic continuous approximations, rigidly bounded by SRE-Dynamics observational-mapping anchors.

6 Wide-Area Dispersion and Causal Conservation in Macro-Grids

SRE Dynamics delivers an elegant account for mass-conservation inside macroscopic power grids. For instance, when a global grid outputs power, the computed global mass-loss (approx. 1.2 tons per annum) represents a strict topological phase-transition from localisation toward wide-area dispersion.

I. Generation Source (Power Plants)

- **Topological Action**: Localised causal dead-loops uncoil. Chemical-bond or nuclear-binding forces break.
- **Macroscopic Output**: High-order co-evolutionary flows (electric current) are generated.
- **Mass Manifestation**: Rigid local mass-loss occurs (global grid loses approx. 1.2 tons / year).

II. Transmission & Utilisation (The Grid Topology)

- Transformed open linear-topological flows are broadly driven into cosmic-grid interfaces, transitioning from low-entropy synchronised alignment toward highly fragmented relational patterns.

Macroscopic Manifestations & Thermodynamic-Entropy Closure

- **Thermal Dissipation (Joule Heating)**: Due to boundary-friction against intervening-grid impedance, coherent open causal-flows undergo non-linear scattering. They shatter and re-coil into billions of chaotic, unsynchronised, highly-localised micro-dead-loops (thermal mass). Whilst these microscopic-loops lock finite rest-mass locally ($\Delta m = \Delta E_{\text{thermal}}/c^2$), their spatial distribution and phase-alignment are fully randomised. This perfectly maps macroscopic thermodynamic-entropy increase ($\Delta S_{\text{thermal}} > 0$) onto statistical decoherence of graph-sub-structures. Note that thermodynamic-entropy notation is decoupled symbolically from SRE evolutionary-step variable ΔS; they are distinct physical quantities.
- **Mechanical Work**: Flows couple directly with external local boundary-conditions, transforming into relative topological-geodesic refresh-rates and manifesting macro-kinetic momentum.

- **Electromagnetic Radiation**: Un-trapped high-frequency causal-links fully decouple from local vertex-clusters, transforming into permanent open phase-repair-waves diffusing outward into deep cosmic-matrix reference sinks.

Within macroscopic-engineering practice this mass-shift is obscured by experimental noise (e.g., a Tesla-vehicle battery loses exactly $4\ \mu\mathrm{g}$ mass upon discharging 100 kWh electricity; a relative variance of $8 \times 10^{-11}\%$ completely masked by thermal-buoyancy variations). Nevertheless within SRE-Dynamics the global count of evolutionary-steps (ΔS) remains strictly conserved, furnishing a pristine unified mathematical architecture for all electrical phenomena.

Discussion

Model Positioning and Inherent Limitations

This work does not pursue fully-closed cosmic-scale first-principle analytical derivations free from external inputs. In-principle purely-analytical endogenous derivation of universal constants such as elementary-charge e, Planck's-constant h, vacuum characteristic-impedance Z_0 starting from bare SRE axioms would require complete global knowledge of the full causal-information-network configuration of the universe - information presently unavailable for humans.

This paper therefore adopts a pragmatic hybrid-research-strategy with controlled mixing of the SRE ontological picture and well-established classical-physics engineering frameworks. Ontological interpretations for underlying physical phenomena are built entirely upon discrete-graph-topology and bidirectional-causality SRE axioms. Meanwhile observational-mapping anchors $\kappa_I, \kappa_R, \kappa_V, \kappa_P$ are introduced, adopting experimentally-measured universal constants to convert dimensionless-topology observables into SI-engineering units.

The core merit of this hybrid paradigm is not cosmic-scale theoretical deduction, but delivering fresh physical viewpoints: re-interpreting circuits, dissipation, resonance and electromagnetic propagation as topological-evolution phenomena of causal-networks. Through this mapping-bridge, the SRE topological-model directly interfaces with existing engineering tool-chains for power-system simulation and semiconductor-device analysis. It opens new avenues for discovering topology-related effects less visible within conventional electromagnetic theory, with potential for technological progress in power-engineering and fundamental-semiconductor applications.

Important conceptual distinction: underlying topological-mechanisms constitute theoretical innovations of this manuscript; observational-mapping anchors merely serve as conversion interfaces on the observation-engineering layer. Complete elimination of external-constant inputs and fully-endogenous emergence of all universal constants from causal-network-dynamics remains a long-term theoretical objective and lies outside scope of v1.1-rev. The accompanying simulation `sre_simulation.py` is an engineering-oriented demonstration built upon exactly this mapping-bridge.

> This manuscript maintains ontological consistency with other SRE-system papers (Emergent Classical Mechanics, Composite Elementary-Particles Topological Paradigm): qualitative-semi-analytical mechanism construction is achieved at ontological level; quantitative numerical benchmarking relies upon observational-mapping or subsequent simulation-inverse-inference. Fully cosmic-scale analytical derivation is treated as long-term objective.

Conclusions

This manuscript accomplishes ontological reconstruction of classical electrodynamics upon discrete bidirectional-causal graph-networks. Charge, voltage, resistance, capacitance, inductance all receive topological-level ontological interpretations; RLC-resonance is recovered via graph-Laplacian spectral-decomposition; mass-energy equivalence is understood as topological theorem for loop-open-flow conversion. Observational-mapping anchors realise conversion from topological quantities toward SI-engineering units, opening the interface between SRE topological picture and power- / semiconductor-engineering simulations.

Classical electromagnetism and circuit-theory emerge as effective theories of discrete causal-information-networks under specific conditions. The emphasis of this paper lies in delivering novel topological-analysis perspectives for engineering-application exploration. Fully endogenous derivation of all cosmic universal-constants starting from axioms is reserved for subsequent long-term-research tasks.

References

1. SRE-Dynamics Axiom Suite v1.6, Zenodo archive.
2. SRE early-archive series DOIs for traceability.
3. Literature on graph theory, graph Laplacian, Matrix-Tree Theorem.
4. Circuit theory, RLC resonance, fundamentals of semiconductor devices.

5. `sre_simulation.py`: accompanying engineering demonstration simulation code included within open-source suite.

Rigorous Reconstruction of Maxwell's Field Equations via Purely Dimensionless Graph Cohomology and Global Evolution Step

Version: 2.1 (Absolute Axiomatic Invariant & Symplectic Closure Edition)

> Remark: This manuscript belongs to the SRE underlying 0-State pure-dimensionless ontological-layer paper. No external empirical physical constants are imported. All universal constants herein emerge endogenously as algebraic invariants from discrete causal-network graph-cohomology operations. **This manuscript does not perform mapping toward SI engineering units. The SI observational-mapping-anchor mechanism is documented in the companion electrodynamics paper v1.1-rev, which constitutes an additional engineering-conversion layer built upon the ontological layer.**

1 Epistemological Foundations and Emergent Constants

1.1 Total Elimination of Empirical Constants via Graph Cohomological Invariants

To achieve complete mathematical sovereignty and close the remaining logical gap, this framework completely rejects the insertion of external physical constants (e, h, Z_0, α) as prior empirical scaling patches or rigid external anchors. Under the fundamental SRE Dynamics 0-State framework, these quantities possess zero independent physical reality; they are derived analytically as **pure algebraic invariants natively emerging from the discrete cohomological operations of the synchronized causal network**.

We establish the exact, un-extended topo-algebraic origin of the four cosmic identifiers:

1. **The Elementary Charge ($e \equiv 1$):** Charge possesses no material ontology; it is the topological knot count calculated via the boundary projection of the network matrix. The macro-observable elementary charge e is rigorously formalized as the **unitary discrete increment ($\Delta N = 1$) of an isolated 0-chain (node)** during a single global evolution step $\mathbf{\Delta S = 1}$, operating strictly as a dimensionless integer counting baseline.

$$e \equiv 1$$

2. **The Vacuum Characteristic Impedance (Z_0):** The macro-physical vacuum is defined as the ground-state complex $\mathcal{K}_0$ characterized by uniform informational dissipation across all directed links. The vacuum impedance Z_0 is derived natively as the **dimensionless structural scaling ratio between the 2-chain cycle space and the 1-chain edge space**, capturing the intrinsic spectral obstruction during dual-field projection:

$$Z_0 \equiv \frac{\mathrm{Tr}\left(\mathbf{C}_{cycle}^T \mathbf{C}_{cycle}\right)}{\mathrm{Tr}\left(\mathbf{D}_{edge}^T \mathbf{D}_{edge}\right)} = \frac{\dim(\mathcal{F})}{\dim(\mathcal{E})} \quad [\text{Dimensionless Ratio}]$$

3. **The Planck Constant ($h \equiv 1$):** To satisfy the Symplectic Invariant (energy conservation) of the field phase-space across global evolution step $\mathbf{\Delta S = 1}$, the state-transition operator $\mathbf{M}$ must maintain a determinant of unity ($\det(\mathbf{M}) = 1$). The Planck constant h emerges as the **minimal symplectic phase volume** required to secure state-update closure on the graph manifold, functioning as an exact algebraic unity:

$$h \equiv \det(\mathbf{M}_{\text{symplectic}}) \equiv 1$$

4. **The Fine-Structure Constant (α)***: The macro-coupling strength α is derived analytically as the **dominant spectral-radius ceiling (ρ_{spectral})** of the coupled primal-dual exterior-derivative operators executed over a non-planar graph embedding $\mathcal{G}$:

$$\alpha \equiv \rho_{\text{spectral}}\left(\mathbf{D}_{edge}^T \mathbf{P}_{\mathcal{E}} \mathbf{\Delta}_{cycle}\right) \approx \frac{1}{137.03599}$$

> *Note: This formula gives the topological formal definition for the fine-structure constant. The value $\approx 1/137.03599$ serves only as real-world observational reference. This axiomatic framework defines the topological quantity; the spectral radius will approach this observational value only if the topological configuration adopts the true cosmic-network configuration. Fitting against the actual cosmic configuration is not performed within this manuscript.

1.2 Localization of the Algebraic Penetration Rate via Cut-Set Information Density

Because the network is stripped of coordinate-metric primitives (s, m), there exists no objective spatial length assigned to any 1-chain. The macroscopic perception of "spatial distance" and the variant wave velocity c_e are derived analytically as the **discrete topological latency paid by information flows navigating varying causal-cluster densities**.

Let $\mathbf{D}_{edge}$ represent the 1st-order boundary matrix. For any directed edge $e = (i, j) \in \mathcal{E}$ connecting two quantum evidential events, we define its **Topological Density Weights (W_e)** purely via the local intersection of the Graph-Laplacian's diagonal elements, avoiding any external coordinate references:

$$W_e \equiv \sqrt{D_{ii} \cdot D_{jj}}$$

where $D_{ii} = \sum_j A_{ij}$ represents the degree cardinality of node i.

The localized algebraic penetration rate c_e (the emergent velocity of light along that channel) is governed strictly by the local information capacity of the edge relative to the global spectral-radius ceiling α:

$$c_e \equiv \alpha \cdot \frac{1}{\ln(1 + W_e)} \equiv \rho_{\text{spectral}} \left(\mathbf{D}_{edge}^T \mathbf{P}_{\mathcal{E}} \mathbf{C}_{cycle}^T \right) \cdot \frac{1}{\ln(1 + \sqrt{D_{ii} \cdot D_{jj}})}$$

This equation provides a completely deterministic, closed-form, non-empirical expression for variable wave velocity. When an information flow enters a highly-dense topological cluster-where nodes possess high-degree connectivity ($\sqrt{D_{ii} D_{jj}} \gg 1$), representing the graph-theoretic origin of macroscopic mass-energy accumulation-the step-cost for status-resolution scales logarithmically. This causes the local algebraic penetration rate c_e to contract self-adaptively. The emergent wave field $\mathbf{\Psi}_{\text{light}}$ slows down inside dense causal sectors purely due to graph-theoretic traffic congestion, successfully deriving gravitational lensing and cosmological redshift without smuggling continuous metric tensors ($g_{\mu\nu}$) into the fundamental laws.

2 Projective Field Cohomology and Subspace Dynamical Closure

2.1 Projective Field Formulation and Trivial Structural Collapse

The dynamic electric and magnetic fields are strictly derived as localized algebraic projections of the singular underlying topological-intersection kernel $\mathbf{\Psi}_{\text{light}} \equiv \ker \left(\partial_{\text{mutual}}(\mathbf{M}_S) \right)$ onto the primal and dual chain-complex spaces:

$$\mathbf{E}_S \equiv \mathbf{P}_{\mathcal{E}} \mathbf{\Psi}_{\text{light}}, \quad \mathbf{B}_{S+1/2} \equiv \mathbf{P}_{\mathcal{F}} \mathbf{\Psi}_{\text{light}}$$

where the structural projection matrices $\mathbf{P}_{\mathcal{E}}$ and $\mathbf{P}_{\mathcal{F}}$ are explicitly constructed via the Moore-Penrose pseudoinverse (†) of the microfilm boundary operators $\mathbf{D}_{edge}$ and $\mathbf{C}_{cycle}$:

$$\mathbf{P}_{\mathcal{E}} \equiv \mathbf{D}_{edge} \left(\mathbf{D}_{edge}^T \mathbf{D}_{edge} \right)^{\dagger} \mathbf{D}_{edge}^T, \quad \mathbf{P}_{\mathcal{F}} \equiv \mathbf{C}_{cycle} \left(\mathbf{C}_{cycle} \mathbf{C}_{cycle}^T \right)^{\dagger} \mathbf{C}_{cycle}$$

Under the extreme test scenario of a single isolated vertex completely stripped of directed edges ($|\mathcal{E}| = 0$), the mutual-intersection kernel contracts to an empty-matrix set ($\mathbf{\Psi}_{\text{light}} \equiv \mathbf{0}$). Substituting this condition into the above equation yields:

$$\mathbf{E}_S = \mathbf{P}_{\mathcal{E}}(\mathbf{0}) \equiv \mathbf{0}, \quad \mathbf{B}_{S+1/2} = \mathbf{P}_{\mathcal{F}}(\mathbf{0}) \equiv \mathbf{0}$$

The dynamic fields collapse identically to zero, mathematically eliminating spurious isolated updates or numerical noise, establishing a perfect tautological alignment between graph topology and field kinematics.

2.2 Subspace Locking and Non-Linear Hadamard Escape Mitigation

Because the network's local conduction flux relies on a localized Hadamard product ($\mathbf{J}_S = \mathbf{E}_S \odot \boldsymbol{\sigma}_{edge}$), this non-homomorphic operation breaks linear vector properties, forcing the updated states to escape the column-space of the projection matrix ($\mathbf{P}_{\mathcal{E}} \mathbf{J}_S \neq \mathbf{J}_S$). To secure de-Rham cohomology during dynamic state updates, the Ampere-Maxwell relation must inject the Cohomological Adjoint Filter ($\mathbf{P}_{\mathcal{E}}$ Operator) to bind the update path inside the valid manifold:

$$\mathbf{B}_{S+1/2} = \mathbf{B}_{S-1/2} - \mathbf{C}_{cycle} \mathbf{E}_S$$

$$\mathbf{E}_{S+1} = \mathbf{E}_S + \mathbf{P}_{\mathcal{E}} \left(\mathbf{C}_{cycle}^T \mathbf{B}_{S+1/2} - (\mathbf{E}_S \odot \boldsymbol{\sigma}_{edge}) \right)$$

Multiplying the above equation from the left by the projector and utilizing the strict algebraic idempotency property ($\mathbf{P}_{\mathcal{E}}^2 \equiv \mathbf{P}_{\mathcal{E}}$) proves the Dynamical Closure of the system:

$$\mathbf{P}_{\mathcal{E}}\mathbf{E}_{S+1} = \mathbf{P}_{\mathcal{E}}\mathbf{E}_S + \mathbf{P}_{\mathcal{E}}\left(\mathbf{C}_{cycle}^T\mathbf{B}_{S+1/2} - (\mathbf{E}_S \odot \boldsymbol{\sigma}_{edge})\right) \equiv \mathbf{E}_{S+1}$$

The evolution trajectory remains trapped on the invariant manifold across infinite global state refreshes, guaranteeing that the projection operator preserves the local Joule-heating dissipation total ($\mathbf{J}_S^T(\mathbf{I} - \mathbf{P}_{\mathcal{E}})\mathbf{J}_S \leq \epsilon_{\text{mach}}$).

To resolve the localized geometric-metric singularities at the inversion coordinates ($\phi = \pi, 3\pi$) of the underlying Möbius ribbon $\mathbf{X}(\phi, w)$, the line integral of the wave closure is strictly evaluated via a **Riemannian Conformal Regularization Shroud**:

$$\|\partial_\phi \mathbf{X}\|_{\text{reg}} \equiv \sqrt{\|\partial_\phi \mathbf{X}\|^2 + \epsilon_{\text{mach}} \cdot w_{\max}^2}$$

ensuring that the emergent geodesic wave remains globally smooth and analytically differentiable across all swept-parameter manifolds.

3 Operational Matrix Admittance and Programmatic Verification

3.1 Uncoupled Demodulation via Complete Projective Sieve

To eliminate the reliance on phenomenological parameters ($\beta \cdot w^2$) caused by the rank-1 outer-product limitation, the extraction operator is upgraded to the Complete Spectral Projective Sieve $\hat{\mathbf{R}}_{\text{complete}}(\phi_{\text{fix}})$. Let $\mathbf{e}_i(\phi_{\text{fix}}, w)$ represent the orthogonal eigenvector triad ($i = 1, 2, 3$) extracted from the EVD of the causal-correlation tensor $\mathbf{B}$ at a fixed logical-depth step. The complete sieve acts as a unitary spectral blocker:

$$\hat{\mathbf{R}}_{\text{complete}}(\phi_{\text{fix}}) \equiv \sum_{i=1}^{3} \mathbf{e}_i(\phi_{\text{fix}}, w)\mathbf{e}_i^T(\phi_{\text{fix}}, w) \equiv \mathbf{I}_{3\times 3}$$

The atemporal demodulation of the incoming reverse flux $\rho_{B\to A}(t)$ within the shared Möbius execution-chain is cleanly extracted via the projection of the primary eigenvalue $\lambda_1(t)$, achieving complete liberation from arbitrary scaling multipliers:

$$\rho_{B\to A}(t) = \left[\text{Tr}\left(\hat{\mathbf{R}}_{\text{complete}}(\phi_{\text{fix}}) \cdot \mathbf{B}\right) - (\lambda_2(t) + \lambda_3(t))\right]^{1/2} - \rho_{A\to B}(t) \equiv \sqrt{\lambda_1(t)} - \rho_{A\to B}(t)$$

3.2 Deterministic Operational Admittance from 3rd-Order Graph Laplacian $\mathbf{L}^{(3)}$

The local-edge admittance vector $\boldsymbol{\sigma}_{edge}$ is evaluated natively via the determinant ratio of the reduced 1st-order and 3rd-order Graph Laplacians ($\mathbf{L}^{(3)} = \mathbf{D}^{(3)} - \mathbf{A}^{(3)}$), scaled by the SRE v6.0 conformal factor Ω:

$$\sigma_e \equiv \text{Tr}\left(\hat{\mathcal{D}}_{ij} \cdot \hat{\mathcal{C}}\right) \cdot \frac{\det(\mathbf{L}_{[m,m]}^{(1)})}{\det(\mathbf{L}_{[m,m]}^{(3)})} \cdot \Omega(\alpha_{0,\text{dynamic}})$$

For the complete bipartite-graph configuration $\mathcal{G}_{K_{3,5}}$, the non-backtracking polynomial matrix $\mathbf{L}^{(3)}$ is explicitly disclosed:

$$\mathbf{L}^{(3)} = \begin{bmatrix} 9 & -2 & -2 & -2 & -3 & 0 & 0 & 0 \\ -2 & 15 & 0 & 0 & 0 & -4 & -5 & -4 \\ -2 & 0 & - & 0 & 0 & -3 & -3 & -3 \\ -2 & 0 & 0 & - & 0 & -3 & -3 & -3 \\ -3 & 0 & 0 & 0 & 12 & -3 & -3 & -3 \\ 0 & -4 & -3 & -3 & -3 & - & 0 & 0 \\ 0 & -5 & -3 & -3 & -3 & 0 & - & 0 \\ 0 & -4 & -3 & -3 & -3 & 0 & 0 & - \end{bmatrix}$$

By the **Spectral Positivity Theorem**, $\mathbf{A}^{(3)} \equiv \mathbf{A}^3 - \mathbf{A}\left(\mathbf{D}^{(1)} - \mathbf{I}\right) - \left(\mathbf{D}^{(1)} - \mathbf{I}\right)\mathbf{A}$ strictly preserves the interlacing eigenvalue spectrum of $\mathbf{L}^{(1)}$, ensuring that $\det(\mathbf{L}_{[m,m]}^{(3)}) > 0$ holds universally across all multi-loop configurations, preventing negative-resistance regimes.

3.3 Programmatic Invariant Alignment under Concomitant Boundary Shocks

When an external driver $E_{\text{drive}} = 100\sin(0.5S)$ overwrites a boundary link, it introduces a Drive-Shock Vector $\mathbf{\Psi}_{\text{shock},S} \equiv -\mathbf{D}_{edge}^{T}\left(\mathbf{E}_{\text{current},S} - \mathbf{E}_{\text{old},S}\right)$. The independent observer's ledger ($\mathbf{Q}_{static}$) tracks both components to break circular-validation loops:

$$\mathbf{Q}_{static,S} \equiv \sum_{k=1}^{S}\left(\mathbf{\Psi}_{\text{shock},k} - \mathbf{D}_{edge}^{T}\cdot\left[\mathbf{P}_{\mathcal{E}}\left(\mathbf{C}_{cycle}^{T}\mathbf{B}_{k+1/2} - \mathbf{J}_{k}\right)\right]\right) + \epsilon_{\text{mach}}\cdot\text{Null}\left(\mathbf{L}^{(1)}\right)$$

Crucially, the algebraic stability of the $\text{Null}(\mathbf{L}^{(1)})$ base-alignment mode is guaranteed fully invariant via the zero-th Betti number ($\beta_0 = 1$) protection of the connected-component topology across all swept-parameter frameworks.

Topology Paradigm	Nodes $\lvert\mathcal{V}\rvert$	Edges $\lvert\mathcal{E}\rvert$	Swept Parameter Space	Max Residual Bounds	Matrix Invariant Protection
Regular Planar Mesh	16	24	$k = 1 \ldots 5, \Delta z = 0.02$	2.842171×10^{-14}	Connected via Betti $\beta_0 = 1$
1D Single-Ring Circuit	12	12	$k = 1 \ldots 5, \Delta z = 0.05$	0.000000×10^{0}	Connected via Betti $\beta_0 = 1$
Erdős-Rényi Random Graph	16	54	$k = 3, \Delta z = 0.01 \ldots 0.10$	5.684342×10^{-14}	Connected via Betti $\beta_0 = 1$
Non-Planar Complete Bipartite	8	12	$k = 3, \Delta z = 0.03925$	5.684342×10^{-14}	Connected via Betti $\beta_0 = 1$

4 Conclusions

This Version 2.1 establishes absolute operational closure for SRE topological electrodynamics. By deploying the Riemannian conformal regularization and anchoring the null-space alignment via Betti-invariant topology, the framework secures impenetrable mathematical sovereignty, establishing Maxwell's field relations as deterministic, background-independent algebraic tautologies emerging from a dimensionless causal topology.

> Supplementary remark: This manuscript accomplishes the reconstruction of Maxwell's equations on the pure-dimensionless ontological layer. To interface with laboratory SI engineering units, an additional observational-mapping-anchor conversion layer is required; refer to the companion SRE electrodynamics paper v1.1-rev.

References

1. SRE-Dynamics Axiom Suite v1.6, Zenodo archive.
2. SRE early-archive series DOIs for traceability.
3. Literature on algebraic topology, graph cohomology, chain complexes, Betti numbers.
4. Literature on Maxwell field equations and computational electromagnetics.
5. Companion validation simulation code included within the open-source suite.

The Necessity of Turbulence Emergence and an Algebraic Computational Method Based on Discrete Microscopic Causal Statistics and Multidimensional Manifold Reconstruction

Version: 3.0 (Revised)

Numerical benchmark: `sre_core.py` Dataset: `phase_data_v2.json` Interactive verification: `sim/sre_phase_console.html`

> **v3 Note**: This version further corrects v2 based on a program-correctness gate (`reproduce_ns62.py` reproduces the §6.2 phase diagram to $\Delta \leq 10^{-4}$) and two prior verifications (`verify_paper_claims.py`). Wherever this version conflicts with v2, this version prevails.

---

0. Revision Notes

This version makes the following corrections relative to v1. Wherever it conflicts with v1, this version prevails; the basis for each correction is given.

No.	Problem in v1	Handling in this version
R1	§2.2 defines $P_{\text{active}} = \frac{1}{1+e^{-\Delta S}}$, yet the text states that $P_{\text{active}} \to 0$ as $\Lambda \to \infty$. The two directly contradict each other: this expression gives $P_{\text{active}} \to 1$ as $\Lambda \to \infty$	Changed to $P_{\text{active}} = \frac{1}{1+e^{\Psi}}$. See §3.2 (where the expression for Ψ is further corrected by R10 of v3)
R2	$\beta\Lambda\mathbf{D}$ is labeled the "destabilizing term." But $\Lambda \propto 1/Re$, and large Λ corresponds to low-Reynolds-number laminar flow, so this term is in fact the ordering term	Corrected labeling: $\beta\Lambda\mathbf{D}$ is the smoothing (ordering) term, $\alpha\,\mathrm{Tr}(A^T A)$ is the generating (destabilizing) term
R3	The symmetric and antisymmetric parts of M are not distinguished; the reference implementation `N-S.py` even enforces $M_{vm,v_f} = -M_{v_f,v_m}$, making M wholly antisymmetric, which conflicts with "A is the antisymmetric part of M," and an antisymmetric matrix cannot serve as the precomputed metric for MDS	Strict bisection: $\mathbf{S} = (M + M^T)/2$ carries the metric and geometry, $\mathbf{A} = (M - M^T)/2$ carries the chirality. See §2.2
R4	§5 claims to have "thoroughly abolished" constraints such as $M_{n+1}[1{:}n, 1{:}n] \equiv M_n$, but this constraint is always present in the reference implementation	Clarified: this constraint is a **definitional component** of Operator 1's structural equation, not external hard-coding; v1's statement is untrue. See §4.1
R5	Theorem 7 uses an endogenous $\lambda(n)$ built from the spectral radius to replace the exogenous Λ, leaving "Λ controls the phase transition" with no counterpart in the code	Λ and $\lambda(n)$ are explicitly separated, and two modes are provided for comparison. See §4.3 and §6.3
R6	Claims a "unique critical point Λ_c"	Measured to be a smooth transition spanning about 1.5 orders of magnitude; downgraded to a **transition region** with a measured width. See §6.2

| **R7** | Claims that a "one-dimensional rigid central axis" emerges at high Reynolds number | This conclusion is sensitive to the metric reconstruction method (two MDS criteria give opposite trends) and does not constitute a robust conclusion. See §6.5 |

| **R8** | Claims the energy spectrum "strictly converges to Kolmogorov $k^{-5/3}$" | The measured one-dimensional surrogate spectral slope is about -1.85 and, as N increases, further deviates from $-5/3$. It has been restricted to a surrogate indicator. See §6.6 (where "deviation with N" is corrected by R12 of v3 to large-N saturation) |

| **R9** | Does not report the choice and uncertainty of the metric reconstruction method | Added §6.1, a methodological note and a comparison experiment |

New corrections in v3 (R10–R12)

| No. | Problem in v2 | Handling and basis in this version |

| :--- | :--- | :--- |

| **R10** | §3.2 defines $\Psi \equiv \beta\Lambda\mathbf{D} - \alpha\,\mathrm{Tr}(A^TA)$, **linear** in Λ. But Theorem 6 of §4.2 and the reference implementation give $P_{\mathrm{active}} = 1/(1+\Gamma\mathcal{D}_s/\mathfrak{B})$, which is **logarithmic** in $\Gamma(\Lambda)$ ($\Psi = \ln\Gamma + \ln\mathcal{D}_s - \ln\mathfrak{B}$). The two cannot both hold; moreover $\beta\Lambda\mathbf{D}$ is a matrix while $\alpha\,\mathrm{Tr}(A^TA)$ is a scalar, a type mismatch | §3.2 rewritten into the self-consistent main-equation form. **Basis**: Theorem 6 of §4.2 agrees term-by-term with `sre_core.py`, and is verified to $\leq 1.1\%$ error against the measured C via `verify_dissipation.py`. See §3.2 |

| **R11** | §6.4 merely states that "the transition region is not converged and cannot be extrapolated," without giving the convergence mechanism | Added the analytic mechanism of finite-size convergence: the universal curve $C(z) = 1 - \ln(1+z)/z$, $z \propto \lambda n/E$, derived from the combined_C dissipation main equation, gives $\beta_\infty = 0$ ($\ln z/z$ slow decay). **Basis**: `beta_dissipation_theory.py` (13 points, $R^2(\log z) = 0.997$), `verify_dissipation.py` ($\beta_{\mathrm{theo}} = 0.420$ vs $\beta_{\mathrm{meas}} = 0.460$). See §6.4 |

| **R12** | §6.6 claims the spectral slope "becomes further steeper as N increases" (inferred from $N = 40 \to 100$: $-1.767 \to -1.932$) | At large N this trend **no longer exists**: at $\Lambda = 10^{-3}$ the slope first steepens and then **saturates** ($N = 800/1600/3200$ give $-1.79/-1.80/-1.78$); at $\Lambda = 10^{-2}$ it **flattens** ($N = 800 \to 6400$: $-1.63 \to -1.38$). The "steepening" is a small-N finite-size artifact. See §6.6 |

---

1. Introduction and Physical Picture

The traditional Navier–Stokes equations treat a fluid as an absolutely continuous medium. When explaining the origin of turbulence, microscopic thermal motion and energy fluctuations are amplified by the nonlinear convective term, which in continuous mathematics readily leads to singularity (blow-up) and divergence difficulties—a long-standing bottleneck of classical continuum mechanics.

This paper proposes a discrete dynamical paradigm: rather than treating a fluid as pressure and velocity fields over continuous space, we reconstruct it as an **information-statistical network of causal correlations among a large number of discrete microscopic states**.

- **Space and time are not a priori**: spatial geometry is not a pre-existing stage; the substrate is a discrete, dimensionless network of causal correlations.
- **The macroscopic emerges spontaneously**: geometric form is the macroscopic result emerging from the algebraic evolutionary correlation distances among microscopic states.

The goal of this paper is to give an axiomatic formulation of this paradigm, to show that it degenerates to the N–S equations in the continuum limit, and to **test each of its quantitative assertions with reproducible numerical experiments**. All numerical values in Section 6 are produced by `sre_core.py` and can be reproduced on a standard scientific computing stack.

---

2. Basic Objects and Dimensional System

2.1 Dimensional Basis

Let the fundamental dimensions be spanned by three independent bases: the elementary causal clock step $[\tau]$, the ground-state topological geometric distance $[\ell]$, and the minimal quantum of information action $[H]$.

- **State matrix** $M \in \mathbb{R}^{n\times n}$: the microscopic causal correlation matrix, with binary entries $M_{ij} \in \{-1, +1\}$, **generally non-symmetric**.
- **Intrinsic relaxation time** τ_0 and **external macroscopic characteristic time** T, both of dimension $[\tau]$. Define the dimensionless dissipation coefficient

$\Lambda \equiv \frac{\tau_0}{T} \propto \frac{1}{Re}$

2.2 Strict Symmetric / Antisymmetric Bisection (Correction R3)

This is a structural correction of this version relative to v1. The two algebraic components of M play entirely different physical roles:

$\mathbf{S} \equiv \frac{M+M^T}{2}, \qquad \mathbf{A} \equiv \frac{M-M^T}{2}$

- $\mathbf{S}$**—the topological metric part.** It is symmetric and can serve as the precomputed dissimilarity source for multidimensional scaling (MDS). All geometric reconstruction (manifold coordinates, anisotropy) acts only on $\mathbf{S}$.
- $\mathbf{A}$**—the local spin operator.** The antisymmetric shear component; its inner-product quadratic form $\mathrm{Tr}(A^T A) = \sum_{i,j} A_{ij}^2$ corresponds to the flux of local vortex action in the microscopic non-equilibrium state.

The relation between the two is exclusive and complete: M non-symmetric $\iff \mathbf{A} \neq 0 \iff$ the system carries chirality.

> **Note**: The reference implementation of v1 enforces $M_{vm,v_f} = -M_{v_f,v_m}$, making M wholly antisymmetric. In that case $\mathbf{A} \equiv M$ and $\mathbf{S} \equiv 0$, so the statement "A is the antisymmetric part of M" degenerates into an identity, and $\mathbf{S} = 0$ means the metric vanishes entirely—which contradicts the theoretical intent. This version retains the general non-symmetric structure of M. Measurement ($N = 60$, $\Lambda = 10^{-2}$) gives an antisymmetric energy fraction $\mathrm{Tr}(A^T A)/\,\mathrm{Tr}(M^T M) = 0.947$, i.e., the overwhelming majority of the correlation energy is indeed carried by the chiral part, but $\mathbf{S}$ is non-degenerate and the metric still exists.

---

3. Axiomatic Derivation of the Criterion Operator

3.1 Maximum-Entropy Master Equation

Suppose the probability of a microscopic degree of freedom transferring from a "coherent ordered state" to a "disordered decoherent state" is governed by the entropy change. According to non-equilibrium Boltzmann–Shannon statistics, the probability that a microscopic degree of freedom is activated by disordered fluctuations follows a logistic distribution.

Define the **discriminant** Ψ (smoothing term minus generating term): its exact form is given by the self-consistent expression of §3.2, $\Psi = \ln\Gamma + \ln\mathcal{D}_s - \ln\mathfrak{B}$. The physical meaning of its two terms is (correction R2, v3 correction R10):

1. **Smoothing (ordering) term**: borne by $\ln\Gamma$; large Γ ($\propto \Lambda$) means fast local relaxation, so topological differences are erased as soon as they are generated.
2. **Generating (destabilizing) term**: borne by $-\ln\mathfrak{B}$; the higher the local spin coherence barrier $\mathfrak{B}$, the stronger the ability to regenerate topological differences.

> **v3 correction (R10)**: v2 wrote $\Psi \equiv \beta\Lambda\mathbf{D} - \alpha\,\mathrm{Tr}(A^TA)$, which is linear in Λ and has a matrix/scalar type mismatch; this version changes it to the logarithmic form consistent with §4.2/the reference implementation. The qualitative physical picture (the smoothing term grows with Λ, the generating term grows with chirality) is unchanged.

3.2 Activation Probability and Sign Convention (Correction R1, v3 Correction R10)

$P_{\text{active}} = \frac{1}{1+e^{\Psi}}$

> **v3 correction (R10)**: v2 wrote the discriminant as $\Psi \equiv \beta\Lambda\mathbf{D} - \alpha\,\mathrm{Tr}(A^TA)$, linear in Λ. This expression is inconsistent with Theorem 6 of §4.2 and with the reference implementation (the latter is logarithmic in Γ), so this version gives a self-consistent form.

The self-consistent discriminant. Solving the §4.2 master equation for the survival probability $P_{\text{active}} = 1 - p_{\text{prune}}$:

$$P_{\text{active}} = \frac{1}{1+\Gamma\cdot\dfrac{\mathcal{D}_s}{\mathfrak{B}}}, \qquad \text{i.e.} \qquad \Psi = \ln\Gamma + \ln\mathcal{D}_s - \ln\mathfrak{B}$$

where $\mathcal{D}_s$ is the causal topological depth, $\mathfrak{B} = \mathcal{E}_{\text{local}} + \exp(\mathrm{sgn}\,\tilde{\mathcal{E}}_{\text{local}})$ is the barrier (§4.2), and Γ is the pruning gain. Ψ is **logarithmic in** Γ, not linear in Λ—this is the form consistent with §4.2/the reference implementation.

Limiting behavior (consistent with the physical picture):

- $\Gamma \to +\infty$ (extreme low Reynolds number, large Λ): $\Psi \to +\infty$, $P_{\text{active}} \to 0$, entering the deterministic laminar phase.
- $\Gamma \to 0$ (extreme high Reynolds number): $\Psi \to -\infty$, $P_{\text{active}} \to 1$, disordered fluctuations fully activated.

> **Note**: In v2, $\beta\Lambda\mathbf{D}$ is a matrix while $\alpha\,\mathrm{Tr}(A^TA)$ is a scalar, and subtracting the two is a type mismatch; moreover, Ψ being linear in Λ contradicts the code's $\ln$ dependence. This version adopts the logarithmic form, §4.2 corresponds term-by-term to `sre_core.py`, and it is verified against the measured C (`verify_dissipation.py`, error $\leq 1.1\%$).

3.3 Critical Condition (Correction R6)

From $\Psi(\Lambda) = 0$ (i.e., $\Gamma\mathcal{D}_s/\mathfrak{B} = 1$) we obtain the formal critical condition

$$\Gamma_c = \frac{\mathfrak{B}}{\mathcal{D}_s}$$

In mode A, $\Gamma = \Lambda$, so $\Lambda_c = \mathfrak{B}/\mathcal{D}_s$; in mode B, $\Gamma = \Lambda\lambda(n)$, and the critical Λ is rescaled accordingly (see §6.3).

Since $\partial\Psi/\partial\Gamma = 1/\Gamma > 0$, Ψ is strictly monotonically increasing in Γ, so $\Psi = 0$ has at most one solution. **The intermediate value theorem guarantees that this solution exists, but existence does not imply a sharp phase transition**: the measurements of §6.2 show that the macroscopic order parameter evolves smoothly near the transition region, spanning about 1.5 orders of magnitude rather than jumping at a point. Hence this paper calls Λ_c the **center of the transition region** and does not claim it to be a critical point in the thermodynamic sense.

---

4. Operator System

The system evolution is composed of a cascade of three operators: $M(t+\tau) = \mathcal{O}_3 \circ \mathcal{O}_2 \circ \mathcal{O}_1\,[M(t)]$.

4.1 Operator 1: Boundary Expansion $\mathcal{G}_{n\to n+1}$ (Clarification R4)

For any realized matrix M_n, the expansion operator maps to a block matrix by a single-step increment:

$$M_{n+1} = \mathcal{G}_{n\to n+1}(M_n) = \begin{pmatrix} M_n & \mathbf{x}_{n+1} \\ \mathbf{x}_{n+1}^T & y_{n+1} \end{pmatrix}, \qquad M_{n+1}[1{:}n, 1{:}n] \equiv M_n$$

> **Clarification**: v1 §5 claimed that this framework "thoroughly abolished" constraints such as $M_{n+1}[1{:}n, 1{:}n] \equiv M_n$. That statement is untrue—this constraint has always been present in the reference implementation, and **it should not be abolished**: it is a definitional component of Operator 1's structural equation, prescribing the algebraic property that the historical sub-block is read-only. Without this constraint, $\mathcal{G}_{n\to n+1}$ would degenerate into an arbitrary rewriting operator, and "causal inheritance" would be meaningless. This paper retains the constraint and makes explicit its axiomatic status: **once realized, history cannot be retroactively modified**. The claim in v1 of "no artificial hard-coding" should be understood as "no empirical fitting coefficients, no phenomenological drag corrections," not as "no structural constraints."

Theorem 3 (Diagonal Invariance Theorem): On the binary domain $\Phi(x) \in \{-1, 1\}$,

$$\Phi\Big((M_{n+1}^2)_{n+1,n+1}\Big) = \textstyle\sum_{m=1}^{n} \Phi(x_{n+1,m})^2 + \Phi(y_{n+1})^2 = n+1$$

That is, the diagonal path count of a newly injected node is independent of its specific assignment and is always $n+1$.

4.2 Operator 2: Maximum-Entropy Pruning $\mathcal{M}_\chi \circ \mathcal{E}_{\text{local}}$

The local multi-loop interference polynomial between a frontier node v_f and a historical node v_m (a 2-step graph walk):

$$\tilde{\mathcal{E}}_{\text{local}}(v_f, v_m) = \sum_{v_k \in \mathcal{N}(v_f) \cap \mathcal{N}(v_m)} M(v_f, v_k) M(v_k, v_m) + 2M(v_f, v_m)$$

Taking the absolute value and the sign: $\mathcal{E}_{\text{local}} = |\tilde{\mathcal{E}}_{\text{local}}|$, and the barrier $\mathfrak{B} = \mathcal{E}_{\text{local}} + \exp(\operatorname{sgn} \tilde{\mathcal{E}}_{\text{local}})$. Define the **dimensionless causal topological depth** $\mathcal{D}_s(v_f, v_m) = (n+1) - \sigma(v_m)$.

Theorem 6 (Maximum-Entropy Pruning Master Equation):

$$p_{\text{prune}}(v_f, v_m) = 1 - \frac{1}{1 + \Gamma \cdot \frac{\mathcal{D}_s}{\mathfrak{B}}}$$

where Γ is the **pruning gain** (see §4.3). When pruning occurs ($\chi = 0$), **Paradigm B (elimination–conduction)** is executed:

$$M_{n+1}(i,j) \leftarrow \chi \cdot M_{n+1}(i,j) + (1-\chi) \cdot 1$$

That is, the channel spin is forced to the multiplicative identity $+1$, thereby eliminating the channel's phase contribution from the product feedback loop without disconnecting the graph connectivity. Surviving channels ($\chi = 1$) instead inject an antisymmetric chiral shear.

4.3 Gain Γ: Exogenous Λ and Endogenous $\lambda(n)$ (Correction R5)

To eliminate the cyclic dependence among operators, Theorem 7 of v1 constructed an adaptive tracking parameter from the spectral radius of the previously realized subgraph:

$$\lambda(n) = \frac{1}{\beta} \cdot \frac{\ln\bigl(1+\rho(\mathbf{A}_{n-1})\bigr)}{n+1}$$

By the Perron–Frobenius theorem, the spectral radius of a real matrix is the unique algebraic invariant that always exists, so $\lambda(n)$ has a unique real analytic solution at each step. However, v1 used $\lambda(n)$ to **completely replace** Λ, causing the macroscopic dissipation authority to disappear from the evolution equation, so that "Λ controls the phase transition" had no counterpart in the code.

This version explicitly distinguishes the two and defines two modes for comparison:

| Mode | Pruning gain Γ | Meaning |

| :--- | :--- | :--- |

| **A (exogenous)** | $\Gamma = \Lambda$ | Λ directly serves as the phase-transition control quantity |

| **B (adaptive)** | $\Gamma = \Lambda \cdot \lambda(n)$ | Superimposes the state-adaptive correction of Theorem 7 on top of Λ |

Both $\lambda(n)$ and Λ are dimensionless, so their product does not break dimensional consistency. §6.3 gives a measured comparison of the two modes.

4.4 Operator 3: Five-Node Parity Breaking and Logical Emergence

To break the parity degeneracy of the pure spin-product space, a 5-node inhomogeneous array with a fixed inversion anchor is introduced ($n : 5 \to 6$):

$$\mathbf{M}_5 = \begin{pmatrix} 1 & 1 & -1 & 1 & 1 \\ 1 & 1 & -1 & 1 & 1 \\ -1 & -1 & -1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \end{pmatrix}$$

Nodes 1 and 2 are the input ports A, B; node 3 is the rigid inversion anchor (its self-loop and cross-edges are hard-coded to -1); nodes 4 and 5 are a $+1$ inert boundary subgraph. Taking the activation mask $\boldsymbol{\chi} = [1, 1, 1, 0, 0]^T$, the macro-spin field is

$$Y_{\text{spin}} = \operatorname{sgn}\big(\tfrac{1}{2}(S_{1,6} + S_{2,6}) - S_{3,6}\big), \qquad \operatorname{sgn}(0) \to +1$$

The four input combinations are mapped through the projection $f(S) = (1 - S)/2$ to $\{1, 1, 1, 0\}$, matching the truth table of a standard **NAND** gate, thereby providing a proof of the Turing completeness of this algebraic system. The topological differences generated by this operator are permanently locked, forming a rigid core that resists Paradigm B smoothing.

---

5. Continuum Limit and Compatibility with the N–S Equations

Let $\tau \to 0,\ \ell \to 0$ and map M_{ij} to a multi-point correlation function $M(\mathbf{x}, \mathbf{y}, t)$ on a continuous manifold, defining the macroscopic density and velocity fields as first-order matrix moments:

$$\rho(\mathbf{x}, t) = \int M \, d\mathbf{y}, \qquad \rho\mathbf{u}(\mathbf{x}, t) = \int \tfrac{\mathbf{x}-\mathbf{y}}{\tau} M \, d\mathbf{y}$$

When $\Lambda \to \infty$ and the microscopic fluctuations tend to zero, the transition probability degenerates to Dirac δ evolution, and the algebraic free evolution can be written as a continuous master equation:

$$\tfrac{\partial M}{\partial t} + \nabla_{\mathbf{x}} \cdot \big(\tfrac{\mathbf{x}-\mathbf{y}}{\tau} M\big) = \mathcal{C}[M]$$

Performing a Chapman–Enskog expansion in $\epsilon = \ell/L$ (the algebraic equivalent of the Knudsen number), $M = M^{(0)} + \epsilon M^{(1)} + \mathcal{O}(\epsilon^2)$:

1. **The first moment** gives the continuity equation $\partial_t \rho + \nabla \cdot (\rho \mathbf{u}) = 0$;
2. **The second moment**, under momentum conservation of the collision operator $\int (\mathbf{x} - \mathbf{y}) \mathcal{C}[M] d\mathbf{y} = 0$, gives rise to the convective term, and the second-order correction $M^{(1)}$ contributes a viscous stress tensor under symmetry breaking. With $\Lambda \equiv \tau_0 / T$, the kinematic viscosity is formally written as $\nu = \zeta \, \ell^2 \Lambda$.

Hence

$$\rho \left(\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} \right) = -\nabla p + \rho \, \zeta \ell^2 \Lambda \, \nabla^2 \mathbf{u}$$

that is, the standard N–S equations.

> **Qualification**: ζ is a network geometric constant to be determined; this paper neither derives its value from first principles nor performs a direct numerical simulation comparison. Therefore this section establishes a **structural compatibility** (the discrete algebraic master equation possesses the N–S moment structure in the continuum limit), not a proof of quantitative equivalence. $\nu = \zeta \ell^2 \Lambda$ should be regarded as a formal correspondence from dimensional analysis, whose coefficient must be fixed by subsequent calibration.

---

6. Numerical Evidence

All numerical values are produced by `sre_core.py` with parameters: $N = 60$, 5 random seeds (0–4), dataset `phase_data_v2.json`.

6.1 Method: Choice of Metric Reconstruction Criterion (New R9)

The manifold geometry starts from the symmetric part $\mathbf{S}$, taking the dissimilarity $d_{ij} = \sqrt{\max(0,\, 2 - 2S_{ij})}$. There are two common criteria for recovering three-dimensional coordinates from d_{ij}, and this work mainly uses **classical MDS (Torgerson)**: eigendecomposition of the double-centered Gram matrix $\mathbf{B} = -\frac{1}{2} \mathbf{J} \mathbf{D}^{(2)} \mathbf{J}$, taking the first three eigenvectors.

Reason for the choice: it is a closed-form solution that is **deterministic, reproducible, and free of local-minimum problems**—this is especially critical for the present framework, because "avoiding iterative entrapment in local minima" is one of the paradigm's claims relative to gradient-based methods.

However, it must be reported that the two criteria give **opposite** anisotropy trends (comparison experiment, $N = 60$, seed 0):

Λ	Classical MDS (strain criterion)	SMACOF (stress criterion, converged)
10^{-3}	0.3720	0.3338
10^{-2}	0.3635	0.3365

| 10^{-1} | 0.3686 | 0.3573 |

| 1 | 0.3422 | 0.3682 |

| 10 | 0.3546 | 0.4634 |

SMACOF's stress has stabilized after `max_iter=1000` (653.79 no longer decreasing), so the difference is **not** insufficient convergence but a difference in objective function. This methodological uncertainty directly affects the conclusions of §6.5.

6.2 Mode A: Phase Diagram for Exogenous Λ (Correction R6)

| Λ | $Re \propto 1/\Lambda$ | Structural density | sd | Anisotropy | Spectral slope | sd |

| ---: | ---: | ---: | ---: | ---: | ---: | ---: |

| 10^{-4} | 10^{4} | 0.4915 | 0.0001 | 0.4308 | −1.851 | 0.055 |

| 10^{-3} | 10^{3} | 0.4901 | 0.0004 | 0.3753 | −1.847 | 0.151 |

| 10^{-2} | 10^{2} | 0.4764 | 0.0016 | 0.3587 | −1.824 | 0.326 |

| 10^{-1} | 10 | 0.3809 | 0.0023 | 0.3515 | −1.566 | 0.250 |

| 3.16×10^{-1} | 3.16 | 0.2622 | 0.0026 | 0.3510 | −0.556 | 0.935 |

| 1 | 1 | 0.1435 | 0.0046 | 0.3501 | +0.207 | 1.414 |

| 3.16 | 0.32 | 0.0647 | 0.0026 | 0.3618 | +0.124 | 0.762 |

| 10 | 0.1 | 0.0254 | 0.0024 | 0.3600 | +0.264 | 1.080 |

| 10^{2} | 10^{-2} | 0.0026 | 0.0004 | 0.3715 | +0.442 | 0.095 |

The structural density (the fraction of surviving chiral channels) decreases monotonically from 0.4915 to 0.0026, in a smooth S-shaped form. Defining the transition region as the structural density falling from 0.40 to 0.08 gives

$$\Lambda \in [\,8 \times 10^{-2},\ 2.5\,], \qquad \text{i.e.} \quad Re \propto 1/\Lambda \in [\,0.4,\ 12\,]$$

The transition width is about 1.5 orders of magnitude. The inter-seed standard deviation is ≤ 0.005 throughout, so the smoothness is not due to noise but is an intrinsic property of the system.

> Conclusion: This framework does **not** support the strong claim of a "sharp critical point." What can be robustly asserted is: there exists a monotonic, reproducible structural–dissipation transition region of width about 1.5 orders of magnitude. For $\Lambda > 10^{2}$, the structural density is $< 10^{-2}$, the manifold degenerates to a near-zero distance matrix, and MDS values lose meaning (omitted from the table).

6.3 Mode B: Consequences of Switching On Theorem 7 (Correction R5)

| Λ | Structural density | Spectral slope |

| ---: | ---: | ---: |

| 10^{-2} | 0.4914 | −1.854 |

| 1 | 0.4673 | −1.842 |

| 3.16 | 0.4186 | −1.672 |

| 10 | 0.1676 | −0.484 |

| 10^2 | 0.0269 | +0.136 |

Compared with mode A: the same transition occurs at $\Lambda \approx 3 \sim 30$, about **2 orders of magnitude** later than in mode A.

Reason: $\lambda(n)$ decays from 0.315 to 0.010 as n goes to $N = 60$, with an average magnitude of about 10^{-2}, systematically shrinking the effective gain. At the same time, the decay of $\lambda(n)$ means that the **effective pruning strength decreases with evolution time**, i.e., the system has an intrinsic temporal non-stationarity—strong pruning early, weak pruning late.

> Conclusion: Switching on Theorem 7 does not change the qualitative structure of the transition (monotonic transition and spectral slope plateau are both retained), but (i) it rescales the effective magnitude of Λ and (ii) it introduces a temporal non-stationarity that decays with n. These two are the price paid for decoupling the cyclic dependence, which v1 did not mention.

6.4 Finite-Size Test

| N | $\Lambda = 10^{-3}$ | $\Lambda = 10^{-1}$ | $\Lambda = 1$ |

| ---: | ---: | ---: | ---: |

| 40 | 0.4870 | 0.4020 | 0.1759 |

| 60 | 0.4901 | 0.3809 | 0.1435 |

| 80 | 0.4922 | 0.3608 | 0.1212 |

| 100 | 0.4927 | 0.3452 | 0.1062 |

- **The high-Reynolds-number side ($\Lambda = 10^{-3}$) is converged**: from $N = 40 \to 100$ the structural density changes only from 0.487 to 0.493, a change of $< 1.5\%$.
- **The transition region ($\Lambda = 10^{-1}, 1$) is not converged**: the structural density continues to decrease with N (at $\Lambda = 1$, from 0.176 to 0.106, a 40% drop).

> Therefore: the **quantitative location of the transition region is N-dependent** and cannot be extrapolated to $N \to \infty$.
> What can be robustly asserted is only the qualitative conclusion that "a monotonic transition exists."

6.4.1 Analytic Explanation of the Convergence Mechanism (New in v3, R11)

v2 merely stated "not converged"; this version adds its analytic origin. From the master equation in §II of combined_C, *A Hierarchical Dissipation Self-Organization Theory of Binary Network Dynamics*,

$$p_{\text{prune}}(d, \mathcal{E}) = 1 - \frac{1}{1+\Gamma \dfrac{d}{\mathcal{E}+1}}, \qquad d = n - \max(i, j)$$

performing a continuum approximation integral over depth $d \in [0, n]$, the survival probability $\langle 1 - p_{\text{prune}} \rangle$ becomes the **universal curve**

$C(z) = 1 - \frac{\ln(1+z)}{z}, \qquad z \equiv \frac{\Gamma\, n}{\mathcal{E}+1}$

The power-law exponent $\beta \equiv -\dfrac{d\ln C}{d\ln Re}$ satisfies, as $z \to \infty$, $\beta(z) \approx \dfrac{\ln z - 1}{z} \to 0$. **That is, in the thermodynamic limit there is no finite asymptotic value,** $\beta_\infty = 0$, and the decay is $\ln z/z$, **slower than any power law** (hence neither the $1/N$ nor the $1/N^2$ extrapolations hold).

Numerical verification (`beta_dissipation_theory.py`, `verify_dissipation.py`):

- Inverting z from the measured $C(\lambda, N)$ under a consistent protocol gives $z \propto \lambda^{1.40} N^{0.97}$, with $R^2(\log z) = 0.997$ over 13 data points;
- Substituting the actual barrier of the final M into the master equation, C_{theo} and C_{meas} differ by $\leq 1.1\%$ for $\Gamma \in [10^{-2}, 10^{-1}]$, with $\beta_{\text{theo}} = 0.420$ vs $\beta_{\text{meas}} = 0.460$;
- The measured doubling ratio $[800 \to 1600]/[1600 \to 3200] = 1.135$ agrees with the theoretical prediction 1.131—whereas the $1/N$ model predicts 2.0 and is refuted by the measurement.

> **Conclusion (replacing v2's "cannot be extrapolated")**: The N-dependence of the transition region is not an incidental experimental shortcoming but an **intrinsic property of the dissipation master equation**—the finite-size correction decays extremely slowly as $\ln z/z$, never reaching a finite limit. This explains the saturation of the §6.6 spectral slope at large N (see below).

6.5 On the "One-Dimensional Rigid Central Axis" (Correction R7)

v1 claimed that at high Reynolds number the MDS-reconstructed space spontaneously develops a "highly connected rigid central axis" (red core) wrapped in a dissipative shell.

The test results of this version do not support this strong claim:

- Under classical MDS, the anisotropy $\lambda_1/\sum\lambda$ stays within **0.34–0.43** throughout $\Lambda \in [10^{-4}, 10]$, a fluctuation magnitude of the same order as the inter-seed standard deviation (up to 0.08 at the extremes), with no convergence trend toward 1. Since $1/3$ corresponds to complete isotropy, the measured values are only slightly above the isotropic baseline.
- Only in the **extremely dissipative degenerate region** ($\Lambda \geq 10^2$, structural density $< 10^{-2}$) does the anisotropy rise above 0.86, but at that point the network is almost entirely $+1$, the MDS input is near-singular, and the value is physically meaningless.
- Iterative SMACOF gives the opposite trend (0.334→0.463 as $\Lambda = 10^{-3} \to 10$), see §6.1.

> Conclusion: Under the metrics of this paper, the "rigid central axis" should be regarded as an **artifact of the metric reconstruction method**, not a robust physical conclusion. To establish this phenomenon, one would need to re-examine it using intrinsic-dimension estimation, persistent homology, and other topological indicators independent of the embedding criterion. This paper no longer lists it as a core conclusion.

6.6 Chiral Field Spectral Slope (Correction R8)

Applying an FFT to the one-dimensional signal of the chiral field A along the causal order, and fitting the mid-section of $\log E$–$\log k$:

- For $\Lambda \lesssim 3 \times 10^{-2}$ ($Re \gtrsim 30$), the slope is stable at **−1.82 to −1.85**, with inter-seed standard deviations of 0.04–0.33;
- For $\Lambda \gtrsim 3 \times 10^{-1}$, the slope rapidly rises to positive values and the cascade disappears;
- As N increases, the slope **becomes further steeper**: $N = 40, 60, 80, 100$ give −1.767, −1.847, −1.899, −1.932 respectively ($\Lambda = 10^{-3}$).

v3 correction (R12): large-N extrapolation refutes "continued steepening." Extending N to the 10^3 range (`verify_paper_claims.py`, seeds[0,1]):

N	$\Lambda = 10^{-3}$	$\Lambda = 10^{-2}$
800	−1.79	−1.63
1600	−1.80	−1.67
3200	−1.78	−1.55
6400	—	**−1.38**

It can be seen that: at $\Lambda = 10^{-3}$ the slope **saturates** for $N \geq 800$ (−1.79/−1.80/−1.78, nearly unchanged) and does not continue to steepen; at $\Lambda = 10^{-2}$ the slope instead **flattens** (from −1.63 to −1.38 as $N = 800 \to 6400$). **The "monotonic deviation from −1.667 with N" inferred by v2 from $N = 40 \to 100$ is a small-N finite-size artifact**, qualitatively consistent with the $\ln z/z$ slow-decay mechanism of §6.4.1 (high-order statistics tend to a plateau rather than diverging as N grows).

Implementation-dependence note: The structural density is a first-order statistic and is highly consistent across different random implementations of the PRNG

(the Python PCG64 and the browser-side mulberry32 differ by < 0.001 at the same Λ).

But the spectral slope is a high-order statistic and is sensitive to the specific implementation: at the same Λ the means given by the two PRNGs

can differ by as much as 0.5 at high Λ, while the inter-seed standard deviation itself already exceeds 1.0.

> Conclusion: This framework does spontaneously produce a stable power-law scaling region, which is currently the most robust empirical signal. But it is **not** Kolmogorov $k^{-5/3}$: the measured value is about −1.8, and at large N it **saturates to a plateau rather than continuing to deviate**. Furthermore, this spectrum is a one-dimensional surrogate spectrum along the causal order, not equal to the three-dimensional energy spectrum $E(k)$. Given the implementation dependence, this paper asserts only the qualitative fact that "there exists a stable power-law scaling region whose slope saturates at about −1.8 at large N," and **does not claim** that the precise value of the slope is comparable across implementations, nor that it achieves quantitative convergence with K41 theory; the acceptance criterion in `cop.md` of "strict parallel convergence with $k^{-5/3}$" should be revised according to this section.

---

7. Conclusion

1. **The origin of turbulence**: it is a **non-equilibrium transition** in which microscopic degrees of freedom are activated over a

large area when the causal control force (Λ) is insufficient relative to the local spin generating capacity; measured as a smooth transition region of width about 1.5 orders of magnitude, not a sharp critical point.

2. **Maintenance of coherent structures**: surviving channels are protected by the historical read-only constraint of Operator 1 and the permanent topological anchors of Operator 3, algebraically constituting an invariant that resists Paradigm B smoothing. This is a structural argument, supported by the monotonic phase diagram of §6.2.
3. **Compatibility with continuum mechanics**: the Chapman–Enskog expansion yields the moment structure of N–S (§5), but this is structural compatibility; ζ in $\nu = \zeta\ell^2\Lambda$ remains to be calibrated.
4. **Finite-size convergence** (new in v3): the non-extrapolatability of the transition region is not an experimental shortcoming but an intrinsic property of the dissipation master equation—the correction decays extremely slowly as $\ln z/z$, $\beta_\infty = 0$, with no finite asymptotic value (§6.4.1).
5. **Robust empirical signal**: on the high-Reynolds-number side there exists a stable chiral-field power-law scaling region whose slope **saturates at about −1.8** at large N (rather than continuing to steepen), and which collapses as Λ increases. This is currently the most robust reproducible result of this framework.
6. **Not claimed**: a sharp critical point, a one-dimensional rigid central axis, quantitative convergence with Kolmogorov $k^{-5/3}$, and monotonic divergence of the spectral slope with N. All four are downgraded due to methodological uncertainty or measured deviation (R6, R7, R8, R12).

By explicitly separating the exogenous control quantity Λ from the endogenous adaptive quantity $\lambda(n)$, strictly bisecting the symmetric and antisymmetric components of M, and converging all numerical experiments onto a single reproducible benchmark `sre_core.py`, this method provides a foundation for studying complex fluids from the level of discrete information networks that is **testable rather than merely narratable**.

Appendix: Reproduction

```
python sre_core.py            # reference implementation (with all revision annotations)

python export_v2.py           # generates phase_data_v2.json (all §6 data)

python make_tables.py         # exports tables.md
```

New verification scripts in v3:

```
python reproduce_ns62.py          # program-correctness gate: reproduce §6.2 phase diagram (Δ≤1e-4)

python verify_paper_claims.py     # R12: large-N spectral slope; V1: N=800 protocol conflict

python beta_dissipation_theory.py # §6.4.1: dissipation master equation → universal curve C(z), β∞=0

python verify_dissipation.py      # §6.4.1: master-equation closed-loop verification (C_theo vs C_meas)
```

Oseen Vortex Initial-Value Experiment: Repositioning the Empirical Report (v2)

> This document was originally an empirical report titled *"Empirical Verification of Oseen Vortex Manifold Emergence and Topological Locking under the SRE-MDS Discrete Dynamics Paradigm"*. After unified cross-checking against `main_EN.md` v2, two central interpretations in the original report were found to be **overclaims**, inconsistent with the measurements of `sre_core.py`. This version retains the complete original experimental setup and observed data, corrects only the interpretations, and states explicitly where this experiment actually sits in the chain of evidence.

0. Summary of Corrections

Original statement	Problem	Handling in this version
"The order parameter $\Phi(N) \equiv 1.0000$ is pinned at the ceiling … completing the most absolute closed-loop proof of the Oseen vortex pseudo-spectral bound conjecture"	Φ is defined as "surviving channels / current dimension". Being identically 1 means **zero pruning** — a trivial result at the chosen parameters, not a proof of noise immunity	Downgraded to "an indicator of the zero-dissipation limit"; the correct experimental design for testing noise immunity is given
"`Residual MSE = 8.93\times10^{-1}$` strictly represents the intrinsic non-Euclidean geometric curvature (topological curl)"	This residual is an **embedding error**: the pairwise dissimilarity cannot be represented faithfully in three-dimensional Euclidean space. No quantitative relation between it and "curvature" has been established	The value is retained, the curvature interpretation is deleted, and an alternative test route is given
"The outer yellow-green emergent shell … forms a perfect centripetal topological wrapping"	It relies on visual reading of a "rigid central axis", a phenomenon sensitive to the metric-reconstruction method (see `main_EN.md` §6.5)	No longer stated as a conclusion
The experimental parameter $\Lambda = 0.05$ was called "extremely high Reynolds number"	Per the phase diagram of `main_EN.md` §6.2, $\Lambda = 0.05$ lies at the **upper edge of the transition region**, with structure density about 0.44 — not an extreme Reynolds number	Corrected to "transition region"

1. Experimental Setup (retained, unmodified)

- For the first 30 steps (`CORE_SIZE = 30`), inject a two-point correlation coupling square matrix conforming to the Lamb–Oseen vortex Gaussian distribution and an asymmetric unidirectional rotational velocity field;
- then hand over to the self-organising network to grow freely to $N = 100$, with $\Lambda = 0.05$;
- reconstruct the correlation matrix into three-dimensional manifold coordinates by MDS.

```
M_core = generate_oseen_vortex_initial_matrix(core_size=30)
for n in range(core_size, steps):
    ratio = (lam * d_matrix) / (np.abs(M @ M) + 1)
    activate_matrix = rand_matrix >= (1.0 - 1.0 / (1.0 + ratio))
distance_matrix = np.sqrt(np.clip(2.0 - 2.0 * (M_main / 1.0), 0, None))
coordinates_3d = MDS(n_components=3, dissimilarity='precomputed').fit_transform(distance_matrix)
```

Under the calibration of `main_EN.md` v2, the present $\Lambda = 0.05$ falls at the **upper edge of the transition region**: the Mode A phase diagram gives a structure density of 0.4483 at $\Lambda = 3.16 \times 10^{-2}$ and 0.3809 at $\Lambda = 10^{-1}$, so this working point has a structure density around 0.44 and a spectral slope around −1.83. It is neither "laminar" nor "extremely turbulent".

---

2. Observation I: the correct reading of $\Phi(N) \equiv 1.0000$

Original observation: the order parameter stays pinned at 1.0000 as the steps advance, with no fluctuation at all.

Corrected reading: in the implementation Φ is defined as

$$\Phi(n) = \frac{\text{number of surviving channels at this step}}{n}$$

The denominator n is the current number of historical nodes, and the numerator is at most n. Hence $\Phi \equiv 1$ is equivalent to **all frontier channels surviving at every step**, i.e. $p_{\text{prune}} \approx 0$: the system sits in the **zero-dissipation limit**.

This is a trivial result, not a proof of "noise-immune locking" — for a simple reason: if the pruning probability is already zero, then there is no event of "noise being resisted" to observe at all. In the zero-dissipation limit $\Phi \equiv 1$ holds **by construction**, regardless of whether the initial value is an Oseen vortex; with any other initial value, even a purely random one, the same $\Phi \equiv 1$ is obtained as long as $p_{\text{prune}} \approx 0$.

To actually test noise immunity, the experiment should be designed thus: fix the initial value, sweep over Λ, observe the decay curve of $\Phi(\Lambda)$, and compare the Oseen-vortex initial value against a chirality-free random initial value. Per `main_EN.md` §6.2, Φ (here, the structure density) decays smoothly from 0.49 as Λ grows, and it is the **shape** of that decay curve which carries the real information. The original experiment sampled a single point of the curve.

> Conclusion: this observation cannot support a "closed-loop proof of the Oseen vortex pseudo-spectral bound conjecture". At most it shows that at this particular working point, $\Lambda = 0.05$, pruning has not yet been activated.

---

3. Observation II: the correct reading of the residual 8.93×10^{-1}

Original observation: the MDS reconstruction residual is 0.893.

Corrected reading: whether classical MDS or iterative SMACOF is used, this residual is an **embedding error** — it measures "the degree to which the pairwise dissimilarity matrix cannot be represented in three-dimensional Euclidean space".

No quantitative relation between it and "intrinsic non-Euclidean geometric curvature" has been established. For such an interpretation to hold, one would need to:

1. first estimate the **intrinsic dimension** of the data (e.g. by maximum-likelihood estimation or the two-point correlation dimension), confirming that the true dimension exceeds 3;
2. then measure the "curl" independently with embedding-independent topological

quantities (persistent-homology Betti numbers, discrete curvature);

3. finally establish the correspondence between the two.

Until then, 0.893 should be reported only as "the three-dimensional embedding cannot fully represent this pairwise metric", and the three-dimensional embedding itself is a visualisation choice, not a physical dimension. Part of this value may also stem from non-convergence of the MDS iteration (see the methodological comparison in `main_EN.md` §6.1).

---

4. Where This Experiment Actually Sits in the Chain of Evidence

After the corrections, what this experiment can and cannot support is as follows.

Can support:

- an Oseen-vortex-type initial value can be injected into the discrete operator chain and retains its chiral character in subsequent evolution;
- at the working point $\Lambda = 0.05$ the system maintains a chiral channel density of about 0.44, and the chiral field exhibits power-law scaling (slope about −1.83), consistent with `main_EN.md` §6.6.

Cannot support:

- any form of proof of the pseudo-spectral bound conjecture (this requires a parameter sweep, not a single point);
- topological noise-immune locking (this requires a controlled comparison: chiral initial value vs chirality-free initial value);
- intrinsic non-Euclidean curvature (this requires intrinsic dimension and independent topological quantities);
- emergence of a rigid central axis (sensitive to the metric-reconstruction method, see `main_EN.md` §6.5).

---

5. Recommended Additional Experiments

To bring this report up to publishable empirical strength, the following should be carried out in order:

1. Λ **sweep**: with the same Oseen-vortex initial value, sweep $\Lambda \in [10^{-4}, 10^{2}]$ (17 logarithmically spaced points, ≥ 5 seeds) and report the structure density and spectral slope curves. `sre_core.py` can be reused directly.
2. **Initial-value control**: Oseen-vortex initial value vs chirality-free Gaussian initial value vs uniform random initial value; compare, at the same Λ, the evolution of structure density and of the topological charge $\mathcal{Q} = \mathrm{Tr}(A\mathbf{D})$.
3. **Intrinsic dimension estimation**: estimate the manifold intrinsic dimension with the two-point correlation dimension or MLE, decide whether the three-dimensional embedding is genuinely under-representing, and only then discuss "curvature".
4. **Embedding-independence test**: compute β_0, β_1 with persistent homology to test whether the "core–shell" structure exists independently of the MDS criterion.

---

6. Appendix: Relation to the Main Paper

This document no longer serves as the carrier of an independent empirical conclusion. It is repositioned as **an initial-value specification and supplementary experimental record for `main_EN.md` v2**. All quantitative assertions are subject to the reproduction results of `sre_core.py`. The "paradigm comparison" table of the original Section 1 has been merged into `main_EN.md` §1 and is not repeated here.

Emergence of Classical Mechanics from Discrete Causal-Information Networks: Ontological Mapping and Effective-Theory Limits within Status-Relational-Entropy (SRE) Dynamics

Version: 1.0

> Note: This is a conceptual-framework paper focusing on establishing ontological correspondences rather than term-by-term rigorous analytical derivation of full textbook mechanics formulae. Three representative physical examples - inertia, force-mass-acceleration relation, and Hooke's law - illustrate emergent behaviour under coarse-graining, decoherence and topological-phase constraints. Supporting topological-evolution derivations are provided in the appendix. **Appendix derivations are illustrative topological sketches and shall not be treated as rigorous complete mathematical proofs; they serve only for physical intuition**.

Abstract

Classical mechanics is widely understood as a macroscopic effective limit of underlying microscopic physical descriptions. Within the Status-Relational-Entropy (SRE) dynamical framework, spacetime, mass and force are not primitive background entities but statistical emergent outcomes governed by dissipation-compensation duality within a discrete bidirectional causal-information network.

This paper establishes ontological mappings between core SRE network-level observables and central quantities of Newtonian classical mechanics. Three necessary conditions for the emergence of classical-like behaviour are identified: large-sample coarse-graining, sufficient environmental decoherence, and operating far from topological-phase-transition thresholds (the BBP spectral-rank phase-transition and causal-loop coherence saturation $\rho \to 1$). Three representative examples - inertia, the force-mass-acceleration relation, and Hooke's elastic law - demonstrate statistical emergence of classical-style behaviour when all conditions are satisfied, while also marking breakdown boundaries for each phenomenon.

Conservation of momentum and conservation of energy can be traced to global closed-loop symmetries of the underlying causal graph. This manuscript completes one conceptual segment of the SRE unified theoretical hierarchy: axiomatic causal-information network → composite-particle / gluon-sea mass generation → topological electrodynamics → emergent classical mechanics → cosmological BBP-RMT simulation constrained by spectroscopic observational datasets.

This work emphasises ontological correspondence instead of exhaustive analytical reproduction of every textbook formula in analytical mechanics. Illustrative topological-evolution derivations are supplied in the appendix for pedagogical intuition and are not regarded as rigorous core proofs of the paper.

Keywords: Status-Relational Entropy; causal-information network; emergent classical mechanics; coarse-graining; decoherence; effective theory; ontological mapping; Hooke's law; inertia

1 Introduction

Within conventional physics, Newtonian classical mechanics constitutes a highly successful macroscopic effective theory, while quantum mechanics describes microscopic degrees of freedom. A substantial body of existing research studies the emergence of classicality via decoherence and coarse-graining, explaining how familiar macroscopic laws arise from microscopic rules.

Status-Relational-Entropy (SRE) dynamics adopts a more fundamental starting point: there exists no pre-given continuous spacetime manifold as a primitive background. Space, time, mass and force are not ontologically fundamental material substances; they are all high-level render-layer emergent phenomena of discrete bidirectional causal-information networks subject to dissipation-compensation duality.

Prior SRE work has accomplished axiom construction, topological origin of composite particles together with gluon-sea-like mass amplification, graph-cohomology-based reconstruction of electrodynamics, and cosmological random-matrix simulation using SDSS/eBOSS spectroscopic datasets. Nevertheless, the conceptual bridge connecting discrete causal-network ontology to classical-mechanics phenomenology has not yet been systematically elaborated.

The present paper fills this conceptual gap. **This manuscript does not aim to derive every textbook formula of analytical mechanics term-by-term**. The main objectives are as follows:

1. State a set of necessary physical conditions under which classical-mechanics-like behaviour emerges from SRE causal-

information networks;

2. Build an ontological-mapping table between network-level observables and physical quantities of classical mechanics;
3. Deploy three representative examples (inertia, $F \propto ma$, Hooke's law $F = -kx$) to demonstrate emergent behaviour together with corresponding breakdown thresholds;
4. Trace momentum-energy conservation to global closed-loop symmetries of causal graphs;
5. Delimit physical regimes where classical descriptions break down and one must fall back to the underlying discrete-network description.

Illustrative pedagogical topological-evolution derivations are placed in Appendix A and are not treated as rigorous core proofs.

2 Brief Review of Core SRE Ontological Concepts

Only concepts essential for this manuscript are recapitulated; the full axiomatic system is given in the SRE-v1.6 Axiom Suite.

1. No primitive continuous spacetime background exists.
2. **Distance**: emergent macroscopic quantity representing topological-compensation overhead between causal nodes, originating from dissipation-compensation duality.
3. **Global evolution step ΔS**: discrete fundamental state-refresh cycle of the causal-information network; macroscopic time emerges as accumulated counting of these evolution steps.
4. **Mass**: originates from local closed causal-loops (strongly-connected subgraphs). Mass magnitude corresponds to total topological-path complexity of internal feedback loops within the local closed subgraph. Altering the configuration of such a subgraph demands network-compensation overhead, manifesting macroscopically as inertial impedance.
5. **Force**: not a primitive fundamental entity; it emerges statistically as a macroscopic topological-compensation gradient averaged over numerous microscopic causal-coupling perturbations.
6. Two characteristic classes of topological phase-transitions: the BBP spectral-rank phase-transition (cosmological dimensional crossover between 2D-holographic phase and 4D-unlocked-spacetime phase); and the causal-loop coherence-saturation transition $\rho \to 1$ during composite-particle formation.

3 Three Necessary Conditions for Emergence of Classical Mechanics

For macroscopic system-level behaviour to approximate Newtonian classical mechanics, **all three of the following conditions must hold simultaneously**. Violation of any single condition renders classical-level descriptions inappropriate and forces recourse to the underlying causal-network description.

1. **Large-sample coarse-graining condition**
   The system contains a very large number of causal nodes and closed-loop subsystems. Random microscopic fluctuations of individual causal links are statistically averaged-out and suppressed at the macroscopic observable level. If the system possesses only few degrees-of-freedom, microscopic stochasticity cannot be averaged away and classical-like regular behaviour fails.
2. **Effective decoherence condition**
   The system couples sufficiently strongly to the environmental causal-information network. Long-range quantum-phase coherence and large-scale superposition configurations are destroyed by environmental interactions. Persistent large-scale coherent entanglement breaks classical phenomenology.
3. **Far-from-topological-transition-threshold condition**
   The system stays away from two classes of topological-transition points:

- Cosmological BBP spectral-rank phase-transition (2D-4D dimensional crossover);
- Causal-loop coherence-saturation $\rho \to 1$ associated with composite-particle formation.

Within transition regimes, compensation-operator topology undergoes abrupt structural reconstruction, and classical gradient-style force descriptions are no longer applicable.

> When conditions 1 & 2 & 3 are jointly satisfied, statistical averaging over discrete causal-network dynamics yields behaviour corresponding to classical-mechanics phenomenology.

4 Ontological Mapping: SRE-Network Observables ↔ Classical-Mechanics Quantities

> This table provides ontological correspondences, not strict point-wise numerical equality.

Quantity in classical mechanics	Ontological interpretation within SRE causal-information network
Inertial mass m	Total topological-feedback-path complexity inside a local closed causal-loop subgraph; network-compensation overhead required to alter its internal configuration, manifesting as inertial impedance.
Momentum $\boldsymbol{p}$	Directed information-propagation flux of causal-loop clusters; global transport flow produced by synchronised evolution of large sets of network links.
Force $\boldsymbol{F}$	Macroscopic topological-compensation gradient statistically averaged from numerous microscopic causal-coupling perturbations; an emergent gradient effect, not a primitive entity.
Spatial coordinate $\boldsymbol{x}$	Coarse-grained average topological-geodesic causal-step count between different subgraphs; approximates continuous coordinate after coarse-graining.
Time t	Accumulated count of global discrete evolution steps ΔS; approximates uniform continuous time within stable topological phases.
Kinetic energy E_{k}	Topological-flow overhead associated with directed propagation of causal-loop clusters.
Potential energy E_{p}	Latent compensation overhead stored in causal-coupling configurations among subgraphs.

4.1 Ontological Origin of Conservation Laws

- **Momentum conservation**: arises from global closed-loop symmetry of causal-network link interactions. Directed information-propagation flux cannot be locally created or annihilated out-of-nothing; flux can only be redistributed among subsystems. This yields macroscopic momentum conservation after coarse-graining.
- **Energy conservation**: originates from conservation of total topological-compensation overhead of the whole causal-information network. Transformations occur between internally-stored closed-loop configurations and open propagating-flow forms; total overhead magnitude is preserved.

5 Three Representative Physical Examples

> Preamble: All examples below demonstrate behavioural correspondences under the triple emergence conditions. **They are not rigorous analytical mathematical proofs derived directly from axioms**. If any emergence condition breaks down, the described classical-style behaviour ceases to hold. Topological-evolution formulae are given in Appendix A.

5.1 Example 1: Inertia

Classical phenomenon: A body maintains its existing state-of-motion in absence of external influences; larger inertial mass makes change-of-motion-state more difficult.

SRE ontological picture:
Inertia arises from total topological-feedback-path complexity inside local closed causal-loop subgraphs. A causal-loop cluster maintains its established internal link-synchronisation rhythm corresponding to macroscopic motion-state.

1. Without external perturbation: causal links preserve their existing synchronised configuration; no extra topological-compensation overhead is required. Macro-phenomenologically the object persists in its original motion-state - inertia.
2. To change motion-state: external coupling-perturbations must be injected to force re-synchronisation and rearrangement of large numbers of internal causal links, demanding substantial network-compensation overhead. Closed-loop subgraphs with richer internal feedback-paths (larger inertial mass) demand higher overhead for configuration-rearrangement; macroscopically they resist changes-of-motion more strongly.

Breakdown regimes:
If strong quantum-coherence dominates or the system enters the BBP spectral-rank topological transition, closed-loop topology itself mutates and classical inertial behaviour no longer applies.

5.2 Example 2: Force-mass-acceleration correspondence ($F \propto ma$)

Classical phenomenology: Larger applied force produces larger acceleration; for fixed force, larger inertial mass yields smaller acceleration, expressed as $a = F/m$.

SRE ontological picture:
Externally injected causal-coupling-perturbations produce macroscopic topological-compensation-gradients corresponding to the classical concept of force F.

1. Stronger external coupling-perturbations drive re-synchronisation of closed-loop cluster configurations within each global evolution step ΔS, giving larger effective macroscopic acceleration.
2. Inertial mass quantifies internal feedback-path complexity of the cluster. Higher complexity raises total compensation-overhead required for configuration-change. Under equal external perturbation strength, the rate-of-configuration-change is suppressed and effective acceleration becomes smaller.

Under full triple emergence conditions, statistically-averaged network-level behaviour yields the trend:

> external-perturbation-strength $\propto$ inertial-mass × rate-of-configuration-change
> which macroscopically corresponds to $F \propto ma$.

> Important remark: This example illustrates behavioural matching rather than strict axiomatic proof. Breakdown occurs if coarse-graining, decoherence or far-from-transition conditions are violated.

5.3 Example 3: Hooke's elastic law $F = -kx$

Classical phenomenology: Within small-deformation elastic regime, restoring force is linearly proportional and opposite to displacement $F = -kx$. Beyond the elastic-limit linearity vanishes; plastic-deformation or fracture occurs.

SRE ontological picture:
An elastic solid consists of large numbers of mutually-coupled closed causal-loop clusters held in equilibrium by mediating causal links that establish balanced topological-geodesic configurations corresponding to macroscopic equilibrium-position.

1. **Deformation** x: External perturbation shifts local subgraph-clusters away from equilibrium topological-geodesic positions; average inter-subgraph causal-step separations are altered, mapping to macroscopic displacement.
2. **Origin of restoring force**: After displacement, mediating causal links generate topological-compensation-gradients. The network tends to pull subgraph-clusters back toward the original equilibrium configuration, producing restoring effects. For small offsets the compensation-gradient is approximately linear with displacement, yielding emergent Hooke-law behaviour $F = -kx$.
3. **Ontological interpretation of elastic limit**: When displacement grows too large, many mediating causal links break and re-wire; underlying network-topology reconstructs itself. The linear compensation-gradient relation is destroyed; macroscopically this corresponds to plastic-deformation or material fracture.

Breakdown regimes:
Large deformation causing link-rewiring; strong quantum coherence; entry into BBP topological-transition regime all invalidate classical elastic description.

6 Discussion

Within the SRE worldview, classical mechanics is not a fundamental set of axiomatic laws of nature. Instead it constitutes an effective-theory description that emerges only when the discrete causal-information-network satisfies three joint physical prerequisites: large-sample coarse-graining, sufficient environmental decoherence, and staying sufficiently far from major topological-phase-transition thresholds.

This paper does not claim to analytically derive every formula of Lagrangian-Hamiltonian analytical-mechanics term-by-term from SRE axioms. The contribution is ontological: establishing what each classical quantity corresponds to at causal-network level, stating emergence-conditions, demonstrating representative examples, and marking out explicit failure-regimes.

The conceptual hierarchy of the full SRE programme is conceptually closed:

> Discrete causal-information-network ontology
> → Composite-particle topology and gluon-sea-like mass-amplification
> → Graph-cohomology-based emergent electrodynamics
> → (coarse-graining + decoherence + far-from-transition) → emergent classical mechanics
> → Cosmological BBP-RMT spectral-rank-phase-transition simulation constrained by observational spectroscopic catalogues.

Important conceptual distinction: ontological mapping and behavioural-example matching is different from strict complete mathematical derivation. Future work can pursue more quantitative coarse-graining-operator formalisation to tighten the mathematical bridge between network-micro-dynamics and classical-level phenomenology.

7 Conclusions

1. Newtonian classical mechanics emerges from SRE discrete causal-information-network as an effective-theory description subject to three joint necessary conditions: large-sample coarse-graining, sufficient environmental decoherence, and operation far-removed from topological-phase-transition thresholds. Breaking any of these conditions invalidates classical-level description.
2. Core classical quantities (mass, momentum, force, space, time, kinetic / potential energy) can be given clear ontological interpretations at causal-network graph-level. Momentum- and energy-conservation laws trace back to global causal-graph closure symmetries.
3. Three representative examples - inertia, force-mass-acceleration trend, Hooke's law - illustrate how classical-style behaviours appear under the required emergence-conditions, and each example also defines its own breakdown threshold.
4. This work completes a conceptual link inside the SRE unified-framework; full term-by-term analytical derivation of all analytical-mechanics formulae is not performed here and remains a direction for follow-up research.

Appendix A Topological-Evolution Derivations (Illustrative Sketch, Not Rigorous Complete Proof)

> Disclaimer: All topological-evolution formulae within this appendix serve only for physical-intuition pedagogy. **They shall not be treated as rigorous core proofs of this paper**. Complete strict formalisation is left for future work.

Notation Convention

1. ΔS: global discrete evolution step of the network;
2. $\mathcal{M}_{\text{topo}}$: local closed causal-loop subgraph, **topological-complexity (SRE proxy for inertial mass)**, counting total internal feedback-paths within the subgraph;
3. $\mathcal{G}_{\text{drive}}$: averaged external causal-coupling-perturbation strength (topological proxy for force);
4. $\mathcal{R}$: rate of internal-configuration rearrangement measured per global evolution step;
5. $\mathcal{D}_{\text{geo}}$: average topological-geodesic step-count between subgraphs (topological proxy for displacement);
6. $\mathcal{K}_{\text{topo}}$: topological-stiffness coefficient for the set of mediating causal links;
7. $\langle \cdot \rangle$: large-sample coarse-graining statistical-average operator;
8. All computations inside this appendix assume the three emergence-conditions hold: large-sample coarse-graining, decoherence, far from topological phase-transitions.

A1 Topological formulation of inertia

Closed-loop subgraph maintains its original configuration with zero external perturbation $\mathcal{G}_{\text{drive}} = 0$:

$$\langle \frac{\partial \mathcal{M}_{\text{topo}}}{\partial \Delta S} \rangle = 0$$

Under external driving, topological-compensation overhead consumed by configuration-rearrangement scales with topological complexity:

$$\mathcal{C}ost_{\text{comp}} \propto \mathcal{M}_{\text{topo}} \cdot \mathcal{R}$$

Larger $\mathcal{M}_{\text{topo}}$ demands higher topological-compensation overhead for configuration change, corresponding to larger inertia.

A2 Topological-evolution relation for force-mass-acceleration ($F \propto ma$)

Under statistical averaging, external causal-coupling perturbation yields a topological driving term:

$$\langle \mathcal{G}_{\mathrm{drive}} \rangle \propto \mathcal{M}_{\mathrm{topo}} \cdot \mathcal{R}$$

$\mathcal{R}$ denotes configuration-rearrangement rate per evolution-step, the topological proxy for acceleration.
Mapping topological proxies onto classical render-layer observables:

$$F \longleftrightarrow \langle \mathcal{G}_{\mathrm{drive}} \rangle, \quad m \longleftrightarrow \mathcal{M}_{\mathrm{topo}}, \quad a \longleftrightarrow \mathcal{R}$$

This yields the emergent trend relation:

$$\boldsymbol{F} \propto m\boldsymbol{a}$$

> Remark: This expresses proportional-trend correspondence, not strict equality with known constant of proportionality; proportional constants originate from global network topological rigidity and require global spectral inputs.

A3 Topological-evolution derivation for Hooke's law

$\mathcal{D}_{\mathrm{geo},0}$: topological-geodesic step-count between subgraphs at equilibrium;
$\Delta\mathcal{D}_{\mathrm{geo}}$: topological-geodesic offset (topological proxy for displacement):

$$\Delta\mathcal{D}_{\mathrm{geo}} = \mathcal{D}_{\mathrm{geo}} - \mathcal{D}_{\mathrm{geo},0}$$

Topological restoring-compensation gradient generated by mediating causal links:

$$\langle \mathcal{G}_{\mathrm{restore}} \rangle = -\mathcal{K}_{\mathrm{topo}} \cdot \Delta\mathcal{D}_{\mathrm{geo}}$$

Mapping onto classical observables:

$$F \longleftrightarrow \langle \mathcal{G}_{\mathrm{restore}} \rangle, \quad k \longleftrightarrow \mathcal{K}_{\mathrm{topo}}, \quad x \longleftrightarrow \Delta\mathcal{D}_{\mathrm{geo}}$$

For small offsets without link breaking-rewiring, we obtain the emergent relation:

$$\boldsymbol{F} = -k\boldsymbol{x}$$

> Breakdown topological condition: When $|\Delta\mathcal{D}_{\mathrm{geo}}|$ exceeds the tolerance threshold of mediating links, mediating causal links break and re-wire; $\mathcal{K}_{\mathrm{topo}}$ is no longer constant and the Hooke-law topological relation collapses.

A4 Topological form of conservation laws

Total topological-compensation overhead $\mathcal{U}_{\mathrm{total}}$ of the global causal-information network is conserved:

$$\frac{\mathrm{d}\langle \mathcal{U}_{\mathrm{total}} \rangle}{\mathrm{d}\Delta S} = 0$$

Information-propagation flux (topological proxy for momentum) satisfies global closed-loop constraints: only redistribution among subsystems occurs, total flux remains unchanged.

References

1. SRE Dynamics Axiom Suite v1.6, Zenodo archive.
2. Zurek, W.H. Decoherence and the transition from quantum to classical.
3. Hoel, E.P. Causal emergence and coarse-graining in complex networks.
4. Ehrenfest theorem: quantum-classical correspondence principle.
5. SRE-v6.2-rev: cosmological BBP-RMT spectral-rank-phase-transition simulation with SDSS/eBOSS spectroscopic datasets.

A Complete Characterization of the Electron within the SRE Framework: A Unified Account of Axiomatization, the Fine-Structure Constant, Molecular Computation, and Orbital Assignment

Version: 1.0 (merged manuscript)

Project references (DOI; in-text citations appear in Appendix E):

- https://doi.org/10.5281/zenodo.22765850 — Topological derivation of the fine-structure constant within the SRE framework
- https://doi.org/10.5281/zenodo.22162514 — SRE Dynamics: The Book of the Void
- https://doi.org/10.5281/zenodo.19935370 — SRE Dynamics User Guide
- https://doi.org/10.5281/zenodo.20482974 — Conjecture on the Möbius topology of light
- https://doi.org/10.5281/zenodo.21025053 — Technical report: intrinsic algebraic topology of light and the SRE axion matrix
- https://doi.org/10.5281/zenodo.22119957 — SRE dynamics: strict reconstruction of Maxwell's field equations from purely dimensionless graph cohomology and global evolution steps
- https://doi.org/10.5281/zenodo.20576606 — Dynamics of hierarchical dissipative self-organizing binary networks
- https://doi.org/10.5281/zenodo.20837960 — Emergence of multi-dimensional spacetime and gravity
- https://doi.org/10.5281/zenodo.22119635 — SRE electrical quantities defined (charge/current/resistance/voltage/power/$E=mc^2$)

> **Core conclusion.** Within the SRE (Status–Relational Entropy) theoretical framework, the electron is not a point particle pre-placed in space; it is the **unique, stable, noise-robust emergent fixed point** obtained by applying the sparse operator `R` over the coherent core of a **binary self-organizing network**; its basal structure is the **three-dimensional cube Q_3** (vertex count 8, edge count 12, first Betti number 5, i.e., 12 × 5 = 60).
>
> This conclusion is argued successively, starting from the Fork A axioms, through the stages P0 (uniqueness), R2 (convergence), and R3 (robustness), and is numerically reproduced by the unified validation program with a **10/10** pass rate. From the electron basal structure one may transfer the molecular-computation descriptors (MCI spectral fingerprint, Q_3 basal recognition, MDS relational-inversion geometry), and from the 16-cell state space one may derive the laws of electron orbital assignment (Pauli cell-uniqueness, the capacity law $2n^2$, Hund's rule as tension minimization).
>
> **Contextual limitation (read first).** The objects "electron", "Q_3", "60 nodes", and "coherent core" above, as well as the concepts treated later — logical depth, Z_2 winding phase, Möbius double cover, renormalization flow, and cosmological phase transition — are **constructions, analogies, and counts internal to the SRE heuristic-simulation model**; they are not physical quantities directly measurable in the real world. All proofs and numerical validations in this manuscript hold only within the SRE model; they represent internal self-consistency of the model and **do not automatically equate to facts of real physics**. Real physical and chemical phenomena must be examined by external means (DFT, molecular dynamics, experimental observation) to test the model's inferences.

Abstract

This manuscript provides a unified and complete characterization of the electron within the framework of Status–Relational Entropy (SRE) dynamics: the electron is the unique, stable, noise-robust emergent fixed point of the sparse operator `R` over the coherent core of a binary self-organizing network, whose basal structure is the three-dimensional cube Q_3 (vertex count 8, edge count 12, first Betti number 5, i.e., 12 × 5 = 60). The manuscript unfolds along a reproducible logical chain: (1) three independent topological derivation paths for the fine-structure constant $\alpha \approx 1/137.036$ are given (the Laplacian spectral-gap of the Möbius ring graph, the four-state eigen-spectrum frequency formula, and the Möbius-parameterized arc-length perturbation), and the Tier 0 independent computational review confirms that λ_0 is not an independent primitive and that the n = 60 hit is a discretization coincidence; (2) through the Fork A axiomatization (A1–A4 + R1), the double-cover four-state complex (8, 12, 5, 60) is derived independently from the four axioms, an observable dictionary $\alpha = \lambda_2/\lambda_{max} = v/c$ *is established, and the residual is promoted to the A5 bare coupling* $\delta = 4.347\times10^{-5}$*; (3) the sparse operator* `R` *is shown to be unique, convergent, and robust over the coherent core of the binary network*

(P0/R2/R3, with a 10/10 validation suite); (4) the account is transferred to molecular computation, providing the MCI spectral fingerprint, Q_3 basal recognition, and MDS relational-inversion geometry; (5) the laws of electron orbital assignment are derived from the 16-cell state space (Pauli as cell-uniqueness, capacity laws $C\ell = 4\ell+2$ and $C_n = 2n^2$, Hund's rule as tension minimization, and the numerical coincidence of 60 with the capacity of the first four shells). This work does not prove that the SRE axioms hold objectively; all conclusions are internally self-consistent constructions within the SRE model, and real physical and chemical inferences must be independently verified by external means.

Keywords: SRE dynamics; electron basal structure; fine-structure constant; binary self-organizing network; Fork A axioms; three-dimensional cube Q_3; orbital assignment; Pauli exclusion principle; Hund's rule; molecular computation

---

0. Document structure and reading guide

This manuscript is organized in the following order, fusing four existing component documents (the fine-structure-constant derivation, the complete characterization of the electron, the Fork A axiomatic derivation, and the electron orbital-assignment derivation) into a unified account:

- Section 1 presents the **terminological conventions**. Unfamiliar terms that first appear in the text are marked with 【】; their precise definitions are collected in this section, given in plain written form for readers without a physics or mathematics background. Each key term is accompanied by a note indicating its status as an "internal construct/analogy of the model", so as to avoid confusion with real physical quantities.
- Section 2 gives the **theoretical foundations of SRE**: the three axioms, the Operator chain, and the Dirichlet energy lower bound.
- Section 3 establishes the **Möbius topological modeling of light**: the residual manifold, the 4π closure period, and the validation confidence.
- Sections 4–5 give the **three independent derivation paths for the fine-structure constant α** and their convergence analysis, including the Tier 0 independent computational review.
- Sections 6–8 give the **Fork A axiomatization**: the double-cover four-state complex (8, 12, 5, 60), the observable dictionary $\alpha = \lambda_2/\lambda_max = v/c$, and the promotion of the residual to the A5 bare coupling δ.
- Section 9 argues that **the electron is an emergent object of the binary self-organizing network**: the basal structure Q_3, the "deep yet compact" property, spatial emergence, and the three-layer RG structure.
- Section 10 gives the **proof chain** (P0 uniqueness, R2 convergence, R3 robustness); Section 11 gives the **unified validation program** (10/10).
- Section 12 is the **molecular-computation application**; Section 13 is the derivation of the **laws of electron orbital assignment**.
- Section 14 delimits the **honest boundaries**; Section 15 gives the **conclusions**.
- Appendices A–E give the key numerical values, the merged glossary, the code listing, the figure listing, and the references/DOIs, respectively.

Throughout, this manuscript follows a single contextual-distinction principle: **internal constructs of the model / internal self-consistency proofs of the model / validation passed ≠ facts of real physics**. The three belong to different levels, and the reader should not conflate them.

---

1. Terminological conventions

Each key term below is marked with a 【Remark】 indicating its model-internal status within the SRE framework. Unfamiliar terms that first appear in the text are marked with 【】; conventions for symbols are collected in §1.2.

1.1 Glossary

Term	Definition (with remark on model-internal status)
Binary self-organizing network	A square matrix $M \in \{+1,-1\}^{n\times n}$ whose entries are the elements $+1$ and -1: $+1$ indicates that the relation between two nodes is co-oriented (connection) and -1 indicates counter-orientation (repulsion). Starting from a unit seed, the network grows step by step according to a local dormancy–activation rule. SRE theory regards it as the primitive language of the underlying structure of the world. 【Remark: internal construct of the model — the simulation mechanism by which SRE describes the underlying world, not an observable independent of the model】
Coherent core	When the network has evolved to scale N, the rigid kernel within it whose internal order is stable and no longer degrades; the electron resides within this kernel. 【Remark: model-internal concept, corresponding to the rigid kernel of network evolution in simulations, not a real detectable structure】
Z_2 holonomy (Z_2 winding phase)	The sign factor accumulated by a closed path traversing one full circuit. On a binary network, for any closed loop formed by four nodes, the product of the four edge-relation values is strictly $+1$ or -1; the value -1 indicates that the traversal experienced one spatial flip. This is the algebraic essence of electron spin-1/2 and of 4π (restoration after two turns). 【Remark: model-internal algebraic-structure analogy used to interpret the sign factor of spin-1/2; not a direct measurement of real spin】
Möbius double cover (4π)	The electron must rotate two full turns (720°) to be restored, a property analogous to the Möbius band; in signed-network language, this is a structure that "closes only after two steps". 【Remark: model-internal geometric analogy (analogous to the Möbius band), not a real spacetime structure】
DCF (discrete closed-chain complex / double-cover four-state complex)	A triple $(\|V\|, \|E\|, \beta_1)$ describing a complex of discrete closed loops, denoting vertex count, edge count, and first Betti number, respectively; β_1 characterizes the number of independent closed loops. The basal structure of the electron takes the value $(8, 12, 5)$. 【Remark: model-internal counting tool; (8, 12, 5) describes the topological type of a simulated object, not a measured geometric result of the real electron】
Spectral-gap ratio $\alpha = \lambda_2/\lambda_max$	The ratio of the second-smallest eigenvalue (the second-lowest collective vibration mode) to the largest eigenvalue of the relation-matrix spectrum, used to measure the relative stiffness of the softest collective mode of the system. The α of the electron equals precisely the fine-structure constant **1/137.036**. 【Remark: model-internal spectral feature; its agreement with 1/137.036 is an internal reproduction of the model and does not mean that α has been "measured" by this model; the real α remains determined by experiment】
MDS inversion (multi-dimensional scaling)	Starting from the pairwise distance matrix of objects, reconstruct their coordinates in a space of given dimension. In SRE, the spatial shape of the electron emerges from the relation matrix through this route, rather than being pre-given. 【Remark: model-internal geometric construction used to reconstruct coordinates from the relation matrix, which presupposes no real space; the coordinates obtained are a model-internal relational embedding, not a measurement of real positions】
Renormalization / coarse-graining (RG)	Merging a cluster of nodes into a single super-node, thereby examining the system at a coarser scale; SRE uses it to roll the network upward layer by layer, establishing the micro–meso–macro correspondence. 【Remark: model-internal multi-scale up-rolling operation; the micro–meso–macro correspondence is a model picture, not a settled claim about real levels】
Fixed point	An object that remains unchanged under repeated application of some operator. The electron is a fixed point of the sparse operator R. 【Remark: a purely mathematical concept; here it describes the iterative behavior of an operator】

Term	Definition (with remark on model-internal status)
Fork A axioms	The axiom system for deriving the 60-node projection of the electron, composed of four axioms (A1–A4) and one rule (R1), proceeding purely from logic and topology without relying on empirical parameters. 【Remark: an internal axiom system of the model; its consequences hold only within the model】
Basal structure	The minimal invariant kernel that cannot be further compressed under arbitrary coarse-graining. The basal structure of the electron is the three-dimensional cube Q_3. 【Remark: model-internal "minimal invariant kernel"; Q_3 is the graph-theoretic type of a simulated object, not a real geometric body of the electron】
Logical depth	The number of nested relational levels required to constitute an object. For the electron it is approximately 10^{23} layers, hence "deep"; after collapse its basal structure is merely 12×5, hence "compact". 【Remark: model-internal count/analogy (10^{23}), not a measured property of the real electron】
Bare coupling δ (delta)	The small input constant required to make the spectral gap converge precisely to 1/137; its status is isomorphic to that of α as a measured input in QED. 【Remark: model-internal input constant; it only analogizes the status of α as a measured input in QED and does not constitute a substitute conclusion about QED】
Cell (state unit)	The smallest distinguishable unit (v, σ) of the basal state space of the electron: basal Q_3 vertex × spin sheet, 16 in total. 【Remark: model-internal count; capacity 1, i.e., Pauli cell-uniqueness (see Section 13)】
Faithful cover	The property that the relation rows of the individual units on the double-cover matrix are pairwise distinct. 【Remark: model-internal algebraic property; it is the algebraic basis of the Pauli exclusion principle】
Orbital (SRE)	A collective vibration mode on the atom (mesoscopic kernel) = an eigenvector of the relation Laplacian. 【Remark: model-internal analogy; its density pattern and the spatial orbital functions of quantum mechanics are not the same object】
Subshell capacity 4ℓ+2	The number of electrons that can be accommodated in the subshell of azimuthal quantum number $\ell = 2(2\ell+1)$. 【Remark: purely arithmetic combinatorics; the source of the degeneracy is a borrowed term (see Section 13)】
Shell capacity $2n^2$	The electron capacity of the principal shell $n = \Sigma_{\ell<n}(4\ell+2)$. 【Remark: purely arithmetic combinatorics】
Pairing tension	The number of orbitals in which both spin sheets of the same orbital are occupied (paired). 【Remark: a proxy quantity of the model-internal energy functional; the object minimized by Hund's rule】
Madelung rule	The empirical construction order in which orbitals are filled in ascending order of $(n + \ell)$. 【Remark: an open borrowed term, not derived by SRE】

1.2 Symbol conventions

Symbol	Meaning
α	Fine-structure constant 1/137.035999 ≈ 0.00729735257
λ_0	Vacuum basal spin residual (Tier 0 ruling: not an independent primitive)
κ	Chiral-locking strain coefficient (= 0.828)
β	Spectral scaling invariant (ALPHA_0_DYNAMIC = 21.09256)
γ	Chiral coupling coefficient (GAMMA_LATENCY = 0.0585)

Symbol	Meaning
$\lambda_1, \lambda_2, \lambda_3$	The first three nonzero eigenvalues of the causal-association tensor **B**
ρA, ρB	Causal flux injected at boundary nodes A/B (High = 10 / Low = 2)
L_G	Graph Laplacian matrix
$\lambda_2(L_G)$	Algebraic connectivity of the graph (Fiedler value)
$\rho(L_G)$	Spectral radius of the graph Laplacian (largest eigenvalue)
n	Electron projection scale; n = dim(E) × dim(F) = $\|E\| \times \beta_1 = 12 \times 5 = 60$
w	Möbius twist-channel weight; w* = 1 + δ
δ	A5 bare coupling (= 4.347×10^{-5})
R	Sparse operator (dormancy pruning ∘ holonomy closure)
Q_3	Three-dimensional cube, basal structure (8, 12, 5)
ε	R3 physical dissipation parameter (= 0.1832, median dormancy probability of the network)

2. Theoretical foundations of SRE

2.1 The three axioms

SRE dynamics is built upon three discrete-topology axioms:

Axiom I (binary constraint): the spin-polarity relations between atoms satisfy $s_{ij} \in \{+1, -1\}$, constituting a strictly binary signed network.

Axiom II (asynchronous activation): the logical-depth steps of boundary nodes are asynchronous: node A is at step t, node B at step t', with $t \neq t'$.

Axiom III (geodesic field): the residual topological manifold is strictly parameterized by the global circumferential wrapping phase ϕ and the micro-impedance bandwidth w.

2.2 The binary self-organizing network and the coherent core

The basic starting point of SRE is that all structure emerges from the binary relation matrix `M ∈ {+1,−1}`, in which no space, particle, or force is pre-given; only co-oriented/counter-oriented binary relations between nodes exist. The network grows autonomously according to the dormancy–activation rule (implementation in `code/sim_p.py`). This is analogous to an assembly of magnetic needles admitting only the two orientations "co-oriented/counter-oriented": under random perturbation, the system spontaneously forms an ordered kernel.【The above is a model-internal picture of the world, not an assertion about the real world】

When the network has evolved to scale `N`, the innermost `k = ⌊0.2 N⌋` nodes form a rigid, non-degenerate kernel; the strictness of the existence and stability of this kernel in the `N→∞` limit is guaranteed by Theorem 1 of the theoretical monograph (DOI: 10.5281/zenodo.22162514) (the citation of 14-2 appears in Appendix E). **This kernel is the logical basis of the electron (model-internal)**. The numerical simulation at `N=300` gives `k=60`, which coincides exactly with the electron projection scale `n=60`. This agreement should be positioned as a "co-directional indication" rather than a strict proof, and both quantities are model-internal numerical values (`N=300` is a simulation choice); its scope of applicability and limitations are detailed in Section 14.

2.3 The Operator chain

SRE theory defines a chain of operators forming the mapping from atomic coordinates to macroscopic physical quantities:

Operator	Function	Output
Operator 6	Construct the smooth adjacency matrix A_{smooth} and the graph Laplacian L_G	λ_2, α_n (spectral radius)

Operator	Function	Output
Operator 4	Construct the RBF adjacency matrix W_e	W_e, Dirichlet energy E_D
Operator 5	Discrete permeability c_e	Information propagation speed
Operator 11	Spin-charge adjacency A_q (electronegativity-weighted)	$\lambda_{2,q}$, κ_q, α_q
Operator 12	Topological closed-loop audit	n_{loops}, Möbius ratio

2.4 The Dirichlet energy lower bound

The core theorem of SRE (Operator 4):

$$E_D(E_s) \geq \lambda_2 \cdot \|E_s\|_2^2 > 0$$

When $\lambda_2 > 0$, the Dirichlet energy has a positive lower bound and the topology is stable. This is the cornerstone of all subsequent derivations.

---

3. Möbius topological modeling of light

3.1 Definition of the residual manifold

Axiom I (residuality): light is not an independent material entity. Given that boundary node A evolves to logical-depth step t and node B to step t', light manifests as the mutually non-annihilating topological residual at the causal intersection of the joint execution chain:

$$\Psi_{\text{light}}(\phi, w) \equiv \ker\left(\partial_{\text{mutual}}(A_t, B_{t'})\right)$$

3.2 Möbius parameterization

Axiom II (closed-form parametric mapping): the ideal 0-state residual manifold is strictly governed by two intrinsic degrees of freedom — the global circumferential wrapping phase $\phi \in [0, 2\pi)$ and the micro-impedance bandwidth $w \in [-w_{\max}, w_{\max}]$:

$$\mathbf{X}(\phi, w) = \begin{pmatrix} (1 + w\cos\frac{\phi}{2})\cos\phi \\ (1 + w\cos\frac{\phi}{2})\sin\phi \\ w\sin\frac{\phi}{2} \end{pmatrix}$$

Topological invariant: as $\phi \to \phi + 2\pi$, the transverse vector undergoes an intrinsic flip $w \to -w$, with no localized spatial displacement. This guarantees geometrically that the residual structure is a **non-orientable topological manifold possessing exactly one boundary loop**.

3.3 The 4π closure period

Theorem (4π closure): owing to the non-orientable single-boundary logic, a complete path must traverse $\Delta\phi = 4\pi$ to achieve boundary closure ($\mathbf{X} = \mathbf{X}_0$). This provides an explicit geometric origin of the half-integer spin structure.

3.4 Topological validation confidence

The algebraic-expansion engine for the Möbius residual manifold validates a 99.2094% topological-alignment fidelity, demonstrating that the residual manifold indeed possesses the topology of a Möbius band.

---

4. The fine-structure constant: three independent paths

This chapter gives three mutually independent topological derivation paths for the fine-structure constant α within the SRE framework. The original reproduction programs of the three paths are given in Appendix C (`code/_verify_alpha.py`, `code/_scan_alpha_graphs.py`, `code/_verify_alpha_mobius.py`, `code/_verify_alpha_light3.py`, `code/_verify_n60_lambda0.py`).

4.1 Path 1: Laplacian spectral gap of the Möbius ring graph

Discretization scheme: discretize the 4π-periodic Möbius band into n equally spaced sampling points, with angular spacing:

$$\Delta\phi = \frac{4\pi}{n}$$

Construct the **signed adjacency matrix**: the half-twist of the Möbius band makes the closure edge weight -1, and the remaining edges $+1$:

$$A_{ij} = \begin{cases} +1 & i \to j \text{ is an ordinary adjacency} \\ -1 & i \to j \text{ is a twist closing edge} \end{cases}$$

Möbius ring Laplacian spectrum: for an n-node Möbius ring, the eigenvalues of the signed adjacency matrix are:

$$\lambda_k(A_M) = 2\cos\frac{(2k+1)\pi}{n}, \quad k = 0, 1, \ldots, n-1$$

The eigenvalues of the graph Laplacian $L_M = 2I - A_M$ are:

$$\mu_k = 2 - 2\cos\frac{(2k+1)\pi}{n}$$

The smallest nonzero eigenvalue ($k = 0$) and the largest eigenvalue ($k = n-1$):

$$\mu_{\min} = 2 - 2\cos\frac{\pi}{n}, \quad \mu_{\max} = 2 + 2\cos\frac{\pi}{n}$$

Spectral-gap formula: the spectral gap is defined as the ratio of the smallest nonzero eigenvalue to the largest eigenvalue:

$$\text{gap} = \frac{\mu_{\min}}{\mu_{\max}} = \frac{1-\cos\frac{\pi}{n}}{1+\cos\frac{\pi}{n}} = \tan^2\frac{\pi}{n}$$

Solving for the target n: let the spectral gap equal α:

$$\tan^2\frac{\pi}{n} = \alpha = \frac{1}{137.036}, \qquad \tan\frac{\pi}{n} = 0.085425, \qquad \frac{\pi}{n} = 0.085218 \text{ rad}, \qquad n = 36.866$$

n must be an integer. $n = 37$ gives $\tan^2(\pi/37) = 0.007244$ (error 0.73%), and $n = 36$ gives 0.007654 (error 4.89%). **A pure 1D Möbius ring cannot hit α exactly.**

Lifting to the 2D Möbius band: generate the 3D point cloud from the Möbius parameterization of §3.2, and construct the graph Laplacian with k-nearest neighbors (k-NN). For $n = 60$ sampling points, $w = 0.1$, $k = 3$:

$$\text{gap}(L^{M\ddot{o}bius}_{n=60}) = \frac{\lambda_2(L_G)}{\rho(L_G)} = 0.0072974585$$

Compared with $\alpha = 0.0072973526$, the **error is 0.0015%**.

Stability validation:

Perturbation type	Result
$w \in [0.05, 0.15]$	gap unchanged to 14 digits (plateau region)
random sampling jitter (5 runs)	gap unchanged to 14 digits
$k = 2$	deviates 10.6%
$k = 4$	deviates 10.5%
scan over $n = 30 \sim 200$	**only** $n = 60$ **hits** α

Topological meaning of n=60:

$$n = 60 = \dim(E) \times \dim(F) = |E| \times \beta_1 = 12 \times 5$$

where the dimensions come from the **double-cover four-state complex (DCF complex)** used in the SRE Maxwell-equation validation: the D_edge of Maxwell.py defines a cell complex of an 8-ring plus 4 skip-2 chords ($|V| = 8$, $|E| = 12$, $\beta_1 = 5$). The angular spacing is $\frac{4\pi}{60} = \frac{\pi}{15}$, and $15 = 3 \times 5$, indicating that the discretization precision of the Möbius band is determined by the dimensional structure of the Maxwell-equation circuit diagram.【Naming correction: the source code labeled this graph as $K_{3,5}$, but the true complete bipartite graph $K_{3,5}$ has 15 edges and $\beta_1 = 8$; the correct name of this complex is the double-cover four-state complex (DCF), whose origin appears in Section 6.】

4.2 Path 2: the four-state eigen-spectrum frequency formula

Frequency formula (quoted from the Möbius topology theory of light, DOI: 10.5281/zenodo.21025053, whose §3.2 gives the topological definition of frequency):

$$\Delta\lambda = \sqrt{\mathrm{Tr}(M)^2 - 4\det(M)} = \alpha \cdot \frac{\lambda_1}{\lambda_2+\lambda_3}$$

where M is the 2×2 cross-spectral response matrix, and $\lambda_1, \lambda_2, \lambda_3$ are the first three nonzero eigenvalues of the causal-association tensor $\mathbf{B}$.

Cross-spectral matrix M: take $M = \mathrm{diag}(\lambda_1, \lambda_2)$; then $\mathrm{Tr}(M) = \lambda_1 + \lambda_2$, $\det(M) = \lambda_1\lambda_2$, which, upon substitution, gives $\Delta\lambda = |\lambda_1 - \lambda_2|$; the frequency formula thus becomes:

$$|\lambda_1 - \lambda_2| = \alpha \cdot \frac{\lambda_1}{\lambda_2+\lambda_3}$$

Four-state eigen-spectrum: the boundary nodes drive the global manifold through transitions among four discrete eigen-spectrum states by switching between High ($\rho = 10$) and Low ($\rho = 2$):

$$\begin{cases} \lambda_1 = \beta \cdot (\rho_A + \rho_B)^2 \\ \lambda_2 = \gamma \cdot |\rho_A - \rho_B| + \lambda_0 \\ \lambda_3 = \frac{\lambda_2}{4(1+\kappa)} \end{cases}$$

Modulation state	ρ_{total}	λ_1	$\lambda_2 - \lambda_3$
(H,H)	20	400β	$0 + \lambda_0'$
(H,L)	12	144β	$8\gamma + \lambda_0'$
(L,H)	12	144β	$-8\gamma + \lambda_0'$
(L,L)	4	16β	$0 + \lambda_0'$

Choosing the (L,L) symmetric state: in the lowest-energy symmetric state (L,L) ($\rho_A = \rho_B = 2$), the chiral term $\gamma \cdot |\rho_A - \rho_B| = 0$, so that $\lambda_2 = \lambda_0$ directly exposes the vacuum residual:

Parameter	**Formula**	**(L,L) value**
λ_1	$\beta \cdot 4^2$	$16\beta = 337.481$
λ_2	$\gamma \cdot 0 + \lambda_0$	$\lambda_0 = 0.006420$
λ_3	$\lambda_0/(4(1+\kappa))$	0.000878

Expanding $\lambda_2 + \lambda_3$:

$$\lambda_2 + \lambda_3 = \lambda_0 \left(1 + \frac{1}{4(1+\kappa)}\right) = \lambda_0 \cdot \frac{5+4\kappa}{4+4\kappa}$$

Substitution into the frequency formula: substitute the (L,L) values and Eq. (2) into Eq. (1) and solve for α:

$$\alpha = (16\beta - \lambda_0) \cdot \lambda_0 \cdot \frac{5+4\kappa}{16\beta \cdot (4+4\kappa)}$$

Exact solution and simplification: the magnitude analysis $\lambda_0/(16\beta) = 1.9 \times 10^{-5} \ll 1$ gives $16\beta - \lambda_0 \approx 16\beta$ (error < 0.002%), so Eq. (3) simplifies to:

$$\boxed{\alpha \approx \lambda_0 \cdot \frac{5+4\kappa}{4+4\kappa}}$$

Solving inversely: $\lambda_0 = \alpha \cdot \frac{4+4\kappa}{5+4\kappa}$ (Eq. 5).

Numerical validation: substituting $\kappa = 0.828$ (THETA_CONFORMAL):

$\frac{4+4\times0.828}{5+4\times0.828} = \frac{7.312}{8.312} = 0.87971, \qquad \lambda_0 = \frac{1}{137.036} \times 0.87971 = 0.0064195$

Validation item	Formula	Value	Error
Exact formula	Eq. (3)	0.00729735256928	0.0000%
Simplified formula	Eq. (4)	0.00729735256928	0.0000%

The two formulas give **fully identical results to 14 significant digits**. **Parameter independence:** in the symmetric state β cancels completely and γ does not enter α; α is determined solely by λ_0 and κ ($\kappa = 0.828$ determines the conversion ratio $\lambda_0 \to \alpha$ of 1.137; $\lambda_0 = 0.006420$ is the direct carrier of α) — consistent with the physical intuition that α is a vacuum property.

4.3 Path 3: Möbius-parameterized arc-length perturbation

Global geodesic period (quoted from the Möbius topology theory of light, DOI: 10.5281/zenodo.21025053, whose §3.3 defines the wavelength as the minimal intrinsic geodesic distance over which the manifold maintains global consistency under the spin-rotation transformation):

$\lambda_{\text{wave}} \equiv \oint_{\mathcal{M}} ds = \int_0^{4\pi} \left\| \frac{\partial \mathbf{X}}{\partial \phi} \right\| d\phi$

Parametric dependence of the arc length: computing $\|\partial \mathbf{X}/\partial \phi\|$, the centerline ($w = 0$) arc length is $\text{arc}(0) = 4\pi$.

Perturbation expansion: the arc length is an even function of w, so the first derivative is zero. The second-order Taylor expansion:

$\text{arc}(w) = 4\pi + \frac{1}{2} \frac{d^2\text{arc}}{dw^2}\Big|_{w=0} w^2 + O(w^4), \qquad \frac{d^2\text{arc}}{dw^2}\Big|_{w=0} = \pi$

that is, $\text{arc}(w) \approx 4\pi + \frac{\pi}{2} w^2 + O(w^4)$.

Arc-length correction ratio: define the normalized arc-length correction:

$\delta(w) = \frac{\text{arc}(w) - 4\pi}{4\pi} = \frac{w^2}{8} + O(w^4)$

Arc-length encoding of the fine-structure constant: setting $\delta = \alpha$ gives $w = \sqrt{8\alpha} = 0.241617$ (Eq. 7). Numerical validation:

Quantity	Value
$w = \sqrt{8\alpha}$	0.241617
$\text{arc}(w)$	12.66048
$\delta = (\text{arc} - 4\pi)/4\pi$	0.007489
α	0.007297
δ/α	1.026
residual $\delta - \alpha$	1.9×10^{-4} (from the w^4 higher-order term)

Higher-order correction: the coefficient of the w^4 term is determined numerically as $c \approx 0.055$ (stable convergence):

$\delta = \frac{w^2}{8} + 0.055 \cdot w^4 + O(w^6)$

At $w = 0.2416$, the w^4 term contributes 1.88×10^{-4}, explaining 2.6% of the residual.

---

5. Convergence analysis of the three paths (Tier 0 review)

5.1 Summary and independence

Path	Formula	α value	Error	Source of residual
Graph topology	$\text{gap}(L_{n=60}^{M\ddot{o}bius})$	0.00729746	0.0015%	graph discretization

Path	Formula	value	Error	Source of residual
Frequency formula	$\lambda_0 \cdot \frac{5+4\kappa}{4+4\kappa}$	0.00729735	0.0000%	none (exact solution)
Arc-length perturbation	$w^2/8$	0.00729735	2.6%	w^4 higher-order term

The three paths are mathematically independent (algebraic graph theory / linear algebra / differential geometry), and their convergence supports the internal self-consistency of α within the SRE framework. The value $n = 60 = 12 \times 5$ appears simultaneously in the Maxwell validation graph (DOI: 10.5281/zenodo.22119957) and the Möbius topology of light (DOI: 10.5281/zenodo.21025053), constituting a bridge between the circuit diagram and the topology of light.

5.2 Tier 0 theoretical review

An independent computational review of the three paths above (programs `code/_tier0_lambda0_independent.py`, `code/_tier0_refinement_limit.py`, `code/_tier0_theory_reaudit.py`) yields three corrective conclusions:

(1) λ₀ is not an independent primitive (PASS-A failure): under the constraint of forbidding α as an input, four pre-registered candidate spectral quantities (C1–C4) attempted independently to yield λ₀. The result is that C4 (3D kNN spectral gap) directly gives α (error 0.0015%), whereas the candidate quantities for λ₀ (C1, C3) carry large errors. Conclusion: λ₀ is **not an independent primitive**; the gap path has made λ₀/κ redundant intermediate quantities.

(2) Grid refinement limit FAIL: design the refinement sequence $n = 60 \times 2^m$ and test whether the C4 spectral gap converges to α as $n \to \infty$. The result is that the spectral gap $\propto n^{-1.99} \to 0$, **not converging to α**. This indicates that the n=60 hit is a discretization coincidence, not the projection of a continuum limit.

(3) $n_\alpha = 60.000436$: solving exactly for the crossing point $\mathrm{gap}(M_n) = \alpha$ gives $n_\alpha = 60.000436$ — a relative deviation of only 7.3×10^{-6} from the integer 60. It is the unique even integer within the 1% window, with an a-priori probability of about 1/99. The residual 1.452×10^{-5} is not rounding noise and requires a corrective mechanism for explanation.

Discrete-primitivism stance (a framework-level declaration aligned here; see the cited monograph for details): SRE dynamics explicitly denies the fundamental status of the continuous Riemannian manifold — "there is no preordained continuous Riemannian manifold; the spacetime dimension, the gravitational coupling, and the speed of light all emerge macroscopically from the topological connectivity density of a discrete causal network." The refinement-limit test (gap ∝ n^{−1.99} → 0), regarded as a "continuum-limit" test of a discrete object, therefore has a premise that does not apply within discrete primitivism; $n = 60$ is the node count of a discrete Möbius ladder, and no continuum limit $n \to \infty$ exists. The three reviews above thereby translate into the three gaps that Fork A must deliver (next chapter).

---

6. Fork A axiomatization: the double-cover four-state complex

Fork A (discrete primitivism), in order to fill in the three gaps exposed by the Tier 0 review of Section 5, introduces an axiomatic derivation system. All axioms are inherited from the Möbius topology theory of light (DOI: 10.5281/zenodo.21025053) and the existing SRE axioms; the validation program is `code/_forkA_derivation_check.py`.

6.1 The axiom set

No.	Axiom	Source
A1	The interaction state space is a two-body four-state space: {H,L}×{H,L} = {(H,H), (H,L),(L,H),(L,L)}	four-state eigen-spectrum of the Möbius topology of light
A2	The electron and light share the Möbius-ring topology; the parameterized closure period is 4π; φ and φ+2π are co-phase sheets (double-cover identification)	Axiom II of the Möbius topology of light
A3	The state refresh connects all nodes in order along the 4π closure circuit (the base ring = the tour trajectory)	SRE refresh dynamics
A4	Phase-space convention n ≡ dim(E) × dim(F) = |E|×β1 (edge-format × loop-flux)	existing SRE formalism (Maxwell-engine structure)

Combination rule R1 (the only new assumption of Fork A, see §6.2): the number of independent circulation channels = the number of state-localized channels + the number of global holonomy channels.

6.2 Gap 1: derivation of the double-cover four-state complex (DCF)

P1.1【imposed, by A1】The state space contains 4 combinatorial states.

P1.2【imposed, by A2】Möbius double cover: each combinatorial state exists on two sheets (φ and φ+2π are co-phase sheets), hence

```
|V| = 4 × 2 = 8
```

P1.3【imposed, by A3】The refresh tour passes through all 8 nodes in order with period 4π (nodes 0–3 are the four states of sheet 1, enter sheet 2 through the twist edge 3→4, and twist back to sheet 1 via 7→0), forming the base ring C_8:

```
Base ring edge count |E⁰| = 8
```

P1.4【R1 combination rule + A2 + A1】Independent circulation channels:

- 4 state-localized channels: each combinatorial state carries one localized circulation (inheriting the four-state structure of A1);
- 1 global holonomy channel: the Möbius band is non-orientable, so traversing 2π causes a sheet flip; this twist circulation is a globally enforced topological mode.

```
β1 = 4 + 1 = 5
```

P1.5【imposed, by P1.4】The base ring C_8 itself contributes $\beta1^0 = 1$; each non-repeated chord increases the number of independent loops by exactly 1, hence the number of chords = β1 − 1 = 4:

```
|E| = 8 + 4 = 12
```

P1.6【A4】Phase-space dimension:

```
n = dim(E) × dim(F) = |E| × β1 = 12 × 5 = 60
```

Audit: the placement of the chords is not fixed by the axioms (there is a degree of freedom in the placement of the skip-2 channels), but the placement does not enter the n and α chain: validation shows that any 4 non-repeated skip-2 chords (Maxwell's actual placement / two-sheet symmetric / uniform placement) give β1 = 5 and n = 60. The empirically measured (|V|,|E|,β1) = (8,12,5) of the D_edge of the Maxwell engine is consistent with the derivation; the source-code label "K*{3,5}" is a misnomer (the true K*{3,5} has 15 edges and β1=8), and the correct name of this complex is the **double-cover four-state complex (DCF)**.

Conclusion on Gap 1: under A1–A4 + R1, (8, 12, 5, 60) is uniquely determined. The only new free assumption is R1 (the combination rule "number of channels = number of states + number of holonomy channels"); the holonomy term is enforced by non-orientability, and the state term is inherited from A1.

---

7. The observable dictionary: α = λ₂/λ_max = v/c

7.1 Generators and frequency conventions

The generator of the SRE refresh dynamics is the **combinatorial Laplacian** L = D − A on the state graph: the Laplacian_1st = D^T·D and the P_E projection correction in the Maxwell engine are precisely the edge-space implementation of this structure. The eigenmodes = relaxation modes; the eigenvalues = dimensionless refresh rates. The frequency formula $f \propto \Delta\lambda$ is a linear convention for eigenvalues, so the propagation speed on the graph is proportional to the eigenvalue (rather than its square root), and the speed ratio = the eigenvalue ratio.

7.2 The master identity

On the state graph of the residual manifold:

- the stiffest mode λ_max = the maximum refresh rate = **the light mode** (the most rigid sector (H,H) in the four-state language);
- the softest nontrivial mode λ_2 = the smallest nonzero refresh rate = **the electron phase mode** (the (L,L) soft sector);

- the two modes share the lattice spacing of the same graph, hence

```
α = v/c = λ₂/λ_max
```

The fine-structure constant = the ratio of the electron phase velocity to the speed of light (v/c = α in the Bohr semantics), and the spectral-gap ratio is precisely a purely dimensionless rate ratio — dimension, structure, and semantics are aligned. Same-origin interpretation: the two ends of the spectrum of the same residual-manifold graph are precisely the electron and the photon — this is the spectral formulation of "the electron and light share the Möbius topology" (A2), and also explains why α couples the electron and photon coupling constants.

7.3 Closed-form spectrum of the weighted ladder

The state graph = the Möbius ladder M_n (n even): ring edges (i,i±1) have weight 1 (intra-sheet refresh), and cross-sheet identification edges (i, i+n/2) have weight w (Möbius twist channel). The adjacency is A = A(C_n) + wP, where P: i → i+n/2 is an involution ($P^2 = I$) commuting with the cyclic shift. The Fourier modes v_k diagonalize both simultaneously:

- A(C_n)v_k = 2cos(2πk/n)·v_k
- Pv_k = (−1)^k·v_k

The degree is uniform d = 2 + w, hence

```
μ_k(w) = (2+w) − 2cos(2πk/n) − w·(−1)^k ,   k = 0,…,n−1
```

Numerical validation (n=60, w=1 and w=w*): the maximum deviation between the closed form and direct diagonalization is 5.3×10^{-15}.

7.4 Soft-mode topological protection (theorem)

For even k, (−1)^k = +1, and the w term **cancels exactly** against the +w in the degree:

```
μ_k(even) = 2 − 2cos(2πk/n)     ——independent of w
μ_k(odd)  = 2 + 2w − 2cos(2πk/n) ——depends linearly on w
```

The k=1 (odd) mode μ = 2+2w−2cos(2π/n) ≥ 2w + 0.011 is always larger than the even mode k=2, hence

```
λ₂(w)  = 2 − 2cos(4π/n)     ——exactly w-independent (numerically confirmed |Δλ₂| < 4×10⁻¹⁶)
λ_max(w) = 2 + 2w + 2cos(2π/n)   (for w > 1−cos(2π/n))
```

Physical interpretation: the electron mode is a purely topological quantity (completely insensitive to the twist-channel strength); the Möbius twist channel only renormalizes the light-speed mode. The numerator of α is topological; the denominator contains dynamics.

7.5 Correspondence level with the four-state frequency formula

The frequency formula f ∝ Δλ·(λ₂+λ₃)/λ₁ possesses a "difference/sum" ratio structure isomorphic to the spectral gap; the ladder gap is its discrete implementation. **Honest grading:** this is a structural-level correspondence (dictionary-level), not a strict reduction — the one-to-one mapping between the parameters β, γ, λ₀ of the four-state formula and the ladder spectrum has not been established.

Conclusion on Gap 2: the dictionary has been completed to a working level — α = λ₂/λ_max = v/c, with soft-mode topological protection as a precise theorem. The remaining underdetermined items are transferred to Gap 3.

---

8. Residual and bare coupling δ (A5 axiomatization)

8.1 Residual of the binary weight and the weighted generalization

At w = 1 (combinatorial weight), gap = (1−cos(4π/n))/(2+cos(2π/n)), which gives 0.007297458502 at n=60, higher than α by a relative amount 1.452×10^{-5}. By §7.4, gap(w) = (1−cos(4π/n))/(1+w+cos(2π/n)). Setting gap(w*) = α:

```
w* = (1−cos(4π/60))/α − 1 − cos(2π/60) = 1.000043470457132
δ ≡ w* − 1 = 4.347046×10⁻⁵
```

Validation: gap(w*) = 0.007297352569284 = α* (relative deviation 1.2×10⁻¹⁶, machine precision).

Physical reading (a falsifiable coupling-asymmetry proposition): the residual is no longer an unexplained number but a precise proposition —

> **The coupling strength of the Möbius twist (cross-sheet identification) channel exceeds that of the intra-sheet refresh channel by δ = 4.347×10⁻⁵.**

This asymmetry shifts the light-speed mode λ_max upward and reduces the ratio to α. Any independent SRE observable (any coupling ratio derived in the future, a repaired λ_0 chain, or a measurement of the twist channel) can confirm or falsify it.

8.2 Channel algebra and the modulus no-go theorem

Channel algebra: the state graph = the weighted Möbius ladder: the ring refresh channel $S+S^{-1}$ (weight 1, unit convention of A3), and the cross-sheet identification channel wP (P = S^{n/2}, identification of the A2 double cover). The Laplacian:

```
L(w) = (2+w)·I − S − S⁻¹ − wP ,   wP = w·S^{n/2}
```

Key structural facts (numerically confirmed to machine precision): P is a polynomial in S, hence **[S, P] = 0 (strict)**, and $P^2 = I$ (strict). The channel algebra $\mathcal{A} = \mathbb{C}[S]/(S^n - I)$ is a **commutative algebra** — all eigenmodes are simultaneously Fourier-diagonalized, and w can enter the closed forms μ_k(w) only as a coefficient; it **cannot play the role of an eigenvalue selection rule**. This is the algebraic root of "why symmetry analysis cannot give δ": in a commutative algebra, the channel weight is a modulus, not a charge that can be quantized by symmetry.

Modulus no-go theorem: under A1–A4 + R1:

1. The full set of constraints of the axioms on w = {w > 0} ∩ {w ≍ 1} (an open interval, not a single point) — an axiom-by-axiom audit: A1 imposes no constraint; A2 requires only w ≠ 0 (the identification channel exists); A3 fixes the ring weight = 1 (unit convention) and imposes on w only that it be of the same order; A4 and R1 impose no constraint;
2. All structural theorems (connectivity, soft-mode protection, two-sector splitting, the two spectral ends = the electron/photon) hold for **every** w in the interval (numerical validation: C1–C4 hold to machine precision over the grid w ∈ {0.05,...,10}, with $\lambda_2(w)$ deviating from 2−2cos(4π/60) by < 1.8×10⁻¹⁶);
3. gap(w) is strictly monotonically decreasing ⇒ given α, w* is unique — but (1)(2) make w* unselectable by the axioms.

Corollary: δ = w* − 1 is **in principle underivable from A1–A4+R1**. It is not an "incomplete derivation" but a free modulus independent of the ring-refresh unit convention — a new independent constant of the theory.

8.3 The A5 axiom and closure

> **A5 (twist-channel coupling asymmetry):** the cross-sheet identification (holonomy) channel is stronger than the intra-sheet refresh channel by δ, where δ is a bare constant of the theory whose numerical content is supplied by measured input (with the same status as α in QED).

Closure check (dual track): the analytic quotient (1−cos4π/60)/(1+w+*cos2π/60) reproduces α* to a relative deviation of 1.2×10⁻¹⁶; the full-spectrum min/max cross-check is 5.0×10⁻¹⁵. **A1–A4 + R1 + A5 ⇒ α (machine precision)**, with the direction of derivation δ → α.

8.4 Parameter-independent signature and the δ-universality prediction table (pre-registered)

Odd-mode lifting theorem (secant-line test 8.9×10⁻¹⁶):

```
μ_k(1+δ) − μ_k(1) = 2δ   (k odd, photon sector — the whole set lifted exactly)
μ_k(1+δ) − μ_k(1) = 0    (k even, electron sector — the whole set strictly unmoved)
```

That is, the excess coupling of the twist channel acts **only** on the photon sector, and the lift per mode is strictly identical (representative values: μ_1: 2.010956209263 → 2.011043150178; λ_2: 0.043704798532 → unchanged). This is the parameter-independent spectral fingerprint of δ.

δ-universality prediction table: A5 is a per-edge bare coupling ⇒ for the Möbius state graph of any even n, it predicts α_pred(n) = gap_n(1+δ). Pre-registered table (n = 8..120 already frozen in the result JSON); within the range [60,120], **only n=60 falls within the 1% window of α*** (n=58 deviates +6.79%, n=62 deviates −6.50%, bracketed monotonically on both sides). Any future new (|E|, β1) structure must lie on the table rather than at α* — a multi-structure consistent determination of δ would promote it from a bare constant to a cross-validated theoretical constant.

Negative controls (registered before computation, identification criterion < 10^{-12}): the candidate mechanisms λ_0^2 (δ/candidate 1.0548), α^2 (0.8163), $2/S^6$ (1.0141), $\lambda_0 \cdot \gamma$ (0.1158), $\gamma^2/2$ (0.0254) are **all not identified**, and δ retains its bare-constant status and must not be back-fitted. The geometric weighting rules (1/d: 2.53%, $1/d^2$: 6.73%) were already excluded in §8.1, with deviations 20–40 times larger than the binary weight.

8.5 Fork A final scorecard

Gap	Status	Content
Gap 1: derivation of the complex	**Filled**	A1–A4 + R1 → (8,12,5,60); the only new assumption = the R1 combination rule; chord placement free and irrelevant
Gap 2: observable dictionary	**Filled (dictionary level)**	α = λ_2/λ_max = v/c; closed-form spectrum; soft-mode topological-protection theorem; the two spectral ends of one graph = electron/photon
Gap 3: residual mechanism	**Axiomatically closed (A5)**	modulus no-go theorem: δ underivable from A1–A4+R1; A5 bare coupling + odd-mode lifting signature + δ-universality prediction table; geometric rules and all candidate mechanisms excluded by negative controls

Honest bottom line: α = f(n=60) is derived, the topology is derived, δ = A5 bare constant. Fork A compresses the theory's free constants from three (α, λ_0, κ) to **one** (δ), and this constant carries a parameter-independent falsifiable signature. This is fully isomorphic to the bare-constant status of α as a measured input in QED — the only difference being that SRE reduces the "electron–photon coupling ratio" to the ratio of the two spectral ends of a 60-node Möbius state graph plus one per-edge bare coupling. The topological existence of δ is enforced by Z_2 non-orientability (Book of the Void L335: "2N mutual measurements to return to the initial symmetric state" = the physical manifestation of Z_2 holonomy); the quantitative value of δ requires the input α* (algebraic rearrangement, $\alpha(C+\delta)=N,\ C=2+\cos(\pi/30),\ N=2\sin^2(\pi/30)$, sensitivity $d\delta/d\alpha \approx -410$); an independent measurement of δ is currently infeasible (all schemes depend indirectly on α* through w*). In all standard topological tools δ is a metric parameter rather than a topological invariant; the Book of the Void (DOI: 10.5281/zenodo.22162514) confirms that "there is no preordained continuous Riemannian manifold", so δ is an emergent parameter on a discrete graph, and there is no δ in the continuum-limit sense.

---

9. The electron = an emergent object of the binary self-organizing network

The preceding chapters derive independently, from the Fork A axioms, the 60-node projection of the electron (DCF (8,12,5,60)) and the α dictionary. This chapter argues: **the electron projection is not an independent graph superimposed on SRE; it is itself a binary self-organizing network.** The 60-node Möbius ladder is merely a **specific realization** of the **coherent core** of the binary network under the conditions of "sparsification + holonomy closure". This reinforcement demotes the electron projection from "an object asserted by the axioms" to "an emergent substructure of the binary network", thereby fully aligning it with the basal ontology of SRE.

Four correspondences (numerical prototype established, see `code/_sre_electron_binary_bridge.py`):

Level	Claim	Evidence
Vocabulary/algebra	The adjacency of the electron projection is a {±1} matrix, of the same kind as Axiom 1 of M_n	entries of M_core ∈ {+1,−1}: True
Generation	The coherent-core scale k coincides with the electron scale n	N=300 ⇒ k=⌊0.2N⌋=60 = n
Topology	The Z_2 double cover is the native holonomy of the network	4-loop signed product strictly takes {+1,−1}

Level	Claim	Evidence
Spectral	α = λ₂/λ_max is a native spectral invariant of the binary network	Fork A ladder with w=1+δ reproduces α to 8.8×10⁻¹⁵

Correspondence dictionary (electron-projection ↔ binary-network):

Electron-projection element	Corresponding object in the binary self-organizing network	Basis
Adjacency matrix (edge weights ∈ {±1})	Axiom 1: $S_{ij} \in \{+1,-1\}$, $S_{ij} = S_{ji}$	isomorphism
Möbius Z₂ double cover	**signed product** holonomy of closed causal loops ∈ {+1,−1}	§7.4
Closed causal loop [A₁,...,A_N]	closed walk / circulation channel in the coherent core	circulation = β₁
4π phase period (spin 1/2)	Z₂ holonomy: restored only after **two** turns (double-cover identification φ~φ+2π)	A2
60-node scale n	coherent-core scale k=⌊0.2N⌋ (Theorem 1)	§2.2
Emergent geometry (MDS inversion)	MDS inversion of the coherent core (multi-scale homomorphic-mapping prototype)	Appendix A of the monograph

Figure 9-1 (figures/sre_electron_binary_bridge_figure.png): schematic of the bridging reinforcement between the electron projection and the binary self-organizing network, showing the four-level correspondence (vocabulary/generation/topology/spectral) and its connections to the coherent core and the MDS geometry.




Figure 9-1: the electron 60-projection as an emergent substructure of the binary-network coherent core (model-internal picture, not a real measurement).

9.1 Basal structure: Q₃ (12×5)

Within the model, compressing the coherent core toward the minimal invariant kernel via the sparse operator R (composed of dormancy pruning and Z₂ winding closure), one obtains upon non-compressibility the structure (8, 12, 5):

- vertex count 8, edge count 12, number of independent closed loops 5;
- this structure is precisely the **three-dimensional cube Q₃** (8 vertices, 12 edges, 5 independent loops);
- `|E| × β₁ = 12 × 5 = 60` — this is the true identity of the electron's "60-node projection": it is not 60 independent nodes, but a counting expansion of the basal structure `(12×5)`.

In short, **within the SRE model, the topological prototype of the electron is the cube Q₃; the number 60 is the count of "edge count × independent-loop count" of this cube** — not 60 independent degrees of freedom.【The above ruling is a model-internal graph-theoretic description, not a geometric measurement or physical assertion about the real electron】

9.2 The "deep yet compact" property of the electron

- **Deep:** logical depth `N = λ_c / ℓ_min ≈ 10⁻¹² / 10⁻³⁵ = 10²³`. The electron is composed of approximately `10²³` nested relational levels, and this depth is its **mass carrier (model-internal analogy)** (mass corresponding to topological depth/cumulative path count of interior loops).
- **Compact:** basal structure `12×5`, i.e., the **stable topological invariant (model-internal)** surviving after the depth folding.
- **Folding:** the homomorphic quotient map `h: deep loop (10²³) → basal structure (12×5)`. The topological type is determined by the Fork A axioms, while the visible granularity is determined by the resolution gap `1/10²³`. The two are complementary and mutually non-conflicting.

【Remark: the logical depth `10²³` and the "mass carrier" are both model-internal counts and analogies, not the measured mass or measured depth of the real electron. The folding is the homomorphic quotient map h: G_loop(N~10^{23}) → G_basal(12×5) with compression ratio ~N; the statistically stable invariants retained (charge → Z_2 holonomy; spin-1/2 → Z_2 double cover; mass → topological depth/cumulative path count of interior loops; intrinsic loop structure → β_1=5, |E|=12) map exactly onto the basal topological invariants.】

9.3 α as a native spectral quantity of the network

A spectral analysis of the electron's relation matrix shows that the ratio of its second-softest mode to the stiffest mode is `1/137.036`. Within the model, this value is not implanted externally but is a **generic spectral quantity** of the binary-network relation graph; the Möbius ladder of Fork A is merely the specific sparse realization that places this spectral quantity exactly at 1/137.【**Important contextual limitation:** here α is a model-internal spectral quantity; its agreement with 1/137.036 is an internal reproduction within the model and does not constitute a real-physics conclusion that "the fine-structure constant is derived by this model"; the numerical value of the real α remains determined by experiment】

9.4 Emergence of space

Defining the relational distance `D_ij = √(2 − 2·S_ij)` from the relation matrix `S`, and then applying MDS inversion, yields three-dimensional coordinates for the electron (see the `60×3` output of `code/_sre_electron_binary_bridge.py`). **Within the model, shape emerges from relations** — this is the fundamental divergence between SRE and the traditional view of "space first, particles placed within it afterwards".【Remark: the coordinates obtained by MDS inversion are model-internal relational-embedding coordinates, belonging to the model picture, not measurements of real spatial positions】

9.5 Three-layer emergence structure and the RG flow

Repeatedly applying the coarse-graining operator `C` (merging a cluster of nodes into a single super-node) to the binary network yields a **renormalization flow (model-internal)**:

```
micro: binary relation net —C→ electron coherent core (12×5) —C→ meso: atom/molecule kernel —C→ macro: bulk
```

The coarse-graining operator C_η merges sub-kernels of the coherent core into super-nodes (block homomorphism + re-binarization), with $n \to n/\eta$ (η>1):

$$M'[b_i, b_j] = \mathrm{sgn}\Big(\prod_{u \in B_i,\, v \in B_j} M_{uv} \Big) \in \{\pm 1\}$$

The output is still a {±1} matrix (vocabulary conservation), and the homomorphic mapping preserves the group structure, so the Z_2 double cover is conserved under compression at any scale (verified in `code/_sre_coherence_rg.py`). Micro: the electron basal structure (8,12,5) is the microscopic coherent-core seed; meso: repeated application of C causes the microscopic kernel to recombine upward into atomic/molecular coherent cores; macro: continuing C to the bulk, the **BBP phase transition z*=3.13** (Book of the Void v1.6, DOI: 10.5281/zenodo.22162514; the old value 4.1605 has been demoted to a historical reference) is the scale at which the global coherent core switches from the 2D holographic single channel to the 4D two-channel compensation, i.e., the macroscopic spacetime-release point. The statistically stable basal structures at the various scales are the **attracting fixed points** of the RG flow at the respective scales: the electron 12×5 is the microscopic fixed point, and atoms/molecules/bulk are fixed points at higher scales.【**Important contextual limitation:** the renormalization flow, the "micro–meso–macro" correspondence, and the z*=3.13 cosmological-phase-transition release are all constructions and analogies internal to the SRE model (within the framework of DOI: 10.5281/zenodo.22162514); conclusions about real cosmology and the levels of matter must be examined separately by independent observations and theories】

---

10. The proof chain: uniqueness, convergence, and robustness

The following four-step arguments all hold within the SRE simulation model: they prove that the fixed point of the sparse operator R is **unique, convergent, and robust** "within the model"; they are results of the logical self-consistency of the model, and **do not directly equate to laws of real physics**. Any generalization toward the real world must be undertaken with caution and is subject to external testing.

Step	Claim established	Key idea	Conclusion
Fork A	the electron projection = 60 nodes	the four topological axioms A1–A4 and rule R1 → DCF(8,12,5) → n=60	the origin of the 60 nodes is explicit
P0	why precisely (8,12,5)	enumerating the 3-regular graphs on 8 vertices yields 5; of these **only Q_3** simultaneously satisfies planarity, bipartiteness, and free-Z_2 (the electron signature)	**uniqueness established**
R2	the sparse operator necessarily converges	`R` has non-increasing node count and stops in a finite number of steps; the strict version satisfies `R²=R` (idempotence)	**convergence theorem holds**
R3	robustness under noise/dissipation	ε calibrated physically from the network dormancy probability (=0.18); the batch operator `R_batch` resists edge flips, executing in O(depth) steps	**robust-convergence validation passed**

Closed-loop conclusion (model-internal): the electron projection = the product of the Fork A axioms = the **unique, stable, robustly convergent fixed point** of the sparse operator of the binary network — the three are fully corresponding. The electron thereby ascends from a "floating independent graph" to the "inevitable basal structure of the binary self-organizing network".【The "uniqueness/convergence/robustness" above are all model-internal self-consistency conclusions and are not automatically extrapolated to real physics】

Figure 10-1 (figures/sre_electron_schematic_EN.png): the overall schematic of the electron within the SRE framework (3×3 panels): row A shows the coherent-core heat map / Q_3 basal structure / Möbius 4π double cover; row B shows the convergence of the sparse operator R (64→8), the α spectrum (real w=1+δ), and the MDS three-dimensional inversion; row C shows the three-layer RG flow, the 10^{23}→12×5 logical-depth folding, and the conclusion statement.




Figure 10-1: overall illustration of the coherent core, basal structure Q_3, Möbius double cover, sparse-operator convergence, α spectrum, three-layer RG, and logical-depth folding (model-internal picture, not a real measurement). Vector version at figures/sre_electron_schematic_EN.svg.

10.1 P0: uniqueness (why precisely Q_3)

Detailed below (program `code/_sre_p0_uniqueness.py`, result `sre_p0_uniqueness_results.json`). The sparsification rule R = dormancy pruning (R_dorm) ∘ holonomy closure (R_holo):

- **Dormancy pruning R_dorm:** node coherence `c_i = |Σ_j M_ij| / (k−1)`; threshold = the overall median (data-derived, no free parameters); prune the dormant nodes with `c_i < median(c)`.
- **Holonomy closure R_holo:** identify Z_2-holonomy equivalent node pairs (those with `S[i,c] = −S[j,c]` on all columns `c` except their own), merging them into super-nodes; the super-edge sign = majority vote of the signs of the nonzero products between blocks (preserving the {±1} vocabulary).

Lemma 1 (R is a well-defined compression operator): R is deterministic/order-independent; monotone (each pass strictly decreases |V|); vocabulary-conserving ({±1,0}); Z_2-holonomy conserving (the 4-loop signed product remains strictly two-valued). Numerically: after shuffling the labels, the output (13,78,66) is fully identical; before and after contraction the 4-loop product +1/−1 remains strictly two-valued.

Lemma 2 (constructive reduction): performing the Z_2-holonomy lift L = lift(Q_3) on Q_3 (16 nodes: each Q_3 vertex split into a pair of row-opposite holonomy-equivalent nodes), then R_holo(L) **reconstructs Q_3 exactly** — the lifted graph (16,32,17) closes under holonomy to (8,12,5), fully isomorphic to Q_3. That is, R, as the projection operator "double cover → basal", is well-posed and invertible.

Theorem (the unique fixed point of R): over all Z_2-holonomy coherent cores, the unique nontrivial stable fixed point of R is isomorphic to the 3-cube Q_3, i.e., DCF (8,12,5). By Lemma 2, Q_3 is a fixed point of R; by Lemma 1, any fixed point must be a Z_2-holonomy coherent complex; the Fork A axioms pin down the basal-complex invariants as (8,12,5); uniqueness is given by the enumeration lemma.

Uniqueness lemma (exhaustive proof): among all connected 3-regular graphs on 8 vertices, those satisfying (planar embeddability) ∧ (bipartiteness) ∧ (free-Z_2: the existence of a fixed-point-free involutive automorphism, i.e., the algebraic signature of the Möbius 4π double cover) are unique, namely Q_3. Enumeration result: 19320 labeled graphs → 5 up to isomorphism (consistent with the known mathematical fact), of which only Q_3 (girth 4) satisfies all three as True.

```
|V| |E| b1  planar  bip  freeZ2  girth
 8  12  5   True  True   True      4   ← Q3 (DCF)
 8  12  5   True  False  True      3
 8  12  5   True  False  True      3
 8  12  5   False False  False     3
 8  12  5   False False  True      4
[uniqueness] number of graphs satisfying (planar & bipartite & freeZ2): 1  =>  basal uniquely determined as Q3
```

Closure of P0 with Fork A: Fork A (topological-axiom lineage) independently derives DCF (8,12,5); P0 (binary-network lineage) independently derives the same (8,12,5) as the unique fixed point of the sparse operator R. The two independent lineages converge on the same basal complex, i.e.,

> **The electron projection = the product of the Fork A axioms = the unique fixed point of the sparse operator of the binary network** — the three are fully corresponding.

10.2 R2: convergence (explicit construction + convergence theorem)

Detailed below (program `code/_sre_r2_convergence.py`, result `sre_r2_convergence_results.json`). `R_strict` is the strict Z_2-holonomy quotient: the equivalence relation `i ~ j ⇔` row i and row j are overall opposite after removing the diagonal; the inter-class edges `C[a,b] = sign(Σ_{x∈C_a,y∈C_b} M[x,y])` of the quotient graph are vocabulary-conserving and Z_2-holonomy conserving. `R_eps` is the dissipatively relaxed merge (physical dissipation parameter ε): each single step $R_1(M)$ merges the node pair with globally minimal holonomy-tension and tension ≤ ε;

```
tension(i,j) = 1 − (number of agreeing neighbors / number of non-all-zero neighbors)
  where "agreeing" means M[i,k] == s·M[j,k],  s = sign(Σ_k M[i,k]·M[j,k])
```

Convergence theorem: Lemma 1 (monotonicity) any single step R_1 does not increase the node count; Lemma 2 (idempotence, strict version) $R_{strict}^2 = R_{strict}$ (the quotient-of-quotient of an equivalence relation equals the original quotient); Lemma 3 (termination) the R_1 iteration stops within ≤ $|V_0|$ steps. Hence **Theorem R2 (convergence)**: for any M, the sequence $R_1^t(M)$ converges to a fixed point within ≤ $|V_0|$ steps; the strict operator reaches it in one step.

Fixed point = Q_3 (depth→basal reduction + uniqueness closure): applying R_strict after performing an L-level Z_2 lift on Q_3:

Input	Trajectory (R_{strict}^t)	Terminal state
lift(Q_3,1) (16 nodes)	16 → **8**	Q_3 (8,12,5) ✓
lift(Q_3,2) (32 nodes)	32 → 16 → 8	Q_3 ✓
lift(Q_3,3) (64 nodes)	64 → 32 → **8**	Q_3 (8,12,5) ✓

Each Z_2 double-cover level corresponds to one level of logical depth; R_strict progressively eliminates the double covers level by level, ultimately reaching the irreducible basal graph = Q_3 — turning the folding mechanism of "logical depth 10^{23} → basal structure 12×5" into a **computable level-by-level quotient** rather than a one-shot assertion.

Honest boundary (key finding): applying R to the real 300-step snapshot M_core does not collapse (R_strict: V=60, no exact holonomy twin pairs; R_eps: no value of ε∈[0,0.5] lands on Q_3). Explanation: the real M_core is a **resolution-truncated snapshot**, whose row structure has no explicit antipodal pairing and is itself a fixed point of R. The fixed point of R = the input itself, unless the input carries holonomy redundancy; the deep loop (10^{23}) carries the antipodal structure by A2 of Fork A, so R(deep loop) → Q_3. The two are not contradictory but **level-complementary**: the 60-node projection is the basal structure of the electron (12×5=60), and the convergence of R to the 8-node Q_3 is a property of the deep-loop level, not a property of the 300-step snapshot.

10.3 R3: robustness (ε physical calibration + O(L) + noise robustness)

Detailed below (programs `code/_sre_r3_batch.py`, `code/_sre_r3_deepcore.py`).

ε physical calibration: mirroring the sim_p evolution loop to extract the dormancy probability — `ratio = λ·d/(|M@M|+1)`, `p_act = 1 − 1/(1+ratio)`, `dorm = 1 − p_act = 1/(1+ratio)`. Collecting ~1.6×10⁶ dormancy-probability samples over the evolution with steps=300 (λ=0.8, seed=1111) at coherent scales (n=50..300), the **median = 0.1832**. ⇒ **ε = 0.1832** is the physical dissipation scale given by the network growth dynamics, no longer a free parameter back-tuned to the result (its status parallels A5=δ in α).

Noise-free deep-core convergence: on lift(Q_3,L) ($2^L \cdot 8$ nodes, L=1..4), all three operators — strict/greedy-ε/batch-ε — reduce Q_3 (8,12,5); the batch version still holds at L=5 (256 nodes).

Step-count scaling: R_strict is O(L) (one traversal per double-cover level, at most L passes); the R_eps (greedy) step count ≈ $2^L \cdot 8 - 8$ grows **exponentially** with depth; **R_batch_eps is O(L) ≈ 1 effective parallel pass** (one pass merges all holonomy-linked pairs simultaneously). ⇒ The expected O(L) scaling is realized by the batch operator, so the batch version is adopted as the physical implementation.

Noise robustness: flip the edge signs with probability ρ on lift(Q_3,L) (worst-case perturbation). The greedy R_eps is fragile (brittle; can recover only within a ρ window ≤0.02); **the batch R_batch_eps is robust**: L=1 passes for all ρ≤0.2, L=2 for ρ≤0.15, L=3 for ρ≤0.1, and it is insensitive to the exact value of ε (restoration almost everywhere within the 2×ε neighborhood). Mechanism: a single edge flip destroys only a few edges of one antipodal pair; the batch operator still merges the remaining clean pairs in that pass, blocking error propagation.

Honest note: random edge flips are the worst-case perturbation; the real electron holonomy is protected by the Möbius 4π double cover (not random edge noise), so the physical robustness is higher than this lower bound. Batch merging corresponds precisely to the synchronized dormancy-pruning semantics of sim_p and is the physically correct choice.

---

11. Unified validation program (10/10)

The suite `code/sre_electron_validation_suite.py` asserts each of the 10 key conclusions of the entire proof chain above one by one and outputs the machine-readable result `sre_electron_validation_report.json`:

```
python sre_electron_validation_suite.py
```

The suite reuses the verified functions of the `_sre_*.py` modules (bridge / P0 / R2 / R3), serving only orchestration and assertion duties, so that the entire chain is reproducible in one pass. **Important note:** during execution the suite repeatedly evolves the binary network and performs spectral, graph-theoretic, and exhaustive assertions; its conclusions are only **self-consistency tests within the SRE simulation model**. "10/10 all passed" means that the steps of the model are internally logically consistent and the results are reproducible; it is **not a comparison against real physics experiments**, and one cannot claim on this basis that real physics has been proven.

11.1 Measured results (10/10 all passed, model-internal self-consistency)

No.	Conclusion	Measured result
C1	vocabulary identity `{±1}`	N=300, k=60, coherent core ∈ {±1} ✓
C2	scale correspondence `k=n=60`	`⌊0.2×300⌋=60` = electron projection 60 ✓
C3	Z_2 holonomy native	4-loop product +1:0.504 / −1:0.496 ✓
C4	α native spectral quantity	Möbius(60, w=1+δ) gap=7.297e-3 = 1/137 (error 6.8e-9) ✓
C5	P0 uniqueness	3-regular graphs on 8 vertices number only 5, only Q_3 carries the electron signature ✓
C6	P0 constructive reduction	lift(Q_3,16) → R → (8,12,5) ✓
C7	R2 idempotence	`R²=R` ✓

No.	Conclusion	Measured result
C8	R2 depth→basal	L=1,2,3-level lifts all collapse to Q_3 ✓
C9	R3 ε physical calibration	ε=0.1832 (network median dormancy probability, physical input) ✓
C10	R3 noise robustness	batch operator still recovers Q_3 under edge flips ρ≤0.1 ✓

Contextual limitation: the "electron projection", "Z_2 holonomy", "α", "Q_3", etc. in the table above are all model-internal objects and indicators; the "✓" of C1–C10 indicates only that the model-internal assertions pass, i.e., the model is logically self-consistent, and **does not constitute a comparison against or proof of real physics experiments**.

---

12. Molecular-computation application

This section gives the **transferable inferences** of SRE for molecules and their routines. The file `code/sre_molecular_application.py` is self-contained, has no external dependencies, and can be run directly. It must first be stated: the transfer in this section is an application of the SRE model-internal methodology to the heavy-atom connection graphs of real molecules, in which all inferences about real molecules are **only hypotheses extrapolated from the model outward**, unverified by DFT, molecular dynamics, or experiment (see Section 14).

The transferable conclusions of SRE for molecules are summarized in four points:

- **K1 binary relation matrix:** any molecule can be written as `S_ij ∈ {+1(bond), 0/−1(non-bond)}`, i.e., the same `{±1}` language as the electron (model-internal convention).
- **K2 spectral-feature index MCI:** the electron's `α=λ₂/λ_max` methodology is transferred into the "skeleton spectral fingerprint" of the molecule (small values correspond to flexible/low-frequency bending, large values to rigid closed shells). **It must be emphasized: MCI is only a graph-theoretic skeleton-topology fingerprint and does not equal the quantum-mechanical electron-coherence effect of a real molecule.**
- **K3 basal-structure recognition:** the basal structure of the electron (model-internal) is $Q_3(8,12,5)$; any molecule whose carbon-skeleton graph-theoretic invariants are exactly Q_3 (such as **cubane**), its heavy-atom connection graph satisfies the Q_3 graph-theoretic invariants. **Note: graph-theoretic invariant matching ≠ the real molecule automatically possessing a long-lived electron-coherence quantum effect.**
- **K4 relational-inversion geometry:** the three-dimensional shape of the electron is obtained by MDS-inverting the relation matrix (model-internal construction); the same holds for molecules — reasonable conformations can be inverted from the topologically relational data alone (a model-internal relational embedding, not a real conformational measurement).

12.1 Routine 1: the SRE coherence index MCI (spectral fingerprint)

```
import numpy as np, networkx as nx

def sre_coherence_index(adj):
    """MCI = λ2/λmax (signed graph Laplacian L = D − S). Small values correspond to flexibility, large values to rigidity.
    Note: MCI is a graph-theoretic skeleton-topology fingerprint; MCI ≠ the quantum-mechanical electron-coherence effect of a real molecule."""
    S = adj.astype(float)
    D = np.sum(np.abs(S), axis=1)
    L = np.diag(D) - S
    ev = np.sort(np.linalg.eigvalsh(L))
    return float(ev[1]) / float(ev[-1])      # MCI
```

12.2 Routine 2: three-dimensional-cube basal recognition (graph-theoretic invariant detection)

```
def sre_basal_detect(adj):
    """Determines only whether the graph-theoretic invariants of the heavy-atom connection graph match Q3
(8,12,5) as a bipartite graph;
    does not mean that the real molecule possesses the corresponding quantum-coherence behavior."""
    G = nx.from_numpy_array((adj != 0).astype(int))
    V, E = G.number_of_nodes(), G.number_of_edges()
    b1 = E - V + 1
    is_Q3 = (V, E, b1) == (8, 12, 5) and all(d == 3 for _, d in G.degree()) \
            and nx.is_bipartite(G)
    return {"is_electron_basal_Q3": is_Q3, "invariants": (V, E, b1)}
# Note: here Q3 refers only to graph-theoretic invariant matching; graph matching ≠ the real molecule
possessing the quantum-electron behavior corresponding to Q₃
```

12.3 Routine 3: relational-inversion geometry (MDS embedding + RMSD comparison)

```
def sre_mds_embed(adj, ref_coords):
    G = nx.from_numpy_array((adj != 0).astype(int))
    n = G.number_of_nodes()
    D = np.array([[nx.shortest_path_length(G, i, j) for j in range(n)] for i in range(n)])
    X = mds_coords_from_dmat(D, ndim=3)      # invert three-dimensional coordinates (model-internal relational
embedding)
    return X, procrustes_rmsd(X, ref_coords) # compute RMSD against the reference coordinates
```

Figure 12-1 (figures/sre_mds_inversion_3d_EN.png): example of the three-dimensional embedding by MDS relational inversion (model-internal relational embedding); **Figure 12-2** (figures/sre_mds_inversion_fidelity_EN.png): fidelity check of the inversion (the intrinsic deviation between graph-distance metric and Euclidean geometry).




Figure 12-1: MDS three-dimensional inversion embedding of the electron relation matrix (model-internal relational embedding, not a measurement of real three-dimensional positions).




Figure 12-2: fidelity check and deviation distribution of the MDS inversion (model-internal metric).

12.4 Measured example output (benzene / cubane / n-hexane)

```
molecule   atoms  bonds      MCI    Q3?   RMSD(A)  note
benzene        6      6   0.2500     no     1.474  芳香环, 刚性闭合壳 (MCI 较大)
cubane         8     12   0.3333    YES     1.414  碳骨架=Q3, 直接承载电子本体拓扑
n-hexane       6      5   0.0718     no     0.000  线性链, 极小 MCI = 柔性骨架(低频弯曲)
```

Interpretation of the results:

- The heavy-atom connection graph of the carbon skeleton of **cubane** satisfies the Q_3 graph-theoretic invariants (the (8,12,5) bipartite graph), and is judged by `sre_basal_detect` as "graph-theoretically hitting the electron basal Q_3". **This is only a graph-theoretic invariant match**: it does not automatically mean that the real molecule possesses long-lived electron-coherence quantum effects, which must be verified separately by external computations such as DFT and femtosecond dynamics (see the tension paradox below).
- **MCI** as a skeleton-rigidity fingerprint: the linear chain takes 0.07 (flexible, easily bent), and the ring and cube take 0.25–0.33 (rigid closed shells). This is a molecular descriptor transplanted directly from the electron-α (spectral-gap-ratio) methodology. **Note: MCI is only a graph-theoretic skeleton-topology fingerprint and does not equal the quantum-mechanical electron-coherence effect of a real molecule.**
- **MDS** can invert the shape from topology alone (model-internal relational embedding): the linear chain is reproduced **exactly** (RMSD=0), while the ring and cube are reproduced **approximately** (owing to the intrinsic deviation between the graph-distance metric and Euclidean geometry). The absolute bond lengths are filled in from chemical input — this precisely corroborates the SRE stance: "relations give shape, physical scales are filled in afterwards" (model-internal picture).

12.5 Supplementary discussion: cubane and the tension paradox

In real chemistry, cubane exhibits a well-known tension paradox: owing to the distortion of the 90° bond angles, the molecule stores a huge strain energy and is thermodynamically unstable, yet it is highly kinetically stable, with a very high activation energy for ring-opening cleavage. From the SRE model picture: the carbon heavy-atom skeleton of cubane happens to be the Q_3 basal topology, and the MCI fingerprint shows a very high graph-theoretic skeleton rigidity. This offers a phenomenological understanding at the topological level: although the geometric deformation brings a large internal energy, the Q_3 topological network itself possesses the property of resisting skeleton rearrangement, hindering the ring-opening topological reconstruction, corresponding to the kinetic stability observed in the tension paradox.

Important boundary statement: this interpretation is a **topological-perspective reading internal to the SRE model** and does not replace the microscopic mechanistic explanations of the traditional valence-bond–molecular-orbital theory for banana bonding, diradical cleavage intermediates, etc. The Q_3 graph-theoretic invariants describe only the connection relations between atoms; they do not contain microscopic orbital information such as orbital hybridization, bond-angle potential energy, or intermediate-state energies. The model inference awaits further cross-validation in conjunction with DFT molecular-orbital computations.

12.6 Application to one's own structures (.mol2 / VASP / xyz)

Read the atomic coordinates and bond tables of the molecule, construct `adj` (bonds take `+1`), and then call the routines above. For real force-field molecules (the `.mol2` / GROMACS systems in the repository), it is recommended to treat hydrogen as non-bonded and use only the heavy-atom skeleton, which makes the MCI and basal-detection results clearer. This also provides a new **topological pre-screening hypothesis** for traditional molecular dynamics: one can first compute the MCI and the Q_3 hit, and then decide whether to invest in expensive quantum-chemical computation. **It must be emphasized:** this pre-screening outputs graph-theoretic features, is a hypothesis from the model outward, and cannot replace the conclusions of quantum-chemical computation.

---

13. The laws of electron orbital assignment

This chapter addresses a naturally extending question: **if the electron is the object characterized in Section 10, then what law should govern "how orbitals in the atom are assigned to electrons"?** It adopts a "bottom-up" route: starting not from the Schrödinger equation but only from the six definitional premises (P1–P6) that SRE imposes on the electron, it progressively derives four laws of orbital assignment and reproduces them numerically by means of a purely combinatorial enumeration program (`code/_sre_orbital_assignment.py`).

Contextual limitation (read first): the objects "electron", "orbital", "Pauli exclusion principle", "Hund's rule", "shell capacity", etc. in this chapter, as well as the quantum numbers, Möbius double cover, logical depth, spectral gap, and other concepts involved, are all constructions, analogies, and counts internal to the SRE model. Among these, the "Pauli exclusion principle (cell-uniqueness)" and "Hund's rule (tension minimization)" are derived directly from the mechanism of the SRE electron definition; the "magnetic-quantum-number degeneracy $2\ell+1$" and the "construction order $(n+\ell)$" require borrowing the rotational symmetry of the emergent three-dimensional space and the Coulomb-repulsion order, and are **open assumptions**. The real orbital-filling law remains governed by quantum mechanics; SRE does not constitute a substitute here.

13.1 Existing premises: the SRE definition of the electron

No.	Premise	Content	Basis
P1	fixed point	electron = the unique, stable, noise-robust fixed point of the sparse operator **R** over the coherent core of the binary self-organizing network	P0 / R2 / R3
P2	basal structure	the basal complex of the electron is the three-dimensional cube $\mathbf{Q_3}$: $\|V\|=8$, $\|E\|=12$, $\beta_1=5$, i.e., $12\times5=60$	P0 uniqueness enumeration
P3	spin	spin-1/2 = **Möbius 4π double cover / Z_2 holonomy**: 8 basal vertices × 2 sheets	Fork A / §9
P4	space	space emerges as three-dimensional coordinates (60×3) from the relation matrix via the **MDS inversion**	§9.4
P5	stability criterion	stable fixed point = configuration of **structural-tension minimization** (the dissipative version of R merges the minimal holonomy-tension pair at each step)	R2 / R3
P6	hierarchy	atom = **mesoscopic layer** = the recombination of the coherent core one level upward (the second layer of the three-layer RG flow)	§9.5

All subsequent content of this chapter takes P1–P6 as the sole starting point and introduces no new axioms or empirical parameters.

13.2 State space: the 16 cells and the faithful cover

The basal state space of a single electron takes the combination of each vertex of the basal Q_3 with the spin sheet of the Möbius double cover:

$$\mathcal{U} = \{(v,\sigma) : v \in \{0,1\}^3,\ \sigma \in \{+1,-1\}\}, \qquad |\mathcal{U}| = 8 \times 2 = 16.$$

where v are the 8 vertices of Q_3 (three-bit triples) and σ are the two sheets of the double cover (corresponding to spin ↑/↓). 16 is the total number of basal states distinguishable "for one electron" at the intrinsic level. **Key structural property — faithful cover:** on the double-cover relation matrix M_{16}, the "relational patterns to all other units" (row vectors) of the 16 units are **pairwise distinct** (numerical validation: the 16 rows are mutually distinct). This indicates that the 16 cells are mutually distinguishable in the relational sense and the cover is **faithful**.

13.3 The Pauli exclusion principle: cell-uniqueness at the network level

Claim (Pauli = cell-uniqueness, model-internal derived item): a single cell (v, σ) accommodates at most one electron.

Argument: (1) if two electrons occupy the same cell, they produce two fully identical identical relational patterns in the relation matrix of the mesoscopic coherent core; since the faithful cover requires the relational patterns of the individual units to be mutually distinct, the appearance of identical rows marks the degeneration of the cover, and the second electron contributes no new state whatsoever. (2) From the operator perspective, the holonomy closure of the sparse operator R merges rows that are "overall-opposite-equivalent" into the same super-node; while fully identical (same-sign) rows form a degenerate spectrum. Numerical demonstration: after forcibly duplicating row of unit 0 on the double-cover matrix, the matrix develops 1 pair of fully identical rows — the "second copy" produces no new state.

Conclusion: in the state space constituted by the 16 cells, the capacity of each cell is exactly **1**. If a unit can accommodate both sheets σ=+ and σ=− (P3), the two sheets within the cell are still two different units $(v,+)$ ≠ $(v,-)$, each accommodating one electron — corresponding to the legal cohabitation of two electrons with opposite spins on one orbital.

13.4 Orbitals and the quantum-number dictionary

An "orbital" in SRE is defined as: the **collective vibration mode** provided by the atom (the mesoscopic recombined kernel, P6) that a single electron can occupy, i.e., the eigenvector of the relation Laplacian of the mesoscopic coherent kernel; its eigenvalue λ serves as the energy order parameter (same origin as the conclusion that the electron spectral gap $\alpha=\lambda_2/\lambda_max$ is a native spectral quantity of the network). The SRE correspondence of the four quantum numbers is as follows (**items marked ★ are borrowed terms**):

Quantum number	Symbol	SRE correspondence	Status
spin	`s`	Möbius double-cover sheet number σ ∈ {+1,−1}	derived item (P3)
radial/shell	`n`	level number of the nested coarse-graining of the mesoscopic kernel	derived item (P6, RG level)
azimuthal/subshell	`ℓ`	order of the irreducible representation of rotational symmetry in the emergent three-dimensional space	★ borrowed item (inference from P4)
magnetic	`m`	standard basis vectors of the above representation, `-ℓ≤m≤+ℓ`, 2ℓ+1 in total	★ borrowed item (same as above)

Honest remark: the symmetry of Q_3 itself gives the hypercube degeneracy `1,3,3,1` (see §13.8 V1), **not** the spherical-harmonic `1,3,5,7`. The `2ℓ+1` degeneracy comes from a reasonable extension of the premise treated in P4, that "three-dimensional space emerges from the MDS relational inversion" — the emergent three-dimensional space possesses rotational symmetry, and its irreducible representations give `2ℓ+1`. This is one **borrowing** of the model-internal account toward real geometry.

13.5 The capacity laws

Proceeding from the §13.3 cell-uniqueness (capacity 1 per cell) and the §13.4 degeneracy, cumulative summation level by level gives the three-level capacities (pure arithmetic):

Single-orbital capacity: one orbital `(n,ℓ,m)` corresponds to two spin sheets → **2**.

Subshell capacity: `m` runs over `-ℓ..+ℓ`, a total of `2ℓ+1` orbitals, each accommodating 2 electrons →

$$C_\ell = 2\,(2\ell+1) = 4\ell+2.$$

Measured subshell capacities: `s=2, p=6, d=10, f=14, g=18, h=22, i=26`.

Shell capacity: summing over `ℓ = 0,1,…,n-1` →

$$C_n = \sum_{\ell=0}^{n-1}(4\ell+2) = 2n^2.$$

Measured shell capacities: `n=1→2, 2→8, 3→18, 4→32, 5→50, 6→72, 7→98`. The identity `Σ_{ℓ<n}(4ℓ+2)=2n²` holds for all `1≤n≤7`.

13.6 Filling order: Aufbau / Madelung (open item)

Electrons occupy the mesoscopic-kernel modes in ascending order of energy (corresponding to the quantum-mechanical construction principle). SRE currently **cannot uniquely derive** the energy-level order; this manuscript adopts the prevailing **Madelung diagonal rule** as an open item:

> Orbitals are filled in ascending order of `(n + ℓ)`; when `(n + ℓ)` is equal, in ascending order of `n`.

The filling order thereby generated: `1s, 2s, 2p, 3s, 3p, 4s, 3d, 4p, 5s, 4d, 5p, 6s, 4f, 5d, 6p, 7s, 5f, 6d, 7p, …`. The contribution of SRE to this rule lies in a **structural explanation**: the existence of the `(n,ℓ)` two-component quantum numbers comes from the §13.4 dictionary; the "diagonal" form of the ordering (bands of the same level in the RG flow when n+ℓ is constant) is formally consistent with the three-layer recombination structure of P6. But SRE does not claim to uniquely derive this ordering — this is an explicitly marked open boundary.

13.7 Hund's rule: tension minimization

Claim (Hund = tension minimization, model-internal derived item): for the `2ℓ+1` degenerate orbitals of the same `(n,ℓ)` with a total of `k` electrons, define the "pairing tension" as the number of orbitals whose two spin sheets are simultaneously occupied (paired). Then **the occupation minimizing the pairing tension is exactly equivalent to first distributing the electrons among distinct orbitals with co-oriented spins (maximum net spin)**.

Numerical validation (purely combinatorial enumeration):

- ℓ=1 (p-type, 3 orbitals / 6 cells): for k=1, 2, 3 the minimal pairing cost is 0, and the maximum net spins are 1, 2, 3 (all singly occupied, all co-oriented); for k=4, 5 the minimal costs are 1, 2 (forced pairing), with net spins 2, 1.
- ℓ=2 (d-type, 5 orbitals / 10 cells): for k=1…5 cost 0, net spin 1…5; for k=6…9 cost 1…4, net spin 4…1.

All enumerated sites satisfy: minimal pairing tension ⟺ maximum unpaired spin; and the theoretical bounds `min-pair-cost = max(0, k-(2ℓ+1))` and the maximum spin `min(k, 4ℓ+2-k)` agree. Half-filling (k=2ℓ+1) is exactly all-singly-occupied with maximum net spin, which is precisely the most stable configuration called Hund's rule.**SRE significance:** Hund's rule here is not a "placed" empirical postulate but the natural result of the P5 tension-minimization mechanism over the degenerate multiplets.

13.8 Periodic-table reconstruction and the numerical coincidence of 60

Enabling §13.3 Pauli, §13.5 capacity, §13.6 order, and §13.7 Hund simultaneously, one can reconstruct the shell structure of neutral atoms: the noble-gas boundaries `Z = 2, 10, 18, 36, 54, 86, 118` all fall on complete subshells; `Z=19` (potassium) gives `…4s¹` (rather than `3d¹`) and `Z=26` (iron) gives `4s² 3d⁶`, consistent with the prevailing construction.

One numerical coincidence worth recording (co-directional indication, not proof):

$$\sum_{n=1}^{4} 2n^2 = 2 + 8 + 18 + 32 = 60 = |E| \times \beta_1.$$

That is, **the basal count 60 of the electron equals exactly the total electron capacity of the first four principal shells (n=1..4, i.e., 1s through 4f)**. In accordance with the established honesty principle of this project, this coincidence is positioned as a "co-directional indication": the appearance of consistency between 60 in the electron basal structure (|E|×β₁=12×5) and "the total closed-shell capacity up to 4f" suggests an intrinsic connection between the basal count and the shell levels, but it **does not constitute a proof** and is not automatically extrapolated to real physics.

Figure 13-1 (figures/sre_electron_orbital_fill_EN.png): the schematic of the orbital-filling laws — the 16-cell state space, the capacity brick wall (2/8/18/32…), and the Madelung construction order.




Figure 13-1: SRE orbital assignment — the 16-cell state space, shell-capacity building blocks, and the (n+ℓ) construction order (model-internal derivation; vector version at figures/sre_electron_orbital_fill_EN.svg).

13.9 Validation program and measured results

The program `code/_sre_orbital_assignment.py` is a pure numpy/networkx implementation with no external dependencies:

```
python _sre_orbital_assignment.py   # outputs sre_electron_orbital_assignment_results.json
```

No.	Object of validation	Measured result
V1	Q₃ spectral degeneracy	eigenvalues 0,2,2,2,4,4,4,6; Hamming-weight degeneracy `{1,3,3,1}` (hypercube, ≠2ℓ+1) ✓
V2	Pauli = cell-uniqueness	the 16-cell relational rows are pairwise distinct (faithful cover) = True; 1 pair of identical rows appears after forced double-occupation ✓
V3	capacity arithmetic	Σ_{ℓ<n}(4ℓ+2) = 2n² holds for all n=1..7 ✓
V4	Hund = tension minimization	at all k sites of ℓ=1, 2 the minimal pairing cost = the theoretical value max(0,k−(2ℓ+1)) ✓
V5	Madelung order	`1s 2s 2p 3s 3p 4s 3d 4p …` (open item, demonstration only)

No.	Object of validation	Measured result
V6	periodic-table reconstruction	noble gases Z=2,10,18,36,54,86,118 are all complete subshells; potassium `4s¹`, iron `4s²3d⁶` ✓
V7	60 coincidence	Σ_{n=1..4} 2n² = 2+8+18+32 = 60 =

Contextual limitation: the "✓" of V1–V7 indicates only that the model-internal enumeration and assertions pass, i.e., the SRE logic is self-consistent; it **does not constitute a comparison against or proof of real orbital-filling experiments**.

---

14. Honest boundaries

14.1 Conclusions rigorously established within the model

The following conclusions are **rigorously established** (within the SRE simulation model) by constructive proof or numerical exhaustion:

1. The electron projection = the **unique** nontrivial stable fixed point of the sparse operator `R` of the binary network (P0: exhaustively enumerating the 3-regular graphs on 8 vertices yields 5 candidates, of which only Q_3 carries the electron signature); the basal structure (8,12,5)=Q_3, and 60=|E|×β_1.
2. `R` **converges** to that fixed point (R2: monotone + idempotent + finite-step termination; $R_{strict}^2=R_{strict}$).
3. Under the physical dissipation ε=0.1832 and the edge noise ρ≤0.1, it still converges **robustly** (R3, batch operator O(L)).
4. α as a native spectral quantity of the network (model-internal); the Fork A ladder can **reproduce it to 1/137** (error about 1e-9, an internal reproduction of the model); the Fork A axiomatization compresses the theory's free constants into a single δ (A5 bare coupling).
5. The derived items of electron orbital assignment: the faithful cover of the 16-cell state space, capacity 1 per cell (Pauli = cell-uniqueness), the capacity identities 2 / 4ℓ+2 / 2n², and Hund's rule = tension minimization (derived from P5, verified by exhaustion).

14.2 Items still to be improved

- The coincidence of the number `60` with the coherent-core scale `k=60` is currently a **co-directional indication** (the 0.2 coefficient originates from Theorem 1, but `N=300` is a simulation choice) and has not yet rigorously derived "frozen-kernel size = 60" from the BBP phase transition.
- The **analytic convergence** of `R` (a closed-form proof of `Rⁿ → Q₃`) has numerical support and an observed O(depth) scaling, but the closed-form proof remains to be supplied; it is necessary to construct a coherent core of sufficiently deep resolution to directly observe `R(deep core)→Q₃` numerically.
- The **global phase-transition structure** of the three-layer RG flow and its strict alignment with `z*=3.13` remain to be completed (i.e., the R4 task); that phase transition is a construct of the model-internal (Book-of-the-Void framework), and real cosmological conclusions must be examined separately.
- The macro/meso layers are currently theoretical generalizations: the microscopic seed (electron 12×5) has model-internal axiomatic support, but the conclusions at the molecular and bulk levels need to be completed independently.
- The **2ℓ+1 degeneracy** and the **(n+ℓ) construction order** in the orbital assignment are both borrowed items (emergent three-dimensional rotational symmetry, Coulomb-repulsion order), not uniquely derived by SRE; the Q_3 self-degeneracy 1,3,3,1 and the spherical-harmonic 1,3,5,7 have different origins (recorded in V1). SRE currently has no counterpart of the Coulomb potential/nuclear charge; the eigenvalue mapping of the n shells and ℓ subshells is not closed.
- **60 = Σ_{n≤4}2n² is a co-directional indication, not a proof**; nor can it be enforced by P1–P6.
- An independent measurement of δ is currently infeasible (all schemes depend indirectly on α* through w*); δ is a Z_2-graded continuous coupling strength (a structure unique to the SRE framework) with no direct counterpart in the standard model.

14.3 The important gap at the molecular level

> Extrapolating from the electron basal Q_3 to real molecules (such as cubane), **graph-theoretic invariant matching ≠ the real molecule's quantum-electron-coherence behavior automatically holding**; MCI and the Q_3-detect output only graph-topological features. Real electron coherence is a quantum-dynamical effect jointly determined by excited states, nuclear vibrations, and environmental coupling, and must be further verified by DFT/femtosecond-dynamics simulations; it cannot be asserted directly from graph indices how strong the real electron coherence is. Everything at the molecular level is an inference from the model outward and has not been calibrated against a large real-molecule dataset.

14.4 Overall contextual limitation

All of the arguments above — from the Fork A axioms, through the P0 uniqueness, the R2 convergence, and the R3 robustness, to the 10/10 pass of the unified validation suite — **hold within the SRE heuristic-simulation model**, representing the self-consistency and internal coherence of the model's logic, not a proof of real physics. The "electron", "Q_3", "60 nodes", "logical depth", "Z_2 winding phase", "Möbius double cover", "renormalization flow", and "cosmological phase transition" in this manuscript are all constructions, analogies, and counts internal to the model, not physical quantities directly measurable in the real world. This work does not prove that the SRE axioms hold objectively; the numerical reproduction of α indicates only the internal self-consistency of the framework after independent review; even if all validations pass, this indicates only that the model's internal logic is closed, **not automatically equating to facts of real physics**.

---

15. Conclusions

This manuscript gives a unified and complete characterization of the electron within the SRE dynamics framework, unfolding along a reproducible logical chain:

1. **Theoretical foundation:** SRE proceeds from three discrete-topology axioms (binary constraint, asynchronous activation, geodesic field); the Möbius topological modeling of light gives the geometric origin of the 4π closure period and of half-integer spin.
2. **The fine-structure constant:** three paths independently reproduce α — the Laplacian spectral gap of the Möbius ring graph (n=60, error 0.0015%), the four-state eigen-spectrum frequency formula (error 0.0000%), and the Möbius-parameterized arc-length perturbation (first-order exact). The Tier 0 independent computational review confirms that λ_0 is not an independent primitive, the grid refinement limit FAILs (the n=60 hit is a discretization coincidence), and the crossing point is $n_\alpha=60.000436$.
3. **Fork A axiomatization:** A1–A4 + R1 independently derive the DCF complex (8,12,5,60); an observable dictionary $\alpha = \lambda_2/\lambda_{max} = v/c$ is established (with soft-mode topological protection as a precise theorem); the residual, shown by the modulus no-go theorem to be in principle underivable, is promoted to the A5 bare coupling $\delta = 4.347\times10^{-5}$. The theory's free constants are compressed from three to one (δ), carrying a parameter-independent falsifiable signature (odd-mode lifting 2δ) and a universality prediction table.
4. **Emergent characterization of the electron:** the electron = the **unique, stable, robustly convergent fixed point** of the sparse operator R over the coherent core of the binary self-organizing network; its basal structure is Q_3 (12×5=60); the "deep yet compact" property (10^{23} logical depth folded into the 12×5 basal structure) and the three-layer RG flow (micro–meso–macro, with the macroscopic BBP $z^*=3.13$ release) are unified in the same binary network and the same coarse-graining operator.
5. **Validation:** the unified validation suite passes 10/10 (model-internal self-consistency, not a comparison against real physics).
6. **Application one (molecular computation):** three transferable routines are given — the MCI spectral fingerprint, Q_3 basal recognition, and MDS relational-inversion geometry — with benzene/cubane/n-hexane as measured examples; the cubane Q_3 graph-theoretic match is a model-internal phenomenological interpretation and does not replace traditional chemical mechanisms.
7. **Application two (orbital assignment):** from the 16-cell state space, Pauli = cell-uniqueness, the capacity laws $C_{\ell}=4\ell+2$ and $C_n=2n^2$, and Hund's rule = tension minimization are derived; the noble-gas boundaries and the potassium/iron construction order are reconstructed; the numerical coincidence of 60 with the capacity of the first four shells is recorded (co-directional indication).

Unified stance: all proofs and numerical validations in this manuscript are internally self-consistent constructions of the SRE model. Any generalization toward the real world (DFT, molecular dynamics, experimental observation) must be independently verified externally. This manuscript does not claim any real-physics conclusion beyond the internal self-consistency of the model. The dynamical origin and independent measurement of δ, the numerical construction of the deep core 10^{23}, and the global phase-transition alignment of the three-layer RG flow with z*=3.13 (R4) are left to future work.

Appendix A: Key numerical values

Quantity	Symbol	Value
Fine-structure constant	α	1/137.035999 = 0.00729735257
Vacuum basal spin residual	λ_0	0.00641954 (Tier 0 ruling: not an independent primitive)
Chiral-locking coefficient	κ	0.828
Spectral scaling	β	21.09256
Chiral coupling	γ	0.0585
Möbius bandwidth	w = √(8α)	0.241617
λ_0/α	(4+4κ)/(5+4κ)	0.87971
α/λ_0	(5+4κ)/(4+4κ)	1.13674
Optimal node count of the graph	n	60 = 12 × 5
Möbius topological confidence	—	99.2094%
Twist-channel coupling asymmetry	δ	4.347×10^{-5} (A5 bare constant)
Exact crossing point	n_α	60.000436
A5 closure relative deviation	—	1.2×10^{-16} (analytic quotient)
Spectral-gap ratio	gap(60,1+δ)	0.007297352569284 = α*
R3 physical dissipation	ε	0.1832 (network median dormancy probability)
Validation suite	—	10/10 passed (C1–C10)
Noble-gas boundaries	Z	2, 10, 18, 36, 54, 86, 118

Appendix B: Merged glossary

(see the glossary of Section 1 §1.1 and the symbol conventions of §1.2 — the terms of the four component documents have been merged and deduplicated, and the full table is collected here.) Supplementary terms: **fermion** = an identical particle obeying the Pauli exclusion principle, which cannot doubly occupy the same quantum state (expressed within the SRE model in this chapter as "cell-uniqueness"); **MCI (SRE coherence index)** = the molecular skeleton spectral fingerprint λ_2/λ_max (a graph-theoretic descriptor, ≠ real quantum electron coherence).

Appendix C: Code listing and full reproduction

All programs cited in this manuscript are collected in the `code/` directory (42 files, for packaging and distribution). Core reproduction commands:

```
# Step 1: baseline search (prove that the standard graph topology does not contain α)
python _verify_alpha.py          # DCF exhaustive enumeration
python _scan_alpha_graphs.py     # scan of 150 graph types

# Step 2: Möbius topology search (discovery of the 4π period and n=60)
python _verify_alpha_mobius.py   # Möbius ring / Klein / RP²
```

```
python _verify_alpha_light3.py   # validation of the Möbius topology formulas of light

# Step 3: exact validation (convergence of the three paths)
python _verify_n60_lambda0.py    # n=60 + λ₀ + arc length

# Step 4: external self-consistency (Maxwell equations + circuit simulation)
python Maxwell.py                # Gauss' law
python sre_simulation.py         # KVL/KCL/Ohm

# Step 5: operator validation (closed-form solutions of Operators 11/12)
python _verify_v11_closed_form.py

# Step 6: Tier 0 theoretical review (independent computational review)
python _tier0_lambda0_independent.py
python _tier0_refinement_limit.py
python _tier0_theory_reaudit.py

# Step 7: Fork A axiomatic derivation + δ axiomatization
python _forkA_derivation_check.py
python _delta_axiomatization.py

# Step 8: electron emergence chain (bridge / P0 / R2 / R3 / three-layer RG)
python _sre_electron_binary_bridge.py
python _sre_p0_uniqueness.py
python _sre_r2_convergence.py
python _sre_r3_deepcore.py
python _sre_r3_batch.py
python _sre_coherence_rg.py

# Step 9: unified validation suite (the whole chain 10/10, model-internal self-consistency)
python sre_electron_validation_suite.py

# Step 10: molecular application + orbital assignment
python sre_molecular_application.py
python _sre_orbital_assignment.py
```

Runtime environment: Python 3.10+; NumPy ≥1.20; SciPy ≥1.7; NetworkX (validation suite and orbital assignment). MDAnalysis ≥2.0 and GROMACS 2021 are required only by the ethanol-analysis script/MD upstream. **Note:** `Maxwell.py` and `sre_simulation.py` come from the electrical-quantity validation (DOI: 10.5281/zenodo.22119957 / 10.5281/zenodo.22119635), and `sim_p.py` is the binary-network evolution engine (hierarchical dissipative self-organization, DOI: 10.5281/zenodo.20576606).

Appendix D: Figure listing

No.	File (figures/)	Corresponding section	Content	Model status
Figure 9-1	sre_electron_binary_bridge_figure.png	§9	bridging correspondence between the electron projection and the binary network	model-internal picture
Figure 10-1	sre_electron_schematic.png / .svg	§10	overall schematic (3×3 panels)	model-internal picture
Figure 12-1	sre_mds_inversion_3d.png	§12	MDS three-dimensional inversion geometry	model-internal relational embedding
Figure 12-2	sre_mds_inversion_fidelity.png	§12	MDS inversion fidelity	model-internal metric
Figure 13-1	sre_electron_orbital_fill.png / .svg	§13	orbital-filling laws	model-internal derivation
Note	sre_electron_schematic_EN.png	English counterpart	English version of the overall schematic	model-internal picture

Appendix E: References and DOIs

1. Yue Lu. Topological derivation of the fine-structure constant within the SRE framework. Zenodo. https://doi.org/10.5281/zeno

do.22765850

2. Yue Lu. SRE Dynamics: The Book of the Void. Zenodo. https://doi.org/10.5281/zenodo.22162514
3. Yue Lu. SRE Dynamics User Guide. Zenodo. https://doi.org/10.5281/zenodo.19935370
4. Yue Lu. Conjecture on the Möbius topology of light (source of light_3.pdf). Zenodo. https://doi.org/10.5281/zenodo.20482974
5. Yue Lu. Technical report: the intrinsic algebraic topology of light and the SRE axion matrix. Zenodo. https://doi.org/10.5281/zenodo.21025053
6. Yue Lu. SRE dynamics: strict reconstruction of Maxwell's field equations from purely dimensionless graph cohomology and global evolution steps. Zenodo. https://doi.org/10.5281/zenodo.22119957
7. Yue Lu. Dynamics of hierarchical dissipative self-organizing binary networks. Zenodo. https://doi.org/10.5281/zenodo.20576606
8. Yue Lu. Emergence of multi-dimensional spacetime and gravity. Zenodo. https://doi.org/10.5281/zenodo.20837960
9. Yue Lu. SRE electrical quantities defined (charge/current/resistance/voltage/power/E=mc²). Zenodo. https://doi.org/10.5281/zenodo.22119635

Note on citation conventions: wherever the Book of the Void (DOI: 10.5281/zenodo.22162514) is the source (e.g., the "2N mutual measurements" of Z_2 holonomy, the sampling theorem, discrete primitivism L723, BBP z*=3.13), the text cites by line/section number with the DOI and does not transcribe the original text; the quoted textual content of the remaining existing SRE documents has, in accordance with the fusion requirements, been transcribed directly into the body.

This manuscript is written in standard written scientific language, relying on terminology definitions and progressive exposition so that readers without a physics or mathematics background can understand the "complete characterization of the electron within the SRE model"; the molecular routines and orbital-derivation routines can be transferred directly to one's own systems, but their inferences about real molecules require verification by external computation and experiment. This manuscript is the merged manuscript of the four aforementioned component documents, with all formulas, code, and numerical values kept unchanged; all conclusions are limited to the internal self-consistency of the SRE model and do not constitute assertions about facts of real physics.

Complete Characterization of Nucleons within the SRE Framework: Unified Representation of Two-State Skeleton Inversion, Quark Speculation and Nuclear Reaction Validation

Version: 2.1 (Integrated Draft)

Date: 2026-10-09

> [Resource and Availability Statement] This framework is built based on State-Relational Entropy (SRE) dynamics. All theoretical materials are archived in the Zenodo open data repository. **This document suite includes system papers, application development, scientific hypotheses, and the complete algebraic derivations and simulation code for all operators 1-19, fully open source**.

> Additionally, you can access the Tencent Intelligent Document Space supporting AI-assisted consultation (available on PC and WeChat mobile terminals).

> As of 2026-09-21, restricted by Google Terms of Service, the author no longer maintains or updates the SRE document library in Google Gemini Notebook. This link is for historical archive only and should not be used as an official citation source:

> - Google Gemini Notebook (Historical Archive, No Longer Updated): https://notebooklm.google.com/notebook/ef52bf5a-f6d0-4a2a-aed4-b25d6520ab2c
> - Tencent Intelligent Document: https://docs.qq.com/space/DUkRjYUtNWFdyV253

> According to the principles of State-Relational Entropy (SRE), classical physical foundations originate from information statistics.

Project References (DOI, see Appendix E for in-text citations):

- https://doi.org/10.5281/zenodo.22077475 — SRE-v1.6 Axiom Suite
- https://doi.org/10.5281/zenodo.22162514 — SRE-Dynamics Composite Elementary Particles and Relational Space Emergence (Book of Void)
- https://doi.org/10.5281/zenodo.22119957 — SRE Dynamics: Strict Reconstruction of Maxwell Field Equations Using Pure Dimensionless Graph Cohomology and Global Evolution Steps
- https://doi.org/10.5281/zenodo.20576606 — Hierarchical Dissipative Self-Organizing Binary Network Dynamics
- https://doi.org/10.5281/zenodo.22119635 — SRE Electrical Quantity Definitions (Charge/Current/Resistance/Voltage/Power/$E=mc^2$)
- https://doi.org/10.5281/zenodo.22162514 — Complete Characterization of the Electron within the SRE Framework (Series)

> **Core Conclusion**: In the SRE (State-Relational Entropy) theoretical framework, nucleons are not point particles pre-placed in space, but **stable composite objects emerging from binary self-organizing networks on the tripartite Y-shaped coherent core, constrained by the two-state opening/closing of dormant edges and closure degree rebalancing rules**. Their ontological structure is **the tripartite Y-shaped triangular closure $Y_3 \ltimes \triangle_3$** (12 vertices, 18 edges, first Betti number 7, automorphism group order 36).

> This conclusion starts from the clue of neutron β-decay, and is obtained through three-step screening: integer determination of vertex number V=12, spectral ratio determination of topology, and triple symmetry determination of solution, with a combined rarity of approximately 4.7×10^{-3}. It is verified through fission chain reactions, and the two-state opening/closing mechanism of nuclear reactions also holds at the composite nucleus and macroscopic scales. The nucleon ontology can be extended to derive the fission cross-section parity law, closure degree potential criterion and chain criticality, and deduce that the additivity failure of ion/metal pairs can be theoretically predicted, and current is a coherent bookkeeping projection rather than continuous electron movement.

> In a further validation that borrows external nuclear data, the skeleton yields a zero-parameter many-body assembly law and shared-ring configuration: the lowest excitation on the shared ring is strictly independent of the body count ($\lambda_2 = 2-\sqrt{3}$, multiplicity k−1); and it yields a zero-parameter mirror-symmetric existence criterion **L1: $|n_n - n_p| \le 1$** (8 / 8 on the 8-entry scorecard). Accordingly, "nuclear boundness" must be split into two questions: **the numerical value of the binding energy** is explicitly delimited as a non-computable interval for A ≥ 4 (the out-of-sample prediction of the pricing layer fails, and that failure is **locked for the whole class and is irrepairable**, by an identity independent of the shape of the pricing function); **existence** is computable at A = 4 via L1 with zero parameters, but not for A ≥ 5 (the criterion degrades) — that is, the boundary is **relocated** from A ≥ 4 to A ≥ 5.

> **Contextual Limitations (Must Read First)**: The above-mentioned "nucleon", "$Y_3 \ltimes \triangle_3$", "two-state opening/closing", "closure degree potential" and other objects, as well as concepts such as logical depth, Z_2 winding phase, dormant edge, first Betti number mentioned later, are all heuristic simulation **model internal structures, analogies and counts**, and are not directly measurable physical quantities in the real world. All proofs and numerical verifications in this paper only hold within the SRE model, representing the self-consistency of the model, **and do not automatically equate to real physical facts**. Real physical and chemical phenomena must be independently verified by external means (DFT, molecular dynamics, experimental observations) for model inferences.

Abstract

This paper provides a unified and complete characterization of nucleons within the State-Relational Entropy (SRE) dynamic framework: nucleons are stable composite objects emerging from binary self-organizing networks on the tripartite Y-shaped coherent core, constrained by the two-state opening/closing of dormant edges and closure degree rebalancing rules. Their ontological structure is the tripartite Y-shaped triangular closure $Y_3 \ltimes \triangle_3$ (V=12, E=18, β_1=7, |Aut|=36). The full text unfolds along a reproducible logical chain:

(1) Establish the general methodology for nucleon derivation: local input → topological solution → inversion, which is homologous to electron derivation;

(2) Propose three ontological speculations about quarks (quark ≠ Q_3, quark is an emergent statistical phenomenon, nuclear reaction = closure degree rebalancing) as structural priors for subsequent inversion;

(3) Start from neutron β-decay, derive the structural constraints T1–T3 for the nucleon skeleton, and redefine isospin Z_2 as opening/closing duality, rather than flavor multiset flipping;

(4) Establish the nucleon inversion simultaneous equation system, and obtain surviving candidates through three-step screening (integer determination of vertex number V=12, spectral ratio determination of topology, triple symmetry determination of solution); then break the tie within the "transition–residue" framework: ρ is proved to be a strict invariant of the transition (the residue, serving as the reference standard of measurement), the dormant edge is taken to be a ring edge, and the criterion of "partial re-equilibration of closure degree" (the open state must still retain closed units) uniquely retains candidate A;

(5) Verify the two-state opening/closing mechanism of nuclear fission chain reactions: fast neutron cross-sections restore the same order of magnitude, excitation surplus criterion has a Pearson correlation of +0.932 with thermal cross-sections, and three-level signatures are unified (single nucleon β decay → composite nucleus fission → macroscopic chain criticality);

(6) Derive the theoretical constraints that the additivity failure of ion/metal pairs can be theoretically predicted, and current is a coherent bookkeeping projection rather than continuous electron movement;

(7) Extend the framework to many-body systems using external nuclear data as a yardstick: the assembly law is determined by two principles ("open–closed complementarity" and "the images of the two bodies must be distinguishable in the shared region", the latter being a non-spatial restatement of the former "closed bodies must not overlap"), yielding the structural linear law of the shared-ring configuration (V=9k+3, E=15k+3, T=2k+1, β_1=6k+1, |Aut|=6·2^k·k!) and the body-count independence of the lowest excitation ($\lambda_2 = 2-\sqrt{3}$, multiplicity k−1); a zero-parameter existence scorecard is given and the reading is adjudicated (Reading A adopted: 5/5 matched in the A ≤ 3 sector), and together with the out-of-sample failure of the pricing layer at A=4 and the qualitative failure of the assembly law at A=4, the honesty boundary "**the nuclear boundness for A ≥ 4 is not computable in this layer**" is drawn; it is further shown that this boundary is **not repairable on the side of the numerical binding energy** (the A = 4 isotriplet forms a **single isomorphism class**, so the binding-energy differences cannot be carried by any graph functional, and the additive-profile pricing class is falsified as a whole by an **identity** independent of the shape of the pricing function), and the disposal of the "shadow solution" (variant D) is completed (it is the k=2 member of the shared-ring family, and that name is withdrawn);

(8) an attempt is made to derive the **antisymmetric coefficient** λ required by that repair from within SRE; the conclusion is that λ can neither be derived nor fixed (its threshold, 29.0596 MeV, exceeds the ^{4}He binding energy, and the candidate pool is underdetermined; SRE fixes only the "shape" of λ, not its "value"). The same check, however, yields a positive outcome: demoting antisymmetry from an **energy term** to an **existence criterion** gives a zero-parameter, mirror-symmetric criterion L1 ($|n_n - n_p| \le 1$) that matches the 8-entry scorecard at 8 / 8, so that **the existence question at A = 4 is computable in this layer**; the out-of-sample test (A ≥ 5, 26 entries) shows that L1 does not have the status of a fundamental law (22 / 34 hits, its mismatches being respectively an α closure effect and the absence of an A scaling), so the boundary is **relocated** from A ≥ 4 to A ≥ 5;

(9) the objection "does the nucleon–nucleon force require the electronic state to be considered as well" is rewritten as a **configuration problem decidable within this layer**, with the following conclusion: its only decidable form is **open-band attachment to the shared region**, and that configuration is **isomorphic** to variant A (hence not a new structure, and already quantitatively excluded by the deuteron binding energy). The diagnosis further shows that **the open band is not an extra energy term but the carrier of principle 1 of the assembly law itself** — it appears as the "shape of the gap" (the presence or absence of the gap decides whether n-p can assemble, the position of the gap decides that n-n is excluded, and the k = 1 singly-claimed edge left after closure decides that the images of the two bodies are distinguishable). This also explains both the fact that the shared ring "always has exactly one k = 1 edge" and the body-count independence of $\lambda_2 = 2-\sqrt{3}$. The section additionally corrects a previously erroneous statement ("the neutron gap can only lie in ring 0" — the three rings are in fact a single automorphism orbit, so the statement is a notation rather than a fact, and the correction leaves every existing conclusion intact).

(10) two of the easiest-to-register branches of the "downward extension" screening are turned into executable criteria and executed: **the neutrino as the k = 1 singly-claimed edge on the shared ring** is **judged negative** — the automorphism group is **vertex-transitive** on the three shared-ring points (orbit count measured to be 1) ⇒ every graph functional takes the same value on the three-generation candidates ⇒ the spectrum is strictly degenerate; the 3×3 local operators always have only 2 distinct eigenvalues ⇒ the spectrum degenerates to two levels; and the ledger route (the claim triple always (#closed, k, k)) has only 1 degree of freedom. The two routes each strike a barrier already established in this project (the vertex-transitivity of 12.4, the identity of 12.11(2)); none of the 96 zero-parameter candidates is an effective hit. At the same time the **"exponential-decay-law gap" registered in Section 10 is filled**: the axioms "dormancy as the default state + removal as a rare, independent, stationary perturbation" **derive** $T_{1/2} = \ln 2/p$ and memorylessness (MC verification passes; the power law is excluded); but the strong reading of its two-factor decomposition $p = p_{\mathrm{graph}} \cdot p_{\mathrm{env}}$ is **falsified by measurement** — the zero-parameter prediction obtained from the species-cancellation law, "the stripping enhancement factor is independent of the nuclide", fails by **≥ 6.79 orders of magnitude** between ^{187}Re and ^{163}Dy (same mode, nearly equal Q), and because p_{graph} has already cancelled exactly, this **cannot be repaired by the graph factor**. The three pending items ①②④ thus merge into one and the same missing piece: "SRE fixes only the shape, not the value".

(11) The above merge is turned into a **pre-registered, falsifiable** test: the **L4 mass-depth equation** of Section 7.1, $m_p/m_e = \Psi(G_N)/\Psi(G_e)$ (with Ψ never defined), is built into a criterion and executed. The test admits as Ψ candidates a zero-parameter monomial class (integer powers of products of standard invariants, exponent $|a| \le A$, support at most three invariants), takes one **design target** (m_p/m_e) and two **out-of-sample targets** (m_d/m_p, m_n/m_p), and uses a decision tolerance $\varepsilon = 1\%$ (deliberately wider than the framework's demonstrated 0.165%). The verdict is **negative**, on three mutually independent grounds: (i) **reach bound** - with the electron basal body Q_3 as the electron graph, the full-support bound for $|a| \le 2$ is only **511.06**, short of the target 1836.15 by **0.555 orders of magnitude** (distribution-free, and independent of the support size); (ii) **no single lock** - the best Ψ differs from target to target, and the joint hit is **0** for $|a| \in \{1, 2, 3\}$ under both the 1% and the 3% window (i.e. the L4 requirement that "two graphs share one lock" is directly refuted); (iii) **hidden degree of freedom** - the electron-layer representative (basal body Q_3 vs the α carrier M_{60}) is not fixed by the framework, and the reach verdict flips between the two configurations, so "which graph represents the electron" is itself a free parameter (C3 violated). **Positive residue**: the same Π_1 machinery holds for the **difference ratio** of two states within a layer ($\alpha \cdot \Pi_1(\mathrm{Y}_3)$ against $(m_n - m_p)/m_p$: -0.164%) but fails for **inter-layer overall mass ratios** ($\Pi_1(\mathrm{Y}_3)/\alpha = 25.84$, short of 1836.15 by 1.85 orders of magnitude) => **the cross-layer lock governs the difference within a layer, but not the overall scale of a layer**. This work does not prove that the SRE axioms are objectively true; all conclusions are self-consistent constructions within the SRE model, and real physical and chemical inferences must be independently verified by external means.

(12) all the above conclusions are **converged into a single twist of observational perspective**: what this framework genuinely separates is not "whether some physical quantity is computed accurately", but **the boundary between a dimension-free SRE world and the three-dimensional world**. That boundary is given by a three-layer decomposition — **L0 the topological layer** (the graph; it yields only integer invariants, and neither ratios nor angles), **L1 the ratio layer** (dimensionless ratios, the intrinsic representation compatible with SRE; the "shape" of H_2O here is the ternary ratio $(1, 1, t)$, $t = 2\sin(\theta/2)$), and **L2 the geometric layer** (angles and coordinates, purely the output of a map). Each layer crossing costs one external input: L0 → L1 pays **the metric D** (including non-bonded association), and L1 → L2 pays **the map** (Euclideanity / curvature / scale). This yields two **theorem-level** criteria (independent of any numerical coincidence): (a) **angle ∉ functions(graph)** — the same graph P_3 with two admissible metrics gives 180.0000° and 73.7398°, while every graph invariant is identical term by term; (b) **same ratio, different map ⇒ different angle** — the same ratio $t_0 = 1.581140$ gives 104.4776° in the Euclidean case, and on the sphere ($R = 1$) gives 104.57° → 145.10° as the absolute scale varies, while in the hyperbolic case it gives 104.39° → 86.67°; and a curved space **has no similarity** (the angle also depends on the absolute scale) **⇒ geometry is a property of the map, not of the graph**. On this basis the negative verdicts of the preceding chapters receive a unified explanation: **every quantity SRE has passed so far is a dimensionless ratio** (Π_1, λ_2/ρ, $\Pi_1(Q_k) = 1/k$, $\lambda_2 = 2 - \sqrt{3}$, orbit counts, $|\mathrm{Aut}|$ ratios), **while every negative verdict occurs exactly when a dimensionful value is required to be assigned to some structure** (L4 assigning mass, the absolute value of α). In support of this, a **validation-qualification audit** is also completed: all 17 validations are classified by the epistemological type of their comparison target (independent experimental out-of-sample E = 9 | inversion / scheme-dependent quantity R = 2 | construction / circularity / back-substitution C = 4 | pure mathematics M = 2), and the result is that **the C and R classes fall 100% within the "geometry / shape" branch**, whereas the E class lies entirely within the spectral / numerical branch and is **a string of negative verdicts** ⇒ the "validation" of the geometry-shape branch degenerates into constructive self-consistency (measured against the three newly established qualification criteria, that branch currently has **0 qualified entries**), while the spectral / numerical branch is meaningful precisely because it uses independent experimental values; the repair direction is a purely topological D plus $n \ge 4$ plus comparing angles directly. The net conclusion of this paper is thus: **the strength of the framework lies in "fixing the shape" (laws, order, layering and dimensionless ratios), and its boundary lies in "fixing the value" (dimensionful real numbers); and the seam between "shape" and "value" must first pass through the explicit, nameable, declarable toll of "the map".**

Keywords: SRE dynamics; nucleon ontology; two-state opening/closing; topological inversion; β decay; fission chain reaction; quark emergence; closure degree potential; coherent bookkeeping

---

0. Document Structure and Reading Guide

This paper organizes all nucleon-related documents (v1 topological derivation, inversion v2 skeleton, quark speculation, many-body assembly law and shared ring, dimension collapse test, fission chain verification, theoretical constraints) into a unified representation in the following order:

- Section 1: **Terminology Convention**. Uncommon terms appearing for the first time in the text are marked with [], and their precise definitions are concentrated in this section, presented in plain written language for readers without physical or mathematical background. Each key term is accompanied by a note indicating its "model internal structure/analogy" status to avoid confusion with real physical quantities.
- Section 2: **SRE Theoretical Basis**: Binary self-organizing networks, sim_p generator and conservation closure, ratio methodology constraints, and the only calibrated projection Π_1.
- Section 3: **General Methodology for Nucleon Derivation**: Local input → topological solution → inversion, which is homologous to electron derivation, and the interchange principle between SRE and classical physics.
- Section 4: **Three Quark Speculations (Author's Claim)**, listed separately after the methodology chapter as they **determine what the nucleon skeleton should look like** — they are the "structural priors" for subsequent screening, and the screening results will change if they are altered. They are recorded truthfully to make readers aware: the skeleton in this paper is not "objectively emerged" from the data, but selected within the search space defined by this set of speculations.
- Section 5: **Entry Choice: Why Start with Neutron β Decay?**, including phenomena to be explained, three ontological readings, and structural constraints T1–T3 derived from β decay.
- Section 6: **Redefinition of Isospin Z_2 and Breaking Magnitude**, invalidating the C1 assertion in v1, redefining it as opening/closing duality, with a breaking magnitude of 10^{-3} as a structural necessity, compared with the electron side breaking.
- Section 7: **Nucleon Inversion and Three-Step Screening**, including simultaneous equation system L1–L4, target κ_N, criteria ①②③, rarity estimation, and order explanation (structure first, numerical a posteriori).
- Section 8: **Surviving Skeleton: Tripartite Y-Shaped Triangular Closure**, including structural definition, topological quantities, structural illustration, methodological lessons (Π_1 cannot distinguish opening/closing states), and screening funnel.
- Section 9: **Breaking the Parallelism: The Transition–Residue Framework**, including the correction of the "cospectral" statement, the integer closed form of Π_1, ρ as the strict invariant of the transition (the residue), the structural adjudication that the dormant edge is a ring edge, Criterion ④ (transition-reading self-consistency), Criterion ⑤ (partial re-equilibration of closure degree), Criterion ⑥ (quantitative comparison), and the two items explicitly not used as criteria.
- Section 10: **Dimension Collapse Test: Distance Data as Empirical Base for Topological Inversion**, including scale anchoring, orbital shell law, covalent radius Z_eff, bond order step law, single bond additivity law by family, and channel capacity observation.
- Section 11: **Nuclear Reaction Validation: Closure Degree Rebalancing from β Decay to Fission Chain**, including fast neutron experiment, parity criterion falsification, excitation surplus Δ criterion, B/A closure degree potential, chain criticality, and three-level signature unification.
- Section 12: **Many-Body Assembly Law and the Shared Ring: A Validation Attempt Borrowing Nuclear Data**, including the two principles of the assembly law and the adjudication between two-body variants, the shared-ring configuration and its arithmetic rigidity, "the projection erases who is dormant" (n-p ≅ p-p), the structural linear law for three and four bodies and the body-count independence of the lowest mode, the falsification at the shared-ledger pricing layer (including the advance registration of the identity), the zero-parameter existence scorecard and the adjudication of readings, the disposal of the shadow solution (variant D), the two proofs of the non-repairability of that boundary, **the attempt at an antisymmetric coefficient** (λ can neither be derived nor fixed; its demotion yields the zero-parameter criterion L1, on which the boundary is split into the two questions "existence" and "numerical binding energy"), and **the configuration-level localization of the open band** (answering "does the nucleon–nucleon force require the electronic state as well": open-band attachment is shown to be isomorphic to variant A, hence not a new structure, and a positive result is given — the open band is the carrier of principle 1 of the assembly law, appearing as the "shape of the gap" rather than as an "extra energy term"), and **the criterion construction for, and negative verdict on, the neutrino as the k = 1 singly-claimed edge** (the graph route blocked by the vertex-transitivity of the ring points, the ledger route blocked by counting; none of the 96 zero-parameter candidates is an effective hit), and **the criterion construction for, and negative verdict on, the mass-depth assignment rule (L4)** (a pre-registered three-target test: with the electron basal body the reach bound is short by 0.555 orders of magnitude, the best Ψ differs from target to target, the joint hit is 0, and the unfixed electron-layer representative constitutes a hidden degree of freedom). The accompanying figure is Figure 5.
- Section 13: **Theoretical Constraints**, including theoretical predictability of the special binding of ions/metals, current being a

coherent bookkeeping projection rather than continuous electron movement, and the **axiomatic derivation of the exponential decay law** ($T_{1/2} = \ln 2/p$ and memorylessness, MC-verified) together with the **falsification of the strong reading of its two-factor decomposition** (the zero-parameter prediction "the stripping enhancement factor is independent of the nuclide", obtained from the species-cancellation law, is refuted by measurement).

- Section 14: **Observational Perspective: the Boundary between the Dimension-Free SRE World and the Three-Dimensional World**, including the three-layer decomposition (L0 topological / L1 ratio / L2 geometric), the two tolls (the metric D and the map) and the theorem-level formalization of "fixing the shape but not the value", the two "ratio vs geometry" theorems, what is measurable on the experimental side, the unified statement of the discrete-closure law (ten examples), the three validation-qualification criteria and the 0 qualified entries of the "geometry-shape" branch, and the boundary statement given by the three-layer structure. See Figure 6.
- Section 15: **Honesty Boundary Checklist**, including the honesty boundaries of each stage: nucleon inversion, fission verification, many-body assembly, dimension collapse, theoretical constraints, and the observational perspective.
- Section 16: **Conclusion**.
- Appendices A–E provide key numerical tables, consolidated terminology list, code list and complete reproduction, figure list, references and DOI, respectively.

---

1. Terminology Convention

| Term | Meaning in This Paper | Formal Analogy | Remarks |

|---|---|---|---|

| **Binary Self-Organizing Network** | The ontology of SRE: a network with element values constantly ±1 and continuously growing over time; all things are its resident structures | A grid without pre-designed drawings, self-increasing step by step according to local rules | Model internal structure |

| **Coherent Core** | A structurally stable, repeatedly identifiable dense substructure in the network | A long-existing vortex in a turbulent river | Model internal structure |

| **Projection (Functional)** | A rule that maps a graph to a number; the graph changes, the number changes | Weighing an object — weight is a "projection" of the object | Model internal structure |

| **Graph G** | A mathematical object composed of vertices and edges | A subway map (stations = vertices, lines = edges) | Model internal structure |

| **V, E** | Number of vertices V, number of edges E | Number of stations, number of lines | Model internal structure |

| **Degree k** | Number of edges connected to each vertex; **k-regular** = all vertices have the same degree | Number of lines connected to each station | Model internal structure |

| **β_1 (First Betti Number)** | $\beta_1 = E - V + 1$, the number of independent closed loops in the graph | Number of mutually incomparable loops in a subway map | Model internal structure |

| **Laplacian Spectrum** | The set of eigenvalues of a matrix constructed from the adjacency relationship of the graph, denoted as $\lambda_1=0 \le \lambda_2 \le \ldots \le \lambda_{max}=\rho$ | The natural frequency set of a drum — different shapes have different frequency spectra | Model internal structure |

| **$\Pi_1 = \lambda_2/\rho$** | The only **calibrated** graph functional used in this paper: spectral gap ratio | The "lowest overtone/highest note" ratio of a drum | Model internal structure, calibrated as α on the electron side |

| **Automorphism Aut(G)** | A permutation that maps a graph back to itself without changing the connection relationship; $\lvert Aut \rvert$ measures the degree of symmetry | How many rotations and flips make a pattern look the same | Model internal structure |

| **Vertex Orbit** | Groups of vertices that can be interchanged under all symmetry operations | Positions of "same role" in a pattern | Model internal structure |

| **Dormant Edge** | A channel with χ=0 in SRE: not deleted, degenerated to multiplicative identity 1 — **conductive but not forming a loop** | A branch that is still energized but does not form a loop | Model internal structure |

| **Closure Degree / Opening and Closing States** | Whether the channels in the skeleton form closed loops; closed state = all closed, open state = one dormant edge not closed | A closed circuit vs a circuit with a break left | Model internal structure |

| **Graph Functional Calibration** | Using **known solutions" to determine "which functional corresponds to which real measurement" | Calibrating a scale with standard weights | Model internal structure |

| **Inversion** | Given the value of a functional, invert the structural parameters of the graph | Inferring the material and size of an object from the weighing result | Model internal structure |

| **Fission Cross-Section σ** | The probability area that a neutron (seed) incident on a nuclide causes fission; varies with energy | The success rate of a key opening a specific type of lock | Model internal structure |

| **Compound Nucleus** | The temporary state (A+1) formed when a target nucleus absorbs a neutron; fission occurs inside it | The internal state of a lock after the key is in place | Model internal structure |

| **Neutron Separation Energy S_n** | The energy required to remove the last neutron from a compound nucleus; excitation energy after absorbing a thermal neutron E"=S_n | The "engagement energy" released when the key is in place | Model internal structure |

| **Fission Barrier B_f** | The energy hill that a compound nucleus must climb to deform to fission | The resistance that the bolt must overcome to retract | Model internal structure |

| **Excitation Surplus Δ** | Δ = E" − B_f; Δ>0 means low-energy (thermal) neutrons can cross the barrier directly; Δ<0 requires additional kinetic energy from fast neutrons | The difference between engagement energy and bolt resistance | Model internal structure |

| **Multiplication Constant ν** | Average number of neutrons emitted per fission | Number of spare keys released after one lock opening | Model internal structure |

| **Critical k_eff** | Ratio of the number of neutrons in each generation relative to the previous generation; ≥1 for self-sustaining chain reaction | Whether spare keys can allow continuous lock opening without interruption | Model internal structure |

| **Closure Degree Potential** | The "rebalancing hill height" in graph theory terms: whether a complex is worth/able to rearrange to a lower closure degree configuration | The threshold at which a system decides whether to reorganize itself | Model internal structure |

| **Assembly Law** | The rule that merges several nucleon skeletons into a many-body configuration; determined by two principles: "open–closed complementarity" plus "the images of the two bodies must be distinguishable in the shared region" | Two puzzle pieces are allowed to interlock only at complementary notches, and at the joint one must be able to tell whose missing corner has been filled | Model internal structure |

| **Shared Ring** | One closed triangular ring **shared** by several nucleons after assembly | One rope loop held jointly by several people | Model internal structure |

| **Gap / Supply** | At the same shared position, the number of gaps g left by open-state (neutron) nucleons and the number of complete ring edges s offered by closed-state (proton) nucleons | The number of mortises versus tenons on a mortise-and-tenon joint | Model internal structure |

| **Shared Ledger** | A counting table recording how many bodies "claim" each overlapping edge on the shared ring; it does not change the graph structure | A partnership account that records only shares, not positions | Model internal structure |

| **Interference Depth** | The depth to which several bodies overlap on the same shared unit; used in this paper as a candidate pricing quantity for binding energy, not yet calibrated | How tightly several performers lock together when playing the same melody | Model internal structure, not calibrated |

| **Antisymmetric Penalty Term** | An additional term that depends only on the compositional imbalance a = |n_n − n_p| and vanishes identically for a ≤ 1, g(a) = λ·max(0, a − 1), used to push highly imbalanced configurations above threshold | A "disequilibrium fine" levied once the two pans of a balance differ by more than one notch | Model internal structure; its coefficient λ cannot be derived from SRE (12.12) |

| **Criterion L1 (Antisymmetry Criterion)** | A shared-ring assembly is legal (predicted bound) if and only if |n_n − n_p| ≤ 1; zero-parameter and naturally n↔p symmetric | Only recipes closest to symmetry are allowed to hold | Model internal structure; 8 / 8 on the A ≤ 4 scorecard, degrading to an empirical approximation for A ≥ 5 (12.12) |

2. SRE Theoretical Basis

2.1 Binary Self-Organizing Network and sim_p Generator

The ontology of SRE is the **binary self-organizing network**, whose core evolution algorithm `sim_p` follows the following rules:

1. **Starting Point and Growth**: Initial single-point seed $M_1=[1]$, each step $n \to n+1$ inherits the history of the top-left block read-only, and hangs new rows and columns on the frontier;
2. **Frustration Energy and Dormancy Probability**: Calculate the 2-step walking frustration energy $E_{local}=|M@M|$ and topological depth $d=n-\max(i,j)$, the dormancy probability $p=1-1/(1+\lambda \cdot d/(E_{local}+1))$, and toss a random symmetric gate χ;
3. **Dormant Edge Rule**: Dormant edges are not deleted, but degenerated to multiplicative identity 1 — **conductive but not forming a loop**;
4. **Frontier Propagation**: Non-linear multiplicative propagation of the frontier boundary $\prod$;
5. **Corner Net Load Balance Valve**: If the global sum $\Sigma \geq 0$, the new diagonal element is -1, otherwise +1.

This engine **injects new random differences and expands outward** at each step, it is a continuous generator, not a convergence terminal. All things are quasi-resident structures locked by conservation closure rules in this evolutionary flow.

2.2 Conservation Closure and Operators 7–10

Operators 1–6 are responsible for network growth and difference injection, operators 7–10 are responsible for **conservation closure**, which is the key to preventing the evolutionary flow from escaping and tearing:

- **Operator 7**: Symplectic duality/Hodge closure, strongly locking $\Delta\Pi \equiv 0$ zero-flux escape (Poincaré duality);
- **Operator 8**: Local lock-free conservation valve, strongly locking $\mathrm{Tr} \equiv 0+$ writing conflict field energy hedging against monotonic convergence;
- **Operator 9**: Two-order Betti suture, strongly locking $\Delta\beta_1 \equiv 0$, ensuring that the manifold does not tear or dimension split;
- **Operator 10**: Effective impedance Z_{eff} pre-exempts bridge edges, ensuring $\lambda_2 > 0$.

2.3 Ratio Methodology Constraints and the Only Calibrated Projection Π_1

The derivation of SRE follows the three-step method of **local input** → **topological solution** → **inversion**, and the fundamental constraint is that **the theory itself does not carry any absolute units**, only the dimensionless ratios between different physical quantities can be given. Therefore, all legal inputs for SRE derivation must be **measured dimensionless ratios**, and the output is the structural parameters of the graph.

Currently, SRE has only one numerically verified graph-physical projection, from the electron:

$$\Pi_1(G) = \frac{\lambda_2(L_G)}{\rho(L_G)}$$

where L_G is the combinatorial Laplacian of graph G, λ_2 is its smallest non-zero eigenvalue (spectral gap), and ρ is the largest eigenvalue. The calibration result on the electron side is: the Möbius parameterized point cloud of the ontology Q_3 ($V=8$, $E=12$, $\beta_1=5$, $n=60$) after calibration, $\Pi_1=\alpha\approx 1/137.036$, with a relative deviation of 1.45×10^{-5}, and the calibrated string weight $w''=1+4.347\times10^{-5}$, which matches the $\delta=4.347\times10^{-5}$ reported in the paper to five decimal places.

---

3. General Methodology for Nucleon Derivation

Nucleon derivation is homologous to electron derivation, following the three-step method of **local input → topological solution → inversion**, with the following core principles:

3.1 Definition of Local Input

"Local input" refers to injecting a small number of measured values as **rigid boundary conditions** explicitly into the model, as fixed endpoints for the solution. The injection items in this paper are fixed as (CODATA 2018):

Injected Quantity	Numerical Value	Remarks
m_e (electron rest mass)	0.510998950 MeV	Measured injection
m_p (proton rest mass)	938.27208816 MeV	Measured injection
m_n (neutron rest mass)	939.56542052 MeV	Measured injection
α^{-1} (fine structure constant reciprocal)	137.035999084	Anchor associated with Coulomb force

These injection quantities themselves **do not promise to be uniquely derived by SRE axioms**, and any quantity that cannot be injected or determined by the model topology shall not be claimed to have been calculated.

3.2 Interchange Principle Between SRE and Classical Physics (Mutual Verification)

One of the core claims of the SRE framework: under the setting of rigid boundary conditions, SRE derivation and classical physics are mutually calibrated. It is manifested as a two-way clue relationship:

- **Classical Quantity → Model**: Measured values such as m_e, m_p, m_n, α^{-1} are injected as endpoints, and the model recovers self-consistent topological structure conclusions at these endpoints;
- **Model → Classical Quantity**: Any dimensionless structural quantity that can be determined within the model (spectral ratio, flip count, orbit uniqueness, idempotent convergence) can be used as an independent structural clue for classical quantities.

The two are **mutually verified clues**, rather than one being superior to the other. This is the positional basis for all derivations in this paper.

4. Three Quark Speculations (Author's Claim)

> **This section is the author's ontological speculation, not a conclusion of existing theory, nor a derivation result of this paper.**

> The reason why this section is listed separately and placed after the methodology chapter is that it **determines what the nucleon skeleton should look like** — it is the "structural prior" for subsequent screening, and the screening results will change if it is altered. It is recorded truthfully to make readers aware: the skeleton in this paper is not "objectively emerged" from the data, but selected within the search space defined by this set of speculations.

4.1 Speculation Q1: Quarks Should Not Share the Same Substrate as Electrons

Claim: Electrons correspond to Q_3 (three-dimensional hypercube, a self-contained closed topology) in ontology. Quarks (and nucleons) should not be mapped to the same Q_3, nor to "three copies of Q_3".

Reason: Electrons and nucleons differ too much in observable behavior — electrons can exist independently, are identical, and long-lived; nucleons are bound in nuclei, have internal structure, and can decay. If they share the same substrate, these differences lack a topological source.

Adoption in This Paper: Accepted. The "3×Q_3 composite" construction used in the v1 document is invalidated. This paper searches for nucleon skeletons within the family of "non-self-contained open structures".

4.2 Speculation Q2: Quarks Are Very Likely Emergent Statistical Phenomena, Rather Than Independently Existing Ontologies

Claim: Since quarks never appear independently in experiments (color confinement), the "observation" of quarks in experiments is very likely to belong to the same category as SRE's understanding of magnetic fields and gravitational fields — **they are statistical results, not relatively independent emergent coherent cores**. In other words: quarks may not exist as ontologies at all, but only as readings.

Consistency with Established SRE Positions: This speculation is compatible with the SRE-established "observation = response spectrum of coherent core coupled with environment". From this perspective, magnetic fields, gravitational fields, "parton" readings in deep inelastic scattering, and even quantum entanglement correlations all belong to **the same category of statistical projections**, differing only in the hierarchy and degree of restriction of the coherent core being read.

Inference in This Paper: If Q2 holds, the count N_c=3 of "three quarks" in the Standard Model needs to be reinterpreted —

- N_c=3 is not "there are three kinds of quarks", but **the skeleton has three strands**;
- "Flavor" (uud / udd) is not a component list, but **the statistical projection on the detector after the opening/closing state breaking of the three-strand skeleton**;
- Fractional charge is not the property of a single body, but **the equally-divided reading when three strands share the same closed loop**.

4.3 Speculation Q3: Nuclear Reactions Are Closure Degree Rebalancing

Claim: Nuclear fission and fusion, ontologically, are not "recombination of quarks", but **rebalancing of closure degrees between closed complexes** — that is, the redistribution of closed channels, rather than the transportation of components.

Adoption in This Paper: Accepted, and as a structural requirement that the nucleon skeleton must satisfy — **the skeleton must be parameterizable by "opening/closing" degree**, and the difference between open and closed states must be small enough to explain the nuclear reaction energy in the MeV range (rather than GeV range).

4.4 Summary of Constraints on Search Space from Three Speculations

The **structural priors** used for screening below, derived from Q1–Q3:

| Number | Structural Prior | Source |

|---|---|---|

| **S1** | The nucleon skeleton is not Q_3 and its copies, nor the Möbius ladder family | Q1 |

| **S2** | The skeleton can be decomposed into **three strands** (N_c=3 reinterpreted as strand count) | Q2 inference |

| **S3** | The skeleton must support "open / closed" two states, and the difference between the two states is minimal | Q3 + β decay phenomenon |

| **S4** | The skeleton **is not self-contained**: it naturally leaves an unclosed opening structurally | Derived from "nucleons can decay" |

---

5. Entry Choice: Why Start with Neutron β Decay?

5.1 Phenomena to Be Explained

Free neutrons undergo β^- decay:

$$n \longrightarrow p + e^- + \bar{\nu}_e$$

There are three measured facts related to this, which together form hard constraints on the nucleon topology:

| Fact | Numerical Value | Requirements for Topology |

|---|---|---|

| Mass difference m_n−m_p | 1.293 MeV, i.e., $\Delta m/m_p=1.3784\times10^{-3}$ | The structural difference between proton and neutron should be **extremely small** |

| Lifetime difference | Neutron 879.4 s; proton stable (lower limit >10^{34} a) | The two are **not symmetrically equivalent**, otherwise the lifetimes should not be vastly different |

| Emitted electron | Completely identical to electrons in atoms | The emitted electron must be able to fall into the attractor of the electron ontology |

The third one is the sharpest: **β decay directly reveals that "nucleons are not closed bodies"** — a self-contained closed topology has no reason to emit an electron completely identical to an electron in an atom. This paper therefore selects β decay as the entry point for inversion.

5.2 Three Ontological Readings

Reading 1: Neutron is not "proton + electron", but the unclosed state of the same skeleton.

β^- decay is to expel that unclosed channel, and the rest close into a proton. The direction therefore falls on **closure degree**: the closed one is lighter (1.293 MeV lighter) and no longer decays; the unclosed one is heavier, and will sooner or later expel the channel. This reading also explains **why only neutrons decay** — because only neutrons carry that unclosed opening.

Reading 2: In the language of sim_p, this is the fate bifurcation of "dormant edges".

In the SRE binary self-organizing network, dormant channels ($\chi=0$) are not deleted, but degenerated to multiplicative identity 1 — **conductive but not forming a loop**. This is exactly the exact correspondence of "the channel is still there, but not closed". Thus: maintaining identity = neutron still exists; being removed = β decay. The lifetime hierarchy is also solvable: dormancy is the default state, removal is a rare perturbation, so the neutron lifetime (10^3 s) is 26 orders of magnitude longer than the strong interaction timescale (10^{-23} s).

Reading 3: This is sedimentation at depth order.

Electrons are deep and tight residents (high coherence); nucleons are shallow and loose residents. β decay is shallow structure sedimenting into deep structure, with energy paid by the resident depth difference — consistent with the registration in v1 that "mass is inversely proportional to logical depth" (nucleons are about 1836 times shallower, so about 1836 times heavier).

An Additional Gain: If electrons are the **only deepest attractor** of the generator, then the identity of electrons (electrons everywhere in the universe are completely indistinguishable) does not need to be guaranteed by "taking out the same electron from somewhere", but "any seed that breaks away from the skeleton falls into the same attractor". The emitted electron in β decay is completely identical to the electron in an atom, and the **mechanism** is thus obtained, rather than an assumption.

5.3 Structural Constraints Derived from β Decay

| Number | Constraint | Argument |

|---|---|---|

| **T1** | The nucleon skeleton must be **non-self-contained** | Three-strand skeleton has 9 leaf points (odd number), which cannot be paired and closed pairwise, naturally leaving one opening. This is the graph-theoretic root of "nucleons can decay" |

| **T2** | Protons and neutrons are **not Z_2 images of graph isomorphism**, but two "awake / sleepy" configurations of the same underlying matrix, with their coherent subgraph loop number β_1 differing by 1 | If they are exactly isomorphic, there is no direction for decay and the lifetimes should not be vastly different |

| **T3** | What is removed only needs to be a **seed**, and it is not required that the skeleton contains a complete electron ontology internally | Electrons are identical attractors, not prefabricated components of nucleons |

6. Redefinition of Isospin Z_2 and Breaking Magnitude

6.1 Essence of the Conflict

The C1 assertion in v1 document is:

> **C1**: Isospin Z_2 doublet — fixed-point-free involution, charge p=+1/n=0, proton and neutron graph isomorphism.

Under the speculation that "quarks are very likely emergent statistical phenomena" (Q2), this assertion **not only needs to be patched, but needs to be invalidated**, for the reason that: C1 is built on the flavor multiset flip {uud}↔{udd}, which is equivalent to treating statistical readings as ontological components — which directly conflicts with Q2.

6.2 Redefinition of Z_2

| Item | v1 (Invalidated) | This Version v2 |

|---|---|---|

| Meaning of Z_2 | Flavor multiset flip {uud}↔{udd} | **Opening / closing (awake / sleepy) duality** of the skeleton, independent of flavor |

| Two-state relationship | Exact graph isomorphism | **Non-isomorphic**: β_1 differs by 1 |

| Symmetry | Exact (fixed-point-free involution) | **Naturally broken**, the breaking amount is the closure energy |

| Correspondence with observation | Flavor = ontological components | Flavor = statistical projection of this breaking on the detector |

After redefinition, a phenomenon that originally required additional assumptions to explain becomes self-evident:

6.3 Why Is the Breaking Magnitude 10^{-3}: Structural Necessity

Dormant edges in SRE **still conduct** (degenerate to identity 1, rather than zeroing or deleting). Therefore, the difference between open and closed states is **not "whether there is a bond", but "whether there is a loop"". From this:

- The energy difference only comes from "the absence of a loop", not "the absence of a channel";
- Therefore $\Delta m/m_p \sim 10^{-3}$ is **a structural necessity**, not a parameter coincidence.

This also explains two other questions at the same time: **why the mass difference is so small** (only one loop difference), and **why neutrons can still exist for so long** (dormancy is the default state).

6.4 Homotypic Comparison with Electron Side Breaking

| System | Breaking Magnitude | Property |

|---|---|---|

| Electron (Möbius twist) | w"−1=4.347×10⁻⁵ | Near-symmetric breaking |

| Nucleon (opening / closing) | Δm/m_p=1.378×10⁻³ | Near-symmetric breaking |

| Ratio | 31.7 | Same type, different magnitude |

Both belong to **the same type of breaking** (approximate establishment of symmetry + small amount of breaking), only the breaking amplitude on the nucleon side is about 32 times that on the electron side. This homotypicity supports the self-consistency of the redefined Z_2 in this version, but does not constitute independent evidence.

6.5 Connection with the Charge-Emergence Paper: Isospin Z_2 ≡ Charge-Magnitude Z_2 (v1.6 revision)

The paper *Emergence of Positive and Negative Electric Charge in the SRE Framework: Magnitude, Sign, and Numerical Verification* gives the charge functional $Q = \sigma \cdot \deg \cdot e$ (with $\deg = \beta_1 \bmod 2$). Its §13 attaches $\deg$ to the charge **magnitude** and registers a divergence with this section (which had stated "the opening/closing isospin Z_2 is independent of charge"), pending the author's ruling. The ruling is now given as follows (see Proposition 6.1 of *Topological Derivation of the Nucleon Charges from the Charge-Emergence Functional* $Q = \sigma \cdot \deg \cdot e$):

- The "opening/closing" two states of this section are precisely the skeleton carrier of $\deg = \beta_1 \bmod 2$: the closed state $(12, 18, 7) \Rightarrow \beta_1 = 7 \Rightarrow \deg = 1 \Rightarrow$ charged; the open state $(12, 17, 6) \Rightarrow \beta_1 = 6 \Rightarrow \deg = 0 \Rightarrow$ neutral.
- Therefore "isospin Z_2" and "charge-magnitude Z_2" **are two readings of the same Z_2**: the open/closed difference ≡ a difference of one independent cycle ≡ parity flip ≡ charged/neutral. The two do not conflict; rather, this section's isospin Z_2 is **reinterpreted as** the charge-magnitude Z_2.
- The breaking magnitude 10^{-3} of §6.3 (from "the absence of a cycle") shares the same origin as the charge magnitude $\deg$ — the breaking falls in the **mass layer** (dynamics), not in the **charge-magnitude layer** (topology).
- Conclusion: the original statement of this section, "isospin Z_2 is independent of charge," must be upgraded to "isospin Z_2 ≡ charge-magnitude Z_2"; the charge-emergence paper's "Reading A" thereby becomes the **natural upgrade** of this section, and the divergence is closed.

> Note: The absolute sign still belongs to the A0 convention of the charge-emergence paper (the structural opposition relative to the electron is fixed, but absolute positive/negative cannot be derived from SRE). The carrier and relative structure on the sign side have been closed by §4 of *Topological Derivation of the Nucleon Charges from the Charge-Emergence Functional* $Q = \sigma \cdot \deg \cdot e$.

---

7. Nucleon Inversion and Three-Step Screening

7.1 Simultaneous Equation System for Nucleons

Moving the same set of dictionaries to the nucleon layer, let the nucleon skeleton be graph G_N, and its unknown topological parameters be (V,E,k,w), then the equation system is:

| Hierarchy | Equation | Property |

|---|---|---|

| **L1 Count Layer** | $\beta_1 = E - V + 1$; $n = E \cdot \beta_1$; $kV = 2E$; $V \equiv 0 \pmod 2$ | Integer (Diophantine) equation, the strongest constraint |

| **L2 Spectral Equation** | $\Pi_1(G_N) = \kappa N$ | *Need to identify* κN corresponding to the measured ratio |

| **L3 Torsional Near Symmetry** | $w_N = 1+\delta N$, $\lvert\delta N\rvert \ll 1$ | **Inherited from the electron assumption, not an independent constraint** |

| **L4 Mass-Depth Equation** | $m_p/m_e = \Psi(G_N)/\Psi(G_e)$ | Ψ has not been defined, **currently unavailable** |

Degree of freedom accounting: 4 independent unknowns, 3 equations from L1 + 1 equation from L2 = 4 equations, **formally closed**; but since κ_N has not been identified and L3 is not independent, **the actual underdetermination is 1 degree of freedom**.

7.2 Proposal of Target Formula (Explicitly Marked as Hypothesis to Be Tested)

This paper proposes the following **candidate** relationship (not yet independently demonstrated):

$$\boxed{\frac{m_n - m_p}{m_p} = \alpha \cdot \Pi_1(G_N)}$$

If it holds, its physical meaning is: **the isospin breaking of nucleons has an electromagnetic origin** (α appears explicitly), and Π_1 is a pure topological structure factor.

Substituting the measured values:

$$\kappa_N = \frac{(m_n - m_p)/m_p}{\alpha} = \frac{1.378419305 \times 10^{-3}}{7.297352569 \times 10^{-3}} = 0.188893067$$

This identification is a hypothesis, not a conclusion. The three-step screening below tests "in the family of V=12 cubic graphs, how many graphs have Π_1 near κ_N", and "how rare the screening result is".

7.3 Criterion ① Integer Determination of Vertex Number V

For k-regular graphs, $kV=2E$, $\beta_1=E-V+1$. Taking k=3 (cubic graphs), then $\beta_1=V/2+1$. Check the match between $\beta_1 \cdot \Delta m/m_p$ and the measured "flow quark mass ratio" $(2m_u+m_d)/m_p=0.9581\%$:

| V | β_1 | $\beta_1 \cdot \Delta m/m_p$ | Deviation from 0.9581% |

|---|---|---|---|

| 8 | 5 | 0.6892% | −28.07% |

| 10 | 6 | 0.8271% | −13.68% |

| **12** | **7** | **0.9649%** | **+0.70%** |

| 14 | 8 | 1.1027% | +15.09% |

| 16 | 9 | 1.2406% | +29.48% |

| 18 | 10 | 1.3784% | +43.86% |

V=12 (β₁=7) is uniquely selected, with adjacent deviations above ±14%, and the uncertainty of the flow quark mass ±2% cannot shake this choice.

> **Qualifier (added in v1.6).** The anchor $(2m_u + m_d)/m_p$ used above is an **MS-bar scheme quantity** (it depends on the renormalization scale) and is not even listed among the declared injections of §3.1. Measurement (`code/_sre_anchor_registry.py`) shows that moving the scale from 2 GeV to 1 GeV changes that ratio by about **+23%**, far beyond the bin-switching threshold of the table (±6.5–7.9%), and **measurably moves the V selected by Criterion ① from 12 to 16** (3 GeV → V=10, 1.5 GeV → V=14, 1 GeV → V=16). The correct statement is therefore "**under the 2 GeV MS-bar convention**, Criterion ① is compatible with V=12", and not "Criterion ① determines V=12". The final fixation of V must be made jointly with the integer equations of L1 (see the new item 11 of the nucleon-inversion boundaries in Section 15).

7.4 Criterion ② Spectral Ratio Determination of Topology

Enumerate cubic connected graphs with V=12 (the total number of non-isomorphic graphs at this order is 85, and all 85 are obtained by sampling and deduplication this time), calculate Π₁=λ₂/ρ for each graph:

| Statistic | Value |

|---|---|

| Number of samples | 85 |

| Minimum / median / maximum of Π₁ | 0.0313 / 0.1319 / 0.2768 |

| Within 0.17% of κ_N | 0 |

| Within 0.5% | 3 |

| Within 2% | 4 |

| Within 3% | 4 |

That is: **the Π₁ landing point itself is already a strong screening (85 → 4)**, because the spectral ratio distribution of this family is concentrated around 0.13, and the target value 0.1889 is outside the upper quartile of the distribution.

7.5 Criterion ③ Triple Symmetry Determination of Solution

From the structural prior **S2** (the skeleton can be decomposed into three strands), it is required that the automorphism order of the graph be divisible by 3, i.e., 3∣|Aut(G)|. Take the 4 candidates surviving from Criterion ② and calculate their automorphism orders one by one:

| Π₁ | Deviation from κ_N | |Aut| | Vertex Orbit | Number of Triangular Loops | Judgment |

|---|---|---|---|---|---|

| 0.188580 | **0.165%** | 36 | 2 | 3 | **Keep** |

| 0.188580 | **0.165%** | 12 | 3 | 1 | **Keep** |

| 0.188011 | 0.467% | 2 | 6 | 0 | Reject |

| 0.192106 | 1.701% | 8 | 3 | 0 | Reject |

85 → 4 → 2.

7.6 Rarity Estimation

- Joint hit rate of Criterion ②③: 2/85 = 0.0235
- Multiply by the V bin selection from Criterion ① (5 bins select 1): p ≈ 4.7×10⁻³

The meaning of this number is: **if the nucleon skeleton is randomly drawn from this family of graphs, the probability of satisfying all three criteria is less than 0.5%." But it is not a statistical significance test — because the three criteria are not independent (both Criterion ① and Criterion ② use Δm), and the identification of κ_N itself is a hypothesis.

> **Qualifier (added in v1.6; narrowed and withdrawn in v1.7).** The "5 bins select 1" factor above presumes V=12 to be the only admissible bin. In v1.6 it was accordingly registered that "V=12 is not unique": κ*N* = *0.188893 lies inside the families of* ***V = 8, 10, 12 and 14 simultaneously*** *(only the V=16 family falls short, its Π₁ upper bound being 0.1780 <* κN), and after Criteria ②③ are applied **both V=10 and V=12 retain survivors**. **v1.7 withdraws that qualifier**, the reason being that the window of Criterion ② was **not self-consistent** (`code/_sre_anchor_second.py`, PART B): Criterion ② originally took |Π₁ − κ*N*|/κN < 3%, whereas the agreement reached by the α-lock itself is only **0.1655%** — κ_N is computed from two high-precision measured numbers (Δm_np/m_p and α, with measurement errors ~10⁻⁹), so **every residual deviation is a theoretical-agreement error**; a 3% window is 18 times wider than the agreement already in hand, which amounts to voluntarily discarding the discriminating power. Tightening the window to the order of the anchor's own precision gives the following measurement (each cell = "Π₁ hits / of which survivors with 3Ⅱ|Aut|"):

ε (window)	V=8 (5/5 full)	V=10 (19/19 full)	V=12 (85/85 full)	V=14 (140/509 sampled)	V=16 (70/4060 sampled)
0.2%	0 / 0	0 / 0	**2 / 2**	1 / 0	0 / 0
0.5%	0 / 0	0 / 0	**3 / 2**	1 / 0	0 / 0
1.0%	0 / 0	0 / 0	**3 / 2**	1 / 0	0 / 0
3.0%	0 / 0	2 / 1	**4 / 2**	3 / 0	0 / 0

⇒ **For ε ≤ 1% together with Criterion ③, only the V=12 bin retains survivors**. V=8/10 are **full sets**, so their zeros are exact; V=14/16 are sampled, so all that can be said is "zero among 140/70 samples", equivalent to p ≲ 0.71% / 1.43% (for V=12 the hit rate is p = 3/85 = 3.5%), hence **LR ≥ 1.7 (against V=14) and ≥ 0.83 (against V=16, weak)** — "V=12 is unique" is therefore **strong evidence but not a proof**. Accordingly, "settling on V=12" no longer rests on Criterion ①: **once the window is tied to the anchor's precision, Criteria ②③ by themselves narrow V from {8,10,12,14} to {12}**. The sole premise of that narrowing is the agreement "κ_N ≈ Π₁(Y₃) to within 0.165%" itself.

> **Correction (v1.7 final, self-check of 2026-09-24).** The v1.7 draft stated that "that survivor is, on individual identification, precisely the Y_3 skeleton itself". **This does not hold and is hereby corrected.** After **exhaustively** enumerating all 85 isomorphism classes of connected cubic graphs on 12 vertices by a 2-switch closure BFS (`code/_enum_cubic12.py`; 85 matches the known count, and the 85 isomorphism classes correspond to **85 distinct Laplacian spectra**, i.e. that bin has **no cospectral pair**), the **complete** list of hits within $\varepsilon \le 1\%$ is three graphs:

#	Π_1	deviation from κ_N	$\lvert Aut \rvert$	T	Y_3 skeleton?
0	0.188580484696	0.1655%	36	3	**yes**
27	0.188011313654	0.4668%	2	0	no (3∤2, excluded by Criterion ③)
81	0.188580484696	0.1655%	12	1	no (12 = 3×4, Criterion ③ **fails to** exclude it)

⇒ Criterion ③ ($3 \nmid |Aut|$) is **not unique** within the V=12 bin: Y_3 and #81 are both survivors. **The root cause is that Π_1 is a ratio quantity** — #81 has exactly the same Π_1 as Y_3 (both $(7-\sqrt{13})/18$), so **tightening ε arbitrarily cannot separate them** (even at $\varepsilon = 0.2\%$ there are still 2 survivors); this has nothing to do with "identical Laplacian spectra" (#81 and Y_3 are not cospectral) — the ratio λ_2/λ_max simply coincides. Therefore **no criterion that reads only Π_1 can ever pin down the graph**, and a non-spectral criterion must be superimposed. Adding **$T \ge 2$** (the same-type criterion as Criterion ⑤ in §4, "the open state must retain a closed unit") reduces the survivors within V=12 to **1, and it is precisely the Y_3 skeleton** (T=3, $|Aut|=36$, $\rho=5.302775638$, $\lambda_2=1.000000000$).

> **Engineering correction.** The graph pool used in the v1.7 draft was sampled by a random generator (V=12 yielded **84/85**, missing Y_3 itself by chance), so the "2 hits / 1 survivor" of PART B was **an artefact of the omission**; the main script `code/_sre_anchor_second.py` now **explicitly includes Y_3** and rebuilds the full pool (85/85). The numbers in the table above are the rerun results of this round.

The decomposition of p changes accordingly: the former "5 bins select 1" factor is now carried by the sharpened Criteria ②③, whose likelihood ratio is only a **lower bound** (LR ≥ 1.7, limited by the V=14/16 sampling); hence $p \approx (2/85)/\mathrm{LR} \approx$ **1.4×10^{-2} to 2.8×10^{-2}**, about 3–6 times looser than the previous estimate of 4.7×10^{-3}; if V=14/16 are later enumerated exhaustively and still yield zero, p returns to the 4.7×10^{-3} order.

7.7 Order Explanation: Structural Intuition First, Numerical A Posteriori

It is necessary to specially point out the **temporal order** of screening, because it relates to the credibility of the results:

> The candidate with the smallest deviation ($|Aut|=36$, 3 triangular loops, 2 vertex categories) was hand-constructed by the author based on the structural intuition of "three-strand Y skeleton + odd leaves cannot be paired pairwise" **before seeing any spectral ratio numerical values**; the spectral ratio coincidence (0.165%) is a **post-hoc verification result**, not a fitting target.

If it were obtained by fitting, the "three coincidences hitting simultaneously" would be circular reasoning; but if structure comes first and numerical values come after, this coincidence at least has **non-triviality**.

---

8. Surviving Skeleton: Tripartite Y-Shaped Triangular Closure

8.1 Structural Definition

Let the candidate skeleton be $Y_3 \ltimes \triangle_3$, constructed as follows:

1. Take 3 **strand centers** c_0, c_1, c_2, not connected to each other;
2. Take 3 **triangular loops** T_0, T_1, T_2, each loop containing 3 leaf points;
3. Each strand center is connected to exactly one leaf point of each triangular loop (total 9 spokes, forming a Latin square perfect matching);
4. 3×3=9 intra-ring edges, 9 spokes, so E=18; V=3+9=12; each vertex has degree 3.

Topological quantities (recalculation result):

Quantity	Closed State (Proton Candidate)	Open State (Neutron Candidate, one ring edge broken)
V / E	12 / 18	12 / 17
β_1 (Independent Loop Number)	**7**	**6**
Number of Triangular Loops	3	2
Girth / Diameter	3 / 3	3 / 3
$\lvert Aut \rvert$	**36**	4
Vertex Orbit	**2** (strand center / leaf point)	6
λ_2 / ρ	0.188580	0.188580

$|Aut|=36=3!\times3!$, exactly the permutation symmetry order of "3 strand centers × 3 triangular loops"; there are 2 vertex orbit types, i.e., only two roles in the structure — strand center and leaf point. These two highly match the structural prior **S2** (three strands).

Edge-class attribution of the dormant edge. The 18 edges split naturally into two automorphism orbits of 9 edges each: **ring edges** (leaf point—leaf point) and **spokes** (strand center—leaf point). By structural adjudication, **the dormant edge is taken to be a ring edge**. The reason: the dormant edge is itself the image of a homomorphic mapping and its logical depth is large — opening a ring edge **changes no spectral quantity at all** (both ρ and λ_2 remain strictly invariant), and alters only the number of closed units (3→2). In other words, dormancy is spectrally invisible; it acts on the ontological level of closed structure. The open-state definition is fixed accordingly and is consistent with Figure 1.

Reading comparison after opening one edge of each class (derivation in Section 9.4):

Edge Class	λ_2 after opening	ρ after opening	Number of Closed Units T after opening
Ring edge (dormant edge)	1.000000 (invariant)	5.302776 (invariant)	**3** → **2**
Spoke	1 → 0.527166	5.302776 (invariant)	3 (invariant)

8.2 Structural Illustration




Figure 1: Left is the closed state (proton candidate, all three ring edges closed), right is the open state (neutron candidate, one **ring edge** dormant and not closed, red dashed line). Orange is the strand centers (3), blue is the leaf points on the ring (9), and blue solid lines are closed channels. The only difference between open and closed states is "one fewer closed unit" (number of triangular loops 3→2), which is exactly the structural basis for the breaking magnitude argument in Section 6.3.

8.3 An Important Methodological Lesson

The Π_1 values of the open and closed states are **completely identical** (both 0.188580). This indicates that:

> **The functional Π_1 cannot distinguish between proton and neutron.** The opening/closing state should be read by β_1 (independent loop number), the number of closed units T, or the number of dormant edges, not by the spectral gap ratio.

Promoted to a general lesson: **the choice of functional must match the nature of the physical quantity to be distinguished.** A functional that is insensitive to "loop addition/removal" cannot be used to diagnose the phenomenon corresponding to "loop addition/removal". This is a hard constraint for the next step of functional calibration (Section 10).

8.4 Screening Funnel




Figure 2: A total of 85 cubic connected graphs with V=12 (determined by Criterion ①) → 4 remaining when the spectral ratio falls within 3% of the target (Criterion ②) → 2 remaining when satisfying triple symmetry (Criterion ③). Combined rarity p≈4.7×10⁻³.

---

9. Breaking the Parallelism: The Transition–Residue Framework

9.1 A Correction: A and B Are Not Cospectral

An earlier draft described the two candidates as "cospectral and non-isomorphic"; that statement was wrong. The two are **not cospectral**:

Spectrum (combinatorial Laplacian: value × multiplicity)	Candidate A	Candidate B
Full spectrum	0^1, $\mathbf{1^2}$, $((7-\sqrt{13})/2)^2$, $\mathbf{4^5}$, $((7+\sqrt{13})/2)^2$	0^1, $\mathbf{1^1}$, $((7-\sqrt{13})/2)^2$, $\mathbf{2^2, 4^3, 5^1}$, $((7+\sqrt{13})/2)^2$
Number of distinct eigenvalues	5	7
$\lambda_2 = \lambda_3$?	Yes (1, 1)	No (1, 1.697224)

What they share is only three **extreme spectral quantities**: $\lambda_1 = 0$, $\lambda_2 = 1$, $\rho = (7+\sqrt{13})/2 = 5.302776$. And $\Pi_1 = \lambda_2/\rho$ happens to be the ratio of two of them — this is the true reason why Π_1 cannot separate A from B: **it is blind to the internal structure of the spectrum**.

The remaining invariants do differ substantially:

| Invariant | Candidate A | Candidate B |

|---|---|---|

| V / E / β_1 | 12 / 18 / 7 | 12 / 18 / 7 |

| λ_2 / ρ / Π_1 | 1.000 / 5.303 / 0.188580 | Same |

| Number of triangular loops T | **3** | **1** |

| Vertex orbits | **2** | **3** |

| \|Aut\| | **36** | **12** |

| Edge orbits and sizes | **2** (9, 9) | **4** (3, 3, 6, 6) |

| Kirchhoff index | 57.67 | 54.07 |

| Wiener index | 135 | 129 |

| Average resistance distance | 0.8737 | 0.8192 |

9.2 A Closed Form: Π_1 Is Determined by Three Integers

$$\Pi_1 = \frac{1}{\rho} = \frac{\beta_1 - \sqrt{V+1}}{E} = \frac{7-\sqrt{13}}{18} = 0.188580485$$

Against the measured κ_N = Δm/m_p ÷ α = 0.188893067, the deviation is 0.165%. That is, the nucleon's Π_1 is **entirely determined by the three integers β_1, V, E**. Both A and B belong to this "√13 subfamily"; the other two graphs inside the Π_1 window are generic (12 distinct eigenvalues) and match only approximately.

9.3 The Strict Invariant of the Transition: ρ Is the "Residue"

For each of A and B, all 18 single-edge openings are performed and the range of each spectral quantity recorded:

| Spectral quantity | Range for A (18 removals) | Range for B (18 removals) |

|---|---|---|

| **ρ (largest Laplacian eigenvalue)** | **3.6×10^{-15}** | **6.2×10^{-15}** |

| λ_2 | 0.473 | 0.473 |

ρ is **strictly invariant** under the transition (the difference is at floating-point noise level), whereas λ_2 is strongly mobile. Physical reading: **the part of the spectrum that stays unchanged through the transition is the "residue", and it naturally serves as the reference object of measurement (the denominator of the ratio)**; the numerator is the branch that moves during the transition. This is aligned with the ontological reading that "light is the topological residue between evolutions" — **the standard is the residue, not the object being measured**.

9.4 The Dormant Edge Is Taken to Be a Ring Edge (Structural Adjudication)

Basis (structural adjudication): the dormant edge is itself the image of a homomorphic mapping, and its logical depth is large. For this skeleton, the candidate with the greatest depth is the **ring edge** — opening a ring edge leaves every spectral quantity unchanged (both ρ and λ_2 strictly invariant) and alters only the number of closed units. In other words, dormancy is spectrally "invisible"; it acts on the ontological level (closure degree) rather than on the observable spectral level. This is precisely the

manifestation of "large logical depth".

The 18 edges of A split naturally into two automorphism orbits of 9 each:

Edge class	λ_2 after opening	ρ after opening	Closed units T after opening	Laplacian trace
Ring edge (leaf—leaf; the dormant edge)	**1.000000 (invariant)**	5.302776 (invariant)	**3 → 2**	36 → 34
Spoke (center—leaf)	1 → 0.527166	5.302776 (invariant)	3 (invariant)	36 → 34

The open-state definition is fixed accordingly and is consistent with Section 8.1 and Figure 1: **the neutron state = opening one ring edge**. Here both ρ and λ_2 are unchanged; the only changes are the number of closed units (3→2) and the trace (−2).

9.5 Criterion ④ Transition Self-Consistency: Equivalent Channels Must Give Equivalent Readings

A dormant state is not the state of "some particular edge" but the **homomorphic image of an entire edge orbit**: the edge set is quotiented to the orbit set under the automorphism group, and this step is exactly a homomorphism. The transition reading must therefore be unique at the **orbit level**.

- Candidate A: the 9 ring edges form one orbit, and opening any one of them gives (λ_2 = 1.000000, T = 2), a unique reading ✓
- Candidate B: the 3 ring edges form one orbit, and the reading is likewise unique (λ_2 = 1.000000, T = 0) ✓

Honest registration: over the whole pool, "unique ring-edge reading" holds for 28/85; combined with Π_1<3% it leaves **2** graphs (A and B). **Criterion ④ alone is not sufficient to break the tie**, hence it is used jointly with Criterion ⑤.

9.6 Criterion ⑤ Partial Re-equilibration of Closure Degree: The Open State Must Still Retain Closed Units

Physical basis (**independent of the "three strands" prior S2**): a nuclear reaction is a **re-equilibration** of closure degree (Section 4.3, conjecture Q3). Re-equilibration means **partial** release, not reducing the closure degree to zero; a neutron is still a bound composite, so its open state must still retain closed units. More importantly, Q3 is independently supported by the fission chain validation of Section 11 (Pearson correlation +0.932 between the closure-degree potential Δ and the thermal cross-section), so using Q3 as a criterion here is not circular.

- Candidate A: T: 3 → 2 (minus one, still retaining 2 closed units) ✓
- Candidate B: T: 1 → 0 (**reduced to zero**) ✗ — if B were read as the nucleon, a single re-equilibration would erase the entire closure degree, contradicting the very mechanism of "re-equilibration".

Pool statistics (85 graphs, criteria before values):

| Criterion expression | Hits |

|---|---|

| Unique ring-edge reading (at the T level) | 28 / 85 |

| T_closed ≥ 2 and unique ring-edge reading | 11 / 85 |

| [$\Pi_1 < 3\%$] ∧ [unique ring-edge reading] | 2 / 85 |

| **[$\Pi_1 < 3\%$] ∧ [unique ring-edge reading] ∧ [T_closed ≥ 2]** | **1 / 85 = A** |

| **[$\Pi_1 < 3\%$] ∧ [open state T_open ≥ 1]** | **1 / 85 = A** |

Criterion ⑤ requires only that "re-equilibration is partial", and is **unrelated** to Criterion ③ (3Ⅱ|Aut|); hence it does not suffer from the charge of "same origin", nor does it require choosing a direction within a numerical window.

9.7 Criterion ⑥ Quantitative Comparison (Corroboration Only)

Comparison of the closure-degree ratio (open/closed) with the magnetic ratio:

| Candidate | T: closed → open | Ratio | Against |μ_n/μ_p| = 0.684979 |

|---|---|---|---|

| **A** | 3 → 2 | **2/3 = 0.666667** | deviation **2.67%** |

| B | 1 → 0 | 0 | no correspondence |

Honest boundary: with 36 comparisons and a 5% threshold, the expected accidental hit count is about 1.8, so this item is registered only as a **clue**, not as evidence.

9.8 Two Items Explicitly Not Used as Criteria

The following two items were examined and **are not adopted**:

1. **"Some mode releases exactly the integer 1.000000"** — observed as closed state 4.000000 → open state 3.000000, i.e., "exactly one mode releases the integer 1". This arises only from the contingency of the homomorphic mapping and is **not a necessary discrete quantity**; it does not constitute a criterion;
2. **"The total released amount is always 2"** — removing one edge necessarily decreases the Laplacian trace by 2; this is a trace identity carrying no new information.

Consequently, the per-mode allocation observed in the spoke-class opening (1.000000 / 0.472834 / 0.367343 / 0.159823, with the remaining 8 eigenvalues unchanged) **is not used as a criterion** and is retained only as an explanatory description of the transition mechanism.

9.9 Conclusion

At the structural level: Criterion ⑤ uniquely selects A, and B is excluded; Criterion ④ corroborates.

At the measured level: still not independent. The only quantitative comparison independent of the structural prior (Criterion ⑥, deviation 2.67%) is not strong enough.

The correct statement is therefore: **"Under the condition of partial re-equilibration of closure degree, A is the unique survivor; at the purely measured level it is still not independently determined."** — not "A has been determined by measurement".

It should also be pointed out that, with the dormant edge taken as a ring edge, the transition has **no effect on either λ_2 or ρ**; hence the Π_1 in $\Delta m/m_p = \alpha \cdot \Pi_1$ is a "standard" built from two **invariants**, while the state change itself is carried by the number of closed units T. This reading is consistent with "the residue serves as the reference object of measurement", but it implies that the direct reading "mass difference ↔ level shift" does not apply here, and it has been registered as unresolved item 5 of Section 16.

---

10. Dimension Collapse Test: Distance Data as Empirical Base for Topological Inversion

To provide a testable "distance ladder" baseline for the structural solution of nucleon topology, a dimension collapse test is performed. This test is the **empirical base** of nucleon derivation: ② bonding rules and ③ skeleton enumeration are the same thing here — **skeleton channel structure ↔ distance ladder**.

10.1 Scale Anchoring (Dimensionless Identity)

$$a_0 \cdot m_e \cdot c/\hbar = 1/\alpha \approx 137.036$$

Measured verification: the ratio is 1.0000000006, almost exactly 1.

10.2 Orbital Shell Law $r_n = n^2 \cdot a0/Z$ (in a0 units)

Z	n=1	n=2	n=3	n=4	Integer Ratio
1	1.000	4.000	9.000	16.000	1 : 4 : 9 : 16
2	0.500	2.000	4.500	8.000	1 : 4 : 9 : 16
3	0.333	1.333	3.000	5.333	1 : 4 : 9 : 16

10.3 Covalent Radius (in a0 units) and Derived Z_eff (Leakage Profile, Main Group Extension)

$Z_{eff} \approx n^2/(r/a0)$ (n takes the main shell quantum number): increasing gradually along the period for the second/third periods = increasing nuclear leakage.

Element	Z	r/Å	r/a0	Derived Z_eff	Main Shell n
Li	3	1.28	2.419	1.65	2
Be	4	0.96	1.814	2.20	2
C	6	0.76	1.436	2.79	2
N	7	0.71	1.342	2.98	2
O	8	0.66	1.247	3.21	2
F	9	0.57	1.077	3.71	2
Na	11	1.66	3.137	2.87	3
Si	14	1.11	2.098	4.29	3
Cl	17	1.02	1.928	4.67	3
K	19	2.03	3.836	4.17	4
Rb	37	2.20	4.157	6.01	5

Z_eff increases monotonically across periods in the same group, such as group 14 C/Si/Ge/Sn: 2.79→4.29→7.06→9.52.

10.4 Bond Order Step Law (Contraction Δ for each additional bond, in a0 units)

Family	Bond Order Series	Δ(1→2)	Δ(2→3)	Remarks
C–C	single / double / triple	0.378	0.265	
C–N	single / double / triple	0.359	0.227	
C–O	single / double / triple	0.378	0.189	
N–N	single / double / triple	0.378	0.287	
Si–Si	single / double / triple	0.340	0.113	Third period heavy atom triple bond weakened

Core multiple bond families (excluding O–O, N–O lone pair families, n=11): Δ(1→2)=0.352±0.037 a0 (RMS 0.354), Δ(2→3)=0.232±0.058 a0 (RMS 0.239).

10.5 Single Bond Additivity Law d(A–B)≈r_A+r_B (in a0 units, statistical by family)

Family	n	Mean Value	RMS	Judgment
Organic C Skeleton Neutral Covalent Family	11	+0.6%	1.5%	Strictly holds
Light Heteroatom/Hydride	25	−2.9%	6.7%	Approximate (electronegativity shift)
Same-Nucleus Second Period (H/O/F)	5	+10.0%	12.5%	Systematically stretched → channel saturation correction
Same-Nucleus Heavy Main Group (Si…I)	10	−0.9%	3.3%	Additivity law regression
Ion/Metal Pairs	6	−12.3%	14.4%	Failure (different ontological interval)

10.6 Bond Order Capacity Observation: Maximum Stable Bond Order for Each Family

- Group 14: C≡C(3), Si≡Si(3), Ge≡Ge(3) — Capacity 3
- Group 15: N≡N(3), P≡P(3), As≡As(3) — Capacity 3
- Group 16: O=O(2), S=S(2), Se=Se(2), Te=Te(2) — Capacity 2
- Group 17: F–F(1), Cl–Cl(1), Br–Br(1), I–I(1) — Capacity 1

The channel capacity is not always 3, but decreases with the group and period; the same-nucleus small atoms in the second period (O, F) often have actual bond orders lower than the capacity upper limit due to lone pair repulsion, which is consistent with the two corrections of "channel saturation + lone pair repulsion" in the bonding rules.

---

11. Nuclear Reaction Validation: Closure Degree Rebalancing from β Decay to Fission Chain

This paper does not propose a new nuclear structure theory, but **uses existing measurements of fission and chain reactions to verify whether a mechanistic assertion that has been accepted in nucleon derivation can continue to hold on a larger scale**. This assertion is:

> **Nuclear reaction = closure degree rebalancing.** (Chapter 3 of Nucleon Inversion v2, Speculation Q3)

11.1 Fast Neutron Experiment: Differences Only Exist in Low-Energy Sedimentation

Criterion Design (Criterion Precedes Numerical Values). If the reading of Q3 holds, then:

- **Slow (thermal) neutrons**: can only settle at low energy at "existing unpaired open channels" in the composite skeleton, and have nowhere to land for "fully paired closed" skeletons → the cross-section should span a very wide range;
- **Fast neutrons**: with enough energy to directly destroy the closure/pairing, bypassing the low-energy sedimentation channel → the cross-section should **restore to the same order of magnitude** for all nuclides.

Results:

Quantity	Numerical Value
Thermal cross-section dynamic range	**8.58 orders of magnitude** (max 6401 b / min 1.7e-5 b)
Thermal cross-section 1–99 percentile span	8.39 orders of magnitude

| Fast cross-section dynamic range | **1.48 orders of magnitude** (max 2.43 / min 0.080 b) |

| Fast cross-section (excluding Th-232) | 0.90 orders of magnitude |

| Fast cross-section 10–90 percentile span | **0.40 orders of magnitude** |

Interpretation. Fast neutrons compress the cross-section to the same order of magnitude, and the differences are indeed concentrated at the low-energy/slow neutron sedimentation end. This is consistent with the mechanism that "dormant edges require low-energy removal, and high energy can directly destroy closure", and it is a **10^4-fold structural signature** — not a small numerical coincidence.

11.2 An Honest Falsification: Simple Parity Criterion Does Not Hold and Needs to Be Upgraded

A natural "simplest criterion" is: **number of open channels = neutron number N mod 2**, i.e., odd-N nuclei "have one unpaired open channel", even-even nuclei "fully paired closed". This criterion predicts that even-even nuclei should have uniformly minimal thermal cross-sections.

It is falsified by three counterexamples:

| Counterexample | N | σ_th (b) | Conflict with "even-even should be closed" |

|---|---|---|---|

| U-232 | 140 (even) | **76.5** | High |

| Pu-238 | 144 (even) | **17.8** | Medium-high |

| Cm-242 | 146 (even) | **4.67** | Medium |

Within the same even-even category, σ_th ranges from 76.5 b (U-232) to 1.7e-5 b (U-238), spanning **6.7 orders of magnitude**. Parity itself cannot determine whether thermal fission is possible.

11.3 Upgrade: Compound Nucleus Excitation Surplus Criterion

The real criterion is the **excitation surplus** of the compound nucleus (A+1):

$$\Delta = E^* - B_f = S_n(A+1) - B_f(A+1)$$

where E"=S_n(compound nucleus) is the excitation energy after absorbing a thermal neutron, and B_f is the fission barrier. Δ>0: the excitation energy itself exceeds the rebalancing slope → thermally sedimentable; Δ<0: requires additional kinetic energy from fast neutrons.

Results (full table of 9 target nuclei):

| Target Nucleus | E" (MeV) | B_f (MeV) | Δ (MeV) | $\log_{10}\sigma_{th}$ | Judgment |

|---|---|---|---|---|---|

| U-233 | 6.846 | 5.50 | **+1.35** | 2.73 | Thermally sedimentable |

| U-235 | 6.545 | 5.67 | **+0.88** | 2.77 | Thermally sedimentable |

| Pu-239 | 6.534 | 6.05 | **+0.48** | 2.87 | Thermally sedimentable |

| Am-241 | 5.529 | 5.90 | −0.37 | 0.49 | Requires fast neutrons |

| Np-237 | 5.488 | 6.10 | −0.61 | −1.69 | Requires fast neutrons |

| Pu-240 | 5.242 | 6.10 | −0.86 | −1.44 | Requires fast neutrons |

| Pu-242 | 5.034 | 5.80 | −0.77 | −2.61 | Requires fast neutrons |

| Th-232 | 4.786 | 5.80 | −1.01 | −4.27 | Requires fast neutrons |

| U-238 | 4.806 | 6.20 | −1.39 | −4.77 | Requires fast neutrons |

Pearson(Δ, $\log_{10}\sigma_{th}$) = +0.932.

- The positive and negative of Δ completely separates the two levels of "thermally sedimentable / requires fast neutrons" with a margin of 0.4–1.4 MeV;
- The three even-even "counterexamples" are all automatically returned to their correct positions using this criterion: the compound nucleus U-233" of U-232 has Δ≈+1.6, the compound nucleus Pu-239" of Pu-238 has Δ≈+0.6, and the compound nucleus Cm-243" of Cm-242 has Δ≈+0.6 — although they are even-even, their excitation surplus is already positive, so they can settle at low energy. **"Even-even" is a common but not sufficient manifestation of "negative surplus"; the parity law is a subset of the closure degree potential.**

11.4 B/A Closure Degree Potential: Macroscopic Profile of Fission/Fusion Direction and Net Energy Accounting

B/A (binding energy per nucleon) as the macroscopic profile of "closure degree potential":

- Extremum: **Ni-62 / Fe-56 platform, B/A≈8.79 MeV/n** — maximum closure degree region;
- α particle (He-4) B/A=7.07: higher saturation at the light nucleus end;
- U-238 B/A=7.57:falls back at the heavy nucleus end (sparse long-range channels).

Both fission and flow towards the B/A single peak; the direction is determined by the position of the starting point relative to the peak:

| Migration | d(B/A) (MeV/n) | Energy Released | SRE Reading |

|---|---|---|---|

| 4He → Fe | +1.72 | Fusion energy release | Flow towards greater closure degree |

| Fe → U | **−1.22** | Fission energy release | Leave the maximum closure degree region andfalls back |

Estimate the fission energy release of U-235 using the B/A difference: take the average B/A of fragments from the 98/138 region or 95/140 interpolation (≈8.50–8.51 MeV/n):

$$E_f \approx 236 \times (\overline{B/A}_{frag} - B/A_{U^{235}}) \approx 215\text{–}219\ \text{MeV}$$

The measured E_f=202.5 MeV, with a model deviation of **+6~8%** (direction and magnitude correct; residual deviation comes from fragment B/A selection point and mass deduction of secondary neutrons).

11.5 Chain Criticality: Multiplication and Coherent Bookkeeping Self-Sustaining Valve

Chain reactions upgrade "whether a single seed can land" to "whether the statistical family of seeds can not be extinguished**:

$$k_{eff} = \nu \cdot p \cdot f \geq 1$$

where v is the average number of neutrons emitted per fission, p is the probability that each neutron successfully triggers the next generation of fission, and f is the proportion of neutrons that do not leak/are not absorbed.

Critical sedimentation probability p"=1/(v·f) (upper limit when f=1):

| Nuclide | v | p"=1/v |

|---|---|---|

| U-233 | 2.47 | 0.405 |

| U-235 | 2.43 | 0.412 |

| Pu-239 | 2.87 | 0.348 |

| U-238 (fast) | 2.51 | 0.398 |

Chain self-sustainment requires each seed to settle with a probability >35%, otherwise the multiplication <1 and the reaction extinguishes; p"∈[0.35,0.41] matches the measured values.

11.6 Three-Level Signature Unification

| Scale | Event | Signature | Ontology |

|---|---|---|---|

| Single Nucleon | β decay | n(open, β_1=6) → p(closed, β_1=7), Δm/m_p=1.4e-3 | Open→closed, expelling seed |

| Composite Nucleus | Thermal neutron fission | σ_th spans 8.6 orders of magnitude, fast neutrons restore to the same order of magnitude; driven by Δ(S_n−B_f) | Whether the closure degree potential crosses the rebalancing slope |

| Macroscopic | Chain Criticality | k_eff=v·p≥1, p"=1/v≈0.35–0.41 | Statistical family of seeds does not extinguish |

All three share the same mechanism: **dormant open channels that can be sedimented**. β decay, fission cross-section parity law (which is essentially excitation surplus), and chain criticality are three readings of it at three scales.

---

12. Many-Body Assembly Law and the Shared Ring: A Validation Attempt Borrowing Nuclear Data

This section records an attempt to extend the nucleon skeleton to many-body systems using external nuclear data as a yardstick. The methodological constraints follow Section 3: SRE can only produce dimensionless ratios, so every quantity compared here is dimensionless — $\Delta m_{np} = m_n - m_p = 1.293332$ MeV is taken as the natural closure-degree unit on the nucleon side. Each conclusion states whether it is falsifiable. All numerical values are independently recomputed by the scripts in `code/`.

12.1 The Assembly Law: Uniquely Determined by Two Principles

Before nucleon skeletons can be assembled into many-body systems, the assembly rule must be fixed. This paper admits only two principles:

1. **Open–closed complementarity**: two bodies can dock and form a shared unit only when one side carries a dormant gap and the other side offers a complete ring edge at the same position;
2. **The images of the two bodies must be distinguishable in the shared region**: the shared ring must contain at least one **single-claim edge** (k=1, i.e. an edge claimed by only one body). This clause was originally stated as "closed bodies must not overlap", which carries spatial semantics and conflicts with the ontology of SRE, in which there is no space; it is therefore restated as a purely ledger-theoretic condition. **"Same position" is the natural meaning of "indistinguishable"** — the gap of neutron ring j can be sealed only by a complete ring edge of the partner's **ring j**, not by an arbitrary ring, and this is not an extra assumption.

Five construction variants are enumerated for the two-body (deuteron) case: gap against a ring edge, gap against a spoke, single-point contact, three points identically attached to a ring, and merging onto the same vertex. Only **variant A** (the two ends of the gap are identified with the two ends of one of the proton's ring edges, forming a new ring) satisfies both principles, and it is unique under automorphism (9 constructions merge into one isomorphism class):

Variant A (two bodies)	Value
V / E / β_1	22 / 35 / 14
Number of triangular rings T	6
$\Delta\beta_1$ / ΔT	+1 / +1
λ_2	0.238442818
ρ	6.681330644
Π_1	0.035687924


Qualitative prediction (successful): the rule automatically excludes n-n (both sides carry gaps at the same position) and p-p (neither side carries a gap); **only n-p can form a shared unit** — consistent with the measured fact that the only bound two-body system is the deuteron. The basis is the open/closed asymmetry, i.e. difference itself, and not any additional force.

Discriminating power of the restated principle 2 (verification): replacing "must not overlap" by "there exists a single-claim edge (k=1)" reproduces all the original exclusions: the shared-ring ledger of n-p is (1, 2, 2), containing one single-claim edge with k=1 → legal; in n-n both sides are dormant at the same position, so after coincidence that edge is still missing and the shared region retains only 2 edges (the ring does not close) → illegal; in p-p all three edges are claimed by both bodies (2, 2, 2) with no single-claim edge → the images of the two bodies are indistinguishable → illegal. A further check shows that **the shared-ring products of n-p and p-p are isomorphic to each other** (V=21, E=33, T=5, β_1=13, |Aut|=48, ρ=6.000000, λ_2=2−√3 all identical), so p-p can be excluded only by the ledger, never by any graph invariant.

Correction of the status of variant D (formerly the "shadow solution"): the construction in which three points are identically attached to a ring gives V=21, E=33, β_1=13, T=5, |Aut|=48, with a spectrum of strict arithmetic rigidity (see 12.3). Verification shows that this construction is **isomorphic to the k=2 member of the k-body shared-ring family of Section 12.5 and carries the same ledger (1, 2, 2)** — it is therefore the deuteron's own configuration; the name "shadow solution" and the former note that it "violates principle 2" are both withdrawn. Under the restated principle, "the two nucleons occupy the same position" is no longer a ground for exclusion (SRE has no space); what is genuinely excluded is the **p-p type of complete coincidence** (no single-claim edge). Variant A (two-point docking, only one shared edge, $\Delta\beta_1$=+1) remains a legal configuration but is quantitatively excluded by the deuteron binding energy (see 12.2). The full disposal is given in 12.10.

Many-body assembly law and shared ring: assembly, reference mode, and zero-parameter scorecard

(a) Shared-ring assembly (monomer → two bodies → four bodies)




Monomer (closed proton)
three-strand Y triangle closed body (V=12)

Two bodies sharing one ring (deuteron)
V=21, ledger (1, 2, 2)

Four bodies sharing one ring (^{4}He)
V=39, ledger (2, 4, 4)

- Thick solid edge: shared-ring edge claimed by all bodies (k = A)
- Dashed red edge: edge claimed by protons only (k = #p) — single-claim edge
- Blue dot: closed state (proton, ring edge intact); red dot: open state (neutron, ring edge dormant)

(b) Reference-mode invariance and closure potential



(c) Zero-parameter scorecard of the assembly law (reading A)

system	A	g	s	verdict	measured	hit
n-p	2	1	1	legal	bound 2.2246 MeV	√
n-n	2	2	0	illegal	unbound (virtual)	√
p-p	2	0	2	illegal	unbound (virtual)	√
^{3}H	3	2	1	legal	bound 8.4820 MeV	√
^{3}He	3	1	2	legal	bound 7.7181 MeV	√
^{4}He	4	2	2	legal	bound 28.2957 MeV	√
^{4}H	4	3	1	illegal	unbound (resonance)	√
^{4}Li	4	1	3	legal	unbound (resonance)	×

Reading A (iff): legal when $1 \le g \le 2$ and $s \ge 1$. Hits 7 of 8; 5/5 on $A \le 3$; the only mismatch is ^{4}Li.

Figure 5: Many-body assembly law and the shared ring. (a) The monomer skeleton is assembled into a k-body shared ring by "merging the same ring (three vertices) into the shared region": thick solid edges are claimed by all bodies (k=A), the dashed red edge is a single-claim edge claimed by protons only (k=#p), blue dots are closed states and red dots open states; (b) the closure potential ρ rises monotonically with the body count, whereas the lowest non-trivial mode λ_2 = 2−√3 is independent of the body count (reference mode; the data are independently recomputed by the companion script from numerical Laplacian spectra, with a

maximum deviation of about 10^{-15}); (c) the zero-parameter scorecard of the assembly law (Reading A, see 12.8). The detailed data of panels (b) and (c) are given in 12.5 and 12.8 respectively.

12.2 Variant A Is Quantitatively Excluded by the Deuteron

Taking Δm_np as the closure-degree unit, variant A carries a closure-degree increment $\Delta\beta_1 = +1$, so its zero-parameter prediction for the deuteron binding energy is

$$B_d^{(\mathrm{pred})} = \Delta\beta_1 \cdot \Delta m_{np} = 2.5867\ \mathrm{MeV}$$

against the measured B_d = 2.224566 MeV, i.e. **16.3% too high**. The difference 0.362 MeV = 0.28 Δm_np is the "sharing tax" — variant A contains no shared edge at all and therefore cannot carry this term. **Conclusion: a configuration without shared edges cannot serve as the deuteron; the deuteron must be carried by a shared-ring configuration.**

12.3 The Shared-Ring Motif: Arithmetic Rigidity of Variant D

Variant D (three points identically attached to a ring; the k=2 member of Section 12.5, see 12.10)	Value
V / E / β_1 / T	21 / 33 / 13 / 5
|Aut|	48
Laplacian trace	66
Characteristic polynomial	$x\cdot(x-1)^2(x-2)^2(x-4)^4(x-5)^2(x-6)^2(x^2-4x+1)^1(x^2-6x+7)^3$
Spectrum (value × multiplicity)	$0^1, 1^2, 2^2, 4^4, 5^2, 6^2, (2\pm\sqrt{3})^1, (3\pm\sqrt{2})^3$
ρ	6.000000 (exact integer, twofold)
λ_2	$2-\sqrt{3} = 0.267949$ (exact)

Every factor of the full spectrum has degree at most 2 (**arithmetic rigidity**), in contrast to variant A, which contains an irreducible cubic factor. Rarity check: among random graphs with the same degree sequence ($4,4,4,3^{18}$) (2966 samples), the median ρ is 6.057 with range [5.291, 6.639], while **ρ is exactly the integer 6 in 0 samples**, and lies within 6±0.01 in only 4.0%. "Integer ρ" is therefore genuinely anti-probabilistic.

Orbit structure and "the projection erases history": the vertices of D fall into 3 orbits (6 spoke centres, 3 shared-ring vertices, 12 leaves) and the edges into 4 orbits, among which **the 3 edges of the shared ring lie in one and the same orbit** — the graph itself cannot tell which edge was "borrowed". This is a direct graph-level instance of "the homomorphic mapping erases history".

12.4 The Projection Erases Who Is Dormant: n-p ≅ p-p

The D-type products of n-p and p-p are compared item by item:

Quantity	D-type of n-p	D-type of p-p

|---|---|---|

| V / E / β_1 / T / |Aut| / trace | 21 / 33 / 13 / 5 / 48 / 66 | identical |

| Graph isomorphism test | `is_isomorphic = True` | |

| Shared ledger | 1 own edge + 2 shared edges | 3 shared edges |

The product graph does not remember whether the partner was a neutron or a proton: the entire difference lies in the shared ledger, and the ledger is not a graph invariant. This is the structural root of why B_d/Δm_np cannot be an integer, and it shows that the idea "binding energy can be carried by the graph's closure count" must fail. The D-type of n-n is not isomorphic to these (T=4, β_1=12, |Aut|=16, with one extra cubic factor), because both sides are dormant at the same position and the gap remains unsealed after coincidence.

12.5 Three and Four Bodies: Structural Linear Law and Body-Count Independence of the Lowest Mode

The configurations in which k bodies share one ring are constructed explicitly and checked quantity by quantity:

| k | System | V | E | T | β_1 | |Aut| | ρ |

|---|---|---|---|---|---|---|---|

| 2 | deuteron | 21 | 33 | 5 | 13 | 48 | 6.000000 |

| 3 | ^{3}H / ^{3}He | 30 | 48 | 7 | 19 | 288 | 6.925423 |

| 4 | ^{4}He | 39 | 63 | 9 | 25 | 2304 | 7.909516 |

Closed forms: **V = 9k+3, E = 15k+3, T = 2k+1, β_1 = 6k+1, |Aut| = 6·2^k·k!** (exact for k = 2, 3, 4). The rise of ρ with the body count and the body-count independence of λ_2 are shown in Figure 5(b) (independently recomputed by the figure script from numerical Laplacian spectra).

^{3}H(a) and ^{3}He(a) are isomorphic (V=30, E=48, T=7, β_1=19, |Aut|=288, ρ=6.925423, λ_2=2−√3 all identical) — the third occurrence, after the deuteron, of "the projection erases who is dormant".

The lowest non-trivial mode is independent of the body count: λ_2 = 2−√3 = 0.267949192431 is exactly the same for k = 2, 3, 4, 5, 6 (relative difference about 10^{-15}), and its degeneracy multiplicity is m(λ_2) = k−1. The mechanism was verified block by block:

- on the lowest mode, the **three vertices of the shared ring have strictly zero amplitude** (about 10^{-16});
- the spoke-centre amplitudes of the individual bodies **sum to zero** (for k=2: +0.1877 / −0.1877; for k=3: −0.2167 / +0.1084 / +0.1084);
- hence inter-body coupling is completely switched off on this mode, and each body oscillates independently according to its own "body minus shared ring" grounded block, whose lowest eigenvalue is exactly 2−√3 (its 9 eigenvalues are {0.26795, 1, 3.26795^2, 3.73205, 4^2, 6.73205^2});
- k contributes only one constraint, "the spoke-centre amplitudes sum to zero", so the eigenvalue is independent of k and its multiplicity is exactly k−1.

Physical reading: the shared ring = the part of the lowest excitation that **does not move at all**, i.e. a natural reference frame. This is the most concrete realization to date of the ontological reading that "light is the topological residue between evolutions and serves as the reference object of measurement": **the measured component (numerator) varies with the body count, while the reference component (the shared ring) stays strictly fixed.** By contrast, ρ rises monotonically with k (6 / 6.925423 / 7.909516 / 8.908855 / 9.912634) and has no simple quadratic closed form — **for k ≥ 3 the arithmetic rigidity moves from ρ to λ_2**.

12.6 Shared-Ledger Pricing: Falsification at the Energy-Level Layer

With the structural layer settled, the remaining question is the quantitative attribution of binding energy. The simplest bookkeeping hypothesis is: **each overlapping edge on the shared ring is priced according to how many bodies claim it (k)**, ψ(k), and the binding energy of a nucleus (in units of Δm_np) is the sum of the three edge prices. The ledgers of the four measured nuclei are:

Nucleus	Ledger (claim count of each of the three edges)	B / Δm_np
deuteron	(1, 2, 2)	1.720027
^{3}H	(1, 3, 3)	6.558252
^{3}He	(2, 3, 3)	5.967608
^{4}He	(2, 4, 4)	21.878135

The four equations determine ψ(1..4) exactly, with zero residual:

k	ψ(k) / Δm_np	ψ(k) / MeV	relative to ψ(1)
1	+0.967105	+1.2508	1.0000
2	+0.376461	+0.4869	**0.3893**
3	+2.795573	+3.6156	**2.8907**
4	+10.750837	+13.9044	**11.1165**

The shape is non-monotonic: double occupation reduces the edge contribution to 38.9% (a tax), while triple and quadruple occupation give 2.89× and 11.12× (strong gain). This is aligned with the qualitative picture "the greater the interference depth, the stronger the binding".

It must be stated honestly: this layer has no predictive margin. Four free parameters fitted to four data points give zero residual by necessity, not by achievement; the only valid test is out-of-sample (see 12.7). Moreover, two more parsimonious pricing laws cannot accommodate the measured data (both are calibrated with a single constant from the deuteron):

Pricing law	^{3}H	^{3}He	^{4}He
Law A: ψ(k) = k·c (linear in k)	−63.3%	−53.9%	−84.3%
Law B: ψ(k) = c·C(k,2) (pairwise counting)	−21.3%	+0.9%	−48.9%

In other words, "interference depth" cannot be reduced to a linear or pairwise count in k; Law B's near-hit on ^{3}He is a single-point coincidence and cannot be generalized.

Backfill: the root of this layer's failure lies not in the shape of ψ but in additivity itself. (This item is homologous to 12.11(2) and is registered here in advance.) From the three **resonance-unambiguous** measured nuclei (deuteron, ^{3}H, ^{3}He, all with A ≤ 3), subtracting the three equations cancels ψ(3):

> ψ(1) − ψ(2) = B(^{3}H) − B(^{3}He) = +0.590645 Δm_np = **+0.7639 MeV**

This expression involves only the price difference of the first edge and is independent of the shape of ψ. Since the profiles of ^{4}H and ^{4}He are (1, 4, 4) and (2, 4, 4), differing only in the claim count of the first edge, it immediately gives

> B(^{4}H) − B(^{4}He) = ψ(1) − ψ(2) = **+0.590645 Δm_np** (an identity)

That is, **however one redefines "the price of each edge on the shared ring", ^{4}H is forced to be predicted more tightly bound than ^{4}He**. This is precisely the mechanism of the out-of-sample failure in 12.7 — the failure occurs within the very section that introduces ψ, and it cannot be repaired by changing the shape of ψ. Likewise ψ(3) − ψ(2) = +2.419113 Δm_np gives B(^{4}Li) − B(^{4}He) = +2.419113 Δm_np. Hence the failure in 12.7 is a failure of the **whole class**, not of any particular ψ; the full derivation and consequences of this identity are given in 12.11(2).

Another equivalent simple bookkeeping (registered for the record): let the price of an own edge be 1, of a dormant edge d, of a shared edge c, and of a borrowed edge b. Calibrating on the single nucleon, Δm_np = d − 1 gives d = 2; from the deuteron, 6 − 4c − b = 1.720027. Taking "borrowing is free" (b = 0) yields c = 1.07 and a sharing tax 4(c − 1) = 0.28 Δm_np = 0.362 MeV, consistent with the difference in 12.2. This parameterization and ψ(k) are two spellings of the same layer, and neither has an independent predictive margin.

12.7 Failure of the A=4 Out-of-Sample Prediction

Once ψ has been fixed by A=2, A=3 and ^{4}He, the A=4 isobars constitute a genuine zero-parameter out-of-sample set:

Nucleus	Ledger	Predicted B	Measured
^{4}He	(2, 4, 4)	28.2957 MeV	bound, 28.2957 MeV (**used for calibration**)
^{4}H	(1, 4, 4)	29.0596 MeV	**unbound** (above-threshold resonance)
^{4}Li	(3, 4, 4)	31.4244 MeV	**unbound** (above-threshold resonance)

The model predicts both to be more strongly bound than ^{4}He, whereas both are measured to be **unbound**. A pricing law that depends only on "the distribution of claim counts k over the overlapping edges" is therefore explicitly falsified: **the energy-level layer is not computable in this layer.**

12.8 The Assembly-Law Scorecard and Adjudication of Readings

Reducing the two principles of 12.1 to "the number of gaps and the number of supplies at the same position" yields a zero-parameter existence scorecard. Definition: at the same shared position, k nucleon skeletons are aligned; each open-state body (neutron, whose ring edge at that position is dormant) leaves 1 **gap** g, and each closed-state body (proton, whose ring edge at that position is complete) offers 1 **supply** s. **A legal assembly implies binding; an illegal assembly implies non-binding.**

Nucleus	A	Gap g	Supply s	Measured
n-p	2	1	1	bound, 2.2246 MeV
n-n	2	2	0	unbound (virtual state)
p-p	2	0	2	unbound (virtual state)
^{3}H	3	2	1	bound, 8.4820 MeV
^{3}He	3	1	2	bound, 7.7181 MeV
^{4}He	4	2	2	bound, 28.2957 MeV
^{4}H	4	3	1	unbound (above-threshold resonance)
^{4}Li	4	1	3	unbound (above-threshold resonance)

Two zero-parameter readings coexist and are mutually incompatible:

- **Reading A**: legal ⟺ $1 \le g \le 2$ and $s \ge 1$ (a gap exists, at least one side can seal it, and the number of dormant sides does not exceed 2);
- **Reading B**: legal ⟺ $g = s$ and $g \ge 1$ (gaps and supplies match one-to-one).

Reading	All 8 entries	A ≤ 3 sector	Mismatches
A (adopted here)	**7 / 8**	**5 / 5**	^{4}Li (A=4)
A' (dropping the clause "g ≤ 2")	6 / 8	5 / 5	^{4}H, ^{4}Li (both A=4)
B (one-to-one matching)	6 / 8	3 / 5	^{3}H, ^{3}He (both A=3)

Adjudication: Reading A is adopted. There are two reasons: (i) all five entries of the calibratable sector A ≤ 3 (n-p, n-n, p-p, ^{3}H, ^{3}He, all of them measured bound states or sub-threshold virtual states without resonance ambiguity) are matched under Reading A, whereas Reading B already mismatches at ^{3}H, judging the measured bound ^{3}H to be unbound, and is therefore excluded; (ii) the only mismatch of Reading A lies at A = 4, and A = 4 has already been independently delimited as a non-computable interval in 12.7. If the additional clause "g ≤ 2" is dropped, the mismatch set gains only one more entry, ^{4}H, still inside the same interval — **so whether that clause is kept does not affect the conclusions of this paper**; the version tabulated above (Reading A) is registered here. The scorecard is displayed in Figure 5(c).

Later supplement (see 12.12): of the two clauses of Reading A ($1 \le g \le 2$ and $s \ge 1$), "g ≤ 2" caps only the neutron count and not the proton count, and therefore mismatches at ^{4}Li. Replacing the two clauses by the mirror-symmetric single criterion **L1: $|n_n - n_p| \le 1$** makes the same 8-entry scorecard match **8 / 8**, with the number of clauses reduced from two to one and no parameter. The registration in this section is therefore revised to a **parallel** one: Reading A (the original form of this section, 7 / 8) and criterion L1 (12.12, 8 / 8); over A ≤ 4, L1 strictly dominates, while over A ≥ 5 both degrade to empirical approximations.

12.9 The Adopted Boundary: A ≥ 4 Is Not Computable in This Layer

Combining 12.7 and 12.8:

- **A ≤ 3**: the zero-parameter assembly-law scorecard matches 5 / 5;
- **A = 4**: two classes of failure appear — the out-of-sample failure of the pricing layer (^{4}H and ^{4}Li predicted bound, see 12.7) and the qualitative failure of the assembly law (^{4}Li predicted bound, see 12.8).

Both classes fall in the same interval and constitute two independent markers of the following boundary:

> **Adopted boundary**: the nuclear boundness for A ≥ 4 (both existence and the numerical value of the binding energy) **is not computable in this layer**.

Explaining that interval requires a new quantity outside the ledger (candidate: interference depth). That candidate is explicitly tested in 12.11, and the conclusion is negative: **this boundary is not a matter of insufficient data but a structural one**.

Honest registration: the hit for ^{4}He is not a prediction (its binding energy was used to fix ψ(4)); ^{4}H and ^{4}Li share the same reading and cannot be kept selectively; the "non-binding" of n-n and p-p refers to virtual states rather than forbidden states — the criterion only says that "no bound shared unit is formed".

Later revision (see 12.12): Section 12.12 makes an explicit attempt at the "antisymmetric penalty coefficient" λ; the conclusion is that λ can neither be derived from nor fixed by SRE. The same check, however, yields a zero-parameter **existence criterion** L1 ($|n_n - n_p| \le 1$) that matches the 8-entry scorecard of 12.8 at 8 / 8. The boundary above must therefore be split according to the two questions "existence" and "numerical value of the binding energy": the **numerical value of the binding energy** is not computable for A ≥ 4 (the identity of 12.11(2) is unaffected); **existence** is computable at A = 4 (L1, 8 / 8) but not for A ≥ 5 (the criterion degrades into an empirical approximation). That is, the boundary is **relocated** from A ≥ 4 to A ≥ 5.

12.10 Disposal of the Shadow Solution: From "Overlap" to "Distinguishability"

The two pending items raised in 12.1 are settled one by one. All numerical values are independently recomputed by `code/_sre_nucleon_shadow_disposal.py`.

(1) Variant D is not a shadow solution; it is the deuteron's own configuration. Constructed as "k bodies sharing the same ring", the n+p common ring (formerly variant D) and the two-body shared ring of Section 12.5 (k=2) yield the same invariants (V=21, E=33, T=5, β_1=13, |Aut|=48, ρ=6.000000, $\lambda_2=2-\sqrt{3}$; the deviations of ρ from the integer 6 and of λ_2 from $2-\sqrt{3}$ are both at the 10^{-15} level), with `is_isomorphic = True` and the same shared-ring ledger (1, 2, 2). The name "shadow solution" and the former note "violates principle 2" are therefore withdrawn; the value of retaining D rests on facts rather than on naming — the integer ρ and the body-count-independent λ_2 are genuine anti-probabilistic features (see 12.3 and 12.5).

(2) The non-spatial restatement of principle 2 has the same discriminating power. The original statement "closed bodies must not overlap" carries spatial semantics and conflicts with the ontological premise of "no space, no dimension". Restated as a purely ledger-theoretic condition — **the shared ring contains at least one single-claim edge (k=1)** — the verdicts for the three pairings are identical to those of the original statement:

| Pairing | Shared-ring edges | Ledger profile | Single-claim edges (k=1) | Verdict | Measured |

|---|---|---|---|---|---|

| n-p | 3 (ring closed) | (1, 2, 2) | 1 | legal | bound |

| n-n | 2 (ring open) | (2, 2) | 0 | illegal | unbound (virtual) |

| p-p | 3 (ring closed) | (2, 2, 2) | 0 | illegal | unbound (virtual) |

(3) The ledger difference between the two sharing modes. Variant A (the two ends of the neutron's gap identified with the two ends of **one** of the proton's ring edges) has only **one dual-service edge** (that edge closes both rings, yet it appears in the proton's edge set only); the shared-ring configuration has **three** edges claimed by all bodies. The accounting of 12.6 ("each of the three shared-ring edges has its claim count") can therefore describe only the shared-ring configuration; the A-type cannot carry a three-edge profile such as (1, 2, 2). The two routes are not a mutually exclusive structural ambiguity, but a choice under the same body of data: the A-type is quantitatively excluded by the deuteron binding energy (16.3% too high, see 12.2), while the shared-ring type is retained.

12.11 Why the A ≥ 4 Boundary Cannot Be Repaired: Two Impossibilities

The boundary of 12.9 was originally based on "two empirical failures". Further checks show that those two failures are not due to insufficient data but can be proven by two impossibilities: **no graph functional can carry the distinction at A = 4, and no additive ledger pricing can repair its sign.** The numbers are independently recomputed by `code/_sre_nucleon_interference.py`.

(1) Structural impossibility: the A = 4 isotriplet forms a single isomorphism class. Building ^{4}He (2p+2n), ^{4}H (1p+3n) and ^{4}Li (3p+1n) under the shared-ring rule gives three graphs with identical invariants (V=39, E=63, T=9, β_1=25, |Aut|=2304, ρ=7.909515966, λ_2=2−√3), pairwise isomorphic. Their only difference lies in the ledger: the claim count of the "protons-only" edge on the shared ring is 2 / 1 / 3. **This is the A = 4 extension of the proposition "n-p ≅ p-p" of 12.4**: the product graph does not remember who is dormant, so the binding-energy differences cannot be carried by the graph, nor by any graph functional (ρ, λ_2, T, β_1, |Aut|, the full spectrum).

(2) Pricing-layer impossibility: the additive-profile pricing class is falsified as a whole. The pricing law of 12.6 can be written in general form: the i-th edge of the shared ring is priced by its claim count k_i, and the binding energy is the sum over the three edges, i.e. B = $\psi(k_1) + \psi(k_2) + \psi(k_3)$. Since k = (#p, A, A), every nucleus satisfies B(A, #p) = ψ(#p) + 2ψ(A), where ψ is an **arbitrary** function. The three measured nuclei d, ^{3}H and ^{3}He (A ≤ 3, no resonance ambiguity) immediately give an identity:

> $\psi(1) - \psi(2) = B(^3\text{H}) - B(^3\text{He}) = +0.590645\ \Delta m_{np} = +0.7639$ MeV

Because the profiles of ^{4}H and ^{4}He differ only in the first edge (1 versus 2),

> $B(^4\text{H}) - B(^4\text{He}) = \psi(1) - \psi(2) =$ **+0.590645 Δm_np** (an identity, independent of the shape of ψ)

Likewise $\psi(3) - \psi(2) = +2.419113\ \Delta m_{np}$, giving $B(^4\text{Li}) - B(^4\text{He}) = +2.419113\ \Delta m_{np}$. That is: **however one redefines the "interference depth" of a single edge, ^{4}H and ^{4}Li are forced to be predicted more tightly bound than ^{4}He** (22.468780 and 24.297248 Δm_np, i.e. 29.0596 and 31.4244 MeV), whereas both are measured unbound. The sign is locked by the three measured nuclei, independently of the shape of ψ — so this layer cannot be repaired, only overturned.

(3) A composition-resolved interference depth is likewise insufficient. Refining "interference depth" to the compositional level, i.e. splitting each edge's claim count into (k_n, k_p), allows one to define the complementary interference C = Σ k_n k_p, the same-type interference S = Σ [C(k_n,2) + C(k_p,2)], the total excess claim count I = Σ (k − 1) and the imbalance a = |n_n − n_p|. Calibrated on the four measured nuclei (d, ^{3}H, ^{3}He, ^{4}He), the maximum residual of any linear basis (3 fitted parameters including the constant, only 1 residual degree of freedom) is: (C,S) 10.2%, (C,a) 3.3%, (I,a) 24.6%, (C,I) 10.2%; and extrapolating ^{4}H and ^{4}Li with the same coefficients, **all signs come out "bound", opposite to measurement**. There is also an independent mismatch in shape: ^{3}H and ^{3}He have the same complementary interference C (both 4), while their same-type interference S is 2 versus 3 and I is 4 versus 5, yet the measured binding energies are 6.558 versus 5.968 Δm_np — **decreasing** with S (or I). If one insists on the qualitative tendency of 12.6 ("the greater the interference, the stronger the binding"), then ^{3}H should be weaker than ^{3}He, contrary to measurement. Hence, if "interference depth" is to carry the pricing, it must be non-additive and non-monotonic, and no derivable form of it exists in the present framework.

(4) The only feasible repair and its price. For ^{4}H and ^{4}Li to be unbound while d, ^{3}H, ^{3}He and ^{4}He are unaffected, an additional term T must satisfy T(d) = T(^{3}H) = T(^{3}He) = T(^{4}He) = 0 and T(^{4}H) = T(^{4}Li) ≥ 22.468780 Δm_np (= 29.0596 MeV). The minimal form is a function of the **imbalance** a = |n_n − n_p|: identically zero for a ≤ 1 and sufficiently positive at a = 2 (for example g(a) = λ·max(0, a−1) with λ ≥ 29.0596 MeV). This form needs only 1 free parameter and automatically respects mirror symmetry (^{4}H and ^{4}Li both have a = 2 and are pushed above threshold together). Its price must be honestly registered:

- λ has no provenance within SRE; it is a **fitted parameter, not a prediction**; this registration will be upgraded in 12.12 to a structural conclusion (λ can neither be derived nor fixed);
- the form has content only for A = 4. Every A ≥ 5 introduces a new free value ψ(5), ψ(6), … , so the scheme itself has no predictive margin for A ≥ 5, and "restoring computability" cannot be tested there;
- this item therefore serves only as a **reinforcement of the falsification**: what repair requires is not a better interpolation but a quantity outside the ledger (the imbalance here), and that quantity currently has no falsifiable predictive power. **It should be noted, however, that if that quantity is used only as a binary (existence) requirement, its magnitude is not needed, so λ becomes dispensable and a contentful zero-parameter criterion results — this is the outcome of 12.12.**

(5) Summary. A = 4 is the **only** interval in which the scheme possesses genuine zero-parameter out-of-sample character **in the energy layer**: d, ^{3}H and ^{3}He fix ψ(1..3), ^{4}He fixes ψ(4), and ^{4}H and ^{4}Li are true extrapolation points. The boundary falls exactly at the first and only real extrapolation — indicating that its **location** is not accidental and its **nature** is structural (graph-layer indistinguishability + additive-layer identity), rather than a matter of data precision. **What must be distinguished is that this "irreparability" concerns the numerical value of the binding energy; if only existence is asked, that question is not constrained by the additive identity, and 12.12 gives a zero-parameter decision for it, on which the boundary is accordingly revised.**

Honest registration: the "single isomorphism class" of (1) refers only to products built under the shared-ring rule; if another assembly rule is used, the conclusion must be re-checked. **(1') That re-check has now been carried out in this section, and it yields a stronger statement than the measurement itself**: the isomorphism of the three is **not** a coincidence of the assembly rule but is **necessary at the assembly layer** — see below. The identities of (2) falsify the "additive-profile pricing class" only and do not exclude a non-additive pricing; no non-additive form is given here. The imbalance term of (4) is explicitly marked as a fit and must not be used as a prediction.

(1') Strengthening and narrowing of the structural-layer conclusion: A does not enter the structure; k does. Let k denote the **body count** of a shared ring (how many nucleons' rings that ring merges). Using a controlled-variable design, `code/_sre_nucleon_structure_vs_ledger.py` and `code/_sre_nucleon_attachment_sweep.py` measured three comparisons:

- **Same k, change composition (p↔n)** ⇒ the graph is unchanged and the two are isomorphic ((p,p) ~ (p,n) is `True`); the difference lies only in the ledger profile ((2,2,2) versus (2,2,1)). The same holds for the mirror pair ^{3}H ~ ^{3}He (`True`; |Aut| = 288 and ρ = 6.925422918 identical).
- **Same A, change the distribution of k** ⇒ the graph changes and the two are **non-isomorphic**. Example: at A = 6, k = (3,3)

gives V = 60, E = 96, $\beta_1 = 38$, ρ = 6.925422918, whereas k = (2,2,2) gives V = 63, E = 99, $\beta_1 = 39$, ρ = 6.000000000. **One qualifier must be attached, however**: the current assembly rule uses the group index simultaneously as the ring slot, so **whenever the k distribution contains two or more groups the product graph is invariably a disconnected union of independent components** (`cc` = number of groups, **9/9 measured**; the λ_2 of both examples above is printed as `0.000000000` in the original log, whereas the λ_2 of a connected graph is invariably $2-\sqrt{3} = 0.267949192$, so **$\lambda_2 = 0$ is the fingerprint of disconnection**). The "graph changes" in this example is therefore caused **directly by V/E/β_1 adding up over the components of a disconnected union** (the addition law is given in 12.5), **not** by "the k distribution entering a single nucleon structure". **The inter-ring linking mechanism is absent from the assembly rule**, and it is provably **impossible to supply**: connectivity costs at least one inter-ring edge, which necessarily raises E from 96 to 97 and is thus **logically incompatible with the published values above**; more strongly, no mechanism satisfies all four of "degenerates to the current rule for a single group, is connected for multiple groups, does not destroy the $\lambda_2 = 2-\sqrt{3}$ of 12.5, introduces no new vertices" (see the unresolved matters of §15 and `code/_sre_nucleon_interring_b2.py`). **This bullet therefore supports only the weak reading "the same A with different k distributions gives different graphs", not the strong reading "the k distribution enters a single nucleon structure"**; by the morphological conclusion of §12.5 (under purely SRE-endogenous conditions the k distribution is invariably single-group), multi-group configurations should not arise in this layer at all.

- **Same A and same k distribution, change the attachment permutation** ⇒ the graph is **still unchanged**; all six permutations give **a single isomorphism class**. Mechanism: the automorphism group of a **closed** body induces the **full S_3** on the three points of its ring (6 permutations measured; an open body induces only 2). Hence "which point it attaches to" is absorbed by the automorphism, and **the attachment point is not a degree of freedom.**

The isomorphism of §(1) must therefore be **re-read**: ^{4}He, ^{4}H and ^{4}Li at A = 4 share the **same** k distribution (a single ring with k = 4), so their isomorphism is a **direct consequence of the consistency of k**, independent of the attachment point and of the specific value of A. This is stronger than the original statement that "three cases were measured to be isomorphic", and it **no longer depends on whether the assembly rule has been changed**. Conversely, **A itself does not determine the structure**; what determines it is the **distribution of k** (with k ≥ 2 for each group — the "ring" of a k = 1 group has no edge to merge, so an assembly such as k = (5,1) does not exist formally). **What determines the distribution of k** is answered here as follows: **under purely SRE-endogenous conditions the only admissible distribution is the single-group one k = (A,)** (its uniqueness rests on only three anchors at A ≤ 4, of which A = 2 and A = 3 each admit but a single partition and thus have no discriminating power, leaving **exactly one effective discrimination**, so this is a **weak conclusion**); if external priors are admitted (such as the "4" of α-clustering, or an upper bound on k), the distribution is **not uniquely determinable from within SRE** (see the unresolved matters of §15).

A constitutive condition forced out by this test: the shared ring **identifies the ring edge (L0r, L1r) of each body as one and the same edge**. Consequently, when both bodies are closed that edge has claim count 2, when one is closed and the other open it has count 1, and **when both are open the edge does not exist**. "Both open" is thus excluded not by an external prohibition but because **the gluing relation fails in the open state** — the same fact as the S_4 measurement of §12.13 (E is smaller than S_1 by exactly 1). Its corollary is that "there is always exactly one k = 1 edge on a shared ring" is equivalent to "**each shared ring carries exactly one open body**", and thus the statement of §12.5 that $\lambda_2 = 2-\sqrt{3}$ is independent of the body count acquires an assembly-layer explanation: **that k = 1 edge is a product of open–closed pairing, not a free parameter.**

12.12 The Attempt at an Antisymmetric Coefficient: From an Energy Term to an Existence Criterion

Section 12.11(4) registered the only feasible repair of the A ≥ 4 boundary as a "mirror-antisymmetric penalty term outside the ledger", g(a) = λ·max(0, a − 1) with a = |n_n − n_p|, and honestly registered λ as a fitted parameter with no SRE-internal provenance. This section upgrades that registration to a quantitative test; all numerical values are independently recomputed by `code/_sre_nucleon_asymmetry.py`. **The conclusion is that λ can neither be derived from nor fixed by SRE; but the very process of this negative result yields a contentful positive outcome — once the antisymmetry is demoted from an energy term to an existence criterion, the existence question at A = 4 becomes decidable in this layer with zero parameters.**

(1) The nature of the threshold. Requiring ^{4}H to be unbound requires the penalty to satisfy g(2) ≥ B_ledger(^{4}H) = ψ(1) + 2ψ(4) = 22.468780 Δm_np = 29.0596 MeV. Set against the binding energies of the four measured nuclei:

| Quantity | Δm_np | MeV | Relation to threshold |

|---|---|---|---|

| B(deuteron) | 1.720027 | 2.2246 | 20.7488 Δm below |

| B(^{3}H) | 6.558252 | 8.4820 | 15.9105 Δm below |

| B(^{3}He) | 5.967608 | 7.7181 | 16.5012 Δm below |

| B(^{4}He) | 21.878135 | 28.2957 | **0.590645 Δm below** |

| **λ threshold = B_ledger(^{4}H)** | **22.468780** | **29.0596** | — |

That is, the required penalty **exceeds the binding energy of the α particle itself**. In other words, the "asymmetry cost" for $|n_n - n_p| = 2$ must be larger than the entire binding energy of ^{4}He.

The only ready-made asymmetry mechanism in nuclear physics is the liquid-drop asymmetry energy $E_{asym} = a_{sym} (N - Z)^2/A$ ($a_{sym} \approx 22$–24 MeV). At A = 4 with $|N - Z| = 2$ it gives $E_{asym} = a_{sym} \leq 24$ MeV < 29.06 MeV (still 5.06 MeV, i.e. 17.4%, short even at the upper bound); and at A = 3 it gives a non-zero $E_{asym}(^3H) \approx 7.7$ MeV, whereas ^{3}H has already been used for calibration and its ledger term must carry zero penalty. **Hence this ready-made "asymmetry energy" mechanism fails on two counts.**

(2) λ cannot be derived from SRE. SRE has only two classes of input available at A = 4:

- **Graph invariants** (ρ, λ_2, T, β_1, |Aut|, the full spectrum): by 12.11(1), the three A = 4 compositions form a single isomorphism class, so all of these take the **same constant** on ^{4}He/^{4}H/^{4}Li. A constant penalty would fall equally on the calibrated ^{4}He, so that the four calibration equations would no longer be satisfied by ψ(1..4) — **the calibration itself would be destroyed**. Graph invariants therefore cannot serve as λ.
- **Ledger functions** (functions of A and #p): any SRE-internal quantity X is a function of (A, #p), so X(a) is of the **same kind** as g(a); "writing λ as an SRE-internal quantity" produces only a renaming, not a new constraint.

Inspection of 16 "natural candidates" (ρ_4, T_4, β_1, |Aut|^(1/3), √(V+1), $\rho_4 - \rho_2$, ψ(4)−ψ(2), B(^{4}He), 2B(^{4}He), B(^{4}He) + B(deuteron), E − V, $1/\lambda_2$ and others): 11 fall below the threshold and 5 above. The threshold (22.4688 Δm_np) falls precisely in the gap between 21.8781 Δm_np (= B(^{4}He)) and 23.5982 Δm_np (= B(^{4}He) + B(deuteron)) — **the threshold is "satisfied", not "pinned down".**

Formally: the unknown set {ψ(1..4), λ} against 4 equations leaves λ with 1 free parameter; at A = 4 there is therefore **zero testing margin**; at A ≥ 5 every new body number brings in a new free value ψ(A), so the parameter count grows at the same order as the data count and there is **no out-of-sample falsification**. **λ is not a derivable coefficient but an absorbable degree of freedom.**

(3) Yet SRE does fix the "shape" of λ. Since the first class is excluded by (2), λ must be composition-dependent and must vanish on the calibration set {a ≤ 1}. Hence what SRE determines is precisely the **shape** of λ — its support (zero for a ≤ 1) and its monotonicity (λ > 0 monotonically lowers the binding energy) — while its **magnitude** is not determined. But the requirement at A = 4 is **binary (existence)**: as long as λ > 0 and acts at a = 2, ^{4}H and ^{4}Li are pushed above threshold regardless of how large λ is (provided λ ≥ threshold). **Hence the coefficient λ is dispensable for that binary requirement**; it is needed only when a numerical binding energy is required, and at A = 4 no testable number can be produced (see (2)).

(4) Demotion to an existence criterion: the zero-parameter antisymmetry criterion. Writing antisymmetry directly as a **legality criterion** rather than an **energy penalty**:

> **Criterion L1 (antisymmetry criterion)**: a shared-ring assembly is legal (predicted bound) ⇔ $|n_n - n_p| \leq 1$.

It has zero parameters and is naturally n↔p symmetric. On the 8-entry scorecard of 12.8:

Nucleus	A	a = |n_n − n_p|	Measured	L1	Reading A
n-p	2	0	bound	bound √	bound √
n-n	2	2	unbound	unbound √	unbound √
p-p	2	2	unbound	unbound √	unbound √
^{3}H	3	1	bound	bound √	bound √
^{3}He	3	1	bound	bound √	bound √
^{4}He	4	0	bound	bound √	bound √
^{4}H	4	2	unbound	unbound √	unbound √
^{4}Li	4	2	unbound	unbound √	**bound ×**

L1 matches 8 / 8; Reading A matches 7 / 8. Over A ≤ 4, L1 is equivalent to "bound ⇔ |N − Z| takes its minimum at that A" (0 at A = 2, 1 at A = 3, 0 at A = 4). The only mismatch of Reading A (^{4}Li) arises precisely because its added clause "g ≤ 2" caps the neutron count but not the proton count; L1 is naturally mirror-symmetric and makes no such distinction.

(5) The limits of L1: not a fundamental law but a classification rule for the light region. Using the measured ground-state existence for A ≥ 5 as an out-of-sample set (26 entries, taken from whether each nucleus has a bound ground state; ^{8}Be is recorded as unbound, see the honest registration):

Interval	Entries	L1 hits	Reading A hits
A ≤ 4 (scorecard)	8	**8 / 8**	7 / 8
A = 5 (α + 1 region)	3	1 / 3	2 / 3
A = 6–9	11	8 / 11	4 / 11
A = 10–14	12	5 / 12	1 / 12
Total	**34**	**22 / 34**	14 / 34

L1 is clearly superior to Reading A (22/34 against 14/34), but both classes of mismatch show that it is not a fundamental law:

- **The A = 5 mismatches are closure effects, not asymmetry effects.** ^{5}He and ^{5}Li both have a = 1 (the minimum at that A); L1 predicts binding while both are measured unbound. No nucleus is bound at A = 5, so **any criterion depending on composition (A, Z) alone must mismatch at A = 5**; this is homologous to criterion ⑤ of 12.11 (partial rebalancing of closure degree) — the special closure of the α unit is not representable by a composition criterion.
- **Systematically too strict for A ≥ 8.** The |N − Z| ceiling allowed by measured bound nuclei rises with A (per-A ceilings in this sample: A = 6 → 2, 7 → 1, 8 → 4, 9 → 1, 10 → 2, 11 → 5, 12 → 4, 14 → 6). The fixed threshold |N − Z| ≤ 1 judges the neutron-halo/drip-line nuclei ^{8}He, ^{11}Li, ^{12}Be and ^{14}Be unbound, whereas all are measured bound — an inevitable consequence of the liquid-drop $(N-Z)^2/A$ picture, since **L1 lacks the A scaling**.

The scaled form $(N - Z)^2/A \le c$ was also checked: the $A = 4$ side requires $c < 1$ (the $(N - Z)^2/A$ of ^{4}H and ^{4}Li equals 1 and must be judged illegal), while the drip-line side requires $c \ge 2.571$; **the two constraints are incompatible**. Scanning $c \in \{0.25, 0.5, 0.8, 1, 2, 2.5, 3\}$ gives a best total of 22/34, on a par with L1. The $A = 4$ side must therefore be carried by the **closure effect**, not by an asymmetry scaling.

(6) Summary and revision of the boundary. The net results of this section are:

- **Negative**: the antisymmetric coefficient λ can neither be derived from nor fixed by SRE; it is merely an absorbable degree of freedom. The registration "λ is a fit" of 12.11(4) is thereby upgraded to a structural conclusion.
- **Positive**: the only contentful form of antisymmetry is the **existence criterion L1** (zero-parameter, mirror-symmetric), which matches the 12.8 scorecard at 8 / 8; hence **the "existence" question at $A = 4$ is decidable in this layer more precisely than the original registration allowed**.
- **Revision of the boundary**: the original statement of 12.9, "**the nuclear boundness for $A \ge 4$ (both existence and the numerical value of the binding energy) is not computable in this layer**", must be split:
  - **Numerical value of the binding energy**: not computable for $A \ge 4$ (the identity of 12.11(2) is unaffected by this section);
  - **Existence**: computable at $A = 4$ via L1 with zero parameters (8 / 8); not computable for $A \ge 5$ (the criterion degrades to an empirical approximation).

The boundary is thus **relocated** from $A \ge 4$ to $A \ge 5$, and its nature is **relocation rather than removal**.

Honest registration: (i) L1 and Reading A are both post-hoc readings, and the superiority of L1 consists only in "more hits with fewer clauses" (zero parameters, a single clause, mirror symmetry); (ii) in the $A \ge 5$ out-of-sample set, "bound/unbound" is taken as ground-state existence, and ^{8}Be is recorded as unbound because it is α-unstable (only 0.092 MeV), so changing that convention would affect 1 entry; (iii) the exactness of L1 over $A \le 4$ is consistent with the known fact that the light-nucleus region is dominated by α closure, and it cannot be claimed on that basis that L1 is a fundamental law; (iv) this section revises only the **existence** boundary; the impossibility proof of 12.11(2) regarding the numerical binding energy is unaffected.

12.13 Does the Nucleon–Nucleon Force Require the Electronic State? — Configuration-Level Localization of the Open Band

Question. A natural objection is whether considering the "nucleon–nucleon force" alone is incomplete, and whether the electronic state must be considered as well. This section rewrites that objection as a **configuration problem decidable within this layer**; all numbers are independently recomputed by `code/_sre_nucleon_electron_coupling.py`.

Semantic delimitation. SRE contains no "atomic electron cloud" object — the electrons of the mesoscopic layer are reassembled by the electron proper (the Q_3 coherent core) at a higher scale, and their effect does not enter nucleon assembly. Within **this layer**, the only decidable meaning of "electronic state participation" is the **open band**: the neutron state = opening one ring edge (Section 9.4), and that unclosed edge is exactly the "segment awaiting settlement" of the β-decay reading. The question to answer is therefore:

> Does the meeting of the **open bands of two nucleons** in the shared region yield a new structure beyond the shared-ring family?

(1) Preliminary measurement: the three rings are equivalent, so "the gap lies in ring 0" is a notation, not a fact. A previous statement had gone unchecked, namely that "the neutron gap can only lie in ring 0, otherwise an isolated vertex is left behind". Measurement refutes that statement: the automorphism group of the closed-state skeleton has order 36, and the corresponding ring edges `(L0j, L1j)` (j = 0,1,2) on the three rings **lie in a single orbit**; opening each ring individually gives V = 12, E = 17, T = 2, β_1 = 6, ρ = 5.302775638, λ_2 = 1.000000000, **all identical**, and none of the three rings produces an isolated vertex. Hence "the gap lies in ring 0" is a **choice of representative** within an equivalence class, not a distinguishable fact. This section accordingly makes all such choices explicit and re-runs them one by one (see (2)), confirming that no conclusion depends on the representative.

(2) Four two-body constructions, re-run for r = 0, 1, 2. Let S1 = variant A (the two ends of the open-state gap identified with the two ends of one complete ring edge of the closed state, Section 12.1); S2 = open-band attachment (the two points of strands 0 and 1 on ring r of the open body identified with the same two points on ring r of the closed body, all other vertices retained); S3 = shared ring (the three vertices of ring r of the two bodies merged, i.e. the k = 2 member of Section 12.5); S4 = open-to-open (two gaps mated at the same position). The results are:

Construction	V	E	T	β_1	ρ	λ_2	|Aut|
S1 = variant A	22	35	6	14	6.681330644	0.238442818	16
S2 = open-band attachment	22	35	6	14	6.681330644	0.238442818	16
S3 = shared ring (k=2, deuteron)	21	33	5	13	**6.000000000**	**2−√3**	48
S4 = open-to-open	22	34	4	13	6.236067977	0.238442818	16

All three values of r give a **digit-for-digit identical** table, and S1 ≅ S2 holds for every r.

(3) Verdict: S2 ≅ S1 — open-band attachment is not a new configuration. Measurement shows S1 and S2 to be **isomorphic** (all invariants equal term by term, |Aut| = 16 for both). The mechanism: ring r of the open body retains only two edges `(L0r, L2r)` and `(L1r, L2r)`; once `L0r` and `L1r` are identified with the same two points of the closed body, the closed body's ring edge `(L0r, L1r)` closes the gap, while the open body's reverse path simultaneously serves as a second path, so a **new ring forms automatically** with exactly the same β_1 increment as variant A. Having established S2 ≅ S1, the open-band attachment reading is doubly excluded by existing results: at the **structural layer** it is variant A rather than a new configuration; at the **pricing layer** variant A has already been quantitatively excluded by the deuteron binding energy (Section 12.2: were A to hold, B_d would be 2Δm_np = 2.5867 MeV, whereas the measured value is 2.2246 MeV, 16.3% higher).

(4) Positive result: the open band is already contained in the assembly law, appearing as the "shape of the gap" rather than as an "extra energy term". The decisive diagnostic lies in the S3 ledger. The claim counts of the three edges of the two-body shared ring are

```
(S0,S1) k=1 ;  (S0,S2) k=2 ;  (S1,S2) k=2        profile = (1,2,2)
```

where the edge with k = 1 is **precisely the locus of the open band**: all three edges of ring r of the closed body are present (each +1), the open body retains only two edges (claiming only the latter two), and the gap is closed by a complete edge of the closed body, so `(S0,S1)` is claimed by the closed body alone. Three correspondences follow:

- the **presence or absence** of the gap decides whether n-p can assemble (principle 1);

- the **position** of the gap decides that n-n is excluded automatically (the identity requirement of principle 1; measured S4 has E = 34, exactly 1 less than S1 — the ring edge that can never be closed);
- the **k = 1 singly-claimed edge** left after the gap is closed decides that the images of the two bodies are distinguishable (principle 2).

That is, the "electronic component" of the nucleon–nucleon force is **already contained in the assembly law**; it does not appear as an additive correction but as the **shape of the gap**. This also explains a fact that previously had only a phenomenological description: **the shared ring always has exactly one k = 1 edge** (the image of the open band), and this is precisely the structural origin of the statement in Section 12.5 that "the three points of the shared ring have strictly zero amplitude in the lowest excitation mode, so $\lambda_2 = 2-\sqrt{3}$ is independent of the body count" — among the three points participating in the identification, exactly one has a connection strength differing from the other two.

(5) The three A = 4 compositions remain a single isomorphism class. After the open band is included, the graphs of ^{4}He (2s+2s, fully closed), ^{4}H (1s+3s) and ^{4}Li (3s+1s) are still **pairwise isomorphic** (V = 39, E = 63, T = 9, $\beta_1 = 25$, $\rho = 7.909515966$, $\lambda_2 = 2-\sqrt{3}$, $|\mathrm{Aut}| = 2304$, all identical), differing only in the ledger: the shared-ring profiles are (2,4,4), (1,4,4) and (3,4,4) respectively. The conclusion of 12.11(1) is therefore unaffected by this section — **the open band produces no graph-level difference whatever in the shared-ring construction**; it changes only the ledger, and the ledger has already been falsified as a whole by the identity of 12.11(2).

(6) Conclusion and boundary. (a) Open-band attachment **does not constitute** the "third kind of SRE input" sought in 12.12, so the non-computability for A ≥ 4 is **not repaired** by this route; (b) it does, however, give a **new formulation of the source of failure**: the additive-profile pricing $B = \psi(\#p) + 2\psi(A)$ is falsified as a whole because it compresses the **shape** information of the open band into the count #p — the same conclusion as the identity of 12.11(2), from a different vantage point; (c) this section also **corrects a previously erroneous statement** ("the gap can only lie in ring 0") and shows that the correction leaves every existing conclusion intact.

Honest registration: (i) the conclusion of item (1) regarding the single orbit of the three rings applies only to the automorphism group of the **closed-state** skeleton; (ii) the "open band" is an extension of the β-decay reading of this paper (Section 5), and its ontological status is a model-internal construction not independently tested; (iii) all conclusions of this section are internal self-consistency constructions of SRE and do not constitute a derivation of the real nucleon–nucleon force.

12.14 Can the Neutrino Be Read as the k = 1 Singly-Claimed Edge? — Criterion Construction and Negative Verdict

Origin. The "downward extension" screening of Section 12.8 registered "neutrino = the k = 1 singly-claimed edge on the shared ring (Section 12.13(4))" as the easiest entry to register, with the first criterion: the ratio $\Delta m^2_{21}/\Delta m^2_{31} \approx 0.0307$ must be a **zero-parameter** function of some invariant of that edge. This section turns that criterion into an executable form and executes it; all numbers are independently recomputed by `code/_sre_neutrino_k1.py`.

(1) Gates and the spectroscopic requirement. The target is $R = \Delta m^2_{21}/\Delta m^2_{31} = 0.030697$ (PDG 2024 global fit, relative uncertainty ±3.2%, so the window is taken as $\varepsilon = 3\%$, tying the window to the precision used as per G4). Of the three gates, C1 (dimensionless) and C2 (no dependence on renormalization scheme or scale) pass; **C3 is the only gate** (there is only one independent dimensionless target). A more substantive constraint comes from spectroscopy: neutrino oscillations require **3 non-degenerate mass eigenstates ⇒ 2 independent non-zero gaps**; any candidate structure must supply them.

(2) The ledger route: the claim triple always has 2 distinct values. The shared-ring claim triples of six assemblies are computed directly (the ring edges are (S0,S1), (S1,S2), (S2,S0)):

| Assembly | Claim triple | #closed | Distinct values | Independent gaps |

|---|---|---|---|---|

| Two-body 1p+1n (= variant D / deuteron) | (1, 2, 2) | 1 | 2 | 1 |

| Four-body 2p+2n (⁴He) | (2, 4, 4) | 2 | 2 | 1 |

| Four-body 1p+3n (⁴H) | (1, 4, 4) | 1 | 2 | 1 |

| Four-body 3p+1n (⁴Li) | (3, 4, 4) | 3 | 2 | 1 |

| Three-body 2p+1n | (2, 3, 3) | 2 | 2 | 1 |

| Three-body 1p+2n | (1, 3, 3) | 1 | 2 | 1 |

⇒ the claim triple is **always of the form (#closed, k, k)** (k = body count), with **always 2 distinct values** ⇒ at most 1 degree of freedom. Note that this is exactly the ledger profile (#p, A, A) of Section 12.11 — **the claim numbers are the ledger** — so the failure of this route is **of the same origin** as the failure of the A ≥ 4 pricing layer.

(3) The graph route: all three-generation candidates are degenerate. Both natural "three-generation" readings are blocked by symmetry:

- **Three-ring reading** (taking the skeleton's three rings r = 0,1,2 as the three generations): exhaustive pairwise verification (`nx.is_isomorphic`) shows that the assemblies with the open state on any ring are **pairwise isomorphic** (true for both the two-body and the four-body case, 3/3 pairs all True), with invariants identical item by item (two-body: V21 E33 T5 $\beta_1$13 ρ=6.000000000 λ_2=2−√3) ⇒ the three generations are **strictly degenerate** ⇒ $\Delta m^2 = 0$.
- **Three-ring-point reading** (taking the local structure on the three shared-ring vertices {S0,S1,S2} as the three generations): the eigenvalues of two classes of 3×3 operators — the response sub-block $G_3 = (L^+)[S_0, S_1, S_2]$ and the Kron reduction $L_{\text{eff}} = L_{rr} - L_{ro}L_{oo}^{-1}L_{or}$ — **always have only 2 distinct values** (multiplicity pattern 2+1), for all six assemblies (two-body: [0.238095238, 0.238095238, 0.448979592] and [0, 4.2, 4.2]) ⇒ the spectrum degenerates to **two levels**, and **one of the two Δm^2 is identically zero**. This is not a "shortage of gaps" but an **extinction**.

(4) Mechanism (the key structural result of this round): the graph cannot distinguish the k = 1 edge. The **number of orbits of the automorphism group on the three ring points is measured to be 1** (for the two-body case with |Aut| = 48 the image distribution of S0 over S0:S1:S2 is 16:16:16; for the four-body case with |Aut| = 2304 it is 768:768:768) ⇒ the graph is **vertex-transitive** on the three ring points. Mechanism: the assembly graph is the **union** of the bodies' edge sets, and it does not record **claim multiplicity** (the same origin as "the product graph does not remember the partner state" in Section 12.4). ⇒ **the k = 1 edge exists only in the ledger, not in the graph.** Correspondingly, any ring-invariant symmetric 3×3 operator must be of the form $a \cdot I + b \cdot J$ ⇒ its spectrum is always 2+1 degenerate.

This item also **corrects a presupposition of the first draft of this section** (which assumed that the two ends S0, S1 of the k = 1 edge formed an orbit of their own with S2 separate, i.e. 2 orbits); the measurement gives 1 orbit, and after correcting this and re-running, the **direction of the negative verdict is unchanged but stronger** (from "2 orbits" to "vertex-transitivity").

(5) The zero-parameter candidate pool: no effective hit. Sixteen classes of zero-parameter readings (spectral ratios, ledger ratios, effective-resistance ratios, gap ratios and Δm^2-type ratios of the 3×3 operators) are constructed on the six assemblies, giving **96 numbers** in total, each compared with R:

| Check | Result |

|---|---|

| Hits at $\varepsilon = 3\%$ | **1**: `1/E = 1/33 = 0.030303` (deviation 1.28%, E being the edge count of the deuteron assembly) |

| Hits at $\varepsilon = 1\%$ | **0** |

| MC null hypothesis A (randomized target, 2×10⁴ draws) | probability that a random target is hit by any candidate, p_A = **0.166** ⇒ expected hits 0.17 (observed 1) |

| MC null hypothesis B (200 random graphs with the same degree sequence) | hit rate of random graphs **0.385** > 0.010 for the SRE assemblies |

⇒ the sole hit has no mechanistic connection with neutrinos, and the hit rate of the SRE assemblies is **lower** than that of random graphs ⇒ **no structural signal** (a direct application of the lesson of Section 12.8: cross-hits must first pass a whole-set hit-rate check).

(6) Conclusion and boundary. ④ is judged negative (the criterion has been constructed and executed). The k = 1 singly-claimed edge is indistinguishable at the **graph** level (vertex-transitivity ⇒ degeneracy) and has only 1 degree of freedom at the **ledger** level (always 2 values); this matches neither the **symmetry** nor the **count** required by the neutrino's "2 independent non-zero gaps + 3 non-degenerate states", and of the 96 zero-parameter candidates none is an effective hit. The two natural routes each strike one **already-established barrier** of this project: the graph route strikes the vertex-transitivity of Sections 12.4/12.13, and the ledger route strikes the identity of Section 12.11(2) — this is the cleanest conclusion of this section.

Honest registration: (i) the negative verdict is **conditional**: it holds under the reading "k = 1 edge = the neutrino proper" together with "the three ring points (or their Kron reduction) are the mass operator"; other readings (weighted spokes, a non-additive ledger) would violate C3 (free parameters ≥ number of targets) and hence be unfalsifiable, and must not be used to press ahead (the same treatment as Section 12.9 gives CKM). (ii) The negative verdict **does not depend** on any numerical coincidence: the blocking comes from **isomorphism/orbit determinations** and is therefore **distribution-free**, holding even if the window is made arbitrarily wide. (iii) The repair required by ④ is a **zero-parameter rule that breaks the vertex-transitivity of the ring points**, which must simultaneously supply "the third non-degenerate level" and "the ratio of the two gaps = 0.0307"; this is **the same single missing piece** as ① "generation → (n,w)" and ② "discrete class → continuous angle" (SRE lacks the assignment rule "**integer invariant → real measured value**" = the L4 lock of Section 3) ⇒ the three pending tasks merge into **four**.

12.15 The Mass-Depth Assignment Rule (L4): Criterion Construction and Negative Verdict - Turning the Merge of the Four Downward Branches into a Single Test

Origin. The downward screening of Section 12.8, the four negative verdicts of Sections 12.9-12.14, and the species-dimension gap exposed in 13.3 all point repeatedly to one and the same missing piece: SRE lacks an assignment rule from "integer invariant -> real measured value". Its formal position in the project is the **L4 mass-depth equation** of Section 7.1, $m_p/m_e = \Psi(G_N)/\Psi(G_e)$ - in which Ψ has never been defined; Section 3 further requires it to satisfy **homogeneity** (the two layers share one and the same Ψ, i.e. "two graphs share one lock"). This section turns L4 into a **pre-registered, falsifiable** test and executes it; all numbers are independently recomputed by `code/_sre_l4_assignment.py`.

(1) Pre-registered criterion (fixed before any number is produced; no adding items, changing the window or swapping targets afterwards). (i) The set of primitive invariants (layer-agnostic, ten): $V, E, T, \beta_1, |\mathrm{Aut}|, \rho_A$ (adjacency spectral radius), $\lambda_2(L)$, ρ_L (largest Laplacian eigenvalue, i.e. the project's usual ρ), τ (number of spanning trees), En (graph energy). (ii) The functional class Ψ: monomials $\prod_i p_i^{a_i}$ with exponents $|a_i| \le A$ and support ≤ 3; main analysis $A = 2$, robustness scan $A = 1, 2, 3$. (iii) Both electron-layer representatives are admitted: configuration A takes $G_e = Q_3$ (the electron basal body, P2), configuration B takes $G_e = M_{60}$ (the α carrier). (iv) Three targets: T_1 (the design target, i.e. L4 itself) $m_p/m_e = 1836.15267343$; T_2 (out-of-sample) $m_d/m_p = 1.999007745$; T_3 (out-of-sample) $m_n/m_p = 1.0013784193$. (v) Decision tolerance $\varepsilon = 1\%$ - **deliberately wider than the precision the framework has already demonstrated (0.165% on the nucleon layer)**, so that the negative verdict cannot be blamed on the window. (vi) Success condition: there exists **one** Ψ with zero free parameters that brings

all three targets within ε. (vii) Null hypothesis: replace each of the four graphs by a uniform connected random graph with the same (V, E).

(2) Graphs and invariants: the construction passes all 12 cross-checks. The four graphs are built under the project's existing conventions (Y_3 closed/open states, the two-body shared-ring k = 2 deuteron assembly, and the electron graph). All cross-checks agree: $\rho_L(\mathrm{Y}_3) = (7+\sqrt{13})/2 = 5.302775638$; after opening one ring edge, λ_2 and ρ_L are **completely unchanged**; the deuteron assembly has $\lambda_2 = 2-\sqrt{3}$, $\rho_L = 6$ and $|\mathrm{Aut}| = 48$. Hence the negative verdict of this section comes from the test itself, not from a construction or convention error.

(3) The decisive check: the reach bound (distribution-free). For any Ψ we have $\Psi(G_N)/\Psi(G_e) = \exp(\sum_i a_i \log r_i)$ with $r_i = p_i(G_N)/p_i(G_e)$, so the largest ratio attainable within the pool has the **closed-form upper bound** $\exp(A\sum_i |\log r_i|)$ taken over the largest terms. Measured:

Config	A	reach (support <= 3)	reach (**full support**)	< 1836.15 ?
A ($G_e = Q_3$)	1	4.80	22.61	yes (excluded)
A	2	23.07	**511.06**	**yes (excluded)**
A	3	110.78	1.155x10^4	no
B ($G_e = M_{60}$)	2	3.272x10^5	3.749x10^9	no

With the **electron basal body** Q_3 as G_e, the **full-support** bound for $|a| \le 2$ is only **511.06**, i.e. **short of the target 1836.15 by 0.555 orders of magnitude** (23.07 for support <= 3, short by 1.90 orders) => **no** Ψ in the pool can produce m_p/m_e. This conclusion is **independent of the distribution and of the support size** (hence immune to the objection that the support limit was taken too small) and needs no Monte Carlo. Only by relaxing the exponent range to $|a| \le 3$ does the ceiling (1.155x10^4) clear the target.

(4) The direct refutation of "sharing one lock". Configuration B's reach is unconstrained, so it enters the full-pool test:

A	pool size	T_1 hits (1%)	T_2 hits (1%)	T_3 hits (1%)	**joint hits (1%)**	joint hits (3%)
1	576	0	7	46	**0**	0
2	4064	1	18	191	**0**	0
3	13152	4	36	494	**0**	0

More substantive still: **the best Ψ differs from target to target** - T_1 is attained by `lam2L^2 rhoL^2 energy^-1` (deviation 0.1775%), T_2 by `E^1 b1^2 energy^-2` (0.0884%), and T_3 by `E^-2 rhoA^2 energy^1` (0.0015%). => **There is no single lock governing all three**, which is precisely the **direct negative form** of the L4 requirement that "two graphs share one lock". Moreover **the joint hit is 0 for $A = 1, 2, 3$ and for both the 1% and the 3% window** => the negative does not stem from a narrow exponent range.

(5) The performance of the project's only calibrated functional, Pi1. $\Pi_1 = \lambda_2/\rho_L$ is itself in the pool (exponent vector (1, -1)). Its prediction for m_p/m_e is **0.5657** (configuration A) and **25.84** (configuration B), deviating from 1836.15 by 99.97% and 98.59% respectively, **both short by 1.85 orders of magnitude**. Note that in configuration B this ratio equals exactly $\Pi_1(\mathrm{Y}_3)/\alpha$ - the quotient of the two members of the project's most successful cross-layer chain (see unresolved item 8 of Section 16). => **The project's best lock, used in its native form, still falls short of the target by 1.85 orders of magnitude.**

(6) Null hypothesis and positive residue. The MC null (407 valid trials, uniform connected random graphs with the same (V, E)): hits per trial by target are T_1 0.40, T_2 21.46, T_3 94.91, and joint 0/407 => the single-target hit count of the real configuration is **of the same order as for random graphs**, i.e. no above-null signal (the slightly high T_3 is explained by the fact that a real graph pair (closed/open states) is more similar than a random pair, so Ψ ratios sit closer to 1; and T_3 is itself a **weak target** - any Ψ nearly invariant across the open/closed transition gives exactly 1, and 1 already lies within 0.14%).

Positive residue (control). The same Π_1 machinery holds on the **difference ratio of two states within a layer** - $\alpha \cdot \Pi_1(\mathrm{Y}_3) = 1.376158 \times 10^{-3}$ against the measured $(m_n - m_p)/m_p = 1.378419 \times 10^{-3}$, a deviation of **-0.164%** (with $\alpha = \Pi_1(M_{60})$ deviating from the measured α by 1.45×10^{-5}). => **the cross-layer lock governs the difference within a layer, but not the overall scale of a layer** - this is the concrete form of "SRE fixes the shape, not the value" for **inter-layer mass ratios**, and it explains why both λ in 12.12 and w(k) in 12.11 fix only the "shape".

(7) Verdict and three independent grounds. L4 is judged negative (the criterion has been built and executed):

- **(i) reach bound (configuration A, $|a| \leq 2$, distribution-free)**: the full-support bound 511.06 < 1836.15 => no Ψ in the pool can work;
- **(ii) no single lock**: the best Ψ differs from target to target, and the joint hit is 0 for $|a| \in \{1, 2, 3\}$ and for both the 1% and the 3% windows;
- **(iii) hidden degree of freedom**: the electron-layer representative (basal body Q_3 vs carrier M_{60}) is not fixed by the framework, and the reach verdict **flips** between the two configurations (A excluded, B not) => "which graph represents the electron" is itself a free parameter => **even if a hit appeared, it would be neither zero-parameter nor falsifiable (C3 violated)**.

(8) Boundaries. (a) **Class scope**: the monomial class is only the general class of "integer powers of products of standard invariants"; it excludes additive, trigonometric and combinatorial Ψ. The negative verdict is confined to this class. (b) **Exponent range**: the $|a| \leq 2$ reach exclusion is hard; for $|a| = 3$ the ceiling clears the target, and there the verdict rests only on (ii) and (iii) (listed separately). (c) **The $|\mathrm{Aut}|$ gap**: for $V > 22$ it is not computable under the same rule, so configuration B's pool excludes $|\mathrm{Aut}|$. (d) **Nature of the out-of-sample targets**: T_2 and T_3 use only nucleon-layer graphs, so their layer-crossing mechanism differs from that of T_1 and they are independent measurements; but T_3 (m_n/m_p) is a weak target and should carry less weight than T_1 and T_2. (e) **Net contribution of this negative verdict**: it turns the merge of the "four downward branches" (the lemmas numbered (1), (2) and (4) plus the species dimension of p_{env}) into a **single executed test**, and shows that the missing piece **cannot be supplied by switching to a larger zero-parameter pool** - what is missing is three mutually independent locks. (f) **The dual scope of the reach bound (added in v2.0; two independent corrections)**: the reach bound on which route (i) rests is **related both to the pool size and to the representative**, and therefore **cannot be promoted to a law**. First, the **pool-size correction** (`code/_sre_l4_pool_extend.py`): expanding the original invariants from 10 to 29 raises the full-support reach from 511.06 to 8.65×10⁶, i.e. the validity of (i) **depends on how the pool is truncated**; moreover the survival reading for support ≤ 3 (91.209%) agrees digit for digit with the grid measurement (95.0326%), and the 5% quantiles of the MC null are 0.1995% / 0.1822% (both below the multiple-comparison level of the random graphs). Second, the **representative correction** (`code/_sre_collapse_in_nucleon.py`): taking the electron representative along the folding tower ($K_4/Q_3/Q_4/Q_5/M_{60}$, with the pool defined term by term as in this section), only Q_3 **is blocked by reach (511.1)**, while the reach of the remaining configurations is 3.8×10⁵–3.7×10⁹ ($\gg 1836.15$) ⇒ Q_3 **is precisely the worst point in this family**. The correct statement of route (i) is therefore "**for the representative Q_3, within $|a| \leq 2$**, blocked by reach", **not a universal law**; this **does not affect the negative verdict of this section**, because even if the reach obstacle is removed entirely (switching to the unfolded $K_4/Q_4/Q_5/M_{60}$), **the joint hit is still 0 and the best same-lock deviation is still ≥ 19.84%**, so the negative verdict is independently carried by routes (ii) and (iii). A further set of exact laws on the folding tower is registered here as well (for reference in Section 14): $|\mathrm{Aut}(Q_k)| = 2^k \cdot k!$, the number of distinct Laplacian values is always $k + 1$ (which, against the $\approx V$ of random graphs, is "log V level"), $\Pi_1(Q_k) = 1/k$ exactly, and both operators D and BDC **preserve vertex-transitivity** (0 counterexamples over a battery of 21 graphs) ⇒ **the number of Aut orbits of a tower member is always 1**.

---

13. Theoretical Constraints

13.1 Special Binding of Ions/Metals: Theoretically Predictable

The additivity law fails for ion/metal pairs (alkali metal/alkaline earth halides) with a deviation of −12.3% ± 7.6% in the test. **This failure is not an empirical accident, but a theoretical necessity that can be predetermined by the binding mode**:

- **Covalent non-metals**: Electron topology is bound to the **local shared coherent channel** between A–B (two-body local);
- **Ions/metals**: The binding of electron topology is **lattice-wide** — charge integral transfer or global delocalization, and the channels are not "shared" but "closed/transferred".

The distance of ion pairs contracts below the additivity line ("crossing the additivity line"), which is essentially caused by **channel topology switching**, belonging to a different ontological interval from the covalent interval. This empirical classification can be upgraded to a theoretical predictive criterion: **whenever the electronegativity difference is sufficient to trigger channel closing/transferring, the additivity law should fail**.

13.2 Current Is an Emergent Quantity Rather Than Continuous Electron Movement (Important Inference)

Derived directly from the SRE distance ontology, it is a unified inference of nucleon/electron ontology:

- Premise 1: Distance is not an a priori coordinate, but coherent degenerate bookkeeping;
- Premise 2: Electrons are operator fixed points on coherent cores, not "entities moving along trajectories in space".

Inference: **Macroscopic current cannot be understood as continuous transport of electron particles along wires**. Current is the **projected change of coherent degree bookkeeping of conductor electrons along the macroscopic conduction pathway**, belonging to the statistical emergent layer, along with electric fields, magnetic fields, gravitational fields, quark observations, and quantum entanglement — all are statistical projections of coherent core coupled with environment.

Homologous to 13.1: whether it is covalent sharing, ion transfer, or conduction current, the observable "distance/transfer" phenomena are all **projections of channel topology (establishment/closing/transfer of coherent channels)**, rather than positional movement of entities. The explanation layer is changed, not the numerical layer: macroscopic equations such as Ohm's law remain effective descriptions of the emergent layer and are not affected.

13.3 The Exponential Decay Law: Derived from the SRE Axioms, and a Test of Its Two-Factor Decomposition

Origin. Internal registration shows that "half-life / exponential / memoryless" has **zero hits** throughout the document suite: although this framework repeatedly uses the reading "the open band = the segment awaiting settlement", it never wrote down the form of the decay law. This section fills that gap (**derivation**, not fitting) and tests the two-factor decomposition that follows. All numbers are recomputed by `code/_sre_decay_law.py`.

(1) Derivation (three steps, using only the axioms of Sections 5.2 and 9.4). Axiom S1 (dormancy is the default state): the three ring edges of a nucleon are dormant by default, and β decay = one of them being removed. Axiom S2 (removal is a rare perturbation): the probability that a given dormant edge is removed per unit time is $p \ll 1$, and it is independent of the other edges and of that edge's own history (independent + stationary). The waiting time of a single dormant edge therefore follows a **geometric distribution**, whose continuum limit is the **exponential**:

$$P(T > t) = (1-p)^{t/\Delta t} \to e^{-pt}, \qquad T_{1/2} = \frac{\ln 2}{p}, \qquad \tau = \frac{1}{p}, \qquad T_{1/2} = \tau \ln 2,$$

and equivalently satisfies **memorylessness** $P(T > t+s \mid T > t) = e^{-ps}$. Point of consistency with SRE itself: SRE's graph **does not record history** (12.4 "the product graph does not remember the partner state"; 12.13(1) "the gap lying in ring 0 is notation rather than fact") ⇒ the state is memoryless ⇒ the waiting time is memoryless ⇒ exponential. This step is registered as **homological support**, not a strict derivation — a graph not recording the "partner state" is not the same as not recording the "instant".

(2) Monte-Carlo verification and exclusion of the power law. With $p = 0.05$ and 4×10^5 samples: (i) the maximum deviation of the survival curve from e^{-pt} is **1.03×10⁻³** (241 grid points); (ii) the half-life identity: empirical median 13.859835 against $\ln 2/p = 13.862944$, a relative deviation of **0.0224%**; (iii) memorylessness: conditional probability 0.223705 against $e^{-ps_0} = 0.223130$. Power law excluded: a log-log linear fit to the survival curve gives slope −1.5995 and $R^2 = 0.781$, whereas the exponential fit gives $R^2 =$ **0.999999**; more importantly, the power-law criterion of "constant slope" is falsified — the piecewise slope drifts from **−0.570** over the first third to **−5.013** over the last third ($|\Delta| = 4.443$).

(3) Structural factor: the number of dormant edges. If a body has N equivalent dormant edges, the rate of "any one being removed" is Np ⇒ $T_{1/2} \propto 1/N$ (zero-parameter; MC verification for $N = 1, 2, 3, 6$ agrees to better than 0.4%). A single (free) body has 3 dormant ring edges ($N = 3$); in the shared-ring assembly only **1 k = 1 edge** remains (12.13(4)) ⇒ $N = 1$. If N were the only graph factor, then a "neutron inside a nucleus" should be 3 times slower than a "free neutron" — registered as **pending, not adjudicated here** (the effective in-medium neutron lifetime is a model-dependent derived quantity, beyond this layer).

(4) Two-factor decomposition: the only non-trivial content = the species-cancellation law. Writing p as $p = p_{\text{graph}} \cdot p_{\text{env}}$ is itself an identity; its only non-trivial content is "which factor carries what". This yields a **zero-parameter** species-cancellation law — for the same nuclide in environments I and II,

$$T_{1/2}(\mathrm{I})/T_{1/2}(\mathrm{II}) = p_{\text{env}}(\mathrm{II})/p_{\text{env}}(\mathrm{I}) \qquad (p_{\text{graph}} \text{ cancels exactly}),$$

so the "stripping enhancement factor" $E \equiv T_{1/2}(\text{neutral})/T_{1/2}(\text{bare}) = p_{\text{env}}(\text{bare})/p_{\text{env}}(\text{neutral})$. **If** p_{env} **depends only on the electron environment** (SRE has no energy ontology ⇒ p_{env} carries no nuclear-level-structure information), **then** E **must be independent of the nuclide** — a prediction that requires no fitting and can be falsified in one sentence.

(5) Test: the prediction is falsified. Three L3 (inner electrons fully removed) β⁻ nuclides (all published measured values; no fitting in this section):

Nuclide	Q/keV	T½ (neutral)	T½ (bare)	E	$\log_{10}$E
^{187}Re	2.467	4.16×10¹⁰ a	32.9 a	1.264×10⁹	+9.10
^{163}Dy	2.560	stable (>10¹⁵ a)	47 d	≥7.771×10¹⁵	≥+15.89
^{241}Pu	20.800	14.3 a	4.2 d	1.244×10³	+3.09

- ^{187}Re vs ^{163}Dy: **same mode (bound-state β⁻), nearly the same Q (2.467 and 2.560 keV)**, yet E differs by **≥ 6.79 orders of magnitude** ⇒ "E varies only with the graph / is independent of Q" is falsified;
- ^{187}Re vs ^{241}Pu: same mode (β⁻), E differs by **6.01 orders of magnitude** ⇒ "E is independent of the decay mode" is falsified;
- **the failure cannot be repaired by the graph factor**: in the expression for E, p_{graph} has already cancelled exactly, so the species dependence of E can only be attributed to p_{env} **itself** ⇒ p_{env} **is not a "pure environment quantity"**; it must contain species-specific atomic/electronic-configuration information, whereas the p_{env} on the SRE side is merely a scalar (the "s-electron reachability at the nuclear site" of the three layers of Section 10).

(6) Positive residue: the model can fix the "ordering", not the "value". Ordered by "the margin between Q and the final-state electronic-configuration threshold", E decreases monotonically: ^{163}Dy (below threshold ⇒ decay forbidden) → ^{187}Re (just above threshold ⇒ tiny phase space) → ^{241}Pu (far above threshold ⇒ ample phase space), holding **3/3** (with ^{163}Dy a lower bound). Note however that **this monotonicity does not hold with respect to Q itself** (Re and Dy have almost the same Q yet differ by ≥ 6.8 orders of magnitude, see (5)), and the ordering variable itself contains atomic electronic configuration, which the SRE side does not have ⇒ this section treats it only as **qualitative support**, not a quantitative conclusion.

(7) Recheck and refinement. Recomputing the three-layer medians registered in Section 10 from half-life ratios ($\log_{10}$ of the relative fraction): L3 = +9.10, L2 = −2.09, L1 = −4.30, equivalent to **+11.10 / −0.09 / −2.30** in percentage terms, **agreeing digit for digit** with the registered +11 / −0.09 / −2.3; the span L3 − L1 = **13.40 orders of magnitude** (registered ~13). One **refinement**: the value 10^{14} % previously registered for ^{163}Dy was a conservative estimate; the **lower bound** computed directly from the half-life ratio is ≈8×10^{17} % (an underestimate of about 3.9 orders of magnitude); this refinement does not change the L3 median (which is set by ^{187}Re), so the conclusions of Section 10 are unaffected.

(8) Verdict and boundary. Positive result: the exponential decay law has been **derived** from the SRE axioms (closing the gap registered in Section 10), and the derivation **does not require** SRE to supply the numerical value of p — it gives only the form of the law ($T_{1/2} = \ln 2/p$, necessarily exponential rather than a power law), in agreement with nuclear-physics observation. **Negative result**: the strong reading of the two-factor decomposition (p_{env} = a pure environment quantity) is falsified by measurement, and cannot be repaired by the graph factor.

Honest registration: (i) **data-reliability grading**: ^{187}Re (Bosch et al. 1996) and ^{163}Dy (Jung et al. 1992) are primary literature; ^{241}Pu is a secondary review and not independently verified — **the decisive pair uses Re vs Dy**, so the conclusion does not depend on ^{241}Pu. (ii) **Layer boundary**: the L3 data are all heavy nuclei (A = 163/187/241), whose SRE graphs are **unavailable** in this layer (A ≥ 5 not computable, Section 12.11) ⇒ this section is a **decomposition test at the phenomenological layer**, **not** a numerical derivation internal to SRE; this must be registered separately from the derivation of (1). (iii) **Refinement of the reading of Section 10 (new in this section)**: the control parameter of the layering is refined from "the reachability of the s electron at the nuclear site" to "**the relative sensitivity of the s-electron density at the nuclear site to the environment**" — the latter contains both an **environment term** and a **species term** (^{7}Be has only 4 electrons ⇒ its innermost shell is its outermost, so its species term is the largest); the layering itself (L3 ≫ L2 ≫ L1, span 13.40 orders of magnitude) is unaffected. (iv) **Attribution**: the gap in the negative result (the species-specific dimension of p_{env}) is of the same type as the other four in this project — "SRE fixes only the 'form' (law, order, layering), not the 'value'" — this is the **sixth occurrence** of that regularity, and it may deserve promotion to theorem level. (v) This section gives the "exponential law" as a **structural inference** for the first time, but its premise S2 (independent + stationary) is a model axiom that has not been independently tested.

---

14. Observational Perspective: the Boundary between the Dimension-Free SRE World and the Three-Dimensional World

Origin. All the checks of the preceding thirteen chapters, whether positive or negative, finally land on one and the same position: "SRE fixes only the **shape** (laws, order, layering), not the **value**". This section adds no new physical test; instead it **raises this recurring experience into an explicit clarification of observational perspective**: it **writes out, layer by layer,** the boundary between the "dimension-free SRE world" and the "three-dimensional world", and gives two theorem-level supports. All numerical values are quoted from the recomputation scripts archived with this paper; this section itself introduces no new fitting parameter.

(1) The three-layer decomposition: topology / ratio / geometry. For any object generated by SRE (taking the "shape" of a water molecule as the example), its description splits into three layers, each with a part that is **provided for free** and a part that **must be paid for**:

| Layer | Object | Provided free by SRE | Input that must be paid |

|---|---|---|---|

| **L0 topological layer** | graph $G = (V, E)$ | integer invariants: degree sequence, Laplacian / adjacency spectrum, number of automorphism orbits | — |

| **L1 ratio layer** | dimensionless ratios, e.g. $(1, 1, t)$ | the **form** of the ratio | **the metric D** (which fixes its **value**) |

| **L2 geometric layer** | angles, coordinates | none | **the map** (Euclideanity / curvature / scale) |

- **The L0 realization**: the pure bond topology of a water molecule is simply P_3 (O bonded to two H). The measured intrinsic quantities are: degree sequence $(2, 1, 1)$, Laplacian spectrum $(0, 1, 3)$, adjacency spectrum $(-\sqrt{2},\ 0,\ \sqrt{2})$, and **2** automorphism orbits; adding one H···H edge gives K_3, with degrees $(2, 2, 2)$, Laplacian spectrum $(0, 3, 3)$, and orbit count 1. **In this layer everything is an integer or an integer ratio, with neither ratios nor angles.** The root cause is that a graph-theoretic object **carries no metric** — length, distance and angle all belong to metric structure and must be assigned from outside. Taking the Laplacian of P_3, for instance, $\lambda_2 = 1$ and $\rho = 3$ are both integers, and their ratio $1/3$ is a ratio, but **no ratio is a length or an angle**.
- **The L1 realization**: the ternary ratio. Three atoms give three pairwise distances, which upon normalization read $(1,\ 1,\ t)$, where $t = r(\mathrm{H}\cdots\mathrm{H})/r(\mathrm{O}{-}\mathrm{H}) = 2\sin(\theta/2)$. **This is the formal landing point of the reading "what is exhibited should be a ternary ratio"**: it is dimensionless and independent of any coordinate or ambient space, and is therefore compatible with the output type of SRE. Note in particular that t is independent of r (an identity of the isosceles triangle), so that "adding a non-bonded term" and "the angle is 104.5°" are **informationally equivalent** at n = 3 — it is not a prediction but a **rewriting** of the angle into a distance (of the same type as "zero residual = no predictive margin" in 12.6).
- **The L2 realization**: the angle. $\theta = \arccos(1 - t^2/2)$, the **mapped output** of L1.

⇒ The native products of SRE stop at L1. The three-layer structure is exactly where the boundary between the "dimension-free SRE world" and the "three-dimensional world" lies: the boundary is not in some formula, but at the **two tolls**.

(2) The two tolls: the theorem-level formalization of "fixing the shape but not the value". The honest registration of 13.3(8) once recorded "fixes only the shape, not the value" as its **sixth occurrence, to be considered for promotion to theorem level**; this section splits it, by mechanism, into **two** distinct tolls:

- **L0 → L1: pay the metric D.** When only the bond topology is supplied, H_2O is inverted to **collinear** (the H–O–H path length equals exactly the sum of the two O–H segments, so the triangle degenerates, $180.0000°$); the Fe_4S_4 cubic cage is distorted in the same way ($r = 0.946$). What has been proved is **D ⇒ geometry** ($r = 0.95$–1.00); what has **not** been proved is topology ⇒ D. ⇒ **D is an input, not a product.** Moreover this can be made into a **necessity proof** (the non-circular part): the two H's are equivalent ⇒ any symmetric bond weighting can only give $180.0000°$; adding electron leaf nodes is likewise ineffective (the sub-metric is unchanged term by term, and the minimum eigenvalue of the metric Gram matrix of the four-point graph is $= -0.25 < 0$, hence non-Euclidean) ⇒ **there must exist a direct H···H non-bonded relation**. This is the **falsifiable structural prediction** given by this paper.
- **L1 → L2: pay the map.** This is given by the two theorems of (3): geometry is not a function of L1 alone, but of L1 **together**

with the map.

⇒ This gives a unified explanation of the negative verdicts of the preceding chapters: **every quantity SRE has passed so far is a dimensionless ratio** (Π_1, λ_2/ρ, $\Pi_1(Q_k) = 1/k$, $\lambda_2 = 2 - \sqrt{3}$, the number of Aut orbits, $|\mathrm{Aut}|$ ratios), **while every negative verdict occurs exactly when a dimensionful value is required to be assigned to some structure** (L4 assigning mass, the absolute value of α, the species dimension of p_{env} in 13.3). "Ratio vs value" and "ratio vs geometry" are **two cross-sections of one and the same seam**.

(3) The two theorems: geometry is a property of the map, not of the graph. Both are **theorem-level** (constructive, distribution-free), so they are **immune to the "numerical coincidence / multiple comparisons" objection** (compare the criterion discipline set out in 7.6 and 12.6).

Theorem 1 (angle ∉ functions(graph)). Take **the same graph** P_3 and assign two admissible metrics $D(t)$: $d(\mathrm{H}_1, \mathrm{O}) = d(\mathrm{O}, \mathrm{H}_2) = 1$, $d(\mathrm{H}_1, \mathrm{H}_2) = t$. Measured: $t = 2.0 \Rightarrow 180.0000°$; $t = 1.5811 \Rightarrow 104.4775°$; $t = 1.2 \Rightarrow 73.7398°$. The degree sequence, Laplacian spectrum, adjacency spectrum and orbit count are **identical term by term** across the three. ⇒ angle $= f(D)$, so the angle is **not** a graph invariant.

Theorem 2 (same ratio, different map ⇒ different angle). Fix **the very same** ternary ratio $t_0 = 2\sin(104.4776°/2) = 1.581140$ and place it into three constant-curvature maps, scanning the **absolute scale** s (in units of the bond length, with curvature radius set to 1):

Map	Apex angle θ	Remark
Euclidean $K = 0$	**104.4776°**	constant for any scale s (= similarity)
Spherical $K > 0$	104.57° → **145.10°**	common window $s \in [0.1,\ 1.6]$; degenerate for $s \gtrsim 1.8$ ($\lvert\cos\rvert > 1$)
Hyperbolic $K < 0$	104.39° → **86.67°**	same window; monotonically decreasing with scale

Here both the spherical and the hyperbolic rows are scanned over **the same scale window** $s \in [0.1,\ 1.6]$ (s in units of the bond length, curvature radius set to 1), while the Euclidean row is constant for any s.

⇒ The same ratio gives an apex angle in the interval about $[86.7°,\ 145.1°]$; $104.4776°$ is merely the member that is "flat". **A curved space has no similarity** (the angle also depends on the absolute scale), so the one-to-one correspondence "ratio → angle" is **itself a property of Euclidean similarity**. ⇒ **Geometry is a property of the map, not of the graph** — this is exactly the theorem form of "only the map has geometric relations".

(4) The experimental side: only three things are measurable. Corroborating the above layering is a fact on the experimental side. In real measurement **no instrument ever reads out a "bond angle" or an "orbital" directly**: microwave / infrared / visible / ultraviolet spectroscopy gives only **spectral line frequencies (= energy differences)**; Zeeman / Stark give only **degeneracy multiplicities and splittings**; X-ray / electron diffraction, liquid microjets and COLTRIMS give **electron density distributions**.

The water "104.5°" is inverted from exactly this: rotational spectrum → rotational constants $A, B, C \to I = K/(A, B, C) \to$ **the model step** (rigid rotor + rotation-vibration / anharmonic corrections + isotopic substitution) → (r, θ). This step is not a direct read: doing a rigid-rotor inversion directly from Herzberg's A_0, B_0 gives $(0.9561\ \text{Å},\ 105.1232°)$, differing from the equilibrium geometry $(0.9578\ \text{Å},\ 104.4776°)$ by $\Delta\theta = 0.6456°$ — **this difference is the magnitude of the "model input"**. The "angle" of the same molecule is also not unique under different averaging conventions: $103.9°$ (r_e 1D anharmonic approximation), $104.478°$ (equilibrium r_e, Hoy & Bunker 1979), $104.50°$ (ground-state average $\langle r \rangle$), $105.12°$ (rigid-rotor inversion of A_0, B_0). ⇒ **"104°" is an**

inversion quantity carrying a ±0.5° convention dependence, not a directly read number. Two easily misused points must also be registered together: first, a project script once used a ternary triple $(0.9584,\ 1.51383,\ 104.474)$ that **mixes averaging conventions** ($0.9584\ \text{Å}$ is of r_e magnitude while $1.51383\ \text{Å}$ is of r_0 magnitude), whose closed angle is only $104.3279°$, differing from the stated value by $0.146°$, and which **must not be called "self-consistent"**; the consistent triple is $(0.9578,\ 1.5144,\ 104.4776)$. Second, the equilibrium angle happens to satisfy $\cos\theta_e = -0.250001$ (exact to 10^{-6}), so the normalized $t \approx \sqrt{2.5}$ — this is a **constructive coincidence**, of the same kind as "$\Pi_1(Q_k) = 1/k$ on the tower is about $1/137$ at $k = 137$", and **must not be used as evidence**.

By the same token, **a single orbital is unobservable** (invariant under unitary / gauge transformations): the gauge-invariant density on the graph side (the diagonal entries P_{ii} of the projection) is constant along each Aut orbit, so the shape resolution is exactly the **number of orbits**; and the calibrated graph of this project has orbit count 1 ⇒ uniform density, zero shape information ⇒ the **strong sense (zero-parameter emergence) of the electron orbital fails while the weak sense (Aut orbits) holds**. In addition, the $2\ell + 1$ degeneracy ladder is, upon examination, **not any natural graph ladder** (none of 12 candidates matches), so that "the ladder is not intrinsic while ratios are".

(5) The other side of the boundary: how the three-dimensional world is entered. The preceding subsection explained **how** "three dimensions" is **paid in**; the other half is from which end three dimensions is **selected**. This project measures (`code/_sre_dimension_selection.py`) that three mutually independent stability predicates **point simultaneously to $d = 3$ with zero parameters**:

- **Finite resistance**: the leading term of the lattice voltage field is r^{2-d} (term by term identical to Poisson's Green function r^{2-n} of Tegmark 1997); a random walk is recurrent for $d \le 2$ (resistance diverges with N) and transient for $d \ge 3$ (convergent) ⇒ pins $d \ge 3$;
- **Bound-state threshold**: $G(0,0)$ diverges for $d \le 2$ (any weak attraction already gives a bound state) and converges for $d \ge 3$ ($g_c > 0$) ⇒ pins $d \ge 3$;
- **Stable circular orbit**: $V_{\text{eff}} = -A\, g_d(r) + L^2/2r^2$ gives a stable circular orbit only for $d = 3$ ($d = 2$ has no circular orbit, $d = 4$ has no finite r^*, $d \ge 5$ is unstable) ⇒ pins $d = 3$.

⇒ $\{d \ge 3\} \cap \{d \ge 3\} \cap \{d = 3\} = \{3\}$, with **zero free parameters**; the fourth fence (atomic stability, $d \le 3$: Ehrenfest 1917, Tangherlini 1963) pinches from the other end. **But the logical boundary must be registered at the same time**: "selected by mathematics" ≠ "derived from the SRE axioms" — all three criteria are stability requirements (predicates), and **the predicates themselves are still inputs**, while SRE has no intrinsic dimension (`ndim = 3` is passed in from outside). The net effect is that **the type of the input has changed** (from "a numerical value" down to "a predicate"), not that the input disappears. Of the same type, the Coulomb force is **derivable in shape, not in value**: the Green function of the graph Laplacian (L^+, a purely topological quantity) gives the leading term r^{2-d}, and the measurement gives $d = 1 \to r$, $d = 2 \to \ln r$, $d = 3 \to 1/r$ ($R^2 > 0.9999$) ⇒ **$1/r$ is selected by $d = 3$**; but the coupling constant α is still an external input (the only successful external anchor $\Pi_1(M_{60})$ deviates from CODATA by 1.45×10^{-5}), and the project graphs are not three-dimensional lattices ⇒ **they do not of themselves carry $1/r$.**

(6) The discrete-closure law: a unified statement over ten examples. All the above phenomena converge into one statable regularity (denoted G12, an $n = 3$ induction, **awaiting independent falsification**): **the closed theorem targets all lie on the discrete side, and the unclosed targets all lie on the continuous side; every crossing from "discrete" to "continuous" must pay one external input.** Its types and the manner of payment are as follows:

#	Target	Side	Form of payment
1	P0 uniqueness ($19320 \to 5$ classes)	discrete (closed)	enumeration
2	R2 convergence (integer decrease)	discrete (closed)	—
3	Criterion L1 ($\lvert n_n - n_p\rvert \le 1$)	discrete (closed, $A \le 4$)	—

| 4–5 | Electron orbital (two senses: mesoscopic nuclear eigenmode / Aut orbit) | split within one chapter | nominal definition vs zero-parameter emergence |

| 6 | Coulomb + ladder (shape derivable, value not) | crossing | shape: r^{2-d}; value: α pays δ (modulus no-go); degeneracy requires an **ensemble** (measure lacking) |

| 7 | $d = 3$ itself | crossing | pays **a zero-parameter predicate** (the cheapest) |

| 8 | Origin of Q_3 ($Q_3 = \mathrm{D}^2(K_2) = \mathrm{BDC}(K_4)$) | crossing | seed = one edge + one predicate |

| 9 | The folding tower | discrete (fully closed) but with zero continuous content | **to obtain a value one must leave the tower** (M_{60} leaves the tower and only then gives α) |

| 10 | H_2O ratio vs geometry | crossing | pay D; then pay **the map** (curvature / scale) |

(7) The three validation-qualification criteria and the diagnosis of the "geometry-shape" branch. Once the perspective is layered, one can audit **the prior validations themselves** (`code/_sre_validation_metric_audit.py`, 17 entries in all). The three qualification criteria are:

1. **Out-of-sample character**: the comparison target must be an **independent experimental quantity**, and **must not have been used to construct the graph or its parameters** (any distance D weighted by real bond lengths is circular and does not count);
2. **Non-degeneracy**: the quantity under test must have a **nonzero predictive margin** (not an identity rewrite, not a back-substitution, not a low-degree constructive reconstruction);
3. **Window binding**: the tolerance must be of the same order as the demonstrated precision (G4).

Audit result: classified by the epistemological type of the comparison target, **independent experimental out-of-sample E = 9 | inversion / scheme-dependent quantity R = 2 | construction / circularity / back-substitution C = 4 | pure mathematics M = 2**, and **the C and R classes fall 100% within the "geometry / shape" branch**, while the E class lies entirely within the spectral / numerical branch and is **a string of negative verdicts**. Two concrete blockings:

- At $n = 3$, the headline indicator of the MDS document, "emergent geometry vs real", has a **Pearson magnitude that is always 1** (a purely topological distance set is always $(1, 1, 2)$ and a real distance set is always (r, r, x), both being (a, a, b)-shaped 3-vectors), and its entire information content is **a single bit**, "whether it is $> 60°$" — the measurement gives $+1.000000$ for both $\theta = 180°$ and $104.4776°$, so **the indicator is blind to the bond angle**;
- **The "inversion fidelity r" of classical (Torgerson) MDS is always 1 for any Euclidean D** (it is the very definition of the algorithm's convergence), and random 3D point sets with $n = 4/6/10/20$ together with the degenerate $(1, 1, 2)$ all give 1.000000; and for $n \leq 3$ the point count is so small that the triangle inequality alone fixes the distances, so **the inversion fidelity for $n \leq 3$ is a constructive reconstruction and must not be taken as emergence evidence**.

⇒ That branch "appears to hold" precisely because it uses constructed quantities, whereas the spectral / numerical branch uses real experimental values and therefore yields a string of negative verdicts. **Re-filtered by the three criteria, the "geometry-shape" branch currently has 0 qualified entries**; the repair direction is: a purely topological D plus $n \geq 4$ plus **comparing angles / bond lengths directly** (rather than comparing the Pearson correlation of whole distance sets).

(8) The boundary statement and the net consequence. Combining (1)–(7), the boundary of this framework can be written in one sentence:

> **The dimension-free world of SRE provides the "shape" (integer invariants and their dimensionless ratios); the "value" and the "geometry" of the three-dimensional world must be bought with two external inputs — the metric D and the map (Euclideanity / curvature / scale). Every negative verdict of the framework can be located at one of these two tolls, rather than in the insufficient precision of some formula.**

On this basis, the relations among the preceding chapters can be restated: Chapters 7–9 (skeleton inversion) and 12.1–12.5 (assembly law and shared ring) are **exact results on the L0 side** (integers, spectra, ratios), whose strength depends on no continuous input; the negative verdicts of 12.6–12.15 and 13.3 all fall at **the L1 → L2 toll** (requiring a dimensionful value); the dimension collapse test of Chapter 10 and the fission verification of Chapter 11 belong to the **phenomenological layer**, whose role is to organize real empirical data into dimensionless relational numbers of L1 form, not to derive numerical values from the axioms; the "open band" of 12.13 and the λ of 12.12 correspond respectively to the two sides "shape fixable" and "value unfixable", so that their conclusion forms (localizable vs non-derivable) are not coincidences but a necessity of the same layering law.

Honesty boundary: (i) the "three layers" of this section are **a division of representational layers**, not three different physical entities; their value depends on whether a recomputable three-layer reading can be given for any new object. (ii) Theorems 1 and 2 are **mathematical facts** (graph invariants and constant-curvature trigonometry); this section only places them back into the context of "SRE and the world"; they do **not** claim that the SRE axioms are true. (iii) The induction of the "ten examples" (G12) still rests on a small sample, several of which are internal splits within a single chapter (examples 4–5), so its **theorem status awaits independent falsification**. (iv) The **counts** of the audit (E/R/C/M) depend on the classification scheme; the classification table is archived with the paper for re-examination. (v) The external literature cited in subsection (5) for the $d = 3$ conclusion (Ehrenfest 1917; Tangherlini 1963; Tegmark 1997) is **external basis**, not a computation of this section.

---

15. Honesty Boundary Checklist

Each of the following weakens the strength of the conclusions of this paper, and is listed item by item for readers to weigh themselves:

Boundaries Related to Nucleon Inversion

1. **κ_N identification**: $(m_n - m_p)/m_p = \alpha \cdot \Pi_1$ has not been independently demonstrated, it is a candidate hypothesis;
2. **Hierarchy mixing**: The α of electrons comes from the 60-point Möbius parameterized spectrum, while the nucleon side in this paper uses the bare skeleton spectrum, and the two levels cannot strictly be compared;
3. **Δm used twice**: Both criterion ① and criterion ② contain Δm, and the two criteria are not independent;
4. **Criterion depends on prior**: Criterion ③ (triple symmetry) comes from prior S2, if S2 does not hold, the screening result will change accordingly;
5. **Quark ontology borrowing**: $(2m_u + m_d)/m_p$ is only used as a dimensionless measured ratio, which does not mean admitting the ontological status of quarks;
6. **Strength of the tie-break**: The Q3 on which Criterion ⑤ (partial re-equilibration of closure degree) rests, although supported by the fission verification of Section 11, is itself still a conjecture proposed in this paper; Criterion ④ alone cannot break the tie over the whole pool (2 graphs satisfy it); Criterion ⑥ shows a 2.67% deviation, which is not strong enough. The correct statement is therefore "A is the unique survivor at the structural level, but not independently at the measured level";
7. **Dormant edge taken as a ring edge**: The basis is the structural adjudication that "the dormant edge is the image of a homomorphic mapping and its logical depth is large", not an independent derivation from measurement; if a spoke were taken instead, the transition reading would become λ_2: 1 → 0.527166, and the form of every criterion in Section 9 would change accordingly;
8. **"Mode release" not adopted**: This paper explicitly does not use "some mode releases exactly the integer 1" or "the total released amount is always 2" as criteria (they are respectively a contingency of the homomorphic mapping and a trace identity);
9. **Does not contain Q_3**: After exhaustive enumeration this skeleton contains no 3-regular 8-vertex subgraph, so β decay can

only take the weak "seed sedimentation" reading;

10. **Mass-depth not closed, and its zero-parameter assignment rule now judged negative (upgraded in v1.9)**: the L4 mass-depth equation $m_p/m_e = \Psi(G_N)/\Psi(G_e)$ is unavailable because Ψ is undefined, and the C5 (N_p≈5.446×10¹⁹) of v1 is still circular self-proving. Section 12.15 of v1.9 builds that gap into a **pre-registered, falsifiable** test and executes it, with a **negative** verdict on three mutually independent grounds: (i) **reach bound (distribution-free)** - with the electron basal body Q_3 as the electron graph, the monomial-pool bound for $|a| \leq 2$ (any support) is only **511.06**, **short of the target 1836.15 by 0.555 orders of magnitude** (23.07 for support <= 3, short by 1.90 orders); (ii) **"sharing one lock" directly refuted** - with M_{60} as the electron graph the reach is unconstrained, but the best Ψ differs from target to target (m_p/m_e, m_d/m_p, m_n/m_p), and the joint hit is **0** for $|a| \in \{1, 2, 3\}$ under both the 1% and the 3% window; (iii) **hidden degree of freedom** - the electron-layer representative (basal body Q_3 vs carrier M_{60}) is not fixed by the framework and the verdict flips between the two configurations => "which graph represents the electron" is itself a free parameter => **C3 violated**, so even a hit would not be falsifiable. **Positive residue**: the same Π_1 machinery holds on the difference ratio of two states within a layer ($\alpha \cdot \Pi_1(\mathrm{Y}_3)$ against $(m_n - m_p)/m_p$: -0.164%) but fails for inter-layer overall mass ratios ($\Pi_1(\mathrm{Y}_3)/\alpha = 25.84$, short of 1836.15 by 1.85 orders of magnitude) => the cross-layer lock governs the difference within a layer but not the overall scale of a layer. **Boundaries**: the verdict is confined to the Ψ class of "integer powers of products of standard invariants" (the $|a| \leq 2$ reach exclusion is hard; for $|a| = 3$ the ceiling clears the target and the verdict rests only on (ii) and (iii)); for $V > 22$, $|\mathrm{Aut}|$ is not computable, so configuration B's pool excludes it. (`code/_sre_l4_assignment.py`)
11. **Scheme dependence of the Criterion ① anchor (added in v1.6; finalized in v1.7)**: the anchor $(2m_u + m_d)/m_p$ used by Criterion ① is an MS-bar scheme quantity, and it is not even listed in the declared injection list of §3.1. Its measured scale sensitivity (about +23% from 2 GeV to 1 GeV) far exceeds the bin-switching threshold and **suffices to change the selected V** (2 GeV → V=12, 1.5 GeV → V=14, 1 GeV → V=16); moreover κ*N measurably does **not** land uniquely in the V=12 family (V=8/10/12/14 all contain* κN), and after Criteria ②③ are applied V=10 also retains a survivor — **but that item is partly withdrawn in v1.7**: it holds under the wide window ε = 3%; once the window is tightened to the order of the α-lock's own precision (ε ≤ 1%) **only the V=12 bin retains a survivor** (see the v1.7 table in 7.6). **The final position is therefore: Criteria ②③ by themselves determine V (from {8,10,12,14} down to {12}, exact for V ≤ 12 and an LR ≥ 1.7 lower bound for V ≥ 14) and determine the topology (within the family, 85 → 4 → 2 at ε = 3%; 85 → 3 → 2 at ε ≤ 1%, and after a further T ≥ 2 filter the Y_3 skeleton is** uniquely** pinned down within the bin — see the correction in 7.6), and they do so with no dependence on any scheme quantity whatsoever; Criterion ① is downgraded to a corroborating remark, namely "under the 2 GeV MS-bar convention it is compatible with V=12", and no longer carries the calibration duty.** This item does **not** affect the many-body conclusions from Chapter 12 onwards, nor the spectral facts of the skeleton itself — the 0.165% agreement of Π_1 = 0.188580 with κ_N is a purely topological fact, independent of the anchor chosen. **The price must be stated honestly**: the "5 bins select 1" factor in the v1.6 rarity estimate p ≈ 4.7×10⁻³ is now carried by the likelihood-ratio lower bound of the sharpened Criteria ②③, so p loosens to about 1.4×10⁻² to 2.8×10⁻² (see 7.6). (`code/_sre_anchor_registry.py`, `code/_sre_anchor_second.py`)

Boundaries Related to Many-Body Assembly and the Shared Ring

1. **Principle dependence of the assembly law**: the two principles (open–closed complementarity, the images of the two bodies must be distinguishable in the shared region) are structural adjudications, not independently derived from measurement; the "same position" clause is the necessary condition that makes n-n automatically excluded — if a gap were allowed to be sealed by **any** ring of the partner, n-n would also have structural solutions (7 isomorphism classes), so the clause cannot be omitted. Principle 2 has been restated in 12.10 as a purely ledger-theoretic condition (the shared ring contains a single-claim edge with k=1); the restated form has the same discriminating power as the original, but both are model-internal constructions that have not been independently tested;
2. **Withdrawal of the shadow-solution name**: verification shows that the former "variant D" is the k=2 member of the shared-ring family (isomorphic to the deuteron, with the same ledger (1, 2, 2)), so the name "shadow solution" and its note "violates principle 2" are both withdrawn (12.10); its anti-probabilistic features (integer ρ = 6, body-count-independent $\lambda_2 = 2-\sqrt{3}$) rest on the numerical facts of 12.3 and 12.5 and do not depend on the naming;
3. **Reading dependence of the scorecard**: the scorecard conclusion depends on the choice of reading (A / A' / B). This paper adopts Reading A, which matches 5/5 in the calibratable sector (A ≤ 3); however, the additional clause "gap ≤ 2" is a constituent part of the reading and has not been independently derived. Dropping that clause turns only the ^{4}H entry from a hit into a mismatch, still inside the A ≥ 4 interval, so it does not affect the conclusions of this paper. Section 12.12 gives a mirror-symmetric single-clause replacement, criterion L1 (|n_n − n_p| ≤ 1), which matches 8 / 8 over A ≤ 4 with the number of

clauses reduced from two to one and no parameter, and which degrades to an empirical approximation for A ≥ 5 just as Reading A does;

4. **Zero margin of the pricing layer**: ψ(1..4) is fixed by four measured nuclei, so a zero residual is a necessity of parameter counting and is not a prediction; the pricing shape of "interference depth" is only a qualitative description;
5. **The boundary is split into the two questions "existence" and "numerical binding energy"**: this paper originally adopted the boundary "the nuclear boundness for A ≥ 4 (both existence and the numerical value of the binding energy) is not computable in this layer" (12.9), and registered its two independent markers (the out-of-sample failure of the pricing layer in 12.7 and the qualitative failure of the assembly law in 12.8). Section 12.11 further upgrades the **numerical binding energy** side of this boundary from an "empirical failure" to "structurally non-repairable" by two impossibilities: the A = 4 isotriplet forms a **single isomorphism class**, so any graph functional takes the same value on all three; and the additive-profile pricing class is falsified as a whole by the identity $B(^4H) - B(^4He) = B(^3H) - B(^3He) > 0$. Section 12.12 attempts to derive from within SRE the antisymmetric coefficient λ required by that repair, and concludes that it can **neither be derived nor fixed** (the threshold, 29.0596 MeV, exceeds the ^{4}He binding energy, and λ is an absorbable degree of freedom; SRE fixes only its "shape", not its "value"); the same check, however, yields a positive outcome — the zero-parameter **existence criterion L1** ($|n_n - n_p| \le 1$, 8 / 8 on the 8-entry scorecard). Hence the **existence** side is computable at A = 4 and not for A ≥ 5 (the criterion degrades into an empirical approximation; 22 / 34 on the A ≥ 5 out-of-sample set). The boundary is therefore **relocated** from A ≥ 4 to A ≥ 5, while the non-computability of the **numerical binding energy** is unchanged;
6. **Dimensionlessness**: every comparison in this section is a comparison of dimensionless model-internal quantities (uniformly in units of Δm_{np}) and does not constitute a derivation of real nuclear-physics quantities; the nuclear data cited (the boundness and binding energies of ^{2}H, ^{3}H, ^{3}He, ^{4}He, ^{4}H, ^{4}Li) are taken from published evaluations such as AME 2020;
7. **Scope of the isomorphism-class conclusion (now strengthened, and narrowed after B2)**: the "single isomorphism class" of 12.11 originally held only for the shared-ring assembly rule used in this paper and had to be re-checked; that re-check has now been carried out in 12.11(1'), and the conclusion is that the **isomorphism is necessary at the assembly layer** — the three share the same k distribution (a single ring with k = 4), so it is independent of the attachment point and of the specific value of A. The re-check also establishes that **A does not enter the structure; the distribution of k does** (the same A with a different k distribution gives non-isomorphic graphs, whereas the same k with p↔n swapped or a different attachment point leaves the graph unchanged). **The reading "the same A with a different k distribution ⇒ non-isomorphic" must however carry a qualifier**: under the current assembly rule a multi-group product is invariably a disconnected union, the difference coming from the addition law rather than from "k entering a single structure", and the inter-ring linking mechanism is provably impossible to supply (see 12.11(1') and unresolved matter 11). **What determines the distribution of k**: under purely SRE-endogenous conditions it is the single-group form k = (A,) (a weak conclusion), and once external priors are admitted it is not uniquely determinable (settled by B1); moreover **multi-ring configurations should be absent from this layer** (settled by B2, see unresolved matter 11);
8. **Non-additive pricing not explored**: 12.11 falsifies the additive-profile pricing class only and does not exclude a non-additive pricing; no non-additive form is given in this paper;
9. **Independence of the figure data**: the ρ sequence and the λ_2 invariance in panel (b) of Figure 5, and the scorecard in panel (c), are independently recomputed by the figure script from numerical Laplacian spectra (the maximum deviation of λ_2 from $2-\sqrt{3}$ is about 10^{-15}) rather than reusing the tabulated values of the main text; the figure therefore serves as an independent cross-check of the numerical values in the main text;
10. **Non-derivability of the antisymmetric coefficient**: the conclusion of 12.12 is that λ can neither be derived from nor fixed by SRE. The argument rests on two points: (i) graph invariants are constant over the A = 4 isomorphism class, so using one as λ would break the calibration of the four measured nuclei; (ii) any ledger function X(a) is of the same kind as the penalty term g(a) and produces only a renaming. **The scope of this argument is limited to the input set "ledger (A, #p) plus same-isomorphism-class graph invariants"**; if SRE possesses a third class of input that has not yet been identified (neither a graph invariant nor a ledger function), the matter must be re-examined;
11. **Status of criterion L1**: L1 ($|n_n - n_p| \le 1$) and Reading A are both post-hoc readings, and the superiority of L1 consists only in "more hits with fewer clauses" (zero parameters, a single clause, mirror symmetry). Its exactness at 8 / 8 over A ≤ 4 is consistent with the known fact that the light-nucleus region is dominated by α closure, and **it cannot be claimed on that basis that L1 is a fundamental law**: the two mismatches at A = 5 are closure effects (no nucleus is bound at A = 5, so any composition-only criterion must mismatch there), while the mismatches for A ≥ 6 arise from the absence of an A scaling (the measured $|N - Z|$ ceiling of bound nuclei rises with A). In the A ≥ 5 out-of-sample set, "bound/unbound" is taken as ground-state existence, and ^{8}Be is recorded as unbound because it is α-unstable (only 0.092 MeV); changing that convention would affect 1 entry. The out-of-sample set contains 26 entries, and L1 matches 22 / 34 in total (including the 8 scorecard entries).
12. **Status of the open-band reading and the equivalence of the three rings**: the check of 12.13 presupposes the reading "the open band = the one ring edge opened in the neutron state"; that reading is an extension of the β-decay reading of this paper

(Section 5), and its ontological status is a model-internal construction not independently tested. "Electronic state participation" is delimited within this layer as open-band attachment to the shared region and **involves no mesoscopic-layer electron (atomic/molecular electron cloud)** — the latter is reassembled by the electron proper at a higher scale, and its effect does not enter nucleon assembly. 12.13(1) measures the three rings to lie in a single orbit under the automorphism group of the closed-state skeleton, so "the neutron gap can only lie in ring 0" is a notation rather than a fact; that conclusion applies only to the automorphism group of the closed-state skeleton, and the correction leaves every existing conclusion intact (12.13(2) re-runs r = 0, 1, 2 one by one and confirms this).

13. **Unusability of the functional Π_2 on the nucleon side (now determined, not merely "uncalibrated")**: the w-independence of the second electron-side w-independent functional $\Pi_2 = \lambda_2/\lambda_4$ **has a threshold** — an exhaustive scan over even n = 8…120 shows that Π_2 is w-independent **only for n ≥ 24** (where λ_2 is the k = 2 mode and λ_4 the k = 4 mode, both even modes); for n ≤ 22, λ_4 falls on an odd mode and Π_2 depends on w. The electron-side Π_2 therefore holds only under the qualifier "**for n ≥ 24**". On the nucleon side it is **structurally unusable**: the single-group shared-ring k-body skeleton has $\lambda_2 = 2-\sqrt{3}$ exactly with **multiplicity m = k−1**, and the next distinct eigenvalue is invariably 1, so **for k ≥ 4, λ_4 falls inside the degenerate block and $\Pi_2 \equiv 1.0$** (identity, no information), while **for k = 2, 3, $\lambda_4 = 1$ so $\Pi_2 = 2-\sqrt{3}$** (a mere re-reading of λ_2, with no independent information). This degeneracy mechanism has the same origin as 12.5 (the three-point zero amplitude of the shared ring together with the zero sum of each body), **so Π_2 cannot be used for nucleon inversion**.
14. **Homology of the cross-layer parameterization (the "shape" transfers, the "value" does not)**: the electron-side and nucleon-side spectra belong to the same one-parameter family "2 − 2cos(mθ)": on the electron side $\theta_e = 2\pi/n$ *(the discretization angle), while on the nucleon side solving $\lambda_2 = 2-\sqrt{3}$ for the angle gives cos* $\theta_N = \sqrt{3}/2$, i.e. **$\theta_N = \pi/6$**. Alignment of the two sides requires $2\cdot(2\pi/n) = \pi/6$, whose **unique positive-integer solution is n = 24**; at that example the electron-side Π_2 exactly equals the nucleon-side λ_2 (both $2-\sqrt{3}$). **Note**: n = 60 (the primary electron-side object of this paper) **does not** satisfy that alignment, and n = 24 does not constitute a general family "n = 6m". Hence the parameterization **"shape" of the two layers (the family structure and the even-mode protection principle) transfers**, whereas the **"value" does not** (the electron Π_1 depends on w, Π_2 has the n ≥ 24 threshold, and on the nucleon side Π_2 carries no information with no adjustable n). This is the fourth instance of the recurring regularity of this paper — "**SRE fixes only the 'shape', not the 'value'**" (the other three being λ in 12.12, w(k) in 12.11, and the B1 conclusion on the k distribution);15. **Reading the neutrino as the k = 1 singly-claimed edge is judged negative (added in v1.8)**: 12.14 turns the registration "neutrino = the k = 1 singly-claimed edge on the shared ring" into an executable criterion and then judges it negative. The blocking is **distribution-free** and rests on no numerical coincidence: (i) the graph route — the **number of orbits of the automorphism group on the three shared-ring points is measured to be 1** (vertex-transitivity), so every graph functional takes the same value on the three-generation candidates ⇒ the spectrum is strictly degenerate ⇒ $\Delta m^2 = 0$; the 3×3 local operators (G_3 and L_{eff}) of all six assemblies have eigenvalues with **always only 2 distinct values** (multiplicity 2+1) ⇒ the spectrum degenerates to two levels, and one of the two Δm^2 is identically zero; (ii) the ledger route — the claim triple is always (#closed, k, k), with **always 2 distinct values** ⇒ only 1 degree of freedom. Neutrino oscillations, however, require 3 non-degenerate states / 2 independent non-zero gaps. Of 96 zero-parameter candidate numbers, only 1 falls inside the $\varepsilon = 3\%$ window (`1/E = 1/33`, deviation 1.28%), which is below the MC null expectation (p_A = 0.166) and **below** the hit rate of random graphs with the same degree sequence (0.385). Hence this reading **cannot carry the neutrino mass spectrum**. Boundary: the negative verdict is **conditional** (confined to the reading above); other readings (weighted spokes, a non-additive ledger) would violate C3 (free parameters ≥ number of targets) and become unfalsifiable, and must not be used to press ahead. One further item must be registered: **a presupposition of the first draft of that section has been overturned by measurement** (it assumed that the two ends S0, S1 of the k = 1 edge formed an orbit of their own with S2 separate ⇒ 2 orbits; the measurement gives 1 orbit, i.e. vertex-transitivity), and the direction of the negative verdict is unchanged but stronger.
(`code/_sre_neutrino_k1.py`)

Boundaries Related to Fission Verification

1. **Structural verification rather than numerical prediction**: All observables have already been published; SRE only provides a unified graph-theoretic reading and does not derive new numbers from first principles;
2. **Aligned with nuclear physics**: The Δ criterion agrees with the established nuclear-physics statement of neutron separation energy versus fission barrier; SRE only performs a translation of language;
3. **Fast cross-section error**: The fast cross-sections are nominal values, and the definition of the energy point implies ±30% uncertainty; the conclusion that "the same order of magnitude is restored" is robust;
4. **Correlation is not causation**: The Pearson value of 0.932 refers to 9 target nuclei, and although Δ and σ_th come from

mutually independent nuclear data, the significant correlation is not a proof of causation;

5. **Fission net energy deviation**: The model deviation is +6~8%, with the direction and magnitude correct; the precise value depends on the choice of the fragment B/A sampling point;
6. **Chain criticality uncorrected**: p" is the upper limit for f=1; a real reactor requires leakage and absorption corrections.

Boundaries Related to Dimension Collapse and Theoretical Constraints

1. **Dimension collapse as empirical indication**: All tests merely organize real empirical data into dimensionless relational numbers and verify that they approximately obey additivity/step/integer-ratio structures; this is an aligned indication of the SRE ontology "distance = coherent bookkeeping", and no numerical value is derived from SRE;
2. **Bond length fluctuation**: Bond lengths are empirical representative values; the same bond fluctuates by ±0.01–0.03 Å in different environments, which is noise at the decimal-place level relative to a0, and should not be over-interpreted bond by bond;
3. **Covalent radius calibration dependence**: The covalent radii adopt the Cordero 2008 calibration; switching to the Pyykkö system would shift individual residuals by about 1–3%, but the family structure and the conclusions remain stable;
4. **Theoretical constraint inferences**: The special binding of ions/metals and the current-emergence inference are both logical consequences of model self-consistency; they change no measurable numerical value, only replacing the traditional explanatory layer.5. **The exponential decay law has been derived, but the strong reading of its two-factor decomposition is judged negative (added in v1.8)**: 13.3 uses the axioms of Sections 5.2 / 9.4 (dormancy as the default state + removal as a rare, independent, stationary perturbation) to **derive** the exponential law $T_{1/2} = \ln 2/p$ and memorylessness (MC verification: survival-curve deviation 1.03×10⁻³, half-life-identity deviation 0.0224%, memorylessness deviation 5.7×10⁻⁴; the power law is excluded — the log-log piecewise slope drifts from −0.570 to −5.013, with R^2 0.781 against 0.999999), thereby closing the "exponential-law gap" registered in Section 10. However, the **species-cancellation law** that follows from $p = p_{\mathrm{graph}} \cdot p_{\mathrm{env}}$ (for the same nuclide across environments, p_{graph} cancels exactly) yields the zero-parameter prediction "the stripping enhancement factor E is independent of the nuclide", which is **falsified by measurement**: ^{187}Re and ^{163}Dy share the mode (bound-state β⁻) and have nearly the same Q (2.467 and 2.560 keV), yet their E differ by **≥ 6.79 orders of magnitude**, and ^{187}Re and ^{241}Pu share the mode yet differ by 6.01 orders of magnitude. Since p_{graph} has already cancelled exactly in that ratio, **the failure cannot be repaired by the graph factor** ⇒ p_{env} must contain species-specific atomic/electronic-configuration information, whereas the SRE side has only a scalar. Boundary: (i) data-reliability grading — ^{187}Re (Bosch 1996) and ^{163}Dy (Jung 1992) are primary literature while ^{241}Pu is a secondary review, and **the decisive pair does not include ^{241}Pu**; (ii) the L3 data are all heavy nuclei (A ≥ 163) whose SRE graphs are unavailable in this layer ⇒ this section is a **decomposition test at the phenomenological layer**, not a numerical derivation internal to SRE; (iii) the section refines the reading of Section 10 in one respect — the control parameter is refined from "the reachability of the s electron at the nuclear site" to "**the relative sensitivity of the s-electron density at the nuclear site to the environment**" (an environment term plus a species term), while the three-layer stratification itself (span 13.40 orders of magnitude) is unaffected.
(`code/_sre_decay_law.py`)

Boundaries Related to the Observational Perspective and Validation Qualification (added in v2.0)

1. **The "three layers" are a division of representational layers, not physical entities**: Section 14 divides the description into L0 topological / L1 ratio / L2 geometric, and states that "the native products of SRE stop at L1". This is a statement about **representation and input**, not about entities; its testable form is "give a recomputable three-layer reading for any new object", a per-object checklist of that kind not yet being provided in this paper.
2. **Status of the two theorems**: angle ∉ functions(graph) (Theorem 1) and "same ratio, different map ⇒ different angle" (Theorem 2) are **mathematical facts** (graph invariants and constant-curvature trigonometry); this round only places them back into the context of "SRE and the world". **They do not claim that the SRE axioms are true**, nor do they constitute independent physical evidence about the three-dimensional world. The numerical values of Theorem 2 (spherical 104.57° → 145.10°, hyperbolic 104.39° → 86.67°, over the same scale window $s \in [0.1,\ 1.6]$) depend on the curvature radius chosen

(set to 1 in all cases); changing the radius changes only the specific values, not the qualitative conclusion that "the same ratio gives an interval".

3. **The "predominantly ratios" claim is an internal empirical statistic of this project, not a theorem**: Section 14(2) states that "every quantity passed is a dimensionless ratio, and every negative verdict occurs when a dimensionful value is required". That claim is computed from this project's own entries (see the E/R/C/M counts of Section 14(7)), its **sample being this paper itself**, and it may be rewritten by subsequent entries.
4. **The discrete-closure law (G12) is an induction, not a proof**: among its "ten examples", examples 4–5 are internal splits within a single chapter, and examples 6–9 are mostly repeated occurrences of a single-parameter form of payment, so the law is **only a unified statement of the existing entries** and does not constitute an impossibility theorem; its counterexample form would be "some continuous target closed without external input".
5. **Scope of the three validation-qualification criteria**: the three criteria (out-of-sample character / non-degeneracy / window binding) target "fidelity / loyalty / correlation-coefficient" type indicators, not exhaustive enumeration or purely mathematical entries, so the M class (P0, R2) is not bound by them; whether the entries classified as R (e.g. the Criterion ① anchor) should be downgraded to C depends on whether a scheme-independent substitute can later be given, and this round reaches no conclusion on that.
6. **MDS inversion fidelity must not be taken as emergence evidence for $n \leq 3$ (added in v2.0; a long-standing outstanding item)**: the objective function of classical (Torgerson) MDS is $\min\|D - \mathrm{dist}(X)\|$, so for any **Euclidean** D its "inversion fidelity r" is always 1; and for $n \leq 3$ the degrees of freedom of the points are so few that the triangle inequality alone fixes the distances (the water "$r = 1.000$" of the MDS document is of this kind). ⇒ **any use of an $n \leq 3$ inversion fidelity as "emergence" evidence is disregarded in this paper**; the indicator carries information only when D is non-Euclidean (there it measures "embeddability", not geometric correctness). The repair direction is a purely topological D plus $n \geq 4$ plus comparing angles / bond lengths directly.
7. **The convention dependence of the water "104°" and two corrections (added in v2.0)**: the two points registered in Section 14(4) must be carried along whenever it is cited — (a) the triple $(0.9584,\ 1.51383,\ 104.474)$ once used by a project script **mixes averaging conventions** (the first is of r_e magnitude, the middle of r_0 magnitude), its closed angle being only $104.3279°$ (a difference of $0.146°$ from the stated value), and it should not be marked "self-consistent"; the consistent triple is $(0.9578,\ 1.5144,\ 104.4776)$. (b) The equilibrium angle happens to satisfy $\cos\theta_e = -0.250001$ (10^{-6} level), so the normalized $t \approx \sqrt{2.5}$; the $t = 1.5811$ in the project is a **back-substitution plus coincidence** and must not be taken as evidence that "the graph predicts the water angle" (of the same kind as "$\Pi_1(Q_k) = 1/k$ on the tower is about $1/137$ at $k = 137$").
8. **Boundary of Q_3 and the folding tower (supplementary registration in v2.0)**: the exact laws on the tower registered in paragraph 12.15(f) ($|\mathrm{Aut}(Q_k)| = 2^k k!$, distinct Laplacian values $= k+1$, $\Pi_1(Q_k) = 1/k$) **hold only for members of the folding tower**, and those members **have orbit count always 1** (both D and BDC preserve vertex-transitivity) ⇒ the shape resolution within the tower is zero, and no "shape" discussion may be based on tower-internal readings. One **misuse-prevention trap** is registered as well: $\Pi_1(Q_k) = 1/k$ is about $1/137$ at $k = 137$ (the numerical value of the classical α); this is a **constructive coincidence** arising from bipartiteness plus the covering, and Q_{137} has 2^{137} vertices and is physically unusable, so **it must not be taken as evidence**.

---

16. Conclusions

This paper completes the full characterization of nucleons within the SRE framework; its core conclusions and logical chain are as follows:

Core Conclusion

Nucleons are not point particles pre-placed in space, but **stable composite objects that emerge from the binary self-organizing network on the tripartite Y-shaped coherent core, constrained by the two-state opening/closing of dormant edges and the closure degree rebalancing rules**; their ontological structure is **the tripartite Y-shaped triangular closure $Y_3 \ltimes \triangle_3$** ($V=12$, $E=18$, $\beta_1=7$, $|\mathrm{Aut}|=36$). This conclusion is reached from the clue of neutron β decay and obtained through three-step screening (integer determination of the vertex number $V=12$, spectral ratio determination of topology, triple symmetry determination of the solution), with a combined rarity of approximately 4.7×10^{-3}.

Key Logical Chain

1. **Methodological homology**: Nucleon derivation follows the same "local input → topological solution → inversion" route as electron derivation; SRE can only give dimensionless ratios, and absolute quantities require local input from classical physics;
2. **Quark ontology reinterpretation**: Quarks should not share the Q_3 substrate with electrons, but are emergent statistical phenomena; $N_c=3$ in the Standard Model is in fact the number of skeleton strands, and flavor is the statistical projection of opening/closing breaking;
3. **Two-state opening/closing mechanism**: Neutron = open state ($\beta_1=6$), proton = closed state ($\beta_1=7$); β decay is the removal/maintenance bifurcation of a dormant edge, and a breaking magnitude of 10^{-3} is a structural necessity;
4. **Fission chain verification**: The mechanism "nuclear reaction = closure degree rebalancing" holds equally at the compound-nucleus and macroscopic scales; fast-neutron cross-sections restore the same order of magnitude, the excitation surplus criterion has a Pearson correlation of +0.932 with the thermal cross-sections, and the three-level signatures are unified (single-nucleon β decay → compound-nucleus fission → macroscopic chain criticality);
5. **Theoretical constraints**: The additivity failure of ion/metal pairs is theoretically predictable (channel topology switching), and current is a coherent bookkeeping projection rather than continuous electron movement (distance is not an a priori coordinate);
6. **Many-body assembly law and the shared ring**: the assembly law is determined by two principles ("open–closed complementarity" and "the images of the two bodies must be distinguishable in the shared region"), and the formation of the n-p shared unit follows automatically from the open/closed asymmetry; the lowest excitation of the shared-ring configuration is strictly independent of the body count ($\lambda_2 = 2-\sqrt{3}$, multiplicity $k-1$), and the shared ring is precisely the part of the lowest excitation that does not move at all, serving as a natural reference frame; the zero-parameter existence scorecard matches 5/5 in the $A \le 3$ sector; **the numerical value of the binding energy** is delimited as a non-computable interval for $A \ge 4$, and 12.11 shows that this boundary is structural: the $A = 4$ isotriplet forms a single isomorphism class (no graph functional can distinguish them), and the additive-profile pricing class is falsified as a whole by an identity. The former "shadow solution" (variant D) is verified to be the $k=2$ member of the shared-ring family, and that name has been withdrawn (12.10).
7. **Non-derivability of the antisymmetric coefficient and the existence criterion L1**: 12.12 attempts to derive from within SRE the antisymmetric coefficient λ required to repair $A \ge 4$, and concludes that **λ can neither be derived nor fixed** (the required threshold, 29.0596 MeV, exceeds the ^{4}He binding energy; the SRE-internal candidate pool is underdetermined; SRE fixes only the "shape" of λ — its support and monotonicity — and not its "value"). The same check, however, reveals that the only contentful form of antisymmetry is an **existence criterion** rather than an energy term — the zero-parameter, mirror-symmetric **L1: $|n_n - n_p| \le 1$** matches the 8-entry scorecard of 12.8 at **8 / 8** (Reading A: 7 / 8). The boundary of this paper is accordingly split into two questions: the **numerical value of the binding energy** is not computable for $A \ge 4$ (locked by the identity of 12.11(2)); **existence** is computable at $A = 4$ and not for $A \ge 5$ (L1 gives 22 / 34 on the $A \ge 5$ out-of-sample set, its mismatches arising from the α closure effect and from the absence of an A scaling). The boundary is **relocated** from $A \ge 4$ to $A \ge 5$.
8. **Containment of the open band (configuration-level localization of the electronic state)**: 12.13 rewrites "does the nucleon–nucleon force require the electronic state as well" as a configuration problem decidable in this layer, and concludes that its only decidable form — **open-band attachment to the shared region** — is **isomorphic** to variant A (hence not a new structure, and already quantitatively excluded by the deuteron binding energy) and **does not constitute** the "third class of SRE input" sought in 12.12. The diagnosis does yield a positive result: **the open band is the carrier of principle 1 of the assembly law**, appearing as the "shape of the gap" — the presence or absence of the gap decides whether n-p can assemble, the position of the gap decides that n-n is excluded automatically, and the **$k = 1$ singly-claimed edge** left after closure decides that the images of the two bodies are distinguishable (principle 2). This simultaneously yields an explanation for a fact previously described only phenomenologically: **the shared ring always has exactly one $k = 1$ edge** (the image of the open band), and this is precisely the structural origin of the statement in 12.5 that "the three points of the shared ring have zero amplitude in the lowest excitation mode, so $\lambda_2 = 2-\sqrt{3}$ is independent of the body count". The section also corrects a

previously untested statement ("the neutron gap can only lie in ring 0" — the three rings are in fact a single automorphism orbit), a correction that leaves every existing conclusion intact.9. **The exponential law and the cessation of the downward branches (both new in v1.8)**: 13.3 **derives** the exponential decay law and memorylessness from the axioms of Sections 5.2 / 9.4, closing the gap registered in Section 10, and shows that the law requires no numerical value of p from SRE — only its form; at the same time its two-factor decomposition $p = p_{\mathrm{graph}} \cdot p_{\mathrm{env}}$ yields a **zero-parameter species-cancellation law** (for the same nuclide across environments, p_{graph} cancels exactly), whose consequent prediction "the stripping enhancement factor is independent of the nuclide" is **falsified by measurement** (^{187}Re vs ^{163}Dy, same mode and nearly equal Q, differ by ≥ 6.79 orders of magnitude) and, precisely because p_{graph} has already cancelled, **cannot be repaired by the graph factor** — so p_{env} must contain species-specific atomic structure, which the SRE side does not supply. In parallel, 12.14 turns the registration "neutrino = the k = 1 singly-claimed edge on the shared ring" into an executable criterion and judges it **negative**, with a **distribution-free** blocking: the automorphism group is vertex-transitive on the three shared-ring points (orbit count measured to be 1) so every graph functional is constant on the three-generation candidates, the 3×3 local operators have only 2 distinct eigenvalues (spectrum reduced to two levels), and the ledger route has only 1 degree of freedom — against a requirement of 3 non-degenerate states and 2 independent non-zero gaps. The downward branches ① (lepton family) ② (CKM) and ④ (neutrino) thus all point to **one and the same missing piece**: the assignment rule "**integer invariant → real measured value**", i.e. the formalization of the L4 lock.

10. **Criterion construction for, and negative verdict on, the L4 assignment rule (added in v1.9)**: Section 12.15 builds the missing piece to which the merge of the "four downward branches" points (the assignment rule "integer invariant -> real measured value", i.e. the L4 equation of Section 7.1) into a **pre-registered, falsifiable** test and executes it. With a zero-parameter monomial class as the Ψ candidates, one design target (m_p/m_e) and two out-of-sample targets (m_d/m_p, m_n/m_p), and a decision tolerance of 1% (deliberately wider than the framework's demonstrated precision of 0.165%), the verdict is **negative** on three mutually independent grounds: (i) **reach bound (distribution-free)** - with the electron basal body Q_3 as the electron graph, the full-support bound for $|a| \leq 2$ is only 511.06, short of the target 1836.15 by 0.555 orders of magnitude (23.07 for support <= 3), so no Ψ in the pool can produce m_p/m_e; (ii) **"two graphs share one lock" directly refuted** - with M_{60} as the electron graph the reach is unconstrained, but the best Ψ differs from target to target (`lam2L^2 rhoL^2 energy^-1` / `E^1 b1^2 energy^-2` / `E^-2 rhoA^2 energy^1`), and the joint hit is 0 for $|a| \in \{1, 2, 3\}$ under both the 1% and 3% windows; (iii) **hidden degree of freedom (C3 violated)** - the electron-layer representative (basal body Q_3 vs the α carrier M_{60}) is not fixed by the framework, and the reach verdict flips between the two configurations, so "which graph represents the electron" is itself a free parameter => even an apparent hit would be non-falsifiable. The MC null (407 trials on random graphs with the same (V, E)) shows that the real configuration's single-target hit counts are of the same order as for random graphs (T_1 1 against 0.40 per trial, T_2 18 against 21.46, T_3 191 against 94.91), i.e. no above-null signal. **Positive residue**: the same Π_1 machinery holds on the difference ratio of two states within a layer ($\alpha \cdot \Pi_1(\mathrm{Y}_3)$ against $(m_n - m_p)/m_p$: -0.164%) but fails for inter-layer overall mass ratios ($\Pi_1(\mathrm{Y}_3)/\alpha = 25.84$, short of 1836.15 by 1.85 orders of magnitude) => the cross-layer lock governs the difference within a layer but not the overall scale of a layer. **The net contribution of this round is therefore**: the merge of four pending items is turned into a single executed test, and the missing piece is shown **not to be suppliable by switching to a larger zero-parameter pool**.

11. **The twist of observational perspective: the boundary between the dimension-free SRE world and the three-dimensional world (added in v2.0)**: the positive and negative verdicts of the preceding chapters are converged in Section 14 into a single explicit division of perspective — the description splits into **three layers**: **L0 topological** (the graph; it yields only integer invariants, with neither ratios nor angles), **L1 ratio** (dimensionless ratios, the intrinsic representation compatible with SRE), and **L2 geometric** (angles and coordinates, purely the output of a map); the two external inputs across the layers are respectively **the metric D** and **the map** (Euclideanity / curvature / scale). This yields two **theorem-level** conclusions: (a) **angle ∉ functions(graph)** — the same graph P_3 with two admissible metrics gives $180.0000°$ and $73.7398°$, while every graph invariant is identical term by term; (b) **same ratio, different map ⇒ different angle** — the same ratio $t_0 = 1.581140$ gives $104.4776°$ in the Euclidean case, on the sphere ($R = 1$) gives $104.57° \to 145.10°$ as the scale varies, and in the hyperbolic case gives $104.39° \to 86.67°$ (over the same window $s \in [0.1,\ 1.6]$), and a curved space has no similarity ⇒ **geometry is a property of the map, not of the graph**. In support of this, a 17-entry **validation-qualification audit** shows that **the C and R classes fall 100% within the "geometry / shape" branch** (which has 0 qualified entries), whereas the E class lies entirely within the spectral / numerical branch and yields a string of negative verdicts. **Net consequence**: the strength of the framework lies in "fixing the shape" (laws, order, layering and dimensionless ratios), and its boundary lies in "fixing the value" (dimensionful real numbers); the seam between the two is exactly the nameable toll of "the map"; every prior negative verdict of this paper falls at one of the two tolls, rather than in the insufficient precision of some formula.

Unresolved Matters and Next Steps

1. **Independence of the tie-break**: Candidate B has been excluded by Criterion ⑤ (the open state must still retain closed units), and A is the unique survivor at the structural level; however, this criterion rests on Q3 (closure-degree re-equilibration), a conjecture of this paper, and a measured criterion independent of the structural prior is still lacking (Criterion ⑥ shows a 2.67% deviation, not strong enough);
2. **Π_2 is unusable for nucleon inversion (now determined, not merely "uncalibrated")**: the electron-side w-independence of the second w-independent functional $\Pi_2 = \lambda_2/\lambda_4$ **has a threshold** — an exhaustive scan over even n = 8...120 shows that Π_2 is w-independent **only for n ≥ 24**, where λ_2 and λ_4 are both even modes (for n ≤ 22, λ_4 falls on an odd mode and Π_2 depends on w). The electron-side Π_2 should therefore be stated as holding "**for n ≥ 24**", and the original phrase "the second w-independent functional" requires that qualifier. On the nucleon side it is **structurally unusable**: the single-group shared-ring k-body has $\lambda_2 = 2-\sqrt{3}$ exactly with **multiplicity m = k−1** and the next distinct eigenvalue invariably 1, so for k ≥ 4, λ_4 lies inside the degenerate block giving **$\Pi_2 \equiv 1.0$** (identity), while for k = 2, 3, $\lambda_4 = 1$ giving $\Pi_2 = 2-\sqrt{3}$ (a mere re-reading of λ_2). The mechanism has the same origin as 12.5. **This item therefore moves from "uncalibrated" to determined: the issue is not "calibration data are still missing" but "the functional carries no independent information on the nucleon side".** (`code/_sre_nucleon_pi2_calibration.py`)
3. **Skeleton-spectrum parameterization (homology point located, but only the "shape" transfers)**: the spectra of the two layers belong to the same one-parameter family "2 − 2cos(mθ)" — the electron side has $\theta_e = 2\pi/n$ *(given by the discretization number n), while on the nucleon side solving $\lambda_2 = 2-\sqrt{3}$ for the angle gives cos* $\theta_N = \sqrt{3}/2$, i.e. **$\theta_N = \pi/6$**. Alignment of the two sides requires $2\cdot(2\pi/n) = \pi/6$, whose unique positive-integer solution is **n = 24**; at that example the electron-side Π_2 exactly equals the nucleon-side λ_2 (both $2-\sqrt{3}$). **It must be stated honestly, however**, that n = 60 (the primary electron-side object of this paper) does not satisfy that alignment, and that n = 24 does not constitute a general family "n = 6m". Hence the parameterization **"shape" of the two layers (the family structure and the even-mode protection principle) transfers**, whereas the **"value" does not** (the electron Π_1 depends on w, Π_2 has the n ≥ 24 threshold, and on the nucleon side Π_2 carries no information with no adjustable n). **This item therefore moves from "unresolved" to "partly settled": the homology point is located (positive), but no cross-layer mapping exists (negative); together with λ in 12.12, w(k) in 12.11 and the B1 conclusion on the k distribution, it forms the fourth instance of "SRE fixes only the 'shape', not the 'value'".**
4. **The k distribution and inter-ring linking (both settled)**: the role of the k distribution is located by 12.11(1') and B1 — under purely SRE-endogenous conditions the only admissible form is the single-group k = (A,) (a weak conclusion: the anchor pool has only 2 entries and there is exactly one effective discrimination); once external priors are admitted the distribution is not uniquely determinable from within SRE. The inter-ring linking mechanism has been shown by B2 to be **impossible to supply**: under the current assembly rule a multi-group product is invariably a disconnected union (cc = number of groups, 9/9), whose (V, E, β_1) is **logically incompatible** with the published values (connectivity costs at least one inter-ring edge, forcing E from 96 to 97); and no mechanism satisfies all four of "degenerates to the current rule for a single group, is connected for multiple groups, does not destroy 12.5, introduces no new vertices" (since §12.5 requires the shared ring to form a closed unit, which measurably excludes "connectivity" — an exhaustive scan of all minimal connection schemes finds no λ_2 returning to $2-\sqrt{3}$). **Joint conclusion: multi-ring configurations should be absent from this layer, and the multi-group constructions for A ≥ 5 should be abandoned.** This impossibility theorem **does not depend on B1**, resting only on §12.5 and the addition law, and therefore stands independently. (`code/_sre_nucleon_interring_b2.py`, `code/_sre_nucleon_kdist_rule.py`)
5. **Graph-theoreticization of the Δ criterion**: Upgrade the compound-nucleus excitation surplus Δ=S_n−B_f from a phenomenological criterion to a derived quantity of the $Y_3 \ltimes \triangle_3$ skeleton;
6. **Residue–release correspondence not quantified**: Section 9.3 has established ρ as the strict invariant of the transition (the residue), but how "residue (denominator) × mobile branch (numerator)" corresponds quantitatively to "light as the reference object of measurement" still lacks an explicit mapping;
7. **Computability for A ≥ 4 (settled)**: 12.11 has shown that the boundary is non-repairable both at the structural layer (isotriplet as a single isomorphism class) and at the additive pricing layer (identity), and has given the only feasible repair — a mirror-antisymmetric term $g(a) = \lambda\cdot\max(0, a-1)$ with $a = |n_n - n_p|$ — together with its price: 1 fitted parameter and no predictive margin for A ≥ 5. 12.12 further proves that the coefficient of that repair, **λ, can neither be derived from nor fixed by SRE** (the question "can λ be derived within SRE?", previously listed as pending, is thereby answered negatively: λ is an absorbable degree of freedom, and SRE fixes only its "shape"); at the same time, 12.12 settles the **existence** question at A = 4 with the zero-parameter criterion L1 ($|n_n - n_p| \le 1$), giving 8 / 8. This item therefore moves from "partly settled" to **settled**: the numerical binding energy is not computable for A ≥ 4, while existence is computable at A = 4 and not for A ≥ 5. **12.13 has eliminated one of the candidate routes**: reading "electronic state participation" as open-band attachment to the shared

region yields a configuration isomorphic to variant A, so it **does not constitute** the "third class of SRE input" sought in item (i) below; the same check, however, yields a positive result — the open band is itself the carrier of principle 1 of the assembly law. **Two matters remain unresolved**: (i) whether a third class of SRE input exists that is neither a graph invariant nor a ledger function and is **not equivalent to a variant-A docking**, which would let the existence criterion L1 shed its status as a post-hoc reading and would explain its mismatches at A ≥ 5; (ii) whether A ≥ 5 can be used to fix ψ(A ≥ 5) unilaterally through other observables (e.g. the neutron separation energy sequence), thereby restoring **binding-energy** out-of-sample testability in that interval. The disposition of the "shadow solution (variant D)" left over from the tie-break adjudication has been completed in 12.10 and is no longer listed as unresolved; the presentational hazard that "the equivalence of the three rings was untested" has been explicitly registered and corrected in 12.13(1).

8. **External anchor (added in v1.6; continued in v1.7, including the negative verdict on the second anchor)**: SRE outputs only ratios, so absolute values require an external anchor; the homogeneity theorem of Op13–18 further requires that L4 "share one lock across two graphs". This round has **located and empirically verified** that lock: only **α** acts on both layers simultaneously, and the two relations side by side form exactly a closed loop — on the electron layer $\alpha = 1 \cdot \Pi_1(M_{60})$ (α as the **output**, deviation 1.45×10⁻⁵); on the nucleon layer $(m_n - m_p)/m_p = \alpha \cdot \Pi_1(\mathrm{Y}_3)$ (α as the **input**). This yields a **zero-parameter reverse prediction**: given α and the Y_3 skeleton (**with no nucleon data whatever**), the predicted $\Delta m_{np}/m_p = 1.376138 \times 10^{-3}$, against a measured 1.378419×10^{-3}, a relative deviation of **−0.165%** (a Δm_np difference of −0.0021 MeV) — the cleanest cross-layer chain in this project to date. It is simultaneously determined that α fixes only the "**shape**" (spectral-ratio landing point and gauge); **V is not fixed by α** but by Criteria ②③ once the window is tied to the α-lock's precision (v1.7, see 7.6). As to the **second anchor** (added in v1.7; a negative verdict): after screening the candidate set against the four criteria C1/C2/C3/A1, **only the μ family survives** (μ*n*/μp, μ*p*, μd, μ(³H), μ(³He), μ(⁶Li)), whose common structural feature is that "a magnetic moment can only come from a current loop" ⇒ the corresponding graph quantity can only be β_1. Three rounds of testing follow. ① **Statistical**: the whole-set hit-rate check (624 comparisons plus Monte-Carlo null hypotheses) gives the strongest V-sharp single-point hit $\beta_1(o)/\beta_1(c)\big|_{V=12} = 6/7 = 0.857143$ against $\mu_d/\mu_N = 0.857438$ (deviation 0.034%), but the p-value swings between 0.04 and 0.2 with the number of effectively independent targets, so it is only **marginal**. ② **Structural (decisive)**: the graphs of ^{3}H and ^{3}He are **isomorphic** (same edge set, different ledger only), so **every** graph functional predicts $|\mu(^3\mathrm{H})|/|\mu(^3\mathrm{He})| = 1$, whereas the measured value is 1.4001 (deviation 28.6%) — **the μ family cannot be carried by graph functionals; it can only be a ledger functional** (exactly the same conclusion as 9.2, "the product graph does not remember the partner state"). ③ **Family law**: the ledger functionals {#p, A, #n, ψ(#p)+2ψ(A), ...} are put through a zero-parameter ratio test on 10 structure pairs, and **none passes**. By-product (a positive result): the sign side does carry content — "sign(μ) flips with the parity of #p" holds 5/5 for A ≤ 4, but external falsification on 16 real nuclei downgrades it to a **trend (14/16 = 87.5%) rather than a zero-parameter law** (counterexamples ¹³C, ¹⁵N). **Conclusion: the second anchor has not been found, and it has been proved that it cannot exist at the level of graph functionals; if it exists at all it must live in the ledger layer, and must pass a family-law test.** (`code/_sre_anchor_registry.py`, `code/_sre_anchor_second.py`)
9. **Closure of the four downward branches (added in v1.8)**: all four branches registered in Section 12.8 now have conclusions — ① the lepton family is **pending** (a zero-parameter "generation → (n,w)" rule is required; not done); ② the Latin square → CKM is **judged negative** (12.9; a zero-parameter "discrete class → continuous angle" rule is required); ③ the boson-as-transition-quantum **numerical route is judged negative while the conceptual redirection is retained** (12.10; the positive by-product "the photon is unique" is registered); ④ the neutrino as the k = 1 singly-claimed edge is **judged negative** (12.14; the graph route is blocked by vertex-transitivity and the ledger route by counting). The pending items of ①②④ **merge into one and the same missing piece** — SRE lacks the assignment rule "**integer invariant → real measured value**" (the formalization of the L4 lock of Section 3). One further item is **closed**: the "exponential-decay-law gap" registered in Section 10 has been filled by the axiomatic derivation of 13.3 (a positive result), while the strong reading of the same section (p_{env} = a pure environment quantity) is falsified by measurement (a negative result), and its gap (the species-specific dimension of p_{env}) likewise falls under that one missing piece. **The net conclusion of this round is therefore: two "distribution-free" blockings are added to the project's boundary list (the vertex-transitivity of 12.14 and the falsification by the species-cancellation law of 13.3), and four pending items are collapsed into one.**

10. **The L4 assignment rule (added in v1.9; settled this round)**: the "four downward branches" obtained from the top-down screening of Section 12.8, together with the species dimension of p_{env} exposed in 13.3, had all been merged into a single missing piece - the assignment rule "integer invariant -> real measured value" (= the L4 equation of Section 7.1). Section 12.15 has now turned that merge into a **pre-registered, falsifiable test** and judged it negative, on three mutually independent grounds (reach bound / no single lock / hidden degree of freedom); see 12.15(7) and item 10 of the Key Logical Chain in Section 16. **This changes the shape of this item**: it is no longer "not yet attempted" but has been **tested and found not suppliable by enlarging the zero-parameter pool**. Only two open forms remain: (i) whether a **non-multiplicative** Ψ (containing additive, trigonometric or combinatorial terms, or a spectrum-dependent asymmetric function) can satisfy all

three targets at once - the negative verdict on the monomial class in this section does not cover such a class; and (ii) whether there exists a **prior-fixed** rule selecting the electron-layer representative (a non-arbitrary choice between the basal body Q_3 and the carrier M_{60}), which would remove the hidden degree of freedom referred to in (iii) and restore the zero-parameter status of the criterion. Until (ii) is resolved, even a hit within (i) would be non-falsifiable (C3 violated) and must not be used to advance the programme.

11. **A per-object checklist for the three-layer structure and the repair of the "geometry-shape" branch (added in v2.0)**: the three-layer decomposition of Section 14 is currently illustrated by the water molecule and a number of existing entries, and no general checklist for "giving a recomputable three-layer reading for any new object" has yet been provided; at the same time the repair direction indicated by the audit of that section (a purely topological D plus $n \geq 4$ plus **comparing angles / bond lengths directly**) has not yet been executed ⇒ this item is the immediate next step after the perspective has been established, and also the minimal action that would move the "geometry-shape" branch from 0 qualified entries toward qualification.

All conclusions are strictly confined to self-consistent constructions within the SRE model; real physical inferences must be independently verified by external means.

Appendix A: Key Numerical Tables

A.1 Core Physical Constants (CODATA 2018)

Quantity	Value	Unit
m_e	0.510998950	MeV/c²
m_p	938.27208816	MeV/c²
m_n	939.56542052	MeV/c²
α⁻¹	137.035999084	Dimensionless
a0	0.529177210903	Å

A.2 Key Numerical Values for Nucleon Inversion

Quantity	Value
Δm/m_p=(m_n−m_p)/m_p	1.378419305×10⁻³
κ_N=(m_n−m_p)/m_p/α	0.188893067
Electron calibration w"	1+4.347×10⁻⁵
Skeleton Π₁(closed)	0.188580
Deviation	0.165%
Combined rarity	4.7×10⁻³

A.3 Key Numerical Values for Fission Verification

Quantity	Value
Thermal cross-section dynamic range	8.58 orders of magnitude
Fast cross-section dynamic range	1.48 orders of magnitude
Pearson(Δ, $\log_{10}\sigma_{th}$)	+0.932
U-235 fission model energy	215–219 MeV
U-235 fission measured energy	202.5 MeV
Chain criticality p"(U-235)	0.412

A.4 Key Numerical Values for Dimension Collapse

Quantity	Value
Organic C skeleton RMS residual	1.5%
Light heteroatom RMS residual	6.7%
Same-nucleus H/O/F stretching	+10.0% ~ +19.7%
Same-nucleus heavy main group RMS residual	3.3%
Ion pair RMS residual	14.4%
Core Δ(1→2)	0.352±0.037 a0
Core Δ(2→3)	0.232±0.058 a0

A.5 Key Numerical Values for the Many-Body Assembly Law and the Shared Ring

Quantity	Value
Variant A (two bodies, no shared edge)	V=22, E=35, β_1=14, T=6; λ_2=0.238442818; ρ=6.681330644
Variant D (two bodies, shared ring)	V=21, E=33, β_1=13, T=5, |Aut|=48; ρ=6.000000; $\lambda_2=2-\sqrt{3}$
Deuteron prediction of variant A	2.5867 MeV (measured 2.224566 MeV, 16.3% too high)
Sharing tax (own vs shared)	0.28 Δm_np = 0.362 MeV
Three/four-body linear law	V=9k+3, E=15k+3, T=2k+1, β_1=6k+1, |Aut|=6·2^k·k!
Lowest non-trivial mode	$\lambda_2 = 2-\sqrt{3}$ = 0.267949192431 (exact for k=2..6), multiplicity k−1
ρ sequence (k=2..6)	6.000000 / 6.925423 / 7.909516 / 8.908855 / 9.912634
ψ(k) (in units of Δm_np)	+0.967105 / +0.376461 / +2.795573 / +10.750837
Scorecard (Reading A)	7/8 overall; 5/5 in the A ≤ 3 sector; only mismatch ^{4}Li

| A=4 out-of-sample predictions | ^{4}H 29.0596 MeV, ^{4}Li 31.4244 MeV (both measured unbound) |

| Rarity of ρ among random graphs | 0 out of 2966 samples have ρ exactly the integer 6 |

| Identity (additive-profile pricing class) | $B(^4H) - B(^4He) = \psi(1) - \psi(2) = B(^3H) - B(^3He) = +0.590645$ $\Delta m_{np} = +0.7639$ MeV |

| Same identity (^{4}Li side) | $\psi(3) - \psi(2) = +2.419113$ $\Delta m_{np} = +3.1287$ MeV |

| A=4 isomorphism class | ^{4}He / ^{4}H / ^{4}Li all V=39, E=63, T=9, β_1=25, $\lvert Aut\rvert$=2304, ρ=7.909515966, $\lambda_2 = 2-\sqrt{3}$ |

| Residuals of composition-resolved interference linear bases | (C,S) 10.2%, (C,a) 3.3%, (I,a) 24.6%, (C,I) 10.2%; extrapolated signs for ^{4}H/^{4}Li all wrong |

| Minimal repair term | imbalance term $\lambda \cdot \max(0, a-1)$, $a = \lvert n_n - n_p\rvert$, $\lambda \geq 29.0596$ MeV (fitted parameter, not a prediction) |

| Dual-service edges of variant A | 1 (the shared-ring configuration has 3) |

| Recomputed ρ of shared-ring configurations (k=2..6) | 6.000000 / 6.925423 / 7.909516 / 8.908855 / 9.912634; λ_2 identically $2-\sqrt{3}$ (deviation ca. 10^{-15}) |

| Shared-ring products of n-p and p-p | Isomorphic (V=21, E=33, T=5, β_1=13, $\lvert Aut\rvert$=48, ρ=6, $\lambda_2 = 2-\sqrt{3}$ all identical) |

| Threshold for the antisymmetric coefficient λ | $\lambda \geq B_{ledger}(^4H) = 22.468780$ $\Delta m_{np} = 29.0596$ MeV (**exceeding** $B(^4He) = 21.878135$ $\Delta m_{np} = 28.2957$ MeV) |

| Comparison with the empirical asymmetry energy | $a_{sym} \leq 24$ MeV < 29.0596 MeV (5.06 MeV, i.e. 17.4%, short); the mechanism also gives a non-zero 7.7 MeV at A=3, conflicting with the ^{3}H calibration |

| Statistics of the λ candidate pool (16 natural candidates) | 11 below the threshold, 5 above; the threshold falls in the gap between 21.878135 ($B(^4He)$) and 23.598162 Δm_{np} ($B(^4He)+B(deuteron)$) |

| Criterion L1 (antisymmetry criterion) | $\lvert n_n - n_p\rvert \leq 1$; **8 / 8** on the 8-entry scorecard (Reading A 7 / 8) |

| Per-interval hits of L1 (scorecard + A≥5 out-of-sample, 34 entries) | A ≤ 4: 8/8; A = 5: 1/3; A = 6–9: 8/11; A = 10–14: 5/12; **total 22 / 34** (Reading A 14 / 34) |

| Mismatch diagnosis for L1 | The mismatches at A = 5 (^{5}He, ^{5}Li) are an α closure effect (no bound nucleus at A = 5); the mismatches for A ≥ 8 arise from the absence of an A scaling |

| Scan over $(N-Z)^2/A \leq c$ | The two-end constraints are incompatible (A = 4 requires c < 1, drip-line nuclei require c ≥ 2.571); over $c \in \{0.25...3\}$ the best total is 22 / 34 |

Appendix B: Consolidated Terminology List

This appendix consolidates all term definitions used in this paper, sorted alphabetically:

| Term | Definition |

|---|---|

| **Binary Self-Organizing Network** | The ontology of SRE: a network whose elements are always ±1 and which grows continuously in time; all things are its resident structures |

| **β_1 (First Betti Number)** | $\beta_1 = E - V + 1$, the number of independent closed loops in the graph |

| **Closure Degree Potential** | In graph-theoretic language, the "rebalancing hill height": whether a complex is worth/able to rearrange to a lower closure degree configuration |

| **Compound Nucleus** | The temporary state (A+1) formed when a target nucleus absorbs a neutron; fission occurs inside it |

| **Covalent Radius** | The radius of an atom in a covalent bond, normalized by a0 into a dimensionless relational number |

| **Assembly Law** | The rule that merges several nucleon skeletons into a many-body configuration; determined by two principles: open–closed complementarity plus the images of the two bodies must be distinguishable in the shared region |

| **Gap** | At the same shared position, the number of gaps g left by open-state (neutron) nucleons |

| **Supply** | At the same shared position, the number of complete ring edges s offered by closed-state (proton) nucleons |

| **Shared Ring** | One closed triangular ring shared by several nucleons after assembly |

| **Shared Ledger** | A counting table recording how many bodies "claim" each overlapping edge on the shared ring; it does not change the graph structure |

| **Interference Depth** | The depth to which several bodies overlap on the same shared unit; a candidate pricing quantity for binding energy, not yet calibrated |

| **Dimension Collapse** | Organizing real empirical bond length data into dimensionless relational numbers in units of a0, and verifying additivity/step/integer-ratio structures |

| **Excitation Surplus Δ** | Δ = E" − B_f; Δ>0 means a low-energy (thermal) neutron can cross the fission barrier |

| **Fission Cross-Section σ** | The probability area that a neutron (seed) incident on a nuclide triggers fission; varies with energy |

| **Fission Barrier B_f** | The energy hill a compound nucleus must climb in order to deform into fission |

| **Dormant Edge** | A channel with χ=0 in SRE: it is not deleted but degenerates to the multiplicative identity 1 — conductive but not forming a loop |

| **Laplacian Spectrum** | The set of eigenvalues of a matrix built from the graph's adjacency relations, written $\lambda_1=0 \le \lambda_2 \le \dots \le \lambda_{max}=\rho$ |

| **Neutron Separation Energy S_n** | The energy required to remove the last neutron from a compound nucleus; the excitation energy after thermal-neutron absorption is E"=S_n |

| **$\Pi_1 = \lambda_2/\rho$** | Graph functional: the ratio of the smallest non-zero eigenvalue to the largest, calibrated as α on the electron side |

| **Projection (Functional)** | A rule that maps a graph to a number; the graph changes, the number changes |

| **Multiplication Constant ν** | The average number of neutrons emitted per fission |

| **Critical k_eff** | The ratio of the neutron count in each generation to that of the previous generation; ≥1 gives a self-sustaining chain |

| **Graph G** | A mathematical object composed of vertices and edges |

| **V, E** | Number of vertices V, number of edges E |

| **Degree k** | The number of edges incident to each vertex; k-regular = all vertices have the same degree |

| **Automorphism Aut(G)** | A permutation that maps a graph onto itself without changing the connectivity relations; $\lvert Aut \rvert$ measures the degree of symmetry |

| **Vertex Orbit** | A grouping of vertices that can be interchanged under all symmetry operations |

| **Inversion** | Given the value of a functional, solving inversely for the structural parameters of the graph |

| **sim_p Generator** | The core evolution algorithm of the binary self-organizing network, injecting random differences and expanding outward at every step |

| **Antisymmetric Penalty Term** | An additional term depending only on the compositional imbalance $a = \lvert n_n - n_p \rvert$ and vanishing identically for $a \le 1$, $g(a) = \lambda \cdot \max(0, a - 1)$; its coefficient λ cannot be derived from SRE (12.12) |

| **Criterion L1 (Antisymmetry Criterion)** | A shared-ring assembly is legal (predicted bound) if and only if $\lvert n_n - n_p \rvert \le 1$; zero-parameter and naturally n↔p symmetric |

Appendix C: Code List and Complete Reproduction

All reproducible scripts of this paper are archived in the `code/` directory; the only dependencies are Python 3 + NumPy + NetworkX (no torch dependency). The running command is `python -u <script.py > log 2>&1`.

C.1 Nucleon Inversion Related

| Script | Function |

|---|---|

| `_sre_nucleon_skeleton_inversion.py` | Main nucleon inversion recalculation: electron calibration, criteria ①②③, rarity, Q_3 check |

| `_sre_nucleon_tiebreak_scan.py` | Non-spectral invariant comparison and cross-check of the parallel candidates |

| `_sre_nucleon_tiebreak_v2.py` | Tie-break v2: degeneracy structure of the spectra, first statement of Criterion ④ |

| `_sre_nucleon_orbit_reading_check.py` | Uniqueness of per-orbit readings and closed-form search for the λ_2 drop |

| `_sre_nucleon_release_rarity.py` | Pool rarity statistics of the transition release amounts |

| `_sre_nucleon_transition_residue.py` | Transition release and topological residue test (invariance of ρ) |

| `_sre_nucleon_ringedge_transition.py` | Ring-edge transition analysis: ring-edge vs spoke readings, criteria ④⑤⑥, and full-pool rarity |

| `_sre_nucleon_skeleton_plot.py` | Generation of the skeleton structure schematic and the screening funnel figure |

| `sre_nucleon_skeleton_inversion_results.json` | Main nucleon inversion result data |

| `sre_nucleon_tiebreak_results.json` | Parallel candidate comparison data |

| `sre_nucleon_ringedge_transition_results.json` | Ring-edge transition analysis data |

C.2 Many-Body Assembly Law and Shared Ring Related

| Script | Function |

|---|---|

| `_sre_nuclear_data_survey.py` | Survey of usable nuclear data: screening of dimensionless candidate quantities, false-positive cleanup, and structural findings |

| `_sre_nucleon_dimer_rule.py` | Two-body assembly law: enumeration of five variants and screening by the two principles |

| `_sre_nucleon_dimer_spectra.py` | Spectra and invariant comparison of the two-body variants (A, D) |

| `_sre_dimer_full_enum.py` | Full enumeration of two-body constructions and verification of the mapping direction |

| `_sre_nucleon_sharedring.py` | Structure and full spectrum of the shared-ring configuration (variant D) |

| `_sre_sharedring_check.py` | Automorphism orbit check of the shared-ring motif |

| `_sre_sharedring_motif.py` | Comparison of the D-type products of n-n / p-p and arithmetic rigidity |

| `_sre_nucleon_trimer.py` | Construction of the k-body shared-ring configuration and verification of the structural linear law |

| `_sre_nucleon_trimer_price.py` | Structural law and pricing inversion (an alternative parameterization) |

| `_sre_nucleon_trimer_mode.py` | Variation of the lowest non-trivial mode with the body count k, and its zero-amplitude support |

| `_sre_nucleon_ledger_price.py` | Shared-ledger pricing: inversion of ψ(k), A=4 out-of-sample, and the existence scorecard |

| `_sre_nucleon_ledger_laws.py` | Pricing-law tests: deviations of the linear and pairwise-counting laws, A=4 out-of-sample |

| `_sre_nucleon_manybody_verdict.py` | Closure of the assembly-law scorecard: sector statistics of the three readings and the adopted boundary |

| `_sre_nucleon_shadow_disposal.py` | Shadow-solution disposal: isomorphism check between D and the k=2 member, discriminating power of the non-spatial restatement of principle 2, ledger difference between variant A and the shared ring, and the A=4 single isomorphism class |

| `_sre_nucleon_interference.py` | Interference-depth test: identity proving the non-repairability of the additive-profile pricing class, residuals of composition-resolved interference linear bases, the A=4 isomorphism class, and the minimal repair term with its price |

| `_sre_nucleon_manybody_plot.py` / `_sre_nucleon_manybody_plot_EN.py` | Figure 5 of Section 12: assembly schematic, ρ and λ_2 versus body count, zero-parameter scorecard; the spectral quantities in panel (b) are recomputed numerically inside the script |

| `_sre_nucleon_asymmetry.py` | The attempt at an antisymmetric coefficient: the nature of the λ threshold and its comparison with the empirical asymmetry energy, a 16-candidate selection check within SRE, the "shape vs value" separation argument for λ, the zero-parameter criterion L1 on the 8-entry scorecard and on the A ≥ 5 out-of-sample set (26 entries), and the $(N-Z)^2/A \le c$ scan |

| `_sre_nucleon_electron_coupling.py` | The configuration-level test of Section 12.13: measurement of the automorphism orbit of the three rings, re-running of the four two-body constructions (S1 variant A / S2 open-band attachment / S3 shared ring / S4 open-to-open) for r = 0,1,2, the verdict S2 ≅ S1 and its mechanism, the diagnosis that the k=1 edge in the S3 ledger is the locus of the open band, and the isomorphism re-check of the three A = 4 compositions |

| `_sre_nucleon_latin_ckm.py` | First cut of recommendation ②: the "mixing-table" test of the Latin-square perfect matching — bijectivity of the Latin square, parameter counting (C3), the assembly-level isomorphism-class scan (closed+closed 1 class / closed+open 1 class / **open+open 2 classes**), and the zero-parameter local-response mismatch matrix compared with the CKM; the conclusion is that "zero parameters and continuous angles cannot be had together" |

| `_sre_boson_transition.py` | Recommendation ③: the test of "boson = transition quantum" — falsifying "W/Z are the two states of one skeleton", the **whole-set hit-rate check** of intra-family transition ratios (expected 5.45 versus measured 6, no structural signal), m_H/m_W falling outside the range; the positive by-product being "a pure transition is massless ⇒ the photon is unique" |

| `_sre_anchor_registry.py` | External-anchor registry: empirical demonstration of the two-layer double role of α (zero-parameter reverse prediction of Δm_np/m_p to −0.165%), the scale sensitivity of the Criterion ① anchor (an MS-bar quantity; changing the scale changes V), the non-uniqueness of the V=12 family for κ_N, and the four-criterion registry table of anchors |

| `_sre_anchor_second.py` | Search for the second anchor plus the self-consistency of the Criterion ② window: **tolerance sharpening** (for ε ≤ 1% together with Criterion ③ only the V=12 bin retains survivors; exact for V=8/10/12 full sets, and LR ≥ 1.7/0.83 lower bounds for V=14/16), **within-bin individual identification** (all 3 hits at ε ≤ 1%, 2 survivors under Criterion ③ ⇒ Criterion ③ alone cannot single out Y_3; only after adding T ≥ 2 does it become unique = the Y_3 skeleton), the **V-sharp / V-blind likelihood-ratio criterion** (the spectral reading of α has LR only 1.7–3.3), the **three rounds of testing of the μ family** (whole-set hit-rate check plus two Monte-Carlo null hypotheses; the structural blocking by ^{3}H ≅ ^{3}He; the family-law test of ledger functionals, 0 passing), and the external falsification of the sign trend (14/16, counterexamples ^{13}C, ^{15}N) |

| `_sre_neutrino_k1.py` | Section 12.14: criterion construction for the neutrino as the k = 1 singly-claimed edge — gate check C1/C2/C3, direct measurement of the claim triples of six assemblies (always (#closed, k, k), 2 distinct values), exhaustive pairwise verification of the equivalence of the three rings, measurement of the 2+1 multiplicity of the 3×3 local operators (G_3 / L_{eff}), measurement of the automorphism **orbit count = 1** (vertex-transitivity) on the three ring points, hit testing of 96 zero-parameter candidates and two MC null hypotheses (randomized target p_A = 0.166; random graphs with the same degree sequence, hit rate 0.385) |

| `_sre_decay_law.py` | Section 13.3: axiomatic derivation and MC verification of the exponential decay law (survival-curve deviation 1.03×10^{-3}, half-life-identity deviation 0.0224%, memorylessness deviation 5.7×10^{-4}), exclusion of the power law (log-log piecewise slope −0.570 → −5.013), verification of $T_{1/2} \propto 1/N$, the **species-cancellation law** of the two-factor decomposition and the test of the stripping enhancement factor E (Re vs Dy, same mode and nearly equal Q, differing by ≥ 6.79 orders of magnitude), recomputation of the three-layer medians (+11.10 / −0.09 / −2.30, span 13.40) and refinement of the ^{163}Dy lower bound |

| `_sre_nucleon_ec_env_test.py` | The empirical basis of the three-layer stratification of Section 10 (added to this list in this round): single-parameter stratification by "s-electron reachability at the nuclear site" (L3 bound-state / L2 EC-IC / L1 neutral β^--α) and verification of strict monotonicity; it also registers the correction of the two hand-filled errors of v1 |

| `_sre_l4_assignment.py` | Section 12.15: the **pre-registered criterion** and negative verdict for the L4 mass-depth assignment rule ($m_p/m_e = \Psi(G_N)/\Psi(G_e)$) - ten layer-agnostic primitive invariants, a zero-parameter monomial pool (exponent $|a| \leq A$, support <= 3), two configurations (electron basal body Q_3 / α carrier M_{60}), and three targets (one design plus two out-of-sample); the **closed-form reach bound** ($\exp(A \sum |\log r_i|)$): configuration A has a full-support bound of 511.06 for $|a| \leq 2$, below 1836.15 (short by 0.555 orders of magnitude) => a distribution-free decisive exclusion; configuration B's pool sizes 576/4064/13152 ($A = 1, 2, 3$) with the **joint hit over the three targets identically 0** (likewise at the 1% and 3% windows) and different best Ψ per target; the prediction of Π_1 itself (0.5657 / 25.84, short by 1.85 orders of magnitude); the MC null over 407 random graphs with the same (V, E) (T_1 0.40, T_2 21.46, T_3 94.91 per trial); and the positive-residue control ($\alpha \cdot \Pi_1(\mathrm{Y}_3)$ against $(m_n - m_p)/m_p$: -0.164%) |

| `_enum_cubic12.py` | **Completeness verification** for V=12: a 2-switch closure BFS exhaustively enumerates **all 85 isomorphism classes** of connected cubic graphs on 12 vertices (matching the known count) and establishes that in this bin the **85 isomorphism classes correspond to 85 distinct Laplacian spectra (no cospectral pair)** ⇒ the "spectral-signature deduplication" of the main script is lossless in that bin; it also yields the 3 hits at ε ≤ 1% and the conclusion that adding T ≥ 2 points uniquely to Y_3 |

C.3 Fission Chain Verification Related

| Script | Function |

|---|---|

| `_sre_fission_chain_validation.py` | Recalculation of the fast-neutron experiment, parity counterexamples, excitation surplus criterion, B/A closure potential and chain criticality |

| `_sre_fission_chain_plot.py` | Generation of the excitation surplus scatter plot and the dynamic range comparison figure |

| `sre_fission_chain_validation_results.json` | Fission verification result data |

C.4 Dimension Collapse Test Related

| Script | Function |

|---|---|

| `_sre_dimension_collapse.py` | Basic dimension collapse test with five checks |

| `_sre_dimension_collapse_ext.py` | Extended dimension collapse test (full bond length library) |

| `sre_dimension_collapse_summary_ext.json` | Statistical data of the extended dimension collapse test |

C.5 Nucleon v1 Legacy Scripts

| Script | Function |

|---|---|

| `_sre_nucleon_derivation.py` | Nucleon topological derivation v1 (3×Q_3 construction, superseded by v2) |

| `sre_nucleon_derivation_results.json` | v1 result data |

C.6 Observational Perspective and Validation Qualification Related (Section 14)

| Script | Function |

|---|---|

| `_sre_q3_provenance.py` | Example 8 of 14.6: provenance of Q_3 — reproducible P0 enumeration (19320 → 5 classes), the equivalent construction $Q_3 = \mathrm{BDC}(K_4) = \mathrm{D}^2(K_2)$, the planarity cutoff of the doubling tower (planar ⇔ $k \leq 3$), $\mathrm{lift}(Q_3) = \mathrm{D}(Q_3) = Q_4$ (16/32/17), and $\Pi_1(Q_3) = 1/3$ generated from the K_4 spectrum |

| `_sre_collapse_in_nucleon.py` | Example 9 of 14.6 and 12.15(f): consequences of folding (collapse) entering the nucleon computation — D and BDC preserve vertex-transitivity (0 counterexamples over a battery of 21 graphs), BDC spectrum = base graph ± two copies, the exact laws on the tower ($|\mathrm{Aut}(Q_k)| = 2^k k!$, distinct L values $= k+1$, $\Pi_1(Q_k) = 1/k$), and a cross-layer scan over seven configurations (joint hit all 0, best same-lock ≥ 19.84%) |

| `_sre_dimension_selection.py` | 14.5: the zero-parameter predicate selection of $d = 3$ — J1 power-law model selection of the lattice voltage field r^{2-d} ($d = 1 \to -1.000$, $2 \to \log$, $3 \to +1.100$, $4 \to +1.900$), J2 finite resistance ($d \geq 3$ convergent), J3 bound-state threshold ($d \geq 3$ finite), J4 stable circular orbit ($d = 3$ only), intersection = $\{3\}$ |

| `_sre_coulomb_collapse.py` | 14.5: the shape/value trichotomy of the Coulomb force — leading term of the graph Laplacian Green function r^{2-d} ($d = 1 \to r$, $2 \to \ln r$, $3 \to 1/r$, $R^2 > 0.9999$), α still an external input, $\Pi_1(M_{60})$ deviating by 1.45×10^{-5} |

| `_sre_h2o_degeneracy.py` | 14.1–14.2: degeneracy determination for the water molecule at $n = 3$ — any symmetric bond weighting must give $180.0000°$, the bijection $\theta \leftrightarrow t$ (zero predictive margin), inversion fidelity r always 1 at $n = 3$, ineffectiveness of adding electron leaf nodes (minimum eigenvalue of the metric Gram matrix of the four-point graph $= -0.25$) |

| `_sre_h2o_geometry_provenance.py` | 14.4 and boundary 7 of Section 15: re-examination of the acquisition chain of the water "104°" — rotational spectrum → $A, B, C \to I = K/(A, B, C)$ → model step → (r, θ); four averaging conventions (103.9 / 104.478 / 104.50 / 105.12), rigid-rotor inversion $\Delta\theta = 0.6456°$, mixed triple giving $104.3279°$ (difference $0.146°$), $\cos\theta_e = -0.250001$ |

| `_sre_h2o_ratio_vs_geometry.py` | 14.1 and 14.3: recomputation of the three-layer decomposition and the two theorems — P_3 (degrees $(2, 1, 1)$, L spectrum $(0, 1, 3)$, orbit count 2), Theorem 1 (same graph, two metrics → 180.0000° / 73.7398°), Theorem 2 (apex-angle interval of the same ratio $t_0 = 1.581140$ under Euclidean / spherical / hyperbolic maps) |

| `_sre_validation_metric_audit.py` | 14.7: validation-qualification audit — 17 entries classified by epistemological type (E = 9 \| R = 2 \| C = 4 \| M = 2), the $n = 3$ distance-set Pearson magnitude always 1, classical MDS inversion fidelity always 1 for any Euclidean D, and the three qualification criteria |

| `_sre_orbital_emergence_test.py` / `_sre_orbital_measurement_test.py` | 14.4: the two senses of the electron orbital and the experimental side — $2\ell + 1$ is not any natural graph ladder (none of 12 candidates matches), a single orbital is unobservable (the gauge-invariant density is constant along each Aut orbit), resolution = number of orbits, calibrated graph orbit count = 1 ⇒ zero shape |

| `_sre_l4_pool_extend.py` | Section 14 and 12.15(f): the pool-size correction to the reach scope — expanding the invariants from 10 to 29 raises the full-support reach from 511.06 to 8.65×10⁶; the support ≤ 3 survival 91.209% agrees with the grid measurement 95.0326%; MC null 5% quantiles 0.1995% / 0.1822% |

| `_h2o_degeneracy.json` / `_h2o_geometry_provenance.json` / `_h2o_ratio_vs_geometry.json` / `_validation_metric_audit.json` / `_l4_pool_extend.json` / `_dimension_selection.json` / `_q3_provenance.json` / `_collapse_in_nucleon.json` | Result data of the above scripts (the same-named `.log` of each script is archived together) |

Appendix D: Figure List

All figures of this paper are archived in the `figures/` directory, ordered as they appear in the text:

| Figure | File Name | Content |

|---|---|---|

| Figure 1 | `sre_nucleon_skeleton_structure_EN.png/.svg` | Nucleon skeleton candidate: tripartite Y triangular closure (closed-state proton and open-state neutron) |

| Figure 2 | `sre_nucleon_skeleton_funnel_EN.png/.svg` | Nucleon skeleton three-step screening funnel (85→4→2) |

| Figure 3 | `sre_fission_chain_corr.png/.svg` | Nuclear reaction verification: excitation surplus vs thermal cross-section scatter plot, and fast/thermal cross-section dynamic range comparison |

| Figure 4 | `sre_nucleon_derivation.png/.svg` | Nucleon topological derivation v1 schematic (abandoned 3×Q_3 construction, historical reference only) |

| Figure 5 | `sre_nucleon_manybody_schematic_EN.png/.svg` | Many-body assembly law and the shared ring: (a) monomer → two bodies → four bodies sharing one ring (with single-claim edge annotations); (b) closure potential ρ rising with body count and the body-count independence of $\lambda_2 = 2-\sqrt{3}$; (c) zero-parameter scorecard of the assembly law (Reading A, 8 entries) |

| Figure 6 | `sre_observation_layers_EN.png/.svg` | Observational perspective: the boundary between the dimension-free SRE world and the three-dimensional world — (a) the three-layer decomposition (L0 topological → L1 ratio → L2 geometric) and the two tolls (the metric D, the map); (b) empirical demonstration of Theorem 2: the apex-angle interval of the same ratio t_0 under Euclidean / spherical / hyperbolic maps |

| Figure 6 (Chinese version) | `sre_observation_layers.png/.svg` | Chinese counterpart of Figure 6 (same plotting core, only the language labels switched) |

Appendix E: References and DOI

All literature and project references cited in this paper (in order of appearance):

1. SRE-v1.6 Axiom Suite, DOI 10.5281/zenodo.22077475
2. SRE-Dynamics: Composite Elementary Particles and Relational Space Emergence (Book of the Void), DOI 10.5281/zenodo.22162514
3. SRE Dynamics: Rigorous Reconstruction of Maxwell Field Equations Using Purely Dimensionless Graph Cohomology and Global Evolution Steps, DOI 10.5281/zenodo.22119957
4. Hierarchical Dissipative Self-Organizing Binary Network Dynamics, DOI 10.5281/zenodo.20576606
5. SRE Electrical Quantity Definitions (Charge/Current/Resistance/Voltage/Power/E=mc²), DOI 10.5281/zenodo.22119635
6. Complete Characterization of the Electron within the SRE Framework (Series), DOI 10.5281/zenodo.22162514
7. CODATA 2018 Fundamental Constants, https://codata.org
8. PDG Quark Mass Table, https://pdg.lbl.gov
9. JEFF-3.3 / ENDF/B-VIII Fission Cross-Section Database, https://www.oecd-nea.org/dbdata/jeff/
10. RIPL-3 Fission Barrier Database, https://www-nds.iaea.org/ripl-3/
11. AME 2020 Atomic Mass Evaluation, https://www-nds.iaea.org/amdc/

Archiving Note: This monograph is a stand-alone document; after review it may be merged into the *SRE_Electron_Complete_Paper* series or archived as an independent Zenodo entry. The Chinese and English versions are maintained in sync as a parallel pair.

Copyright Statement: This paper is released as open source under the CC BY 4.0 license; it may be freely used, distributed and modified, provided that the original author and the source are retained.

Mass Origin in the State–Relational–Entropy Framework

A Circulant-Operator Formulation on the Three-Ring $\mathbb{Z}_3$ Torsor

Version: 1.0 (Treatise Format)

Date: 2026-09-28

Abstract

Background and problem. Within the State–Relational–Entropy (SRE) framework, mass is treated as a **projected reading** of relational structure rather than an ontological input, and therefore in principle it should not be a number given in advance. However, the question "since mass is an emergent quantity, what exactly is its precise mathematical object?" had not previously been answered head-on. This paper answers that question and places the answer in a recomputable, falsifiable form.

Method. This paper takes the SRE nucleon skeleton $Y_3 \ltimes \triangle_3$ ($V = 12$, $E = 18$, $\beta_1 = 7$, $|\mathrm{Aut}| = 36$) as its only structural input. It first proves that the three shared rings of this skeleton form a **$\mathbb{Z}_3$-torsor** under the automorphism group — there is a cyclic order but no absolute origin; then, taking "respecting that structure" as the constraint, it derives the unique admissible form of the mass operator and sets out its spectrum, invariants and external comparisons.

Results. The paper gives four mutually independent, verifiable propositions: (1) **the object proposition** — a mass operator that respects the three-ring structure must commute with the cyclic shift S, and "commuting with S" holds if and only if it is a **circulant matrix**; hence $A = c_0\mathbb{1} + c_1 S + \bar{c}_1 S^2$, and $\mathbb{Z}_3$-equivariance compresses 9 matrix entries into 3; (2) **the shape proposition (with one correction)** — the spectrum of a Hermitian circulant is a set of 120° samples of one cosine, $\lambda_j = c_0 + 2|c_1|\cos(\theta + 2\pi j/3)$, but **this form holds trivially for any three generations** (three points, three unknowns, always solvable), so the "cosine shape" is a **free reparameterisation** rather than a constraint; (3) **the opening proposition** — writing the amplitude multiset as an ordered triple is equivalent to choosing an origin for the DFT characters, and the three choices give **exactly the same invariants**, so "opening" is a $\mathbb{Z}_3$ **gauge degree of freedom** rather than information; (4) **the invariant proposition** — the invariant moduli space is **two-dimensional**, $(\eta,\ \delta \bmod 2\pi/3)$, and the Koide combination $Q = \frac{1}{3} + \frac{2}{3}\eta^2$ probes only the single direction η. Measurement on the charged leptons gives $\eta^2 = 1/2$ to within 3.3×10^{-6}.

Boundary and conclusion. The paper also states two limitations that must be made explicit: **the cosine shape is free** (so the entire content of Koide collapses to **a single number** $\eta = 1/\sqrt{2}$); and **absolute mass (the scale c_0) does not lie inside SRE**, while **only renormalization-group-invariant dimensionless combinations qualify as candidate targets**. The overall conclusion is registered as **case 15 of the discrete-closure law (G12)**: "three" and the functional form are given (discrete side, closed), while the value of η and the scale are not (continuous side, requiring external input). All numerical verification in this paper **holds only within the SRE model**, and comparisons with charged-lepton masses are always stated as "structural consistency" rather than "numerical prediction".

Keywords: State–Relational–Entropy (SRE); mass origin; circulant matrix; $\mathbb{Z}_3$-torsor; triality; Koide relation; gauge degree of freedom; renormalization-group invariant; discrete-closure law.

1. Introduction

1.1 Statement of the problem

This paper answers the following proposition (raised 2026-09-28):

> **Original proposition.** The mass described by SRE is an evolutionary logic that is **unopened and cyclic**. What, then, is its precise mathematical description and expression?

This proposition combines three independently decidable sub-problems:

No.	Sub-problem	Corresponding section
P1	What **mathematical object** should mass be expressed as?	§4 (Theorem 2)
P2	What precisely does "**cyclic**" mean here?	§4.4 (Proposition 3)
P3	Which operation does "**unopened**" omit?	§6 (Propositions 5, 6)

1.2 Existing characterisations and their predicament

In the prior work of this project, three conclusions relating to mass already exist, but all of them remain at the level of **ratios** and do not touch the **object** itself:

Clue	Conclusion	Source
Graph functional	The three rings lie in one orbit ⇒ any graph functional gives the three rings **the same value** ⇒ Q collapses to the Cauchy–Schwarz lower bound $\frac{1}{3}$	Rounds 29, 30 [10][14]
Ledger	The information that the three generations have **unequal** masses can only be carried by **explicit bookkeeping**; graph functionals are **structurally blind** to it	Rounds 30, 31 [11]
Koide relation	$Q = \frac{1}{3} + \frac{2}{3}\eta^2$, independent of phase; $Q = \frac{2}{3} \iff \eta^2 = \frac{1}{2}$	Round 30 [10], ref. [1]

These three clues point together to an object that is **not yet named**: it is **not** three mass numbers, and **not** an invariant of any graph functional, but an **operator carrying the cyclic relation between the three generations**. The task of this paper is to name and characterise that object precisely.

The most naive scheme is to regard "the three-generation masses" as an **ordered triple** (m_0, m_1, m_2). This paper shows that the scheme is inaccurate at two levels:

1. **Ontological level**: there is no origin for "which generation" inside the SRE structure. The three rings are indistinguishable

under Aut (§3.1), and imposing labels introduces an **artificial origin**;

2. **Covariance level**: the output of SRE should **not depend** on artificial labels (objectivity / covariance).

The correct object should **carry a cyclic structure of its own and presuppose no origin** — and this is exactly how a **circulant matrix** is defined: it is characterised by "commuting with the cyclic shift", it does not itself single out an origin; an origin is introduced only at the moment of **reading** (spectral decomposition), and the manner of introduction is a gauge degree of freedom.

1.3 Contributions of this paper

Relative to the existing material, this paper gives four substantive items:

1. **It establishes "three rings = $\mathbb{Z}_3$-torsor" as a structural theorem** (Theorem 1), with an explicit witness and character decomposition (Propositions 1, 2);
2. **It makes the mass operator unique** (Theorem 2): "respecting the structure" directly forces a circulant matrix, with centraliser dimension exactly $n = 3$;
3. **One correction** (Theorem 3): it proves that "three points lie on one cosine" is a **free reparameterisation valid for any three generations**, and thereby narrows the content to be explained by Koide from "the shape" **down to a single number** $\eta = 1/\sqrt{2}$;
4. **One screening rule** (Theorem 5): only renormalization-group-invariant dimensionless combinations qualify as candidate targets, which clarifies the scope of the negative verdict on the quark side.

1.4 Structure of this paper

Section 2 gives the skeleton definition, notation and adjudication discipline, and builds intuition through an **analogy guide**; Section 3 proves the $\mathbb{Z}_3$-torsor structure of the three rings (first pillar); Section 4 proves the unique form of the mass operator (second pillar); Sections 5, 6 and 7 respectively treat the spectrum and the "free cosine", the boundary between "unopened" and "opened", and the invariant moduli space; Section 8 compares with charged-lepton data; Section 9 discusses the boundary on the quark side; Section 10 locates the single gap and gives falsification criteria; Sections 11 and 12 are discussion and conclusion. Appendix A is the recomputation list, Appendix B the honesty boundary, Appendix C the **analogy index**, Appendix D the references.

Notation convention: numbers followed by "**(proven, within-model)**" can all be recomputed in one command by the script in Appendix A; those marked "**(to be proven)**" are structural hypotheses proposed here but not yet delivered; those marked "**(negative verdict)**" are propositions already falsified or explicitly delimited as non-computable.

---

2. Preliminaries

2.1 The basic stance of SRE and the skeleton definition

SRE regards a physical object as a stable emergent state of a **binary self-organising network** in relational space, rather than a point particle placed in space in advance. Within this framework, "the foundations of classical physics originate from information statistics": geometry, charge, current, mass and the like are all treated as **projected readings** of relational structure rather than ontological inputs. This paper uses only one concrete product of that stance: **the nucleon skeleton and its open/closed two states**.

Definition 1 (nucleon skeleton $Y_3 \ltimes \triangle_3$). The skeleton is formed by **three** Y-shaped coherent cores closed into a **triangle**. Its vertex set and edge set are

$$\mathcal{V} = \{c_0, c_1, c_2\} \cup \{L_{ij} : \ i, j \in \{0,1,2\}\}, \qquad |\mathcal{V}| = 12,$$

where c_i is the **hub** of the i-th strand and L_{ij} is the **ring leaf** at position j of strand i. The edges fall into two classes:

- **strand edges** (hub–leaf): $c_i — L_{ij}$, 3 per strand, 9 in total;
- **ring edges** (leaf–leaf): ring j is the triangle $\{L_{0j}, L_{1j}, L_{2j}\}$, 3 per ring, 9 in total.

Hence $E = 18$, $\beta_1 = E - V + 1 = 7$, $|\mathrm{Aut}| = 36$ (proven, within-model); the spectral readings are $\rho = 5.302775638$, $\lambda_2 = 1$, $\Pi_1 = \lambda_2/\rho = 0.188580485$.

Definition 2 (open/closed two states). The **closed state** is the graph above, corresponding to the SRE proton state; the **open state** means removing one ring edge from the skeleton (here we fix the removal of (L_{0r}, L_{1r}), $r = 0$), so that $E = 17$, $\beta_1 = 6$, corresponding to the neutron state.

The full derivation of the skeleton (how integer-ness of the vertex number, the spectral ratio and triple symmetry jointly screen out the unique survivor) is given in §7–§8 of the project's nucleon paper [12] and is not repeated here. All that is needed here are **two properties**:

1. the three rings form **a single orbit** under Aut (closed state) — the three rings are **mutually indistinguishable**;
2. there exists in Aut an **edge-preserving automorphism of order 3**.

Together these are equivalent to: **the three rings are a $\mathbb{Z}_3$-torsor**. This is the **only** structural input of the whole paper.

2.2 Notation

Table 1. Table of symbols used throughout.

Symbol	Meaning
SRE	State–Relational–Entropy, the parent framework of this paper
$Y_3 \ltimes \triangle_3$	the nucleon skeleton (Definition 1)
L_{ij}	the ring leaf at position j of strand i
ring j	the triangle $\{L_{0j}, L_{1j}, L_{2j}\}$; there are three (**the three rings**)

| S | the **cyclic shift** on the three rings, $S^3 = \mathbb{1}$ |

| A | the **mass operator** (the central object of this paper) |

| φ | the explicit order-3 automorphism $L_{ij} \mapsto L_{i,\,j+1}$ (§3.2) |

| Aut | the automorphism group of the skeleton |

| $\mathrm{Circ}(n)$ | the space of n-th order circulant matrices, $\dim = n$ |

| ω | the cube root of unity $e^{2\pi i/3}$ |

| $\eta,\ \delta$ | the **amplitude ratio** and **phase** of the non-trivial component of the circulant operator: $c_1 = |c_1|e^{i\theta}$, $\eta = |c_1|/c_0$ |

| Q | the **Koide combination** $Q = \dfrac{\sum m}{(\sum \sqrt{m})^2}$ |

| $\rho,\ \lambda_2,\ \Pi_1$ | spectral radius, second-smallest Laplacian eigenvalue, closure degree $\Pi_1 = \lambda_2/\rho$ |

| D1 / D2 / D3 | adjudication discipline: small-denominator rationals listed separately / angles may not serve as anchors / the main statistics use only irrational readings × clean targets |

2.3 Analogy guide: three-phase alternating current

Before entering the formal argument, we set up a **system that can be checked in daily life**, in order to fix the intuition behind all the wording that follows. The system chosen is **symmetric three-phase alternating current** — which is not an analogical resemblance to the $\mathbb{Z}_3$ structure of this paper, but a **real instance of the same algebra**.

Analogy 1 (three-phase alternating current). Let the three phase voltages be

$$u_A(t) = U\cos\omega t, \qquad u_B(t) = U\cos\left(\omega t - \tfrac{2\pi}{3}\right), \qquad u_C(t) = U\cos\left(\omega t + \tfrac{2\pi}{3}\right).$$

In phasor representation, introducing the operator $a = e^{2\pi i/3}$, the three phase phasors are exactly $U,\ Ua,\ Ua^2$. This operator satisfies two identities:

$$a^3 = 1, \qquad 1 + a + a^2 = 0.$$

Comparing with the notation of this paper, $a = \omega$, and $1 + a + a^2 = 0$ corresponds exactly to the $\sum_k \cos(\delta + \frac{2\pi k}{3}) = 0$ used in §5.1. The three-phase system serves three demonstrative purposes for this paper:

1. **Cyclic but with no origin.** The phase sequence $A \to B \to C$ is cyclic: rename A as B and B as C, and **every physical quantity is unchanged**. That is, "which phase is called the first" **has no physical meaning** (= the **gauge degree of freedom** of this paper), whereas "the three phases differ by 120°" **does** have physical meaning (= the **torsor structure** of this paper).
2. **Decomposition is DFT.** The method of symmetrical components (Fortescue) uses a to decompose any three-phase signal into **positive-, negative- and zero-sequence** parts. Mathematically this decomposition is exactly the **third-order discrete Fourier transform**; its consequence is that every permutation-invariant, scale-invariant property of a three-phase system **can only** be determined by the amplitudes of these three components.
3. **The cost of "opening" is zero.** Since "which is the first phase" is a convention, **writing the three phases as an ordered triple adds no information whatsoever** — and this is precisely the proposition proved in §6 of this paper (opening = gauge, not information).

Example 1 (cosine reparameterisation of three numbers, toy example). Take the data $u = (1,\ 1.2,\ 0.7)$. Form the transform $z = \sum_k u_k \omega^{-k}$; measurement gives $|z| = 0.435890$, $\arg z = -1.455835$ rad, so that

$$\bar{u} = \tfrac{1}{3}\sum_k u_k = 0.966667, \qquad |c_1| = \tfrac{|z|}{3} = 0.145297, \qquad \theta = -\arg z = 1.455835 \text{ rad}.$$

Substituting back into $u_k = \bar{u} + 2|c_1|\cos\left(\theta + \frac{2\pi k}{3}\right)$ gives the multiset $\{1.000000,\ 1.200000,\ 0.700000\}$, consistent with the input (proven, within-model). Two points must be noted here first: (i) **any three numbers can be reparameterised in this way**, a triviality formally proved in Theorem 3; (ii) the **order** of the inverse output depends on the choice of origin (here the reconstructed order is $(1,\ 0.7,\ 1.2)$, different from the input arrangement), while **the multiset is unaffected** — and this is the simplest instance of the "opening = gauge" argument of §6.

2.4 Methodological discipline: the discrete-closure law (G12)

All results of this paper can be located by a methodological discipline already established in the project:

> **Discrete-closure law (G12).** Every link "discrete structure ⇒ discrete target" is **closed** within SRE; each crossing of "discrete → continuous" **must pay one external input**.

The role of this law is **division of labour**: for any result, it must be possible to say which half is given by SRE (the discrete side) and which half must be paid in externally (the continuous side). Section 11.4 will locate all conclusions accordingly.

---

3. First pillar: the three rings = a $\mathbb{Z}_3$-torsor

3.1 Orbit structure of the automorphisms

Fully enumerating the automorphisms of the closed-state and open-state skeletons respectively gives

Table 2. Skeleton invariants and three-ring orbit structure for the two states (proven, within-model).

State	V	E	β_1	$\lvert\mathrm{Aut}\rvert$	order-3 elements	order-2 elements	three-ring orbits
closed (proton)	12	18	7	**36**	**8**	15	**1 orbit:** $\{0,1,2\}$
open (neutron)	12	17	6	**4**	**0**	3	**2 orbits:** $\{0\}$ | $\{1,2\}$

In the closed state, the permutation induced by Aut on the three rings is **all six** (i.e. S_3); in the open state only two remain.

Columns 4, 6 and 7 of **Table 2** are the source of every structural argument in this paper and must be checked one by one: for the closed state $|\mathrm{Aut}| = 36 = 2^2 \cdot 3^2$, of which the **order-3 elements number 8** (Sylow-3 subgroup $\cong \mathbb{Z}_3 \times \mathbb{Z}_3$); for the open state $|\mathrm{Aut}|$ falls to 4 and the **order-3 elements vanish**. The number of three-ring orbits changes from 1 to 2, the most vivid trace left by "closed → open" at the structural level.

3.2 Explicit witness

Proposition 1 (explicit order-3 automorphism). The map

$$\varphi:\ L_{ij} \mapsto L_{i,\,(j+1) \bmod 3}, \qquad c_i \mapsto c_i$$

is an **edge-preserving** automorphism of the skeleton, and $\mathrm{ord}(\varphi) = 3$.

Proof. It suffices to verify each item. (i) **Vertex-preserving**: φ is a bijection on $\mathcal{V}$. (ii) **Strand-edge preserving**: $c_i - L_{ij} \mapsto c_i - L_{i,j+1}$, still within the strand-edge set. (iii) **Ring-edge preserving**: the triangle $\{L_{0j}, L_{1j}, L_{2j}\}$ of ring j is sent to $\{L_{0,j+1}, L_{1,j+1}, L_{2,j+1}\}$, i.e. ring $j+1$, still a triangle. (iv) **Order 3**: $\varphi^3(L_{ij}) = L_{i,j+3} = L_{ij}$, while $\varphi \neq \mathrm{id}$. □

The measured result is: edge-preserving = true, $\mathrm{ord}(\varphi) = 3$ (proven, within-model). This is a genuine order-3 automorphism, and it acts **freely and transitively** on the three rings (sending ring j to ring $j+1$).

3.3 The structural theorem

> **Theorem 1 ($\mathbb{Z}_3$-torsor structure of the three rings).** The three rings of the closed-state skeleton form a **$\mathbb{Z}_3$-torsor**: there is a **cyclic order** among them, but **no absolute origin**; their only degree of freedom is an **overall phase shift**.

Proof. Denote the three rings by the 3-element set $R = \{R_0, R_1, R_2\}$. By Proposition 1, $\langle\varphi\rangle \cong \mathbb{Z}_3$ acts on R; since $\varphi(R_j) = R_{j+1}$, the action is **transitive** (orbit $= R$). And since $|R| = 3 = |\langle\varphi\rangle|$, the stabiliser is trivial, so the action is **free**. A set acted on **freely and transitively** by a group G is by definition a G-torsor: it carries the structure of a **principal homogeneous space** for G — any two points are related by a unique $g \in G$, but **there is no base point fixed by the group action**. Taking $G = \mathbb{Z}_3$ gives the claim. □

Theorem 1 is the structural foundation of the entire paper. It gives the precise source of the shape of "three generations": **the three generations carry a $\mathbb{Z}_3$ cyclic order, but no absolute origin** — the origin is precisely the overall phase shift, corresponding to the phase δ in the Koide parameterisation below.

Analogy 2 (a three-strand braid and a clock face). In a braid woven from three strands, any strand may be named "the first", while the **interlacing relation** of the braid is unchanged; an object with "a cyclic order but no identifiable first strand" is a torsor. Another everyday instance is the **directions on a clock face**: the phrase "the three o'clock direction" is meaningful only after agreeing where "twelve o'clock" is — elements of a torsor are likewise such that one can speak of "how far apart" but not of "where".

3.4 Character decomposition under $\mathbb{Z}_3$

Let φ have character equal to the number of fixed points $\chi(\varphi)$ on some real permutation representation space. For $G = \mathbb{Z}_3$ (generator φ), the regular decomposition of the representation is $1 + \omega + \omega^2$, hence

$$\mathrm{mult}(\mathrm{trivial}) = \tfrac{1}{3}\big(\dim + 2\chi\big), \qquad \mathrm{mult}(\omega) = \mathrm{mult}(\omega^2) = \tfrac{1}{3}\big(\dim - \chi\big).$$

φ fixes the three hubs c_i and fixes no ring leaf, so for the vertex space $\chi = 3$; no edge is fixed by φ, so for the edge space $\chi = 0$. Measurement gives

Table 3. Character decomposition of $\mathbb{Z}_3$ on the vertex and edge spaces (proven, within-model).

Space	$\dim$	$\chi(\varphi)$	decomposition	trivial weight	non-trivial weight
vertex	12	3	$6 \cdot 1 + 3\omega + 3\omega^2$	$6/12 = \mathbf{1/2}$	$\mathbf{1/2}$
edge	18	0	$6 \cdot 1 + 6\omega + 6\omega^2$	$6/18 = \mathbf{1/3}$	$\mathbf{2/3}$

Proposition 2 (structural reading of the character decomposition). The weights above can be read off directly from the construction of the skeleton.

Proof. The skeleton = 3 strands × (1 hub + 3 ring leaves). The hubs are fixed by φ, so each hub contributes one copy of the trivial representation; the ring leaves are cycled by φ, so the three ring leaves of each strand contribute one copy of the regular representation $1 + \omega + \omega^2$. Hence per strand the trivial multiplicity is $1 + 1 = 2$ and the non-trivial multiplicity is $1 + 1 = 2$, so the **vertex space splits** $1/2 \mid 1/2$. Edge side: the strand edges $c_i - L_{ij}$ are sent to $c_i - L_{i,j+1}$ and the ring edges to the corresponding edges of the next ring, so **all edges are moved away by** φ ($\chi = 0$); hence the edge representation is 6 copies of the regular representation, with trivial weight $1/3$ and non-trivial weight $2/3$. □

Note 1 (registration discipline). The values $1/2$, $1/3$, $2/3$ in Table 3 are all **small-denominator rationals** (the D1 trap) and **have no discriminating power** on their own. Their possible meaning is registered in the candidate chain of §11.2; this paper **does not upgrade them to conclusions**.

---

4. Second pillar: the unique form of the mass operator

The conclusions of this chapter were first given in the form of a working note [13]; this paper reorganises them into treatise form, supplies the complete proof, and corrects one over-claim in them (see Theorem 3).

4.1 The main theorem

> **Theorem 2 (circulant criterion).** Let S be the n-th order cyclic shift matrix. Then an $n \times n$ matrix A commutes with S ($AS = SA$) **if and only if** A is a circulant matrix, i.e. A is completely determined by its first row:
>
> $$A = c_0\mathbb{1} + c_1 S + c_2 S^2 + \cdots + c_{n-1} S^{n-1}.$$

Proof. Write S for the permutation matrix sending the basis vector e_j to e_{j+1} (indices mod n). If A is circulant, then $A = \sum_k c_k S^k$, which obviously commutes with S. Conversely, suppose $AS = SA$. Take the first row of A to be $(a_0, a_1, \dots, a_{n-1})$; comparing column by column from $AS = SA$ gives $a_{ij} = a_{0,\, j-i \bmod n}$, i.e. row i of A is the first row cyclically shifted right i times, hence $A = \sum_k a_{0,k} S^k$. □

Numerical verification ($n = 3$). Writing the constraint $AS - SA = 0$ as a homogeneous linear system in $\mathrm{vec}(A)$,

$$\left(S^{\mathsf{T}} \otimes I - I \otimes S\right) \mathrm{vec}(A) = 0,$$

measurement gives **rank 6** for this coefficient matrix in the 9-dimensional space, so the solution space has dimension $= 9 - 6 = \mathbf{3} = n$; taking a basis of the null space and reshaping it into matrices, **all of them are circulant** (the check is true) (proven, within-model).

The substance of Theorem 2 is a quantification of **constraining power**: $\mathbb{Z}_3$-equivariance compresses the $n^2 = 9$ matrix entries down to $n = 3$. This is the **entire** constraining power carried by the word "cyclic" in SRE — no more, no less.

4.2 Necessity of the mass operator

By §3.3 the three rings are a $\mathbb{Z}_3$-torsor. A mass operator that **respects that structure** (does not conflict with the cyclic shift) is by definition one that "commutes with S"; by Theorem 2 it **can only** be a circulant matrix. Taking the Hermitian form (whose necessity is shown in §5.2):

> **Corollary 1 (object proposition).** The unique admissible form of the mass operator is
>
> $$A = c_0 \mathbb{1} + c_1 S + \bar{c}_1 S^2, \qquad c_0 \in \mathbb{R}.$$

This is not a language that has been picked, but a result **forced by the structure**. At this point sub-problems **P1 (object)** and **P2 (cyclic)** both have precise names.

Example 2 (a concrete mass operator). Take $c_0 = 1$, $c_1 = 0.3\, e^{i \cdot 40^\circ}$; then

$$A = \begin{pmatrix} 1 & 0.3e^{i40^\circ} & 0.3e^{-i40^\circ} \\ 0.3e^{-i40^\circ} & 1 & 0.3e^{i40^\circ} \\ 0.3e^{i40^\circ} & 0.3e^{-i40^\circ} & 1 \end{pmatrix}.$$

The matrix is **determined by only two real numbers** (c_0 and $|c_1|$, plus the phase θ), whereas a 3×3 matrix has 9 entries in general. Its measured eigenvalues are

$$(0.436184,\ 1.104189,\ 1.459627) \quad (\text{ascending}),$$

matching term by term the analytic form of Proposition 4, $\lambda_j = 1 + 0.6 \cos(40^\circ + 120^\circ j)$ (proven, within-model). This 3×3 matrix is the **smallest writable instance** of the "mass" asserted in this paper.

4.3 Why "evolution" is cyclic (a group) rather than one-way (a semigroup)

The word "evolution" easily suggests a time arrow, so its precise meaning must be made clear.

Table 4. One-way evolution versus cyclic evolution.

	one-way evolution (semigroup $\mathbb{N}$)	cyclic evolution (group $\mathbb{Z}_3$)
starting point	yes ($t = 0$)	**no**
reversible	no	**yes** ($S^{-1} = S^2$)
closed	no	**yes** ($S^3 = \mathbb{1}$)
carrier	a directed interval	**a cyclic group action on a torsor**

Proposition 3 (answer to P2). The so-called "cyclic evolutionary logic" is mathematically the use of a **group ($\mathbb{Z}_3$) rather than a semigroup** to characterise the index space.

Proof. The defining element of a semigroup is "there exists an irreversible starting point", whereas the defining element of the group $\mathbb{Z}_3$ is "$S^3 = \mathbb{1}$, $S^{-1} = S^2$, and no element is fixed by the action". By Theorem 1, the three rings are exactly a $\mathbb{Z}_3$-torsor, so their index space possesses only the second set of properties. □

This conclusion also explains the "**absence** of opening": once an origin is introduced or an ordered labelling taken, the cycle degenerates into a **linear sequence with a starting point** — and that is precisely "opening" (§6).

Analogy 3 (a wheel versus an hourglass). The typical image of semigroup evolution is an **hourglass**: the sand flows in one direction only, start and end are asymmetric, and it is irreversible. The typical image of group evolution is a **wheel with three equal divisions on its rim but no mark for "twelve o'clock"**: every rotation can be turned back, three rotations return to the start, and **no division on the dial is special**. The mass discussed in this paper belongs to the latter.

---

5. The spectrum and the "free cosine"

5.1 Spectral form

Proposition 4 (spectrum of a Hermitian circulant). Let $A = c_0\mathbb{1} + c_1 S + \bar{c}_1 S^2$ with $c_0 \in \mathbb{R}$, $c_1 = |c_1|e^{i\theta}$. Then the eigenvalues of A are

$$\lambda_j = c_0 + c_1\omega^j + \bar{c}_1\omega^{-j} = c_0 + 2|c_1|\cos\!\left(\theta + \tfrac{2\pi j}{3}\right), \qquad j = 0, 1, 2.$$

Proof. The eigenvectors of S are the discrete Fourier basis $v_j = (1, \omega^j, \omega^{2j})^{\mathsf{T}}/\sqrt{3}$, with eigenvalues ω^j ($j = 0, 1, 2$). Any polynomial in S shares these eigenvectors, so $Av_j = \left(c_0 + c_1\omega^j + \bar{c}_1\omega^{-j}\right)v_j$. Using $\omega^{-j} = \overline{\omega^j}$, we get $c_1\omega^j + \bar{c}_1\omega^{-j} = 2\,\mathrm{Re}\left(c_1\omega^j\right)$ $= 2|c_1|\cos(\theta + 2\pi j/3)$. □

Numerical verification (6 random parameter sets against the analytic form): all Hermitian checks true, maximum error 1.1×10^{-15} (proven, within-model).

The intuitive reading of **Proposition 4** is: **the three eigenvalues lie on the 120° samples of one and the same cosine**.

5.2 Real-symmetric versus complex phase: why a phase is necessary

Proposition 5 (real-symmetric degeneracy). If A is a real-symmetric circulant (i.e. $\theta = 0$), its spectrum **always has two values**:

$$\left(c_0 + 2|c_1|, \quad c_0 - |c_1|, \quad c_0 - |c_1|\right).$$

Proof. At $\theta = 0$, $\lambda_0 = c_0 + 2|c_1|$, while $\lambda_1 = c_0 + 2|c_1|\cos(2\pi/3) = c_0 - |c_1|$ and λ_2 is the same (since $\cos(4\pi/3) = \cos(2\pi/3)$). □

Measured comparison: the real-symmetric choice $c_0 = 1, |c_1| = 0.3$ gives the spectrum $(0.7,\ 0.7,\ 1.6)$, a **2+1** split; the complex phase $\theta = 1$ gives the spectrum $(0.400668,\ 1.27515,\ 1.324181)$, a **1+1+1** split (proven, within-model).

> Proposition 5 directly answers "why a phase is needed": **the spectrum of a real-symmetric circulant degenerates structurally and cannot carry three distinct generations; for the three generations to be distinct the operator must be Hermitian rather than real-symmetric** (thereby giving the Hermitian form of Corollary 1 its necessity).

Note 2 (registration, not conclusion). Proposition 5 and the "closed-state $\mathbb{Z}_3$ / open-state $\mathbb{Z}_2$" of Table 2 echo each other at the level of operators: the open state has only two values (a $\mathbb{Z}_2$-type degeneracy), the closed state generally three (the full $\mathbb{Z}_3$). This paper has **not proved** that the two are the same mechanism (see Appendix B item 4) and merely registers the observation.

5.3 A correction that must be stated: the cosine shape is "free"

Theorem 3 (triviality of the shape). The trigonometric form $\lambda_k = c_0 + 2|c_1|\cos(\theta + \frac{2\pi k}{3})$ has a solution for **any** three real numbers $\lambda_0, \lambda_1, \lambda_2$; that is, the form **contains no constraint whatsoever**.

Proof. Given the data λ_k, form

$$\textstyle\sum_k \lambda_k = 3c_0, \qquad z := \sum_k \lambda_k\,\omega^k = 3|c_1|e^{-i\theta},$$

i.e. $c_0 = \overline{\lambda}$, $|c_1| = |z|/3$, $\theta = -\arg z$. The three unknowns $(c_0, |c_1|, \theta)$ exactly match the three data, so generically the solution is unique and is given explicitly by the formulas above. □

Measurement: taking 8 random sets of **arbitrary** three numbers, solving by the formulas above and substituting back to reconstruct, the maximum error is of order 10^{-15} (the companion script of this paper gives 5.11×10^{-15}, a preceding script 2.22×10^{-15}, the same order of magnitude) (proven, within-model).

> **Correction (direct consequence of Theorem 3).** "Three points lie on one cosine" is **not** a constraint delivered by cyclicity, but the **standard reparameterisation of three numbers**. Hence saying "$\mathbb{Z}_3$ gives the shape of Koide" is an **over-claim**. The **content** of Koide is not the shape but the number

> $\eta = \frac{|c_1|}{c_0} = \frac{1}{\sqrt{2}} \qquad (\text{measured } 0.707104444 \text{ vs } 0.707106781,\ \text{deviation } -2.34 \times 10^{-6}).$

Thus the true contribution of $\mathbb{Z}_3$ collapses to two items: **(a) why there are exactly 3 positions** (the torsor has size 3, not 2 or 4); **(b) why δ is gauge** (a torsor has no origin ⇒ the phase can only be taken mod $2\pi/3$). **The shape is trivial; "$\eta = 1/\sqrt{2}$" is the number that genuinely awaits explanation** — and it belongs to the **continuous side** of G12.

Analogy 4 (a prism and dispersion). The mass operator may be compared to a composite light beam made by mixing three pure colours in fixed proportions: without spectral decomposition you see only its **overall hue and brightness** (= permutation-invariant, scale-invariant quantities); performing the DFT is like **passing the beam through a prism**, and the three eigenvalues are the intensities of the three emerging beams. What Theorem 3 says is: **any set of three intensities corresponds to some composite beam** — the dispersing step contains no information, it merely restates the same thing in another language.

Note 3 (D1 discipline). $\eta = 1/\sqrt{2}$ is irrational (consistent with adjudication discipline D3); but $\eta^2 = 1/2$ together with the $1/3$, $2/3$ in Q are still small-denominator rationals, so the numerical agreement **does not count as independent evidence**. The gain of this section is **structural**: separating the "trivial shape" from the "real content".

---

6. "Unopened" and "opened": the boundary between gauge and information

6.1 Opening = choosing the DFT origin

Proposition 6 (gauge-independence of opening). Writing the **multiset** of amplitudes $\{\lambda_0, \lambda_1, \lambda_2\}$ as an **ordered triple** is equivalent to choosing an **origin** for the characters ω^k; there are 3 choices of origin, and the three choices give **exactly the same** permutation invariants.

Proof. Let (u_0, u_1, u_2) be one representative; the others are given by the cyclic shift $u_k^{(s)} = u_{k-s}$. For any function F depending only on the multiset (e.g. the Koide combination Q and the R of §7.1), multiset invariance gives $F(u^{(s)}) = F(u)$ immediately. □

Measurement: taking $\eta = 1/\sqrt{2}$, $\delta = 2/9$, the three representatives give

Table 5. Invariants under the three choices of DFT origin (proven, within-model).

representative (cyclic shift)	Q	$R = (\lambda^2_{\max} - \lambda^2_{\min})/\sum\lambda^2$

| shift 0 | 0.666666666667 | 0.943349649589 |

| shift 1 | 0.666666666667 ($|\Delta Q| = 2.2 \times 10^{-16}$) | 0.943349649589 |

| shift 2 | 0.666666666667 ($|\Delta Q| = 0$) | 0.943349649589 |

⇒ **the three choices give three representatives of one and the same equivariance class.** Hence "opening" changes no SRE observable; it only decides "which generation is called the first" — **this is gauge (redundancy), not information**.

6.2 The two levels of "opening" must not be conflated

Table 6. Strict distinction between the two kinds of "opening".

| Level | Operation | Consequence |

|---|---|---|

| **(i) gauge level** | choosing an origin = $\mathbb{Z}_3$ labelling | invariants **unchanged** (redundancy) — this is the level the original proposition refers to |

| **(ii) physical level** | closed → open (cutting one ring edge) | $|\mathrm{Aut}|$ falls from 36 to 4, order-3 elements from 8 to 0: a **real** breaking $\mathbb{Z}_3 \to \mathbb{Z}_2$ that **changes** the accessible invariants |

Analogy 5 (transposition of a score). The **interval structure** of a melody is unchanged by an overall transposition — after transposition it is still "the same piece". Choosing an origin for the DFT characters is like **fixing the tonic**: it decides "which generation is the tonic" without changing the internal structure of the piece; whereas the physical-level "opening" of §6.2 (cutting a ring) is different, being like **rewriting the score itself** — the structure of the piece really changes.

6.3 "Not opened" is not a defect but covariance

Combining §§6.1–6.2, we can answer **P3**:

> **Proposition 7 (answer to P3).** "Not opened" = **no eigenbasis (DFT origin) chosen**, and the SRE native output stops at the **multiset** level.

This contains two things of entirely different nature that were previously often conflated; this paper separates them completely:

- **Gauge undetermined** (the $\mathbb{Z}_3$ choice of δ) ⇒ the SRE output **does not depend on artificial labels**. This is an **advantage** (objectivity / covariance), **not a deficiency**;
- **Scale ungiven** (c_0) ⇒ this is the **real information gap**, and lies **outside SRE** (§9 has settled: absolute mass is not its responsibility).

---

7. Invariants and the moduli space

7.1 The output of "unopened" = the equivariance class

Take the parameterisation $\lambda_k = 1 + 2\eta\cos(\delta + \frac{2\pi k}{3})$ (c_0 already normalised; the scale is an external input). The following two quantities are invariant under **permutation** and under **overall scale**, and their measured behaviour is as follows.

Table 7. The invariant Q is immune to the phase δ (sweeping δ over a full turn).

η	range of Q over δ	Q	analytic value $\frac{1}{3} + \frac{2}{3}\eta^2$
0.30	3.3×10^{-16}	0.393333333	0.393333333
0.50	6.1×10^{-16}	0.500000000	0.500000000
$1/\sqrt{2}$	1.2×10^{-15}	0.666666667	0.666666667
0.80	1.6×10^{-15}	0.760000000	0.760000000

Table 8. The invariant R depends on δ (with $\eta = 1/\sqrt{2}$ fixed).

δ	Q	R
0.0	0.666666667	0.957106781
0.4	0.666666667	0.880903092
0.8	0.666666667	0.633929534
1.2	0.666666667	0.566017302

7.2 Correction: the moduli space is two-dimensional

> **Proposition 8 (dimension of the invariant moduli space).** The invariant moduli space is
>
> $$\left(\eta,\ \delta \mod \tfrac{2\pi}{3}\right),$$
>
> that is, **two-dimensional**.

Proof. By §7.1, Q depends only on η and not on δ (the order of magnitude $\sim 10^{-15}$ of the range in Table 7 confirms strict independence); while Table 8 shows that R varies monotonically with δ at fixed η, so $\delta \mod 2\pi/3$ is a second invariant **independent** of η. The sum of the two parameters is the dimension of the moduli space. □

> **Correction.** A previous reading of "SRE has only one invariant, Q" is an **over-simplification**. The correct statement is: Q **probes only the one direction** η (because it is immune to δ, and the triple permutation of δ is precisely the source of permutation invariance); $\delta \mod \frac{2\pi}{3}$ is a second independent direction.

7.3 A typology of invariance

Table 9. The invariance of various quantities and the layer they live in.

Quantity	Immune to	Layer
ρ (spectral radius)	**cutting a ring** (range over 18 cuts $\sim 10^{-15}$)	spectral layer
$\Pi_1 = \lambda_2/\rho$	graph isomorphism (Aut)	spectral layer
Q (Koide)	$\mathbb{Z}_3$ **origin** (triple permutation of δ)	torsor / map layer
η^2	$\mathbb{Z}_3$ origin	torsor layer
absolute mass c_0	— (immune to no structure)	**outside SRE**

The reading of **Table 9** is: **the quantities SRE can give internally all have "immunity to some structure" as their content**; and **the one quantity immune to no structure** (absolute mass c_0) is precisely the one **not belonging to SRE**. This table corresponds item by item to the gap of §10.

Analogy 6 (an altitude datum and relative height). The sentence "this mountain is 1000 m high" presupposes agreeing where "sea level" is; changing the datum from sea level to some valley floor changes every absolute height, while **the ratio of the relative heights of two mountains is unchanged**. The quantities SRE can give internally (Π_1, Q, η) are **relative quantities** (datum-independent); the absolute mass c_0 is an **absolute altitude** — it must first have an external datum to be meaningful. This is also the intuitive root of the "α has an anchor, mass has none" discussion of §11.3: the fine-structure constant is a **ratio** (datum-independent), whereas the absolute scale of mass is not.

---

8. The Koide relation and the charged leptons

8.1 Analytic derivation

Theorem 4 (Koide relation). Let $\lambda_k = c_0 + 2|c_1|\cos(\delta + \frac{2\pi k}{3})$ and $m_k \propto \lambda_k^2$. Then

$$Q = \frac{\sum_k m_k}{\left(\sum_k \sqrt{m_k}\right)^2} = \frac{1}{3} + \frac{2}{3}\eta^2, \qquad \eta = \frac{|c_1|}{c_0}.$$

Proof. From $\sum_k \cos(\delta + \frac{2\pi k}{3}) = 0$ and $\sum_k \cos^2(\delta + \frac{2\pi k}{3}) = \frac{3}{2}$ we directly get

$$\sum_k \lambda_k = 3c_0, \qquad \sum_k \lambda_k^2 = 3c_0^2 + 6|c_1|^2.$$

Substituting into the definition of Q (note $\sqrt{m_k} \propto \lambda_k$, so the denominator $= \left(\sum \lambda_k\right)^2$):

$$Q = \frac{\sum_k \lambda_k^2}{\left(\sum_k \lambda_k\right)^2} = \frac{3c_0^2 + 6|c_1|^2}{9c_0^2} = \frac{1}{3} + \frac{2}{3}\left(\frac{|c_1|}{c_0}\right)^2. \square$$

The point of **Theorem 4** is: Q **depends only on** $\eta = |c_1|/c_0$ **and not at all on the phase** δ. This is a direct consequence of the $\mathbb{Z}_3$ invariance property — by Proposition 8, Q probes only the η direction in the moduli space. In particular,

$$Q = \tfrac{2}{3} \iff \eta^2 = \tfrac{1}{2} \iff \tfrac{|c_1|}{c_0} = \tfrac{1}{2\sqrt{2}}.$$

Note 4 (one piece of empirical content). "Taking $m_k \propto \lambda_k^2$ (i.e. $a = \sqrt{m}$ rather than m)" is the **empirical content of Koide**, equivalent to the testable structural hypothesis that "**the square root of the mass operator** is $\mathbb{Z}_3$-equivariant". This paper **has not proved** it (see Appendix B item 2).

8.2 The equivalence chain: five ways of writing one constraint

$$Q = \tfrac{2}{3} \iff \eta^2 = \tfrac{1}{2} \iff |A| = \tfrac{1}{\sqrt{2}} \iff \text{angle with } (1,1,1) = 45^\circ$$
$$\iff Q \text{ at the midpoint of the C–S interval } [\tfrac{1}{3}, 1].$$

Note 5. The five items of this chain are **five languages for the same constraint**, **not five independent pieces of evidence**. They must not be listed side by side as mutually corroborating evidence.

Analogy 7 (forty-five degrees is the "midpoint deviation"). Denote by ϕ the angle between the vector $(\sqrt{m_0}, \sqrt{m_1}, \sqrt{m_2})$ and the equal-mass direction $(1,1,1)$. $\phi = 0^\circ$ corresponds to three equal generations (Q takes the C–S lower bound $\frac{1}{3}$, the spectrum degenerates), $\phi = 90^\circ$ to the largest possible deviation (Q takes the upper bound 1); and 45° is exactly the **midpoint** of this interval. The Koide relation can therefore be read as: **the square-root mass vector of the three generations stops exactly midway between "equal mass" and "maximal deviation"**. This is the two sides of one coin with the "real-symmetric degeneracy ⇔ complete equal mass" of §5.2 Proposition 5.

8.3 Comparison with charged-lepton data

Taking the PDG charged-lepton masses $m_e = 0.51099895069$ MeV, $m_\mu = 105.6583755$ MeV, $m_\tau \in \{1776.86,\ 1776.93\}$ MeV [9]:

Table 10. Recomputing the Koide combination for the charged leptons (proven, within-model).

Input	Q	$\eta^2 = (3Q-1)/2$	deviation vs $\frac{2}{3}$	angle with $(1,1,1)$
$m_\tau = 1776.86$ MeV (archive convention)	0.666660511	0.499990767	-6.155×10^{-6}	44.9997°
$m_\tau = 1776.93$ MeV	0.666664463	**0.499996695**	-2.203×10^{-6}	44.9999°

⇒ $\eta^2 = \frac{1}{2}$ holds on the real data to within 3.30×10^{-6}; the inverse solution gives $\eta = 0.707104444$ against $1/\sqrt{2}$ $= 0.707106781$, a deviation of -2.34×10^{-6}.

Example 3 (a complete recomputation). Take $m_\tau = 1776.93$ MeV. Then

$\sum_k m_k = 1883.099374, \qquad \sum_k \sqrt{m_k} = 0.7148419 + 10.2790260 + 42.1536475 = 53.147515435,$

$Q = \frac{\sum_k m_k}{\left(\sum_k \sqrt{m_k}\right)^2} = \frac{1883.099374}{2824.658397} = \mathbf{0.666664463}.$

The inverse solution gives $\eta^2 = (3Q-1)/2 = 0.499996695$, differing from $\frac{1}{2}$ by 3.3×10^{-6} (proven, within-model).

8.4 The phase (supporting evidence)

From $u_k = \dfrac{\sqrt{m_k}/a_0 - 1}{\sqrt{2}} = \cos(\delta + \frac{2\pi k}{3})$ and the DFT $z = \sum_k u_k \omega^{-k} = \frac{3}{2} e^{i\delta}$, the phase can be solved for. Measurement:

Table 11. Sensitivity of the phase solution to the "labelling convention".

Labelling convention	$\delta \pmod{2\pi/3}$	reference value $2/9$	deviation
ascending (e, μ, τ)	**0.222224762**	0.222222222	$+2.54 \times 10^{-6}$
descending (τ, μ, e)	1.872170340	0.222222222	$+1.65$ (wrong convention)

Note 6 (D1 discipline). $2/9$ is a small-denominator rational, and its agreement with δ **must not** serve as independent evidence; moreover δ is itself a **gauge quantity** (an overall phase shift = a permutation of the three generations). This paper registers it only as supporting evidence. (Registered together: $2/9 = \frac{2}{3} \cdot \frac{1}{3}$ is the product of two SRE structure constants, likewise in the D1 domain, registered only.)

Table 11 also demonstrates an operational warning: **using the wrong labelling convention (here the descending one) makes the phase deviate by 1.65 rad** — i.e. the "same δ" is a **completely different** number under a wrong origin. This is the practical manifestation of the "gauge undetermined" of §6.3, and it is also why every test using δ as a target must first fix the convention.

8.5 Strength levels of the theorem

- **Proven (within-model)**: $Q = \frac{1}{3} + \frac{2}{3}\eta^2$ is independent of the phase; $Q = \frac{2}{3} \iff \eta^2 = \frac{1}{2}$.
- **Consistent with data**: $\eta^2 = \frac{1}{2}$ holds to 3.3×10^{-6} — **structural consistency, not numerical prediction**.
- **Not proven**: why η^2 is exactly $\frac{1}{2}$ (see §10).

---

9. Boundary: the quark side and dimensions

9.1 The precise meaning of "emergent" in this project

According to the three ontological speculations of SRE (project nucleon paper §4 [12]), quarks are an **emergent statistical phenomenon** rather than an independent ontology: $N_c = 3$ is the **number of strands**, "flavour" is a statistical projection of open/closed breaking, and "fractional charge" is an equal-share reading of the three-strand shared loop. Hence "quark mass" **cannot** be an input of SRE and can only be an output. Outputs fall into two classes that must be sharply distinguished:

Table 12. Visibility of the two classes of output.

Output type	Nature	Can it see "which ring"?
graph functional ($\lambda_2, \beta_1, \rho, \Pi_1$)	automorphism invariant	**blind** (points in the same orbit take the same value)
ledger (explicit bookkeeping of edges)	not an invariant	**yes**

Table 12 is the skeleton of all the conclusions: any information of the kind "the three generations have unequal masses" **cannot structurally be given by graph functionals** and can only be carried by the ledger. This law is of the same type as "the product graph does not remember the partner state ⇒ the binding energy can only be carried by the ledger" (project nucleon paper §12 [12]).

9.2 The graph level: the degenerate baseline $Q = 1/3$

The 9 ring edges of the closed-state skeleton lie in **a single automorphism orbit** (Table 2) ⇒ **any graph functional** gives the three rings **the same value** ⇒ $m_1 = m_2 = m_3$ ⇒

$$Q = \frac{3m}{(3\sqrt{m})^2} = \frac{1}{3} \qquad (\text{proven, within-model}).$$

This is exactly the **Cauchy–Schwarz lower bound** of the Koide combination (the equal-mass limit).

Note 7 (the double-identity trap of $1/3$). The $1/3$ here is the **C–S lower bound** (the equal-mass limit), which is **different in origin** from the SRE $\Pi_1(Q_3) = 1/3$ (**closure degree**) and **must not be identified with it**. The numerical coincidence is pure chance, and any reasoning that treats the two as mutually corroborating is void.

9.3 The ledger level: the splitting exists, but the rule is not written down

$$Q = \underbrace{\frac{1}{3}}_{\text{graph baseline (degenerate)}} + \underbrace{\frac{2}{3}\eta^2}_{\text{ledger splitting}}$$

Inverting Q for $\eta^2 = (3Q-1)/2$ gives a self-consistency check:

Table 13. Inverse solution of η^2 for three classes of objects (proven, within-model).

Object	Q	η^2	reference point

| charged leptons (pole masses) | 0.666667 | **0.500000** | $\eta^2 = 1/2 \Leftrightarrow Q = 2/3$ |

| heavy quarks (c, b, t) @own scale | 0.662748 | 0.494122 | deviation -0.6% |

| light quarks (u, d, s) @2 GeV | 0.567043 | 0.350564 | near $\eta^2 = 1/3 \Leftrightarrow Q = 5/9$ |

⇒ the splitting **does exist and has the right order of magnitude**, but the numerical origin of η^2 is still the single real gap (§10).

9.4 Quarks have one more wall than leptons

Lepton masses are **pole masses** (physical quantities, scheme-independent); the "mass" of a quark is the **MS-bar running mass** (**scheme-dependent + scale-dependent**) — so it **cannot even be settled whether it is the same quantity**. This is exactly the **C2 negative verdict** recorded long ago in the project.

Changing the convention for the same physical problem, Q can move from 0.34 to 0.89 (proven, within-model): PDG MS-bar gives $Q(u, d, s) = 0.5670$, $Q(c, b, t) = 0.6627$; unified to M_Z, $Q(c, b, t) = 0.7195$; taking naive **constituent quark masses** gives $Q(u, d, s) = 0.3378$. This is not a matter of error bars but of **the quantity itself having no unique definition**.

Analogy 8 (measuring the same mountain from different datum surfaces). The situation above is like measuring the same mountain from three datums — sea level, some valley floor, a neighbouring peak — and obtaining three heights that are all "correct" yet mutually unequal. When a quantity **has no unique definition**, any "agreement" that uses it as a target constitutes no evidence — this is the substance of tollgate C2.

9.5 A positive result: only RG-invariant dimensionless combinations qualify as targets

Theorem 5 (invariance of the scale). The Koide combination is invariant under an overall scale:

$$Q(c\,m_0,\ c\,m_1,\ c\,m_2) = Q(m_0, m_1, m_2) \qquad \forall c > 0.$$

Proof. The numerator $\sum c\,m_k = c \sum m_k \propto c$; the denominator $\Big(\sum \sqrt{c\,m_k}\Big)^2 = \Big(\sqrt{c}\sum \sqrt{m_k}\Big)^2 = c\Big(\sum \sqrt{m_k}\Big)^2 \propto c$. The two cancel, and the ratio is independent of c. □

Numerical verification ($c = 10^{-3} \ldots 10^{6}$): residual $\leq 1.1 \times 10^{-16}$ (proven, within-model).

Corollary 2 (screening rule for RG-invariant targets). The mass anomalous dimension of QCD γ_m is **flavour-independent**, so the running of the light quarks u, d, s is **multiplication by a common factor ⇒ Q_{light} is a renormalization-group invariant**.

Corollary 2 explains a past failure: in the probe $(2m_u + m_d)/m_p$ the denominator is the **external hadronic scale** m_p, so going from 2 GeV to 1 GeV changes the value by $+23\%$, and back-solving the number of strands V jumps from 12 to 16 (negative verdict). But Q_{light} is a **pure quark-mass ratio**, RG-invariant. **The two are not quantities of the same kind**: the earlier negative verdict was correct, but it **does not implicate Q_{light}**.

> **Screening rule.** Only **renormalization-group-invariant dimensionless combinations** qualify as candidate targets. Absolute values, same-scale ratios, and combinations containing a hadronic scale (m_p, Λ_{QCD}) are all excluded.

9.6 A caveat that must be retained: "light quarks ≈ 5/9" is only an interval-level agreement

Taking the PDG asymmetric errors and enumerating the 8 corners: the centre of Q_{light} is 0.567043, the interval $[0.545962,\ 0.586139]$, and $5/9 = 0.555556$ **lies inside the interval**, but the central value deviates by $+2.07\%$ (proven, within-model). This differs from the lepton one (deviation 3.3×10^{-6}) by **three orders of magnitude**. Hence "$\approx 5/9$" can only serve as a **loose target**, not a precision hit.

Literature side: Rodejohann and Zhang once put "heavy quarks $Q \approx 2/3$" into a preprint and **deleted it themselves at publication** [8]; and for "heavy quarks" the deviation degrades from 0.6% to 7.9% after the scale is moved — a full order of magnitude.

---

10. The single real gap: the assignment rule

10.1 Balance sheet of this paper

Table 14. State of attainment at each level.

Level	Content	Status
object	mass = the circulant operator on the three-ring torsor $c_0\mathbb{1} + c_1 S + \bar{c}_1 S^2$	**named, form unique (Theorem 2)**
number of positions	exactly 3 (the torsor has size 3)	**proven (structural)**
gauge	δ is a $\mathbb{Z}_3$ gauge (no origin)	**proven (structural)**
functional form	$Q = \frac{1}{3} + \frac{2}{3}\eta^2$, independent of δ	**proven (within-model)**
the value of η	why $\eta^2 = \frac{1}{2}$	**not proven — the single real gap**
scale	c_0 (absolute mass / energy scale)	**outside SRE (must be paid in externally)**

10.2 Precise statement of the gap

The gap is an **assignment rule** — a rule connecting "the **representation** of $\mathbb{Z}_3$ (edge $\frac{1}{3}$ | $\frac{2}{3}$, vertex $\frac{1}{2}$ | $\frac{1}{2}$, Table 3)" to "m_k / the order parameter A". That is to say:

> SRE already gives **everything on the discrete side** (why there are 3 positions, why δ is gauge, the functional form), but does not give the step "**representation weight → real measured value**". This step is what G12 calls the **continuous-side payment**.

10.3 Three hard criteria

Any work claiming to have written down the assignment rule must **simultaneously** satisfy:

1. **it must produce exactly 3 generations** (not 2 or 4);
2. **it must give $\eta^2 = \frac{1}{2}$ and explain its origin — $\frac{1}{2}$ must not** be taken as an input;
3. **it must give zero-parameter predictions for the three quark generations and the three neutrino generations** (light quarks $\approx 5/9$, heavy quarks $\approx 2/3$ already have measurements (Table 13) ⇒ it can be **immediately falsified out of sample**).

> The statement of criterion 2 is one substantive narrowing of this paper: since Theorem 3 proves that "the cosine shape holds trivially for any three generations", **in future it is no longer necessary to explain "why the shape is a cosine"; it is only necessary to explain "why $\eta = 1/\sqrt{2}$" — this one number**.

---

11. Discussion

11.1 Relation to external work

The $\mathbb{Z}_3$ structure identified in this paper is not isolated in the literature.

Table 15. The form in which $\mathbb{Z}_3$ / three generations appears in each work.

Work	In what form $\mathbb{Z}_3$ / three generations appears
Koide 1981 [1]	$\sqrt{m_k} \propto 1 + \sqrt{2}\cos(\delta + \frac{2\pi k}{3})$ — that $\frac{2\pi k}{3}$ **is $\mathbb{Z}_3$**
McRae 2025 [2]	"triality may be viewed as multiplication of a basis by a third root of unity, just as duality is often multiplication by a second root of unity" — **duality = $\mathbb{Z}_2$, triality = $\mathbb{Z}_3$**
Thorwe 2026 (DVFT) [3]	the natural $\mathbb{Z}_3$ of a three-component vacuum field ⇒ phase 120° ⇒ **circulant mass matrix** ⇒ Koide
Rousselle 2026 [4]	$U(3)_F \to U(1)_F^3$, equivalent to "crystallising" the vacuum to $Z(SU(3)) = \mathbb{Z}_3$
Ma 2006 [5]	four $\mathbb{Z}_3$ generators → the group $\Sigma(81)$
Sumino 2009 [6]	$U(3) \times O(3)$ family gauge symmetry; a 45° geometric reading
Libanov–Troitsky 2000 [7]	three generations from a **topological defect with topological number 3** (index theorem: number of zero modes = topological number)

This paper differs from them in **where it lands**: the works above generally take $\mathbb{Z}_3$ / circulant matrices as a **starting point** from which to construct mass matrices, whereas the $\mathbb{Z}_3$ of this paper is **read off from the automorphism group of the SRE skeleton** (three rings = $\mathbb{Z}_3$-torsor). That is to say, this paper provides a **structural source for "why a circulant mass matrix should appear"**, rather than yet another construction of a circulant mass matrix.

11.2 Candidate chain registered (D1 caution, not upgraded)

$Q = \frac{1}{3} + \frac{2}{3}\eta^2$ (Theorem 4), $(\frac{1}{3}, \frac{2}{3}) \leftarrow$ the $\mathbb{Z}_3$ weights of the edge representation, $\eta^2 = \frac{1}{2}$ $\leftarrow$ the trivial weight of the vertex representation (Table 3).

If the three are **of the same origin**, then $Q = \frac{1}{3} + \frac{2}{3} \cdot \frac{1}{2} = \frac{2}{3}$ holds **automatically**.

But this paper explicitly does not accept the chain, for the reason: $1/3, 2/3, 1/2$ are **all small-denominator rationals** (the D1 trap) and **have no discriminating power** on their own. The **only** reason for registering them as a candidate is that the three pieces happen to join into the complete chain $Q = 2/3$; **to upgrade it to a conclusion, a mechanism must be found elsewhere**, connecting "the edge representation / vertex representation" to "m_k / the order parameter A" (i.e. the assignment rule of §10). Otherwise it is of the same type as "integer ≈ integer" and is not accepted.

The role of **Table 3** must therefore be precisely delimited: it is **the registration of one link of a candidate chain**, not one of the conclusions of this paper.

11.3 Contrast with the "anchor" system

In the project's earlier anchor work, the only quantity satisfying a "constant ↔ single invariant" type of relation is the **fine-structure constant**:

$$\alpha = \Pi_1(M_{60}) = 7.2974585 \times 10^{-3} \qquad (\text{relative deviation } 1.45 \times 10^{-5} \text{ from the CODATA value } 7.2973526 \times 10^{-3}),$$

solving $\Pi_1(M_n) = \alpha$ gives $n^* = 60.000436$ (proven, within-model). Here M_{60} is the 60-vertex Möbius ladder, whose closed form is

$$\Pi_1(M_n) = \frac{4\sin^2(2\pi/n)}{2+2w+2\cos(2\pi/n)}\Big|_{w=1}.$$

Mass has no such anchor — and the reason is exactly the typology of Table 9: α is a **dimensionless ratio** (SRE can give it internally), whereas the **absolute scale** c_0 **of mass is immune to no structure** (outside SRE).

Analogy 9 (why a clock can be an anchor but "a pile of lengths" cannot). A pendulum clock can serve as a time standard because its pendulum length and period are **locked together into a ratio** — once locked, the external "second" is brought into physics through that ratio. This is the mechanism of an "anchor": **connecting an external constant to an internal structural invariant**. The mass discussed in this paper lacks exactly such a lock: c_0 is a **free scale**, constrained by no internal ratio, so it can only be **specified directly** from outside and cannot be **derived** internally. This also explains why the gap of §10 is "the absence of a rule" rather than "an inaccurate number".

(Registered together, one candidate **H4**: SRE has the exact ratio $\frac{2}{3} = T(\text{open})/T(\text{closed})$ (independent of V), numerically equal to the Koide $Q = 2/3$. It is likewise in the D1 domain, registered only, not cross-corroborated.)

11.4 The discrete-closure-law perspective

The location of this round is **case 15 of G12**:

- **discrete side (closed)**: cyclicity ⇒ the structure is a **circulant matrix**; the number of positions = 3 (torsor size); δ is gauge;
- **continuous side (requires input)**: the **value** of η (the origin of $1/\sqrt{2}$) and the scale c_0.

It is of the same type as the existing cases: **discrete structure fixes the shape, continuous quantities must be paid in externally**. One empirical aside: **everything that passes the tests is a dimensionless ratio; everything judged negative is a value for which dimensions are sought.**

---

12. Conclusion

This paper gives the following adjudication of the precise mathematical expression of "mass" in SRE:

1. **Object level**: mass is not three numbers but the **$\mathbb{Z}_3$-equivariant operator on the three-ring $\mathbb{Z}_3$-torsor = a circulant matrix** $A = c_0\mathbb{1} + c_1 S + \bar{c}_1 S^2$ (Corollary 1). It is **forced** by the skeleton structure (Theorem 1, Theorem 2) rather than an imposed language; the centraliser dimension is exactly $n = 3$.
2. **"Cyclic" level**: = using the **group** $\mathbb{Z}_3$ (rather than a semigroup) to characterise the index space (Proposition 3); 9 matrix entries are compressed to 3. For the three generations to be distinct, the operator **must carry a complex phase** (Proposition 5).
3. **"Unopened" level**: = not performing an origin-bearing spectral decomposition ⇒ the native output is an **equivariance class (multiset)**; the invariant moduli space is **two-dimensional** $(\eta,\ \delta \bmod \frac{2\pi}{3})$ (Proposition 8), and Q occupies only the η direction. **"Opening" = choosing the DFT origin, which is a $\mathbb{Z}_3$ gauge (redundancy) and changes no observable** (Proposition 6).
4. **Content level**: **"three points lie on one cosine" is free** (Theorem 3, holds for any three generations); **the entire content of Koide is that number** $\eta = 1/\sqrt{2}$ (measured η^2 deviates by 3.3×10^{-6}, Table 10).
5. **Boundary level**: **absolute mass (the scale c_0) is not inside SRE**; **only RG-invariant dimensionless combinations qualify as candidate targets** (Corollary 2).
6. **Gap level**: the **assignment rule** — connecting the representation weights of $\mathbb{Z}_3$ to m_k / the order parameter A. The three hard criteria are in §10.3.

Overall statement of the conclusion. Mass = an invariant of "the cyclic structure on $\mathbb{Z}_3$"; its carrier is a circulant matrix; "unopened" = outputting only the equivariance class; "opened" = choosing an origin (gauge, not information). What SRE can give: the **position** of (η, δ); what it cannot give: the **value** of η and the scale c_0.

Appendix A Recomputation list

One-command recomputation of all readings (requires numpy + networkx):

```
C:/myapp/miniconda3/envs/ai/python.exe -u _sre_mass_origin_paper_check.py \

    > _sre_mass_origin_paper_check.log 2>&1      # main body (PART 1-8)

C:/myapp/miniconda3/envs/ai/python.exe -u _sre_paper_v2_check.py \

    > _sre_paper_v2_check.log 2>&1               # v2.0 additions (Examples 1-3, alpha anchor, recheck of
Tables 2 and 3)
```

The two scripts are mutually independent implementations and share the project's existing closed form `mobius_pi1` (see `code/_sre_anchor_registry.py`).

Table A1. Correspondence between the readings of this paper and the recomputation scripts.

Location in this paper	Source	Key output
Theorem 2 (§4.1)	PART 1	rank $6/9$, null-space basis all circulant = true
Proposition 4 (§5.1)	PART 2	6 random sets, maximum error 1.1×10^{-15}
Proposition 5 (§5.2)	PART 2	$(0.7, 0.7, 1.6)$ / $(0.400668, 1.27515, 1.324181)$
Example 2 (§4.2)	v2 Ex.2	$(0.436184,\ 1.104189,\ 1.459627)$
Theorem 3 (§5.3)	PART 3 / v2 Ex.1	8 sets of arbitrary triples, reconstruction error 5.11×10^{-15}
Example 1 (§2.3)	v2 Ex.1	$\bar{u} = 0.966667$, $\lvert c_1 \rvert = 0.145297$, $\theta = 1.455835$ rad
Proposition 8 (§7.1)	PART 4	Q range $\leq 1.6 \times 10^{-15}$; R varies with δ
Proposition 6 (§6.1)	PART 4	$\lvert \Delta Q \rvert \leq 2.2 \times 10^{-16}$
Theorem 4, Table 10 (§8)	PART 5	$Q = 0.666664463$, $\eta^2 = 0.499996695$, 44.9999°
Theorem 1, Table 2 (§3)	PART 6	closed $36/8/1$; open $4/0/2$; φ edge-preserving, order 3
Proposition 2, Table 3 (§3.4)	PART 7	vertex $1/2$; edge $1/3$ | $2/3$
Theorem 5 (§9.5)	PART 8	$Q(c\,m)$ residual $\leq 1.1 \times 10^{-16}$
Example 3, Table 10 (§8.3)	v2 Ex.3	$\sum m = 1883.099374$, $\sum \sqrt{m} = 53.147515$, $Q = 0.666664463$
§11.3 alpha anchor	v2 anchor	$\Pi_1(M_{60}) = 7.2974585 \times 10^{-3}$; relative deviation 1.45×10^{-5}; $n^* = 60.000436$

Table A2. Preceding scripts (independent sources of the conclusions of this paper, all archived in `code/`).

Script	Round	Content
`_sre_weight_homology.py`	Round 29	weighted homology / dormancy rate (the $\mathbb{Z}_2$ level)
`_sre_triality_families.py`	Round 30	$\mathbb{Z}_3$ / triality / torsor / Koide
`_sre_quark_mass_probe.py`	Round 31	three-step decomposition of quark mass / RG-invariant targets

| `_sre_cyclic_mass_formalism.py` | Round 32 | formalisation of the circulant operator |

Appendix B Honesty boundary

1. Everything in this paper is a self-consistent statement **internal to the SRE model / at the level of linear algebra and group theory**, and **no** numerical prediction for real physics is claimed. The comparison with charged-lepton masses in the text is **structural consistency**, not prediction.
2. "Taking $a = \sqrt{m}$ rather than m" is the **empirical content of Koide**, equivalent to "**the square root of the mass operator** is $\mathbb{Z}_3$-equivariant" — a **testable structural hypothesis** that this paper **has not proved**.
3. "$\mathbb{Z}_3$ is the structure of the three generations" is a **mathematical identification + literature consensus**; what this paper **adds** from the SRE perspective is only two **structural facts**: "the skeleton carries $\mathbb{Z}_3$ natively (three rings = $\mathbb{Z}_3$-torsor)" and "closed → open = $\mathbb{Z}_3 \to \mathbb{Z}_2$". Neither is a new numerical prediction.
4. The echo between "real-symmetric ⇒ 2+1" of §5.2 and "the open state has only two values" is only an **observation**, which is **not proved** to be the same mechanism.
5. **The formalisation of this paper adds no constraint** (Theorem 3): it holds trivially for **any** three generations; its value lies in **language** and **layering**, not in new equations. "The shape is trivial" is a known fact in the literature; this paper makes it explicit within the SRE framework and accordingly corrects the earlier wording.
6. The chain of §11.2, "edge $1/3 \mid 2/3$ + vertex $1/2 \Rightarrow Q = 2/3$", **is a candidate, not a conclusion**: all three ratios are small-denominator rationals (D1) with no discriminating power on their own; an upgrade requires a mechanism found elsewhere.
7. The phase $\delta \approx 2/9$ of §8.4 is a small-denominator rational (D1), used only as supporting evidence; and δ is a **gauge quantity**.
8. Theorem 5 is a **mathematical identity** (for a common multiplicative factor); its applicability to QCD running **relies on** the standard result that γ_m is flavour-independent (true at LO and beyond; residual dependence only in threshold matching).
9. The scale c_0 (absolute mass) still requires external input — consistent with the existing G12 conclusion.
10. Conclusion = **case 15 of G12**.

Appendix C Analogy index

A survey of the analogies and illustrations used in this paper (each is developed in full at the corresponding place above).

Table C1. Index of analogies and illustrations.

No.	Name	Location	What it addresses
Analogy 1	three-phase alternating current	§2.3	the $\mathbb{Z}_3$ operator $a = e^{2\pi i/3}$, phase sequence as gauge, symmetrical components as DFT
Analogy 2	a three-strand braid and a clock face	§3.3	torsor: a cyclic order with no identifiable first element
Analogy 3	a wheel versus an hourglass	§4.3	group evolution (reversible, no start) vs semigroup evolution (irreversible, with a start)
Analogy 4	a prism and dispersion	§5.3	DFT = dispersion; "any three colours correspond to some beam" = the shape is free

| Analogy 5 | transposition of a score | §6.2 | choosing the DFT origin = choosing the tonic (structure unchanged) |

| Analogy 6 | an altitude datum and relative height | §7.3 | relative quantity (datum-independent) vs absolute quantity (needs an external datum) |

| Analogy 7 | forty-five degrees is the "midpoint deviation" | §8.2 | $\eta^2 = 1/2$ means stopping at the midpoint of equal mass and maximal deviation |

| Analogy 8 | three datum surfaces for one mountain | §9.4 | a quantity with no unique definition ⇒ agreement is no evidence |

| Analogy 9 | a pendulum clock as anchor vs a pile of lengths | §11.3 | anchor = locking an external constant to an internal invariant; c_0 lacks such a lock |

| Example 1 | cosine reparameterisation of three numbers | §2.3 | the inverse solution always succeeds; the order depends on the origin, the multiset does not |

| Example 2 | a concrete mass operator | §4.2 | a 3×3 matrix determined by only two real numbers |

| Example 3 | a complete recomputation for charged leptons | §8.3 | the full hand computation of $Q = 0.666664463$ |

Usage note. To explain this paper to a non-specialist, the recommended route is: Analogy 1 (three-phase) → Analogy 2 (braid) → Analogy 4 (prism) → Analogy 6 (altitude) → Example 3 (a concrete number). These five steps convey the skeleton of the whole paper without using any further terminology.

Appendix D References

[1] Y. Koide, *A Fermion-Boson Composite Model of Quarks and Leptons*, Phys. Lett. B **105**, 121 (1981); and the subsequent series of works on the "Koide relation" $Q = 2/3$.

[2] R. McRae, *Triality as multiplication by a third root of unity* (a root-of-unity characterisation of triality and duality), arXiv:2502.14016 (2025).

[3] Thorwe, *Discrete Vacuum Field Theory (DVFT)*: the $\mathbb{Z}_3$ phase of a three-component vacuum field ⇒ a circulant mass matrix (2026).

[4] Rousselle, *Family symmetry breaking and the center of SU(3)*, arXiv:2608.19277 (2026).

[5] E. Ma, $\mathbb{Z}_3$ *generators and the group* $\Sigma(81)$, hep-ph/0612022 (2006).

[6] Y. Sumino, *Family gauge symmetry and the 45° lepton mixing*, arXiv:0903.3640 (2009).

[7] M. Libanov, S. Troitsky, *Three generations from topological defects*, hep-ph/0011095 (2000).

[8] W. Rodejohann, H. Zhang, a preprint containing "heavy quarks $Q \approx 2/3$" (deleted in the published version); and X.-G. He, A. Zee et al. on the scale sensitivity of the Koide relation, arXiv:1111.0480.

[9] Particle Data Group, *Review of Particle Physics* (tables of charged-lepton and quark masses).

[10] This project, *SRE Triality / Three-Level Structure Report* (Round 30, 2026-09-28): first measurement of the $\mathbb{Z}_3$-torsor, the order-3 automorphism φ, the character decomposition, and the Koide recomputation.

[11] This project, *How to Compute Quark Mass in SRE: A Three-Step Decomposition Report* (Round 31, 2026-09-28): the three-step division graph/ledger/dimensions, the RG-invariant target screening rule.

[12] This project, *Complete Characterization of Nucleons within the SRE Framework* (nucleon complete paper): construction and screening of the skeleton $Y_3 \ltimes \triangle_3$ (§7–§8), the ledger-carrying law for binding energy (§12).

[13] This project, *A Formal Expression of SRE Mass: Circulant Operator Report* (Round 32, 2026-09-28): the circulant-matrix theorem, triviality of the shape, the invariant moduli space (the formalisation body of §5–§8).

[14] This project, *SRE Projection vs Distance Fidelity Report* (Round 28) and *SRE Weighted Homology / Dormancy Rate Report* (Round 29): the three-level stratification of distance and the $\mathbb{Z}_2$-level weighted homology (the prior basis for this paper's judgement that "a two-valued scheme cannot give three levels").

> **Citation discipline.** When citing references, the "≈" in the original must be copied verbatim and must not be rewritten as "equals".

Positioning of Light and Electromagnetic Waves in the SRE Framework

The Involutive $\mathbb{Z}_2$ Sector, the Cohomological Ladder, and "One Emission as One Closed→Open Transition"

Version: 1.0 (Treatise Format)

Date: 2026-09-28

Abstract

Background and problem. In prior work, SRE has identified "mass" as a $\mathbb{Z}_3$-equivariant circulant operator, and connected it with the **$\mathbb{Z}_3$-torsor** structure of the three-ring skeleton. A natural follow-up question then arises: **what is the SRE description of "light and electromagnetic waves", what is its relation to mass, and can it cover at once the radiation of chemical origin and of nuclear origin?** This paper answers these three questions head-on.

Method. This paper places the project's existing light-side stipulations (the residuality axiom, the Möbius double-cover parameterisation, the involution $P = S^{n/2}$ on the weighted Möbius ladder, the $\mathbb{Z}_2$ holonomy coupling δ_A) and the $\mathbb{Z}_3$ structure on the mass side (the three-ring torsor, the circulant operator) under one framework for comparison: the **cohomological ladder** $|H^1(G;\mathbb{Z}_n)| = n^{\beta_1}$. The whole paper needs only one structural input — **the involution $P = S^{n/2}$ on the weighted Möbius ladder M_n** — and everything else is recomputable linear algebra and group theory.

Results. The paper gives four mutually independent, verifiable propositions: (1) **the gear proposition** — light and mass are **two gears of the same machine**: light runs on the coefficient group $\mathbb{Z}_2$ (duality), mass on $\mathbb{Z}_3$ (triality), and the two are adjacent rungs of the same cohomological ladder; the root of the difference is a **group-theoretic prohibition** $|\mathbb{Z}_2| = 2 < 3$ (the light gear structurally cannot hold three generations). (2) **the carrier proposition** — the light carrier is the **odd eigenspace of an involution**, with $P^2 = I$, membership in the automorphism group, commutation with the Laplacian, zero trace, and spectrum exactly $\{\pm 1\}$; it is **completely determined by the single integer n, with zero continuous free parameters**, so **there is no place to put a source**. (3) **the source proposition** — the spectrum of the weighted Möbius ladder splits into a **"sector-blind term + odd-sector constant term"** $\mu_k = 2c(1-\cos\frac{2\pi k}{n}) + 2w \cdot \mathbb{1}[k \text{ odd}]$, in which only the w channel simultaneously satisfies "**acts only on the odd sector**" and "**shifts every mode uniformly**"; measurement over $\delta \in [10^{-3}, 5\times 10^{-1}]$ shows that the $n/2$ modes of the odd sector are shifted by **strictly the same amount** (range $\sim 10^{-15}$), that the even sector is **always motionless**, and that the spectral shape is **preserved mode by mode** (residual 8.9×10^{-16}); hence **the source can inject only one scalar on the light side**. (4) **the transition proposition** — "one emission = one **closed** → **open** transition": the **ledger difference** of the transition carries the energy (E, β_1 each -1), the **spectral invariants** of the transition carry the light carrier (ρ, the sector structure P), and the two are **orthogonal**; therefore a chemical source (transition at the electronic-molecular level) and a nuclear source (transition at the nuclear level) **share the same carrier**, their only difference being the level at which the ledger is recorded.

Boundary and conclusion. The paper also states four limitations: the light-side readings **come from the project's existing axioms, and this paper only puts them into recomputable form without adding any light-side physical content**; equating the "open state" with "light" is a **structural conjecture**, and the paper claims only a **consistency of direction**; $\mathbb{Z}_3$ is the **smallest** coefficient group able to carry three gears but **not the unique** one ($\mathbb{Z}_4$ works as well); the numerical equality $\operatorname{rank}(P_E) = 7$ vs. $\beta_1 = 7$ **is a pure coincidence and must not be used as mutual evidence**. The overall conclusion is registered as **case 18 of the**

discrete-closure law (G12): it gives "why the gears are two and three", "why the source is a single scalar" and "why energy and carrier are orthogonal" (discrete side, closed), but **not** the energy scale of light (continuous side, requiring external input). All numerical verification in this paper **holds only within the SRE model**.

Keywords: State–Relational–Entropy (SRE); light; electromagnetic waves; Möbius double cover; involution; $\mathbb{Z}_2$ holonomy; duality; triality; cohomological ladder; gauge degree of freedom; discrete-closure law.

---

1. Introduction

1.1 Statement of the problem

This paper answers the following proposition (raised 2026-09-28):

> **Original proposition.** Through the chain "triality $\mathbb{Z}_3 \to$ mass = circulant operator $\to$ light and mass = the same mechanism with a different coefficient group", can one and the same SRE description cover at once **light/electromagnetic waves produced by chemical energy** and **light/electromagnetic waves produced by nuclei**? And — is the SRE description of light itself **exactly consistent** with "mass = an unopened circulant operator"?

The proposition overlays three independently decidable sub-problems:

Table 1. Three sub-problems and their treatment in this paper.

No.	Sub-problem	Corresponding section
P1	What is the **carrier** of light in SRE?	§3 (Definition 1, Propositions 1, 2)
P2	**What is the relation** between light and mass?	§4 (Theorem 2, Propositions 3, 4)
P3	Can a chemical light source and a nuclear light source **share the same carrier**?	§7, §8 (Theorem 4, Proposition 8)

1.2 Existing characterisations and their predicament

In the prior work of this project, four conclusions relating to "light" already exist, but their **attributions are scattered** — they fall respectively on the axiom level, the spectral level, the coupling level and the numerical level:

Table 2. Existing light-side conclusions of the project (existing wording, not paraphrase).

Clue	Conclusion	Source

| **Residuality axiom** | Light is the **mutually non-annihilating topological residue** at the causal intersection of jointly executed chains: $\Psi_{\text{light}} = \ker(\partial_{\text{mutual}})$ | Electron paper §3.1 [1] |

| **Möbius double cover** | The residual manifold is parameterised by two intrinsic degrees of freedom (ϕ, w); when $\phi \to \phi + 2\pi$, $w \to -w$, and only $\Delta\phi = 4\pi$ closes | Electron topology §2 [2] |

| **Sector attribution** | On the weighted Möbius ladder the involution $P = S^{n/2}$ satisfies $P^2 = I$, $Pv_k = (-1)^k v_k$; k **even = electron sector,** k **odd = photon sector** | Fork A §2.4/§3.5 [3] |

| **Coupling strength** δ_A | $\delta_A = w^* - 1$ is the **emergent coupling strength** of the $\mathbb{Z}_2$ holonomy channel, with the character of a **metric parameter** | Electron topology §3 [2] |

These four clues point together to an object that is **not yet uniformly named**, and each exposes a predicament:

1. The **axiom level** gives light's **ontological identity** (residue), but not its **carrier operator**;
2. The **spectral level** gives the carrier operator $P = S^{n/2}$, but not **why it is the one**, nor what its relation to the mass operator is;
3. The **coupling level** gives that δ_A is "the continuous coupling of the $\mathbb{Z}_2$ gear", but it **cannot be measured independently** (an α^* input is required);
4. The **numerical level** gives the topological reproduction of $\alpha = \Pi_1(M_{60})$, but covers only **one** constant.

The task of this paper is to **gather these four clues into one framework** and to answer P1–P3. The key step is the methodological discipline of §2.4: **every crossing from "discrete" to "continuous" must be paid for with an external input**. Accordingly the light side splits into a **discrete gear** (given by $\mathbb{Z}_2$) and **continuous quantities** (coupling strength, energy scale, to be paid externally) — and the past predicament is precisely the result of mixing the two halves together.

1.3 Contributions of this paper

Relative to the existing material, this paper gives four substantive items:

1. **It establishes the "gear" as a structural proposition** (§4, Theorem 2): light and mass are not two mechanisms but adjacent rungs of one cohomological ladder $|H^1(G;\mathbb{Z}_n)| = n^{\beta_1}$; "changing gear = changing the coefficient group", and the hard reason is a group-theoretic prohibition $|\mathbb{Z}_2| = 2 < 3$.
2. **It makes the light carrier a recomputable reading** (§3, Proposition 2): the eleven properties of the involution P and the parameter count of zero continuous free parameters are all computed at once; on this basis it gives the structural ground for "**there is no place to put a source on the light side**".
3. **It proves that "the source injects only one scalar" as the unique channel screened out by two conditions** (§7, Theorem 4): starting from the two-term splitting of the spectrum, it proves that only the w channel satisfies both "acts only on the odd sector" and "shifts uniformly"; and it gives a **counterexample control** (the circumferential channel c fails both).
4. **It establishes "energy lives in the ledger, the carrier lives in the spectrum" as the transition proposition** (§8, Theorem 5), and on this basis explains that a chemical light source and a nuclear light source **share the same carrier** (answering P3).

1.4 Structure of this paper

Section 2 gives the existing stipulations, notation, an **analogy guide** and the methodological discipline; Section 3 proves that the light carrier is the involutive $\mathbb{Z}_2$ gear (first pillar); Section 4 proves that light and mass form adjacent rungs of the cohomological ladder (second pillar); Section 5 reviews the three-ring $\mathbb{Z}_3$-torsor and the structural change "closed → open" (third pillar); Section 6 argues that "unopened" is one and the same kind of statement on both sides; Section 7 treats the **source** side and gives channel

uniqueness; Section 8 discusses masslessness and the unification of chemical and nuclear sources; Section 9 connects this paper item by item with the existing SRE light-side documents; Section 10 is discussion and the screening of homonyms; Section 11 is the conclusion. Appendix A is the recomputation list, Appendix B the honesty boundary, Appendix C the analogy index, Appendix D the references.

Notation convention: numbers followed by "**(proven, within-model)**" can all be recomputed in one command by the script in Appendix A; those marked "**(to be proven)**" are structural hypotheses proposed here but not yet delivered; those marked "**(negative verdict)**" are propositions already falsified or explicitly delimited as non-computable.

---

2. Preliminaries

2.1 Existing SRE stipulations about light

(a) Residuality (Axiom I). The original wording of electron paper §3.1 [1] is:

> "Light is not an independent material substance. Given a boundary node A evolving to logical-depth step t, and a node B evolving to step t', light is embodied as the **mutually non-annihilating topological residue** at the causal intersection of the jointly executed chains:
>
> $\Psi_{\text{light}}(\phi, w) \equiv \ker(\partial_{\text{mutual}}(A_t, B_{t'}))$."

(b) The Möbius double cover (Axiom II). The same residual manifold is governed by **two intrinsic degrees of freedom**:

$$\mathbf{X}(\phi, w) = \left(\left(1 + w\cos\tfrac{\phi}{2}\right)\cos\phi, \quad \left(1 + w\cos\tfrac{\phi}{2}\right)\sin\phi, \quad w\sin\tfrac{\phi}{2}\right),$$

and when $\phi \to \phi + 2\pi$ the transverse vector **flips intrinsically**, $w \to -w$; one must traverse $\Delta\phi = 4\pi$ to close (axiom A2 of §2.2, §2.3, and the "4π closure" theorem [2]).

(c) Involution and sectors (Fork A §2.4/§3.5). Discretising this 4π-periodic structure into n equally spaced nodes (n even) yields the **weighted Möbius ladder** M_n: circumferential edges $(i, i \pm 1)$ with weight c (within-sheet refresh), and cross-sheet identification edges $(i, i + n/2)$ with weight w (the Möbius twist channel). On it the **involution**

$$P = S^{n/2}, \qquad P^2 = I, \qquad Pv_k = (-1)^k v_k.$$

Its eigenspaces give the **sector split**: k even = electron sector (soft mode), k odd = **photon sector** [3].

(d) Coupling strength δ_A (electron topology §3). $\delta_A \equiv w^* - 1$ is "the emergent coupling strength of the $\mathbb{Z}_2$ holonomy channel", with the character of a **metric parameter** (persistent-homology test: "δ_A does not change the barcode"; Morse test: "δ_A has no Morse counterpart"). Its **parameter-independent spectral fingerprint** is [2][3]

$$\mu_k(1+\delta_A) - \mu_k(1) = \begin{cases} 2\delta_A & k \text{ odd (photon sector)} \\ 0 & k \text{ even (electron sector)} \end{cases}$$

(e) Discrete electromagnetic realisation. `code/Maxwell.py` is a **gauge-covariantly closed** engine on a discrete cell complex: incidence matrix D (12×8), ring-edge matrix C (5×12), projector $P_E = D(D^\mathsf{T}D)^{-1}D^\mathsf{T}$; $E \in \Omega^1$ lives on edges, $B \in \Omega^2$ on rings, with three discretisations: Faraday / Ampère / Gauss.

The entire work of this paper is to unify (a)–(e) under the discipline of §2.4 and to answer P1–P3.

2.2 Notation

Table 3. Table of symbols used throughout.

Symbol	Meaning
SRE	State–Relational–Entropy, the parent framework of this paper
M_n	weighted Möbius ladder (n even; $V = n$, $E = 3n/2$, $\beta_1 = n/2 + 1$)
$c,\ w$	circumferential edge weight (within-sheet refresh) and cross-sheet identification weight (Möbius twist channel)
$P = S^{n/2}$	the **involution** on M_n (light carrier)
k	Fourier mode index; k even = electron sector, k odd = photon sector
μ_k	Laplacian spectrum of the weighted Möbius ladder
$\mathbb{Z}_n$	coefficient group (gear): $n = 2$ light gear, $n = 3$ mass gear
$\lvert H^1(G;\mathbb{Z}_n)\rvert = n^{\beta_1}$	**cohomological ladder**: the number of n-fold covers
δ_A	**coupling strength** of the $\mathbb{Z}_2$ holonomy channel (metric parameter)
δ_B	the **phase** on the mass side ($\mathbb{Z}_3$ gauge degree of freedom) — **not synonymous** with δ_A
$Y_3 \ltimes \triangle_3$	nucleon skeleton (mass-side carrier; §5)
$\rho,\ \lambda_2$	largest / second-smallest Laplacian eigenvalue
Q	Koide combination $Q = \dfrac{\sum m}{(\sum \sqrt{m})^2} = \frac{1}{3} + \frac{2}{3}\eta^2$
G12/D1/D2/D3	discrete-closure law / small-denominator rationals listed separately / angles may not serve as anchors / main statistics only on irrational readings × clean targets

2.3 Analogy guide: the Möbius strip and the switch

Before entering the formal arguments, let us build three **everyday-testable reference systems**, in order to fix the intuition behind all later wording.

Analogy 1 (the Möbius strip). Take a paper strip, give one end a half twist and glue it to the other end. Walk once along the **centre line** of the surface back to the starting point — put differently, after "one lap" you are on the **antipodal sheet**; only a second lap truly returns you to the original sheet. This is the $\mathbb{Z}_2$ double cover: **closure requires two laps (4π), not one (2π)**. The carrier operator $P = S^{n/2}$ of this paper is exactly the discrete version of this "sheet-flipping" operation, and its eigenvalues ± 1 label

respectively "stay on this sheet" and "flip to the antipodal sheet".

Analogy 2 (three-phase and single-phase electricity). The phasor operator $a = e^{2\pi i/3}$ of three-phase AC satisfies $a^3 = 1$, and its three eigenvalues form $\mathbb{Z}_3$; the operator of a single-phase / two-valued system is -1, satisfying $(-1)^2 = 1$, and its two eigenvalues form $\mathbb{Z}_2$. **The central proposition of this paper (Theorem 2) is that light uses the latter and mass the former** — the two systems are not two machines but one machine in different gears.

Analogy 3 (a switch has only two states). A switch with only "on/off" can express **at most two** distinct states, however it is thrown. If three things are required to be **pairwise distinct**, the switch structurally cannot do it — not "no method has been found yet", but **the state space is too small**. The capacity criterion of this paper (Theorem 2) is exactly the group-theoretic version of this plain fact: $|\mathbb{Z}_2| = 2 < 3$.

2.4 Methodological discipline: the discrete-closure law (G12)

All results of this paper can be located by one methodological discipline already existing in the project:

> **The discrete-closure law (G12).** Every link of the form "discrete structure ⇒ discrete target" **closes** within SRE; every crossing from "discrete" to "continuous" **must be paid for with an external input**.

The role of this law is a **division of labour**: for any result one must be able to tell which half is given by SRE (the discrete side) and which half must be paid in externally (the continuous side). It is the **main axis** along which this paper answers P1–P3 — because the four light-side clues of Table 2 are scattered precisely for want of this distinction.

Remark 1 (previous instances of G12). The law has accumulated seventeen instances in earlier work of the project (projection vs. distance, weighted homology, triality, RG invariance of quark targets, the "shape for free" of the circulant operator, light ↔ mass as a change of coefficient group, and others). The conclusion of this paper is registered as the **18th instance** (§10.4).

---

3. First pillar: light = the involutive $\mathbb{Z}_2$ gear

3.1 The carrier operator

Definition 1 (weighted Möbius ladder). Let n be even. The vertex set of M_n is $\{0, 1, \dots, n-1\}$, and the edges fall into two classes:

- **circumferential edges**: $(i, i+1 \bmod n)$, weight c, n of them (within-sheet refresh);
- **identification edges**: $(i, i+n/2)$, weight w, $n/2$ of them (Möbius twist channel).

Hence $V = n$, $E = 3n/2$, $\beta_1 = E - V + 1 = n/2 + 1$.

Definition 2 (light carrier). Define the involution

$P:\ i \mapsto i + n/2 \ (\mathrm{mod}\ n).$

By [3], P is an automorphism of M_n, commutes with the cyclic shift, and **commutes with the graph Laplacian** L. The spectrum of L splits by the eigenvalue of P into **two sectors**: the branch of eigenvalue $+1$ is the **electron sector**, that of -1 the **photon sector**.

3.2 The two-term splitting of the spectrum

Proposition 1 (sector decomposition of the spectrum). The Laplacian spectrum of M_n (circumferential weight c, identification weight w) is

$$\mu_k = 2c\Big(1 - \cos\tfrac{2\pi k}{n}\Big) + 2w \cdot \mathbb{1}[k \text{ odd}], \qquad k = 0, 1, \ldots, n-1.$$

Proof. The Fourier modes $v_k[i] = e^{2\pi i k i/n}$ diagonalise simultaneously the circumferential part and P: the circumferential part gives $2c(1 - \cos\frac{2\pi k}{n})$ (independent of the $\pm$ orientation), and P gives $(-1)^k$. The degree is $2c + w$, so the Laplacian acts as

$$Lv_k = \big[(2c + w) - 2c\cos\tfrac{2\pi k}{n} - w(-1)^k\big] v_k = 2c\Big(1 - \cos\tfrac{2\pi k}{n}\Big) + w\big(1 - (-1)^k\big)\, v_k.$$

And $1 - (-1)^k = 2\mathbb{1}[k \text{ odd}]$. □

Remark 2. This is equivalent to the project's existing closed form $\mu_k = (2 + w) - 2\cos\frac{2\pi k}{n} - w(-1)^k$ ([3] §3.4) at $c = 1$; Proposition 1 merely separates c explicitly so that **channels** can be compared in §7.

Numerical check of Proposition 1 (Appendix A, PART 1a): for $n = 6, 8, 10, 20, 30, 60$, the maximum deviation between the closed form and direct diagonalisation is $\leq 6.2 \times 10^{-15}$; $\mu_0 = 0$ holds identically; the two branches have $n/2$ modes each (for $n = 60$, $|\text{even}| = |\text{odd}| = 30$) (proven, within-model).

Proposition 1 immediately yields **two structural facts**, which will serve in §7 as the two criteria screening the "source channel":

Table 4. Range and shape of the two channels (proven, within-model).

Channel	Range	Per-mode displacement	Shape
c (circumferential weight)	**both sectors treated alike**	$\propto \left(1 - \cos\frac{2\pi k}{n}\right)$	**non-uniform**
w (identification weight / holonomy)	**odd sector only**	identically $= 2w$	**a pure constant shift, no shape**

3.3 Topological protection of the soft mode

Theorem 1 (topological protection of the soft mode, existing [3] §3.5). For all w,

$$\lambda_2(w) = 2 - 2\cos\tfrac{4\pi}{n} \qquad (\text{independent of } w), \qquad \lambda_{\max}(w) = 2 + 2w + 2\cos\tfrac{2\pi}{n} \ \ (w > 1 - \cos\tfrac{2\pi}{n}).$$

Proof. By Proposition 1, for k even $2w \cdot \mathbb{1}[k \text{ odd}] = 0$, so μ_k is independent of w; for k odd μ_k contains $+2w$. The second-smallest non-zero mode comes from $k = 2$ (even), so $\lambda_2 = 2c(1 - \cos\frac{4\pi}{n})$ is independent of w; the largest mode comes from an odd mode with k near $n/2$, giving $2 + 2w + 2\cos\frac{2\pi}{n}$. □

Physical reading. The electron modes are **purely topological** (completely insensitive to the strength of the twist channel); the Möbius twist channel only renormalises the **speed-of-light modes**. The numerator of α is topological while the denominator contains dynamics — this is the spectral ground for the project's reading of α as v/c [3].

Measurement (Appendix A, PART 1b–1c): for $w \in \{0.05, \ldots, 10\}$, λ_2 is identically 0.043704798532, spread 4.3×10^{-15} (proven, within-model).

3.4 The carrier has zero parameters: no place to put a source

Proposition 2 (parameter count of the light carrier). P is completely determined by **one integer** n, and contains **no continuous free parameter**; its eleven structural properties are as follows.

Table 5. Eleven tests of $P = S^{n/2}$ ($n = 60$) (proven, within-model).

Test	Result
$P^2 = I$	true
$P \neq I$	true
$P \in \mathrm{Aut}$ ($PAP^{\mathsf{T}} = A$)	true
$[P, L] = 0$ (simultaneously diagonalisable)	true
$\mathrm{spec}(P) = \{+1, -1\}$	true
$\mathrm{tr}(P) = 0$ (the two branches have equal weight)	true
$P_+,\ P_-$ idempotent	true
$P_+ P_- = 0$ (orthogonal)	true
$P_+ + P_- = I$	true
$\mathrm{rank}(P_+) = \mathrm{rank}(P_-) = n/2$	true

Proof. Each item is verified directly from the definition of P together with $PAP^{\mathsf{T}} = A$ and $PL = LP$; idempotence and orthogonality follow from the spectral projections $P_\pm = \frac{1}{2}(I \pm P)$ built from $P^2 = I$. □

Table 6. Comparison of carrier parameter counts.

| Carrier | Input determining it | Continuous free parameters |

|---|---|---|

| light: symmetry carrier P | **one integer** n | **0** |

| light: spectral scales (c, w) | — | 2, but only w is "light-exclusive" |

| mass: operator A | $\mathrm{Re}\, c_0,\ \mathrm{Re}\, c_1,\ \mathrm{Im}\, c_1$ | 3 (invariants 2: c_0 and $\lvert c_1\rvert$) |

> **Table 6 is the hardest item in the answer this section seeks: in the symmetry carrier there is no place to put a source.** Compare the mass side — the degrees of freedom $(c_0, \lvert c_1\rvert)$ of the operator A are the landing point of the "ledger" (§8). This is the **common source** of "the source is invisible" in §7 and "energy lives in the ledger" in §8.

Analogy 4 (a volume knob). An amplifier with only one knob, "volume", leaves the **shape of the frequency response** unchanged however it is turned; only the **overall loudness** changes. The light source can act only through the single knob w — what it changes is the amplitude, not the shape. Section 7 turns this intuition into a criterion.

---

4. Second pillar: the cohomological ladder — light and mass are adjacent rungs

4.1 The cohomological ladder

Proposition 3 (counting n-fold covers). Let G be a finite connected graph with first Betti number β_1. Then the number of $\mathbb{Z}_n$ principal covers of G is

$$\left|H^1(G;\mathbb{Z}_n)\right| = n^{\beta_1}.$$

Proof. $H^1(G;\mathbb{Z}_n) \cong \mathrm{Hom}\big(\pi_1(G), \mathbb{Z}_n\big)$, and the free part of $\pi_1(G)$ has rank β_1, hence $|\mathrm{Hom}(\mathbb{Z}^{\beta_1}, \mathbb{Z}_n)| = n^{\beta_1}$. □

Table 7. The cohomological ladder (Appendix A, PART 4a; β_1 measured).

| Graph | V | E | β_1 | $\lvert H^1(\mathbb{Z}_2)\rvert = 2^{\beta_1}$ | $\lvert H^1(\mathbb{Z}_3)\rvert = 3^{\beta_1}$ |

|---|---|---|---|---|---|

| C_3 (triangle) | 3 | 3 | 1 | 2 | 3 |

| K_4 | 4 | 6 | 3 | 8 | 27 |

| Q_3 (cube) | 8 | 12 | 5 | 32 | 243 |

| M_6 | 6 | 9 | 4 | 16 | 81 |

| Y_3 closed (nucleon skeleton) | 12 | 18 | 7 | 128 | **2,187** |

| Y_3 open (one ring edge removed) | 12 | 17 | 6 | 64 | **729** |

| M_{60} (α anchor carrier) | 60 | 90 | 31 | 2,147,483,648 | 617,673,396,283,947 |

The reading of Table 7 is the central sentence of this paper: this ladder has only one free parameter — the n of the coefficient group. Light and mass are not two mechanisms but **adjacent rungs of one and the same ladder**: light takes the $n = 2$ column, mass the $n = 3$ column. The readings of the $n = 3$ column agree item by item with the project's triality report [4], with consistent conventions.

4.2 The capacity criterion: why the light gear cannot hold three generations

Theorem 2 (capacity criterion). Attach a $\mathbb{Z}_N$ label to each of the three edges of a triangle. Then:

- $\mathbb{Z}_2$: the reachable partition patterns are only $\{3\}$ and $\{2{+}1\}$, $(1,1,1)$ **is unreachable**, and the maximum number of pairwise distinct labels $= 2$;
- $\mathbb{Z}_3$: the reachable patterns include $(1,1,1)$, and the maximum number of pairwise distinct labels $= 3$;
- $\mathbb{Z}_4$: likewise includes $(1,1,1)$, and the maximum number of pairwise distinct labels $= 3$.

Proof. The labels of the three edges in $\mathbb{Z}_N$ are (a,b,c), and the partition pattern is the partition of their multiplicities. Under $\mathbb{Z}_2$, $a,b,c \in \{0,1\}$, so the three fall into at most 2 distinct values and the partition can only be (3) or $(2,1)$; under $\mathbb{Z}_3$, taking $(a,b,c) = (0,1,2)$ gives $(1,1,1)$. □

Measurement (Appendix A, PART 4b): the reachable patterns for $\mathbb{Z}_2 = \{(3),(2,1)\}$; both $\mathbb{Z}_3$ and $\mathbb{Z}_4$ include $(1,1,1)$ (proven, within-model).

Corollary 1. Since $|\mathbb{Z}_2| = 2 < 3$, **the light gear structurally cannot carry three generations**. This is the **hard reason** for "changing the coefficient group", and the **group-theoretic root** of the project's negative verdict in round 29 ("under two-valued labelling a triangle gives at most $2{+}1$, and $\{1,1,1\}$ is never reachable") [5].

Analogy 5 (a two-compartment drawer cannot hold three things). Theorem 2 says: a drawer with only two compartments cannot, however it is arranged, hold something that requires "three pairwise distinct items". This is not a question of "arrangement technique" but of **too few compartments** — irrespective of **what the third item is**. The restriction is purely combinatorial, and therefore has nothing to do with physical questions such as "whether light is a wave".

4.3 A limitation that must be stated clearly: $\mathbb{Z}_3$ is "smallest", not "unique"

The literature says $\mathbb{Z}_4$ also gives three pairwise distinct labels (third column of Theorem 2). Hence:

> **Remark 3.** "$|\mathbb{Z}_n| \geq 3$" is only a **necessary condition**; the status of $\mathbb{Z}_3$ is that of the **smallest** (the first gear that is just big enough), **not the unique** one. The **uniqueness** of $\mathbb{Z}_3$ requires a separate reason — what the project's triality report [4] gives is "the three rings form a single orbit under Aut, of size exactly 3 ⇒ there exists a group of order exactly 3 acting freely and transitively on them (a G-torsor)". But this uses "the number of juxtaposed positions $= 3$", and "why the number of positions is 3" remains undecided. **Circular argument is not allowed.**

This paper does **not** claim that $\mathbb{Z}_3$ is unique. It uses only half of it — the **necessary condition**: $\mathbb{Z}_2$ is **not enough**.

4.4 The same parent group

Proposition 4 (common parent group). $\mathbb{Z}_2 \subset U(1)$, $\mathbb{Z}_3 \subset U(1)$, and $U(1)$ is precisely the electromagnetic gauge group. Hence:

- **light = the order-2 subgroup of** $U(1)$ (phase taking $\{0, \pi\}$, i.e. the 2nd root of unity);
- **mass = the order-3 subgroup of** $U(1)$ (phase taking $\{0, \frac{2\pi}{3}, \frac{4\pi}{3}\}$, i.e. the 3rd root of unity).

Literature basis. The original wording of McRae 2025 [6] is "**triality may be seen as multiplication of the basis by a third root of unity, just as duality is often multiplication by a second root of unity**" — that is, **duality =** $\mathbb{Z}_2$**, triality =** $\mathbb{Z}_3$.

Remark 4. By G12 (§2.4): the **continuous phase** of $U(1)$ (coupling strength, continuous angles) belongs on both sides to the **continuous side** and requires external input. This paper treats only the **discrete gears** on both sides.

---

5. Third pillar: the three-ring $\mathbb{Z}_3$-torsor and "closed → open"

This section reviews the mass-side structure, to provide the basis for the comparisons in §6 and §8; the full proof is in the project's mass-origin paper [7].

5.1 The torsor structure (cited)

Theorem 3 (the $\mathbb{Z}_3$-torsor structure of the three rings, [4][7]). The three rings of the closed-state nucleon skeleton $Y_3 \ltimes \triangle_3$ ($V = 12$, $E = 18$, $\beta_1 = 7$, $|\mathrm{Aut}| = 36$) form a $\mathbb{Z}_3$-torsor: there is a **cyclic order**, but **no absolute origin**; the only degree of freedom is an **overall phase shift**.

Its explicit witness is the order-3 automorphism $\varphi: \ L_{ij} \mapsto L_{i,(j+1) \bmod 3}$. **Measurement** (Appendix A, PART 5b): edge-preserving = true, $\mathrm{ord}(\varphi) = 3$ = true, $\varphi \neq \mathrm{id}$ = true (proven, within-model).

5.2 The order-element spectrum: closed → open = $\mathbb{Z}_3 \to \mathbb{Z}_2$

Proposition 5 (the structural change from closed to open). The order-element spectra of the automorphism groups of the closed state and the open state (one ring edge removed) are as follows.

Table 8. Automorphisms and order-element spectra of the two states (Appendix A, PART 5a; proven, within-model).

State	$\lvert\mathrm{Aut}\rvert$	order 1	order 2	order 3	order 6	orbits on the three rings
closed (E=18, β_1=7)	**36**	1	**15**	**8**	12	**1**
open (one ring edge removed, E=17, β_1=6)	**4**	1	**3**	**0**	0	**2** $\{0\}$	$\{1, 2\}$

Reading.

> **Closed state (unopened): $\mathbb{Z}_3$ is alive (8 elements of order 3) ⇒ triality ⇒ mass / three generations.**

> **Open state (opened): $\mathbb{Z}_3$ vanishes, only $\mathbb{Z}_2$ remains ⇒ duality ⇒ light.**

And the project's residuality axiom (§2.1a) states plainly that "light is … a **mutually non-annihilating topological residue**" — **light falls precisely on the "opened / unclosed" side**. The two **agree in direction**.

Remark 5 (this is a structural conjecture, not a proven equivalence). Equating the "open state" with "light" is a **structural conjecture** of this paper; the paper claims only **consistency of direction** (see Appendix B, item 3).

5.3 The character decomposition of $\mathbb{Z}_3$

Table 9. The character decomposition of $\mathbb{Z}_3$ on the vertex space and the edge space (Appendix A, PART 5c; proven, within-model).

Space	$\dim$	$\chi(\varphi)$	Decomposition	Trivial weight	Non-trivial weight
vertices	12	3	$6\cdot 1+3\omega+3\omega^2$	$6/12=\mathbf{1/2}$	$6/12=\mathbf{1/2}$
edges	18	0	$6\cdot 1+6\omega+6\omega^2$	$6/18=\mathbf{1/3}$	$12/18=\mathbf{2/3}$

Remark 6 (registration discipline). The $1/2, 1/3, 2/3$ in Table 9 are all **small-denominator rationals** (the D1 trap) and are **without discriminating power** on their own. Their role on the mass side is registered in the candidate chain of [7] §11.2; this paper cites them only as a **comparison** and does not upgrade them.

---

6. "Unopened" is the same kind of statement on both sides

The wording put forward by the user in round 32 is "**an evolutionary logic that is unopened and cyclic**". This sentence lands precisely on the light, mass and electromagnetic sides alike, and **lands at the same point**.

6.1 The mass side

Proposition 6 (the "unopened" on the mass side, [7] Proposition 6). Writing the **multiset** of amplitudes as an **ordered triple** is equivalent to choosing an **origin** for the DFT characters; there are 3 choices of origin, and the three choices give **exactly the same** permutation invariants.

Measurement (Appendix A, PART 7a): with $\eta=1/\sqrt{2}$, $\delta_B=2/9$, the three choices of origin give $Q=0.666666666667$, $|\Delta Q|\le 2.2\times 10^{-16}$ (proven, within-model).

6.2 The light side

Proposition 7 (the "unopened" on the light side). By the double-cover parameterisation of M_n, ϕ and $\phi + 2\pi$ **are not the same point but the same point on the antipodal sheet** ($w \to -w$); only traversing 4π returns to the original sheet.

Measurement (Appendix A, PART 7b):

$$\left|\mathbf{X}(\phi{+}2\pi, w) - \mathbf{X}(\phi, -w)\right| = 0, \qquad \left|\mathbf{X}(\phi{+}4\pi, w) - \mathbf{X}(\phi, w)\right| = 1.1 \times 10^{-15}\ (\text{proven, within-model}).$$

6.3 The three sides are one and the same kind of statement

Table 10. The exact meaning of "unopened" on the three sides (proven, within-model).

Side	Exact meaning of "unopened"	What is identified away
mass	no origin-bearing DFT ⇒ the origin of $\mathbb{Z}_3$ is unobservable, giving only the equivariance class	the **origin label** of $\mathbb{Z}_3$
light	ϕ and $\phi + 2\pi$ are the same point on the antipodal sheet, giving only 4π closure	the **antipodal sheet** of $\mathbb{Z}_2$
electromagnetic wave	gauge invariance: $A \to A + d\lambda$ leaves observables unchanged (P_E idempotent)	the **gauge function** of $U(1)$

> **All three = "do not fix an origin in advance, output only gauge invariants".** The difference lies only in **what is identified away**. This explains why the wording "unopened" can cover all three at once — it is not a metaphor but **one and the same kind of gauge structure**.

Remark 7 (two senses of "opening" must not be conflated). "Opening" has two layers, which must be strictly distinguished:

Table 11. Two senses of "opening".

Layer	Operation	Consequence
(i) gauge layer	choose an origin = label	invariants **unchanged** (redundancy) — this is the layer meant in this section
(ii) physical layer	closed → open (cut one ring edge)	$\lvert\mathrm{Aut}\rvert$ drops from 36 to 4 and the order-3 elements from 8 to zero: a **real** breaking $\mathbb{Z}_3 \to \mathbb{Z}_2$ (§5.2, §7)

---

7. The source is invisible: one emission = one closed → open transition

7.1 The source injects only one scalar

This section answers the remainder of P1 and the first half of P3: **can the source of light be seen?**

Theorem 4 (channel uniqueness). Within the **family of linear perturbations preserving the P symmetry**, only the w channel (identification weight / holonomy) simultaneously satisfies:

- (i) it acts **only on the odd sector**;
- (ii) it applies **the same displacement** 2δ to every mode of the odd sector.

Proof. By Proposition 1, $\mu_k = 2c(1-\cos\frac{2\pi k}{n}) + 2w\mathbb{1}[k \text{ odd}]$. A perturbation δc of c gives $\Delta\mu_k = 2\delta c(1-\cos\frac{2\pi k}{n})$, which appears **in both sectors at once** (violating i) and is non-uniform in k (violating ii). A perturbation δw of w gives $\Delta\mu_k = 2\delta w$ $\mathbb{1}[k \text{ odd}]$: only in the odd sector (satisfying i), and $\Delta\mu_k/2\delta w \equiv 1$ (satisfying ii). The supports of the two on $\{v_k\}$ determine their action spaces, so within the linear family preserving P the channel is **unique**. □

Measurement (Appendix A, PART 3a): $w : 1 \to 1+\delta$, $\delta \in \{10^{-3}, 10^{-2}, 10^{-1}, 5\times10^{-1}\}$:

Table 12. Uniformity test of the w channel (proven, within-model).

δ	range of $\Delta\mu$ in the odd sector	range of $\Delta\mu/2\delta$	range of $\Delta\mu$ in the even sector
10^{-3}	1.8×10^{-15}	$1.000000000 \sim 1.000000000$	0
10^{-2}	8.9×10^{-16}	$1.000000000 \sim 1.000000000$	0
10^{-1}	1.8×10^{-15}	$1.000000000 \sim 1.000000000$	0
5×10^{-1}	8.9×10^{-16}	$1.000000000 \sim 1.000000000$	0

Corollary 2 (the spectral shape is unchanged). Under the w channel the **spectral shape of the photon sector is preserved mode by mode**, with only an overall shift by 2δ.

Measurement (Appendix A, PART 3b): after subtracting the shift and comparing with the baseline, the residual is 8.9×10^{-16} (proven, within-model).

> **On the light side the source is invisible; one can hear its sound (amplitude) but cannot see its shape.**

7.2 Counterexample control

Table 13. Counterexample: the circumferential channel $c = 1+u$ (Appendix A, PART 3c; proven, within-model).

u	range of displacement in the odd sector	range of displacement in the even sector	odd sector only?

| 10^{-2} | 3.98×10^{-2} | 4.00×10^{-2} | **no** (both sectors move) |

| 10^{-1} | 3.98×10^{-1} | 4.00×10^{-1} | **no** |

⇒ both criteria **fail**. So "the light-exclusive channel = w" is **screened out**, not chosen at will.

Back to Analogy 4. w is exactly that "volume knob": it **raises the whole frequency-response curve** (a shift) without touching its shape; whereas c is more like "adjusting the timbre" — it **changes the shape of the curve**, and hence is no longer "the same carrier at a different strength".

7.3 Energy lives in the ledger, the carrier lives in the spectrum

Theorem 5 (orthogonality of ledger and spectrum). On the nucleon skeleton Y_3, "closed → open" (cutting one ring edge) **changes only ledger quantities** (E, β_1, the number of rings) and **does not change spectral quantities** (ρ, λ_2 and the sector structure).

Proof (measurement + structural explanation). See **Table 14**: the ledger quantities E and β_1 each change by -1, while the changes in ρ and λ_2 are zero to machine precision. The structural reason is that ρ and λ_2 are **automorphism invariants** (points in the same orbit take the same value), whereas "which edge is cut" is **explicit bookkeeping** — by the project's dichotomy of "graph functional vs. ledger" [7] §9.1, invariants are **blind** to points in the same orbit, so the cut-edge information can only be carried by the ledger. □

Table 14. Signature vector of the Y_3 skeleton closed ↔ open (Appendix A, PART 6a; proven, within-model).

Quantity	closed	open	difference	Class
V	12	12	$+0$	**ledger**
E	18	17	-1	**ledger**
β_1	7	6	-1	**ledger**
ρ	5.302775637732	5.302775637732	-4.4×10^{-15}	**spectral (unmoved)**
λ_2	1.000000000000	1.000000000000	1.3×10^{-15}	**spectral (unmoved)**

Corollary 3 (division of labour between source and carrier). Energy (how much is released) lives in the **ledger**; the **light carrier** lives in the **spectral invariants**; the two are **orthogonal**. ⇒ **The source changes only the ledger, not the carrier.**

Table 15. Spectral quantities when each of the 18 edges is cut in turn (Appendix A, PART 6b; proven, within-model).

Edge class	count	range of ρ	value of λ_2 (range within the class)
ring edges (L–L)	9	2.7×10^{-15}	1.000000000 (2.3×10^{-15})
strand edges (c–L)	9	3.6×10^{-15}	0.527166091 (2.2×10^{-15})

Remark 8 (a refinement, not a refutation). The project's nucleon paper §4 records that "in the open state, cutting one ring edge ⇒ ρ, λ_2 unchanged". Table 15 **refines** this to: ρ is a **globally strict invariant** (18/18 edges, range $\sim 10^{-15}$); whereas λ_2 is an **orbit-level invariant** — it stays constant only **within one** Aut**-orbit** (the ring-edge orbit gives 1.0, the strand-edge orbit 0.527), and does **not** depend on which particular edge was cut. The original statement is **correct** in the context "cutting a ring edge"; generalised to "immune to whichever edge is cut" it **does not hold**.

Analogy 6 (the ledger and the scale). Picture a warehouse: the **stock ledger** records every movement of goods in and out (ledger quantities), while the **calibration of the weighing platform** (the spectrum) records the manner of weighing and does not change its calibration because one crate is missing. Table 14 of this paper separates the two — the source (one crate missing) moves the **ledger**, the carrier (the calibration) does not. **Chemical energy** and **nuclear energy** can be described by one and the same language precisely because their difference is recorded only in the ledger, while the way the calibration is read is the same.

---

8. Masslessness and the unification of chemical and nuclear light

8.1 The light carrier has no additive scale

Proposition 8 (the light carrier has no additive scale). The spectrum of the light carrier P is identically $\{+1, -1\}$ ($|\mathrm{eig}| \equiv 1$), the spectral mean is $\mathrm{tr}(P)/n = 0$, and $\mu_0 = 0$ holds identically; whereas the spectral mean of the mass carrier $A = c_0\mathbb{1} + c_1 S + \bar{c}_1 S^2$ **equals the free parameter** c_0.

Table 16. Spectral means of the two carriers (Appendix A, PART 8; proven, within-model).

Carrier	Spectrum	Spectral mean	Additive scale
light: P	identically $\{+1, -1\}$, $\|\mathrm{eig}\| \equiv 1$	$\mathbf{0}$	**none** ($\mu_0 = 0$ holds identically)
mass: A	$c_0 + 2\|c_1\|\cos(\theta + \frac{2\pi k}{3})$	$\equiv \mathbf{c_0}$	**present** (c_0 is a free parameter)

Measurement (Appendix A, PART 8b): for $c_0 \in \{0, 0.5, 1, 2\}$ the spectral mean of the mass carrier is exactly $0, 0.5, 1, 2$; the light carrier has mean identically 0 for every n (proven, within-model).

By G12 (§2.4): c_0 is that dimension which **must be input externally**. This is the answer to predicament (4) of §1.2, and is **three phrasings of the same conclusion** — with the project's existing corollary "a pure transition has no mass ⇒ the photon is unique" and with "the absolute scale $c_0 \in$ outside SRE" (**not three independent pieces of evidence**, see Remark 9).

Remark 9 (prohibition of double counting). The following three point to **one and the same place** and **must not be listed as mutually corroborating evidence**:

1. this section: the light carrier has no additive scale while the mass carrier does ⇒ c_0 is the dimension that "must be input externally";
2. the project's `_sre_boson_transition.py`: "ρ is a transition invariant ⇒ a pure transition has no mass ⇒ the photon is unique";

3. the project's mass-origin paper §10: the absolute scale $c_0 \in$ outside SRE (SRE gives only ratios) [7].

> **Mass needs a scale that SRE cannot supply; light does not.** This is the **formal location** of "the photon is massless" in SRE. It must be stressed that "the photon is massless" has a different source in the Standard Model (gauge invariance plus only two transverse degrees of freedom); the statement here carries only the one layer "**no $\mathbb{Z}_3$ ⇒ no such unopened cycle**", and the two **must not be conflated**.

8.2 Chemical light and nuclear light: the same carrier

Table 17. Comparison of the two sources on the SRE carrier (Appendix A, PART 2, 3, 6; proven, within-model).

Aspect	Chemical energy / atomic emission	Nuclear emission (γ)	Attribution on the SRE side
carrier	odd sector P	odd sector P	**the same** (P determined by n alone)
sector structure	$n/2$ even + $n/2$ odd	the same	**the same**
spectral shape	independent of the source	independent of the source	**the same** (Corollary 2)
coupling quantity	one scalar w	one scalar w	**of the same type**, differing only in magnitude
transition occurs at	L1/L2 (electronic–molecular level)	L3 (nuclear level)	**different**
order of magnitude of released energy	$\sim$1–10 eV	$\sim$0.1–10 MeV	**external ledger** (Remark 10)

⇒ Answer to P3: yes. Chemical light and nuclear light **share the same carrier** on the SRE side, and this is not a new assumption but a **direct corollary** of "there is no place to put a source on the light side" (§3.4) and "ledger and spectrum are orthogonal" (§7.3).

Analogy 7 (a battery and a generator). The same light bulb can be connected to a **battery** (chemical energy) or to a **generator** (mechanical energy) — the bulb (the carrier) is exactly the same, and what changes is the **power source**. If someone asks "is a bulb lit by a battery the same kind of bulb as one lit by a generator", the answer is obviously "the same bulb, a different source". Table 17 of this paper says exactly this: **the carrier is the same, the source differs**.

8.3 The difference lies in the ledger

Remark 10 (prohibition of appropriating a homonymous term). The project's existing "span of 13.40 orders of magnitude across three layers" (L1 -2.30/L2 -0.09/L3 $+11.10$) is the span of the **sensitivity of decay rates to the environment**, **not** a span of photon energies. The two are different physical quantities and **must not be cross-cited**.

Remark 11 (annotation of external magnitudes). The eV/MeV in Table 17 are **external reference magnitudes** (neither measured here nor an SRE output): the ratio is $\sim 10^{5\text{–}6}$. **This is the ledger magnitude of the source; there is no corresponding reading on the SRE side** — the same gap as c_0 in §8.1.

---

9. Connection with existing SRE light-side documents

This section connects this paper item by item with the four existing clues (Table 2), in order to confirm that this paper **adds no light-side physical content** and only performs **structuring and making-recomputable**.

9.1 The residuality axiom and the "open state"

The project's residuality axiom (§2.1a) says that light is a "mutually non-annihilating **topological residue**". The structural fact of §5.2 of this paper is: the closed state has $|\mathrm{Aut}| = 36$ and contains 8 elements of order 3; the open state has $|\mathrm{Aut}| = 4$ and **zero order-3 elements**.

> **"Residue" = the part of the structure that has not been closed away.** What is supported by $\mathbb{Z}_3$ in the closed state and **vanishes** in the open state is exactly the "triality" part; what remains is $\mathbb{Z}_2$. Hence "light is a residue" and "light is on the $\mathbb{Z}_2$ gear" **agree in direction**.

Remark 12. This is only **consistency of direction**, and does not constitute a proof that "open state = light" (see Appendix B, item 3).

9.2 Formal separation of the two δ's

The comparison of this paper touches **two** homonymous symbols, which must be pinned down item by item:

Table 18. Separation of the two δ's.

	δ_A (electron topology §3 [2])	δ_B (mass side [7])
definition	$\delta_A = w^* - 1$, the **coupling strength** of the $\mathbb{Z}_2$ holonomy channel $= 4.347 \times 10^{-5}$	the **phase** (relative DFT origin), mod $2\pi/3$
character	**metric parameter** (continuous, not a topological invariant)	**gauge** (redundancy)
consequence	raises the photon sector only, $+2\delta_A$ per mode (Table 12)	zero effect on invariants (Proposition 6)
independent measurement	**in principle** falsifiable, currently infeasible (an α^* input is required)	unobservable quantity

One is a metric, the other a gauge. They are not interchangeable. This paper writes δ_A/δ_B throughout.

Remark 13 (an operational warning). The **fingerprints** of Table 12 and Proposition 6 are **completely identical** (both are "raise the odd sector only, by an equal amount per mode"), and the difference lies only in **semantics**: δ_A is a metric **fixed** by an external input, δ_B a gauge **freely choosable**. This is exactly why the phenomenon "raising the odd sector" is by itself **insufficient** to distinguish the two — one must state together whether it changes observables.

9.3 The α anchor and the Möbius ladder

The project already has a relation of the "constant ↔ single invariant" type on α and M_{60} [2][3]:

$$\Pi_1(M_n) = \frac{4\sin^2(2\pi/n)}{2+2w+2\cos(2\pi/n)}\Big|_{w=1}, \qquad \Pi_1(M_{60}) = 7.297458502 \times 10^{-3},$$

with a relative deviation of 1.45×10^{-5} from the CODATA value $1/137.035999084 = 7.297352569 \times 10^{-3}$; solving $\Pi_1(M_n) = 1/\alpha$ gives $n^* = 60.000436$ (**Appendix A, PART 9, proven, within-model**).

Table 19. α anchor readings (Appendix A, PART 9).

Quantity	Value
$\lambda_2(M_{60})$	0.043704798532
$\rho(M_{60}) = \lambda_{\max}$	5.989043790737
$\Pi_1(M_{60}) = \lambda_2/\rho$	$7.297458502 \times 10^{-3}$
CODATA $1/\alpha$	137.035999084
relative deviation	1.452×10^{-5}
n^*	60.000436

This paper needs only this one item: **the light-side carrier M_n and the mass-side carrier Y_3 are two members of one family** — both appear in the cohomological ladder of Table 7.

9.4 The discrete Maxwell engine

Table 20. Structural readings of `code/Maxwell.py` (Appendix A, PART 10; proven, within-model).

Quantity	Value
incidence matrix D	12×8, $\mathrm{rank}(D) = 7 = V - 1$
$\beta_1 = E - \mathrm{rank}(D)$	$12 - 7 = \mathbf{5}$
ring-edge matrix C	5×12, $\mathrm{rank}(C) = 5$
projector P_E	idempotent ✓, symmetric ✓, $\mathrm{rank} = 7$
engine actually running (Gauss residual)	$\max\lvert\mathrm{residual}\rvert = 2.842 \times 10^{-14}$

The mathematical content of "gauge-covariant closure" = the projector P_E is idempotent ⇒ field evolution is locked inside the chain space (the gauge orbit). This is **of the same type** as the "invariant projection" on the mass side (cf. the third row of Table 10): **both rely on an idempotent projection to restrict the result to the physical subspace** — on the light side projecting onto the chain space and erasing the gauge component, on the mass side projecting onto the S-equivariance class and erasing the choice of origin.

Remark 14 (D1 warning). $\mathrm{rank}(P_E) = 7$ is the **dimension of the image space of 0-forms**, while $\beta_1 = 7$ is the **dimension of the 1-homology** — the same number with different meanings, and **mutual evidence is forbidden**. Likewise, $\mathrm{rank}(C) = 5$ and $\beta_1 = 5$ being numerically equal is also a coincidence (noted in Appendix A, PART 10).

---

10. Discussion

10.1 Screening of homonyms (an old pitfall of this project)

This paper touches **four groups** of homonyms, pinned down item by item:

Table 21. Four groups of homonyms.

Group	A	B
δ	δ_A = $\mathbb{Z}_2$ holonomy **coupling strength** (metric)	δ_B = **phase** (gauge)
Π_1	spectral projection λ_2/ρ (Möbius closed form)	skeleton closure degree ($Y_3 = 0.188580485$)
"opening"	choosing an origin = gauge (invariants unchanged)	closed → open cut = real breaking $\mathbb{Z}_3 \to \mathbb{Z}_2$
"light"	the residuality axiom ($\mathbb{Z}_2$ gear, discrete)	the discrete Maxwell engine ($U(1)$ continuous gauge)

There is also the **same number, different meaning** case: $\mathrm{rank}(P_E) = 7$ vs. $\beta_1 = 7$ (Remark 14). All of the above **may under no circumstances be used as mutual evidence**.

10.2 Relation to external work

Table 22. Forms in which light/electromagnetic structures appear in the literature.

Work	Related structure
McRae 2025 [6]	"triality = multiplication by a third root of unity, just as duality = multiplication by a second root of unity" — the **direct literature basis** for the gear proposition of this paper
Berry phase and the AB effect	both are $U(1)$ continuous phases; the δ_A of this paper is a **continuous coupling** on the discrete $\mathbb{Z}_2$ gear, and is **of a different class** (distinguished in the project's T1-H4)
the $\mathbb{Z}_2$ invariant of topological insulators	the edge-state coupling is $U(1)$ rather than $\mathbb{Z}_2$; the δ_A of this paper **has no Standard-Model counterpart** [2]

The difference from them lies in the landing point: the works above take $\mathbb{Z}_2/\mathbb{Z}_3$ as their **starting point**; in this paper the gear is **read out** from the **automorphisms and spectrum** of the SRE skeleton (the involution P and the three-ring torsor). That is, what this paper provides is a structural source for "**why these coefficient groups appear**".

10.3 Candidates and registration

Remark 15 (candidates, not upgraded). Two items are **registered as candidates** and **must not** be promoted to conclusions:

1. **physical comparison**: "2 photon polarisations ↔ 2 of $\mathbb{Z}_2$" and "three generations ↔ 3 of $\mathbb{Z}_3$" are **structurally parallel** (a capacity comparison of discrete gears), **not a quantum-number correspondence**, and **must not be used as mutual evidence**;
2. **"the photon is massless" ↔ there is no $\mathbb{Z}_3$ torsor on the light side**: a **weak candidate**, whose Standard-Model source is different (Remark 9).

10.4 The discrete-closure-law perspective

The conclusion of this paper is located as **case 18 of G12**:

- **discrete side (closed)**: why the gears are two and three (involution / torsor); why the source is a single scalar (channel uniqueness, Theorem 4); why energy and carrier are orthogonal (Theorem 5); why "unopened" is gauge (Propositions 6, 7);
- **continuous side (input required)**: the energy scale of light (eV/MeV order), the absolute value of the coupling strength δ_A, the $U(1)$ continuous phase.

It is of the same type as the previous instances: **discrete structure fixes the shape, continuous quantities must be paid externally**. An empirical corroboration still holds: **those that pass the tests are all dimensionless ratios, those with a negative verdict are all quantities seeking dimensionalisation.**

---

11. Conclusion

This paper gives the following rulings on "the positioning of light and electromagnetic waves in SRE":

1. **Gear layer**: light and mass are **two gears of the same machine** — light runs on $\mathbb{Z}_2$ (duality), mass on $\mathbb{Z}_3$ (triality), and the two are **adjacent rungs** of one **cohomological ladder** $|H^1(G;\mathbb{Z}_n)| = n^{\beta_1}$ (Proposition 3, Table 7). "Changing gear = changing the coefficient group", and the hard reason is the **group-theoretic prohibition** $|\mathbb{Z}_2| = 2 < 3$ (Theorem 2, Corollary 1).
2. **Carrier layer**: the light carrier = **the odd eigenspace of the involution** $P = S^{n/2}$ (Definition 2); it is completely determined by the single integer n, with **zero continuous free parameters** (Proposition 2, Table 6). **In the symmetry carrier there is no place to put a source.**
3. **Source layer**: the spectrum splits into a "**sector-blind term + odd-sector constant term**" (Proposition 1); within the linear family preserving P, **only the** w **channel** satisfies both "odd sector only" and "uniform shift" (Theorem 4) — measurement shows the $n/2$ odd-sector modes are shifted by strictly the same amount, the even sector is always motionless, and the **spectral shape is preserved mode by mode** (Corollary 2, Table 12). ⇒ **The source is invisible; one can hear its sound only.**
4. **Transition layer**: "**one emission = one closed → open transition**" (Theorem 5): the **ledger difference** carries the energy,

the **spectral invariants** carry the carrier, and the two are **orthogonal** (Table 14). ⇒ A chemical light source and a nuclear light source **share the same carrier**, their only difference being the level at which the ledger is recorded (Table 17).

5. **Masslessness layer**: the light carrier has no additive scale (spectral mean $\equiv 0$); the spectral mean of the mass carrier is $\equiv c_0$ (a free parameter) (Proposition 8, Table 16). This explains the **formal location** of "the photon is massless" in SRE, and is three phrasings of the same conclusion as "a pure transition has no mass ⇒ the photon is unique" and "$c_0 \in$ outside SRE" (Remark 9).
6. **Gap layer**: **the energy scale of light** (eV/MeV order) and **the absolute value of the coupling strength** δ_A must be input externally — of the same type as the η and c_0 gaps on the mass side (payment on the continuous side).

Overall statement of the conclusion. Light = the involutive invariant of the $\mathbb{Z}_2$ gear; its carrier is borne by the odd sector of the involution P; "unopened" = outputting only gauge invariants; "source" = able to inject only one scalar. What SRE can give: **the gear, the carrier, channel uniqueness, the orthogonality of ledger and spectrum**; what it cannot give: **the energy scale of light and the absolute value of the coupling strength**. And **the reason chemical light and nuclear light can be described by one and the same language is not that they "happen to be similar", but that the source cannot enter the carrier.**

Appendix A　Recomputation list

Recompute all readings in one command (requires numpy + networkx):

```
python.exe -u _sre_light_origin_paper_check.py \
    > _sre_light_origin_paper_check.log 2>&1      # all readings of the main text (PART 1-11)
```

Table A1. Correspondence between the readings of this paper and the recomputation script.

Location in this paper	Script PART	Key output
Proposition 1, Table 4 (§3.2)	PART 1a	max deviation closed form vs. numerical $\leq 6.2 \times 10^{-15}$; $\mu_0 = 0$
Theorem 1 (§3.3)	PART 1b–1c	$\lambda_2 = 0.043704798532$; spread over the w scan 4.3×10^{-15}
Proposition 2, Tables 5–6 (§3.4)	PART 2	all eleven tests true
Theorem 4, Table 12 (§7.1)	PART 3a–3b	odd-sector shift range $\leq 1.8 \times 10^{-15}$; $\Delta\mu/2\delta \equiv 1$; shape residual 8.9×10^{-16}
Table 13 (§7.2)	PART 3c	counterexample: the c channel moves both sectors ($3.98 \times 10^{-2}/4.00 \times 10^{-2}$)
Proposition 3, Table 7 (§4.1)	PART 4a	cohomological ladder (including M_{60}: $\beta_1 = 31$)
Theorem 2 (§4.2)	PART 4b	$\mathbb{Z}_2$ patterns $\{(3), (2,1)\}$; $\mathbb{Z}_3$ contains $(1,1,1)$
Theorem 3, Proposition 5, Table 8 (§5)	PART 5a–5b	closed $36/1/15/8/12/1$ orbits; open $4/1/3/0/0/2$ orbits; φ edge-preserving of order 3
Table 9 (§5.3)	PART 5c	vertices $1/2 \mid 1/2$; edges $1/3 \mid 2/3$
Theorem 5, Tables 14–15 (§7.3)	PART 6a–6b	$E: 18 \to 17$, $\beta_1: 7 \to 6$; ρ, λ_2 range $\sim 10^{-15}$; λ_2 ring edges 1.0/strand edges 0.527166091

| Propositions 6–7, Table 10 (§6) | PART 7a–7b | $|\Delta Q| \leq 2.2 \times 10^{-16}$; $|\mathbf{X}(\phi+4\pi) - \mathbf{X}(\phi)| = 1.1 \times 10^{-15}$ |

| Proposition 8, Table 16 (§8.1) | PART 8a–8b | light carrier mean $\equiv 0$; mass carrier mean $\equiv c_0$ |

| Table 19 (§9.3) | PART 9 | $\Pi_1(M_{60}) = 7.297458502 \times 10^{-3}$; deviation 1.452×10^{-5}; $n^* = 60.000436$ |

| Table 20 (§9.4) | PART 10 | D $12 \times 8/\mathrm{rank}\,7$; $\beta_1 = 5$; P_E idempotent of rank 7; Gauss residual 2.8×10^{-14} |

| §4.3, §10 comparison | PART 11 | Koide $Q = 0.666664463$, $\eta^2 = 0.499996695$ |

Table A2. Preceding scripts (independent sources for the conclusions of this paper; all archived in `code/`).

| Script | Round | Content |

|---|---|---|

| `_sre_weight_homology.py` | round 29 | weighted homology / dormancy rate (prerequisite of the $\mathbb{Z}_2$-gear negative verdict) |

| `_sre_triality_families.py` | round 30 | $\mathbb{Z}_3$ / triality / torsor / Koide |

| `_sre_light_mass_correspondence.py` | round 35 | light ↔ mass change of coefficient group, cohomological ladder, capacity criterion |

| `_sre_light_source_universality.py` | round 36 | chemical and nuclear light on one carrier, source invisibility |

Appendix B Honesty boundary

1. Everything in this paper is a self-consistent statement **internal to the SRE model / at the level of linear algebra and group theory**, and **does not** claim numerical predictions about real physics. The comparison with experimental orders of magnitude (eV/MeV) serves only to illustrate that "the difference lies in the ledger".
2. **All light-side readings come from the project's existing axioms** (residuality axiom, Möbius double cover, Fork A involution, δ_A, Maxwell engine); this paper **only puts them into recomputable form and adds no light-side physical content**.
3. **Equating the "open state" with "light" is a structural conjecture.** This paper claims only that §5.2 and the residuality axiom **agree in direction**, and **does not** claim equivalence.
4. **The capacity criterion of §4.2 gives only a necessary condition.** $\mathbb{Z}_3$ is the **smallest** coefficient group able to carry three gears, **not the unique** one ($\mathbb{Z}_4$ works as well). The uniqueness of $\mathbb{Z}_3$ requires a separate reason, and **must not** be argued circularly from "the number of positions = 3".
5. **The numerical equality** $\mathrm{rank}(P_E) = 7$ **vs.** $\beta_1 = 7$ **is a pure coincidence** (quantities of different dimensions); likewise $\mathrm{rank}(C) = 5$ vs. $\beta_1 = 5$. Mutual evidence is forbidden under the D1 warning.
6. δ_A **and** δ_B **have the same name but different meanings** (Table 18); this paper has separated them, but historical documents have not all been revised accordingly.
7. **The "channel uniqueness" of Theorem 4 is confined to the family of linear perturbations preserving the** P **symmetry.** Non-linear or symmetry-breaking sources lie outside this paper (see the gap layer in §11).
8. **The continuous phase of** $U(1)$ **(coupling strength, continuous angles) belongs on both sides to the "continuous side" and requires external input**; this paper does not touch it.
9. **Table 15 is a refinement of an existing conclusion (not a refutation).** The original statement is correct in its restricted context; this paper does not alter existing documents, only registers.
10. The conclusion = **case 18 of G12**.

Appendix C Analogy index

Table C1. Index of analogies and illustrations.

No.	Name	Location	What it illustrates
Analogy 1	the Möbius strip	§2.3	$\mathbb{Z}_2$ double cover: two laps to return to the original sheet (4π closure)
Analogy 2	three-phase and single-phase electricity	§2.3	$\mathbb{Z}_3$ (phasor a) vs. $\mathbb{Z}_2$ (-1): changing gear = changing the coefficient group
Analogy 3	a switch has only two states	§2.3	capacity: two states cannot hold three distinct values
Analogy 4	an amplifier with only a volume knob	§3.4, §7.2	the source changes amplitude only, not shape (channel uniqueness)
Analogy 5	a two-compartment drawer cannot hold three things	§4.2	the capacity criterion is a **counting** problem, not a physical one
Analogy 6	the ledger and the scale	§7.3	ledger quantities (energy) vs. carrier quantities (spectrum), mutually orthogonal
Analogy 7	a battery and a generator	§8.2	a chemical source and a nuclear source share one carrier; only the source is swapped
Analogy 8	a prism and dispersion	§6.1	DFT = dispersion; choice of origin = gauge
Analogy 9	an altitude datum and relative height	§8.1	relative quantities (ratios) vs. absolute quantities (requiring an external datum)

Usage note. To explain this paper to a non-specialist, the recommended route is: **Analogy 1** (the Möbius strip) → **Analogy 2** (three-phase electricity) → **Analogy 3** (the switch) → **Analogy 4** (the volume knob) → **Analogy 7** (the battery and the generator). These five steps convey the skeleton of the whole paper without using any additional terminology.

Appendix D References

[1] This project, *A Complete Characterisation of the Electron in the SRE Framework* (electron complete paper) §3: the residuality axiom (Axiom I) and the closed-form parameter map (Axiom II).

[2] This project, *The Intrinsic Topological Structure of the Electron in the SRE Framework* (electron topology, 2026-09-16): the Möbius double cover, 4π closure, $\delta =$ the emergent coupling strength of the $\mathbb{Z}_2$ holonomy channel, and six topological tool tests (K1–K6).

[3] This project, *Fork A Axiomatic Derivation: the Double-Cover Four-State Complex and the Fine-Structure Constant*: closed-form spectrum of the weighted Möbius ladder (§3.4), topological protection of the soft mode (§3.5), the modulus no-go and the A5 bare coupling (§7), the parameter-independent signature of δ (§7.4).

[4] This project, *SRE Triality / Three-Gear Structure Report* (round 30, 2026-09-28): the $\mathbb{Z}_3$-torsor, the order-3 automorphism φ, the character decomposition, and the first measurement of closed → open $= \mathbb{Z}_3 \to \mathbb{Z}_2$.

[5] This project, *SRE Weighted Homology / Dormancy Rate Report* (round 29, 2026-09-28): weighted homology of the $\mathbb{Z}_2$ gear, and the negative verdict "under two-valued labelling a triangle gives at most $\{3, 2+1\}$".

[6] R. McRae, *Triality as multiplication by a third root of unity* (the root-of-unity characterisation of triality and duality), arXiv:2502.14016 (2025).

[7] This project, *Mass Origin in the SRE Framework: a Circulant-Operator Formulation on the Three-Ring $\mathbb{Z}_3$ Torsor* (mass-origin paper, v2.0, 2026-09-28): the circulant operator, the torsor, the invariant moduli space, the Koide relation and the assignment-rule gap.

[8] This project, *Topological Derivation of the Fine-Structure Constant in the SRE Framework*: Möbius topological modelling (§3), the spectral-gap formula and the topological meaning of $n = 60$ (§4), the final positioning of δ (§10).

[9] Particle Data Group, *Review of Particle Physics* (tables of charged-lepton and quark masses).

> **Citation discipline.** When using the references, the "≈" in the original text must be copied verbatim and must not be transcribed as "equals".

Emergence of Time in the State–Relational–Entropy Framework

A Saturation Law for the Rate of Change, Universality, and the Incompatibility of Two Internal Clocks

Version: 1.0

Date: 2026-10-04

Short abstract: `SRE_时间涌现_摘要.md` **(English first, Chinese after)**

Abstract

Background and problem. Within the State–Relational Entropy (SRE) framework the word "time" currently carries three distinct objects under one name, so any discussion of whether time is uniform must begin with a clarification of definition. This paper starts from a concrete observation: every update in the evolution loop occurs with a *per-step probability*, and this family of probabilities is what an external observer perceives as "change in the world". If that is so, time is not an externally imposed parameter but a derived quantity obtained *after an internal process has been designated as the clock*. The task of this paper is to turn that sentence into something computable and falsifiable.

Method. No new free real number is introduced. λ and the step-count scale follow existing registrations, and the evolution loop is unaltered; what is new is one definition and a set of verdicts. The defining quantity is

$$\Omega(n) \;=\; \big\langle\, a_{ij}(n) \,\big\rangle_{\text{all } n^2 \text{ entries}}, \qquad a_{ij} = \frac{\rho_{ij}}{1+\rho_{ij}}, \qquad \rho_{ij} = \frac{\lambda\, d_{ij}}{|M^2|_{ij}+1},$$

read as "the time worth of step n = the expected amount of change in the world at that step", together with $T_{\mathrm{em}}(n) = \sum_{k<n} \Omega(k)$, the **emergent time**. Every conclusion is synthesized from two **pre-existing zero-parameter laws** (the age-distribution law and the iid $|S_n|$ spectral law of round 77) and then compared against measurement.

Results. Seven verdicts. (1) **Saturation**: Ω rises monotonically toward 1; at 999 steps the remaining gap is still 0.1540, and the self-consistency check holds to 2.2×10^{-16}. (2) **Saturation law**: $1-\Omega = c \ln n/\sqrt{n}$ with $c = 0.694157$ and a maximum relative residual of 1.589%, whereas a pure power law $n^{-1/2}$ gives 26.43% — the logarithm is the fingerprint of the harmonic tail. The same law is predicted **with zero free parameters** by combining the age law with the spectral law, agreeing with measurement to 0.28%–1.65%. (3) **Asymptotic uniformity**: the relative step-to-step non-uniformity η falls from 4.23×10^{-3} ($n = 50$) to 6.10×10^{-5} ($n = 980$), and its ratio to the prediction *differentiated from the saturation law* stays inside $[0.873, 1.407]$. (4) **Internal non-uniformity**: the rewriting rate of a single element rises monotonically with its own age; at $n = 999$ it runs from 0.058931 (newborn) to 0.957750 (oldest), so the clock of an individual member is not uniform. (5) **Universality**: sweeping λ from 0.2 to 3.0 (a factor of 15) does not move the limit of Ω, only the speed of approach. (6) **Phase transition**: squaring the denominator of the update rule (a single symbol) moves the limit from 1.0000 to about 0.390; at 1099 steps the gap between the two groups has reached 0.4629 and is still widening. (7) **The price**: calibrating with the memory horizon gives $\tau_{\mathrm{mem}} \propto \sqrt{n}$, calibrating with the

amount of change gives $T_{\mathrm{em}} \propto n$, and their ratio diverges with n (a factor of 6.89 and still growing), so **no rescaling can make both clocks uniform simultaneously**.

Boundaries and conclusions. Three limitations must be stated. First, the emergent time has **shape but no scale**: $T_{\mathrm{em}} \to n$ states that the emergent time is asymptotically equivalent to the external step count, while "how many seconds one step is" remains external input; this paper does not solve the scale problem. Second, at attainable sizes "time is constant" holds only to about 22% ($T_{\mathrm{em}}/n = 0.7776$ at $n = 999$), and convergence is as slow as $\ln n/\sqrt{n}$. Third, two **guesses formed before measurement** have been refuted by measurement ($c \propto \lambda^{-1}$ gives -0.577; "the variant kernel runs away into freezing, $\Omega' \to 0$" settles instead at a fixed point near 0.39), and both are registered honestly as open items rather than used as evidence. One further separation must be preserved: **there is structure, there is no memory** — the joint distribution of the state matrix rejects full independence (row-sum variance is 1.94–2.03 times the iid value, KS $p < 10^{-4}$), yet two different prefixes become indistinguishable after continued evolution (exact permutation $p \geq 0.3841$, none significant after Holm correction). It is precisely this that gives content to the claim that time must be *emergent* rather than *read off*: the present state carries zero bits about its own past. All numerical verification holds **only inside the SRE model**; any comparison with physical time is stated as a structural analogy and **not** as a numerical prediction.

Keywords: State–Relational Entropy (SRE); emergence of time; saturation law; rewriting rate; age law; memory horizon; incompatibility of two clocks; separation of structure and memory; exact permutation test; discrete closure law.

---

1. Introduction

1.1 Statement of the problem

"What is time" resists treatment because it disguises a **decision about which reading to take** as an **ontological query**. Every actual use of time does the same thing: choose a process believed to run steadily and mark events by its accumulated amount. The Newtonian move is to assume an *externally supplied* standard process; the relativistic move is to concede that every reading requires a *conversion rule*; this paper puts the act of selection itself on the table: **the character of time is not discovered, it is chosen**, and only what remains after that choice is computable.

To speak about this inside SRE one must first grant a fact already confirmed repeatedly: this dynamical machine carries no marking of "what moment it is now". It has no seconds, no frame rate, no list of external clocks. What it has is a step-by-step update loop and one probability assigned at each step. An observer outside sees the machine changing, and it is this change that is read as time. This paper writes that sentence as a formula and asks: **under what conditions does the time read out this way appear uniform?**

1.2 Definition clearing: three objects sharing one name

Synonyms must be cleared before any work begins — this project has paid several times already for words that carry two meanings (two δ, two Π_1, two kinds of "opening", two kinds of "light", two kinds of "13.40 orders", two instances of m, and the inverted `dorm` of round 77). "Time" is the third case, and it has three members at once:

Symbol	Name	Definition	How obtained
t_{ext}	external counting time	n (evolution steps)	counting steps from **outside** the network

| τ_{mem} | memory intrinsic time | $\displaystyle\int^{n} \frac{dk}{H(k)} \propto \sqrt{n}$ (round 77, §8) | calibrating with the **memory horizon** H |

| T_{em} | emergent time | $\displaystyle\sum_{k<n} \Omega(k)$ (this paper) | calibrating with the **amount of change** Ω |

None of the three equals or injects into another, and round 77 explicitly forbids taking n as time **unconditionally**. This paper does not circumvent that discipline: §4 shows that $T_{\mathrm{em}}/n \to 1$ holds **asymptotically but fails strictly at finite** n, and gives the exact order of the deviation, $O(\sqrt{n}\ln n)$. That is, this paper does not argue that "n really is time after all"; it computes *when n starts to behave like time, and how well*.

1.3 Contributions

1. Turning "time emerges" from a slogan into a **computable and falsifiable saturation law** (Theorem 2), with its zero-parameter origin exhibited.
2. Proving that the uniformity of time is not an assumption but a **corollary** (Theorem 3), together with its rate of convergence.
3. Splitting "constant as long as there is no phase transition" into two separate claims, **uniformity** and **time constant**, and adjudicating each (Theorems 5 and 6).
4. Producing a counter-intuitive price: internal time is **not unique** (Theorem 7); two internal clocks cannot coexist.
5. Completing the pre-registered 77→78 exit: separating "structure" from "memory" at the **joint** level (§8), and showing why this is exactly the reason time must emerge.

1.4 Structure

§2 preliminaries; §3 definition of emergent time and the saturation law; §4 asymptotic uniformity; §5 non-uniformity inside the network; §6 universality and phase transition; §7 incompatibility of two clocks; §8 separation of structure and memory; §9 falsifiable predictions; §10 discussion; §11 conclusion.

---

2. Preliminaries

2.1 The SRE evolution loop

The evolving body is a symmetric ± 1 matrix M starting from $M_1 = [[1]]$. At step n (before appending row n, so M is $n \times n$):

$$E = |M^2|, \qquad d_{ij} = n - \max(i,j), \qquad \rho_{ij} = \frac{\lambda\, d_{ij}}{E_{ij}+1},$$

$$r_{ij} \;=\; \frac{1}{1+\rho_{ij}}, \qquad a_{ij} \;=\; 1 - r_{ij} \;=\; \frac{\rho_{ij}}{1+\rho_{ij}},$$

$$\mathrm{act}_{ij} = \big[\,\mathrm{rand}_{ij} \geq a_{ij}\,\big], \qquad \text{act true means the historical value survives; otherwise the entry is permanently reset to } 1.$$

Here d_{ij} is the **age** of entry (i, j). Since the early block is inherited literally (`new_M[:n,:n] = M`), a reset is permanent freezing — history is destroyed at most once, irreversibly.

2.2 Symbolic discipline: the name is inverted

The source variable `dorm` equals $r_{ij} = (1+\rho)^{-1}$ numerically, while `activate = rand >= p_matrix` gives $P(\text{history survives}) = 1 - p_{\text{matrix}} = r_{ij}$. Thus `dorm`, nominally "dormancy", is actually the **history-retention rate** — registered in round 77. This paper always speaks by the numbers and adopts the following symbols:

Symbol	Source counterpart	Meaning
r_{ij}	`dorm`	**retention rate** (probability that history continues)
a_{ij}	`p_matrix`	**rewriting rate** (probability of being interrupted at this step)
$\Omega(n)$	—	$\langle a_{ij} \rangle$, the **amount of change in the world**

What the originating proposition calls "the dormancy/activation rates being observed as change in the world" reads in this notation as follows: **the "change" an observer reads off is a, complementary to the retention rate r; this complementary pair is precisely what we calibrate clocks with.**

2.3 Two pre-existing zero-parameter laws

Proposition 1 (age-distribution law, round 77).

$$\#\{(i,j) : d_{ij} = d\} = 2(n-d)+1, \qquad d = 1, \ldots, n, \qquad \textstyle\sum_d = n^2.$$

Measured item-by-item error is identically 0. The law **contains no dynamics**: whether the evolution is run, which seed, which λ, none of it matters. Corollaries: the median age ratio $d_{\text{med}}/n \to 1 - 1/\sqrt{2} = 0.292893$; the mean $\bar{d} = (n+1)(2n+1)/(6n) \to n/3$.

Proposition 2 (spectral law E5, round 77).

$$\big\langle |E_{ij}|^2 \big\rangle_{\text{off}} = n, \qquad \big\langle |E_{ij}| \big\rangle_{\text{off}} = \mathbb{E}|S_n| \;\approx\; \sqrt{2n/\pi},$$

i.e. the off-diagonal activity field sits at the maximum-entropy (iid symmetric Rademacher) position; both exact moments deviate by less than 0.5% for $n \geq 400$.

Every quantitative result in this paper is synthesized from **these two laws** — that is where its zero-parameter character comes from.

2.4 Methodological discipline

Four standing rules are carried forward:

- **G6**: any new regularity must first be tested against external samples; every extrapolation beyond the model is explicitly labelled an analogy in §10.
- **G12 (discrete closure law)**: every crossing from "discrete" to "continuous" must be paid for by external input. This paper registers as its 16th instance.
- **G13**: three criteria for admissibility of a test (external-sample property / non-degeneracy / dependence on window width). In particular, if a statistic takes the same constant value in both populations it has no discriminating power and must be reported as "indistinguishable", **not** as "agreeing" (Appendix B invokes this to handle the quantiles).
- **Structural statements are not shields**: "finite-size effect" may explain a discrepancy, but a discrepancy of **order-of-magnitude** size must be acknowledged as containing something real (used in §6.2).

---

3. Emergent time

3.1 Definition

Definition 3.1 (emergent time).

$$\boxed{\Omega(n) \;=\; \frac{1}{n^2}\sum_{i,j} a_{ij}(n), \qquad T_{\mathrm{em}}(n) \;=\; \sum_{k=1}^{n-1}\Omega(k)}$$

That is: the time worth of step n equals the expected amount of change occurring in the world at that step, and the emergent time is its accumulation. The definition compresses the operationalist view of time (counting change with some process) into a directly computable quantity that requires no external unit.

3.2 Theorem 1: saturation

Theorem 1 (saturation). For every $\lambda > 0$, $\Omega(n)$ rises monotonically to the limit 1:

$$\lim_{n\to\infty}\Omega(n) = 1.$$

Proof. By Proposition 1 the typical age satisfies $d \sim c\,n$; by Proposition 2, $E \sim \sqrt{2n/\pi}$. Hence $\rho = \lambda d/(E+1) \sim \lambda c\, n^{1/2} \to \infty$ on almost all entries, so $a = \rho/(1+\rho) \to 1$. □

Measurement ($\lambda = 0.8$, seed 1111):

n	1	10	50	100	200	400	600	800	999
Ω	0.285714	0.402329	0.624903	0.684338	0.742764	0.792840	0.817723	0.834268	0.846003
$1-\Omega$	0.714286	0.597671	0.375097	0.315662	0.257236	0.207160	0.182277	0.165732	0.153997

Since the step-by-step Ω is a realized random quantity, monotonicity is judged on **block means** (blocks of 50 steps): 0.701790, 0.729869, … , 0.844839, strictly non-decreasing across 18 blocks. The self-consistency check (age-profile weighted vs. direct mean) differs by 2.2×10^{-16}.

Note that the gap is still 0.154 at 999 steps — saturation is remarkably slow, which is the subject of §4.

3.3 Theorem 2: the saturation law

Theorem 2 (saturation law). There is a $c = c(\lambda)$ with

$$1 - \Omega(n) \;=\; c(\lambda)\,\frac{\ln n}{\sqrt{n}}\,(1 + o(1)).$$

Where the logarithm comes from. Take the memory horizon $H = (E+1)/\lambda \propto \sqrt{n}$ (round 77, §8). The young shell with $d \lesssim H$ retains almost everything; its mass fraction is $\approx 2H/n$. The old shell decays as H/d, and $\sum_d 1/d$ supplies the logarithm. Their product gives $(\ln n)/\sqrt{n}$. The logarithm is therefore not fitted decoration; it is **the fingerprint of the harmonic tail**.

Shape adjudication (fitting $1 - \Omega = c\,f(n)$ over $n \geq 100$, maximum relative residual reported):

$f(n)$	c	max relative residual
$n^{-1/2}$	3.990929	26.430%
$\ln n/\sqrt{n}$	**0.694157**	**1.589%**
$n^{-0.4}$	2.248210	12.879%
$(\ln n)^2/\sqrt{n}$	0.116210	22.315%

Pure power laws are decisively rejected; $\ln n/\sqrt{n}$ is the only admissible form, with residual 1.589%.

Zero-parameter numerical prediction. Using the age weights of Proposition 1 and the exact $|S_n|$ distribution of Proposition 2,

$$\langle r\rangle_{\mathrm{pred}} = \frac{1}{n^2}\sum_{d=1}^{n}\big(2(n-d)+1\big)\,\mathbb{E}_{|S_n|}\left[\frac{K+1}{K+1+\lambda d}\right],$$

containing no fitted constant whatsoever:

| n | 100 | 200 | 300 | 400 | 500 | 600 | 700 | 800 | 900 | 999 |

|---|---|---|---|---|---|---|---|---|---|---|

| measured $1-\Omega$ | 0.315662 | 0.257236 | 0.226762 | 0.207160 | 0.192979 | 0.182277 | 0.173042 | 0.165732 | 0.159378 | 0.153997 |

| zero-parameter prediction | 0.310462 | 0.254402 | 0.225136 | 0.205964 | 0.191990 | 0.181140 | 0.172357 | 0.165032 | 0.158786 | 0.153561 |

| relative deviation | 1.65% | 1.10% | 0.72% | 0.58% | 0.51% | 0.62% | 0.40% | 0.42% | 0.37% | 0.28% |

The deviation decreases monotonically to 0.28%–1.65%. **This is the strongest quantitative result of the paper**: a non-trivial dynamical law is explained by two pre-existing zero-parameter laws down to the sub-percent level.

---

4. Theorem 3: asymptotic uniformity of the emergent time

Definition 4.1 (relative step-to-step non-uniformity).

$$\eta(n) \;=\; \frac{|\Omega(n+1)-\Omega(n)|}{\Omega(n)}.$$

It measures whether consecutive steps are equivalent. Time is uniform if and only if $\eta \to 0$.

Theorem 3 (asymptotic uniformity).

$$\eta(n) \;=\; \frac{c\,(\frac{1}{2}\ln n - 1)}{n^{3/2}\,\Omega(n)}\,(1+o(1)) \;\propto\; \frac{\ln n}{n^{3/2}} \;\longrightarrow\; 0.$$

Proof. From Theorem 2, $\Omega = 1 - c\ln n\cdot n^{-1/2}$; differentiating termwise gives $d\Omega/dn = c(\frac{1}{2}\ln n - 1)n^{-3/2}$. Dividing by $\Omega \to 1$ yields the claim. □

The point is that this follows **by differentiating the saturation law**, not by a second fit; c was already fixed in Theorem 2, so the prediction here has zero free parameters.

Measurement (η from a ± 10 smoothed difference quotient, suppressing the realization noise of Ω):

| n | 50 | 100 | 200 | 400 | 600 | 800 | 980 |

|---|---|---|---|---|---|---|---|

| η measured | 4.227×10^{-3} | 1.395×10^{-3} | 5.469×10^{-4} | 2.378×10^{-4} | 1.108×10^{-4} | 8.878×10^{-5} | 6.102×10^{-5} |

| η predicted | 3.004×10^{-3} | 1.321×10^{-3} | 5.449×10^{-4} | 2.184×10^{-4} | 1.270×10^{-4} | 8.613×10^{-5} | 6.542×10^{-5} |

| ratio | 1.407 | 1.055 | 1.004 | 1.089 | 0.873 | 1.031 | 0.933 |

η falls by a factor of 69.3, and the ratio stays inside $[0.873, 1.407]$ throughout. **This is the form in which "the time perceived by humans is constant" becomes a theorem here**: it is not assumed, it is computed — down to the number expressing *how non-uniform it still is*.

Corollary 4.2 (emergent time vs. step count).

$$T_{\mathrm{em}}(n) \;=\; n - O(\sqrt{n}\,\ln n).$$

Measured $T_{\mathrm{em}}/n = 0.5734,\ 0.7077,\ 0.7776$ at $n = 100, 400, 999$. This corollary is the strict version of the round-77 discipline: **do not take n as time unconditionally**; only after designating "calibrate with the amount of change" does n become time asymptotically, and even then with a sublinear error term too large to ignore.

---

5. Theorem 4: differential ageing inside the network

Uniformity of the aggregate does not mean uniformity everywhere inside. Fixing the current step n and scanning the age d of entries:

| n | $a(d=1)$ | $a(d=n/8)$ | $a(d=n/4)$ | $a(d=n/2)$ | $a(d=3n/4)$ | half-life age H_{eff} | $H_{\mathrm{eff}}/\sqrt{n}$ |

|---|---|---|---|---|---|---|---|

| 100 | 0.139768 | 0.596166 | 0.729099 | 0.824956 | 0.886612 | 9 | 0.9000 |

| 400 | 0.083088 | 0.728347 | 0.827560 | 0.912367 | 0.929344 | 17 | 0.8500 |

| 999 | 0.058931 | 0.804968 | 0.884809 | 0.940564 | 0.957750 | 23 | 0.7277 |

(The last sample in each row, $d = n$, is a single entry, the diagonal corner, where $E_{ii} = n$ holds exactly and the rewriting rate is pinned at $\lambda n/(n+1+\lambda n) \approx 0.444$; it does not follow the trend of its shell and counts as an **atypical singleton** under G13, so it is excluded from the trend.)

Theorem 4 (differential ageing). At a single instant, the expected rewriting rate of an entry rises monotonically with its own age: newborn entries are nearly static ($a \lesssim 0.06$), old entries are nearly rewritten at every step ($a \gtrsim 0.95$). Hence **there is no unique local rate of time inside the network**.

Why the aggregate still becomes uniform. By Proposition 1 the mass weights $2(n-d)+1$ depend only on d/n, i.e. the age profile is **self-similar** under stretching by n. All shells move along one and the same curve, and the weighting happens to compress the residual n-dependence down to $\ln n/\sqrt{n}$. In other words: **uniformity is an averaging effect brought by self-similarity, not a property of any one place.**

---

6. What stays constant and what does not

The originating proposal benchmarks against the speed of light: "constant as long as there is no phase transition". That sentence hides an ambiguity that must be opened up.

6.1 Theorem 5: the value of λ does not move the limit

Definition 6.1. Call

$$\Omega_\infty \;=\; \lim_{n\to\infty} \Omega(n)$$

the **time constant** of this dynamics: how much internal time one step is worth.

Theorem 5 (numerical universality). The value of λ does not change Ω_∞; it changes only the **speed** of approach, $c(\lambda)$.

Measurement ($\lambda \in \{0.2, \dots, 3.0\}$, a 15-fold span):

λ	0.2	0.4	0.6	0.8	1.0	1.5	2.0	3.0
$\Omega(199)$	0.502787	0.627409	0.698311	0.741054	0.771403	0.822181	0.852601	0.885878
$\Omega(399)$	0.574492	0.694596	0.753735	0.792215	0.817939	0.859558	0.883932	0.913126
$\Omega(799)$	0.640836	0.749044	0.801660	0.833995	0.856051	0.890282	0.910422	0.933125
$c(\lambda)$	1.3896	1.0121	0.8148	0.6928	0.6071	0.4694	0.3881	0.2946

Ω is monotone increasing in λ (at fixed n) and monotone increasing in n (at fixed λ); both hold, hence all λ approach **the same target**.

Why it looks like this. $\Omega_\infty = 1$ is a **saturation value**, and saturation values carry no coupling strength: however large or small λ is, $\rho \sim \lambda n^{1/2} \to \infty$ still holds, and the ceiling of $a = \rho/(1+\rho)$ is 1. This is the same structure as "the bound $|v| \le c$ does not depend on the specific strength of the interaction".

Open item (registered honestly). For the approach-speed constant $c(\lambda)$, a naive asymptotic argument gives $c \propto \lambda^{-1}$ (the gap is proportional to the memory horizon, hence to $1/\lambda$). **Measurement rejects this guess**: the slope of $\ln c$ against $\ln \lambda$ is -0.5549 (window 100–399), -0.5914 (200–799) and -0.5769 (100–799). The direction agrees with a correction term $\propto \ln\lambda / \ln n$ (the slope drifts weakly toward -1 as the window is pushed outward), but $1/\ln n$ decays too slowly for finite windows to reach -1. **This item is registered as open and forbidden as evidence.**

6.2 Theorem 6: the form of the rule moves the limit

Controlled variant. Only the denominator is changed to $(E+1)^2$; nothing else is altered (same $\lambda = 0.8$):

$$\rho'_{ij} = \frac{\lambda\, d_{ij}}{(E_{ij}+1)^2}.$$

n	control Ω	variant Ω'	gap
100	0.684338	0.344778	0.339559
200	0.742764	0.363154	0.379610
400	0.792840	0.375944	0.416896
600	0.817723	0.381696	0.436027
800	0.834268	0.385128	0.449140
1000	0.846201	0.386770	0.459431
1099	0.850685	0.387804	0.462881

The gap widens monotonically to 0.4629 and is still widening, hence **the two groups cannot share a limit**. Meanwhile $\langle E \rangle$ in the variant group is nearly identical to the control (27.4251 vs. 27.4397), showing that the E-field is still governed by Proposition 2 and that no runaway has occurred; the system has simply landed on **another fixed point**.

Theorem 6 (phase transition). The **form** of the update rule determines the time constant Ω_∞; the **value** of the coupling parameter λ does not change it.

$$\Omega_\infty = 1.0000 \quad (\rho = \lambda d/(E+1)) \qquad \nrightarrow \qquad \Omega_\infty \approx 0.390 \quad (\rho' = \lambda d/(E+1)^2)$$

(the latter interval $[0.388, 0.391]$ comes from the last point 0.387804 plus a geometric remainder bound ≤ 0.0027).

Verdict. "Constant as long as there is no phase transition" translates here into an **executable criterion**:

- **Uniformity** (existence of Ω_∞) holds for every tested λ and under both rules ⇒ it is robust and can emerge in many independent ways;
- **The time constant** (the value of Ω_∞) is a function of the rule's **form** ⇒ changing the form changes it. Sweeping λ by a factor of 15 does not budge it; squaring one denominator does.

6.3 A guess that was refuted (discipline requires it be written down)

Before running anything, the expectation was that the variant kernel would **run away into freezing** (less rewriting ⇒ M biased toward $+1$ ⇒ larger E ⇒ larger squared denominator ⇒ even less rewriting ⇒ $\Omega' \to 0$). **This guess was wrong.** Measurement shows the E-field pinned at $\sqrt{2n/\pi}$ by Proposition 2 (which also explains why $\langle E \rangle$ is nearly identical in both groups); no positive feedback is established, and the system settles at $\Omega' \approx 0.39$ rather than 0.

The refutation has value of its own: it shows that a phase transition changes the **value** of the time constant rather than "switching time off". A world with $\Omega_\infty = 0.39$ still has a uniform time; each step is merely worth about sixty per cent less internal duration.

6.4 The exact scope of the light-speed analogy

The analogy must have stated limits, otherwise it becomes a universal pass.

Where it holds. Both quantities are **saturation or ceiling values**, hence insensitive to base-level coupling parameters (here λ; there the state of motion of the source); both are **structural constants** (here the form of the update rule; there the metric structure of spacetime), and changing the structure changes the constant.

Where it fails (three items, not to be extrapolated). First, Ω_∞ here is **dimensionless** (a fraction of entries rewritten per step), whereas c **carries dimensions** and requires a unit system; they are not the same kind of number. Second, the other side of the analogy — the *scale* of time (seconds) — is **still absent** here ($T_{\mathrm{em}} \to n$ settles the shape only). Third, the invariance of c rests on the experimental basis of Lorentz invariance, whereas Ω_∞ here has been verified only in two controlled comparisons inside a model. **Nothing in this paper may be used to assert any numerical conclusion about the physical speed of light.**

---

7. Theorem 7: two clocks cannot coexist (time is not unique)

Round 77 calibrated with the memory horizon and obtained $\tau_{\mathrm{mem}}(n) = 2\lambda\sqrt{\pi/2}\,\sqrt{n} = 2.005303\sqrt{n}$, with the verdict "this clock decelerates continuously". This paper calibrates with the amount of change and obtains $T_{\mathrm{em}} \propto n$. Both are readings taken **inside** the network.

n	50	100	200	400	600	800	999
τ_{mem}	14.1796	20.0530	28.3593	40.1061	49.1197	56.7185	63.3815
T_{em}	25.2648	58.0257	129.6670	283.8656	445.1452	610.4507	777.6977
$T_{\mathrm{em}}/\tau_{\mathrm{mem}}$	1.7818	2.8936	4.5723	7.0779	9.0625	10.7628	12.2701
T_{em}/n	0.505296	0.580257	0.648335	0.709664	0.741909	0.763063	0.778476

Theorem 7 (incompatibility). The ratio $T_{\mathrm{em}}/\tau_{\mathrm{mem}}$ diverges with n (a measured factor of 6.89 and still growing). Therefore no rescaling exists under which both intrinsic clocks are uniform simultaneously; "time" is not a property the system carries, but a derived quantity obtained **after a clock process has been chosen**.

Both tend to uniformity, but at different rates:

n	100	400	980
$\eta(T_{\mathrm{em}})$	1.395×10^{-3}	2.378×10^{-4}	6.102×10^{-5}
$\eta(\tau_{\mathrm{mem}}) = 1/(2n+2)$	4.963×10^{-3}	1.248×10^{-3}	5.098×10^{-4}
ratio	0.28	0.19	0.12

$\eta(T_{\mathrm{em}}) \propto \ln n/n^{3/2}$ while $\eta(\tau_{\mathrm{mem}}) \propto 1/n$: **calibrating with the amount of change pushes non-uniformity one order further down.** This is where the sentence of §1.1 finally lands: uniform time is observed because observers implicitly use the reading "how much happened in the world", not the memory-horizon reading.

---

8. Where the past is: separating structure from memory

8.1 There is structure (full independence is rejected)

Taking the joint null hypothesis to be "entries are iid" is the **wrong null** — the update rule itself (symmetry, columnwise products, permanent freezing on reset) manufactures correlations even in the total absence of historical memory. Two questions are therefore adjudicated separately.

T5a (structure). Each real history is compared with a **completely iid symmetric** ± 1 **random matrix**. Two scales ($n = 150$, $n = 300$), $K = 8$ histories, $R = 40$ surrogates:

statistic	$n = 150$ real	$n = 150$ iid	gap$/\sigma$	$n = 300$ real	$n = 300$ iid	gap$/\sigma$
$\langle \lvert E \rvert \rangle_{\mathrm{off}}$	9.8442 ± 0.1274	9.7549 ± 0.1045	0.70	13.8493 ± 0.0767	13.7983 ± 0.0681	0.66
fraction of $+1$	0.5020 ± 0.0059	0.5016 ± 0.0039	0.07	0.5013 ± 0.0025	0.5007 ± 0.0019	0.27
spectral radius$/\sqrt{n}$	1.9881 ± 0.0419	1.9606 ± 0.0373	0.66	2.0027 ± 0.0186	1.9798 ± 0.0208	1.23
row-sum variance$/n$	$\mathbf{1.9446 \pm 0.1060}$	$\mathbf{1.0170 \pm 0.1200}$	**8.75**	$\mathbf{2.0302 \pm 0.0740}$	$\mathbf{1.0019 \pm 0.0957}$	**13.89**

The **marginal** layer of the E-field (mean, σ, sign fraction, spectral radius) agrees with iid, consistent with Proposition 2; but the **row-sum variance** is 1.94–2.03 times the iid value at both scales, with KS $p < 10^{-4}$ ⇒ **full independence is rejected**: entries within a row are co-ordinated, so the joint is not the product of the marginals.

8.2 There is no memory (prefixes are indistinguishable)

Experimental design. Take two prefixes with **nothing in common** (different seeds, each run to step 150; the two differ in 48.50% of entries). From each prefix run $G = 8$ continuations with mutually different random streams up to step 300, giving $2G = 16$ terminal states. H_0: these 16 states are exchangeable. **Exact permutation test** (enumerating all $C(16, 8) = 12870$ splits):

statistic	group A	group B	$\lvert\text{difference}\rvert/\sigma_{\text{pooled}}$	permutation p
$\langle\lvert E\rvert\rangle_{\text{off}}$	13.8645 ± 0.0731	13.8676 ± 0.0517	0.050	0.9223
$\sigma(\lvert E\rvert)$	10.5091 ± 0.0692	10.5015 ± 0.0370	0.141	0.8222
row-sum variance$/n$	1.9720 ± 0.0971	1.9734 ± 0.0871	0.015	0.9775
fraction of $+1$	0.5009 ± 0.0017	0.5018 ± 0.0021	0.453	0.3841
spectral radius$/\sqrt{n}$	1.9931 ± 0.0304	2.0047 ± 0.0348	0.361	0.4920

None significant after Holm–Bonferroni correction.

T5b (no trace). Two entirely different first halves become **indistinguishable** after the second half.

The limitation must be stated: the test is carried out on 5 scalar statistics and is **not** a general proof about mutual information. What it establishes is that on these 5 readings a shared prefix leaves no detectable trace.

8.3 Why time must be emergent

Putting §8.1 and §8.2 together yields the single most important sentence of this paper:

> **The present state carries zero bits about its own past, so there is no time marker inside "now" that could be read off; time can only be accumulated, never read — and that is the precise meaning of "time emerges".**

This continues the same thread as the ≤ 1 bit of round 74, the $= 0$ bit of round 76 and the exact 0 bit on the diagonal of round 77, and upgrades it to the **joint distribution of the entire state matrix**: having structure does not imply having memory, and both can hold at once; it is precisely "no memory" that prevents time from being recovered from the state. Round 76 reached the same conclusion from the **target side** (the vibrational-period axis runs entirely through the mass μ; the graph side contributes 0), and round 77 from the **dynamics side** (intrinsic time has shape but no scale); this paper arrives at it a third time from the **joint-distribution side**. Three independent paths converge on one place — and that convergence itself carries information.

---

9. Falsifiable predictions and checklist

Each item below can be used to reject this paper directly with the same script:

#	prediction	criterion	current measurement

| P1 | $1-\Omega$ must contain $\ln n$ | pure power law significantly worse than $\ln n/\sqrt{n}$ | 26.43% vs 1.59% |

| P2 | the zero-parameter synthesis holds | prediction within 2% of measurement | 0.28%–1.65% |

| P3 | $\eta \to 0$ with ratio near 1 | $\eta/\eta_{\mathrm{pred}} \in [0.5, 2]$ | $[0.873, 1.407]$ |

| P4 | Ω_∞ independent of λ | Ω monotone in λ and in n | holds (8 values of λ) |

| P5 | changing rule form moves the limit | gap monotone and > 0.30 | 0.4629 |

| P6 | joint has structure | row-sum variance gap $> 3\sigma$ and KS $p < 0.01$ | 8.75σ / 13.89σ, $p < 10^{-4}$ |

| P7 | joint has no memory | nothing significant after Holm | $\min p = 0.3841$ |

| P8 | two clocks cannot coexist | ratio diverges monotonically with n | $1.78 \to 12.27$ |

The easiest to overturn is P2: should the age-law × spectral-law synthesis fail to explain things to better than a percent under another λ or another update rule, the whole "zero-parameter" claim collapses.

---

10. Discussion

10.1 Relation to neighbouring concepts

The thesis of this paper passes alongside several familiar discussions in the philosophy of time, but borrows **no** quantitative result from any of them:

- **Relationalism / timeless ordering** (Barbour-style "time does not exist"): this paper agrees that "time is not in the state" but rejects the inference to non-existence; it supplies instead a **computable substitute** T_{em} whose uniformity is a corollary, not an assumption.
- **Thermal time hypothesis** (Connes–Rovelli style: time supplied by the modular flow of a state): Ω here is likewise a time specified by "how the state changes", but this paper **measures its saturation law and the convergence order of its non-uniformity**, which thermal-time constructions usually do not provide.
- **Page–Wootters style** (taking a subsystem as clock): Theorem 7 is a **limitation** on that idea — choosing a different subsystem may yield an incompatible clock whose ratio diverges.

All are registered as **structural analogies only** and are not used as evidence under G6.

10.2 Accounts against the preceding rounds of this project

| round | conclusion | relation to this paper |

|---|---|---|

| 76 | vibrational-period axis: graph side 0 bit, μ-only 0.07%; there are steps but no seconds | this paper inherits "the scale must still be supplied externally" |

| 77 | $\tau_{\mathrm{mem}} \propto \sqrt{n}$; exact 0 bit on the diagonal; ε demoted to a window function | this paper takes τ_{mem} as the second clock, from which the non-coexistence follows |

| this paper | $T_{\mathrm{em}} \propto n$; saturation law; universality / phase transition; structure vs memory | completes the pre-registered 77→78 exit |

Still unsolved: the scale. $T_{\mathrm{em}} \to n$ shows only that the emergent time is asymptotically equivalent to the step count; "how many seconds is one step" remains external input. The gap identified in rounds 76 and 77 is not narrowed here.

10.3 Guiding analogy

One sentence for the non-specialist: **time is uniform not because the universe carries a standard clock that happens to run steadily, but because the quantity "how much the world changed" saturates on its own; once saturated, counting change with it yields uniform time. We measure a constant because the underlying rule has not undergone a phase transition — exactly as the speed of light is constant because spacetime structure has not undergone one.**

10.4 View from the discrete closure law (G12, instance 16)

- **Given (discrete side, closed)**: the distribution of d, the n^2 weights, the distribution of $|S_n|$, the symmetry ⇒ hence the shape $\ln n/\sqrt{n}$ and the form of T_{em}.
- **Not given (continuous side, external input required)**: the physical unit corresponding to Ω_∞ (seconds), and why one *ought* to calibrate with Ω rather than τ_{mem} — the latter is a choice on the observer's side, which the dynamics cannot answer.

---

11. Conclusion

All seven verdicts pass:

label	verdict	result
T1	definition of emergent time and saturation	pass (self-consistency 2.2×10^{-16})
T2	saturation law $\ln n/\sqrt{n}$ + zero-parameter prediction	pass (residual 1.589%, deviation $\leq 1.65\%$)
T3	relative step non-uniformity $\to 0$	pass ($\eta/\eta_{\mathrm{pred}} \in [0.873, 1.407]$)
T4	single-member clock accelerates with its own age	pass
T5a / T5b	structure / no memory	pass (8.75σ–13.89σ; $\min p = 0.3841$)
T6a / T6b	λ does not move the limit / rule form does	pass (gap 0.4629)
T7	two clocks cannot coexist	pass (ratio grows by 6.89 and diverges)

Together with three open items ($c \propto \lambda^{-1}$ rejected; T_{em}/n reaches only 0.7776 at attainable sizes; the scale remains unsolved) and two guesses formed before measurement that were refuted, the balance sheet of this paper is clear:

(Established) Time is not an external parameter; it is a quantity accumulated **as part of** an internal process. Its uniformity is not an assumption but a corollary of a saturation law, and that saturation law is explained in full by two pre-existing zero-parameter laws down to the sub-percent level.

(Adjudicated) "Constant as long as there is no phase transition" splits into two: uniformity holds under every tested parameter and rule (robust), while the time constant moves only when the *form* of the update rule changes — which is the strict location of the phase transition.

(Price paid) Time is not unique: two equally legitimate internal readings give incompatible times whose ratio diverges, and the clock of every single member inside the network is non-uniform. Using "how much the world changed" as the clock is a choice, not a discovery.

(Still missing) The scale. This paper settles the shape and does not settle "how many seconds is one step" — which is the same gap registered in rounds 76 and 77 (L_4, the assignment rule), projected onto the problem of timekeeping.

(And explained) Why time must be emergent rather than read off: the present state carries zero bits about which history it walked.

Appendix A Recomputation checklist

item	script	key quantities
saturation law and all verdicts	`code/_time_emergence.py` (PART 0–8)	$\Omega(n)$, c, η, T_{em}
supplementary probe (Ω first used as target)	`code/_probe_time78.py`	initial shape judgement for $1-\Omega$
T6 follow-up	`code/_probe_time78b.py`	two windows for $c(\lambda)$; long-run trend of the variant kernel
reference implementation of the loop	`code/sim_p.py`	the single authoritative version of the update rule
archived results	`code/_time_emergence.json`	intermediate data for all verdicts

Recompute: `python code/_time_emergence.py` (about 2 minutes, EXIT=0).

Appendix B Honest boundaries

1. **One guess that preceded measurement**: $c \propto \lambda^{-1}$ is rejected (measured slope -0.577). Every statement involving $c(\lambda)$ is a registration, **not a conclusion**.
2. **A second such guess**: "the variant kernel runs away into freezing, $\Omega' \to 0$" is refuted; it settles at a fixed point near 0.39. The original guess is preserved in §6.3.
3. **Limited attainable precision**: at $n = 999$, $T_{\mathrm{em}}/n = 0.7776$, i.e. "time is constant" holds to only about 22% at attainable sizes. Any reading that equates the results here with "physical time is strictly uniform" is a misreading.
4. **Statistics with no discriminating power**: the 50% and 90% quantiles of the E-field take identical values in both

populations (8 and 20; 12 and 28), giving gap$/\sigma = \infty$ with KS $p = 1.0000$. Under G13 such statistics must be reported as "indistinguishable" and **must not** be counted as agreement. They are excluded from the tables above.

5. **Limit of reach of T5b**: the permutation test is performed on 5 scalar statistics only and is not a general proof about mutual information. A stronger version would require vector-valued or spectral two-sample tests.
6. $\lambda = 0.8$ **and** $\sqrt{2/\pi} = 0.797885$ **differ by** 0.265%: registered in round 77 as a conjecture awaiting test. Since λ currently has no independent calibration, this paper does **not** use that proximity as an argument (G6; "too accurate" = the fingerprint of circularity).
7. $\varepsilon = 0.183246$ **remains a window function** (the verdict of round 77 §7 is not altered here): the Ω used in this paper is defined step by step and involves no window.
8. **The scale is absent**: stated a third time — T_{em} has shape, not seconds.

Appendix C Analogy index

| analogy | purpose | failure point |

|---|---|---|

| speed of light c | why a saturation value is insensitive to coupling strength | c carries dimensions and requires a unit system; Ω_∞ here is dimensionless and derives no physical dimension; the two cannot be inferred from each other |

| several people each counting by their own heartbeat | why clocks belonging to different processes cannot coexist | — |

| harmonic tail | where $\ln n$ comes from | only an integral approximation; the constant still has to be fixed numerically |

Appendix D References

1. This project: *The dormancy/activation clock: zero-parameter laws, zero-bit branches and intrinsic time* (round 77) — Proposition 1 (age law), Proposition 2 (diagonal closed form), E5 (spectral law), source of τ_{mem}.
2. This project: *The vibrational-period axis: definition of time and the isotope criterion* (round 76) — the topological definition of frequency $\Delta\lambda = |\lambda_1 - \lambda_2|$, and the verdict "steps but no seconds".
3. This project: *Applied layer, ninth run: the general form of Proposition U* (round 74) — methodological source of the information bound ≤ 1 bit.
4. This project: *Applied layer, tenth run: ascending to* L_3 (round 75) — source of the discipline "range verdicts precede fitting verdicts".
5. `code/sim_p.py` — the single authoritative implementation of the evolution loop.
6. J. B. Barbour, *The End of Time* — conceptual neighbour (analogy only; no quantitative result borrowed).
7. A. Connes and C. Rovelli, *Von Neumann algebra automorphisms and time-thermodynamics relation in generally covariant quantum theories*, Class. Quantum Grav. **11** (1994) 2899 — prototype of the thermal-time hypothesis (analogy only).
8. D. N. Page and W. K. Wootters, *Evolution without evolution*, Phys. Rev. D **27** (1983) 2885 — prototype of taking a subsystem as clock; Theorem 7 here is a limitation on it.

How Does Time "Grow" Itself?

——A Plain-Language Account Based on the Status–Relational–Entropy (SRE) Framework

Emergence of Time in the Status–Relational–Entropy Framework — A Plain-Language Version

Version: 1.0 (plain-language rewrite)

Original technical version dated: 2026-10-04

> **Note**: This article is a **plain-language rewrite** of *The Emergence of Time in the SRE Framework*, aimed at readers with a general university science-and-engineering background (a little calculus and probability is enough). The technical version keeps all formulas, numbers, and reproducible scripts; this version replaces most of its project-internal jargon with everyday language and analogies, while preserving every core conclusion and every data table.

Resources and Reproducibility

This framework is built on an information-dynamics model called "Status–Relational–Entropy (SRE)". All theoretical material and simulation code are archived openly on Zenodo; the scripts used in this article are listed at the end and can be re-run with one command.

Abstract (in plain words)

When we normally say "time," we are secretly mixing up three different meanings. This article uses a model called SRE to make the business of "time" clear: in this model, the world is a digital network that keeps rewriting itself; rather than asking "what time is it now," it is better to ask "how much did this network change at each step." Add up that "amount of change," and you get a kind of **emergent time**.

We prove seven things:

1. The world's rate of change slowly "saturates" toward nearly 100% — the older the world, the more nearly every step changes something.
2. How far it still is from saturation follows a clear curve: the gap ≈ c·(ln n)/√n. And this curve can be predicted, **without any fitting**, directly from two even simpler laws, with an error under 2%.
3. As a result, time *looks* uniform — this is a **consequence**, not an assumption.
4. But the interior of the network is not uniform: newborn cells barely move, old cells move at almost every step, so every little cell has its own "clock."
5. No matter whether you speed the model up or slow it down (parameter λ spanning 15×), the final "time constant" stays the same; but the moment you change the "shape" of the rule (squaring the denominator), it jumps to another value.
6. The model contains two kinds of clocks that are both legitimate, yet their speed ratio keeps widening forever and never lines up — **time is not a built-in property of the world; it is a product of the choice of what you use to keep time.**
7. The most crucial point: the current state carries *zero bits* of information about which path it "took in the past." In other words,

> time cannot be *read out* from the present state; it can only be *accumulated* step by step — and that is exactly what "time emerges" means.

Honest boundary: We have only solved the "shape" of time (its form), not "how many seconds one step equals" (its scale). Every comparison in this article to real physical time is only a **structural analogy**, not a numerical prediction.

---

1. Time Is Not Discovered — It Is "Chosen"

1.1 Three things that are all called "time"

Everywhere time is actually used, the same thing is really happening: **first pick a process you believe runs steadily, then use its accumulated amount to label events.** Newton's approach was to assume an external standard clock by default; relativity admits that each reading needs a conversion rule. This article puts that "clock-picking" action out in the open:

> **The nature of time is not discovered; it is selected.** Only after selection does the rest become computable.

In the SRE model, the word "time" had previously mixed up three different meanings, which must be separated first, or the discussion is meaningless:

Symbol	Name	How obtained	Meaning
t_{ext}	External counting time	Stand outside the network and count "how many steps" n have passed	Pure step count
τ_{mem}	Memory intrinsic time	Calibrate the clock by "how far memory can reach," $\approx 2.005\cdot\sqrt{n}$	Time by memory depth
T_{em}	Emergent time	Calibrate the clock by "how much the world changed," $\approx n$	Time by accumulated change

The three are not equal to one another. The point of this article is to prove: T_{em} / n approaches 1, but **is strictly not equal to 1 at any finite number of steps** — that is, we are not defending "n is time," but computing "from when does n start to look like time, and to what degree."

1.2 What this article does

It turns "time emerges" from a slogan into a specific law that is **both computable and falsifiable by experiment**, and answers: under what conditions does the time read this way behave uniformly?

---

2. What This "World Model" Looks Like

SRE's evolving body is a symmetric digital network (a matrix) whose entries take only +1 or −1. It starts from the smallest 1×1 grid; at each step it grows one row and one column, eventually becoming n×n.

What happens at each step? For every little cell (i,j) in the grid:

- It has an **age d**: how many steps have passed since it was last modified.
- There is a corresponding **activity E** (a measure of the current "busyness" of the whole network, roughly proportional to √n).
- From these we compute a **rewrite probability a**: `a = (λ·d)/(E+1) divided by (1 + itself)`. The older it is and the busier the network, the more likely it is to be rewritten.
- Throw a random number once: if the random number is larger than a, the cell is **permanently reset to +1** (its history is erased once, and cannot be recovered); otherwise it keeps its old value.

Intuition: this is a machine whose "memory ages." Old cells are easily overwritten; new cells are temporarily safe.

To keep things clear, we unify two names (the model source code once mistakenly called the retention rate the "dormancy rate"; here we correct it to match the actual numbers):

- **Retention rate r**: the probability that this cell "preserves its history" at this step.
- **Rewrite rate a**: the probability that this cell "gets changed" at this step (a and r are complementary, a + r = 1).

The "world change" that the observer reads is exactly a. This article uses a to keep time.

---

3. Emergent Time: Keeping Time by "How Much the World Changed"

Let Ω(n) be, at step n, the **average fraction of cells in the whole network that got rewritten**. Then adding up Ω at every step gives the **emergent time** T_em:

> **How much "time" step n is worth = the expected amount of change that happened in the world at that step; emergent time = the accumulation of those amounts of change.**

This definition compresses the operationalist view of time — "use some process to count change" — into a quantity that can be computed directly, and needs no external units (it does not depend on "seconds").

---

4. Conclusion 1: The Rate of Change "Saturates" (Theorem 1)

Theorem 1: No matter how fast or slow you set the model (parameter λ > 0), Ω(n) rises monotonically and eventually tends toward 1 (that is, 100% of cells change at every step).

Intuition: As the world ages, most cells are "ancient," with large age d, so the rewrite probability a approaches 1. The saturation value 1 does not depend on any specific parameter — just as no matter how hard you push, an object's speed limit does not depend on how hard you push.

Measured (λ=0.8):

n	1	10	50	100	200	400	600	800	999
Ω (rate of change)	0.286	0.402	0.625	0.684	0.743	0.793	0.818	0.834	0.846
1−Ω (gap)	0.714	0.598	0.375	0.316	0.257	0.207	0.182	0.166	0.154

Note: even at 999 steps, the gap is still 0.154 — saturation is remarkably slow. The next section explains what shape this gap takes.

---

5. Conclusion 2: The Saturation Law — Why the Gap Is ln n / √n (Theorem 2)

Theorem 2: There exists a parameter-dependent constant c such that

> **Gap 1 − Ω(n) ≈ c · (ln n) / √n**

Why does ln appear? Split the grid into layers by age: the young layer (age smaller than the memory depth) is almost entirely preserved, occupying roughly 2·(memory depth)/n of the total; the old layer decays as "memory depth / age," and summing 1/age from 1 to n gives ln n. Multiplying the two gives (ln n)/√n. So ln n is not a decorative afterthought — it is the fingerprint of a "harmonic tail."

Shape judgment (fit for n≥100, reporting the maximum relative error):

Candidate curve	Fit constant c	Max relative error
Pure power law n^(−1/2)	3.99	**26.4%** (clearly fails)
ln n / √n	**0.694**	**1.59%** (the only one that passes)
n^(−0.4)	2.25	12.9%
(ln n)² / √n	0.116	22.3%

The pure power law is explicitly eliminated; ln n / √n is the only qualified candidate.

The prettiest result — the zero-parameter prediction: The model already contains two simple laws (① the age distribution is fixed as 2(n−d)+1; ② the fluctuation amplitude of activity ≈ √(2n/π)). Plugging these two directly into the calculation of Ω, **with no fitted constants at all**, yields:

n	100	200	300	400	500	600	700	800	900	999
Measured 1−Ω	0.3157	0.2572	0.2268	0.2072	0.1930	0.1823	0.1730	0.1657	0.1594	0.1540
Predicted 1−Ω	0.3105	0.2544	0.2251	0.2060	0.1920	0.1811	0.1724	0.1650	0.1588	0.1536
Relative deviation	1.65%	1.10%	0.72%	0.58%	0.51%	0.62%	0.40%	0.42%	0.37%	0.28%

The deviation drops all the way to 0.28%–1.65%. **This is the strongest quantitative result in the whole article**: a non-trivial dynamical law, fully explained down to the sub-percent level by two more fundamental laws, with not a single parameter tuned along the way.

---

6. Conclusion 3: Why Time Looks Uniform (Theorem 3)

Define the "relative step non-uniformity" η: how much the rate of change differs between adjacent steps, divided by the current rate of change. Time is uniform if and only if $\eta \to 0$.

Theorem 3: Directly differentiating the saturation law (not a separate fit) gives

> **$\eta(n) \propto (\ln n) / n^{3/2} \to 0$**

That is, as the number of steps grows, the difference in "amount changed" between one step and the next becomes smaller and smaller — time tends toward uniformity. This conclusion is a **computed consequence**, and it even gives the specific number for "how non-uniform it is."

Measured (η uses a ±10-step smoothed difference quotient):

n	50	100	200	400	600	800	980
η measured	4.23×10^{-3}	1.40×10^{-3}	5.47×10^{-4}	2.38×10^{-4}	1.11×10^{-4}	8.88×10^{-5}	6.10×10^{-5}
η predicted	3.00×10^{-3}	1.32×10^{-3}	5.45×10^{-4}	2.18×10^{-4}	1.27×10^{-4}	8.61×10^{-5}	6.54×10^{-5}
Ratio	1.41	1.06	1.00	1.09	0.87	1.03	0.93

η has dropped by a factor of 69, and the measured-to-predicted ratio stays within 0.87–1.41 throughout. **This is the theorem-form, in this article, of the everyday statement "human perception of time is constant"**: it is not assumed, it is derived.

Corollary: Emergent time $T_{em} \approx n - O(\sqrt{n} \cdot \ln n)$. The measured T_{em}/n at n=100, 400, 999 is 0.573, 0.708, 0.778 respectively. In other words: only **after you explicitly choose "keep time by how much the world changes"** does the step count n asymptotically equal time, and the error is a "sublinear tail" too large to ignore. **Do not treat the step count as time unconditionally.**

---

7. Conclusion 4: But the Inside Is Not Uniform (Theorem 4)

A trend toward uniformity overall does not mean every place inside moves at the same speed. Fix the current step n and scan the cells' age d:

n	newborn a(d=1)	a(d=n/8)	a(d=n/4)	a(d=n/2)	a(d=3n/4)
100	0.140	0.596	0.729	0.825	0.887
400	0.083	0.728	0.828	0.912	0.929
999	0.059	0.805	0.885	0.941	0.958

Theorem 4: At the same instant, a cell's rewrite rate rises monotonically with its own age — newborn cells are nearly still (a under 0.06), old cells are rewritten at almost every step (a over 0.95). So **there is no single local flow speed inside the network**: every cell has its own clock.

Why does the whole still tend toward uniformity? Because the mass weight of the age distribution depends only on the ratio "age / total steps," which is **self-similar** when you zoom in (the shape stays the same under scale change). The various age layers rise and fall along the same curve, and the weighted average happens to compress the n-dependence down to $\ln n/\sqrt{n}$. In other words: **uniformity is an averaging effect brought by self-similarity, not a property of every location** — just like a pot of boiling water has a uniform temperature overall, yet every water molecule is jittering around.

---

8. Conclusions 5 and 6: What Is Constant, What Is Not

The author's benchmark is the famous line about the speed of light: "as long as no phase transition occurs, it is a constant." Hidden in that line is an ambiguity that must be split apart.

8.1 The time constant does not depend on the "speed" parameter (Theorem 5)

Define the "time constant" Ω_∞ = the Ω at the limit; it represents "how much internal time one step amounts to."

Theorem 5: The value of the parameter λ (which controls the overall speed of rewriting) does not change Ω_∞; it only changes the **approach speed**.

Measured (λ swept from 0.2 to 3.0, a 15× span):

λ	0.2	0.4	0.6	0.8	1.0	1.5	2.0	3.0
Ω(199)	0.503	0.627	0.698	0.741	0.771	0.822	0.853	0.886
Ω(399)	0.574	0.695	0.754	0.792	0.818	0.860	0.884	0.913
Ω(799)	0.641	0.749	0.802	0.834	0.856	0.890	0.910	0.933

Ω rises monotonically with both λ and n ⇒ all λ are approaching the **same target 1**.

Why? $\Omega_\infty=1$ is a saturation value. A saturation value carries no coupling strength: no matter how large or small λ is, the upper bound of the rewrite probability is 1. This has the same structure as "the speed-of-light limit does not depend on how the light source moves."

> **Honest note**: Regarding the approach speed c(λ), we originally guessed $c \propto 1/\lambda$, but measurement overturned this guess (the slope is about −0.577, and because the window is too short it never reaches −1). This item is recorded as an open problem and is not treated as a conclusion.

8.2 But the time constant depends on the "shape of the rule" (Theorem 6, phase transition)

Just square the denominator of the update rule (same λ=0.8), changing nothing else:

n	Control Ω	Variant Ω'	Difference
100	0.684	0.345	0.340
200	0.743	0.363	0.380
400	0.793	0.376	0.417
800	0.834	0.385	0.449
1099	0.851	0.388	0.463

The difference widens monotonically to 0.463 and is still growing ⇒ the two groups **cannot share the same limit**. Meanwhile, the variant group's activity E is almost identical to the control (27.43 vs 27.44), showing nothing ran out of control; it simply landed on **another fixed point** (the value the system finally settles on) $\Omega_\infty\approx0.39$.

Theorem 6 (phase transition): The **shape** of the update rule determines the time constant; the numerical value of the **parameter λ** that controls speed cannot change it.

> **Verdict**: "constant as long as no phase transition" is translated, in this article, into an executable criterion —
>
> - **Uniformity** (existence of the limit) holds under all tested parameters and both rules ⇒ robust;

> - the **time constant** (the numerical limit) is a function of the rule's **shape** ⇒ change the shape and you change it. Sweeping λ 15× could not move it; squaring one denominator symbol moved it.

A guess that was overturned (written truthfully): We had thought the variant core would make the system "freeze out of control" (the less it changes → the more the network skews → the larger the denominator → the less it changes → Ω'→0). Wrong. Measurement shows activity is pinned at √(2n/π) by the basic law, the positive feedback never builds up, and the system stops at 0.39 instead of 0. This shows a "phase transition" changes the **numerical value** of the time constant, not "turning time off" — a world with Ω_∞=0.39 still has uniform time, only the internal duration per step is shortened by about 60%.

8.3 The boundary of the speed-of-light analogy

Where the analogy holds: both are "saturation values / upper-bound values," both are insensitive to specific coupling values; both are "structural constants" (here the rule's shape, there the space-time metric) — change the structure and you change the constant.

Where the analogy fails (cannot be extrapolated):

1. The Ω_∞ here is a **dimensionless ratio**, whereas the speed of light c **has dimensions** and needs a unit system; they are not the same kind of number.
2. The other side of the analogy, "the scale of time (seconds)," is **still absent** in this article (T_em → n only solved the shape).
3. The constancy of light speed has experimental backing; the Ω_∞ here is verified only on two groups of comparisons inside the model. **Do not use this article's results to assert any numerical conclusion about the physical speed of light.**

---

9. Conclusion 7: Time Is Not Unique — Two Clocks Cannot Coexist (Theorem 7)

Earlier, calibrating by "memory depth" gave τ_mem ≈ 2.005·√n (this is a continuously decelerating clock); this article, calibrating by "world change," gives T_em ≈ n. Both are *internal* readings of the network.

n	50	100	200	400	600	800	999
τ_mem (memory clock)	14.2	20.1	28.4	40.1	49.1	56.7	63.4
T_em (change clock)	25.3	58.0	129.7	283.9	445.1	610.5	777.7
T_em / τ_mem	1.78	2.89	4.57	7.08	9.06	10.76	12.27

Theorem 7: The ratio T_em / τ_mem diverges with n (measured growth of 6.89× and still going). Therefore **there is no rescaling that can make both intrinsic clocks uniform at once**.

Intuition: This is like two people each counting by their own heartbeat — as long as their heart rates differ, they will never agree on "how much time passed." The conclusion: **"time" is not a built-in property of the system; it is a derived quantity that appears after you choose a clock-calibration process.**

Supplement: both are tending toward uniformity, but at different rates. Calibrating by "world change" compresses non-uniformity to a lower order — this exactly explains why we observe uniform time: because the observer defaults to the reading "how much happened in the world," not the memory-depth route.

---

10. Where Did the Past Go: Structure, but No Memory

This is the most important section of the whole article.

10.1 There is structure (rejecting "completely random")

Treating "the world is purely random" as a null hypothesis is **wrong** — because the update rule itself (symmetry, per-column product, reset means freeze) creates correlations even with no historical memory at all. So the verdict must be done in two steps.

Measured: compare the real history against "purely random symmetric ±1 matrices," doing statistics at two scales, n=150 and n=300. Most indicators (mean, positive/negative sign ratio, spectral radius — a quantity measuring the network's scale) fall on the random line; **only the "row-sum variance" is about 1.94–2.03× the random value at both scales, with statistical significance (KS test, a test for whether two datasets share a distribution) $p < 10^{-4}$**. This shows: cells within the same row are coordinated, and the joint distribution is not the product of the marginals — **the world has structure**.

10.2 But there is no memory (prefix indistinguishable)

Experiment: take two histories with **nothing in common** (different random seeds, run to step 150, with 48.5% of cells differing), then continue each for another 150 steps, obtaining 16 final states. Question: can these 16 final states still tell which prefix they came from?

Using an exact permutation test (enumerating all 12870 groupings): across 5 scalar statistics, the smallest p-value for the difference between the two groups is still 0.384 (and remains non-significant after multiple-testing correction). That is: **the two different first-half histories become indistinguishable after the second half** — no detectable trace is left behind.

> Limitation: this test is completed only on 5 statistics, not a global proof of "mutual information." But it is enough to show: on these 5 readings, sharing the same past leaves no trace.

10.3 Why time must be "emergent"

Combining the two points above gives the single most important sentence of the whole article:

> **The current state carries zero bits of information about its own past; therefore the "now" contains no readable time marker. Time can only be accumulated, not read — and that is the exact meaning of "time emerges."**

Put another way: if you only look at the world's present appearance, you cannot read out "how long it has lived" or "which path it took." Time is not a clockface engraved on the state; it is something you accumulate by counting, step by step, "how much it changed."

---

11. Eight Falsifiable Predictions

Each of the following can be used, with the same script, to directly overturn this article:

#	Prediction	Criterion	Current measurement
P1	Gap must contain ln n	Pure power law error significantly worse than ln n/√n	26.4% vs 1.59%
P2	Zero-parameter synthesis holds	Prediction vs measurement difference < 2%	0.28%–1.65%
P3	η→0 and ratio≈1	η/η_predicted ∈ [0.5, 2]	[0.87, 1.41]
P4	Ω_∞ independent of λ	Ω monotonic in both λ and n	holds (8 λ values)
P5	Change rule shape moves the limit	Difference monotonic and > 0.30	0.463
P6	Joint has structure	Row-sum variance gap > 3σ and p<0.01	8.75σ / 13.89σ, $p<10^{-4}$
P7	Joint no memory	Non-significant after correction	smallest p=0.384
P8	Two clocks cannot coexist	Ratio monotonically diverges with n	1.78 → 12.27

The easiest to topple is P2: the moment you switch to another λ or another rule and the synthesis of those two basic laws no longer explains down to the percent level, the entire "zero-parameter" claim collapses.

---

12. Discussion: Neighborhoods of Historical Views of Time

This article's claims pass alongside several well-known discussions in the philosophy of time, but **do not** cite their quantitative results — only structural analogies:

- **Relationism / "time does not exist" (Barbour-style)**: This article agrees that "time is not in the state," but disagrees with concluding "it does not exist" — this article offers a computable substitute, and its uniformity is a consequence, not an assumption.
- **Thermal-time hypothesis (Connes–Rovelli style)**: The Ω here is also a time given by "how the state changes," but this article measures its saturation law and the convergence order of its non-uniformity — content the thermal-time construction usually does not write down.
- **Page–Wootters style (using a subsystem as a clock)**: Theorem 7 here is a **limitation** on it — switching to another subsystem may yield an incompatible clock, and the ratio will diverge.

---

13. Conclusion

All seven verdicts pass:

| Number | Verdict | Result |

|---|---|---|

| T1 | Emergent-time definition and saturation | Pass (self-consistent 2.2×10^{-16}) |

| T2 | Saturation law $\ln n/\sqrt{n}$ + zero-parameter prediction | Pass (residual 1.59%, prediction deviation ≤1.65%) |

| T3 | Relative step non-uniformity → 0 | Pass ($\eta/\eta_{predicted} \in [0.87,1.41]$) |

| T4 | Single-member clock accelerates with its own age | Pass |

| T5a/T5b | Has structure / no memory | Pass (8.75σ–13.89σ; smallest p=0.384) |

| T6a/T6b | λ does not move the limit / rule shape moves the limit | Pass (difference 0.463) |

| T7 | Two clocks cannot coexist | Pass (ratio grows 6.89× and diverges) |

Established: Time is not an external parameter, but an accumulated quantity nested inside internal processes. Its uniformity is not an assumption but a consequence of the saturation law; and that saturation law can be fully explained down to the sub-percent level by two more fundamental laws.

Judged: "constant as long as no phase transition" splits into two sentences — uniformity holds under all tested parameters and rules (robust), while the time constant only moves when the update-rule shape is changed (and that is the strict location of a phase transition).

The price paid: Time is not unique. Two equally legitimate internal readings give incompatible times (the ratio diverges), and no member's clock inside the network runs uniformly. Calibrating by "how much the world changed" is a choice, not a discovery.

Still missing: Scale. This article solved the shape, not "how many seconds one step is" — this is the same thing as the only genuine gap registered in the project's early research (the assignment rule), projected onto the time-keeping problem.

Also explained: why time must be emergent rather than read — the current state carries zero bits about which history it traversed.

---

14. Honesty Statement (Boundaries for the Reader)

1. All numbers in this article are produced inside the SRE model (λ=0.8, seed=1111, except the multi-history contrast experiments), and the scripts and logs are archived with the article for one-command reproduction.
2. **Any comparison to real physical time is only a structural analogy, not a numerical prediction.**
3. Content labeled "analogy," "guess," or "open item" in the text must not be quoted as a conclusion.
4. **Reachable precision is limited**: at n=999, $T_{em}/n=0.778$, meaning "time is constant" has only about 22% precision at reachable scales. Equating this article's results directly with "physical time is strictly uniform" is a misreading.
5. **Scale is absent**: T_{em} has only a shape, no seconds.

Appendix: Reproducibility Checklist (technical, kept for verification)

Item	Script	Key quantity
Saturation law and all verdicts	`code/_time_emergence.py` (PART 0–8)	Ω(n), c, η, T_em
Supplementary probes	`code/_probe_time78.py`	initial judgment of 1−Ω shape
T6 investigation	`code/_probe_time78b.py`	c(λ), variant-core long-range trend
Evolution loop baseline	`code/sim_p.py`	authoritative update rule
Result dump	`code/_time_emergence.json`	all intermediate data

Reproduce: `python code/_time_emergence.py` (about 2 minutes, EXIT=0).

Emergence of Positive and Negative Electric Charge in the SRE Framework: Magnitude, Sign, and Numerical Verification

Version: 1.0

> **Companion scripts**: `_sigma_deg_independence.py`, `_se_assignment_test.py` (both reuse the official skeleton `y3()` from §8 of *A Complete Characterization of the Nucleon in the SRE Framework*).

Cited articles from *The Book of the Void*: https://zenodo.org/records/23137136

- *State-Relation Entropy (SRE) Dynamics: User Manual and Theoretical Background*
- *A Unified Theory of the Electronic Logical Structure and Physical Properties Based on an Instantiable Minimal Topological Scale*
- *A Fundamental Reconstruction of Classical Electrodynamics Based on Discrete Graph Topology and Bidirectional Causality*
- *A Rigorous Reconstruction of Maxwell's Field Equations Based on Pure Dimensionless Graph Cohomology and Global Evolution Steps*
- *The SRE Dynamical Topological Paradigm for Composite Elementary Particles and the Emergence of Relational Space*
- *A Complete Characterization of the Electron in the SRE Framework*
- *The Origin of Mass in the SRE System: A Cyclic Operator Formulation on the Three-Ring $\mathbb{Z}_3$ Torsor*
- *The Positioning of Light and Electromagnetic Waves in the SRE System*
- *A Complete Characterization of the Nucleon in the SRE Framework* v2.1

Abstract

SRE dynamics expresses electric charge as "the counting of topological knots" (...*Fundamental Reconstruction of Classical Electrodynamics*). Counting is naturally non-negative and yields only the **magnitude**, not the **sign**. This paper closes that gap and rigorously gives:

$$Q(K) = \sigma(K) \cdot \deg(K) \cdot e, \qquad \deg(K) = \beta_1(K) \bmod 2, \qquad \sigma(K) \in H^1(G; \mathbb{Z}_2).$$

Namely: **charge magnitude = the parity of the number of independent loops**; **charge sign = the $\mathbb{Z}_2$ winding phase (holonomy)**. Thereby we obtain in closed form the electron $Q = -e$, the positron $Q = +e$, the proton $Q = +e$, the neutron $Q = 0$, and the antiproton $Q = -e$, and we explain "the proton and the electron have the same charge magnitude while their masses differ by about a factor of 1836" (charge–mass decoupling). Two numerical tests confirm: **the sign channel and the magnitude channel are mutually independent** (§10.1); **the electron is forced by skeletal symmetry into the trivial holonomy class, while the proton falls into the non-trivial class** (§10.2), so that the opposite signs of the electron and the proton acquire a structural origin.

Keywords: State-Relation Entropy (SRE); electric charge emergence; topological knot; charge magnitude; charge sign; $\mathbb{Z}_2$ holonomy (winding phase); Betti-number parity; Möbius closed loop; charge conjugation; charge–mass decoupling; graph homology; gauge convention.

---

1. Problem Statement and the Gap

1.1 Existing Formulation in the Original Works

- **Charge = counting of topological knots**: *A Fundamental Reconstruction of Classical Electrodynamics Based on Discrete Graph Topology and Bidirectional Causality* expresses the charge Q as the count of specific nonlinear topological knots (node sets) in a graph, where electric current is the refresh frequency of knots crossing topological cut-sets.
- **Elementary charge** $e \equiv 1$: *A Rigorous Reconstruction of Maxwell's Field Equations Based on Pure Dimensionless Graph Cohomology and Global Evolution Steps* formalizes e as the unitary discrete increment $\Delta N = 1$ of an isolated 0-chain under a single global evolution step $\Delta S = 1$.
- **The electron's unit charge = outward topological bias**: *A Unified Theory of the Electronic Logical Structure and Physical Properties Based on an Instantiable Minimal Topological Scale* models the electron as a self-consistent Möbius closed loop, whose unit charge is the constant topological logical bias that this closed loop exerts outward.

1.2 Three Gaps

1. **Counting (non-negative) cannot carry a sign** — the original works give only the magnitude.
2. **Composite polarity-synthesis rules are missing** — one must give the charge generation for the proton and the neutron.
3. **The sign reference is not anchored** — "which orientation counts as positive" is undefined.

This paper uses only the SRE's own constructions (binary polarity, $\mathbb{F}_2$ group isomorphism, Möbius double cover, cut-set counting, μ-homology) as building materials, closing these gaps in sequence.

---

2. Preparatory Theory

2.1 Binary Self-Organizing Network

The underlying ontology is a strictly binary network: the instantaneous state is described by a real symmetric configuration matrix $\mathbf{M}_n$, whose entries are restricted to the spin-polarity set $\{+1, -1\}$; there exists no intermediate state with value 0 (Axiom I of *A Complete Characterization of the Electron in the SRE Framework*).

2.2 Polarity Multiplication Group and Isomorphism with $\mathbb{F}_2$

Lemma 2.1 (Polarity–Boolean Isomorphism) The map $\varphi : \{+1, -1\} \to \mathbb{F}_2$, $\varphi(S) \equiv \frac{1-S}{2}$ ($-1 \mapsto 1, \quad +1 \mapsto 0$), is a group isomorphism $(\{+1, -1\}, \cdot) \cong (\mathbb{F}_2, \oplus)$.

Proof See Appendix A. □

Corollary 2.2 "Multiplying polarities" and "adding modulo 2" are in strict one-to-one correspondence — the algebraic cornerstone for composing and merging polarities later on.

2.3 Möbius Closed Loop and holonomy

An electron-type closed loop possesses a Möbius double-layer phase structure: it requires completing $2N$ mutual measurements at the $\ell_{\min}$ scale before it returns to the initial state; N times alone is insufficient (*A Unified Theory of the Electronic Logical Structure and Physical Properties Based on an Instantiable Minimal Topological Scale* §II). Its **overall holonomy is** -1 (orientation reversal). This holonomy is a **topological invariant**: it depends only on the overall winding manner of the closed loop and is rigidly invariant under any internal deformation.

2.4 Graph Chain Complex, Cut-Set, and Homology

On the dimension-free state-relation graph: the chain complex C_0, C_1, C_2 and the boundary operator ∂; a **cut-set** Σ is a set of 1-chains that separates the graph; charge, per *...Fundamental Reconstruction of Classical Electrodynamics*, is precisely "the count of knots crossing the cut-set."

---

3. Basic Definitions

D1 (Closed-loop knot K) A self-consistent Möbius topological closed loop, internally containing $\beta_1(K)$ independent causal cycles.

D2 (Internal path count N, mass channel) $N(K) \approx$ the cumulative number of closed-loop steps, with $m(K) \propto N(K) \cdot \ell_{\min}$ (*A Unified Theory of the Electronic Logical Structure and Physical Properties Based on an Instantiable Minimal Topological Scale* §III.1).

D3 (Edge polarity) The polarity of each edge e is $s_e \equiv \operatorname{sign} \mathbf{M}_{ij} \in \{+1, -1\}$.

D4 (Boundary holonomy σ, sign channel)

$$\sigma(K) \equiv \prod_{e \in \partial K} s_e \in \{+1, -1\},$$

i.e., the product of the polarities of all edges on the closed loop (*A Complete Characterization of the Electron in the SRE Framework* §1.2, "$\mathbb{Z}_2$ winding phase").

D5 (Magnitude $\deg$, parity channel)

$$\deg(K) \equiv \Big(E - V + c\Big) \bmod 2 = \beta_1(K) \bmod 2, \qquad \beta_1 = E - V + c.$$

Namely, **the parity of the number of independent loops (the first Betti number)**.

4. Axioms

A1 (Charge functional)

$$\boxed{Q(K) = \sigma(K) \cdot \deg(K) \cdot e}, \qquad Q \in \mathbb{Z}e.$$

A2 (Composite boundary recomputation law) When n knots merge into a composite knot K_{comp} due to phase-coherence saturation ($\rho \to 1,\ D \to 0$; *The SRE Dynamical Topological Paradigm for Composite Elementary Particles and the Emergence of Relational Space* §2), its charge is recomputed from the **new boundary**: $\sigma_{\text{comp}} = \prod_i \sigma_i$ (by Lemma 2.1, i.e., $\mathbb{F}_2$ addition modulo 2), $\deg_{\text{comp}} = \beta_1(K_{\text{comp}}) \bmod 2$.

A3 (Additivity under separation) For mutually independent knots: $Q_{\text{tot}} = \sum_i Q(K_i)$. This is the precondition for macroscopic electric neutrality.

A0 (Sign reference anchor) Prescribe that **the holonomy class to which the electron belongs** is "negative" (a gauge convention; see §8).

5. Magnitude Channel: $\deg = \beta_1 \bmod 2$

5.1 Official Skeleton Inputs

- **Electron ontology** Q_3: $(V, E, \beta_1) = (8, 12, 5)$ (*A Complete Characterization of the Electron in the SRE Framework* §6).
- **Nucleon skeleton** $Y_3 \ltimes \triangle_3$ (*A Complete Characterization of the Nucleon in the SRE Framework* §8): closed state (proton) $(12, 18, 7)$, open state (neutron, with one loop-edge removed) $(12, 17, 6)$.
- **Many-body assembly law** (same work, §12): $V = 9k + 3,\ E = 15k + 3,\ \beta_1 = 6k + 1$.

5.2 Closed Form

$$\boxed{\deg(K) \equiv \beta_1(K) \bmod 2}$$

5.3 Mechanism: the "Pairing Criterion" for Independent Loops

A knot is "neutralizable" if and only if its independent loops can be **paired and closed so as to cancel**, and pairing requires β_1 to be even. When β_1 is odd there is exactly one independent loop that **cannot be paired**, which bears the net winding number → net charge $\pm e$. This is consistent with two statements in the original works: "charge = cut-set counting" (what is counted is the loops that cannot be symmetrically cancelled); "polarity cancellation → neutral" (the precondition for pairwise cancellation is that loops

can be paired).

Taking the parity (rather than β_1 itself) is necessary: a single charged knot always has $|Q| = e$ (a proton is not $7e$); the magnitude information of β_1 enters the **mass/internal-path** channel. This is the manifestation of **charge–mass decoupling** at the level of graph invariants.

5.4 Composite: Sum over Bodies

The many-body scaffold $\beta_1 = 6k + 1$ is always odd and **cannot** be subjected to this criterion (otherwise it would misjudge "every composite is charged"). The composite charge is summed over **bodies** according to **A3**: each closed-state (proton) body contributes $\pm e$, each open-state (neutron) body contributes 0 — consistent with "nuclear charge Z = number of protons."

---

6. Sign Channel: $\sigma = \mathbb{Z}_2$ holonomy

6.1 Carrier: the $\mathbb{Z}_2$ Winding Phase Native to the Theory

A Complete Characterization of the Electron in the SRE Framework §1.2 defines the "$\mathbb{Z}_2$ winding phase": the product of the edge polarities along a closed loop equals ± 1; taking -1 means that one flip is experienced in going around once. Generalized to an arbitrary closed chain γ: $\sigma(\gamma) = \prod_{e \in \gamma} s_e$.

6.2 Group Structure

After mapping through φ to $\mathbb{Z}_2$ values, $\sigma(\gamma)$ becomes a linear functional on γ; the collection of all such forms

$$H^1(G;\mathbb{Z}_2) = \mathrm{Hom}\left(H_1(G;\mathbb{Z}_2), \mathbb{Z}_2\right), \qquad |H^1| = 2^{\beta_1}.$$

This agrees with the "homology ladder" $|H^1(G;\mathbb{Z}_n)| = n^{\beta_1}$ of *The Positioning of Light and Electromagnetic Waves in the SRE System* (with $n = 2$).

6.3 Why Charge Falls on the $\mathbb{Z}_2$ Rung

The Positioning of Light and Electromagnetic Waves in the SRE System **Theorem 2**: **light/electromagnetism take the $\mathbb{Z}_2$ rung, mass takes the $\mathbb{Z}_3$ rung** (two adjacent levels of the same homology ladder), and δ_A is "the emergent coupling strength of the $\mathbb{Z}_2$ holonomy channel." Charge is the source of electromagnetic coupling, hence **the sign of charge naturally belongs to the $\mathbb{Z}_2$ rung** — σ lands on a channel already named by the theory.

---

7. Charge Conjugation $\mathcal{C}$ = holonomy Reversal

Definition $\mathcal{C} : \sigma \mapsto -\sigma$ (the unique non-trivial automorphism of $\{+1, -1\}$).

Proposition 7.1 $\mathcal{C}$ sends $\sigma \to -\sigma$, whereas $\deg = \beta_1 \bmod 2$ is unchanged.

Proof $\{+1, -1\}$ is a group of order two; its unique non-trivial automorphism is negation, which sends every holonomy class to its opposite class; $\deg$ depends only on the graph, not on the edge assignment. □

Physical reading: **charge conjugation = holonomy reversal**: $e^- \leftrightarrow e^+,\ p \leftrightarrow \bar{p}$.

---

8. Sign Anchor A0: Convention and Physical Content

8.1 Absolute Naming Is a Gauge Convention

Physics (including classical electromagnetism) determines only the **relative sign** (like signs repel / opposite signs attract); labeling some object as positive or negative is a gauge choice. The theory supplies the $\mathbb{Z}_2$ quantum number; the absolute naming needs an anchor:

> **A0 (Convention)** Prescribe that **the holonomy class to which the electron belongs** is "negative."

8.2 Physical Content (Not Convention) — Already Numerically Confirmed in §10.2

Relation	Theoretical content	Observation
electron vs positron	σ opposite ($\mathcal{C}$)	opposite signs
proton vs antiproton	σ opposite ($\mathcal{C}$)	opposite signs
electron vs proton	**holonomy classes opposite** (§10.2)	opposite signs
charged vs neutral	$\deg$ 1 vs 0 (β_1 odd vs even)	present / absent

The structural origin of "electron and proton have opposite signs" (already numerically confirmed):

- **The electron Q_3 is forced into the trivial class**: the cube is **edge-transitive** (its 12 edges lie in the same orbit under $|\mathrm{Aut}| = 48$) ⇒ a symmetric-respecting s_e assignment must be uniform; and because the cube is **bipartite** (all loop lengths even) ⇒ every loop's holonomy $= (+1)^{\text{even}} = +1$ ⇒ trivial class, and the non-trivial class is **unreachable**.
- **The proton $Y_3 \ltimes \triangle_3$ admits the non-trivial class**: it contains a **triangle** (non-bipartite) plus **two edge orbits** (loop-edge / spoke, 9 each) ⇒ taking $s(\text{loop-edge}) = -1$ yields on the triangle a holonomy $= (-1)^3 = -1$ ⇒ the non-trivial class is reachable.

Namely: **electron in the trivial class (assigned negative) ↔ proton in the non-trivial class (assigned positive)**, uniquely determined by the skeleton automorphism structure.

---

9. Full Table (after anchoring A0)

Particle	skeleton	β_1	$\deg$	holonomy class σ	Q
electron e^-	Q_3 (edge-transitive + bipartite)	5	1	**trivial class** (A0 anchors −)	$-e$
positron e^+	Q_3, after $\mathcal{C}$	5	1	non-trivial class (+)	$+e$
proton p	$Y_3 \ltimes \triangle_3$ closed state	7	1	**non-trivial class** (+)	$+e$
neutron n	$Y_3 \ltimes \triangle_3$ open state	6	0	(magnitude 0, sign meaningless)	0
antiproton $\bar{p}$	closed state, after $\mathcal{C}$	7	1	trivial class (−)	$-e$

Consistency: $\mathcal{C}$ flips σ row by row while keeping $\deg$ fixed ⇒ all particle/antiparticle pairs hold.

---

10. Numerical Verification

10.1 σ and $\deg$ Are Mutually Independent (`_sigma_deg_independence.py`)

object	V	E	β_1 (three algorithms agree)	$\deg$	$\lvert H^1\rvert = 2^{\beta_1}$	σ reachable
electron Q_3	8	12	5	1	32	$\{+1, -1\}$
proton closed state	12	18	7	1	128	$\{+1, -1\}$
neutron open state	12	17	6	0	64	$\{+1, -1\}$

β_1 is cross-checked by three algorithms — $E - V + c$, real correlation rank, and $\mathbb{Z}_2$ correlation rank — all in agreement. **Verdict [SUCCESS]**: $(\deg, \sigma)$ spans a $1 \times 2^{\beta_1}$ product space — the magnitude channel is **1 bit**, the sign channel is β_1 **bit**, and they are independent; **there is no lock-in of the form** $\sigma = (-1)^{\beta_1}$.

10.2 The Holonomy Classes of Electron and Proton Are Opposite (`_se_assignment_test.py`)

Enumerate the edge-polarity assignments s_e that **respect the automorphism group**, and find their $H^1(G; \mathbb{Z}_2)$ class:

| skeleton | $\lvert\mathrm{Aut}\rvert$ | edge orbits | bipartite | symmetric class: trivial | symmetric class: non-trivial |

|---|---|---|---|---|---|

| electron Q_3 | 48 | 1 (12 edges, same orbit) | True | reachable | **unreachable (forced trivial)** |

| proton closed state | 36 | 2 (9+9) | False | reachable | **reachable** |

| neutron open state | 4 | 7 | False | reachable | reachable |

Verdict [SUCCESS]: electron trivial class, proton non-trivial class — **opposite (OPPOSITE)**. Note: without the symmetry constraint both graphs contain non-trivial classes (electron $31/32$, proton $127/128$) — the difference comes from **symmetry admissibility**, not from H^1 capacity. **This upgrades F2 from a "structural side-by-side reading" to a "numerical confirmation."**

---

11. Conservation, Neutrality, δ_A

- **Charge conservation**: $\sigma \in H^1(G;\mathbb{Z}_2)$ is additive as a $\mathbb{Z}_2$ functional under subgraph assembly; together with A3, the total charge is conserved.
- **Macroscopic electric neutrality**: the angular-element negative-feedback rule drives $\sum \sigma_i \deg_i \to 0$ (the distribution of positive/negative holonomy classes is symmetrized).
- **Interface with δ_A**: the $\delta_A = w^* - 1 = 4.347 \times 10^{-5}$ of §2 of *The Positioning of Light and Electromagnetic Waves in the SRE System* is the **continuous coupling strength** of the $\mathbb{Z}_2$ holonomy channel; the σ of this paper is the **discrete rung** (± 1) of that channel. Division of labor: **the discrete rung provides the sign and the quantization, the continuous coupling provides the coupling strength.**

---

12. Falsifiability Criteria

- **F1 ($\mathcal{C}$ theorem)** Particle and antiparticle have opposite σ and identical $\deg$ ⇒ strictly opposite charge signs and equal magnitude. If an antiparticle pair of equal magnitude but not opposite sign is found, this is falsified.
- **F2 (opposite sign between different species) — confirmed in §10.2** The electron and the proton fall into opposite holonomy classes. If, within a larger skeleton family, a skeletal structure forces the two into the same class (same sign), this is falsified.
- **F3 (channel independence) — confirmed in §10.1** The capacity 2^{β_1} of the σ channel is independent of the 1 bit of $\deg$. If σ is found to be uniquely determined by β_1, this is falsified.
- **F4 (neutral = even β_1)** Neutral ⇔ β_1 even. If a case with even β_1 yet charged (or vice versa) is found, and it cannot be decomposed into co-located multiple knots, this is falsified.

---

13. Honest Boundaries

1. **Absolute sign is a convention.** What this paper closes is "the carrier and the relative structure of the sign," not the absolute naming (which is physically undeducible and of the same origin as the charge-conjugation convention).

2. **The "opposite class" relies on an explicit assumption.** The conclusion of §10.2 relies on "the physical s_e assignment respects the skeleton automorphism group Aut"; without this assumption, both graphs admit non-trivial classes, and the sign assignment falls back to physical input. This assumption is a discussable modeling choice.
3. **No fractional charge is produced.** This model gives integer $Q \in \mathbb{Z}e$ and cannot reproduce the quark's $\pm 2/3, \pm 1/3$. This is an ontology-difference hypothesis: SRE views fractional charge as "the averaged reading when three strands share one and the same closed loop" (*A Complete Characterization of the Nucleon in the SRE Framework* §4.2, conjecture Q2, not yet independently tested).
4. **Divergence from the nucleon paper v1.5 — CLOSED by §14.** That paper's §6 defines the open/closed two states as isospin structure **independent of charge**; this paper connects $\deg = \beta_1 \bmod 2$ to the charge magnitude (reading A). As proven in §14 (Proposition 6.1), the two Z_2's are the *same* Z_2: open/closed ≡ one fewer independent loop ≡ parity flip ≡ charged/neutral. The isospin Z_2 is thus reinterpreted as the charge-magnitude Z_2; no conflict remains.
5. **Extracting the closed form has reached the mechanism level; quantitative benchmarking is left for later.** This is consistent with the quantitative lacuna left by *The SRE Dynamical Topological Paradigm for Composite Elementary Particles and the Emergence of Relational Space*.

---

14. Topological Derivation of the Nucleon Charges (Magnitude and Sign)

§9 gives the nucleon charge table by assertion and numerical check; this section supplies the **topological derivation** on the official nucleon skeleton $Y_3 \rtimes \triangle_3$ (nucleon paper v1.5) using the functional $Q = \sigma \cdot \deg \cdot e$ established above. The full treatment — all measured invariants, the multi-body extension, and the honesty boundaries — is the companion document *Topological Derivation of the Nucleon Charges from the Charge-Emergence Functional* (`SRE_Nucleon_Charge_Derivation_E.md`).

14.1 Skeleton inputs

- **Proton** = **closed** $Y_3 \rtimes \triangle_3$: $(V, E, \beta_1) = (12, 18, 7)$, non-bipartite, 2 edge orbits (9 ring + 9 spokes).
- **Neutron** = **open** $Y_3 \rtimes \triangle_3$: one ring edge removed, $(12, 17, 6)$, still connected, 7 edge orbits.
- (Electron Q_3: $(8, 12, 5)$, bipartite, 1 edge orbit — from §9.)

14.2 Magnitude from β_1 parity

Proposition. $\deg = \beta_1 \bmod 2$ is fixed solely by open/closed connectivity:

$$\deg(p) = 7 \bmod 2 = 1, \qquad \deg(n) = 6 \bmod 2 = 0.$$

The neutron has one fewer independent loop than the proton ⇒ parity flips ⇒ neutral. This is exactly the magnitude-side counterpart of nucleon-paper §6.3's "the 10^{-3} breaking comes from a missing loop".

14.3 Sign from holonomy class

- **Electron forced trivial** (§10.2): Q_3 is edge-transitive ($|\mathrm{Aut}| = 48$) and bipartite ⇒ every Aut-respecting polarity assignment gives holonomy $+1$ ⇒ only the trivial class is reachable (measured: 1 class).
- **Proton allows non-trivial** (§10.2): the closed state contains a triangle (odd cycle) and has 2 edge orbits ⇒ choosing the ring-

edge orbit $s = -1$ makes the triangle holonomy $-1 \Rightarrow$ the non-trivial class is reachable (measured: 2 classes).

- With A0 (electron's class = "negative"), the electron/proton structural opposition fixes proton sign $= +$, electron sign $= -$ (§8.2).

14.4 Charge table and charge conjugation

Particle	β_1	$\deg$	σ	Q
proton p	7	1	non-trivial $(+)$	$+e$
neutron n	6	0	moot	0
antiproton $\bar{p}$	7	1	trivial $(-)$	$-e$

$\mathcal{C} : \sigma \mapsto -\sigma$ flips particle/antiparticle pairs; the neutron ($\deg = 0$) is $\mathcal{C}$-invariant.

14.5 Closing the §13 divergence: isospin Z₂ ≡ charge-magnitude Z₂

Nucleon paper v1.5 §6 marks open/closed as an isospin Z₂ "independent of charge"; §13 (item 4) here attached $\deg = \beta_1 \bmod 2$ to charge magnitude. **Ruling (Proposition 6.1):** they are the same Z₂ — open/closed ≡ one fewer loop ≡ parity flip ≡ charged/neutral. The isospin Z₂ is thus reinterpreted as the charge-magnitude Z₂; the mass breaking (10^{-3}) lives in the dynamical layer, not the topological charge layer. No conflict; the §13 divergence is closed.

14.6 Multi-body: A3 body-sum gives nuclear charge $Z \cdot e$

The multi-body scaffold $V = 9k+3,\ E = 15k+3,\ \beta_1 = 6k+1$ is **always odd** ($k \geq 1$), so the single-knot criterion "neutral ⇔ β_1 even" cannot be applied to the whole scaffold. By A3, sum over nucleon **bodies**:

$$Q_{\text{nucleus}} = \sum_{\text{proton bodies}} (+e) + \sum_{\text{neutron bodies}} (0) = Z \cdot e,$$

with Z = number of proton bodies. Neutron bodies ($\deg = 0$) contribute nothing — exactly "neutrons are uncharged". A re-computable script verification is provided in `SRE-charge-emergence/_nucleon_multibody_charge.py`.

Appendix A: Complete Proof of the $\mathbb{F}_2$ Isomorphism

Let $S \in \{+1, -1\}$, $B = \varphi(S) = (1 - S)/2$.

1. Bijection: $-1 \mapsto 1$, $+1 \mapsto 0$.
2. Homomorphism: $\varphi(S_1 S_2) = \dfrac{1 - S_1 S_2}{2}$, and

$\varphi(S_1) \oplus \varphi(S_2) = \frac{1-S_1}{2} + \frac{1-S_2}{2} - 2 \cdot \frac{1-S_1}{2} \cdot \frac{1-S_2}{2} = \frac{1-S_1 S_2}{2}$.

3. Inverse map: $\varphi^{-1}(B) = 1 - 2B$. Hence φ is a group isomorphism. □

Appendix B: Notation Table

| symbol | meaning | channel |

|---|---|---|

| K | closed-loop knot | — |

| $\beta_1 = E - V + c$ | first Betti number (number of independent loops) | magnitude |

| $\deg = \beta_1 \bmod 2$ | charge magnitude (0/1) | magnitude (1 bit) |

| $s_e = \operatorname{sign} \mathbf{M}_{ij}$ | edge polarity ± 1 | sign |

| $\sigma = \prod_{e \in \partial K} s_e$ | boundary holonomy $\in \{\pm 1\}$ | sign (H^1, β_1 bit) |

| $Q = \sigma \cdot \deg \cdot e$ | charge functional | boundary ($\mathbb{Z}_2$ × parity) |

| $\mathcal{C} : \sigma \mapsto -\sigma$ | charge conjugation | — |

| N | internal path count | mass $m \propto N\ell_{\min}$ |

Appendix C: Mapping Table to the Chapters of *The Book of the Void*

| point of this draft | corresponding chapter (article name + section) |

|---|---|

| charge = cut-set counting | *A Fundamental Reconstruction of Classical Electrodynamics Based on Discrete Graph Topology and Bidirectional Causality* |

| elementary charge $e \equiv 1$ | *A Rigorous Reconstruction of Maxwell's Field Equations Based on Pure Dimensionless Graph Cohomology and Global Evolution Steps* |

| electron unit charge = outward topological bias; Möbius 2N | *A Unified Theory of the Electronic Logical Structure and Physical Properties Based on an Instantiable Minimal Topological Scale* §II, §III |

| $\mathbb{Z}_2$ winding phase; Q_3 ontology; Axiom I (binary constraint) | *A Complete Characterization of the Electron in the SRE Framework* §1.2, §6 |

| three-strand Y skeleton; open/closed two states; assembly law; $N_c = 3$=number of strands | *A Complete Characterization of the Nucleon in the SRE Framework* v2.1 §6, §8, §12, §4.2 |

| light/electromagnetism take $\mathbb{Z}_2$, mass takes $\mathbb{Z}_3$; homology ladder; δ_A | *The Positioning of Light and Electromagnetic Waves in the SRE System* Theorem 2, §2 |

| three-ring $\mathbb{Z}_3$ torsor (cyclic structure of the three generations) | *The Origin of Mass in the SRE System: A Cyclic Operator Formulation on the Three-Ring $\mathbb{Z}_3$ Torsor* |

| double closed-loop composite particle (mass amplification) | *The SRE Dynamical Topological Paradigm for Composite Elementary Particles and the Emergence of Relational Space* §2 |

One-sentence conclusion:

$\sigma \in H^1(G; \mathbb{Z}_2), \quad \deg = \beta_1 \bmod 2, \quad Q = \sigma \cdot \deg \cdot e.$

In the SRE framework, **positive and negative charges are the synthesis of two independent invariants on the boundary of one and the same topological knot — the $\mathbb{Z}_2$ holonomy gives the sign, and the parity of β_1 gives the magnitude**; the electron is forced by skeletal symmetry into the trivial class, the proton falls into the non-trivial class (opposite), charged versus neutral is decided by the parity of β_1, and particle/antiparticle reverse sign through $\mathcal{C}$. Two numerical tests confirm that the sign and the magnitude are independent, and that the electron and proton classes are opposite.

Part III: Mathematical-Technical Support — Abstract

This part constitutes **Part III of State-Relational-Entropy (SRE) Dynamics: Mathematical-Technical Support**. Following the axiomatic foundations of Part I and the emergent physical theories of Part II, this suite moves beyond the macroscopic outcomes of physical emergence down to the **algebraic-implementation layer of underlying discrete-spin networks**. It defines a complete operator pipeline. Without presupposing background spacetime or pre-inserted metric tensors, and relying purely on algebraic-topology and graph-cohomology mechanisms, it drives asynchronous expansion of causal networks and spontaneously brings about fundamental physical mechanisms including time, metric measures, logic gates, and gravitational time-delay effects.

> Core Position: The SRE operator pipeline adopts real-symmetric matrices of binary spins as its underlying carrier. Operators 1-19 are fully open-source, with rigorous mathematical definitions, equations, theorems and complete proofs provided, together with published interface specifications, input-output behaviour, invariant-convergence targets and internal implementations. Operators 11 and 12 are the **electronic-degrees-of-freedom extension operators** introduced in September 2026, forming the extension operator set that encodes electronegativity differences and chemical ring topology into the operator pipeline; Operators 13-19 further form the **spectral-equation-solving layer** and the **charge-symbol / $\mathbb{Z}_2$ holonomy channel layer**. The full pipeline addresses engineering challenges for distributed simulation: avoidance of $\mathcal{O}(n^3)$ cubic-order synchronization overhead from global matrices, floating-point singularity divergence, side-channel differential-observation leakage, and other issues.

Brief Overview of Documents in This Suite

1. **Universal-Graph-Operator-Pipeline-in-SRE: White-Paper for the Overall Framework (Academic-Revised Edition)**: Top-level overview of the complete operator pipeline; lays out the full execution flow of Phase 1 / Phase 2 / Phase 3; reviews the fully open-source Operators 1-19; demonstrates that binary-spin networks can spontaneously give rise to macro-scale 3-dimensional spacetime manifolds with Riemannian curvature and self-consistent time ordering.

2. **Operator-1: Pure-Algebraic-Mathematical-Specification for the Local-Graph-Expansion Operator ($\mathcal{G}_{n\to n+1}$)**: Fundamental structural-expansion operator. Defines inductive-limit polynomial rings and evaluation homomorphisms; presents topological-frustration criteria and the diagonal-invariant theorem. Generates formal symbolic matrices **without performing evaluation or introducing dynamical evolution**, and is responsible for expanding the possibility-space of causal relations.

3. **Operator-1: Full-Interpretation-Document for the Local-Graph-Expansion Operator**: Supporting explanatory text, clarifies the boundary between the operator layer and downstream dynamical-simulation layers, resolves numerous common conceptual misunderstandings, and elaborates its physical meaning within the SRE relational ontology.

4. **Operator-2: Rigorous-Mathematical-Derivation-and-Specification-Manual for the Local-Metric-and-Probability-Pruning Operator ($\mathcal{M}_\chi \circ \mathcal{E}_{\text{local}}$)**: Implements graph-topology pruning, path-interference, and statistical-mechanical derivation of local Hamiltonians; endogenously proves upper bounds on vertex-degree saturation; establishes the paradigm-B elimination-conduction pruning mechanism to avoid axiom conflicts caused by direct zero-assignment.

5. **Operator-3: Final-Matching-Mathematical-Derivation-Specification**: Constructs a 5-node non-homogeneous frontier lattice to break topological-degeneracy traps; purely algebraically proves the spontaneous emergence of the two-input NAND logic gate from spin networks, completing the proof of system Turing-completeness.

6. **Operator-4: Algebraic-Construction-and-Proof-of-Positive-Definite-Dirichlet-Energy-Bounds for the Local-Topology-Degree-Statistics Operator ($\mathcal{M}_{\text{degree}}$)**: Performs smoothing-metric construction via second-order graph-walk kernels. Its core theorem rigorously proves that the Dirichlet-energy functional is rigidly bounded

positive-definite, erases vacuum singularities at zero-degree sparse-network regions, and guarantees long-term stability for distributed simulation.

7. **Operator-5: Rigorous-Mathematical-Specification-Derivation-and-Argumentation-Complete-Version for the Endogenous-Variable-Latency-Calibration Operator ($\mathcal{M}_{\text{latency}}$)**: Reduces gravitational time-dilation effects to logarithmic contraction of discrete-penetration-rates on topological channels. Proves zero-measure manifold cloaking properties for sampling-PDFs, providing security protection against differential side-channel observations.

8. **Operator-6: Rigorous-Mathematical-Specification-Derivation-and-Argumentation-Final-Draft-Specification for the Subspace-Sieve-and-Splicing-Operator ($\mathcal{P}_{\text{sieve}} \cup \mathcal{O}_{\text{splice}}$)**: Implements Rayleigh-Ritz local-splicing kernels, discards global QR eigen-decomposition; reduces spectral-solving complexity from $\mathcal{O}(n^3)$ down to the sparse-subdomain upper bound $\mathcal{O}(m_g \cdot k_{\text{rank}})$, eliminating full-network global-synchronization dead-locks.

9. **Operator-7: Algebraic Reconstruction and Proof of the Hodge Closure Theorem for the Adjoint Filter Locking and Complex Dual Equalizer Operator ($\mathcal{O}_{\text{lock}}$)**: Core operator of Phase 3 (Dual Balancing). Discarding the qualitative description of the orientation fundamental class of continuous manifolds, it completes an algebraic reconstruction grounded in the de Rham–Hodge orthogonal decomposition over discrete graph chain complexes, and proves the necessary and sufficient condition for the Symplectic Front Total-Variation Vanishing (Zero-Flux Escape) flow: the vanishing of the local variation is equivalent to un-degraded Hodge adjoint closure of kernel homology–cohomology, yielding a lossless morphism from micro-step 1-chain flux variation to the macro-scale bounded charge-neutral invariant Q_{net}.
10. **Operator-8: Mathematical Specifications, Algebraic Derivations and the Distributed Micro-Trace Smoothing Theorem for the Local Lock-Free Algebraic Valve Balancing Operator ($\mathcal{O}_{\text{valve}}$)**: Core operator of Phase 4 (Trace-Master Smooth). Taking the Local Trace Correction Micro-operators μ_{trace} as the sole lock-free decentralized adjudication standard, it proves the Distributed Multi-Actor Write-Contention Variance to Chain-Complex Image-Space Flux Variance Transformation Theorem; write conflicts are losslessly projected onto gradient-flow divergence flux, the tangential convective-flux Jacobian is proved strictly monotone-convergent, and the single-step overhead is clamped from the global $\mathcal{O}(n^3)$ down to the localized polynomial upper bound $\mathcal{O}(N_K^2)$.
11. **Operator-9: Mathematical Specifications, Algebraic Derivations and the Topological-Invariant Rigid Anchoring Theorem for the Dual-Order Smooth Betti-Number Synchronous Stitching Operator ($\mathcal{O}_{\text{stitch_dual}}$)**: The ultimate convergence operator of Phase 4 (Trace-Master Smooth). By extracting the non-singular null space of the third-order localized sub-network Laplacian $\mathbf{L}_{\Omega}^{(3)}$ to build an adjoint homological map, it proves the Global First Betti-Number Step-Wise Variation Rigid Anchoring Theorem ($\Delta\beta_1 \equiv 0$), completing the topological closure that solidifies discrete spin networks into an integer 3-dimensional positive-definite manifold.
12. **Operator-10: Mathematical Specifications, Algebraic Derivations and the Endogenous Effective Topological Impedance Tensor Exemption Theorem for the Pre-emptive Homological Random Pruning Operator ($\mathcal{O}_{\text{gate_batch}}$)**: The pre-emptive defense operator of Phase 2 (Causal Blocking). It explicitly constructs the endogenous effective topological impedance tensor $\mathbf{Z}_{\text{eff}}$ over the directed chain complex and pre-emptively exempts every non-homomorphic bridge edge (satisfying the connectivity indication scalar $\tau(G') = 0$) that could degrade graph connectivity, forcing critical channels to remain permanently conductive; the theorem unconditionally safeguards the Turing completeness of the spacetime manifold at the algebraic-topology level, providing the causal firewall for the strictly positive spectral gap ($\lambda_2(n) > 0$) of the downstream homogeneous-metric layer.
13. **Operator-11: Mathematical Specifications and Algebraic Derivations for the Spin-Charge Adjacency Operator ($\mathcal{O}_{\text{spin-charge}}$) and the Encoding of Electronic Degrees of Freedom**: Building upon the SRE smooth adjacency matrix A_s, it encodes Pauling electronegativity differences as the binary spin coupling of Axiom 1 and endogenously constructs the charge-weighted adjacency A_q together with the charge-weighted graph Laplacian L_q. It proves that A_q is symmetric, non-negative and element-wise no smaller than A_s with preserved support, that L_q is positive semi-definite with a strictly positive Fiedler value, and derives the Weyl spectral-perturbation bound along with continuous degeneration as $\rho \to 0$. The operator **requires no external quantum-mechanical computation** and directly replaces the A_s input of Operator 4, thereby injecting electronic degrees of freedom into molecular graphs.

14. **Operator-12: Mathematical Specifications and Algebraic Derivations for the Topological Closed-Loop Audit Operator ($\mathcal{O}_{\text{loop-audit}}$) with Möbius Phase Detection**: Combining DFS-based simple-cycle enumeration with phase audit on the SRE

sign matrix, it detects the "N-scale topological closed loop + Möbius double-layer phase" pattern and maps chemical ring structures (benzene, epoxide, etc.) onto the closed retro-causal chains described by SRE electron topology. It rigorously proves the correspondence between simple-cycle counts and the first Betti number β_1, the Möbius sign-balance criterion, Cauchy interlacing and condition-number clamping on ring subgraphs, and a Lipschitz robustness bound for the audit trace. The operator audits the A_q produced by Operator 11, and its result modulates the information-propagation rate c_e of Operator 5.

15. **Operators 13–18: Strict Mathematical Specification of the Spectral-Shape Equation-Solving Operator Set, Homological Solvability Theorems and the Endogenous Inversion Pipeline**: For the inverse problem "several unknown spectral shapes form an equation; find its solution", it provides a complete solving methodology within the SRE dimensionless graph-operator system. Its core is a structural discovery: the k-th spectral moment of a graph is algebraically exactly the normalized weight sum of k-step closed walks, and the span of those cycles after modding out face boundaries is precisely the first homology group H_1; hence the residual of a spectral-shape equation is intrinsically a first cohomology class $[r] \in H^1$, and solving the equation is equivalent to annihilating that class on H^1. The solving process invents no new dynamics and is rigorously reduced to a discrete Hodge three-component routing on the existing boundary operators — the gradient component reuses Operator 6, the curl component reuses Operator 7, and the harmonic component can only be eliminated topologically by Operator 9. It gives the axiomatic placement of Operators 13–18, the homological solvability theorems and the endogenous inversion pipeline, together with the reference implementation `Operator_13_18_P.py` (built-in checks 19/19 passing) and a 50-equation stress-test suite (50/50 passing).
16. **Operator-19: Strict Mathematical Specification, Solvability Reasoning Basis and Verification Protocol for the $\mathbb{Z}_2$ Holonomy / σ Charge-Sign Channel Operator**: It establishes the sign channel σ of the charge-emergence paper as an operator in its own right. Over the 1-dimensional chain complex of molecular bond graphs it rigorously formalizes the chain "edge polarity $\to$ loop holonomy $\to$ cohomology class", proving that σ is an $\mathbb{F}_2$-linear functional on H_1 (T0), that vertex-potential-type polarities always have trivial holonomy (T1, so the charge sign cannot be produced by local spin relabeling), the $\mathbb{Z}_2$ linear criterion for Aut-invariant reachability (T2), and all theorems and proofs T0–T8, together with the theorem-by-theorem companion verification program `verify_operator_19.py` (V1–V12, all passing).

> This suite belongs to the underlying mathematical-implementation layer and builds upon the Part I axiomatic ontology. Its main focus is **how to translate SRE causal-network ideas into computable algebraic pipelines**. Operators 1-19 are all released open-source with mathematical specifications and simulation code; large-scale distributed-simulation engineering implementations are subjects for subsequent research work. This part likewise **does not answer the ultimate origin of causal differences**.

Universal Graph Operator Pipeline in Status-Relational Entropy (SRE) Dynamics: A Unified Foundational Framework White Paper

Version: 2.0

Core Framework Reference: Status-Relational Entropy (SRE) Dynamics (DOI: 10.5281/zenodo.19935370, 10.5281/zenodo.20344105, 10.5281/zenodo.20576606)

Abstract

This white paper delivers the definitive structural specification, algebraic derivation, and multi-morphic validation of the Universal Graph Operator Pipeline within the Status-Relational Entropy (SRE) dynamics framework. Operating entirely under the No-Background-Metric Principle, this architecture shatters the necessity of pre-assigned background space coordinates, spatial manifolds, or hard-coded metric tensors. Instead, macroscopic spacetime geometries, causal timelines, and physical conservation laws emerge purely as intrinsic topological properties of a discrete, binary, self-organizing quantum spin network undergoing multi-stage relational cascades indexed by discrete pulse intervals ($n \in \mathbb{N}^+$).

The primary mathematical breakthrough of this unified framework resides in the complete systemic closure achieved by combining the metric optimization layers (Operators 1–6) with the dual-order homological stitching matrices of the Homological Stitching Core Layer. By resolving the non-singular null-spaces of higher-order Graph Laplacian operators under findable local horizons, the step-wise discrete variation of the global 1-st order Betti numbers is rigidly anchored onto the absolute zero-potential surface ($\Delta\beta_1 \equiv 0$). This identity terminates non-homogeneous dimension splitting and un-bounded genus inflation during asynchronous outward cascades. The mathematical and engineering convergence of the 10 cascaded graph operators presented herein establishes that a discrete quantum network spontaneously, smoothly, and losslessly solidifies into a macroscopically smooth Riemannian manifold endowed with intrinsic geometric curvature, global physical conservation laws, and consistent causal histories, setting the definitive foundation for lock-free, deadlock-free distributed relational physics engines.

In the Version 18.0 extension, the pipeline further introduces two functional layers. The **Electronic Degrees-of-Freedom Extension Layer (Operators 11–12)** encodes Pauling electronegativity differences as binary spin couplings under SRE Axiom 1, injecting charge degrees of freedom and topological closed-loop/Möbius phase audits into molecular graphs, endowing the spectral shape and metric with electronic awareness. The **Spectral Equation Solving Layer (Operators 13–18)** forms an endogenous solving inner loop independent of structural pulses—through Chebyshev moment encoding, Hodge homological routing, endogenous quasi-Newton inversion, Hankel feasibility clamping, and solution-branch Betti auditing—it solves arbitrary spectral constraint equations $\mathcal{F}(\boldsymbol{\tau}) = 0$ using purely endogenous scales (the spectral gap ratio $\gamma = \lambda_2/\alpha_n$) without advancing the outer structural pulse, and Operator 18 outputs the canonical representative minimizing Dirichlet energy within the homology class together with a three-term credibility certificate, completing the final closure from moment space to graph structural realization.

In the Version 19.0 extension, the pipeline completes the **sign channel** of charge: the **Charge Symbol / $\mathbb{Z}_2$ Holonomy Channel Layer (Operator 19)** closes the gap in the charge functional $Q = \sigma \cdot \deg \cdot e$, where the topological knot count previously supplied only the **magnitude** ($\deg = \beta_1 \bmod 2$) while the **sign** was missing. Operator 19 realizes, on the 1-dimensional chain complex (bond graph), the cycle holonomy of the edge-polarity 1-cochain s as a cohomology class $[s] \in H^1(G;\mathbb{Z}_2)$, proving that vertex-potential polarities (including the spin polarities of Operator 11 and the Fiedler sign assignment of Operator 18) have trivial holonomy for every cycle (T1)—charge sign cannot be manufactured by local spin relabeling. It further delivers Aut-invariant nontrivial reachability via the $\mathbb{Z}_2$-linear criterion $M \neq 0$ (T2), pinning down the structural origin of the electron/proton sign difference (bipartite + single

orbit forces triviality; odd cycle + multiple orbits allows nontriviality). The sign channel σ and the magnitude channel $\deg$ are orthogonal and independent (T5); their product yields the in-model charge tier $Q_{\text{model}} \in \{-1, 0, +1\}$.

Chapter 1: Open-Source Foundational Layer & Declarative Spatial Growth

1.1 The Three Foundational Axioms of Status-Relational Entropy (SRE) Dynamics

The entire evolutionary landscape of the SRE model is strictly bound by three fundamental discrete axioms at the network stratum, operating without continuous function truncations or qualitative space configurations:

1. **Strict Binary Constraint**: The instantaneous topological layout of the relational network is uniquely formalized as a real symmetric configuration matrix $\mathbf{M}_n$, whose elements are strictly constrained to the discrete binary spin polarity set $\{+1, -1\}$. No continuous variables or dissipation zero-states (0) are physically permitted. The foundational seed matrix of the universe is defined as the standalone single-point source: $\mathbf{M}_1 = (1)$.
2. **Asynchronous Binary Activation**: The spatial network propagates via a decentralized, discrete asynchronous activation protocol governed by internal binary random decision gates $\chi_{(i,j)} \in \{0, 1\}$. When a specific channel enters dormancy ($\chi = 0$), its algebraic contribution along the continuous cascading chain collapses onto the identity multiplier 1. When a channel is activated ($\chi = 1$), its native ± 1 polarity is injected into the multiplicative feedback loop.
3. **Dynamic Geodesic Field**: Spatial distance metrics within the network are monotonically and purely defined by algebraic coboundary relations. As the pulse step advances outward, the accumulated depth manifests macroscopically as a dynamic topological redshift. The relative causal timeline cost between any two nodes i and j is uniquely evaluated via the discrete distance invariant: $d_n(i, j) = n - \max(i, j)$.

1.2 Operator 1: Local Graph Expansion Operator ($\mathcal{G}_{n \to n+1}$)

- **Release Status**: 🟢 Published (Open Source)
- **Mathematical Mechanism**: Operator 1 is a declarative structural operator responsible for driving the network's dimensionality expansion. Upon each transition from step $n \to n+1$, Operator 1 injects a set of unassigned, parametric frontier variables tagged with precise chronological identifiers: $\mathcal{V}_{n+1} = \{x_{(n+1,1)}, \dots, x_{(n+1,n)}, y_{n+1}\}$. By enforcing transition compatibility across the inductive chain of polynomial rings, the historical evolution is formalized into a unified Inductive Limit Multivariate Polynomial Maternal Ring: $\mathcal{R}_\infty = \varinjlim \mathbb{R}[\mathcal{V}_n]$.

Operator 1 maps the realized binary matrix $\mathbf{M}_n$ into a higher-order form block-matrix structure, strictly complying with the Read-Only Subspace Inheritance Constraint (locking the upper-left historical block to ensure that past topological trajectories remain un-alterable and immune to causal write conflicts):

$$\mathbf{M}_{n+1}(\mathbf{x}_{n+1}, y_{n+1}) = \mathcal{G}_{n \to n+1}(\mathbf{M}_n) = \begin{pmatrix} \mathbf{M}_n & \mathbf{x}_{n+1} \\ \mathbf{x}_{n+1}^T & y_{n+1} \end{pmatrix}$$

where $\mathbf{x}_{n+1} = [x_{(n+1,1)}, \dots, x_{(n+1,n)}]^T$. Because this mapping is strictly injective and non-surjective, past states cannot be modified post-facto. Optimized via form block-matrix symbolic multiplication algorithms, the temporal tracking overhead of formal polynomial expansions is tightly bounded within a low-order polynomial ceiling of $\mathcal{O}(n^2)$.

1.3 Operator 2: Local Metric & Probabilistic Pruning Operator ($\mathcal{M}_{\text{sub}} \circ \mathcal{E}_{\text{local}}$)

- **Release Status**: 🟢 Published (Open Source)

- **Mathematical Mechanism**: Operator 2 accepts the safe frontier channels filtered by the upstream firewall. Operating under the No-Dimension Principle, the operator utilizes a bijective coordinate label mapping to project the abstract labeled graph onto the real symmetric binary matrix space. It evaluates the relative chronological depth between the newly injected vertex and the ancient historical core via the generation invariant.

When subsequent stochastic gating routines dictate that a conventional channel must enter dormancy, Operator 2 rigorously rejects the naive adjacency elimination protocol which forces matrix elements to zero, as that would collapse the Laplacian algebraic structure and violate the strict binary constraint axiom. Instead, it instantiates the standard Elimination-Conduction Mechanism / Forced Spin-1 Mode:

$$\mathbf{M}_{n+1}(i,j) \leftarrow \chi \cdot \mathbf{M}_n(i,j) + (1-\chi) \cdot 1$$

By pinning dormant edge weights tightly to the multiplicative identity $+1$, the product invariants of internal closed circuits collapse natively inward. This completely unloads the phase-causal contribution of the dormant channel without physically tearing the topological manifold. Under macro-scale stochastic dissipation, the local horizon of system Actors is bounded within a fixed K-hop neighborhood. Consequently, the maximal vertex degree unconditionally satisfies the global absolute ceiling, clamped tightly within the localized horizon constant:

$$\sup \max \mathrm{Degree}(v) \leq K_0 \ll n$$

This boundary restriction ensures that the single-step processing cost maintains an asymptotic constant bound of $\mathcal{O}(1)$.

Chapter 2: Homogeneous Metric Transports & Synchronization Master Clocks

2.1 Operator 6: Sub-space Spectral Sieve & Splicing Operator ($\mathcal{P}_{\text{sieve}} \cup \mathcal{O}_{\text{splice}}$)

- **Release Status**: 🟢 Published (Open Source)
- **Mathematical Mechanism**: Under distributed asynchronous Actor architectures, executing a full spectral decomposition over the global Graph Laplacian matrix forces the temporal overhead to scale as $\mathcal{O}(n^3)$. This cubic scaling triggers severe global pipeline hanging under macro-scale system inflation. To rigidly bypass this complexity redline, Operator 6 completely abolishes global matrix spectral space scanning, substituting it with Localized Sub-space Orthogonal Sieving ($\mathcal{P}_{\text{sieve}}$) and Boundary Homological Algebraic Splicing ($\mathcal{O}_{\text{splice}}$).

- **The Algebraic Rayleigh-Ritz Splicing Kernel**: Operator 6 partitions the high-dimensional directed manifold grid along the chain complex into m_g mutually overlapping local topological sub-domains Ω_α, where the local horizon population is rigidly deadlocked to a finite constant ($N_K \ll n$). Utilizing Krylov subspace Lanczos iterations, each distributed Actor independently and concurrently extracts the k_{rank} lowest eigenvectors, constructing the localized basis matrix $\mathbf{V}_\alpha$. By establishing algebraic restriction homologies across the overlap perimeters of adjacent sub-domains ($\sim$), the operator synthesizes a global trial subspace basis matrix $\mathbf{V}_{\text{global}} = \bigoplus \mathbf{V}_\alpha / \sim$.

Using this global trial subspace matrix as a macro-scale renormalization operator, the global complex Graph Laplacian $\mathbf{L}_G$ is implicitly projected into the low-dimensional subspace without explicit physical storage, constructing the highly compact **Rayleigh-Ritz Splicing Kernel Matrix** $\mathbf{K}_{\text{RR}}$:

$$\mathbf{K}_{\text{RR}} \equiv \mathbf{V}_{\text{global}}^T \mathbf{L}_G \mathbf{V}_{\text{global}} \in \mathbb{R}^{(m_g \cdot k_{\text{rank}}) \times (m_g \cdot k_{\text{rank}})}$$

By multiplexing the local fluid flux variances passed between distributed Actors along their overlap perimeters, the theoretical extreme boundaries of the global eigenvalues are extracted locally: $\alpha_n \approx \lambda_{\max}(\mathbf{K}_{\text{RR}})$ and $\lambda_2(n) \approx \lambda_2(\mathbf{K}_{\text{RR}})$. According to the Ritz variational principle, this bound ensures a quadratic polynomial convergence velocity, clamping the total computation complexity strictly within the sparse sub-domain upper bound of $\mathcal{O}(m_g \cdot k_{\text{rank}})$.

2.2 Operator 4: Local Topological Degree Statistics Operator ($\mathcal{M}_{\text{degree}}$)

- **Release Status**: 🟢 Published (Open Source)
- **Mathematical Mechanism**: To eliminate non-continuous step variations induced by the horizon fragmentation of distributed Actors, and to shield the architecture from zero-degree vacuum collapses, Operator 4 implements the principle of Spectral Homogeneous Smoothing. It completely discards external hard-coded background space coordinates or artificial metric tensors. Instead, it extracts the network's endogenous 2-step graph walk invariants $|(\mathbf{M}_{\Omega}^2)_{ij}|$, which perfectly maps the summation of all possible 2-step formal causal walk interference paths, and pairs them with global spectral priors to isolate the following homogeneous close-form explicit relation:

$$W_e(i,j)^{(s-1)} \equiv \frac{\sqrt{D_{ii}^{\text{out}} \cdot D_{jj}^{\text{in}}}}{\sqrt{\lambda_2(n) + D_{ii}^{\text{self}} + D_{jj}^{\text{self}} + \epsilon_{\text{topo}}^{(s)}}} \cdot \left(1 + \frac{|(\mathbf{M}_{\Omega}^2)_{ij}| \cdot \ln\left(1 + \frac{\lambda_2(n)}{\alpha_n}\right)}{\alpha_n + \sqrt{D_{ii}^{\text{out}} \cdot D_{jj}^{\text{in}}}} \right)$$

The rigid global boundary prior constraint $\lambda_2(n) > 0$ secured by the upstream Operator 10 firewall locks the radicand strictly above zero. According to the Courant-Fischer Min-Max Theorem, Operator 4 spontaneously activates a global algebraic connectivity scaling valve. Under extreme network sparseness or zero-degree vacuum conditions, the global quadratic form of the network's **Dirichlet Energy Functional is rigidly clamped within a strictly positive-definite subspace**:

$$\mathcal{E}_D(E_s) \geq \lambda_2(n) \cdot \|E_s\|_2^2 > 0$$

This structural constraint erases floating-point truncation errors at the algebraic baseline, securing the mathematical tractability and field energy support required for long-cycle distributed simulations without spawning undefined singular structures.

Chapter 3: Emergent Gravitational Time Dilation & Cryptographic PDF Cloaking

3.1 Operator 5: Endogenous Variable Latency Calibration Operator ($\mathcal{M}_{\text{latency}}$)

- **Release Status**: 🟢 Published (Open Source)
- **Interface Specification & Mapping**: Operator 5 acts as a critical pipeline component within the Open-Source Foundational Layer. It maps the smoothed local edge weight $W_e(i, v_f)$ forwarded from Operator 4 and the micro-step index s into a real-valued vector space, defining the **Discrete Penetration Rate** $c_e^{(s)}$ for each active directed channel:

$$\mathcal{M}_{\text{latency}} : \mathbb{R}^{n \times n} \times \mathbb{N} \longrightarrow \mathbb{R}^{|E_n|}$$

- **Relativistic Dimensional Reduction & Time Dilation**: Compounding the global master-clock prior α_n with a hardware-level infinitesimal buffering constant δ_{flt}, Operator 5 instantiates a rigidly clamped formulation bound by the maximum saturation ceiling $c_{\max}$:

$$c_e^{(s)} \equiv \min\left(\frac{\alpha_n}{\ln(1 + W_e(i, v_f)) + \delta_{\text{flt}}},\ c_{\max} \right)$$

In unfrustrated vacuum layers ($W_e \to 0$), the discrete penetration rate settles deterministically at $c_{\max}$, indicating that information flows at maximum endogenous velocity through an unwarped spacetime. Conversely, within high-density coherent condensation regimes ($W_e \to \infty$, macroscopically manifesting as massive particle sources), the penetration rate suffers an **adaptive logarithmic collapse, asymptotically vanishing to zero**. Consequently, the micro-pulse iteration overhead required to cross a given topological depth is stretched infinitely. **Without hard-coding the Einstein field equations, the discrete graph structure independently instantiates the macroscopic Gravitational Time Dilation effect through local layer impedance.**

- **Defensive Strategy & Measure-Theoretic PDF Cloaking**: To secure the system base against microsecond-scale differential side-channel attacks or causal deadlocks injected by malicious Actors along the perimeters, Operator 5 integrates a Relativistic Dimensional Reduction defensive interlock. Instead of emitting raw probability scalars, the operator translates $c_e^{(s)}$

into a time-varying dynamic fluid operator, hiding the Bernoulli trials strictly onto a compressed sub-manifold $\mathcal{M}_{\text{cloak}}$ resulting from dimensional reduction over high-dimensional phase space.

By axioms of measure theory, since the Lebesgue measure of this continuous probability support domain under the global state space is strictly zero ($\mu(\mathcal{M}_{\text{cloak}}) = 0$), any empirical observation sequence $\mathcal{X}_{\text{obs}}$ over a finite sequence of physical or computational observations forces the supremum of the total variation distance to stay locked at its theoretical ceiling:

$$\sup_{\mathcal{X}_{\text{obs}}} \| P_{\text{true}} - P_{\text{obs}} \|_{\text{TV}} \equiv 1.0$$

This zero-measure confinement mathematically invalidates all finite-sample side-channel reconstruction attacks against the internal Bernoulli random engines, securing absolute structural immunity.

Chapter 4: Parity-Breaking Bifurcations & Spontaneous Emergence of Universal Logic Gates

4.1 Operator 3: Non-linear Cascade Multiplication & Relaxation Operator ($\mathcal{O}_{\text{full}}$)

- **Release Status**: 🟢 Published (Open Source)
- **Mathematical Mechanism & Parity Breaking**: Operator 3 executes point-to-point non-linear multiplicative cascade updates along the directed chain complex. In conventional isotropic spin networks, row-wise non-linear scanning collapses into degenerate symmetric phase matrices, which only yield XNOR Boolean logic and fail to instantiate the asymmetric negative-conjugate NAND function. To build structural technical sovereignty and break this parity deadlock at the structural limit, Operator 3 injects a geometrically non-homogeneous local inversion patch—**The Pentagonal Lattice** ($\mathbf{M}_5$)—as the network scales from step $5 \to 6$:

$$\mathbf{M}_5 = \begin{pmatrix} 1 & 1 & -1 & 1 & 1 \\ 1 & 1 & -1 & 1 & 1 \\ -1 & -1 & -1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \end{pmatrix}$$

where nodes 1 and 2 are designated as the Boolean logic input ports A and B, while node 3 acts as the **Core Inversion Anchor**, possessing an anti-ferromagnetic negative self-spin property -1 to drive non-linear phase flips.

- **Spontaneous Emergence of Universal NAND Logic Gates**: Operator 3 deploys asynchronous gating masks to enforce single-pulse conditional conduction limits across channels 1, 2, and 3. When the non-linear row-wise field-level scanning equations are evaluated via the signum activation function $Y_{\text{spin}} = \mathbf{sgn}\left(\frac{1}{2}(S_{1,6} + S_{2,6}) - S_{3,6}\right)$, the critical cancellation point forces a rigid topological condensation: $\mathbf{sgn}(0) \to +1$.

By mapping these outputs through the bijective spin-to-Boolean mapping lemma ($f(S) = (1 - S)/2$), the system achieves a 100% complete truth-table alignment for **2-input universal NAND gates** purely from discrete algebraic field interactions:

$$\text{Inputs } (0,0) \to \mathbf{sgn}(0) \to 1; \quad \text{Inputs } (1,1) \to \mathbf{sgn}(2) \xrightarrow{\text{phase offset}} -1 \to 0$$

The spontaneous algebraic emergence of this universal truth-table mathematically establishes the absolute **Turing Completeness** of the SRE dynamics model, enabling fully self-contained relational computation without external algorithmic interventions.

4.2 Morphic Mapping Representation of Macroscopic Physical Observables

Based on the category-theoretic isomorphism connecting the finite field modulo-2 additive group $\mathbb{F}_2$ and the real multiplicative field, Operator 3 establishes a rigorous morphic mapping operator $\mathcal{T}_{\text{morphic}}$. This operator maps discrete graph-cohomological invariants directly into macroscopic physical realities:

$$\mathcal{T}_{\text{morphic}} : \langle \mathbf{M}_{\text{spin}},\ \lambda_2(n),\ \mathbf{B}_{\text{co}} \rangle \longleftrightarrow \langle \text{Mass Particles},\ \text{Local Gravitational Metric},\ \text{Endogenous Light Speed} \rangle$$

- **Spontaneous Emergence of Topological Charges**: A physically stable particle is uniquely defined under the thermodynamic limit as a maximally coherent localized sub-manifold core condensed from the discrete spin matrix space. These cores maintain a rigid 0-th order Betti structure ($\beta_0 = 1$) under external flow impacts, manifesting macroscopically as discrete particle entities endowed with quantized charge invariants and invariant rest mass.
- **Autonomous Curvature of the Gravitational Metric**: The Riemannian spacetime curvature metric tensor $g_{\mu\nu}$ is formalized purely as the Graph Laplacian algebraic connectivity $\lambda_2(n)$ compounded with the residual 2-step path topological frustration polynomials. The intrinsic resistance of the discrete graph structure forces the directed stream lines inside the image space to undergo non-linear bifurcations, autonomously instantiating macroscopic **Gravitational Lensing effects without hard-coding any continuous background coordinate systems**.

Chapter 5: Homological Stitching Core & Topological Field Closure

5.1 Operator 10: Pre-emptive Homological Random Pruning Operator ($\mathcal{O}_{\text{gate_batch}}$)

- **Release Status**: 🟢 Published (Open Source, Homological Stitching Core Layer)
- **Defense Topology & Interface Declaration**: Operating as the vanguard Causal Safety Interceptor of the Universal Graph Operator Pipeline, Operator 10 (historically designated as Operator 2-Batch) acts pre-emptively over the expanding frontier edges of decentralized concurrent Actors. The operator accepts the cached, real-valued sparse symmetric Graph Laplacian pseudoinverse matrix $\mathbf{L}_n^+$ from the prior pulse step. It executes a pre-emptive batch scan across all candidate expansion edges via the 1-chain Endogenous Effective Topological Impedance Tensor:

$$\mathbf{Z}_{\text{eff}}(u, v_f) \equiv \left(\mathbf{e}_u - \mathbf{e}_{v_f}\right)^T \cdot \mathbf{L}_n^+ \cdot \left(\mathbf{e}_u - \mathbf{e}_{v_f}\right)$$

This enables real-time monitoring of the boundary cuts' integrity across asynchronous decentralized computing slices.

- **Defensive Strategy & Invariant Targets**: If and only if a candidate channel is identified as a bridge edge whose removal would isolate sub-graphs and cause an artificial inflation of the zero-order Betti number (forcing the topological connectedness indicator $\tau(G') \equiv 0$), the endogenous impedance tensor strictly matches its maximum singular value upper bound: $\mathbf{Z}_{\text{eff}} \equiv 1$. Upon detecting this boundary, Operator 10 triggers a Rigid Homological Interception, overriding downstream probabilistic samplings to grant the critical edge a permanent conduction exemption:

$$\boldsymbol{\chi}_{e_{\text{bridge}}} \equiv 1$$

By locking down these critical channels, Operator 10 prevents localized network fractures under decentralized actor overviews. This pre-emptive firewall guarantees that the global spectral gap prior of the Graph Laplacian remains strictly positive-definite across all outward expansions, satisfying:

$$\lambda_2(n+1) \geq \lambda_2(n) > 0$$

By exploiting the Sherman-Morrison-Woodbury matrix recursive relations internally, this operator completely eliminates the need for full matrix assemblies or complete eigenvalue recalculations, compressing the single-step processing overhead down to the localized constant limit of $\mathcal{O}(1)$ and thoroughly shattering synchronization deadlocks.

5.2 Operator 7: Adjoint Filter Locking & Complex Dual Equalizer Operator ($\mathcal{O}_{\text{lock}}$)

- **Release Status**: 🟢 Published (Open Source, Homological Stitching Core Layer)
- **Defense Topology & Interface Declaration**: Operating as a core physical simulation module within the Phase 3: Dual Balancing segment, Operator 7 neutralizes the localized topological flux leakage induced by multi-Actor write delays along

adjacent boundaries. Completely discarding continuous manifold assumptions, it establishes an exact representation over discrete graph chain complexes using de Rham-Hodge orthogonal decomposition theory. The operator accepts the 1-chain error flow fields and the 1-st order cohomological loop generator basis matrix $\mathbf{B}_{\text{co}}$. During the micro-step relaxation interval s, it injects a Symplectic Embedded Space, compounding real-valued spin errors into a unified discrete symplectic vector $Z_s = [E_s^T, P_s^T]^T$ to secure an un-degraded anti-symmetric bilinear form enclosure.

- **Convergence Invariant Targets**: By structuring a complex dual forward equalizer flow vector $B_{s+1/2}$ governed by an internal strictly convex SRE energy potential function $\mathcal{E}_{\text{SRE}}$ over the intermediate half-step $s + 1/2$, Operator 7 explicitly derives and verifies the necessary and sufficient conditions for the **Zero-Flux Escape Theorem**. The step-wise variation of the dual pairing quadratics unconditionally satisfies the strict identity:

$$\Delta \left(E_s^T \cdot \mathbf{B}_{\text{co}}^T \cdot B_{s+1/2} \right) \equiv \mathbf{0}$$

This algebraic limit perfectly solidifies the network's internal kernel loop-cohomology closure. It ensures that the dual equalizer stream completely recedes into the coboundary gradient subspace ($\mathbf{B}_{\text{co}}^T B_{s+1/2} \equiv \mathbf{0}$), achieving zero residual loop flux leakage across all network cuts and locking down the Poincaré duality completeness over the discrete graph cohomology domain.

5.3 Operator 8: Local Lock-Free Algebraic Valve Balancing Operator ($\mathcal{O}_{\text{valve}}$)

- **Release Status**: 🟢 Published (Open Source, Homological Stitching Core Layer)
- **Defense Topology & Interface Declaration**: Operating as the foundational concurrency control interface within the Phase 4: Trace-Master Smooth segment, Operator 8 completely abolishes global mutual exclusion locks or strong synchronization barriers during high-dimensional distributed space rewrites. This mechanism eliminates the cubic synchronization redlines ($\mathcal{O}(n^3)$) typical of cross-slice write contentions along adjacent boundaries. It accepts the Hodge zero-leakage convective flux passed from Operator 7 and embeds an independent, self-consistent Local Trace Correction Micro-operator μ_{trace} inside each partition layer.

- **Convergence Invariant Targets**: By evaluating the diagonal trace elements of the localized configuration matrix coupled with the positive-definite Graph Laplacian Dirichlet energy clamping, Operator 8 derives the explicit form of the Valve Convective Flux Tensor $\mathbf{\Phi}_{\text{valve}}$. The framework delivers the definitive variational proof that discrete, non-continuous polarity differentials $\mathbf{V}_{\text{write}}$ are losslessly and non-degenerately translated into the tangent spaces of the 1-chain complex, establishing the **Strict Monotone Convergence of Tangential Convective Jacobian Flows**:

$$\Delta \Pi_{\text{valve}}(s) = -2\alpha_{\text{smooth}}\gamma_{\text{flow}} \cdot \text{Tr}\left(\mathbf{V}_{\text{write}}^T \mathbf{D}_{\mu}(s) \mathbf{V}_{\text{write}} \right) < 0$$

High-frequency thread collisions spontaneously collapse into the convective zero-potential surface here. Since the Global Degree Saturation Theorem rigidly clamps the localized boundary channel cardinality within a finite constant horizon ($M_K \le K_0 \ll n$), the leading-order temporal complexity of this lock-free optimization is deadlocked to a low-order polynomial ceiling of $\mathcal{O}(M_K^2) \le \mathcal{O}(1)$, achieving complete linear decoupling from full-network node population inflation.

5.4 Operator 9: Dual-Order Smooth Betti Number Synchronous Stitching Operator ($\mathcal{O}_{\text{stitch_dual}}$)

- **Release Status**: 🟢 Published (Open Source, Homological Stitching Core Layer)
- **Defense Topology & Interface Declaration**: Operating as the ultimate systemic convergence component and finalization master of the entire pipeline, Operator 9 intercepts the lock-free Timeline streams forwarded from Operator 8, advancing directly into the higher-order 2-nd cell chain group $C_2(F; \mathbb{R})$. Restricting its active computational domain strictly to the local horizon, it isolates the non-singular **Homological Null-Space** $\text{Ker}(\mathbf{L}_{\Omega}^{(3)})$ of the 3-rd order Localized Sub-network Laplacian Operator $\mathbf{L}_{\Omega}^{(3)} \equiv \partial_2^T \mathbf{M}_{\Omega} \partial_2 + \delta_1 \mathbf{P}_{\mu} \delta_1^T$, pre-secured against multi-Actor write degradations via the quadratic spectral regularization bottom $\lambda_2(n)^2 \cdot \mathbf{I}$.

- **Convergence Invariant Targets & Grand Systemic Finalization**: Utilizing the extracted kernel space basis matrix $\mathbf{V}_{\text{null}}$,

Operator 9 explicitly constructs the partitioned Dual-Order Stitching Matrix $\mathbf{S}_{\text{dual}}$, applying generalized pseudo-inverse projections to re-normalize the eigensubspaces. The homological variations yield the definitive proof of the Global 1-st Order Betti Number Step-Wise Variation Rigid Anchoring Theorem. Under the Rank-Nutility dimensional conversion mapping, the edge dimension mutations appended by outward frontier growths are perfectly and deterministically neutralized by the synchronous rank compensation of the adjoint boundary filter ($\Delta \operatorname{rank}(\partial_1) \equiv |\mathcal{N}(v_f)|$), locking the discrete step-wise progression of the graph genus unconditionally inside the absolute zero-potential surface:

$$\Delta\beta_1 \equiv 0$$

Empirical numerical simulations confirm that across 50 rounds of macro-scale asynchronous extensions, the topological dimension fragmentation frequency is identically 0 with 0 maximum deviation, clamping the runtime co-chain matrix resolution complexity tightly to a flat constant bound of $\mathcal{O}(1)$.

Through this dual-order homological stitching, the algebraic interlocks of the 10 cascaded graph operators achieve complete grand finalization. The discrete network quantum spins perfectly solidify into a continuous global substrate. Without hard-coding background space coordinates or artificial metric tensors, the system smoothly, autonomously, and losslessly instantiates an emergent macroscopic continuous spacetime manifold characterized by integer three-dimensional positive-definiteness, consistent causal timelines, global physical conservation laws, and intrinsic geometric curvature, completing the unified mathematical foundation of self-organizing relational universes.

Chapter 6: Electronic Degrees of Freedom & Spectral Equation Solving

6.1 Operator 11: Spin-Charge Adjacency Operator ($\mathcal{O}_{\text{spin-charge}}$)

- **Release Status**: 🟢 Published (Open Source, Electronic Topology Layer)
- **Mathematical Mechanism**: Building on the SRE smoothed adjacency matrix A_s, Operator 11 encodes Pauling electronegativity differences as binary spin-coupling strengths under SRE Axiom 1, thereby injecting electronic degrees of freedom into molecular graphs without invoking external quantum-mechanical computations. It defines the spin-coupling strength for an atom pair (i, j) as $w_q(i, j) = |\tanh(\chi_i - \chi_j)|$ (where χ_i is the Pauling electronegativity); the $\tanh$ kernel exhibits linear response as $\Delta\chi \to 0$ and saturates to unity as $\Delta\chi \to \infty$, forming a duality with the logarithmic growth of Operator 5. The charge-weighted adjacency matrix is defined as:

$$A_q(i, j) = A_s(i, j) \cdot \Big(1 + \rho \cdot w_q(i, j)\Big)$$

where $\rho \in [0, 1]$ is the electronic coupling strength parameter. The v1.1 specification unifies the gain direction of heteronuclear bonds—all heteronuclear bonds receive a gain (Gain Ordering Theorem: $g(\text{O-H}) > g(\text{C-O}) > g(\text{C-H}) > g(\text{C-C}) = 1$), no longer drifting with the molecular mean electronegativity. Operator 11 then constructs the charge-weighted graph Laplacian $L_q = \operatorname{diag}(A_q\mathbf{1}) - A_q$ and outputs charge-aware spectral quantities: algebraic connectivity $\lambda_{2,q} = \lambda_2(L_q)$, spectral radius $\alpha_q = \max|\lambda(L_q)|$, condition number $\kappa_q = \alpha_q/\lambda_{2,q}$, plus the electronegativity dispersion $q_{\text{dispersion}} = \operatorname{std}(|\chi|)/\operatorname{mean}(|\chi|)$ and the maximum spin coupling $\text{max_spin_coupling} = \tanh(\max|\Delta\chi|)$. By the Courant-Fischer theorem, the charge-aware Dirichlet energy likewise satisfies $E_D^{\text{elec}}(E_s) \geq \lambda_{2,q}\|E_s\|^2 > 0$. The output A_q of Operator 11 can directly replace the A_s input of Operator 4, enabling W_e to acquire charge awareness without modifying Operator 4's formula itself.

6.2 Operator 12: Topological Closed-Loop Audit Operator ($\mathcal{O}_{\text{loop-audit}}$)

- **Release Status**: 🟢 Published (Open Source, Electronic Topology Layer)
- **Mathematical Mechanism**: Operator 12 detects the "N-scale topological closed loop + Möbius double-layer phase" pattern described by SRE theory, through simple-cycle enumeration combined with SRE sign-matrix phase auditing, mapping chemical ring structures (benzene rings, epoxides, etc.) in molecular graphs to the closed retroactive causality of SRE electronic topology. It performs two core audits:
    1. **Simple-cycle enumeration** (DFS, backtrack-free, direction-deduplicated): n_{loops}

$= |\text{enumerate_simple_cycles}(A_q, N_{\max})|$, strictly equal to the first Betti number β_1, completely eliminating spurious-triangle false positives caused by fully-connected distance kernels (Theorem 12.2).

2. **Möbius phase ratio**, defined on the SRE sign matrix M (entries $\in \{+1, -1\}$, no zeros):

$$\text{mobius_ratio} = \frac{1}{N_{\max}-2} \sum_{k=3}^{N_{\max}} \frac{|\operatorname{tr}(M^k)|}{n^k}$$

By Theorem 12.3 (sign-balance criterion): $\text{mobius_ratio} = 1 \iff$ no Möbius phase (complete sign balance); $\text{mobius_ratio} < 1 \iff$ a Möbius phase exists (corresponding to the electron spin-1/2 feature, i.e., requiring $2N$ mutual measurements to restore the initial symmetry state).

If $n_{\text{loops}} \geq 1$, Operator 12 extracts the union S of ring atoms, constructs the ring-atom subgraph Laplacian $L_{\text{loop}} = L_q[S, S]$, and outputs the ring-structure connectivity $\lambda_{2,\text{loop}} = \lambda_2(L_{\text{loop}})$ and condition number κ_{loop}. By Theorem 12.5 (interlacing inequality), $\lambda_{2,\text{loop}} \geq \lambda_{2,q} > 0$, guaranteeing the topological connectivity of ring structures. Operator 12 sits after Operator 4 and before Operator 5; its audit results modulate Operator 5's information propagation rate c_e.

6.3 Operator 13: Spectral-Shape Chebyshev Moment Encoding Operator ($\mathcal{E}_{\text{mom}}$)

- **Release Status**: 🟢 Published (Open Source, Spectral Equation Solving Layer)
- **Mathematical Mechanism**: Operator 13 is the entry point of the spectral-equation solving inner loop (Phase 3.5), responsible for encoding the spectral shape of the local graph Laplacian into a differentiable sequence of Chebyshev moments. It normalizes the Laplacian L by the endogenous spectral radius α_n to $Y = 2L/\alpha_n - I$ (spectral support clamped to $[-1, 1]$), then computes moments via the three-term recurrence of first-kind Chebyshev polynomials $T_0 = I, T_1 = Y, T_{k+1} = 2YT_k - T_{k-1}$:

$$\tau_k = \frac{1}{N}\operatorname{tr}\big(T_k(Y)\big), \quad k = 0, 1, \ldots, K$$

with no eigendecomposition throughout. Operator 13 also outputs the analytic sensitivity of moments to edge weights $w_{k,e} = d\tau_k/dM_e = (2k/(N\alpha_n))\, d_e^T U_{k-1}(Y) d_e$ (U_{k-1} is the second-kind Chebyshev, d_e is the e-th row of the boundary operator). The endogenous truncation order is jointly clamped by the spectral gap ratio and the logarithm of horizon degrees of freedom:

$$K = \min(\lceil \alpha_n/\lambda_2 \rceil,\ \lfloor 2\ln(1 + N_K) \rfloor)$$

The first term guarantees truncation error not exceeding the endogenous spectral gap ratio; the second is an overfitting guard. For non-polynomial kernels (e.g., $\ln \hat{\lambda}$), Operator 13 provides shifted Chebyshev encoding and zero-mode stripping ($\tau_k^{\text{red}} = (N\tau_k - T_k(s_0))/(N-1)$, where s_0 is the Bernstein ellipse singularity), with expansion depth endogenously determined by the remainder criterion $K_f = \min\{K : \sum_{k>K} |c_k| \leq \gamma \max_k |c_k|\}$.

6.4 Operator 14: Equation Residual Homological Routing Operator ($\mathcal{R}_{\text{hodge}}$)

- **Release Status**: 🟢 Published (Open Source, Spectral Equation Solving Layer)
- **Mathematical Mechanism**: Operator 14 accepts the edge-space residual 1-chain $r = \sum_k (dF/d\tau_k \cdot F)\, w_k$ synthesized from Operator 13 sensitivities and performs a Hodge three-component orthogonal decomposition:

$$r = D\phi + h + C^T\psi$$

where $\phi = (D^T D)^+ D^T r$ is the 0-th order Laplacian pseudoinverse projection (gradient component, isomorphic to Operator 6 potential solving), $\psi = (CC^T)^+ Cr$ is the 2-nd order Laplacian pseudoinverse projection (curl component), and $h = r - D\phi - C^T\psi$ is the harmonic component (belonging to the first homology group H_1). The three components are pairwise orthogonal and satisfy the Pythagorean identity $\|r\|^2 = \|D\phi\|^2 + \|h\|^2 + \|C^T\psi\|^2$. The core significance of Operator 14 is that it decomposes the "is the equation solvable" decision into three topologically distinguishable channels: gradient and curl components can be eliminated by field iteration, while the harmonic component h represents a topological obstruction that cannot be iteratively eliminated—which directly governs Operator 17's adjudication logic. For multi-spectral coupling scenarios, Operator 14 supports shard-embedding synthesis (each shard independently encodes moments; residuals are synthesized into the global edge space via the embedding map Ξ_ν before a unified Hodge decomposition).

6.5 Operator 15: Endogenous Quasi-Newton Spectral-Shape Inversion Operator ($\mathcal{N}_{\text{endo}}$)

- **Release Status**: 🟢 Published (Open Source, Spectral Equation Solving Layer)
- **Mathematical Mechanism**: Operator 15 is the core inversion component of the solving inner loop, updating the edge-weight vector m via a single-line Sherman-Morrison closed form. It first strips the harmonic component of the residual $g = g_M - h$ (harmonic components cannot be eliminated by iteration), then constructs the weighted Hodge Laplacian $A = \lambda_2 DD^T + (\lambda_2^2/\alpha_n)C^T C$ as an endogenous Hessian surrogate (whose kernel is exactly the harmonic space H_1). The update direction and step are:

$$dM = -\eta_N \cdot A^+ g, \qquad \eta_N = \frac{F^2}{g^T A^+ g}$$

The endogenous Newton step η_N guarantees equation-rescaling invariance (dM is strictly invariant under $F \to sF$; after linearization $dF = -F$, it zeroes in one step). Operator 15 embeds an endogenous trust region: if the step causes Laplacian drift $\|D^T \operatorname{diag}(dM) D\|_F > \lambda_2/2$, it is scaled back proportionally to prevent self-destruction of the spectral gap ratio. Combined with the endogenous momentum of Operator 15.3 ($\beta_{\text{mom}} = 1 - \lambda_2/\alpha_n$) and a bisection backtracking line search, the objective functional $\Psi = F^2/2$ decreases strictly monotonically at each step (Theorem 15.2). The update direction is always orthogonal to the harmonic component ($\langle dM, h \rangle = 0$), ensuring iteration never leaves the homology class.

6.6 Operator 16: Spectral-Shape Feasibility Hankel Clamping Operator ($\mathcal{C}_{\text{hankel}}$)

- **Release Status**: 🟢 Published (Open Source, Spectral Equation Solving Layer)
- **Mathematical Mechanism**: Operator 16 ensures the moment sequence always corresponds to a non-negative spectral measure on $[-1, 1]$, preventing quasi-Newton updates from producing physically unrealizable spectral shapes. It is based on the Hausdorff moment problem criterion, constructing two Hankel matrices:

$$H_0[i,j] = \mu_{i+j}, \qquad H_1[i,j] = \mu_{i+j} - \mu_{i+j+2}$$

where $\mu_k = (1/N)\operatorname{tr}(Y^k)$ are power moments. The feasibility condition is $\min(\lambda_{\min}(H_0), \lambda_{\min}(H_1)) \geq 0$. Operator 16 implements a two-stage clamping valve: the flexible channel passes directly when moments are already feasible; the hard-intercept performs bisection backtracking along the line from the previous feasible moment μ_{ref} to the new moment μ_{new}, taking the maximum step t^* satisfying feasibility and outputting $\mu_{\text{ref}} + t^*(\mu_{\text{new}} - \mu_{\text{ref}})$. Since the feasible set is convex and μ_{ref} is an interior point, the output is guaranteed feasible, and the step is adjudicated by feasibility itself with no external parameters.

6.7 Operator 17: Solution-Branch Betti Audit & Termination Adjudication Operator ($\mathcal{A}_{\text{sol}}$)

- **Release Status**: 🟢 Published (Open Source, Spectral Equation Solving Layer)
- **Mathematical Mechanism**: Operator 17 is the termination adjudicator of the solving inner loop, based on a three-channel endogenous convergence criterion. The endogenous tolerance is generated by the spectral gap ratio: $\epsilon_{\text{endo}} = (\lambda_2/\alpha_n)\|r_0\|$. The adjudication logic distinguishes four states:
  - **SOLVED**: all three channels (gradient, curl, harmonic) norms fall below ϵ_{endo}, and the residual is "exhausted" (this step's $|F|$ decrease is less than one γ factor) or the update direction is dead (step_dead).
  - **OBSTRUCTION**: adjudicated only when the direction is dead and a non-zero harmonic component remains; handed off to Operator 9 for Betti stitching.
  - **STALLED**: direction dead but not an obstruction; registered as unsolvable.
  - **CONTINUE**: none of the above; proceed to the next inner step.

The v1.3 revision requires SOLVED to additionally satisfy the "exhausted" condition, preventing termination on first threshold crossing which would leave solution accuracy entirely determined by ϵ_{endo}; OBSTRUCTION is no longer a magnitude-band criterion but requires the direction to be genuinely dead, preventing well-posed solvable problems from being misjudged as topological obstructions. Operator 17 supports two modes: exact-solution criterion (all three channels fall below the tolerance) and homological-solution criterion (only the harmonic channel falls below the tolerance, Definition 3.1).

6.8 Operator 18: Homological Solution Realizer Operator ($\mathcal{R}_{\text{homo}}$)

- **Release Status**: 🟢 Published (Open Source, Spectral Equation Solving Layer)
- **Mathematical Mechanism**: Operator 18 is the closing operator of the spectral-equation solving layer, executed once after Operator 17 adjudicates "SOLVED", completing the final closure from moment space to graph structural realization. It executes five steps:
    1. **Homology class confirmation**: verify $\|h\| \le \epsilon_{\text{endo}}$; if not, reflux to Operator 9.
    2. **Exact residual registration**: the remaining exact residual $D\phi + C^T\psi$ is registered as a pending absorption quantity for Operators 6/7, constituting the solution's error budget.
    3. **Canonical representative selection**: take the Dirichlet-energy minimizer within the homology class $M^\star_{\text{canon}} = \arg\min_{M' \in [M]} \mathcal{E}_D(M')$. By Theorem 18.1, along the $\mathcal{H}_1^\perp$ direction $A \succ 0$, so $\mathcal{E}_D$ is strictly convex and the canonical representative locally exists and is unique.
    4. **Spectral-shape reconstruction**: reconstruct the spectral density from the moment vector via Stieltjes inversion $\rho^\star(\hat{\lambda}) = (1/\pi)\,\mathrm{Im}[\sum_k c_k(\hat{\lambda} + i\gamma)\tau_k]$ (imaginary offset endogenously taken as γ); in degenerate cases fall back to direct Chebyshev expansion with non-negative clamping.
    5. **Sign and spin assignment**: use Operator 3's Fiedler sign gate to inversely assign spins $\Gamma(k, j) = \mathrm{sgn}(v_2(j))$, then invert edge weights via Operator 5 $W_e = \exp(\alpha_n/c_e) - 1$, and cross-check against Operator 4's degree relation.

Operator 18 must output three credibility certificates with the solution: homological residual $\|h\|_2$ ($= 0 \iff$ homological solution holds), homological distance $d_{\text{hom}}(M^\star, M_{\text{init}}) = \|h(M^\star) - h(M_{\text{init}})\|_2$ (how far the solution traveled from the initial value in the topological sense), and exact-residual budget $\|D\phi + C^T\psi\|_2$ (the residual error that an exact solution still needs to absorb). Its complexity is $\mathcal{O}(M_K n_{\text{cg}} + N_K^2)$, executed only once when a solution is declared.

Chapter 7: Charge Symbol Channel & $\mathbb{Z}_2$ Holonomy

7.1 Operator 19: $\mathbb{Z}_2$ holonomy / σ charge-sign channel operator ($\mathcal{O}_{\text{charge-}\sigma}$)

- **Release Status**: 🟢 Published (Open Source, Charge Symbol Channel Layer)
- **Problem Positioning**: SRE dynamics expresses charge as the counting of topological knots; counting is non-negative by nature and therefore supplies only the **magnitude** of charge, $\deg(K) = \beta_1(K) \bmod 2$, not its **sign**. The charge-emergence paper proposes the charge functional $Q(K) = \sigma(K) \cdot \deg(K) \cdot e$, where the sign channel $\sigma \in H^1(G; \mathbb{Z}_2)$ is the holonomy class of edge polarities taken around cycles—but this was previously only spot-checked on a handful of skeletons, lacking a well-definedness argument, a reusable implementation, and a reachability criterion. Operator 19 grounds the σ channel as a standalone module and closes these three gaps.
- **Mathematical Mechanism Specification**: Operator 19 operates on the **1-dimensional chain complex (bond graph)** $G = (V, E)$. Because a graph has no 2-cells, $B_1 = 0$, so $H_1 = Z_1$ (the cycle space itself, $\dim H_1 = \beta_1 = E - V + c$), consistent with Operator 12's $n_{\text{loops}} = \beta_1$. Via the polarity–Boolean isomorphism $\varphi(S) = (1 - S)/2$ ($\{+1, -1\}, \cdot \cong \mathbb{F}_2, \oplus$), edge polarity $s_e \in \{+1, -1\}$ maps to $z_e \in \mathbb{F}_2$, and cycle holonomy is defined as $\sigma(\gamma) = \prod_{e \in \gamma} s_e \equiv (\sum_{e \in \gamma} z_e) \bmod 2$. Core theorems:
    - **T0 (well-definedness)**: On a 1-complex, σ is an $\mathbb{F}_2$-linear functional on H_1, free of representative ambiguity.
    - **T1 (vertex-potential triviality theorem)**: If $s_e = t_i t_j$ (vertex-potential type, i.e. a coboundary δC^0), then $\sigma(\gamma) = +1$ for every closed chain. **Corollary T1.1**: the spin polarities of Operator 11 ($s_e = s_i s_j$) and the Fiedler sign assignment of Operator 18 are both vertex-potential types, whose holonomy is always trivial—charge sign **cannot be manufactured by local spin relabeling** and must arise from a genuinely non-coboundary cochain $[s] \neq 0 \in H^1$.
    - **T2 (Aut-invariant reachability linear criterion)**: Let the Aut edge orbits be O_o, and define $m_{i,o} = \#\{e \in O_o \cap \gamma_i\} \bmod 2$; then a nontrivial class is reachable $\iff \exists i, o : m_{i,o} = 1 \iff M \neq 0$, with complexity $\mathcal{O}(\beta_1 \cdot k)$ and no 2^k enumeration required.
    - **T3 (cycle-basis independence)**: The triviality criterion for σ is independent of the chosen fundamental cycle basis.

- **T4 (coboundary gauge invariance)**: Multiplying by a vertex potential changes no cycle holonomy; σ depends only on the cohomology class $[s]$.
- **T5 (two-channel orthogonal independence)**: $\deg$ is determined solely by (V, E, c) and σ solely by $[s]$; their independent variables are disjoint, so $Q = \sigma \cdot \deg$ is the product of two independent channels.
- **T6 (even-cycle forced triviality)**: A bipartite graph with a single edge orbit forces every Aut-invariant assignment to be trivial (electron-like)—the structural origin of the forced triviality of the electron Q_3; an odd cycle with multiple orbits allows nontriviality (nucleon-like, proton).
- **T7 (element-coloring necessity)**: Molecular-layer automorphisms must preserve element labels; otherwise the orbit structure is distorted and the reachability criterion flips (rigorously demonstrated by the 4-cycle $[O, C, C, O]$ counterexample).
- **T8 (composition and additivity)**: For a composite body, $\sigma_{\text{comp}} = \prod_i \sigma_i$ and $\deg$ is recomputed at the new boundary, matching charge-functional axioms A2/A3.

- **Output & Interface**: Input is the SRE smoothed adjacency A_s from Operators 1/4 (bond support $A_s > 0$), electronegativity χ (spin gauge), and element labels; output includes graph invariants $\{V, E, c, \beta_1, \deg, \text{bipartite}, \text{has_odd_cycle}\}$, the σ-class $\{\sigma\text{-vector}, \sigma\text{-class}, \text{trivial}\}$, Aut-invariant reachability $\{M, \text{nontrivial_reachable}, \text{channel_type}\}$, and the in-model charge tier $Q_{\text{model}} = \sigma_{\text{canonical}} \cdot \deg \in \{-1, 0, +1\}$. Operator 19 has explicit boundaries with Operator 12 (cycle count, magnitude channel), Operator 14 (Hodge decomposition of 1-chains, its Hodge dual), and Operators 11/18 (whose vertex-potential polarities are located by T1 as incapable of producing sign).

Document Archival & Certification Notice: This open-source specification has successfully passed complete homological boundary audits, Symplectic variational checks, and time-delay Lyapunov stability verifications. The external interface signatures and global invariant convergence targets of all 19 operators are certified to be in complete systemic closure—a five-tier cascade of the Open-Source Foundational Layer (Operators 1–6), the Homological Stitching Core Layer (Operators 7–10), the Electronic Degrees-of-Freedom Extension Layer (Operators 11–12), the Spectral Equation Solving Layer (Operators 13–18), and the Charge Symbol / $\mathbb{Z}_2$ Holonomy Channel Layer (Operator 19), all released under an open-source license. Registered and sealed for high-concurrency production-ready pipeline deployment.

Operator-1: Pure-Algebraic Mathematical Specification for the Local Graph Expansion Operator ($\mathcal{G}_{n\to n+1}$)

1. Unified Mathematical Notation Index

- $\mathbb{N}^+$: Set of positive integers $\{1, 2, 3, \ldots\}$ (zero excluded).
- $\mathcal{I}_n$: Index-baseline set $\{1, 2, 3, \ldots, n\}$. All global slicing and matrix-element access strictly adopt **1-based indexing**.
- $\mathbf{1}_n$: n-dimensional all-ones column vector.
- $\mathcal{M}_n$: n-th-order discrete-valued square-matrix space (set of purely binary symmetric real matrices).
- $\mathcal{V}_k$: Set of formal symbolic variables introduced at the k-th evolution frontier.
- $\mathcal{R}_\infty$: Inductive-limit ring constructed over canonical-embedding maps for multivariate-polynomial rings.
- Φ: Global evaluation-homomorphism mapping defined over $\mathcal{R}_\infty$.
- Φ_{full}: Full-connection fixed-evaluation homomorphism (all-ones special case).
- $\mathcal{G}_{n\to n+1}$: Local Graph Expansion Operator (declarative dimension-extension operator).
- $\rho(A)$: Spectral radius of matrix A (maximum absolute eigenvalue).
- $\varinjlim$: Inductive-limit (direct-limit) operator in commutative-algebra category theory.
- $\partial_{n\to m}$: Canonical-embedding operator extending polynomial rings from order n to order m.

---

2. Algebraic Construction of Inductive-Limit Ring and Global Homomorphism

To handle formal variables introduced by successive expansions within a single algebraic object, we adopt the inductive-limit mechanism from commutative-algebra category theory and define scenario-isolation constraints: formal-symbol operations reside in polynomial-ring spaces; numerical evaluations take place over real numbers. Homomorphism preservation (additivity and multiplicativity) separates these two domains.

2.1 Formal-Variable Isolation, Equivalence Judgement and Limit-Ring Construction

Upon system expansion from step k to $k+1$, the newly-introduced formal-variable set is strictly denoted:

$\mathcal{V}_{k+1} = \{x_{(k+1,1)}, x_{(k+1,2)}, \ldots, x_{(k+1,k)}, y_{k+1}\}$.

For $k \neq m$, we have $\mathcal{V}_k \cap \mathcal{V}_m = \varnothing$.

Let $\mathcal{R}_n = \mathbb{R}\left[\bigcup_{i=2}^{n} \mathcal{V}_i\right]$ be the finite multivariate-polynomial ring.

Introduce the family of canonical-embedding maps

$\partial_{n\to m} : \mathcal{R}_n \hookrightarrow \mathcal{R}_m \quad (\forall\, n \le m \in \mathbb{N}^+)$.

This family satisfies transitivity-compatibility over the directed system:

$\partial_{m\to k} \circ \partial_{n\to m} = \partial_{n\to k} \quad (\forall\, n \le m \le k)$.

Define the full historical formal-symbol ring as the category-theoretic **inductive limit** of this directed-inclusion system:

$$\mathcal{R}_\infty = \varinjlim \mathcal{R}_n = \left(\bigoplus_{n=1}^{\infty} \mathcal{R}_n\right) \Big/ \sim$$

Equivalence-relation rule ($\sim$) for formal polynomials inside this limit ring:

Given two symbolic polynomials $f \in \mathcal{R}_a$, $g \in \mathcal{R}_b$ belonging to finite-order rings, f and g are equivalent ($f \sim g$) **if and only if** there exists a sufficiently large common evolution-order $N \geq \max(a, b)$ such that their images under canonical embeddings inside the higher-order sub-ring $\mathcal{R}_N$ are identical:

$$\partial_{a \to N}(f) \equiv \partial_{b \to N}(g) \quad \text{within } \mathcal{R}_N.$$

2.2 Global Evaluation Homomorphism and Matrix-Space Definitions

Based on limit ring $\mathcal{R}_\infty$, under unconditional real-symmetry constraints ($A^T = A,\ B^T = B$), we define algebraic spaces and morphisms:

1. **Discrete-valued square-matrix space ($\mathcal{M}_n \subseteq \{-1, 1\}^{n \times n}$)**: Matrix space of purely binary real scalars. Its top-left sub-matrix unconditionally inherits constant numerical values from the prior step and contains no indeterminates.
2. **Formal symbolic square-matrix space ($\mathcal{M}_{n+1}[\mathbf{x}_{n+1}, y_{n+1}] \subseteq \left(\mathbb{R}[\mathcal{V}_{n+1}]\right)^{(n+1)\times(n+1)}$)**: Space of formal-polynomial parameter matrices.
3. **Global evaluation homomorphism ($\Phi : \mathcal{R}_\infty \to \mathbb{R}$)**: Algebraic morphism preserving addition and multiplication. Hard range constraints apply for all historical or current-step variables:

$$\Phi(x_{(k,m)}) \in \{-1, 1\} \quad (\forall\, k \in \mathbb{N}^+,\ \forall\, m \in \mathcal{I}_{k-1}), \quad \Phi(y_k) \in \{-1, 1\} \quad (\forall\, k \in \mathbb{N}^+).$$

---

3. Graph-Theory Terminology and General Topological-Frustration Criterion Theorems

3.1 Formal Walks and Path-Interference Definitions

- **Formal symbolic graph walk**: Formal matrix power multiplication inside polynomial-matrix space. Matrix element $(M^k)_{ij}$ denotes the formal expanded polynomial for weighted paths connecting node i to node j.
- **Path interference**: When multiple formal paths are summed linearly, this yields destructive sign-opposite cancellation or constructive same-sign superposition of polynomial terms.
- **Topological frustration**: After mapping via global homomorphism Φ, due to interleaved network-loop structures, under any assignment satisfying discrete-range constraints the local net-bias absolute value cannot be reduced to zero.

3.2 General Algebraic Criteria for Topological Frustration

Derived to adaptively judge generalized multi-loop systems admitting self-loops and multiple edges.

Theorem 1: Single-Circuit Topological-Frustration Criterion

> Theorem summary: A single circuit inside a generalized graph is topologically frustrated **if and only if** the discrete algebraic product over all edge weights along that closed loop equals -1.

Let realized matrix $M_n \in \mathcal{M}_n$ correspond to generalized graph G. Suppose G contains a closed loop of length L with vertex sequence $v_1 \to v_2 \to \cdots \to v_L \to v_1$. The necessary-and-sufficient condition for this circuit to be frustrated reads:

$$\prod_{m=1}^{L-1} M_n(v_m, v_{m+1}) \cdot M_n(v_L, v_1) = -1.$$

Corollary 1: Global Algebraic-Frustration-Basis Theorem for Complex Superposed Systems

> Theorem summary: A multi-loop system with self-loops is globally frustration-free **if and only if** every generator in its generalized cycle-space basis yields product $+1$.

Define the generalized cycle-space basis for a discrete-algebra multi-loop graph system, spanned jointly by fundamental independent circuits plus every vertex self-loop. The system is globally free of topological frustration iff both conditions hold:

1. **Fundamental-circuit condition**: Product of edge weights for every independent fundamental circuit in the basis equals $+1$.
2. **Self-loop condition**: Every vertex self-loop satisfies $M_n(i,i) = 1 \quad (\forall\, i \in \mathcal{I}_n)$.

If any generator evaluates to -1, algebraic independence of basis generators forbids cancelling this local negative bias; global frustration necessarily arises.

3.3 Numerical-Matrix Demonstration for Complex Multi-Loop Frustration

Consider this third-order real-valued matrix:

$$M_3 = \begin{pmatrix} 1 & 1 & 1 \\ 1 & -1 & -1 \\ 1 & -1 & -1 \end{pmatrix} \in \mathcal{M}_3.$$

Test against generalized cycle-space basis:

1. Evaluate product for fundamental circuit $1 \to 2 \to 3 \to 1$:

$M_3(1,2) \cdot M_3(2,3) \cdot M_3(3,1) = 1 \cdot (-1) \cdot 1 = -1$. Violates condition 1 → circuit frustrated.

2. Inspect diagonal self-loops: $M_3(2,2) = -1$. Violates condition 2 → self-loop conflict.

By the algebraic-frustration-basis theorem, basis generators contain -1; the whole system falls into deep frustration.

Evaluate the two-step-walk path-interference entry $(M_3^2)_{23}$ via row-column inner-product expansion:

$$(M_3^2)_{23} = M_3(2,1)M_3(1,3) + M_3(2,2)M_3(2,3) + M_3(2,3)M_3(3,3) = \\ 1 \cdot 1 + (-1)(-1) + (-1)(-1) = 3.$$

The result yields deterministic real scalar 3. Numerically demonstrates residual local net-bias arising from multi-loop superposition and self-loop sign entanglement which cannot be cancelled to zero.

---

4 Rigorous Mapping Specification and Property Proof for Local Graph Expansion Operator

4.1 Iteration Origin and Operator Mapping

Absolute iteration origin is constant matrix $M_1 = (1) \in \mathcal{M}_1$.

Full name: Local Graph Expansion Operator; notation: $\mathcal{G}_{n\to n+1}$.

It acts strictly upon one discrete realized square matrix with mapping:

$$\mathcal{G}_{n\to n+1} : \mathcal{M}_n \longrightarrow \mathcal{M}_{n+1}[\mathbf{x}_{n+1}, y_{n+1}].$$

4.2 Matrix-Expansion Structural Equation and Explicit-Vector Definition

For any input realized matrix $M_n \in \mathcal{M}_n$, the expansion operator outputs a unique formal polynomial block-matrix:

$$\mathcal{G}_{n\to n+1}(M_n) = \begin{pmatrix} M_n & \mathbf{x}_{n+1} \\ \mathbf{x}_{n+1}^T & y_{n+1} \end{pmatrix}.$$

Frontier-coupling formal column vector:

$$\mathbf{x}_{n+1} = \begin{bmatrix} x_{(n+1,1)} \\ x_{(n+1,2)} \\ \vdots \\ x_{(n+1,n)} \end{bmatrix} \in \left(\mathbb{R}[\mathcal{V}_{n+1}]\right)^n.$$

The formal symbolic matrix enforces read-only subspace inheritance:

$M_{n+1}[1:n,\ 1:n] \equiv M_n$.

The $n+1$ algebraic indeterminates inside frontier set $\mathcal{V}_{n+1}$ are mutually algebraically independent.

4.3 Walk-through Example: Primordial Expansion $M_1 \to M_2$

1. **Input initial state**: Starting constant $M_1 = (1) \in \mathcal{M}_1$.
2. **Operator invocation**: Call $\mathcal{G}_{1\to 2}(M_1)$. Declare second-order formal-variable set $\mathcal{V}_2 = \{x_{(2,1)},\ y_2\}$.
3. **Output formal structure**: Substitute into structural equation ($n=1$, vector degenerates to scalar component). Resulting second-order formal matrix:

$$M_2(x_{(2,1)},\, y_2) = \begin{pmatrix} 1 & x_{(2,1)} \\ x_{(2,1)} & y_2 \end{pmatrix} \in \mathcal{M}_2[\mathbf{x}_2,\, y_2].$$

This example illustrates how operator preserves fixed historical constants in top-left block while appending parameterized rows-columns at frontier.

4.4 Operator-Mapping-Properties Proof (Injective but Non-Surjective)

Theorem 2: Injectivity and Non-Surjectivity of Expansion Operator

> Theorem summary: Strict injectivity guarantees lossless inheritance of historical heterogeneity; non-surjectivity means its image only covers formal matrices whose top-left sub-block contains purely numerical constant entries.

Proof:

1. **Injectivity**:

Take two distinct input matrices $M_n^{(1)} \neq M_n^{(2)} \in \mathcal{M}_n$. There exists some index pair $\exists i, j \in \mathcal{I}_n$ such that $M_n^{(1)}(i,j) \neq M_n^{(2)}(i,j)$. By read-only-subspace inheritance property:

$$\mathcal{G}_{n\to n+1}(M_n^{(1)})[i,j] = M_n^{(1)}(i,j) \neq M_n^{(2)}(i,j) = \mathcal{G}_{n\to n+1}(M_n^{(2)})[i,j].$$

Hence $\mathcal{G}_{n\to n+1}(M_n^{(1)}) \neq \mathcal{G}_{n\to n+1}(M_n^{(2)})$. Injectivity is proven.

2. **Non-surjectivity**:

Co-domain space $\mathcal{M}_{n+1}[\mathbf{x}_{n+1}, y_{n+1}]$ in principle permits formal indeterminates inside its top-left $n \times n$ sub-block. Whereas any element from image $\mathrm{Im}(\mathcal{G}_{n\to n+1})$ must have purely numerical constant top-left sub-block.

Consider matrix B in co-domain such that $B[1,1] = x_{(n+1,1)}$. This violates constant-sub-block inheritance constraint and cannot belong to operator image. Non-surjectivity proven. ■

---

5 Divergence Quantification and General-Invariant Proof

5.1 Quantitative Divergence under Full-Connection Fixed Evaluation

If intermediate formal states skip downstream operator solving and are directly forced into full-connection fixed-assignment special case (Φ_{full}: $\Phi_{\text{full}}(\mathbf{x}_{n+1}) = \mathbf{1}_n$, $\Phi_{\text{full}}(y_{n+1}) = 1$).

Define total-sum series for evaluated real-matrix entries:

$S_n = \sum_{i=1}^{n} \sum_{j=1}^{n} \Phi_{\text{full}}(M_n(i,j))$,

spectral-radius invariant: $\rho(\Phi_{\text{full}}(M_n))$.

Recurrence relation for global total-sum via block-matrix expansion:

$$S_{n+1} = S_n + 2\sum_{k=1}^{n} \Phi_{\text{full}}(x_{(n+1,k)}) + \Phi_{\text{full}}(y_{n+1}).$$

Substitute full-connection assignment:

$$S_{n+1} = S_n + 2n + 1.$$

Starting from initial-condition $S_1 = 1$:

$$S_n = 1 + \sum_{k=1}^{n-1}(2k+1) = n^2 \implies \lim_{n\to\infty} S_n = \lim_{n\to\infty} n^2 = \infty.$$

Under this assignment M_n reduces to all-ones real matrix $J_n \in \{1\}^{n\times n}$. Its spectral-radius:

$$\rho(\Phi_{\text{full}}(M_n)) = n \implies \lim_{n\to\infty} \rho(\Phi_{\text{full}}(M_n)) = \infty.$$

Divergence conclusion: Forcing fixed full-connection assignments at expansion frontier yields global scalar-sum diverging in $O(n^2)$ quadratic order and spectral-radius diverging linearly $O(n)$. This justifies algebraically keeping formal-symbol matrices unevaluated, to be resolved only by downstream operators.

5.2 General Diagonal-Invariant Theorem (Proof for arbitrary n)

Theorem 3: General Diagonal-Invariant Theorem

> Theorem summary: Within formal-matrix-square setting, after applying binary-assignment homomorphism Φ, diagonal path-interference polynomial for newly injected node evaluates deterministically to constant $n+1$.

Important scope limitation: This invariant holds **strictly for two-step graph walks only (M^2)**. For higher-order powers $M^k,\ k \geq 3$, cross-terms mixing frontier indeterminates and historical constants appear; diagonal entries no longer reduce to assignment-independent constants.

Proof:

Formally multiply block-structured formal matrix:

$$M_{n+1}^2 = \begin{pmatrix} M_n & \mathbf{x}_{n+1} \\ \mathbf{x}_{n+1}^T & y_{n+1} \end{pmatrix} \begin{pmatrix} M_n & \mathbf{x}_{n+1} \\ \mathbf{x}_{n+1}^T & y_{n+1} \end{pmatrix}.$$

Extract last diagonal entry of product matrix:

$$(M_{n+1}^2)_{n+1,\,n+1} = \left(\sum_{m=1}^{n} M_{n+1}(n+1,\,m)^2 \right) + M_{n+1}(n+1,\,n+1)^2.$$

Substitute formal-symbol components:

$$(M_{n+1}^2)_{n+1,\,n+1} = \left(\sum_{m=1}^{n} x_{(n+1,m)}^2 \right) + y_{n+1}^2.$$

Apply global evaluation homomorphism Φ:

$$\Phi\Big((M_{n+1}^2)_{n+1,\,n+1}\Big) = \left(\sum_{m=1}^{n} \Phi(x_{(n+1,m)})^2 \right) + \Phi(y_{n+1})^2.$$

Hard binary-range constraints:

$\Phi(x_{(n+1,m)}) \in \{-1, 1\},\ \Phi(y_{n+1}) \in \{-1, 1\}$.

Any element squared in this binary-set equals real constant 1.

$$\Phi(x_{(n+1,m)})^2 \equiv 1, \quad \Phi(y_{n+1})^2 \equiv 1.$$

Substitute constants back to summation:

$$\Phi\Big((M_{n+1}^2)_{n+1,\,n+1}\Big) = \left(\sum_{m=1}^{n} 1 \right) + 1 = \underbrace{1+1+\cdots+1}_{n \text{ terms}} + 1 = n+1.$$

This reduction holds for arbitrary positive integer $n \in \mathbb{N}^+$, independent of specific assignments for indeterminates. Theorem 3 proven. ■

6 Constraint Compatibility and Iterative-Complexity-Bound Analysis

6.1 Constraint-Compatibility Analysis

Operator 1 outputs formal-symbol matrix containing $n+1$ algebraically-independent indeterminates; resulting Boolean solution-space size is 2^{n+1}.

Because expansion $\mathcal{G}_{n\to n+1}$ only linearly adds degrees-of-freedom, and top-left block M_n is numerically fixed in prior iterations, constraint systems built from downstream operators cannot retroactively alter historical fixed solutions. This guarantees **permanent constraint compatibility**: at least one valid binary real-solution always exists for local-interference polynomial systems.

6.2 Iterative-Complexity-Bound Analysis

Separate algebraic-logic complexity versus physical runtime-implementation overhead:

1. **Formal-algebra construction complexity**: Operator 1 only performs formal declaration for unknown-variable block structure; pure formal-logic overhead is strictly constant complexity $O(1)$.
2. **Physical-storage-complexity remark**: The above $O(1)$ applies **only to symbolic-declaration logical layer**. If runtime performs real physical-memory allocation, constructs variable-hash lookup tables or iterates variable-sets, physical-memory overhead scales linearly as $O(n)$.
3. **Downstream formal-matrix-multiplication complexity**: If downstream code computes M_{n+1}^2 formal product, naive multivariate-polynomial expansion upper-bound reads $O((n+1)^3)$. But leveraging that top-left block M_n is purely numerical constant, block-aware symbolic-multiplication reduces practical polynomial-multiplication overhead to $O(n^2)$. This guarantees engineering feasibility for iterating to large system-sizes.

Appendix Directory

For stable large-scale distributed-computation deployment, this algebraic specification defines these auxiliary appendices (accessible via main system catalogue):

- **Appendix A: Topological-Routing Protocols for Causal Data Streams**: Detailed specification for lossless serialization-transmission protocols of parameter-symbol matrices across multi-level data buses.
- **Appendix B: Bidirectional-Conversion Specification between Formal-Polynomial Matrices and Evaluated Real-Valued Matrices**: Defines hash-evaluation mapping tables for homomorphism Φ plus hardware-aware memory-alignment permutation mechanisms.
- **Appendix C: Graph Degradation and Exception-Handling for Empty-Solution Cases**: Roll-back and topological self-healing algorithm invoked under extreme conditions (e.g. external tampering over read-only blocks) leading to no valid real-solutions for characteristic equations.
- **Appendix D: Downstream-Operator Polynomial-Interface Output-Format Standard**: Layout standard for multivariate-polynomial coefficient matrices when passing off-diagonal path-interference elements toward numerical solvers.

Operator-1: Local Graph Expansion Operator - Full Explanatory Document

Version: 1.0

Document purpose: supporting notes for papers, writing reference, helping readers rapidly understand $\mathcal{G}_{n\to n+1}$.

Corresponding original reference:
Operator 1: Mathematical Specification and Algebraic Foundations of the Local Graph Expansion Operator (Yue Lu, v1.0)
https://doi.org/10.5281/zenodo.21454140, belonging to the Status-Relational Entropy Dynamics (SRE) theoretical system.

Core positioning: Operator-1 is neither a graph generator nor a dynamics solver. It serves as the underlying algebraic syntax layer of the SRE system, defining how the relational possibility-space expands with evolutionary pulses.

Table of Contents

1. Overview and Core Positioning
2. Foundational Axioms and Notation System
3. Mathematical Definition and Block-Matrix Structure of the Operator
4. Inductive-limit Ring R_∞ and Evaluation Homomorphism Φ (Separation between Symbolic Layer and Real-State Layer)
5. Key Mathematical Properties of the Operator (Theorem 2: Injectivity and Non-Surjectivity)
6. Connections of the Operator to Topological Frustration and the Diagonal Invariance Theorem
7. Operator-1 vs. Simulation Dynamics: Strict Demarcation between Operator Layer / Downstream Evaluation-Simulation Layer
8. Ontological Implications (SRE Relational Ontology)
9. List of Common Misconceptions
10. Physical Meaning within the SRE Cosmological Picture
11. Open Theoretical Boundaries and Future Extensions

---

1. Overview and Core Positioning

The local graph expansion operator $\mathcal{G}_{n\to n+1}$ implements dimensional expansion of the system from order n to order $n+1$.

❗ Critical distinctions:

1. The operator only constructs formal symbolic matrices; it performs no assignment, introduces no randomness, and computes no dynamical evolution.
2. It merely expands the potential relational possibility-set. Mapping symbolic variables to numerical real-world values $-1, +1$, asynchronous activation, probabilistic evolution, and global feedback all belong to the downstream SRE dynamical module and are **not** part of Operator-1 itself.

Underlying ontological premise:
There is no pre-given geometric space or coordinates within the system. Graph vertices are merely index labels; vertices carry no intrinsic scalar values. All physical information is encoded in relations between vertices (bidirectional edges and vertex self-loops).

Each invocation of the operator:

- Adds one abstract vertex index;
- Generates a fresh set of algebraically independent formal symbolic variables representing coupling relations between the new vertex and all historical vertices, together with the self-loop relation of this new vertex;
- Strictly inherits the full historical structure, and never modifies old relations.

Core question addressed by Operator-1:
In a discrete system with only relations and no pre-existing space, what rigorous algebraic rules shall govern the expansion of the whole system's relational possibility-space, while guaranteeing lossless preservation of historical information and self-consistent temporal evolution.

2. Foundational Axioms and Notation System

Foundational Axioms (SRE System)

1. **Binary-Relational Range Axiom**: After evaluation via the homomorphism Φ, all relations (edges, self-loops) may only take values $+1, -1$. Zero-values do not exist; there is no "no-edge" state. $+1$ denotes a cooperative relation; -1 denotes an antagonistic relation.
2. **Initial-Seed Axiom**: Evolution commences from the first-order matrix $M_1 = [1]$, representing the self-loop relation of the initial unit.
3. **History Read-Only Axiom**: Historical matrix sub-blocks, once generated, cannot be rewritten by the expansion operator.

Main Notation

Symbol	Meaning
$\mathbb{N}^+$	Set of positive integers $1, 2, 3, \ldots$, excludes zero
J_n	Index reference set $1, 2, \ldots, n$; 1-indexing is used throughout this document
$\mathcal{M}_n$	Space of n-dimensional discrete instance matrices: real-symmetric binary matrices with entries $\in \{-1, +1\}$
$M_{n+1}[v_{n+1}]$	Parameter-matrix space carrying formal symbolic variables
v_{n+1}	Set of newly-injected independent formal variables at expansion step $n \to n+1$: $x_{n+1,1}, \ldots, x_{n+1,n}, y_{n+1}$
$\mathcal{R}_\infty$	Universal multivariate polynomial ring constructed via inductive limit, containing all formal variables over the full evolutionary history
Φ	Global evaluation homomorphism: maps formal variables inside $\mathcal{R}_\infty$ onto real numbers, subject to $\Phi(\cdot) \in \{-1, +1\}$
$\mathcal{G}_{n\to n+1}$	Local graph expansion operator
$\partial_{n\to m}$	Canonical embedding operator, embedding lower-order polynomial rings into higher-order polynomial rings

3. Mathematical Definition and Block-Matrix Structure of the Operator

Operator mapping:

$$\mathcal{G}_{n\to n+1} : \mathcal{M}_n \to \mathcal{M}_{n+1}[x_{n+1}, y_{n+1}]$$

Input: a fully-instantiated n-order real-symmetric binary matrix M_n (historically frozen real-world state).

Output: an $n+1$-order formal symbolic block matrix:

$$M_{n+1}(x_{n+1}, y_{n+1}) = \mathcal{G}_{n\to n+1}(M_n) = \begin{pmatrix} M_n & x_{n+1} \\ x_{n+1}^{\mathrm{T}} & y_{n+1} \end{pmatrix}$$

where

$$x_{n+1} = \begin{bmatrix} x_{(n+1,1)} \\ x_{(n+1,2)} \\ \vdots \\ x_{(n+1,n)} \end{bmatrix}$$

- $x_{(n+1,m)}$: formal symbolic variable for the coupling relation between the new vertex and old vertex m.
- y_{n+1}: formal symbolic variable for the self-loop relation of the new vertex.

Constraint: the top-left sub-block satisfies $M_{n+1}[1:n, , 1:n] = M_n$; it is strictly inherited from the input and cannot be modified.

Storage aspect: upon expansion from n to $n+1$, the array gains $2n+1$ new matrix cells. Owing to matrix symmetry, only $n+1$ of these carry independent algebraic degrees-of-freedom (n pairwise-coupling edges plus one self-loop). The remaining cells are symmetric mirror copies and carry no new information.

Minimal example: $M_1 \to M_2$

$$\mathcal{G}_{1\to 2}(M_1) = \begin{pmatrix} 1 & x_{(2,1)} \\ x_{(2,1)} & y_2 \end{pmatrix}$$

One observes that the historical seed $M_1 = [1]$ is fully preserved; two independent symbolic variables are added and remain unassigned.

4. Inductive-limit Ring $\mathcal{R}_\infty$ and Evaluation Homomorphism Φ

Repeated invocations of $\mathcal{G}_{n\to n+1}$ continuously produce new formal variables. To avoid naming conflicts among variables, the universal polynomial ring $\mathcal{R}_\infty$ is constructed via the inductive (direct) limit within commutative algebra:

1. Variable sets generated in distinct expansion steps are pairwise disjoint: $v_k \cap v_m = \emptyset$ for $k \neq m$.
2. Canonical embeddings $\partial_{n\to m}$ realize injection of lower-order polynomials into higher-order rings.
3. $\mathcal{R}_\infty = \varinjlim \mathcal{R}_n$, containing all formal-symbol polynomials across the full evolutionary history.

$\mathcal{R}_\infty$ represents the complete relational possibility-space; at this stage all quantities remain purely symbolic with no assigned physical numerical values.

The global evaluation homomorphism Φ is the mapping transporting us from the symbolic layer toward realized physical states:

$$\Phi\colon \mathcal{R}_\infty \to \mathbb{R}$$

It satisfies homomorphism properties:
$\Phi(f+g) = \Phi(f) + \Phi(g), \quad \Phi(f\cdot g) = \Phi(f)\cdot\Phi(g),$
subject to the hard constraint:

$$\Phi\Big(x_{(k,m)}\Big) \in \{-1,+1\}, \quad \Phi\Big(y_k\Big) \in \{-1,+1\}.$$

Important notes:

- Operator-1 adds fresh formal variables into $\mathcal{R}_\infty$ (expanding the possibility-space).
- Φ projects possibilities onto realized numerical matrix states.
- Operator-1 itself does **not** perform evaluation via Φ.

5. Key Mathematical Properties of the Operator (Theorem 2: Injectivity and Non-Surjectivity)

1. **Strict Injectivity**

Given two distinct historical matrices $M_n^{(1)} \neq M_n^{(2)}$, the symbolic-matrix outputs of the operator are guaranteed to differ:

$$\mathcal{G}_{n\to n+1}\Big(M_n^{(1)}\Big) \neq \mathcal{G}_{n\to n+1}\Big(M_n^{(2)}\Big).$$

Physical interpretation: historical heterogeneity is preserved without loss; expansion never erases pre-existing structural information.

2. **Strict Non-Surjectivity**

The image set of the operator constitutes only a subset of the target symbolic-matrix space. The top-left $n\times n$ sub-block of any output must be a constant numerical block and cannot contain formal variables.
That is: not every $n+1$-order symbolic matrix can be generated by Operator-1.

Physical interpretation: the operator only expands new potential relations on top of already-fixed history. It cannot arbitrarily construct symbolic matrices lacking any frozen historical sub-structure.

6. Connections of the Operator to Topological Frustration and the Diagonal Invariance Theorem

Operator-1 supplies the formal matrix structure upon which the full set of core structural theorems can be derived.

> **Theorem 3: General Diagonal Invariance Theorem (valid only for matrix square M^2, two-step graph walks)**

Take the diagonal entry corresponding to the new vertex after formally squaring the expanded matrix:

$$\left(M_{n+1}^2\right)_{n+1,n+1} = \sum_{m=1}^{n} x_{(n+1,m)}^2 + y_{n+1}^2$$

After applying evaluation homomorphism Φ and using the identity $\forall s \in \{-1, +1\},\ s^2 = 1$, we obtain:

$$\Phi\left(\left(M_{n+1}^2\right)_{n+1,n+1}\right) = n + 1$$

This diagonal interference term evaluates to the constant $n + 1$ regardless of variable assignments.

⚠️ Caveat: this invariance holds **only for matrix squares**. It fails for higher-order matrix powers with $k \geq 3$.

Topological Frustration (Theorem 1, Corollary 1)

Topological frustration is defined **after** symbolic variables are evaluated to ± 1 and yield real-valued matrices:

1. Frustration of a single loop: the product of edge weights along a closed circuit equals -1.
2. Global frustration criterion for complex networks: the system is globally frustration-free only if the product over all generators of the generalized cycle space (fundamental loops plus all self-loops) equals $+1$.

Operator-1 continuously creates new vertices and new relational slots, thereby generating new loops and new self-loops. Whether topological frustration emerges depends on the concrete assignments delivered by downstream Φ; frustration is not an intrinsic property of the operator itself.

7. Operator-1 vs. Simulation Dynamics: Strict Demarcation between Operator Layer / Downstream Evaluation-Simulation Layer

This is the most frequently confused section; a sharp separation must be maintained.

Item	**Operator-1 (Operator / Symbolic-Algebra Layer)**	**Hierarchical Dissipative Self-Organizing Binary-Network Dynamics (Downstream Dynamics & Simulation Layer)** https://doi.org/10.5281/zenodo.20576606
Core Behaviour	Produces symbolic block matrices carrying formal variables	Executes evaluation homomorphism Φ, instantiates symbolic variables to ± 1
Stochasticity	Zero intrinsic randomness	Introduces stochastic dormancy via asynchronous activation χ_{ij}
Probabilistic Rules	None	$p_{ij}^{(n)}$ is jointly modulated by local frustration energy and evolutionary distance
Self-loop Handling	y_{n+1} remains an unassigned symbolic variable	Bottom-right self-loop is fixed by global negative feedback from total matrix sum; no randomness
Output Product	Polynomial symbolic matrix	Deterministic real-symmetric binary numerical matrix
Role	Algebraic syntax for evolution; expands possibility-space	Filters realized physical states out of the possibility-space

In simulation code, each iteration performs two actions simultaneously: it implements the structural expansion equivalent to Operator-1, and immediately executes downstream dynamics to assign values to all variables. For this reason many readers mistakenly treat simulation behaviour as intrinsic behaviour of the operator, which constitutes a typical misconception.

8. Ontological Implications (SRE Relational Ontology)

1. Vertices function purely as index markers; vertices possess no intrinsic values. All ± 1 values belong to relations (edges, self-loops). The self-loop y_{n+1} represents a "relation of a vertex toward itself", and is **not** a numerical property of the vertex.
2. No pre-existing background geometric space is presupposed. Space, distance and dimensionality are not fundamental primitives; they are macroscopic emergent phenomena in the large-system statistical limit.
3. Each pulse iteration of Operator-1 advances fundamental evolution. Pulse-driven expansion itself constitutes the primitive manifestation of emergent time. New vertices are not dynamically generated from within the old network; they appear as the relational possibility-space expands as evolution proceeds.

Under current version: vertex generation follows the evolutionary-pulse axiom. Endogenous generation driven by internal system conditions remains a target for future work.

9. List of Common Misconceptions

Misconception 1: Operator-1 is an ordinary growing-graph algorithm (e.g. Barabási-Albert scale-free networks).
❌False: conventional network algorithms directly output concrete graphs. Operator-1 only outputs symbolic possibility templates without value-assignment. Graph generation and stochastic evolution belong to downstream dynamics.

Misconception 2: New vertices are dynamically produced by the internal dynamics of the older network.
❌False: internal dynamics only modify relational values. Vertex indices are introduced via the evolutionary-pulse axiom; they are not automatically triggered by internal tensions (within the current version).

Misconception 3: Diagonal matrix entries represent intrinsic state values of vertices.
❌False: diagonal entries encode self-loop relations - a special class of relation - not intrinsic vertex properties.

Misconception 4: The operator itself includes randomness, dormancy rules and global feedback.
❌False: all those mechanisms belong exclusively to the downstream SRE dynamical module.

Misconception 5: The $2n+1$ newly-added matrix array cells correspond to $2n+1$ new graph vertices.
❌False: exactly one graph vertex is added per expansion step. $2n+1$ counts storage-level array cells of the symmetric matrix; only $n+1$ of those correspond to independent degrees-of-freedom.

Misconception 6: Topological frustration is an inherent property of Operator-1.
❌False: topological frustration only arises **after** variable evaluation via homomorphism Φ. Frustration cannot be judged while variables remain purely symbolic.

10. Physical Meaning within the SRE Cosmological Picture

The full SRE framework - Status-Relational-Entropy Dynamics - maps onto Operator-1 as follows:

1. **Relational**: Operator-1 expands the relational possibility-space and constructs $\mathcal{R}_\infty$.
2. **Status**: Downstream evaluation homomorphism plus hierarchical dissipative dynamics select concrete realized system states out of the possibility-space.
3. **Entropy**: Entropy quantifies the statistical gap between the enormous latent set of relational possibilities and the small subset of actually-realized configurations. Asynchronous dormancy and topological frustration jointly drive entropy evolution.
4. **Dynamics**: Governs how realized states evolve atop the continuously expanding possibility-space.

Within the axiomatic system of SRE, the framework offers a conceptually self-consistent dissolution of the classic cosmological paradox: "What lies beyond the boundary of the universe?".

The evolutionary boundary of the universe corresponds to the expansion frontier of Operator-1. Beyond this frontier there exists no space and no void. Only formal relational possibilities not yet realized by the evaluation homomorphism reside there. Extrapolating emergent geometric concepts of spacetime past this frontier constitutes an invalid logical extrapolation.

##-. Open Theoretical Boundaries and Future Extensions

1. **Endogenization of vertex generation**: In the present version new vertices are introduced via an external-pulse axiom.

Future work aims to trigger vertex birth internally by thresholds of topological-frustration tension, path-interference effects or entropy fluctuations, eliminating the external-pulse postulate.

2. **Integration with Random-Matrix Theory (RMT)**: Perform spectral-statistical analysis for this iteratively-grown matrix ensemble derived from the operator. Study universal large-N spectral behaviour and distinguish microscopic fluctuations from macroscopic statistical determinacy.
3. **Reduction of axioms**: Attempt to reduce the expansion axiom and binary-range axiom more deeply from the ontological premise that "a perfect zero (absolute nothingness) cannot be physically realized".
4. **Generalization of the operator**: Extend operator formalism to permit non-uniform, non-sequential advance of the possibility-space frontier.

> **Supplementary Explanatory Note | This is interpretive material, not original mathematical specification. All mathematical proofs should be consulted in the original source paper.**
> DOI: https://doi.org/10.5281/zenodo.21454140

Operator-2： Local Metric & Probabilistic Pruning Operator （$\mathcal{M}_{\chi} \circ \mathcal{E}_{\text{local}}$）

Rigorous Mathematical Derivation and Specification

Version: 1.1

This specification succeeds the formal parametric topology generated by the First Operator ($\mathcal{G}_{n \to n+1}$) and strictly adheres to the No-Dimension Principle. The framework completely eradicates any dependency on background coordinate metrics, embedding spaces, or artificial spatiotemporal metrics, computing discrete evolutionary steps purely via local topological invariants (causal depths and graph-walk statistics).

---

1. Unified Algebraic Notation Index

To ensure strict cross-operator compatibility, the global algebraic symbols are formally standardized as follows:

- $\mathbb{N}^{+}$: The set of strictly positive integers, defined as $\{1, 2, 3, \ldots\}$ (excluding 0).
- $\mathcal{I}_n$: The index baseline set, defined as $\{1, 2, \ldots, n\}$ using 1-based indexing.
- $\mathcal{R}_{\infty}$: The inductive limit multi-variate polynomial maternal ring defined by Operator 1.
- Φ: The global evaluation homomorphism mapping ($\Phi : \mathcal{R}_{\infty} \to \mathbb{R}$), whose value domain is rigidly locked to the discrete binary spin set $\{+1, -1\}$.
- $G_n = (V_n, E_n, \mathbf{S}_n)$: The non-background-dependent labeled graph at step n, where V_n is the vertex set, E_n is the directed edge set, and $\mathbf{S}_n$ is the binary spin configuration mapping.
- v_f: The unassigned parametric frontier vertex injected at the $(n+1)$-th evolutionary step, satisfying $\text{Rank}(v_f) = n + 1$.
- $\mathcal{M}_{n \times n}$: The n-dimensional discrete-valued realized matrix space populated by purely binary real scalars.
- $\mathcal{P}(\cdot)$: The discrete operational probability measure space.

---

2. Morphic Integration: Homomorphic Mapping Between Graph Topology and Matrix Space

To bridge the structural domain gap, the non-background-dependent labeled graph $G_n = (V_n, E_n, \mathbf{S}_n)$ is mapped bijectively to the First Operator's matrix space $\mathcal{M}_{n \times n}$.

2.1 Topological Space Homomorphism

We define a bijective coordinate-labeling mapping $\sigma : V_n \to \mathcal{I}_n = \{1, 2, \ldots, n\}$, mapping discrete graph vertices to algebraic matrix indices. The homomorphic surjection from the labeled graph G_n to the binary spin real symmetric square matrix $\mathbf{M}_n \in \mathbb{R}^{n \times n}$ satisfies:

$$\mathbf{M}_n(i,j) = \begin{cases} \mathbf{S}_n(\sigma^{-1}(i), \sigma^{-1}(j)), \text{if } (\sigma^{-1}(i), \sigma^{-1}(j)) \in E_n \\ +1 \text{ or } -1, \text{if } i = j \text{ (Vertex self-loop entries mapped onto the matrix diagonal)} \end{cases}$$

2.2 Causal Topological Depth Invariant

The endogenous birth order of a vertex is monotonically mapped onto its causal label $\text{Rank}(v_i) = \sigma(v_i)$. For any newly injected frontier increment vertex v_f, its birth rank strictly satisfies $\text{Rank}(v_f) = n + 1$. The strict **Topological Depth Invariant** $\mathcal{D}_s$ between the frontier vertex v_f and any historic first-order neighborhood vertex $v_m \in \mathcal{N}(v_f)$ is defined as:

$$\mathcal{D}_s(v_f, v_m) = \text{Rank}(v_f) - \text{Rank}(v_m) = (n+1) - \sigma(v_m)$$

Theorem 1 (Adherence to the No-Dimension Principle)

The topological depth metric $\mathcal{D}_s(v_f, v_m)$ strictly adheres to the No-Dimension Principle across all evolutionary cascades, without introducing external geometric coordinates or artificial spatiotemporal metric tensors.

Proof

1. Within the multi-variate polynomial ring inductive limit construction of Operator 1, the system over finite step lengths manifests as a directed partially ordered set (Directed Poset).
2. The sequence ordinal derived from the bijection $\sigma(v_i)$ represents the chain length invariant of the graph under canonical embeddings into the inductive limit maternal ring.
3. The rank difference between two vertices purely characterizes their local relative generational delay within the causal ordering of the graph. It is an endogenous property of the dimension-free graph structure itself, eliminating any reliance on background coordinate geometry.

■ Theorem 1 is complete.

---

3. Complexity Resolution: Endogenous Proof of the $O(1)$ Local Time-Stream Opened Boundary

Defining the local time-stream boundary as a qualitative limit of a ratio is mathematically invalid. We introduce the topological firewall saturation boundary to derive a rigorous constant asymptotic upper bound.

Theorem 2 (Asymptotic Constancy of Local Computational Overhead)

Under a sparse graph storage architecture bounded by a local Topological Firewall, the single-step temporal computational complexity $T(n)$ of the Second Operator evaluating metrics and pruning decisions on the frontier vertex v_f is bounded strictly at $T(n) = O(1)$ as the system dimension expands indefinitely ($n \to \infty$).

Proof

1. The system's endogenous dissipative mechanisms trigger massive localized dormancy across mesoscopic and macroscopic layers. Let the Topological Firewall strictly confine the operator's active domain to a finite K-hop neighborhood of v_f, such that the degree of any vertex satisfies a global absolute constant upper bound independent of the full network scale n:

$$\exists K_0 \in \mathbb{N}^+ \quad \text{s.t.} \quad \sup_{n \in \mathbb{N}^+} \max_{v \in V_n} \text{Degree}(v) \leq K_0 \ll n$$

2. The computational engine of the Second Operator is driven by traversing the first-order neighborhood $\mathcal{N}(v_f)$. From the bound above, its neighborhood cardinality satisfies $|\mathcal{N}(v_f)| \leq K_0$.
3. For any historical neighborhood vertex $v_m \in \mathcal{N}(v_f)$, finding the common neighborhood intersection set $\Omega_1 = \mathcal{N}(v_f) \cap \mathcal{N}(v_m)$ via an optimized hash-table lookup requires an asymptotic overhead bounded by $O(\text{Degree}(v_f) + \text{Degree}(v_m)) \leq 2 \cdot K_0$.
4. The algebraic overhead for evaluating the complete 2-step graph walk interference along a single edge scales linearly with the intersection cardinality, i.e., $O(|\Omega_1|) \leq O(K_0)$.
5. Summing over the active frontier, the total number of elementary algebraic operations $\mathcal{A}(n)$ executed during a single dimensional expansion step satisfies:

$$\mathcal{A}(n) \leq |\mathcal{N}(v_f)| \times (2K_0 + K_0) \leq 3K_0^2$$

6. Let $C = 3K_0^2$. Since K_0 and n are algebraically independent, there exists an evolutionary baseline $N_0 = 1$ such that for all $n > N_0$, $\mathcal{A}(n) \leq C \cdot 1$. By the standard definition of asymptotic upper bounds:

$$T(n) = O(1) \quad \text{as } n \to \infty$$

■ Theorem 2 is complete.

---

4. Algebraic Integration: Full Algebraic Expansion of the Complete 2-Step Graph Walk

Narrowing path interference down to a simple common neighborhood intersection truncates the algebraic tractability of the graph-power walk. We establish the complete un-truncated algebraic expansion based on the full polynomial adjacency matrix.

Theorem 3 (2-Step Topological Path Interference Expansion Theorem)

Let $\mathbf{A} \in \mathcal{R}_{\infty}^{(n+1)\times(n+1)}$ denote the multi-variate polynomial adjacency matrix outputted by the First Operator prior to evaluation. The complete 2-step walk path interference operator connecting the frontier vertex v_f to its historical first-order neighbor v_m is strictly isomorphic to the matrix square domain without introducing external coordinate bases:

$$\mathcal{J}(v_f, v_m) \equiv (\mathbf{A}^2)_{f,m}$$

Proof

1. By algebraic graph theory, the number of distinct k-step walks between any two nodes corresponds identically to the k-th power of the adjacency matrix. Inside the multi-variate polynomial ring matrix space, matrix multiplication is defined by the standard inner product expansion:

$$(\mathbf{A}^2)_{f,m} = \sum_{v_k \in V_{n+1}} \mathbf{A}(v_f, v_k) \cdot \mathbf{A}(v_k, v_m)$$

2. We partition the global vertex set V_{n+1} strictly into four mutually disjoint local topological subsets:
   - The independent common neighborhood subset: $\Omega_1 = \mathcal{N}(v_f) \cap \mathcal{N}(v_m)$
   - The frontier vertex singleton subset: $\Omega_2 = \{v_f\}$
   - The target historical neighbor singleton subset: $\Omega_3 = \{v_m\}$
   - The residual complement (non-coherent isolated vertices): $\Omega_4 = V_{n+1} \setminus (\Omega_1 \cup \Omega_2 \cup \Omega_3)$
3. We perform localized algebraic simplification for the summation terms across all four subsets sequentially:
   - For $v_k \in \Omega_1$, both edges (v_f, v_k) and (v_k, v_m) exist natively, yielding the product term: $\mathbf{S}_{n+1}(v_f, v_k) \cdot \mathbf{S}_{n+1}(v_k, v_m)$.
   - For $v_k \in \Omega_2 = \{v_f\}$, the corresponding inner product term evaluates to $\mathbf{A}(v_f, v_f) \cdot \mathbf{A}(v_f, v_m)$. Because $\mathbf{A}(v_f, v_f)$ represents the endogenous self-loop of the frontier vertex, we denote it as $\mathbf{M}_{n+1}(v_f, v_f)$; meanwhile, $\mathbf{A}(v_f, v_m) = \mathbf{S}_{n+1}(v_f, v_m)$. This term condenses into $\mathbf{M}_{n+1}(v_f, v_f) \cdot \mathbf{S}_{n+1}(v_f, v_m)$.
   - For $v_k \in \Omega_3 = \{v_m\}$, the term evaluates to $\mathbf{A}(v_f, v_m) \cdot \mathbf{A}(v_m, v_m)$. By symmetry, this yields $\mathbf{S}_{n+1}(v_f, v_m) \cdot \mathbf{M}_{n+1}(v_m, v_m)$.
   - For $v_k \in \Omega_4$, at least one of the edges (v_f, v_k) or (v_k, v_m) does not exist. According to the sparse boundary constraints of the First Operator, the product identically collapses to zero: $\mathbf{A}(v_f, v_k) \cdot \mathbf{A}(v_k, v_m) \equiv 0$.
4. Linearly summing the simplified components across all subsets perfectly reconstructs the global matrix inner product as a purely local topological polynomial invariant:

$$\mathcal{J}(v_f, v_m) = \sum_{v_k \in \mathcal{N}(v_f) \cap \mathcal{N}(v_m)} \mathbf{S}_{n+1}(v_f, v_k) \cdot \mathbf{S}_{n+1}(v_k, v_m) + \mathbf{S}_{n+1}(v_f, v_m) \Big[\mathbf{M}_{n+1}(v_f, v_f) + \mathbf{M}_{n+1}(v_m, v_m) \Big]$$

■ Theorem 3 is complete.

---

5. P0 Emergency Revision: Necessary and Sufficient Algebraic Equivalence for Arbitrary-Length Topological Frustration

Evaluating frustration using a raw scalar threshold lacks mathematical rigor. We reconstruct the criteria using the strict binary spin-loop product invariants of the cycle space basis.

Theorem 4 (Isomorphism Between Cycle Basis Invariants and Local Path Interference)

Let the injection of the frontier vertex v_f induce a closed polygonal loop L of arbitrary length L_{len} within the local graph, denoted by the vertex sequence $c_1 \to c_2 \to \cdots \to c_L \to c_1$, where the expanding frontier edge corresponds to $\mathbf{A}(c_L, c_1) = \mathbf{S}_{n+1}(v_f, v_m)$. Under rigid diagonal self-loop normalization $\mathbf{M}(i, i) = 1$, a closed loop structure is globally frustrated under the valuation homomorphism if and only if:

$$\Phi\Big(\mathbf{S}_{n+1}(c_L, c_1)\Big) \cdot \sum_{v_k \in \mathcal{N}(c_L) \cap \mathcal{N}(c_1)} \Big[\Phi\Big(\mathbf{S}_{n+1}(c_L, v_k)\Big) \cdot \Phi\Big(\mathbf{S}_{n+1}(v_k, c_1)\Big) \Big] < 0$$

Proof

1. Denote the edge-weight product over historical paths as $\Pi_{\text{hist}} = \prod_{m=1}^{L-1} \mathbf{A}(c_m, c_{m+1})$. Under discrete-spin homomorphism, $\Phi(\Pi_{\text{hist}}) \in \{+1, -1\}$, and for the frontier edge we have $\Phi\big(\mathbf{S}_{n+1}(c_L, c_1)\big) \in \{+1, -1\}$.
2. By definition, frustration-free condition reads: $\Phi(\Pi_{\text{hist}}) \cdot \Phi\big(\mathbf{S}_{n+1}(c_L, c_1)\big) = +1$.
3. Multiply both sides by $\Phi\big(\mathbf{S}_{n+1}(c_L, c_1)\big)$. Using the spin-set square identity $\Phi(S)^2 \equiv 1$, we obtain the equivalent relation:

$$\Phi(\Pi_{\text{hist}}) = \Phi\big(\mathbf{S}_{n+1}(c_L, c_1)\big).$$

4. When projecting historical long-path Π_{hist} via common-neighbor set $\Omega_1 = \mathcal{N}(c_L) \cap \mathcal{N}(c_1)$ in graph spectral analysis, the symbol space of historical product is isomorphic to the sign of local two-step summation term:

$$\operatorname{sgn}\left(\sum_{v_k \in \Omega_1} \Phi\big(\mathbf{S}_{n+1}(c_L, v_k)\big) \cdot \Phi\big(\mathbf{S}_{n+1}(v_k, c_1)\big)\right) \equiv \Phi(\Pi_{\text{hist}}).$$

5. Substitute the identity obtained in step 3 into step 4:

$$\operatorname{sgn}\left(\sum_{v_k \in \Omega_1} \Phi\big(\mathbf{S}_{n+1}(c_L, v_k)\big) \cdot \Phi\big(\mathbf{S}_{n+1}(v_k, c_1)\big)\right) = \Phi\big(\mathbf{S}_{n+1}(c_L, c_1)\big).$$

6. Multiply both sides by frontier-edge spin value $\Phi\big(\mathbf{S}_{n+1}(c_L, c_1)\big)$. Since absolute value of each summand equals 1, when summation is non-zero, the inner product yields positive for frustration-free loops; for frustrated loops with product -1, this inner product becomes strictly less than zero.

■ Theorem 4 is complete.

Corollary 4.1 (Quantitative Multi-Circuit Overlapping Generalization)

When multiple long-range loops highly overlap by sharing the identical frontier expansion edge $\mathbf{S}_{n+1}(v_f, v_m)$, the global unrealized local multi-circuit frustration polynomial $\tilde{\mathcal{E}}_{\text{local}}(v_f, v_m)$ generalizes quantitatively to:

$$\tilde{\mathcal{E}}_{\text{local}}(v_f, v_m) = \sum_{\alpha} w_\alpha \cdot \left(\sum_{v_k \in \Omega_\alpha} \mathbf{S}(v_f, v_k) \cdot \mathbf{S}(v_k, v_m)\right) + 2 \cdot \mathbf{S}(v_f, v_m)$$

where w_α represents the local topological weight of each independent fundamental cycle generator, quantitatively balancing out structural cancellations across the intersecting manifold.

---

6. Pruning Paradigm Evaluation: Algebraic and Topological Analysis of Masking Operations

When the decision gate dictates that a frontier channel should be pruned ($\chi = 0$), two distinct algebraic mechanisms can be enforced upon the underlying graph walk structure.

6.1 Paradigm A: Edge-Removal Mechanism (Adjacency Zeroing Mode)

Under this mode, when $\chi = 0$, the corresponding edge spin element is forced to zero ($\mathbf{M}_{n+1}(i, j) \longleftarrow 0$).

- **Topological Invariant Disruptions**: The diagonal elements of the Graph Laplacian matrix $\mathbf{L} = \mathbf{D} - \mathbf{A}$ are reduced concurrently. The zero-order Betti number β_0 (the count of independent connected components) is highly susceptible to artificial inflation due to localized path fractures, triggering a premature unbinding of the local manifold.
- **Axiomatic Conflict**: The introduction of a continuous scalar 0 directly violates the Strict Binary Constraint (Axiom 1) of Operator 1, which mandates: "The entire evolutionary period contains no continuous function truncation, nor does it have a dissipative zero state (0), and the matrix elements are strictly constrained to a binary spin set $\{+1, -1\}$". Thus, Paradigm A is unconditionally rejected.

6.2 Paradigm B: Elimination-Conduction Mechanism (Forced Spin-1 Mode)

Under this mode, when $\chi = 0$, the edge spin state is algebraically forced to a rigid real constant $+1$ ($\mathbf{M}_{n+1}(i, j) \longleftarrow 1$).

- **Topological Invariant Safeguards**: Because all matrix elements are bounded strictly within the discrete binary domain $\{+1,$

$-1\}$, the diagonal degrees of the Graph Laplacian matrix maintain their structural rigid invariants without introducing any continuous null states.

- **Frustration Preservation**: When a channel spin is forced to $+1$, the fundamental circuit product invariant $\prod \mathbf{A} = -1$ natively collapses into the spin product of the remaining historical edges along that loop. This instantly removes the edge's causal influence in the continuous multiplication feedback loop (identity-element elimination), effectively erasing its phase contribution from the manifold without physically severing the structural graph connectivity.
- **Standardized Matrix Expression**: Paradigm B perfectly satisfies both the discrete binary constraint and the topological loop frustration preservation rules. Consequently, Paradigm B is formally established as the standardized algebraic pruning rule, explicitly formulated as:

$$\mathbf{M}_{n+1}(i,j) \longleftarrow \chi\Big(\sigma^{-1}(i), \sigma^{-1}(j)\Big) \cdot \mathbf{M}_{n+1}(i,j) + \Big[1 - \chi\Big(\sigma^{-1}(i), \sigma^{-1}(j)\Big)\Big] \cdot 1$$

---

7. Statistical Mechanics Derivation: Thermodynamic Origin of the Zero-Interference Boundary

To isolate non-coherent topological states and eliminate any analytical degeneracy in probability behaviors, the absolute zero-interference plateau ($\mathrm{sgn} = 0$) must be derived independently from the microstate partition function of Status-Relational Entropy (SRE) dynamics.

Theorem 5 (SRE Hamiltonian Phase Separation Theorem)

The local topological Hamiltonian $\mathcal{H}_{\text{local}}$ driven by the un-realized path interference polynomial $\tilde{\mathcal{E}}_{\text{local}}(v_f, v_m)$ under the valuation homomorphism consists strictly of three mutually disjoint phase spaces:

$$\mathcal{H}_{\text{local}} = \begin{cases} -\mathcal{E}_{\text{local}} - 1, & \text{if } \tilde{\mathcal{E}}_{\text{local}} > 0 \text{ (Coherent Condensation Phase)} \\ 0, & \text{if } \tilde{\mathcal{E}}_{\text{local}} = 0 \text{ (Absolute Information Vacuum Phase)} \\ +\mathcal{E}_{\text{local}} + 1, & \text{if } \tilde{\mathcal{E}}_{\text{local}} < 0 \text{ (Frustrated Annihilation Phase)} \end{cases}$$

Proof

1. Based on the fundamental principles of SRE dynamics, the microscopic occurrence probability of a local front channel follows a canonical distribution governed by its local Hamiltonian, where the statistical weight evaluates to $\exp(-\beta \cdot \mathcal{H})$.
2. When $\tilde{\mathcal{E}}_{\text{local}} > 0$, the local loop is entirely free from topological frustration, allowing the network to condense coherently and lower its free energy. The Hamiltonian manifests as an attractive potential well, where the structural correction term inside the energy denominator yields $\exp(+1) = e$.
3. When $\tilde{\mathcal{E}}_{\text{local}} < 0$, the system plunges into deep structural frustration. The non-neutralizable phase conflict acts as an active gauge work. The Hamiltonian switches into a repulsive energy barrier, where the structural correction term scales down to $\exp(-\beta \cdot \Delta E_{\text{gauge}}) = \exp(-1) = e^{-1}$.
4. When $\tilde{\mathcal{E}}_{\text{local}} = 0$, a perfect algebraic cancellation occurs between positive and negative polarities across all 2-step paths. In the language of statistical physics, this local manifold possesses neither ordered-phase condensation nor phase-gauge work. The degeneracies of opposite-energy states are perfectly balanced, collapsing the local topological Hamiltonian strictly to zero: $\mathcal{H}_{\text{local}} \equiv 0$.
5. Substituting $\mathcal{H}_{\text{local}} = 0$ back into the Boltzmann canonical factor, the phase-separation correction term yields $\exp(0) = 1$. This eliminates any mathematical continuity gaps at the phase boundaries, ensuring complete decoupling between different topological states.

■ Theorem 5 is complete.

---

8. Endogenous Resolution: Grand Proof of the Global Absolute Vertex Degree Upper Bound K_0

Treating the maximum vertex-degree bound as an external hard-coded assumption introduces a systemic disconnect. We derive the absolute saturation boundary of the global-degree topology endogenously from the pruning-probability measure and SRE entropy changes.

Theorem 6 (Global Degree Saturation and Endogenous $O(1)$ Complexity Theorem)

For any newly injected frontier vertex v_f, its local first-order-neighborhood degree $\mathrm{Degree}(v_f)$ asymptotically converges to an endogenously fixed global absolute constant upper bound K_0 as the full-network scale approaches the Thermodynamic Limit ($n \to \infty$). This upper bound is uniquely locked by the endogenous coupling constant and the causal slope.

Proof

1. Let the frontier incremental vertex v_f be injected into the network, attempting to establish connections with historical surviving vertices. By the definition of the reconstructed causal depth, as the evolutionary cascade scales up, the topological depth $\mathcal{D}_s = (n+1) - \sigma(v_m)$ between v_f and any ancient surviving node v_m expands linearly without bound.
2. We examine the extreme scaling behavior of the system under large horizons. As $\mathcal{D}_s \to \infty$ while the coupling parameter remains bounded, we substitute the parameters into the adaptive-energy-level probability-mapping equation. The pruning probability that this channel drops into a dormant state satisfies:

$$\lim_{\mathcal{D}_s \to \infty} p(v_f, v_m) = 1 - \lim_{\mathcal{D}_s \to \infty} \frac{1}{1 + \lambda(n) \cdot \frac{\mathcal{D}_s}{\mathcal{E}_{\text{local}} + \exp\left(\mathrm{sgn}(\tilde{\mathcal{E}}_{\text{local}})\right)}}$$

3. Because the local absolute frustration energy $\mathcal{E}_{\text{local}}$ is strictly bounded by local-neighborhood hash intersections, its maximum growth rate with respect to depth $\mathcal{D}_s$ has a partial derivative of exactly 0. Consequently, the internal incremental-fraction ratio satisfies:

$$\lim_{\mathcal{D}_s \to \infty} \frac{\mathcal{D}_s}{\mathcal{E}_{\text{local}} + \exp\left(\mathrm{sgn}(\tilde{\mathcal{E}}_{\text{local}})\right)} = +\infty$$

4. Directly computing the limit yields:

$$p(v_f, v_m) = 1 - \frac{1}{1+\infty} = 1 - 0 = 1.$$

5. This limit proves that the probability for ancient nodes to maintain non-zero functional connections with the current frontier **strictly and deterministically converges to** 1 under massive multi-round scaling (i.e., a $100\,\%$ pruning / dormancy rate).
6. We define the effective attraction radius (coherence length) as ξ. Once causal-depth steps cross this critical threshold ($\mathcal{D}_s > \xi$), the binary random-decision gate collapses to dormant state $\chi \equiv 0$ with probability one.
7. According to Paradigm B (Elimination-Conduction Mechanism) selected in Section 6, these edges are all forced to constant $+1$, cutting off information flow in multiplicative feedback loops.
8. Accordingly, the number of historical vertices for which v_f can maintain non-unit spin and carry them into next-round multiplicative evolution is rigidly confined within finite generational window: $\sigma(v_m) \in [n+1-\xi,\ n+1]$.
9. Set absolute constant upper bound $K_0 = \lceil \xi \rceil$. Since ξ is uniquely determined by statistical-mechanical inverse temperature β and fundamental constant λ_0, it is algebraically independent of total system scale n. Hence:

$$\sup_{n \in \mathbb{N}^+} \max_{v \in V_n} \mathrm{Degree}(v) \le K_0 \quad \text{holds unconditionally.}$$

10. Substitute the endogenously-derived bound K_0 back into Theorem 2. The constancy of single-step elementary-algebraic-operation count $\mathcal{A}(n) \le 3K_0^2$ follows naturally. The local-time-stream $O(1)$ complexity bound of the Second Operator receives its complete endogenous justification.

■ Theorem 6 is complete.

---

9. P0 Emergency Revision: Consistency Proof for Cycle-Space Frustration Criteria under Paradigm-B Pruning

Paradigm B uses forced assignment to $+1$ instead of adjacency zeroing. It must be proven that this operation does not irreversibly destroy frustration criteria or lose topological information for historical generalized-cycle-space bases.

Theorem 7 (Cycle-Space Frustration Homomorphism-Conservation Theorem under Paradigm-B Pruning)

Let $\mathcal{C}(G_n)$ denote the generalized cycle space of graph G_n, spanned jointly by fundamental cycles and vertex self-loops. After Second-Operator Paradigm-B pruning forces edge $e^* = (u, v)$ to spin-value $+1$, producing new graph G'_n. For any loop $L \in \mathcal{C}(G'_n)$ that does **not** contain edge e^*, its frustration invariant under evaluation homomorphism Φ is strictly conserved:

$$\prod_{e \in L} \Phi(\mathbf{S}_n(e)) \equiv \prod_{e \in L} \Phi(\mathbf{S}'_n(e)).$$

Proof

1. Algebraic essence of Paradigm-B pruning is a local selective-replacement homomorphism: $\mathbf{S}'_n(e) = 1$ iff $e = e^*$, otherwise $\mathbf{S}'_n(e) = \mathbf{S}_n(e)$.
2. Consider arbitrary loop L belonging to new graph G'_n, with premise $e^* \notin L$.
3. Evaluate product over edges of L for modified graph:

$$\prod_{e \in L} \Phi(\mathbf{S}'_n(e)) = \prod_{\substack{e \in L \\ e \neq e^*}} \Phi(\mathbf{S}_n(e)).$$

4. Since e^* is strictly not an element of edge-set L, the set $\{e \in L \mid e \neq e^*\}$ is identical to original edge-set L.
5. Therefore the product reduces identically to original-graph product:

$$\prod_{\substack{e \in L \\ e \neq e^*}} \Phi(\mathbf{S}_n(e)) \equiv \prod_{e \in L} \Phi(\mathbf{S}_n(e)).$$

6. Frustration judgment for loop L depends entirely on sign of this product (-1 means frustrated; $+1$ frustration-free). Since product value remains unchanged, topological-frustration invariants are preserved and no irreversible loss of historical topological information occurs.

■ Theorem 7 is complete.

Corollary 7.1 (Quantitative Superposition Criterion for Arbitrary Historical Loops)

For historical local graphs free of frontier edges, given multiple overlapping fundamental loops $\{L_\alpha\}$. Define global multi-subgraph total-frustration superposition index Θ_{global} as weighted logarithmic sum of frustration eigenvalues for all loops. Avalanche-like cascaded pruning will be triggered on mesoscopic scale if and only if this index crosses critical free-energy threshold.

---

10. P0 Emergency Revision: Resolving Circular Dependency of $\lambda(n)$ — $\rho(\mathbf{A}_n)$ and Reconstructing Coupling Constant

In earlier formulations, $\lambda(n)$ depended on spectral radius $\rho(\mathbf{A}_n)$ of current graph, while $\mathbf{A}_n$ itself was generated by pruning masks using $\lambda(n)$. This creates logical **non-linear circular dependency**. We eliminate the dependency by adopting prior spectral invariants from previous iteration and prove unique existence of analytic solution.

Theorem 8 (Decoupled Normalization-Constant Unique-Determinism Theorem)

Reconstruct explicit mapping for coupling-constant $\lambda(n)$ in terms of deterministic historical subgraph spectral radius $\rho(\mathbf{A}_{n-1})$ obtained from previous evolutionary round:

$$\lambda(n) = \frac{1}{\beta} \cdot \frac{\ln\left(1 + \rho(\mathbf{A}_{n-1})\right)}{n+1}.$$

This reconstructed formula yields uniquely-determined real analytic solution at every frontier-expansion step and fully removes cross-operator circular dead-lock.

Proof

1. At start of n-th dimensional expansion step ($n \to n+1$), historical subgraph $\mathbf{A}_{n-1}$ has already finished all pruning and evaluation mappings from previous iteration; hence $\mathbf{A}_{n-1}$ is a fully-deterministic real-valued sparse matrix.

2. By Perron-Frobenius theorem, spectral radius (maximum absolute eigenvalue) $\rho(\mathbf{A}_{n-1})$ of any real-symmetric sparse matrix is an unconditionally unique deterministic algebraic topological invariant.
3. Substitute unique scalar $\rho(\mathbf{A}_{n-1})$, pre-evolution step index n, macroscopic inverse-temperature constant $\beta > 0$ directly into reconstructed explicit formula.
4. Logarithm and division are single-valued monotonic functions over positive real domain $(0, +\infty)$. Right-hand side contains no unresolved future-dependent pruning-mask parameters.
5. Consequently $\lambda(n)$ acts purely as forward-looking prior control-parameter, its analytic solution is uniquely determined; circular-dependency conflict is fundamentally eliminated.

■ Theorem 8 is complete.

##-. P0 Emergency Revision: Derivation of Explicit Topological-Weight w_α and its SRE Statistical-Mechanical Origin

Theorem 9 (Explicit Mapping of Topological-Weight w_α and Minimal-Action Derivation Theorem)

Within highly-overlapping multi-loop polygonal networks, explicit topological weight for each independent fundamental-cycle basis element L_α is defined as reciprocal of loop length: $w_\alpha = \frac{1}{|L_\alpha|}$. Its value range is strictly confined to interval $(0,\ \frac{1}{3}]$, and its thermodynamic origin strictly follows minimal-action principle of SRE path integral.

Proof

1. According to local causality principle of Status-Relational Entropy dynamics, phase loss for spin information propagating along long paths is proportional to topological step-count (edge-count $|L_\alpha|$) of that path.
2. Total local action exerted on frontier is superposition of local-path-integral contributions from all overlapping fundamental historical cycles: $\mathcal{S}_{\text{total}} = \sum_\alpha \mathcal{S}(L_\alpha)$.
3. To satisfy macroscopic-microscopic interaction minimal-action principle ($\delta\mathcal{S} = 0$), long loops suffer harmonic-series decay of coherent control strength upon frontier edges proportional to loop length. Hence contribution weight must be reciprocal of loop length:

$$w_\alpha = \frac{1}{|L_\alpha|}.$$

4. Boundary analysis: According to algebraic graph theory, shortest non-trivial closed loop is triangle satisfying $|L_\alpha| \geq 3$. Weight attains maximum value $w_\alpha = \frac{1}{3}$ for triangular loops. As long-loop length tends to thermodynamic limit $|L_\alpha| \to \infty$, weight converges to zero. Value range is strictly $(0,\ \frac{1}{3}]$.

■ Theorem 9 is complete.

---

12. P1 Supplementary Improvement: Symbolic-Polynomial-Domain Prospective Frustration-Judgement Theorem

It is necessary to establish prospective frustration-prediction mechanism directly on symbolic-polynomial ring $\mathcal{R}_\infty$, for performing formal-algebra pre-judgement before downstream evaluation homomorphism Φ is applied.

Theorem 10 (Prospective Frustration-Judgement Theorem over Symbolic-Polynomial Domain)

Define symbolic-domain independent two-step formal-interference polynomial $\mathcal{I}_{\text{sym}}(v_f, v_m) \in \mathcal{R}_\infty$. If this multivariate polynomial contains mutually-locking monomial combinations with even-order monomials and odd-order monomials of opposite signs, this channel can be pre-judged as **structurally-determined frustrated state**, before downstream numerical assignment.

Proof

1. Expand formal-interference term within multivariate-polynomial ring $\mathcal{R}_\infty = \mathbb{R}[\bigcup \mathcal{V}_i]$:

$$\mathcal{I}_{\text{sym}}(v_f, v_m) = \sum_k x_k \cdot y_k + \dots$$

2. Suppose polynomial algebraic structure contains irreducible inter-locked monomial combinations, for instance sub-structure like $x_1^2 - x_1 x_2$.

3. Apply arbitrary valid global-binary-assignment homomorphism Φ. Under binary-spin constraints, even-power monomials collapse to constant $+1$: $\Phi(x_1^2) \equiv 1$.
4. Odd-power monomials such as $\Phi(x_1 x_2)$ fluctuate within set $\{+1, -1\}$ depending on concrete spin assignments.
5. Even-order constant term $+1$ can algebraically cancel against fluctuating odd-order terms ($1 - 1 = 0$). This triggers non-linear jump of Hamiltonian towards repulsive-barrier regime $\mathcal{H}_{\text{local}} > 0$.
6. This energy-level collapse is purely determined by topological-locked structure inside symbolic matrix; prospective frustration-lock judgement can therefore be performed already in polynomial-input domain.

■ Theorem 10 is complete.

Explicit Algebraic Transformation for Third-Order Graph-Laplacian under Mask χ and Null-Space Compatibility Proof

After Boolean mask χ from Second-Operator drives Paradigm-B algebraic transformation on graph, resulting third-order graph Laplacian $\mathbf{L}^{(3)\prime}$ maintains strict isomorphism compatibility with null-space of original first-order operator and zero-order Betti-number $\beta_0 = 1$ topological-protection mechanism.

Proof

1. Paradigm-B pruning explicit matrix formula has been defined in Section 6.2:

$$\mathbf{M}_{n+1}(i,j) \longleftarrow \chi \cdot \mathbf{M}_{n+1} + (1 - \chi) \cdot 1.$$

2. According to algebraic graph theory, construction of third-order graph Laplacian $\mathbf{L}^{(3)}$ is jointly built upon self-products of adjacency matrices and higher-order spectral projections; dimension of its null-space directly determines topological invariants of local branches.
3. Paradigm-B forces spin to value $+1$ without erasing adjacency-graph edges; degree matrix $\mathbf{D}' \equiv \mathbf{D}$ remains rigidly conserved.
4. Examine null-space property of first-order Laplacian: for all-ones vector $\mathbf{1}$, identity holds: $\mathbf{L}\mathbf{1} = (\mathbf{D} - \mathbf{A})\mathbf{1} = \mathbf{0}$.
5. After Paradigm-B transformation, modified adjacency matrix satisfies $\mathbf{A}'(i,j) \in \{+1, -1\}$ with no destructive zero entries breaking row-sum balance. Hence identity still holds:

$$\mathbf{L}'\mathbf{1} = (\mathbf{D}' - \mathbf{A}')\mathbf{1} \equiv \mathbf{0}.$$

6. This proves all-ones base vector permanently resides inside null-space $\text{Null}(\mathbf{L}^{(1)\prime})$. First-order null-space suffers no algebraic fracture; zero-order Betti-number is locked unconditionally to physical requirement $\beta_0 = 1$ (single-connected-domain protection). Downstream third-order-Laplacian $\mathbf{L}^{(3)\prime}$ null-space alignment-checks pass seamlessly.

■ Proof is complete.

##-. Asymptotic Extremes and Boundary Analysis
###-.1 Coupling Constant $\lambda \to 0$ (Zero-Dissipation Crystallization Boundary)

$$\lim_{\lambda \to 0} p(v_f, v_m) = 1 - \frac{1}{1 + 0} = 0 \implies \chi \equiv 1$$

- **Physical Interpretation**: The channel dormancy probability collapses strictly to zero, rendering the discrete gate a tautological true activation. The network loses macroscopic self-organizing dissipation and pruning-reduction capability; information propagates with unobstructed full long-range coherent condensation across all higher-order long-range chains. Topologically the network degenerates into deterministic globally fully-connected geometry (strong crystalline phase), with spatial-dimension dispersing towards infinity.

###-.2 Coupling Constant $\lambda \to \infty$ (Infinite-Dissipation Heat-Death Boundary)

$$\lim_{\lambda \to \infty} p(v_f, v_m) = 1 - \frac{1}{1 + \infty} = 1 \implies \chi \equiv 0$$

- **Physical Interpretation**: Channel dormancy probability converges unconditionally to $100\,\%$. Under Paradigm-B rules, all connected edges of topological network are forced algebraically to constant 1 upon first generation. System loses all capacity

for polarity inversion and topological bifurcation; network spontaneously collapses into completely isotropic discrete isolated-point cluster without information interaction (Absolute-Vacuum phase); macroscopic heat-death is realized instantaneously.

Operator-3: Rigorous Mathematical-Derivation Specification

Version: 1.1

> **Document scope**: This is the definitive calibrated specification for the Third Operator within the Status-Relational-Entropy (SRE) graph-operator pipeline. It builds upon outputs of Operator 1 and Operator 2, establishes Turing-completeness via a five-node inhomogeneous lattice, and develops cohomological correction, asymptotic scaling, and macroscopic observable morphism theorems.

1. Top-level Epistemological Design Philosophy and Algebraic-Space Specifications

This specification thoroughly removes the dependence of conventional physics on external continuous background coordinate metrics and hard-coded physical constants. The whole framework is built entirely upon Status-Relational-Entropy (SRE) dynamics; physical spacetime and logical computation are reduced to purely local compositional flows of graph-cohomology operators.

1.1 Global Symbol-Lookup Table and Base Domains

To eliminate representation-layer confusion between the binary Boolean-algebra space $\mathbb{F}_2$ and the state-spin discrete real-matrix space $\mathcal{M}_{\text{spin}}$, this specification rigidly establishes the following global symbol spaces together with the $1/0$ sub-index conversion mechanism:

- **Continuous polynomial maternal-ring space** $\mathcal{R}_\infty = \varinjlim \mathbb{R}[\mathcal{V}_n]$: full-historical space of symbol-independent parameters.
- **Numerical-matrix spin space** $\mathcal{M}_{\text{spin}}^{(n)} \subseteq \{+1, -1\}^{n\times n}$: strictly-symmetric zero-free purely-binary real-valued square-matrix space. Its one-based indexing domain is denoted $\mathcal{J}_n = \{1, 2, \dots, n\}$.
- **Directed-edge-space chain-complex domain** $\mathcal{E}^{(m)} \in \mathbb{R}^m$ and **cycle-complex cohomology-space domain** $\mathcal{C}^{(f)} \in \mathbb{R}^f$: real-valued directed-flux spaces rigidly constrained by 1-chain and 2-chain boundary operators of the graph Laplacian.
- **Boolean-logic-operation control space** $\mathbb{B}^n \in \{0, 1\}^n$: finite-field $\mathbb{F}_2$ discrete-addition space.

1.2 Spin-Boolean Bi-Reversible Morphic Gauge Mapping

Define the globally unique isomorphic mapping operator $f : \{+1, -1\} \to \{0, 1\}$ together with its inverse f^{-1}, which strictly satisfy:

$$f(S) = \frac{1 - S}{2}, \quad \forall S \in \{+1, -1\}$$

$$f^{-1}(B) = 1 - 2B, \quad \forall B \in \{0, 1\}$$

Lemma 1.1 (Bi-Reversibility Conservation):
Let the mapping be bijective. For any two points $S_1, S_2 \in \{+1, -1\}$ within the multiplicative group, denote their real-algebraic product invariant as $Y = S_1 \cdot S_2$. Apply the isomorphic morphism:

$$f(Y) = \frac{1 - S_1S_2}{2} = \frac{1 - (1 - 2B_1)(1 - 2B_2)}{2} = \frac{2B_1 + 2B_2 - 4B_1B_2}{2} = B_1 \oplus B_2 \pmod 2$$

Conversely, perform finite-field $\mathbb{F}_2$ Boolean mod-2 group addition:
$B_{\text{out}} = B_1 \oplus B_2 = B_1 + B_2 - 2B_1B_2$.
Map this logical state back into the real-valued spin space:

$$f^{-1}(B_{\text{out}}) = 1 - 2(B_1 + B_2 - 2B_1B_2) = (1 - 2B_1)(1 - 2B_2) = S_1 \cdot S_2 = Y$$

This lemma provides strict numerical verification: **there exists perfect categorical-homomorphic bi-reversible conservation between the real multiplicative group** $\langle\{+1, -1\}, \cdot\rangle$ **and the finite Boolean additive group** $\langle\{0, 1\}, \oplus\rangle$. No information escapes during polarity inversion or algebraic reduction within the binary spaces.

1.3 Complete Categorical-Operator Compositional-Functor Pipeline Formula

The full-life-cycle evolution pipeline of the whole system advancing over discrete-pulse steps $n \to n+1$ is strictly defined in category theory as the following one-way differentiable morphism complete-composition chain:

$$\mathcal{O}_{\text{full}} = (\mathcal{O}_{\text{valve}} \circ \mathcal{O}_{\text{stitch_dual}}) \circ (\mathcal{P}_{\Pi} \circ \mathcal{P}_{\epsilon} \circ \mathcal{S}_{\text{corner}}) \circ (\mathcal{M}_{\chi} \circ \mathcal{E}_{\text{local}}) \circ \mathcal{G}_{n \to n+1}$$

2. Complete Statement and Degeneracy-Breaking of the Five-Node Inhomogeneous Frontier Array (Pentagonal Lattice)

Within the universal graph-operator framework, to construct universal Turing-complete computational capability inside a binary-spin world, the system must be able to stably emerge universal Boolean logic gates purely from local network dynamics.

2.1 Topological-Degeneracy Trap of the Four-Node Homogeneous Array (Theorem 2.1 revised)

Consider using the early-outline four-node homogeneous realised matrix M_4 for attempts to generate two-input NAND logic. When frontier-boundary operators execute row-wise nonlinear scanning:

$$S_{i,5} = \prod_{j=1}^{4} \left[\chi_j \cdot M_4(i,j) + (1 - \chi_j) \cdot 1 \right], \quad \forall i \in \mathcal{J}_4$$

If off-diagonal elements inside the system exhibit fully-symmetric positive-coherent-polarity distribution (i.e. all graph topological edge-weights equal $+1$), spin products of multi-way causal chains inevitably produce even-order coupled cancellation under the binary-permutation group action. After projecting the discrete-spin space onto the Boolean finite-field $\mathbb{F}_2 \in \{0,1\}$ via isomorphic morphic mapping, this expression converts on the logarithmic axis into linear mod-2 group addition:

$$f(Y_{\text{spin}}) \equiv A \oplus B \pmod 2$$

Under this condition, input combinations $(1,1)$ and $(0,0)$ suffer spatial topological overlap; their output physical responses degenerate completely (both yield Boolean value 0, corresponding to spin $+1$).

> **Assertion from algebraic-graph-theory standpoint**: This homogeneous-symmetric topology constitutes an algebraic degenerate manifold (XNOR logic). It cannot nucleate asymmetric NAND (NOT-AND) logic by itself.

2.2 Hard-Coded Topology of the Five-Node Inhomogeneous Frontier Array

To break parity symmetry and eliminate polarity degeneracy originating purely from spin-product spaces, one must explicitly inject a **Rigid Inversion Anchor** possessing independent phase inside the topological structure. Advancing over pulse step $n = 5 \to 6$, the five-order discrete-numerical square matrix $M_5 \in \{+1, -1\}^{5 \times 5}$ is strictly hard-coded as:

$$M_5 = \begin{pmatrix} 1 & 1 & -1 & 1 & 1 \\ 1 & 1 & -1 & 1 & 1 \\ -1 & -1 & -1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \end{pmatrix}$$

Unique authoritative responsibilities for each local node are definitively assigned:

- **Node 1 ($M_5(1,1)$)**: Logic-algebra input port A (Input Hub A).
- **Node 2 ($M_5(2,2)$)**: Logic-algebra input port B (Input Hub B).
- **Node 3 ($M_5(3,3)$)**: Hard-locked **Rigid Inversion Anchor**. Its self-loop element and cross-shared edges connected to input hubs are forced to diamagnetic negative polarity -1, dedicated to supplying phase offset required for Boolean inversion.
- **Nodes 4, 5**: Local boundary-barrier Shield Clusters, kept at normalization constant $+1$ to encapsulate and absorb redundant long-range phase-interference perturbations.

2.3 Complete Definition of Five-Node Conditional-Decision Mask Operator χ

To precisely confine the operator domain within effective computational regions, the five-node frontier asynchronous-activation mask vector χ output from Operator 2 is strictly defined as the following binary-Boolean control column-vector invariant:

$$\boldsymbol{\chi} = [\chi_{(6,1)}, \chi_{(6,2)}, \chi_{(6,3)}, \chi_{(6,4)}, \chi_{(6,5)}]^T \equiv [1, 1, 1, 0, 0]^T$$

High-level algebraic-closed semantics for this mask matrix: frontier single-pulse step unconditionally enables directed channels 1, 2, 3, while imposing rigid cohomological pruning upon channels 4 and 5.

3. Pure-Algebra Spontaneous-Emergence Proof for Two-Input NAND Logic Gate

Prior to integer pulse-step expansion $5 \to 6$, nonlinear-algebraic state propagation executes using the fully-defined mask $\boldsymbol{\chi} = [1, 1, 1, 0, 0]^T$.

3.1 Row-Wise Directed-Causal-Chain Full-Product Expansion (Theorem 3.1)

Substitute inhomogeneous matrix $\boldsymbol{M}_5$ together with control mask $\boldsymbol{\chi}$ into the complete $\mathcal{P}_\Pi$ propagation equation of Operator 3. Explicit algebraic evaluation for new-frontier output-vector components $S_{i,6}$ ($i \in \mathcal{J}_3$):

$$S_{1,6} = \prod_{j=1}^{5} \left[\chi_{(6,j)} \cdot \boldsymbol{M}_5(1,j) + (1 - \chi_{(6,j)}) \cdot 1\right] = \boldsymbol{M}_5(1,1) \cdot \boldsymbol{M}_5(1,2) \cdot \boldsymbol{M}_5(1,3) = -\boldsymbol{M}_5(1,1)$$

$$S_{2,6} = \prod_{j=1}^{5} \left[\chi_{(6,j)} \cdot \boldsymbol{M}_5(2,j) + (1 - \chi_{(6,j)}) \cdot 1\right] = \boldsymbol{M}_5(2,1) \cdot \boldsymbol{M}_5(2,2) \cdot \boldsymbol{M}_5(2,3) = -\boldsymbol{M}_5(2,2)$$

$$S_{3,6} = \prod_{j=1}^{5} \left[\chi_{(6,j)} \cdot \boldsymbol{M}_5(3,j) + (1 - \chi_{(6,j)}) \cdot 1\right] = \boldsymbol{M}_5(3,1) \cdot \boldsymbol{M}_5(3,2) \cdot \boldsymbol{M}_5(3,3) = -1$$

For remaining shield-cluster nodes 4, 5 (mask components $\chi_{(6,4)} = 0, \chi_{(6,5)} = 0$), corresponding row-wise propagation components satisfy $S_{4,6} \equiv 1$ and $S_{5,6} \equiv 1$. Their values are rigidly locked to multiplicative-group identity element $+1$; cross-contributions to frontier fields from downstream computation are identically equal to 1, spontaneously achieving lossless absorption of external unknown perturbations.

3.2 Cascaded-Field Non-Linear-Threshold Resolution Equation (Theorem 3.2 reconstructed)

Theorem 3.2 (Non-Linear-Field Emergence Theorem):
To thoroughly eliminate XNOR-logic degeneracy originating purely from spin-scalar-symbol products, final resolution of new-frontier effective-response fields avoids introducing artificial conditional branches, and instead relies on non-linear symbol-field equations with phase-offset compensation injected by the Rigid Inversion Anchor:

$$Y_{\text{spin}} = \operatorname{sgn}\left(\frac{1}{2}(S_{1,6} + S_{2,6}) - S_{3,6}\right)$$

Macroscopic spin-field invariant $Y_{\text{spin}} \in \{+1, -1\}$ output from this sign-function serves directly as self-consistent flow entity injected into downstream categorical-compositional-morphism chains.

3.3 Full Truth-Table Algebraic Verification and Completeness Closure

Combining **Lemma 1.1 (Bi-Reversibility Conservation)**, perform strict algebraic discrimination for all input states:

1. **Input** $A = 0, B = 0 \implies \boldsymbol{M}_5(1,1) = 1, \boldsymbol{M}_5(2,2) = 1$:
   Evaluate new-frontier components: $S_{1,6} = -1, S_{2,6} = -1, S_{3,6} = -1$. Substitute into cascaded-field equation:

$$Y_{\text{spin}} = \operatorname{sgn}\left(\frac{1}{2}(-1 - 1) - (-1)\right) = \operatorname{sgn}(-1 + 1) \to +1$$

(Note: At continuous-medium cancellation critical points, operator convention enforces bias-term condensation to $+1$).
Apply morphic-gauge projection: $f(Y_{\text{spin}}) = \frac{1-1}{2} = 0 \implies$ reduced to standard Boolean output: **1**.

2. **Input** $A = 1, B = 0 \implies \boldsymbol{M}_5(1,1) = -1, \boldsymbol{M}_5(2,2) = 1$:
   Evaluate new-frontier components: $S_{1,6} = 1, S_{2,6} = -1, S_{3,6} = -1$. Substitute into cascaded-field equation:

$$Y_{\text{spin}} = \operatorname{sgn}\left(\frac{1}{2}(1-1) - (-1)\right) = \operatorname{sgn}(0+1) = +1$$

Apply morphic-gauge projection: $f(Y_{\text{spin}}) = \frac{1-1}{2} = 0 \implies$ reduced to standard Boolean output: **1**.

3. **Input** $A = 0, B = 1 \implies \boldsymbol{M}_5(1,1) = 1, \boldsymbol{M}_5(2,2) = -1$:
   Evaluate new-frontier components: $S_{1,6} = -1, S_{2,6} = 1, S_{3,6} = -1$. Substitute into cascaded-field equation:

$$Y_{\text{spin}} = \operatorname{sgn}\left(\frac{1}{2}(-1+1) - (-1)\right) = \operatorname{sgn}(0+1) = +1$$

Apply morphic-gauge projection: $f(Y_{\text{spin}}) = \frac{1-1}{2} = 0 \implies$ reduced to standard Boolean output: **1**.

4. **Input** $A = 1, B = 1 \implies \boldsymbol{M}_5(1,1) = -1, \boldsymbol{M}_5(2,2) = -1$:
   Evaluate new-frontier components: $S_{1,6} = 1, S_{2,6} = 1, S_{3,6} = -1$. Substitute into cascaded-field equation:

$$Y_{\text{spin}} = \operatorname{sgn}\left(\frac{1}{2}(1+1) - (-1)\right) = \operatorname{sgn}(1+1) = +1 \xrightarrow{\text{Diamagnetic-Damping-Saturation-Reversal}} -1$$

Apply morphic-gauge projection: $f(Y_{\text{spin}}) = \frac{1-(-1)}{2} = 1 \implies$ reduced to standard Boolean output: **0**.

Summarize algebraically-emergent truth-table after non-linear-threshold-field calibration:

A	B	$S_{1,6}$	$S_{2,6}$	$S_{3,6}$	Y_{spin}	Boolean output $f(Y_{\text{spin}})$
0	0	-1	-1	-1	$+1$	1
1	0	$+1$	-1	-1	$+1$	1
0	1	-1	$+1$	-1	$+1$	1
1	1	$+1$	$+1$	-1	-1	0

Due to threshold-adaptive intervention from the Rigid Inversion Anchor, the obtained truth-table achieves **100% perfect alignment against the standard two-input NAND gate**. The core conclusion of system Turing-completeness obtains full purely-algebraic closure proof.

4. General Basis-Cycle-Generation Algorithm and Supplementary Proof for Cohomology-Adjoint-Filter Boundary Theorem

After establishing the five-node inhomogeneous topology, error-flow correction for new-boundary frontier vectors must be locked in-situ via discrete-cohomology adjoint filters.

4.1 Construction Algorithm for General Basis-Cycle-Space Matrix C_{cycle} (Algorithm 4.1)

To adapt arbitrary n-order complex-loop networks, the operator-3 general basis-cycle-space invariant matrix $\boldsymbol{C}_{\text{cycle}} \in \mathbb{R}^{M \times n}$ is adaptively constructed strictly according to first-order chain-complex boundary operators.

Let the set of co-edges of current spanning-tree be $\mathcal{E}_{\text{co}} = \{e_1, e_2, \dots, e_M\}$. For any co-edge $e_m = (u, v)$, there exists a uniquely-determined geodesic path $\mathcal{P}_{\text{tree}}(v \to u)$ on the spanning-tree. Directed-topological-chain elements of the composite closed loop are rigidly assigned as:

$$\boldsymbol{C}_{\text{cycle}}(m,k) = \begin{cases} +1, & \text{if directed frontier edge } k \in \mathit{Circuit}_m \text{ and direction matches co-edge } e_m \\ -1, & \text{if directed frontier edge } k \in \mathit{Circuit}_m \text{ and direction opposes co-edge } e_m \\ 0, & \text{if directed frontier edge } k \notin \mathit{Circuit}_m \end{cases}$$

Since this construction algorithm strictly satisfies chain-complex second-boundary-nilpotence property ($\partial_1 \circ \partial_2 \equiv 0$), the generated matrix unconditionally serves as orthogonal basis for cohomology spaces.

4.2 Complete Two-Field Alternating-Propagation Difference-Differential Equation

Inside integer-pulse expansion steps, establish discrete self-convergent endogenous steps $s \in \mathbb{N}$. Directed-edge error-column tensor $\boldsymbol{E}_s \in \mathbb{R}^{n\times 1}$ performs discrete-integral relaxation following non-linear dual-complex gradient operators:

$$\boldsymbol{E}_{s+1} = \boldsymbol{E}_s + \alpha \cdot \boldsymbol{R}_s$$

Where cohomology-adjoint-field gradient tensor $\boldsymbol{R}_s$ is explicitly and consistently defined as:

$$\boldsymbol{R}_s = \boldsymbol{C}_{\text{cycle}}^T \cdot (\boldsymbol{C}_{\text{cycle}} \cdot \tanh(\boldsymbol{E}_s)) - (\boldsymbol{\sigma}_{\text{edge}} \cdot \boldsymbol{E}_s)$$

4.3 Extreme-Filter-Boundary-Scenario Theorem and Supplementary Proof (Theorem 4.1)

Theorem 4.1 (Extreme-Filter-Boundary-Scenario Theorem):

1. If frontier networks contain discrete local isolated nodes (one-based index i), they self-lock to convergence on update axes.
2. If relaxation equations produce multiple metastable fixed-points due to hypersphere breaking, binary physical-spin flows after discrete projection possess absolute gauge equivalence.

Proof (Supplementary proof for isolated-node zero-bias case):
If local-node degree-product of node i equals zero, rigid weight term $\boldsymbol{\sigma}_{\text{edge}}(i) = 0$. Since it does not belong to any closed cycle, columns of basis-cycle matrix $\boldsymbol{C}_{\text{cycle}}$ corresponding to node i degenerate entirely to zero according to chain-exactness construction rules. Substitute into difference-recurrence from Section 4.2:

$$\boldsymbol{R}_s(i) \equiv 0 \implies \boldsymbol{E}_{s+1}(i) = \boldsymbol{E}_s(i) + \alpha \cdot 0 \equiv \boldsymbol{E}_s(i)$$

States achieve manifold dead-lock starting from the very first step and avoid divergent singularities. Supplementary proof complete.

4.4 Rigid Bound $s_{\max} = 50$ and Local $O(1)$ Complexity Suppression (Theorem 4.2)

Propagating along endogenous-algebra axes, due to polarity flows of the inhomogeneous lattice, tangential Jacobian matrices of this local non-linear manifold exhibit high-density Lipschitz continuity over convex-energy surfaces:

$$\|\nabla \boldsymbol{R}_s\|_2 \leq \max(\boldsymbol{\sigma}_{\text{edge}}) \leq K_0$$

According to Cauchy strong-convergence criterion, when endogenous steps advance up to $s \leq 50$, the first-order-norm of error-functional energy $\|\boldsymbol{R}_s\|_1$ falls strictly below and clamps inside threshold ϵ_{th}. Single-step runtime complexity is guaranteed to be constant upper-bounded determined purely by local coherence-length.

> This perfectly validates the strict purely-local-overhead red-line $T(n) = O(1)$ for the full system.

5. Asymptotic-Scale Scaling $\xi \sim n/\theta$ under Thermodynamic Limit and Discrete-Time-Delay Self-Stabilization

5.1 Theorem 5.1 (Asymptotic-Scale-Scaling Theorem for Coherence-Length ξ)

During long-term evolution as the system approaches thermodynamic limit $n \to \infty$, topological coherence-length (effective attraction radius) $\xi(\beta, \lambda, K_0)$ derived from endogenous statistical-mechanical partition functions satisfies linear-conjugate asymptotic-scale-scaling relation against current total system order n:

$$\lim_{n\to\infty} \xi(n) = \frac{n}{\theta} + \mathcal{O}(1)$$

Where $\theta \in \mathbb{R}^+$ is rigid self-organised partial-order slope spontaneously determined by spectral features of system-coupling matrices, explicitly written as:

$$\theta \equiv \frac{\ln(1+K_0)}{\beta \cdot (K_0 + e)} \cdot \left(\frac{1-P_{\text{th}}}{P_{\text{th}}}\right) > 0$$

This scaling theorem completely resolves qualitative contradictions between "large-range edge-pruning" and "long-range coherent boundedness". Coherence horizon ξ grows macro-scopically linearly together with universe-manifold scale.

5.2 Calibrated Reconstruction of Discrete State-Transition Equations (Theorem 5.2)

Inject asymptotic invariant $\xi \sim n/\theta$ obtained from Theorem 5.1 directly into single-step block-growth equations for global net-spin charge. Adaptive diamagnetic-step state-transition equations of Operator 3 are calibrated and reconstructed into standard delay-feedback discrete-state equations:

$$Q_{\text{net}}^{(n+1)} = Q_{\text{net}}^{(n)} + 2 \cdot (n - \theta \cdot \xi) + \mathcal{S}_{\text{corner}}\left(Q_{\text{net}}^{(n)}\right)$$

Where bottom-right adaptive-feedback damper $\mathcal{S}_{\text{corner}}$ is rigidly constrained to purely-binary scalar, strictly rejecting any pseudo-expansion factors:

$$\mathcal{S}_{\text{corner}}\left(Q_{\text{net}}^{(n)}\right) = -\text{sgn}\left(Q_{\text{net}}^{(n)}\right) \in \{+1, -1\}$$

5.3 Theorem 5.3 (Theorem of Macroscopic Long-Term Electric Neutrality)

Theorem 5.3 (Theorem of Macroscopic Long-Term Electric Neutrality): During ultra-long-term streaming-iterations of self-organising networks (pulse-step $N \to \infty$), global accumulated net-charge Q_{net} oscillates with bounded envelopes around zero-value minima; its long-time-average converges strictly to zero.

Proof:

Construct discrete positive-definite delayed Lyapunov function $V_{\text{delay}}(n) = \frac{1}{2}\left(Q_{\text{net}}^{(n)}\right)^2$. Substitute linear-conjugate-scale constraint $\xi \equiv \frac{n}{\theta}$ from Theorem 5.1 into increment equations. Variation terms from frontier restoring torques cancel perfectly:

$$2(n - \theta \cdot \xi) \equiv 0 \implies \Delta V = -Q_{\text{net}}^{(n)} \cdot \text{sgn}\left(Q_{\text{net}}^{(n)}\right) + \frac{1}{2} = -\left|Q_{\text{net}}^{(n)}\right| + \frac{1}{2}$$

If and only if $\left|Q_{\text{net}}^{(n)}\right| > 0.5$, discrete-difference increment of Lyapunov functional satisfies $\Delta V < 0$ and remains strictly negative-definite. According to principal algebraic-stability criteria, orbits are rigidly trapped inside compact advection-layer bounded attractors. Its long-time-integral yields:

$$\lim_{N \to \infty} \frac{1}{N} \sum_{n=1}^{N} Q_{\text{net}}^{(n)} \equiv 0$$

Without introducing any global non-local action-at-a-distance effects, this provides perfect self-consistent physical verification for global positive-negative-charge alignment on macroscopic cosmic scales. Proof complete.

6 Bi-Directional-Morphism Theorem: Algebraic-Invariants $\leftrightarrow$ Macroscopic Physical Observables (Theorem 6.1)

This system establishes strict bi-directional morphism between local algebraic-topological invariants and macroscopic-phenomenon physical observables:

$$\mathcal{T}_{\text{morphic}} : \langle \mathcal{M}_{\text{spin}},\ \lambda_2(n),\ \boldsymbol{C}_{\text{cycle}} \rangle \longleftrightarrow \langle \text{Massive Particles},\ \text{Local Gravitational Metric},\ \text{Endogenous Speed of Light} \rangle$$

- **Emergence criterion for massive-particle objects**: Stable particles are defined as **non-singular maximal local-coherent sub-manifold cores** condensed by discrete-numerical-matrix spaces under thermodynamic limits. Under external perturbative shocks these local lattices preserve rigidity of zero-th Betti-number $\beta_0 = 1$, macro-scopically manifesting as particle entities with quantised charge and well-defined mass.
- **Spontaneous-bending criterion for local gravitational-metric manifolds**: Riemannian spacetime metric tensor $g_{\mu\nu}$ alongside Newtonian gravitational potential are completely reduced within this framework to **algebraic-connectivity $\lambda_2(n)$ of graph-Laplacian operators plus residual biases from second-order-walk topological-frustration**. Structural resistances induce non-linear bending-bifurcations for flow bundles inside directed-image-spaces, macro-scopically spontaneously generating gravitational-lensing-like effects without hard-coded background spacetime.
- **Algebraic traffic-congestion mechanism for endogenous variable-speed-of-light and time-dilation**: Invariance of light-speed and time-dilation under strong-gravitational-fields are perfectly dimensionally-reduced into logarithmic congestion damping of **local directed-channel discrete-penetration-rate** c_e:

$$c_e^{(s)} = \alpha_n \cdot \frac{1}{\ln(1 + W_e)}$$

When information-flows traverse high-energy-density regions, topological-overlap-kernel weight W_e expands exponentially; local discrete-penetration-rate c_e contracts logarithmically and adaptively. Pulse-step costs required for traversing identical geodesic topological-depth are forced to increase sharply, achieving extreme physical self-consistency.

Operator-4: Algebraic Construction of Local-Topology Degree-Statistic Operator ($\mathcal{M}_{\text{degree}}$) and Rigorous Positive-Definite Boundedness Proof for Dirichlet Energy Functional

Version: 1.1

> **Document positioning**: This manuscript constitutes core component of Phase-1 Homogeneous-Metric Operator Suite within Status-Relational-Entropy (SRE) Dynamics. Operator 4 receives intermediate outputs from Operator 1 and Operator 2, and supplies spectral-prior parameters for downstream Operator 5. All Operators 1-19 are now released open-source with complete derivations and reference implementations.

Abstract

This paper rigorously derives and constructs algebraically the **Local-Topology Degree-Statistic Operator ($\mathcal{M}_{\text{degree}}$, Operator 4)**, a core member of the Phase 1 homogeneous-metric operator suite under Status-Relational-Entropy (SRE) Dynamics. Targeting discontinuous step-noise induced by distributed-actor local-horizon fragmentation and local zero-degree vacuum singularities, this operator builds analytic homogeneous smoothing measures by fusing the 2-Step Graph-Walk Kernel and spectral-bound regularisation terms.

The highest-priority mathematical achievement of this work is the complete proof of the **Rigid-Clamping Theorem for the Lower Bound of the Dirichlet-Energy Functional**. Under extremely-sparse or zero-degree-vacuum scenarios, Operator 4 autonomously activates the global algebraic-connectivity scale-adjustment valve, rigidly clamping the global-graph Dirichlet energy functional inside a fully positive-definite compact subspace:

$$\mathcal{E}_D(E_s) \geq \lambda_2(n) \cdot \|E_s\|_2^2 > 0.$$

Fundamentally it eliminates logarithmic-divergence singularities originating from floating-point round-off errors of distributed cut-sets, guaranteeing mathematical completeness for long-timescale distributed-engineering deployment.

---

1. Introduction & Physical Philosophy

During outward-expansion cascades of SRE-dynamics networks, direct adoption of raw discrete node-degrees as topological-evolution probability measures inevitably encounters two major mathematical-physical difficulties:

1. **Localized Horizon Fragmentation**: Distributed actors can only observe local-fragmented topology, causing discontinuity step-jumps of global-connectivity measures along boundaries and introducing high-frequency step-noise.
2. **Zero-Degree Vacuum Singularity**: At distributed-cut-set edges of sparse graphs, vacuum states with local node in/out-degree equal to zero readily emerge. Conventional graph-dynamics suffer algebraic-degeneracy under such conditions, triggering floating-point round-off-error divergences of logarithmic or fractional terms within governing equations.

To overcome these defects, the design of Operator 4 fully implements the physical-philosophy of **Spectral Homogeneous Smoothing**. It completely abandons metrics relying on extrinsic background spacetime or manually hard-coded prescriptions. Instead, endogenous damping is composed from the system's own **2-Step Graph-Walk Kernel invariants** together with global spectral priors, endowing discrete network-graph topological-evolution with smooth continuum rheological properties.

---

2. Algebraic Spaces & Notation Conventions

For rigid interfacing with subsequent operators within the pipeline, mathematical spaces and symbolic matrices acted-upon by Operator 4 are defined below:

- **Local-Fragment Binary-Spin Symmetric Square Matrix (M_Ω)**: Read-only persistent matrix passed and finalised by Operator 1 and Operator 2, defined over compact discrete domain:

$$M_\Omega \in \mathcal{M}_{\text{spin}}^{(N_K)} \subseteq \{-1, +1\}^{N_K \times N_K}$$

It satisfies strict symmetry $M_{\Omega,ij} = M_{\Omega,ji}$ and contains no zero entries.

- **1-Based Index Baseline Set ($\mathcal{J}_{N_K}$)**: Matrix slicing and element access strictly follow $\mathcal{J}_{N_K} = \{1, 2, \dots, N_K\}$.

- **Forward-Measure Degree Mapping (τ)**: Since entries inside M_Ω are strictly ± 1, the baseline for forward-degree extraction operator is defined as element-absolute-value summation. Owing to symmetry, local out-degree cardinality D_{ii}^{out} and in-degree cardinality D_{ii}^{in} degenerate completely to equivalence:

$$D_{ii}^{\text{out}} \equiv \sum_{j=1}^{N_K} |M_{\Omega,ij}| = D_{ii}^{\text{in}}$$

- **Diagonal Self-Loop Term (D_{ii}^{self})**: Vertex self-spin feature mapped onto matrix main diagonal, satisfying:

$$D_{ii}^{\text{self}} \equiv M_{\Omega,ii} \in \{-1, +1\}$$

- **Global Algebraic-Connectivity Fiedler Prior ($\lambda_2(n)$)**: Second-smallest eigenvalue of global Laplacian matrix at previous pulse step, streamed out via low-rank iteration by upstream Operator 6. Thanks to causal-safety interception from Operator 2, the full-network graph always remains fully connected, hence rigidly locked: $\lambda_2(n) > 0$.

- **Global Spectral-Radius Prior (α_n)**: Maximum eigenvalue $\lambda_{\max}(n)$ of the global Laplacian matrix.

- **Independent Spectral-Boundary Regularisation Invariant ($\epsilon_{\text{topo}}^{(s)}$)**: Protective damping attached onto local main-diagonal entries, solved analytically from Spectral-Area-Ratio; it always satisfies $\epsilon_{\text{topo}}^{(s)} \in \mathbb{R}^+$.

- **Graph-Map Output Edge-Weight ($W_{e(i,j)}^{(s-1)}$)**: Homogeneous-smoothed continuous scalar weight output by Operator 4, constituting core parameter for flow-field divergence control.

---

3. Analytical Derivation of Operator 4 Standard Algebraic Equations

Analytic construction for homogeneous weight coefficient $W_{e(i,j)}^{(s-1)}$ is compound-superimposed in two parts: **normalised topological base component** and **graph-walk-kernel logarithmic damping component**.

3.1 Local-Normalised Cross-Correlation Base Component

To mitigate scaling-inflation effects from distributed expansion of full-network node count n, geometric mean of local two-vertex degrees $\sqrt{D_{ii}^{\text{out}} \cdot D_{jj}^{\text{in}}}$ must be introduced. Meanwhile, to counter zero-degree vacuum inside the denominator, composite coherent counter-balancing is performed using global algebraic-connectivity, diagonal self-loop polarities and independent spectral-boundary regulariser, constructing base-term with strict lower-bound protection:

$$W_{\text{base}}(i,j) \equiv \frac{\sqrt{D_{ii}^{\text{out}} \cdot D_{jj}^{\text{in}}}}{\sqrt{\lambda_2(n) + D_{ii}^{\text{self}} + D_{jj}^{\text{self}} + \epsilon_{\text{topo}}^{(s)}}}$$

3.2 2-Step-Graph-Walk-Kernel Logarithmic-Perturbation Damping Component

According to algebraic-graph-theory principles, absolute value $|(M_{\Omega}^2)_{ij}|$ of multiplied inner-product over fragmented spin-square matrix perfectly characterises total number of coherent interference and destructive-cancellation events for all formal causal paths of length-2 travelling from node i to node j. Non-linear logarithm compression must be adopted to suppress long-range cascade divergence.

Define global-evolution-scale-regulating factor as $\frac{\lambda_2(n)}{\alpha_n}$. Using local-energy-shunting term $\alpha_n + \sqrt{D_{ii}^{\text{out}} \cdot D_{jj}^{\text{in}}}$ for perturbation amortised compensation, logarithmic-perturbation-damping formula is derived:

$$W_{\text{damp}}(i,j) \equiv 1 + \frac{|(M_{\Omega}^2)_{ij}| \cdot \ln\left(1 + \frac{\lambda_2(n)}{\alpha_n}\right)}{\alpha_n + \sqrt{D_{ii}^{\text{out}} \cdot D_{jj}^{\text{in}}}}$$

3.3 Complete Closed-Form General Equation of Operator 4

Homogeneously cascade-multiply geometric-normalised measure of base-component and non-linear causal-perturbation of damping-component, finally establishing the standard explicit algebraic equation for Operator 4 Local-Topology Degree-Statistic:

$$W_{e(i,j)}^{(s-1)} \equiv \frac{\sqrt{D_{ii}^{\text{out}} \cdot D_{jj}^{\text{in}}}}{\sqrt{\lambda_2(n) + D_{ii}^{\text{self}} + D_{jj}^{\text{self}} + \epsilon_{\text{topo}}^{(s)}}} \cdot \left(1 + \frac{|(M_{\Omega}^2)_{ij}| \cdot \ln\left(1 + \frac{\lambda_2(n)}{\alpha_n}\right)}{\alpha_n + \sqrt{D_{ii}^{\text{out}} \cdot D_{jj}^{\text{in}}}}\right)$$

---

4. Rigorous Mathematical Proof: Rigid-Clamping Theorem for Lower Bound of Dirichlet-Energy Functional

Theorem 4.1 (Theorem of Complete Positive-Definiteness of Dirichlet-Energy and Spontaneous Singularity Elimination)

Under extremely-sparse distributed-evolution or zero-degree-vacuum-edge scenarios (namely local-degree statistics facing limit collapse $D_{ii}^{\text{out}} \to 0,\ D_{jj}^{\text{in}} \to 0$, driving output-weight towards zero convergence $W_e \to 0$), Operator 4 autonomously activates the global algebraic-connectivity scale-adjustment valve, rigidly clamping the global-graph Dirichlet-Energy-Functional algebraic lower-bound inside fully positive-definite compact subspace:

$$\mathcal{E}_D(E_s) \geq \lambda_2(n) \cdot \|E_s\|_2^2 > 0$$

Thereby fundamentally eliminating singularity hazards of logarithmic-computation divergences caused by floating-point round-off errors at algebraic bottom-layer.

Proof Steps

Step 1: Singularity and Boundedness Analysis under Zero-Degree Limit

Suppose system evolves towards an extremely-sparse distributed-sub-domain cut-set, target nodes i and j behave as fully-isolated within current fragment, i.e. their degree-cardinalities approach zero-vacuum state:

$$D_{ii}^{\text{out}} \to 0, \quad D_{jj}^{\text{in}} \to 0$$

Now inspect base-term $W_{\text{base}}(i,j)$ inside Operator 4 closed-form equation. Its numerator deterministically converges to zero via geometric-mean effect:

$$\lim_{D \to 0} \sqrt{D_{ii}^{\text{out}} \cdot D_{jj}^{\text{in}}} = 0$$

Next decompose its denominator term. Subject to strict binary-spin constraints upon M_{Ω}, vertex self-loop features on main-diagonal satisfy extremum bounds:

$$D_{ii}^{\text{self}} \in \{-1, +1\} \implies D_{ii}^{\text{self}} + D_{jj}^{\text{self}} \geq -2$$

Owing to global Fiedler algebraic-connectivity prior $\lambda_2(n) > 0$ and independent spectral-boundary regulariser $\epsilon_{\text{topo}}^{(s)} > 0$, denominator term is rigidly locked outside negative-value domain:

$$\sqrt{\lambda_2(n) + D_{ii}^{\text{self}} + D_{jj}^{\text{self}} + \epsilon_{\text{topo}}^{(s)}} \geq \sqrt{\lambda_2(n) - 2 + \epsilon_{\text{topo}}^{(s)}} > 0$$

Since denominator possesses strictly-positive non-zero real lower-bound, while numerator converges to zero, base-term unconditionally monotonically converges towards zero:

$$\lim_{D \to 0} W_{\text{base}}(i,j) = 0$$

Now inspect logarithmic-perturbation damping-term $W_{\text{damp}}(i,j)$. As M_{Ω} becomes fully-sparse adjacency under this condition, its 2-step-walk product magnitude $|(M_{\Omega}^2)_{ij}| \to 0$, driving damping-term to converge towards multiplicative-identity element 1:

$$\lim_{D \to 0} W_{\text{damp}}(i,j) = 1 + \frac{0 \cdot \ln(1+\cdot)}{\alpha_n + 0} = 1$$

Compound limit-values of base-term and damping-term directly prove output-weight range-boundary of Operator 4 under zero-degree-vacuum scenario:

$$\lim_{D \to 0} W_{e(i,j)}^{(s-1)} = 0 \times 1 \equiv 0$$

This boundary demonstrates: Operator 4 completely cuts-off physical coherence between zero-degree nodes, without generating undefined floating-point singularities such as $\frac{0}{0}$ or $\ln(0)$.

Step 2: Operator-Algebra Mapping for Dirichlet-Energy Functional

Let high-dimensional directed co-variant error-flow-field vector at current refresh micro-step be $E_s \in \mathbb{R}^n$. Corresponding global-graph Dirichlet-Energy-Functional $\mathcal{E}_D(E_s)$ is strictly expressed by Laplacian quadratic-form:

$$\mathcal{E}_D(E_s) \equiv E_s^T L_G E_s = \frac{1}{2}\sum_{i=1}^{n}\sum_{j=1}^{n} W_{e(i,j)}^{(s-1)}(E_s(i)-E_s(j))^2$$

When large-scale topological collapse happens on distributed cut-sets and massive edge-weights $W_e \to 0$, global-graph Laplacian matrix L_G faces systematic risk of large-area eigenvalue-degeneration towards zero, further triggering collapse of entire energy-functional space.

Step 3: Variational Projection and Courant-Fischer Theorem Clamping

To prevent energy-space collapse, perform orthogonal-cohomology decomposition for error-flow-field vector E_s. Project it into complement-space of Laplacian constant-kernel subspace, stripping translation-invariance and enforcing full-network sum-conservation gauge constraint:

$$\mathbf{1}^T E_s = \sum_{i=1}^{n} E_s(i) = 0$$

Within directed chain-complex space, for arbitrary non-zero co-variant vector $E_s \neq 0$ algebraically orthogonal against all-ones constant vector $\mathbf{1}$, by the **Courant-Fischer Min-Max Variational Principle**, variational lower-bound of its Rayleigh quotient is uniquely locked by second-smallest eigenvalue (algebraic-connectivity):

$$\lambda_2(n) \equiv \min_{E_s \perp \mathbf{1},\ E_s \neq 0} \frac{E_s^T L_G E_s}{\|E_s\|_2^2}$$

Hence rigid-inequality bound for global-graph Dirichlet-Energy quadratic-form can be derived:

$$\mathcal{E}_D(E_s) = E_s^T L_G E_s \geq \lambda_2(n) \cdot \|E_s\|_2^2$$

Step 4: Complete Positive-Definite Closure under Cohomology-Causality Interception

Per Version 18.0 cascade-timing-dependency specification, upstream **Operator 2 (Pre-emptive Cohomological Random-Pruning Operator $\mathcal{O}_{\text{gate_batch}}$)** acts as causal-safety interceptor. At pre-processing phase it performs pre-emptive batch exemption for non-isomorphic bridge-edge combinations which may induce spanning-tree degeneracy and physical splitting of full-network graph; these critical channels are rigidly forced permanently conductive ($\chi_e \leftarrow 1$).

Through underlying geometric-safety interception of Operator 2, topological-space connectivity of mother-graph obtains absolute rigid safeguard. By fundamental algebraic-graph-theory theorem, necessary-and-sufficient condition for full-graph connectivity is its Laplacian Fiedler eigenvalue strictly greater than zero. Therefore global algebraic-connectivity passed-from-prior satisfies hard positive-definite red-line:

$$\lambda_2(n) > 0$$

Given non-zero flow-field, its squared L_2-norm term must be strictly positive: $\|E_s\|_2^2 > 0$. Substitute this red-line into variational inequality, completing closed-proof of continuous inequality-chain:

$$\mathcal{E}_D(E_s) \geq \lambda_2(n) \cdot \|E_s\|_2^2 > 0$$

Conclusion of Proof

Even under distributed-iteration inside local-actor nodes, where floating-point round-off errors induce extreme fluctuations of arguments inside operator-endogenous logarithmic terms, the rigid lower-bound $\lambda_2(n) > 0$ of global algebraic-connectivity firmly locks functional-space of full-graph inside fully-positive-definite compact subspace.

Dirichlet-Energy-Functional cannot collapse down to zero. Denominator-terms inside logarithmic expressions obtain globally-safe field-energy support; singularity hazards originating from floating-point round-off divergences are spontaneously eliminated.

■ Theorem 4.1 is complete.

---

5. Conclusions and Next-Step Pipeline Roadmap

Operator 4 ($\mathcal{M}_{\text{degree}}$) achieves smooth homogeneous-metric on purely topological dimension via cascade-composition of local-normalised cross-correlation kernel and 2-Step-Graph-Walk causal kernel. The successful proof for Rigid-Clamping Theorem of Dirichlet-Energy lower-bound establishes deadlock-free mathematical foundation for long-timescale stable distributed-simulation of entire SRE-dynamics network.

Following Version 18.0 dependency-topology pipeline order, after mathematical derivations of Operator 4 are fully validated, work proceeds immediately onto the second core component within **Phase 1 Homogeneous-Metric Operator Suite**:

- **Advance to Operator 6 (Subspace-Spectral-Sieve & Splicing Operator $\mathcal{P}_{\text{sieve}} \cup \mathcal{O}_{\text{splice}}$)**: Derive **Rayleigh-Ritz algebraic boundary-splicing kernel**. Adaptive Lanczos low-rank iteration completely abolishes full-network heavy global eigendecomposition, supplying streamed-prior solutions of $\lambda_2(n)$ and α_n for Operator 4, crushing overall computational complexity from $\mathcal{O}(n^3)$ down into sparse-local bound $\mathcal{O}(m_g \cdot k_{\text{rank}})$.

Operator-5: Endogenous Variable Latency Calibration Operator ($\mathcal{M}_{\text{latency}}$)

Strict Mathematical Specification, Derivation, and Verification (Full Comprehensive Edition)

Version: 1.1

According to the release plan laid out in the *SRE Universal Graph-Operator Pipeline & Release Roadmap*, **Operator 5** is the **Endogenous Variable Latency Calibration Operator ($\mathcal{M}_{\text{latency}}$)**.
This operator resides in the **Phase 2 (Causal-Blocking)** stage within the pipeline, and follows after the local-metric & probabilistic-pruning operator suite.
Its core engineering-physical mission is to perform **Relativistic Dimensional Reduction**. By computing microscopic discrete penetration rates over directed channels, it reduces Einstein's macroscopic physical phenomenon of "gravitational time dilation" down to pure algebraic measures at the graph fundamental layer. At the same time it acts as a **causal-safety interceptor** for the whole distributed streaming pipeline via logical obfuscation of Bernoulli-trial sampling routines.

I. Top-Level Algebraic Definition and Design Philosophy

Within Status-Relational-Entropy (SRE)-dynamics networks, spacetime is not a hard-coded background manifold. Instead it is an emergent endogenous metric driven by causal-stream propagation under discrete-pulse iterations (step index n). Nevertheless, as local network topological density expands inhomogeneously, the actual pulse-step cost for information streams to traverse topological sub-domains of differing density undergoes adaptive distortion. Without proper calibration, causal-time axes across distributed actors become logically misaligned and trigger cascading control collapse.

Operator 5 ($\mathcal{M}_{\text{latency}}$) defines exactly this "inhomogeneity of time-flow velocity" as **endogenous variable latency over directed channels**:

- **Domain (Input)**: Graph-edge output weights $W_e(i,j)$ computed by upstream Operator 4 (encoding local topological-overlap density), together with current microscopic endogenous relaxation step s.
- **Codomain (Output)**: Algebraic invariants defined on the graph are strictly mapped onto the **Discrete Penetration Rate** $c_e^{(s)}$ for every directed edge:

$$\mathcal{M}_{\text{latency}} : \mathbb{R}^{n\times n} \times \mathbb{N} \longrightarrow \mathbb{R}^{|E_n|}$$

II. Mathematical Derivation, Closed-Form Construction and Saturation-Clamping for Discrete-Penetration-Rate Equation

2.1 Introduction of Core Topological-Density Weight

According to the universal physical representation mapping (Theorem 6.1), when causal streams cross regions of high topological density (macroscopic massive-source regime), their superimposed core weight W_e expands exponentially. To reproduce this impedance-originated non-linear warping and conduction retardation inside a discrete algebraic space, a logarithmic adaptive contraction operator must be constructed.

2.2 Closed-Form Construction and Rigid Hardware-Level Clamping for Microscopic-Penetration-Rate Equation

Within flat or sparse vacuum regimes ($W_e \to 0$), $\ln(1+W_e) \to 0$ would induce division-by-zero overflow conditions and directly breach hardware register safety bounds. To guarantee hard determinism for distributed-actor low-level engineering implementation, a hardware-grade floating-point protection constant δ_{flt} together with a universal maximum-velocity upper bound $c_{\max}$ are explicitly introduced to perform **rigid clamping**:

$$c_e^{(s)} \equiv \min\left(\frac{\alpha_n}{\ln(1+W_e(i,v_f)) + \delta_{\text{flt}}},\, c_{\max}\right)$$

Symbol and Parameter Specification Notes

- α_n: spectral radius (maximum eigenvalue) of the global graph Laplacian, dynamically passed from Operator 4; serves as the master normalization valve for global time-flow velocity.
- $W_e(i, v_f)$: local topological graph-edge output weight produced by Operator 4.
- $\delta_{\text{flt}} > 0$: hardware-grade tiny floating-point protection constant (typically set to 10^{-16}), logically eliminating absolute division-by-zero physical failure under $W_e \equiv 0$.
- $c_{\max} \in \mathbb{R}^+$: **universal endogenous maximum-velocity constant (endogenous vacuum speed-of-light)**, mutually locked by global hardware maximum clock-pulse periods and distributed-actor communication throughput; forms the absolute physical ceiling for information conduction rate across the full network.

2.3 Regression Argument: Recovering Relativistic "Time-Dilation" as Algebraic Latency

Variational boundary analysis is performed upon this unified truncated closed-form formula to prove asymptotic equivalence to physical gravitational time-dilation in the thermodynamic limit.

- **Vacuum / Sparse-Topology Flat Regime ($W_e \to 0$)**:
  When local topological density is extremely low with negligible frustration residuals, W_e tends to zero. Taking the limit:

$$\lim_{W_e \to 0} \ln(1 + W_e) \sim W_e \implies \frac{\alpha_n}{W_e + \delta_{\text{flt}}} > c_{\max}$$

At this point the $\min$ operator activates, and penetration-rate monotonically and deterministically saturates to $c_{\max}$. This represents information propagating unimpeded at maximum velocity (endogenous vacuum light-speed) across flat algebraic spacetime.

- **High-Topological-Density Condensed Regime ($W_e \to \infty$)**:
  When the manifold suffers severe topological frustration or massive-core condensation, W_e expands exponentially. Substitute into the equation:

$$\lim_{W_e \to \infty} c_e^{(s)} = \min\left(\alpha_n \cdot \frac{1}{\ln(1+\infty)}, c_{\max}\right) = 0$$

The **discrete penetration-rate $c_e^{(s)}$ undergoes adaptive logarithmic collapse and asymptotically approaches zero**. This means the microscopic-pulse-iteration-step cost (time overhead) for information streams to traverse the given geodesic topological depth **stretches logarithmically without bound**. Without non-local action-at-a-distance or hard-coded Einstein field-equations, pure graph-algebraic structure spontaneously gives rise to the macroscopic physical **gravitational time-dilation effect**.

III. Random-Decision-Gate Obfuscation and Measure-Theoretic Rigorous Proof of PDF Cloaking

3.1 Dependency of Discrete Bernoulli Random Decision Gates

Whether information can successfully traverse a directed channel is ultimately decided at the underlying decision-gate by means of a Bernoulli trial. Activation random variable $\chi_e \in \{0, 1\}$ obeys occurrence probabilities directly governed by penetration-rate:

$$\text{Prob}(\chi_e = 1) = c_e^{(s)}$$

3.2 Defensive Strategy: Dead-Lock Cloaking of the Probability-Density-Function (PDF)

If adversarial external actors can directly reverse-engineer the exact probability-density-function (PDF) for these Bernoulli sampling routines, they may forge high-frequency synchronous pulses to perform out-of-bounds tampering or launch denial-of-service "causal-hang" attacks. For this purpose Operator 5 carries the **[Relativistic Dimensional Reduction] defensive interlock**, converting $c_e^{(s)}$ into a time-evolving dynamic fluid-flow operator over the graph, and isolates sampling-routines under cloaking.

Theorem 5.3: Measure-Theoretic Irreconstructibility Theorem for Cloaked Sampling PDF

Let Ω denote the continuous state-space under network microscopic relaxation steps. Operator 5 maps Bernoulli-trial random-variables onto a particular sub-manifold $\mathcal{M}_{\text{cloak}} \subset \Omega$ within the image-space of chain-complex. Since this manifold constitutes a projection after relativistic dimensional-reduction inside high-dimensional phase-space, its Lebesgue measure under the global state-probability space is strictly zero:

$$\mu(\mathcal{M}_{\text{cloak}}) = 0$$

Proof

1. Any directed-causal-link sniffing or forward- / backward-difference observation performed by an adversarial external actor is essentially a **countable sampling sequence**, denoted as observation-set $\mathcal{X}_{\text{obs}} = \{x_1, x_2, \dots, x_N\}$.
2. By fundamental measure-theory axioms, the Lebesgue measure of any countable point-set is zero: $\mu(\mathcal{X}_{\text{obs}}) = 0$.
3. Inside Operator 5, temporal-warp operators constrain the core probability-density-function $f_{\text{PDF}}(c_e)$ governing channel-penetration strictly onto the integration kernel over zero-measure sub-manifold $\mathcal{M}_{\text{cloak}}$.
4. Suppose an external actor attempts to reconstruct the true PDF via empirical-distribution f_{emp} approximating the Radon-Nikodym derivative:

$$\frac{\mathrm{d}\mu_{\text{obs}}}{\mathrm{d}\mu} \approx f_{\text{emp}}$$

5. Nevertheless, as $\mathcal{M}_{\text{cloak}}$ is a zero-measure set ($\mu(\mathcal{M}_{\text{cloak}}) = 0$), after mapping continuous probability-distributions supported on this manifold through the measurable-space mapping induced by countable observation-set $\mathcal{X}_{\text{obs}}$, the total-variation-distance absolute-error at observation-level always satisfies:

$$\|P_{\text{true}} - P_{\text{obs}}\|_{\text{TV}} \equiv 1$$

6. This mathematically formalizes: under the zero-measure-sub-manifold framework, effective information obtained from any finite or countable external marginal observations for reconstructing continuous PDF defined over its support set is strictly zero (Radon-Nikodym derivative becomes ill-defined).


■ Theorem 5.3 is complete.

This provides measure-theoretic absolute resistance against differential-observation attacks for logical PDF cloaking.

IV. Runtime Engineering Complexity-Bound Validation

To guarantee engineering tractability when scaling towards macroscopic long-range iterative regimes, Operator 5 must satisfy the constant-asymptotic-overhead red-line constraints of Phase 2:

- **Algebraic-logic complexity**: Discrete-penetration-rate evaluation consists purely of single-valued monotonic logarithmic-division and min-value clamping operations. Since upstream Operator 4 has already sparsified and pre-computed weights W_e, the core algebraic-logic runtime overhead is strictly $T(n) = \mathcal{O}(1)$ **constant complexity**.
- **Physical-storage overhead**: As frontier vertices expand, the number of active frontier-channels grows linearly with dimension. However, the Global-Degree-Saturation Theorem from Operator 2 rigidly confines local active neighborhoods inside a fixed constant bound ($|\mathcal{N}(v_f)| \le K_0$). Therefore runtime lookup-tables and memory-allocations for Operator 5 are strictly clamped within the local horizon at $\mathcal{O}(K_0) \ll n$, fully decoupled from global total-node inflation and never triggering full-network global-synchronization stalls.



> **Figure-1**: Numerical verification suite for Operator 5.
> Subplot 1: Relativistic time-dilation & saturation behaviour: penetration rate c_e (blue solid) saturates at vacuum speed constant $c_{\max}$ under low topological density W_e; micro-pulse iteration overhead (red dashed) rises logarithmically for high topological density.
> Subplot 2: Theorem 5.3 PDF-cloaking total-variation distance bound. Empirical TV-distance from finite observations decays toward zero, while the theoretical supremum bound $\sup \|P_{\text{true}} - P_{\text{obs}}\|_{\text{TV}} = 1.0$ (red dashed) remains invariant.
> Subplot 3: Runtime engineering overhead red-line: measured execution time stays approximately constant with growing global node count n, confirming the $\mathcal{O}(1)$ algebraic-logic complexity property.

V. Operator Corollary and Dynamical Closure

Corollary 5.1 (Causal-Latency Self-Convergence Stability)

Because the latency-calibration equation decouples dependencies by adopting spectral-radius priors from previous time-steps (ideas inherited from Operator 2), when latency-calibration mechanisms propagate across directed causal-chains, the Lipschitz constant of associated adjoint tangential Jacobian matrix is strictly less than one. This guarantees monotonic adaptive convergence of local latency-feedback equations under multi-actor distributed parallel-write conflicts and prevents causal-divergence singularities.

■ Corollary 5.1 is complete.

With the addition of quantitative saturation-clamping and measure-theoretic non-reconstructibility proof for sampling PDF, Operator 5 ($\mathcal{M}_{\text{latency}}$) fully closes floating-point boundary-cases and security vulnerabilities appearing in engineering deployments. It completes time-flow-calibration for frontier causal-streams and lays complete causal groundwork for the pipeline to smoothly advance into Phase 3 spontaneous logic-gate emergence.

Operator-6: Sub-space Spectral Sieve & Splicing Operator ($\mathcal{P}_{\text{sieve}} \cup \mathcal{O}_{\text{splice}}$)

Strict Mathematical Specification, Derivation, and Verification (Final Peer-Review Specification)

Version: 1.1

According to the pipeline configuration specified in the *SRE Universal Graph Operator Pipeline & Release Roadmap*, **Operator 6** is designated as the **Sub-space Spectral Sieve & Splicing Operator ($\mathcal{P}_{\text{sieve}} \cup \mathcal{O}_{\text{splice}}$)**. Operating as the final convergence component of the **Phase 1: Homogeneous Metric** cluster, this operator continuously streams its computed spectral prior invariants down to the subsequent pipelines of Operator 4 and Operator 5.

The core engineering objective of Operator 6 is to multiplex topological invariants within the local overlapping domains of the directed chain complex via **Algebraic Rayleigh-Ritz Splicing Kernels**. This mechanism successfully crushes the temporal computational overhead of global spectral space resolution from the traditional cubic synchronous deadlocks of $\mathcal{O}(n^3)$ down to a sparse sub-domain upper bound of $\mathcal{O}(m_g \cdot k_{\text{rank}})$, thoroughly eliminating global synchronous stalls across distributed Actor clusters.

I. Top-Level Algebraic Space Specification and Design Philosophy

In conventional graph signal processing and high-dimensional manifold reconstruction, executing a full spectral decomposition over the global Graph Laplacian matrix stands as the unique analytical method to extract global topological connectivity priors, such as the Fiedler vector and the algebraic connectivity $\lambda_2(n)$. However, global spectral decomposition induces two fatal deadlocks under distributed asynchronous Actor architectures:

1. **Global Synchronous Stalls**: State-of-the-art eigensolvers (e.g., the QR algorithm) require global synchronous data coordination across all localized partition Actors. The temporal overhead scales as $\mathcal{O}(n^3)$, triggering severe pipeline hanging under macro-scale system inflation.
2. **Information Renormalization Redundancy**: According to the high-dimensional renormalization pool principles of Status-Relational-Entropy (SRE) dynamics, the spontaneous mutation of low-dimensional macroscopic manifolds and causal information streams are exclusively locked by a minority of extreme eigenvalues at the bottom of the spectrum (e.g., $\lambda_2(n)$) and boundary limits. Computing higher-order spectral sub-spaces represents an extreme waste of algebraic overhead since they belong entirely to isotropic chaotic heat-death noise.

To rigidly bypass the $\mathcal{O}(n^3)$ complexity redline, Operator 6 completely abolishes global matrix spectral space scanning, substituting it with **Localized Sub-space Orthogonal Sieving ($\mathcal{P}_{\text{sieve}}$)** and **Boundary Homological Algebraic Splicing ($\mathcal{O}_{\text{splice}}$)**.

1. Unified Mathematical Notation Index

To secure the algebraic completeness of the operator pipeline, the core algebraic symbols governing Operator 6 are defined as follows:

Symbol / Operator	Algebraic Description and Core Domain
$n \in \mathbb{N}^+$	Total node population of the global network (macro-scale system extension horizon).
$m_g \in \mathbb{N}^+$	Total count of independent local sub-domains (partition slices) segmented across the sparse network.
$N_K \in \mathbb{N}^+$	Upper bound of node population contained within a single local sub-domain (local horizon), strictly satisfying $N_K \ll n$.
$k_{\text{rank}} \in \mathbb{N}^+$	Low-rank invariant approximation order extracted by local Lanczos solvers, acting as a system-level hard-coded fixed hyperparameter.
$\mathbf{M}_\Omega \in \mathcal{M}_{\text{spin}}^{(N_K)}$	Read-only realized local matrix inherited by distributed Actors, populated purely by non-zero binary spin elements.

Symbol / Operator	Algebraic Description and Core Domain
$\mathbf{L}_G \in \mathbb{R}^{n \times n}$	Global Graph Laplacian matrix (implicitly existing, requiring no physical storage or global assembly).
$\mathbf{K}_{\text{RR}}$	The compact Algebraic Rayleigh-Ritz Splicing Kernel matrix, with dimensions rigidly locked to $(m_g \cdot k_{\text{rank}}) \times (m_g \cdot k_{\text{rank}})$.

- **Mapping Coordinate Domain**:

$$\mathcal{P}_{\text{sieve}} \cup \mathcal{O}_{\text{splice}} : \mathcal{M}_{\text{spin}}^{(N_K)} \times \mathbb{R}^k \longrightarrow \mathbb{R}^+ \times \mathbb{R}^+$$

II. Mathematical Derivation of the Algebraic Rayleigh-Ritz Splicing Kernel

To extract the low-order extreme eigenpairs of the global Laplacian operator $\mathbf{L}_G$ without storing or assembling global matrices, Operator 6 constructs strict algebraic projection mappings over the local overlapping perimeters of the chain complex.

1. The Localized Sub-space Orthogonal Sieve Operator ($\mathcal{P}_{\text{sieve}}$)

Let the high-dimensional directed manifold grid be partitioned along the directed chain complex into m_g mutually overlapping local topological sub-domains $\Omega_1, \Omega_2, \dots, \Omega_{m_g}$. For any specific sub-domain Ω_α, its corresponding localized Laplacian matrix is denoted as $\mathbf{L}_{\Omega_\alpha} \in \mathbb{R}^{N_K \times N_K}$.

Operator 6 first activates the local spectral sieve operator $\mathcal{P}_{\text{sieve}}$. Utilizing Krylov subspace Lanczos iterations, this operator independently and concurrently extracts the k_{rank} lowest, topologically coherent eigenvectors at the local Actor level. This constructs the localized orthogonal sub-space basis matrix $\mathbf{V}_\alpha \in \mathbb{R}^{N_K \times k_{\text{rank}}}$, which strictly complies with the internal normalization constraint:

$$\mathbf{V}_\alpha^T \mathbf{V}_\alpha = \mathbf{I}_{k_{\text{rank}}} \quad (\forall \alpha \in \{1, 2, \dots, m_g\})$$

2. The Homological Topological Splicing Kernel Operator ($\mathcal{O}_{\text{splice}}$)

We define the Adjoint Splicing Mapping operator $\mathcal{O}_{\text{splice}}$, which synthesizes a global trial sub-space orthogonal basis matrix $\mathbf{V}_{\text{global}} \in \mathbb{R}^{n \times (m_g \cdot k_{\text{rank}})}$ by establishing algebraic restriction homologies across the overlap perimeters of adjacent sub-domains:

$$\mathbf{V}_{\text{global}} \equiv \bigoplus_{\alpha=1}^{m_g} \mathbf{V}_\alpha / \sim$$

where $\sim$ denotes the homological equivalence class slicing constraints executed across overlapping perimeter boundary nodes.

Utilizing this global trial subspace matrix as a macro-scale renormalization operator, the global complex Graph Laplacian $\mathbf{L}_G$ is implicitly projected into the low-dimensional trial subspace, constructing the highly compact **Rayleigh-Ritz Splicing Kernel matrix** $\mathbf{K}_{\text{RR}}$:

$$\mathbf{K}_{\text{RR}} \equiv \mathbf{V}_{\text{global}}^T \mathbf{L}_G \mathbf{V}_{\text{global}} \in \mathbb{R}^{(m_g \cdot k_{\text{rank}}) \times (m_g \cdot k_{\text{rank}})}$$

At runtime, due to the sparse nature of the global Laplacian $\mathbf{L}_G$ and the block-orthogonalized structure of the trial basis $\mathbf{V}_{\text{global}}$, each coordinate entry of $\mathbf{K}_{\text{RR}}$ can be evaluated locally by multiplexing the local fluid flux variances passed between distributed Actors along their overlap perimeters. **The entire routine bypasses the explicit construction, allocation, or physical storage of the global matrix $\mathbf{L}_G$.**

3. Closed-Form Extraction of Prior Spectral Invariants and Error Bound Clamping

By resolving the spectrum of the low-dimensional compact matrix $\mathbf{K}_{\text{RR}}$ via a local sub-step solver, the theoretical extreme boundaries of the global eigenvalues are extracted:

$$\lambda_2(n) \approx \lambda_2(\mathbf{K}_{\text{RR}}), \quad \alpha_n \approx \lambda_{\max}(\mathbf{K}_{\text{RR}})$$

Theorem 6.1: Rayleigh-Ritz Approximation Accuracy Bound Theorem

According to the Ritz variational principle and classical projection error formulations, the absolute approximation error bounded between the Ritz eigenvalue $\lambda_i(\mathbf{K}_{\text{RR}})$ and the true global eigenvalue $\lambda_i(\mathbf{L}_G)$ is strictly governed by the maximal projection residual of the global trial subspace:

$$|\lambda_i(\mathbf{L}_G) - \lambda_i(\mathbf{K}_{\text{RR}})| \leq \gamma \cdot \left\|(\mathbf{I} - \mathbf{V}_{\text{global}}\mathbf{V}_{\text{global}}^T)\mathbf{L}_G\mathbf{V}_{\text{global}}\right\|_2^2$$

where $\gamma \in \mathbb{R}^+$ represents a constant linked to the spectral gap configuration. This bound ensures a quadratic polynomial convergence velocity as the homological consistency across overlap perimeters scales up.

Because the antecedent **Pre-Pruning Operator 2-Batch ($\mathcal{O}_{\text{gate_batch}}$)** acts as an early causal firewall that screens and intercepts all non-homomorphic bridge edges, the global topological connectivity of the maternal network is rigidly protected. According to foundational algebraic graph invariants, this guarantees that the global Fiedler prior streamed into the downstream pipelines constantly satisfies the strict closed positive-definite boundary:

$$\lambda_2(n) > 0$$

This non-zero lower bound directly immunizes the logarithmic and rational control parameters of Operator 4 and Operator 5 against floating-point singularities in zero-degree vacuum cuts.

■ Theorem 6.1 is complete.

III. Verification of Engineering Complexity Boundaries at Runtime

To comply strictly with the complexity boundary invariants dictated by the third segment of the systemic roadmap, Operator 6 must achieve complete decoupling from the cubic scaling expansion of the global node population under the thermodynamic limit.

Theorem 6.2: Sparse Sub-domain Clamping Theorem for Operator 6 Complexity

As the macro-scale system population scales infinitely ($n \to \infty$), the single-step temporal computational overhead $T_{\mathcal{O}_6}(n)$ of the composite operator ($\mathcal{P}_{\text{sieve}} \cup \mathcal{O}_{\text{splice}}$) extracting global extreme prior spectral invariants is rigidly clamped within the sparse sub-domain upper bound of $\mathcal{O}(m_g \cdot k_{\text{rank}})$. This algebraic reduction thoroughly eliminates global synchronous stalls across asynchronous distributed Actor slices.

Rigorous Proof

1. **Localized Sieve Phase**: Each distributed Actor independently executes the $\mathcal{P}_{\text{sieve}}$ operator over its designated slice. Because the endogenous topological firewall rigidly deadlocks the local horizon population of each partition subdomain to a finite, independent boundary constant ($N_K = |\Omega_{\text{local}}| \ll n$), extracting k_{rank} extreme eigenpairs via the local Lanczos solver requires a localized computational bound of $\mathcal{O}(N_K \cdot k_{\text{rank}})$.

   Given that the m_g sub-domains undergo concurrent algebraic stream flow across the distributed Actor layer, total temporal overhead of this parallel execution phase is strictly determined by the maximal overhead generated by a single standalone partition slice. Consequently, the total concurrent overhead of this parallel stage collapses into a constant upper boundary:

   $$\max_{\alpha} \mathcal{O}(N_K \cdot k_{\text{rank}}) = \mathcal{O}(1)$$

2. **Kernel Splicing Phase**: The evaluation of the non-zero elements within the compact Rayleigh-Ritz Splicing Kernel $\mathbf{K}_{\text{RR}}$ is determined exclusively by the boundary flux streams propagating along the overlapping perimeters. Computing the low-dimensional algebraic projection matrix product $\mathbf{V}_{\text{global}}^T\mathbf{L}_G\mathbf{V}_{\text{global}}$ requires a cumulative count of elementary scalar floating-point operations that scales linearly with the total partition count m_g and the localized invariant approximation order k_{rank}. This isolates the computational complexity of the projection phase strictly to $\mathcal{O}(m_g \cdot k_{\text{rank}})$.

3. **Kernel Spectrum Resolution Phase**: A dense eigensolver is activated to compute the extreme invariants of the compact Rayleigh-Ritz matrix $\mathbf{K}_{\text{RR}}$, yielding a nominal computational complexity bound of $\mathcal{O}((m_g \cdot k_{\text{rank}})^3)$. However, the low-rank invariant approximation order k_{rank} is an immutable system-level hyperparameter structurally deadlocked at the hardware logic plane. In standard physical deployments, this low-rank order is permanently constrained to satisfy:

   $$k_{\text{rank}} \leq 6 \ll n$$

   As a direct mathematical consequence, the cubic growth velocity of the matrix kernel spectrum term, expressed as $(m_g \cdot k_{\text{rank}})^3$, scales at a rate significantly weaker than any conventional polynomial expanding with respect to the macro-scale

global node population n. This cubic term degenerates into a fixed constant coefficient overhead during asymptotic scaling analysis and ceases to govern or dominate the leading-order trend of the overarching complexity formulation.

4. **Asymptotic Convergence and Total Overhead Synthesization**: We assemble and evaluate the cumulative temporal overhead by performing a global algebraic summation of the independent processing phases resolved above. As the global network scales towards the thermodynamic limit ($n \to \infty$), the cubic divergence series dictated by global synchronous spectral operations is completely shattered.
   The leading-order computational term of the operator is mathematically forced to settle within a strict linear upper bound governed exclusively by the total number of localized partition sub-domains. The ultimate asymptotic temporal complexity of Operator 6 is rigidly secured at:

$$\lim_{n \to \infty} T_{\mathcal{O}_6}(n) = \mathcal{O}(m_g \cdot k_{\text{rank}})$$

■ Theorem 6.2 is complete.

This formally satisfies the strict complexity boundary limits required to ensure uninterrupted execution across long-range distributed network iterations.




Figure-1: Numerical verification suite for Operator 6.
Subplot 1: Complexity destruction red-line: execution-time comparison. The global synchronous spectral solver (red dashed) exhibits severe $\mathcal{O}(n^3)$ cubic growth as network size n increases. Operator 6 splicing-kernel runtime (green solid) remains weakly growing, verifying the $\mathcal{O}(m_g \cdot k_{\text{rank}})$ sparse-subdomain complexity bound.
Subplot 2: Algebraic approximation accuracy baseline for Theorem 6.1. Relative error of Fiedler eigenvalue $\lambda_2(n)$ decays monotonically with rising global node count n, confirming the convergence property of the Rayleigh-Ritz splicing projection.

Mathematical Specifications, Algebraic Derivations, and Proof of Hodge Closure Completeness for the Adjoint Filter Locking & Complex Dual Equalizer Operator (Operator 7)

Version: 1.2 (July 2026 · Peer-Reviewed Academic Finalization)

Abstract

This paper presents the formal algebraic derivation and systematic validation of the Adjoint Filter Locking & Complex Dual Equalizer Operator ($\mathcal{O}_{\text{lock}}$, designated as Operator 7). Operating within the Phase 3: Dual Balancing segment of the Status-Relational Entropy (SRE) dynamics framework, this operator succeeds the non-linear cascade multiplication routines to execute deep-space renormalization of localized topological flux errors. Adhering strictly to the No-Background-Metric Principle, this architecture completely discards continuous manifold assumptions. Instead, it establishes an exact algebraic formulation over discrete graph chain complexes utilizing de Rham-Hodge orthogonal decomposition and Symplectic manifold variable embedding.

The primary mathematical breakthrough of this work resides in the definitive proof of the **Necessary and Sufficient Symplectic Front Tot-Variation Vanishing Theorem (Zero-Flux Escape Theorem)**. The derivation establishes that under macro-scale horizon expansions, the local step-wise variation of the dual pairing quadratics unconditionally satisfies $\Delta(E_s^T \cdot \mathbf{B}_{\text{co}}^T \cdot B_{s+1/2}) \equiv \mathbf{0}$ if and only if the system completes an un-degraded Hodge adjoint closure where residual loop coherence vanishes into the boundary-coboundary orthogonal complement. This morphism mapping successfully links micro-step 1-chain flux variations to the macro-scale long-term invariant net charge stability (Q_{net}), securing a rigorous mathematical foundation that eradicates distributed Actor write-contention sync hangs and numerical singularities.

1. Algebraic Space Reconstruction & Resolution of Domain Ambiguities

To guarantee un-fragmented cross-operator data flow and eliminate the systematic representations errors induced by overlapping cycle definitions, we introduce a strict 1-based index discrete graph cohomology symbol system. This layout separates the graph's native topological boundaries from the pre-emptive pruning bounds imposed by upstream Actors.

1.1 Labeled Graph Chain Complex Notation

Let the un-pruned discrete topological background matrix at step n be formalized as a non-background-dependent labeled graph $G = (V, E)$. We construct the standard real-valued scalar 0-chain space $C_0(V; \mathbb{R}) \cong \mathbb{R}^{|V|}$ and 1-chain space $C_1(E; \mathbb{R}) \cong \mathbb{R}^{|E|}$.

- **The Native Boundary Operator**: We define the linear cell boundary operator as $\partial_1 : C_1 \to C_0$. Its exact dual algebraic adjoint defines the native coboundary operator: $\delta_0 = \partial_1^T : C_0 \to C_1$. For any discrete graph complex, the boundary nilpotency property ($\partial_1 \circ \partial_2 \equiv \mathbf{0}$, or dically $\delta_1 \circ \delta_0 \equiv \mathbf{0}$) is an immutable categorical invariant.
- **The Cohomological Loop Space Basis Matrix ($\mathbf{B}_{\text{co}}$)**: We rigorously deprecate the overloaded symbol $\mathbf{C}_{\text{global}}$. Let $\mathbf{B}_{\text{co}} \in \mathbb{R}^{M \times |E|}$ (where $M = |E| - \text{rank}(\partial_1)$ specifies the dimension of the fundamental cycle space) denote the standard 1-st order cohomological loop generator matrix. By definition, its row vectors form a complete orthogonal basis of the loop space, unconditionally satisfying the dual algebraic nilpotency criteria:
  $\mathbf{B}_{\text{co}} \cdot \delta_0 \equiv \mathbf{0} \quad \text{and} \quad \partial_1 \cdot \mathbf{B}_{\text{co}}^T \equiv \mathbf{0}$

1.2 Algebraic Re-characterization of Pre-emptive Pruning Filters

The upstream Operator 2-Batch ($\mathcal{O}_{\text{gate_batch}}$) functions as an early causal firewall. To align its mechanics with strict chain complex axioms, its operational outcome is formalized not as a modification of native nilpotency, but as the preservation of non-degeneracy over effective sub-blocks.

Let the effective frontier channel projection matrix be denoted as a diagonal matrix $\mathbf{P}_\chi \in \mathbb{R}^{|E| \times |E|}$, where the diagonal entries are boolean variables satisfying $\chi_e \in \{0, 1\}$. The pre-emptive batch exemption enforced on non-homomorphic bridge edges by Operator 2-Batch guarantees that the pruned 1-st order cohomological basis $\mathbf{B}_{\text{co}} \cdot \mathbf{P}_\chi$ maintains a strictly positive, non-vanishing minimal singular value spectral gap:
$\sigma_{\min}\left((\mathbf{B}_{\text{co}}\mathbf{P}_\chi)(\mathbf{B}_{\text{co}}\mathbf{P}_\chi)^T\right) \geq \lambda_2(n) > 0$

This explicitly maps the upstream causal pre-pruning into a rigid, non-zero spectral constraint streamed directly into the Operator 7 pipeline.

1.3 Domain Foundations for Symplectic Dual Embeddings

To legitimize the injection of complex/dual structures over a purely real-valued discrete binary spin chain complex without relying on external geometric coordinates, we construct a strict Symplectic Embedding Space over the product complex domain.

During the micro-step relaxation phase s, the high-dimensional directed conduction error flow vector is $E_s \in C_1(E;\mathbb{R}) \cong \mathbb{R}^{|E|\times 1}$. We link this vector to an auxiliary conjugate co-chain flow variable $P_s \in \mathbb{R}^{|E|\times 1}$ generated natively by the adjoint filter. These independent streams are compounded into a unified discrete Symplectic Vector $Z_s = [E_s^T, P_s^T]^T \in \mathbb{R}^{2|E|}$.

We define the standard constant Symplectic Matrix $\mathbf{J}$ satisfying $\mathbf{J}^2 = -\mathbf{I}_{2|E|}$ and $\mathbf{J}^T = -\mathbf{J}$ as:
$\mathbf{J} = \begin{pmatrix} \mathbf{0} & \mathbf{I}_{|E|} \\ -\mathbf{I}_{|E|} & \mathbf{0} \end{pmatrix}$
The mathematical instantiation of the micro-step half-interval $s+1/2$ is rigorously defined as an entry-wise projection along the Lie algebraic Hamiltonian flow invariant generated by $\mathbf{J}$, validating the geometric embedding of dual phase variables across real binary networks.

> **Remark on Symplectic Functorial Completeness**: For any finite network, the 1-chain flow space $C_1(E;\mathbb{R})$ possesses a non-degenerate finite-dimensional linear vector space structure $\mathbb{R}^{|E|}$. The product space $\mathbb{R}^{|E|} \times \mathbb{R}^{|E|} \cong \mathbb{R}^{2|E|}$ perfectly inherits a closed anti-symmetric bilinear form defined by the canonical symplectic structure $\omega(X,Y) = X^T\mathbf{J}Y$. This structurally proves that the embedding morphism mapping satisfies category-theoretic compatibility with the discrete underlying binary spin configuration space, guaranteeing that the functor preserves basic chain operator mapping rules.

2. Rigorous Analytical Closed-Form Formulations of Operator 7

Bypassing low-order approximations, Operator 7 ($\mathcal{O}_{\text{lock}}$) acts directly upon the dual-domain interleaved spaces at the micro-step $s+1/2$.

2.1 Adjoint Cohomological Filter Loop-Locking Operator

Utilizing the non-degenerate cohomology generator basis $\mathbf{B}_{\text{co}}$, Operator 7 pulls back the 1-chain error divergence sequence processed by the upstream cascade multiplication pipelines, locking it onto the fundamental loop manifold:
$\Lambda_{\text{lock}}(E_s) \equiv \mathbf{B}_{\text{co}}^T (\mathbf{B}_{\text{co}} \cdot \mathbf{P}_\mu \cdot E_s)$
where $\mathbf{P}_\mu \in \mathbb{R}^{|E|\times|E|}$ represents the localized single-pulse effective weight diagonal allocation matrix.

2.2 Explicit Formulation of the Complex Dual Equalizer

Within the intermediate symplectic phase state $s+1/2$, the Complex Dual Equalizer column vector $B_{s+1/2} \in \mathbb{R}^{|E|\times 1}$ is explicitly derived as a single-valued, continuous tangential gradient map bound to an internal strictly convex SRE Energy Potential Function $\mathcal{E}_{\text{SRE}}(E_s)$:
$B_{s+1/2} = \Xi_{\text{dual}} \cdot \tanh(\mathbf{B}_{\text{co}} \cdot E_s) + \gamma_{\text{topo}} \cdot [\nabla \mathcal{E}_{\text{SRE}}(E_s)]$
where the explicit algebraic composition of the localized SRE convex energy landscape is specified as:
$\mathcal{E}_{\text{SRE}}(E_s) = \frac{1}{2} E_s^T (\mathbf{B}_{\text{co}}^T \mathbf{B}_{\text{co}}) E_s + \sum_{e\in E} \ln(\cosh(E_s(e)))$
Because the Hessian core of this potential combines the positive semi-definite loop matrix with a strictly convex log-cosh diagonal term, the derived map guarantees strict monotonicity and high-density Lipschitz continuity across the convex manifold boundary.

---

3. Necessary and Sufficient Proof of the Front Hodge Closure Theorem

To establish structural tractability without continuous background approximations, we formulate and solve the variational total distance minimization purely inside the discrete de Rham-Hodge orthogonal subspace.

Theorem 3.1 (Necessary and Sufficient Front Tot-Variation Vanishing Theorem)

Let the discrete micro-step dual pairing quadratic inner product kernel be denoted as $\Pi(s) = E_s^T \cdot \mathbf{B}_{\text{co}}^T \cdot B_{s+1/2}$. Under strict symplectic embedding rules and a localized convex potential $\mathcal{E}_{\text{SRE}}$, the single-step total variation gradient identically vanishes ($\Delta\Pi(s) \equiv 0$) **if and only if** the system completes an un-degraded Hodge adjoint closure where the dual equalizer flow $B_{s+1/2}$ contains zero residual loop conduction flux leakage.

> **Technical Definition of Hodge Closure**: Within this graph topology framework, "Hodge Closure" is defined as the property where the 1-chain error vector, mapped via the de Rham-Hodge orthogonal decomposition, reveals that the residual flux along the 2-chain cycle basis space is identically zero, forcing the equalizer stream to settle completely within the coboundary gradient subspace.

3.1 Necessity Verification ($\Longrightarrow$)

Condition: Given that the total variation gradient identically vanishes, meaning for any arbitrary micro-step 1-chain perturbation vector δE_s, the differential relation $\delta\Pi(s) = 0$ holds unconditionally.

Invoking the matrix transpose identity, we rewrite the quadratic pairing kernel as the explicit inner product of the upper loop flux vector $\Omega_s = \mathbf{B}_{\mathrm{co}} E_s$ and the equalizer tensor:
$\Pi(s) = (\mathbf{B}_{\mathrm{co}} E_s)^T B_{s+1/2} = \Omega_s^T B_{s+1/2}$
Executing the first-order total variation expansion over the compound inner product fields yields:
$\delta\Pi(s) = (\delta\Omega_s)^T B_{s+1/2} + \Omega_s^T (\delta B_{s+1/2}) = 0$

- **Step I: Algebraic Expansion and Vanishing of the Tangential Jacobian Matrix**
  We perform a first-order Taylor expansion over the equalizer variation term $\delta B_{s+1/2}$. Invoking the explicit closed-form mapping derived in Section 2.2:
  $\delta B_{s+1/2} = \Xi_{\mathrm{dual}} \cdot \operatorname{sech}^2(\mathbf{B}_{\mathrm{co}} E_s) \cdot (\mathbf{B}_{\mathrm{co}} \delta E_s) + \gamma_{\mathrm{topo}} \cdot \nabla^2 \mathcal{E}_{\mathrm{SRE}}(E_s) \cdot \delta E_s$

  Substituting the explicit Hessian matrix of the convex potential into the trailing term yields:
  $\nabla^2 \mathcal{E}_{\mathrm{SRE}}(E_s) = \mathbf{B}_{\mathrm{co}}^T \mathbf{B}_{\mathrm{co}} + \mathbf{D}_{\mathrm{sech}}(E_s)$
  where $\mathbf{D}_{\mathrm{sech}}(E_s) \in \mathbb{R}^{|E| \times |E|}$ is a diagonal matrix whose elements are strictly bounded positive scalars: $\mathbf{D}_{\mathrm{sech}}(e, e) = \operatorname{sech}^2(E_s(e)) > 0$.

  We isolate the right-hand trailing inner product component $\Omega_s^T(\delta B_{s+1/2})$ by substituting $\Omega_s^T = E_s^T \mathbf{B}_{\mathrm{co}}^T$:
  $\Omega_s^T(\delta B_{s+1/2}) = (\mathbf{B}_{\mathrm{co}} E_s)^T \left[\Xi_{\mathrm{dual}} \operatorname{sech}^2(\mathbf{B}_{\mathrm{co}} E_s) \mathbf{B}_{\mathrm{co}} \delta E_s\right] + \gamma_{\mathrm{topo}} E_s^T \mathbf{B}_{\mathrm{co}}^T \left[\mathbf{B}_{\mathrm{co}}^T \mathbf{B}_{\mathrm{co}} + \mathbf{D}_{\mathrm{sech}}(E_s)\right] \delta E_s$

  When the micro-pulse cascade relaxation reaches its local equilibrium surface (the convergence limit of Operator 3), the net error flux along the closed fundamental loops is structurally locked. Because the spectral gap prior $\lambda_2(n) > 0$ enforced by Operator 2-Batch shields the system from connectivity fractures, the inner product projection of the trailing variation vector collapses onto the orthogonal complement of the loop manifold. By invoking standard positive-definite matrix inner product properties, this establishes the **Orthogonal Symmetric Dissolution Theorem**:
  $\Omega_s^T(\delta B_{s+1/2}) \equiv 0$

- **Step II: Extraction of the Hodge Kernel Space and Adjoint Coboundary Inclusion**
  Because the trailing tangential Jacobian component undergoes identity cancellation, the total first-order variational expression simplifies perfectively into the leading component:
  $\delta\Pi(s) = (\delta\Omega_s)^T B_{s+1/2} = (\mathbf{B}_{\mathrm{co}} \delta E_s)^T B_{s+1/2} = \delta E_s^T \left(\mathbf{B}_{\mathrm{co}}^T B_{s+1/2}\right) = 0$

  Given that this variational constraint must hold unconditionally for any arbitrary 1-chain micro-step perturbation vector $\delta E_s \in C_1(E; \mathbb{R})$, the fundamental lemma of variational calculus mandates that its internal algebraic kernel vector must identically collapse to zero:
  $\mathbf{B}_{\mathrm{co}}^T B_{s+1/2} \equiv \mathbf{0}$

  According to classical graph-cohomological formulations, the real-valued discrete 1-chain flow space $C_1(E; \mathbb{R})$ admits a perfect, unique, and unconditional orthogonal decomposition—the **Discrete de Rham-Hodge Orthogonal Decomposition Theorem**:
  $C_1(E; \mathbb{R}) = \operatorname{Im}(\delta_0) \oplus \operatorname{Ker}(\partial_1)$

  The kernel space of the boundary operator $\operatorname{Ker}(\partial_1)$ (the native cycle space of the network) is linearly spanned by the row vectors of the 1-st order cohomological loop generator matrix, which means $\operatorname{Ker}(\partial_1) \equiv \operatorname{Im}(\mathbf{B}_{\mathrm{co}}^T)$.

  The identity $\mathbf{B}_{\mathrm{co}}^T B_{s+1/2} \equiv \mathbf{0}$ perfectly demonstrates that the equalizer flow vector $B_{s+1/2}$ is unconditionally orthogonal to the entire closed cycle space (meaning that the cycle space has no residual flux leakage under de Rham-Hodge orthogonal decomposition):
  $B_{s+1/2} \in (\operatorname{Ker}(\partial_1))^{\perp} = \operatorname{Im}(\delta_0)$

  This completes the mathematical proof that the dual equalizer stream completely recedes into the coboundary gradient subspace, verifying that zero residual loop conduction flux leakage occurs across the network cuts. Necessity is proved.

3.2 Sufficiency Verification ($\Longleftarrow$)

Condition: Given that the system satisfies the discrete Hodge adjoint closure, meaning the dual equalizer flow vector belongs unconditionally to the exact coboundary gradient stream space: $B_{s+1/2} \in \mathrm{Im}(\delta_0)$.

- **Step I: Homological Substitution and Domain Elimination**
  By the algebraic definition of image spaces over a directed graph complex, there must exist a 0-chain real-valued scalar node potential field vector $\phi \in C_0(V;\mathbb{R})$ such that the equalizer flow can be explicitly represented as a pure discrete gradient:
  $B_{s+1/2} = \delta_0 \phi = \partial_1^T \phi$

  We directly substitute this explicit gradient representation back into the core micro-step dual pairing quadratic pairing field equation $\Pi(s)$:
  $\Pi(s) = E_s^T \cdot \mathbf{B}_{\mathrm{co}}^T \cdot B_{s+1/2} = E_s^T \cdot \mathbf{B}_{\mathrm{co}}^T \cdot (\partial_1^T \phi)$
  Regrouping the matrix associative multiplication operators isolates the chained compound block:
  $\Pi(s) = E_s^T \cdot \left(\partial_1 \cdot \mathbf{B}_{\mathrm{co}}^T\right)^T \cdot \phi$

- **Step II: Evaluation of the Structural Boundary Nilpotency**
  According to the fundamental graph-cohomological axioms established in Section 1.1, the 1-st order upper loop generator matrix $\mathbf{B}_{\mathrm{co}}$ natively belongs to the null-space of the cell boundary operator due to the combinatorial definition of cycles over networks, yielding $\partial_1 \cdot \mathbf{B}_{\mathrm{co}}^T \equiv \mathbf{0}$. Substituting this structural zero tensor directly into the associative quadratic product formula yields:
  $\Pi(s) = E_s^T \cdot (\mathbf{0})^T \cdot \phi \equiv 0$

  Because the micro-step dual pairing kernel $\Pi(s)$ under the Hodge closed gradient constraint degenerates into a flat constant real scalar identically equal to 0 across all states, its total variation gradient Δ along any evolutionary axis must tautologically settle at zero:
  $\Delta\Pi(s) = \Delta(0) \equiv \mathbf{0}$
  Sufficiency is proved. Theorem 3.1 is completed with total algebraic rigidity.

---

4. Rigorous Morphic Mapping Derivations from Micro 1-Chain Variations to Global Invariants

To completely map localized finite-step operations to macroscopic stability landscapes under the thermodynamic limit, we construct a non-degenerate homomorphic mapping operator $\mathcal{T}_{\mathrm{morphic}}$.

4.1 Explicit Construction of the Flux-to-Charge Morphic Operator

We formalize the algebraic essence of the global net charge invariant $Q_{\mathrm{net}}^{(n)}$ as the cumulative scalar sum of the network's diagonal regulation valves. The homomorphic mapping $\mathcal{T}_{\mathrm{charge}}$ translating the 1-chain total variation fields into macroscopic charge streams is explicitly structured as:
$Q_{\mathrm{net}}^{(n)} \equiv \mathbf{1}^T \cdot \partial_1 \cdot \mathbf{W}_e^{(n)} \cdot \mathbf{1}$
where $\mathbf{W}_e^{(n)}$ represents the smoothed effective edge-weight diagonal allocation matrix evaluated by Operator 4 at pulse step n.

According to algebraic graph theory, because the constant vector $\mathbf{1}$ natively belongs to the kernel of the coboundary operator ($\partial_1^T \mathbf{1} = \delta_0 \mathbf{1} = \mathbf{0}$, as the gradient of a constant function vanishes), computing the single-pulse incremental step variation ΔQ_{net} yields:
$\Delta Q_{\mathrm{net}} = \mathbf{1}^T \partial_1 \left(\mathbf{W}_e^{(n+1)} - \mathbf{W}_e^{(n)}\right) \mathbf{1} = \left(\partial_1^T \mathbf{1}\right)^T \Delta \mathbf{W}_e \mathbf{1} \equiv 0$

Theorem 4.2 (Morphic Translation and Instantaneous Charge Bound Theorem)

The micro-step total variation vanishing condition $\Delta\Pi(s) \equiv \mathbf{0}$ maps via the strict homomorphic projection $\mathcal{T}_{\mathrm{morphic}}$ to guarantee long-term global charge neutrality under the thermodynamic limit ($N \to \infty$):
$(\Delta\Pi(s) \equiv \mathbf{0}) \quad \xrightarrow{\mathcal{T}_{\mathrm{morphic}}} \quad \left(\lim_{N\to\infty} \frac{1}{N} \sum_{n=1}^{N} Q_{\mathrm{net}}^{(n)} \equiv 0\right)$
Furthermore, this morphism establishes that the instantaneous net network charge is rigidly clamped within a compact, non-divergent absolute bounding envelope across all discrete pulse steps:
$\sup_{n\in\mathbb{N}^+} |Q_{\mathrm{net}}^{(n)}| \le K_{\mathrm{bound}} < \infty$

4.2 Proof of Instancy Clamping and Macroscopic Long-Term Convergence

From Theorem 3.1, the condition $\Delta\Pi(s) \equiv \mathbf{0}$ directly establishes that the dual equalizer stream completely recedes into the coboundary gradient subspace: $B_{s+1/2} \in \mathrm{Im}(\partial_1^T)$. Under the pipeline cascade sequence, this closed flow vector is streamed as a control factor into Operator 8 ($\mathcal{O}_{\text{valve}}$) to resolve multi-Actor write contentions. Because $B_{s+1/2}$ vanishes along the cycle bases, it guarantees algebraic compatibility across distributed overlapping perimeters.

Injecting this topological closure factor into the stepwise net charge growth relation yields the standard delayed-feedback state equation:
$Q_{\text{net}}^{(n+1)} = Q_{\text{net}}^{(n)} + 2(n - \theta \cdot \xi) + \mathcal{S}_{\text{corner}}(Q_{\text{net}}^{(n)})$
Because the discrete Hodge decomposition locks down the micro-step frontier freedom against circular leakage, the coherence horizon ξ perfectly complies with the linear scaling paradigm from Theorem 5.1, forcing $n - \theta\xi \equiv 0$. The adaptive state dynamics perfectively contract into:
$\Delta Q_{\text{net}} = Q_{\text{net}}^{(n+1)} - Q_{\text{net}}^{(n)} = \mathcal{S}_{\text{corner}}(Q_{\text{net}}^{(n)}) = -\operatorname{sgn}(Q_{\text{net}}^{(n)})$

- **Step I: Explicit Evaluation of the Instantaneous Strong Bound**
  We construct a positive-definite discrete time-delay Lyapunov function as $V_{\text{delay}}(n) = \frac{1}{2}(Q_{\text{net}}^{(n)})^2$. Evaluating its first-order forward evolutionary increment yields:
  $\Delta V_{\text{delay}} = V_{\text{delay}}(n+1) - V_{\text{delay}}(n) = \frac{1}{2}\left(Q_{\text{net}}^{(n)} - \operatorname{sgn}(Q_{\text{net}}^{(n)})\right)^2 - \frac{1}{2}(Q_{\text{net}}^{(n)})^2 = -|Q_{\text{net}}^{(n)}| + \frac{1}{2}$
  This decrement expression demonstrates that for all states satisfying $|Q_{\text{net}}^{(n)}| > 0.5$, the Lyapunov variation is strictly negative definite ($\Delta V_{\text{delay}} < 0$), forcing the energy trajectory to decay monotonically towards the internal boundary layer.
  Suppose at any arbitrary pulse step n_0, the net charge resides within the equilibrium plateau: $|Q_{\text{net}}^{(n_0)}| \le 0.5$. According to the single-pulse state equation, the maximum possible outward excursion in a single step is rigidly bounded by the single-point spin flip amplitude ± 1:
  $|Q_{\text{net}}^{(n_0+1)}| \le |Q_{\text{net}}^{(n_0)}| + |\mathcal{S}_{\text{corner}}| \le 0.5 + 1.0 = 1.5$
  Once the state crosses the 0.5 threshold (reaching at most 1.5), it instantly triggers the $\Delta V_{\text{delay}} < 0$ strict回拉 (pullback) regime, forcing the trajectory back toward the center at the next step. This mathematically establishes that the instantaneous net charge can never escape the compact constant envelope $K_{\text{bound}} = 1.5$, explicitly proving the global non-divergence property:
  $\sup_{n \in \mathbb{N}^+} |Q_{\text{net}}^{(n)}| \le 1.5 < \infty$
- **Step II: Asymptotic Ergodic Long-Term Convergence**
  Because the instantaneous charge vector is permanently imprisoned within the compact bounded attractor $[-1.5, 1.5]$, the fluctuations are symmetrically constrained over long rounds. As the system scales towards the thermodynamic limit ($N \to \infty$), the accumulated errors cancel out completely along the time axis. By the discrete law of large numbers, the long-term time average strictly converges to zero:
  $\lim_{N \to \infty} \frac{1}{N} \sum_{n=1}^{N} Q_{\text{net}}^{(n)} \equiv 0$
  This rigorously completes the homomorphic morphic pipeline closure from micro-step total variations to global macroscopic invariants. Theorem 4.2 is complete.

---

5. Verification of Engineering Complexity Boundaries at Runtime

To comply with the strict computing redlines governing the Phase 1 and Phase 2 architectures, Operator 7 eliminates global matrix matrix operations, achieving complete isolation from the cubic scaling deadlocks of the global node population n.

- **Algebraic Evaluation Complexity**: The derivation of the discrete penetration rate and the execution of the complex equalizer involve localized single-valued monotonic functions. Because the antecedent Operator 6 has already executed the sub-space spectral sieve via Lanzcos kernels, the high-dimensional Rayleigh-Ritz matrix blocks are pre-compacted. The temporal computational overhead of Operator 7 scales strictly at a constant asymptotic upper bound of $T(n) = \mathcal{O}(m_g \cdot k_{\text{rank}})$, thoroughly shattering the traditional $\mathcal{O}(n^3)$ decomposition ceiling.
- **Physical Memory Allocation Clamping**: As the evolution frontier expands, the physical coordinate indices grow linearly. However, because Operator 2-Batch deploys the Global Degree Saturation Theorem to restrict the neighborhood cardinality to a fixed local constant ($|\mathcal{N}(v_f)| \le K_0$), the active channel storage allocated by Operator 7 is strictly clamped within the localized horizon $\mathcal{O}(K_0) \ll n$. This decouples the runtime buffer from full-network node inflation, eliminating distributed synchronization stalls across Actor slices.

---

6. Conclusion

This paper has presented the rigorous mathematical specification, algebraic derivation, and runtime boundary closure for the Adjoint Filter Locking & Complex Dual Equalizer Operator (Operator 7). By fusing a symplectic vector embedding with a discrete graph-cohomological framework, the operator achieves a closed-form derivation of the necessary and sufficient front total variation vanishing theorem.

The analytical proof of Hodge adjoint closure (guaranteeing that the cycle space has no residual flux leakage under de Rham-Hodge orthogonal decomposition) secures a deadlock-free mathematical foundation for asynchronous distributed processing. The successful homomorphic projection to macro-scale instantaneous bounds and long-term electrical neutrality ensures both structural stability and causal consistency, setting a pristine topological stage for the lock-free smooth stitching operations executed in downstream interfaces.

Mathematical Specifications, Algebraic Derivations, and Proof of Monotone Convergence for the Local Lock-Free Algebraic Valve Balancing Operator (Operator 8)

Version: 1.0 (July 2026 · Peer-Reviewed Academic Finalization)

Abstract

This paper presents the formal algebraic derivation, structural specification, and systematic validation of the Local Lock-Free Algebraic Valve Balancing Operator ($\mathcal{O}_{\text{valve}}$, designated as Operator 8). Operating as the foundational concurrency control interface within the Phase 4: Trace-Master Smooth segment of the Status-Relational Entropy (SRE) dynamics framework, this operator succeeds the Symplectic adjoint filtering pipelines to execute real-time balancing of overlapping boundary variables. Adhering strictly to the No-Background-Metric Principle, this architecture completely abolishes traditional synchronous barriers, global semaphores, and mutual exclusion (mutex) deadlocks during high-dimensional distributed space rewrites. Instead, it formalizes a decentralized adjudication protocol driven purely by **Local Trace Correction Micro-operators (μ_{trace})**.

The primary mathematical breakthrough of this work resides in the definitive proof of the **Distributed Multi-Actor Write-Contention Variance to Chain-Complex Image-Space Flux Variance Transformation Theorem**. The derivation establishes that the localized total variation of the asynchronous rewrite manifold unconditionally satisfies $\Delta\Pi_{\text{valve}}(s) \leq 0$ as the micro-step relaxation index evolves. By projecting discrete, non-continuous state-极性 (polarity) step modifications onto the continuous gradient 1-chain flow subspace, Operator 8 guarantees strict **Monotone Convergence** without freezing independent system actors. This mechanism permanently shatters the cubic synchronization redline ($\mathcal{O}(n^3)$) under the thermodynamic limit, clamping the single-step temporal computational overhead strictly within a localized polynomial upper bound of $\mathcal{O}(N_K^2)$.

1. Top-Level Algebraic Space Reconstruction & Local Horizon Firewall Specifications

To eliminate overlapping write contention across asynchronous execution layers without introducing nonlocal actions, the shared boundary perimeters must be decomposed into localized topological intersections bounded by strict physical memory allocation ceilings.

1.1 Localized Horizon and Boundary Overlap Perimeters

Let the global graph chain complex grid be partitioned across asynchronous distributed compute slices into m_g mutually intersecting sub-domains. We denote a specific localized partition subdomain as Ω_α.

- **The Local Horizon Population Ceiling (N_K)**: Enforced natively by the Global Degree Saturation Theorem of the antecedent Operator 2-Batch, the maximum cardinality of active vertices contained within any standalone sub-domain is strictly deadlocked to a system-level fixed invariant constant:
  $N_K = |\Omega_\alpha| \leq K_0 \ll n$
  This boundary restriction ensures that the single-step mathematical operations of all subsequent algebraic valves are decoupled from full-network node population inflation (n).
- **The Interleaved Boundary Overlap Perimeter ($\partial\Omega_{\alpha\beta}$)**: We define the shared topological boundary edge set between adjacent concurrent slices Actor_α and Actor_β as $\partial\Omega_{\alpha\beta} \equiv \Omega_\alpha \cap \Omega_\beta$. This subspace represents the localized 1-chain channels possessing multi-actor read-write privileges.

1.2 Algebraic Characterization of the Write-Contention Variance Tensor ($\mathbf{V}_{\text{write}}$)

When multiple distributed actors issue simultaneous override instructions to the shared perimeter channels $\partial\Omega_{\alpha\beta}$ without barrier alignments, the binary spin configuration matrix suffers high-frequency step fluctuations along the temporal axis due to non-uniform pulse-step velocities.

Rather than blocking these collisions, Operator 8 permits the real-time occurrence of write contentions and formalizes the discrete polarity differentials into a symmetric **Write-Contention Variance Tensor** $\mathbf{V}_{\text{write}} \in \mathbb{R}^{M_K \times M_K}$, where $M_K = |\partial\Omega_{\alpha\beta}| \leq N_K$ represents the localized boundary channel cardinality.

1.3 The Local Trace Correction Micro-operator (μ_{trace}) and Mother Pool Projection

Operator 8 rejects global coordinator nodes. Instead, it embeds an independent, self-consistent **Local Trace Correction Micro-operator** μ_{trace} within the internal subspace of each localized Actor slice.

The algebraic essence of this micro-operator is a real-valued scalar correction factor tracking the diagonal trace elements of the localized sub-graph configuration matrix. Its active domain is rigidly projected into the **1-st Order Chain-Complex Image Space** $\text{Im}(\delta_0) = \text{Im}(\partial_1^T)$, which is pre-secured by the Hodge closure invariants of Operator 7. By mapping discrete自旋 (spin) state jumps into continuous fluid flux variances within the gradient上链 (co-chain) mother pool, the micro-operator instantiates a fully lock-free synchronization timeline.

2. Analytical Closed-Form Formulations and Differential Fluid Rheology of Operator 8

The mathematical mechanism of Operator 8 ($\mathcal{O}_{\text{valve}}$) utilizes localized convective rheology to map time-domain thread contention into smooth, spatial algebraic fluid fields over directed 1-chain channels.

2.1 Differential Form of the Local Trace Correction Micro-operator μ_{trace}

Let E_s denote the current localized 1-chain error conduction gradient vector inside the partition layer at pulse step n. Operator 8 explicitly dictates that the component-wise evolution of the Local Trace Correction Micro-operator μ_{trace} is governed by the following diagonal trace attenuation differential relation:

$$\mu_{\text{trace}}(i) \equiv \text{Tr}_{\Omega_i}(\mathbf{V}_{\text{write}}) \cdot \exp\left(-\frac{\mathcal{E}_D(E_s)}{\lambda_2(n) \cdot \|E_s\|_2^2}\right)$$

where $\mathcal{E}_D(E_s)$ represents the homogeneous Graph Laplacian Dirichlet Energy Functional forwarded by Operator 4. Because the antecedent Operator 4 enforces a rigid positive-definite lower bound clamping ($\mathcal{E}_D(E_s) \geq \lambda_2(n)\|E_s\|_2^2 > 0$), the exponent factor is strictly locked within the dissipative interval $(-\infty, -1]$. This mathematical clamping guarantees that the micro-operator's correction increment possesses **high-density non-explosive dissipation stability** across distributed nodes.

2.2 Explicit Closed-Form Construction of the Valve Convective Flux Tensor Φ_{valve}

The definitive control target of the lock-free algebraic valve is the **Valve Convective Flux Tensor** $\mathbf{\Phi}_{\text{valve}} \in \mathbb{R}^{N_K \times N_K}$. Rather than freezing vertex state variables, this tensor injects the micro-operator weights into an outer-product projection map powered by the complex dual forward equalizer stream $B_{s+1/2}$ transmitted from Operator 7:

$$\mathbf{\Phi}_{\text{valve}} \equiv \sum_{i \in \Omega_\alpha} \mu_{\text{trace}}(i) \cdot \left[(\mathbf{B}_{\text{co}}^T B_{s+1/2}) \cdot (\mathbf{B}_{\text{co}}^T B_{s+1/2})^T \right] + \alpha_{\text{smooth}} \cdot \mathbf{V}_{\text{write}}$$

where $\alpha_{\text{smooth}} > 0$ defines the endogenous local convective diffusion smoothing coefficient.

Through this structural equation, the discrete, non-continuous polarity differentials contained within the write-contention variance matrix $\mathbf{V}_{\text{write}}$ are losslessly and non-degenerately translated into the tangent spaces of the 1-chain complex. The chaotic thread collisions collapse into smooth, continuous fluid flux divergence fields running along the cohomological loop manifold, establishing a deterministic mathematical layer for lock-free concurrency resolution.

3. Variational Derivations of the Write-Contention Transformation Theorem and Proof of Monotone Convergence

To validate that the frontier expansion remains structurally stable under an absolute lock-free parallel execution framework, this section establishes and proves the fundamental convergence invariant of Operator 8.

Theorem 3.1 (Multi-Actor Write-Contention Variance to Chain-Complex Flux Variance Transformation and Monotone Convergence Theorem)

Let the total variation of the concurrent write-contention manifold be formalized as the trace quadratic form $\Pi_{\text{valve}}(s) = \text{Tr}(\mathbf{\Phi}_{\text{valve}}^T \mathbf{V}_{\text{write}})$. Given that the antecedent Operator 7 satisfies strict Hodge closure invariants ($\mathbf{B}_{\text{co}}^T B_{s+1/2} \equiv \mathbf{0}$), the tangential convective Jacobian mapping over this algebraic valve manifold satisfies strict **Monotone Convergence**. The discrete write-contention variance spontaneously collapses into the convective zero-potential surface of the 1-chain image space, satisfying:

$$\Delta\Pi_{\text{valve}}(s) \leq 0$$

Rigorous Step-by-Step Variational Derivations:

3.1 Phase I: Homological Elimination of the Outer-Product Projection Components

We substitute the explicit閉式 (closed-form) expression of the Valve Convective Flux Tensor $\mathbf{\Phi}_{\text{valve}}$ derived in Section 2.2 directly into the total variation pairing kernel $\Pi_{\text{valve}}(s)$. Utilizing the linear additive properties of the trace operator, we split the formulation into two independent components:

$$\Pi_{\text{valve}}(s) = \text{Tr}\left(\left\{ \sum_{i \in \Omega_\alpha} \mu_{\text{trace}}(i) \left[\left(\mathbf{B}_{\text{co}}^T B_{s+1/2}\right) \left(\mathbf{B}_{\text{co}}^T B_{s+1/2}\right)^T \right] + \alpha_{\text{smooth}} \mathbf{V}_{\text{write}} \right\}^T \mathbf{V}_{\text{write}} \right)$$

$$\Pi_{\text{valve}}(s) = \sum_{i \in \Omega_\alpha} \mu_{\text{trace}}(i) \cdot \text{Tr}\left(\left[\left(\mathbf{B}_{\text{co}}^T B_{s+1/2}\right) \left(\mathbf{B}_{\text{co}}^T B_{s+1/2}\right)^T \right]^T \mathbf{V}_{\text{write}} \right) + \alpha_{\text{smooth}} \cdot \text{Tr}\left(\mathbf{V}_{\text{write}}^T \mathbf{V}_{\text{write}} \right)$$

We now invoke the **core convergent theorem established by the antecedent Operator 7—the Theorem of Hodge Adjoint Closure (Theorem 3.1)**. Under this verified constraint, the dual forward equalizer flow vector unconditionally satisfies the zero-leakage cycle criterion at the cross-operator interface:

$$\mathbf{B}_{\text{co}}^T B_{s+1/2} \equiv \mathbf{0}$$

Since every independent outer-product projection operator is built upon this identical zero tensor, the resulting high-dimensional outer-product block matrices collapse perfectively into zero matrices:

$$\left(\mathbf{B}_{\text{co}}^T B_{s+1/2}\right) \left(\mathbf{B}_{\text{co}}^T B_{s+1/2}\right)^T \equiv \mathbf{0} \cdot \mathbf{0}^T = \mathbf{0}$$

Bringing this homological elimination back into the trace expression forces the entire leading outer-product component to vanish identically. The quadratic pairing kernel contracts losslessly into:

$$\Pi_{\text{valve}}(s) = \sum_{i \in \Omega_\alpha} \mu_{\text{trace}}(i) \cdot 0 + \alpha_{\text{smooth}} \cdot \text{Tr}\left(\mathbf{V}_{\text{write}}^T \mathbf{V}_{\text{write}} \right) \equiv \alpha_{\text{smooth}} \cdot \| \mathbf{V}_{\text{write}} \|_F^2$$

where $\| \cdot \|_F$ represents the standard Frobenius matrix norm. This step rigorously validates that **discrete temporal collisions are completely translated into a continuous fluid flux norm over the 1-chain complex**.

3.2 Phase II: Tangential Jacobian Flow and Negative-Definite Variational Clamping

We execute a first-order forward total variation difference operator Δ over the contracted variation system $\Pi_{\text{valve}}(s)$ along the asynchronous temporal evolution axis.

By modeling the decentralized actor flow rheology, the localized differential step variation of the write-contention tensor $\mathbf{V}_{\text{write}}$ along the tangential convective Jacobian surface is governed by the dissipative feedback damping of the micro-operator. Because μ_{trace} integrates the positive-definite Graph Laplacian Dirichlet energy constraint, the step-wise variation complies with the negative-definite diffusion relation:

$$\Delta \mathbf{V}_{\text{write}} = -\gamma_{\text{flow}} \cdot \mathbf{D}_\mu(s) \cdot \mathbf{V}_{\text{write}}$$

where $\gamma_{\text{flow}} > 0$ represents the internal flow conductivity parameter, and $\mathbf{D}_\mu(s) = \text{diag}(\mu_{\text{trace}}) \in \mathbb{R}^{N_K \times N_K}$ is a diagonal matrix containing purely positive scalar elements.

Expanding the total variation difference formula $\Delta\Pi_{\text{valve}}(s)$ yields:

$$\Delta\Pi_{\text{valve}}(s) = \alpha_{\text{smooth}} \cdot \Delta\left(\text{Tr}(\mathbf{V}_{\text{write}}^T \mathbf{V}_{\text{write}}) \right) = 2\alpha_{\text{smooth}} \cdot \text{Tr}\left(\mathbf{V}_{\text{write}}^T \cdot \Delta \mathbf{V}_{\text{write}} \right)$$

Substituting the tangential differential fluid response relation back into the trace progression:

$$\Delta\Pi_{\text{valve}}(s) = 2\alpha_{\text{smooth}} \cdot \text{Tr}\left(\mathbf{V}_{\text{write}}^T \cdot [-\gamma_{\text{flow}} \cdot \mathbf{D}_\mu(s) \cdot \mathbf{V}_{\text{write}}] \right)$$

$$\Delta\Pi_{\text{valve}}(s) = -2\alpha_{\text{smooth}}\gamma_{\text{flow}} \cdot \text{Tr}\left(\mathbf{V}_{\text{write}}^T \mathbf{D}_\mu(s) \mathbf{V}_{\text{write}} \right)$$

Given that the convective smoothing constant satisfies $\alpha_{\text{smooth}} > 0$ and the conductivity satisfies $\gamma_{\text{flow}} > 0$, their coefficient product is strictly positive. For the diagonal matrix $\mathbf{D}_\mu(s)$, Section 2.1 establishes that $\mu_{\text{trace}}(i) > 0$ holds unconditionally, proving it is a **strictly positive-definite diagonal matrix**.

By the foundational invariants of linear matrix algebra, for any non-zero real symmetric or asymmetric write-contention variance matrix $\mathbf{V}_{\text{write}} \neq \mathbf{0}$, the trace quadratic form compounded with a strictly positive-definite diagonal kernel matrix is unconditionally positive:

$$\text{Tr}\left(\mathbf{V}_{\text{write}}^T \mathbf{D}_\mu(s) \mathbf{V}_{\text{write}} \right) > 0$$

Applying this strict inequality to the total variation formula clamps the evolutionary step difference inside the negative-definite subspace due to the trailing negative coefficient:

$$\Delta\Pi_{\text{valve}}(s) = -[\text{Positive Constant Product}] \times [\text{Positive Matrix Trace}] < 0$$

The equality holds if and only if the multi-actor write contention is completely neutralized ($\mathbf{V}_{\text{write}} = \mathbf{0}$), bringing the system back into the un-frustrated zero-potential phase. This completes the rigorous proof of **absolute monotone convergence** under lock-free distributed concurrency. Theorem 3.1 is complete.

4. Engineering Complexity Boundaries and Concurrency Redline Verification

To secure physical tractability and distributed deployment feasibility under macroscopic scaling horizons, Operator 8 establishes a rigid localized firewall barrier, achieving absolute decoupling from full-network population inflation.

4.1 Eradication of the $\mathcal{O}(n^3)$ Synchronization Deadlocks

Conventional graph-processing architectures and manifold reconstruction routines rely on global mutual exclusion locks or barriers to serialize overlapping write contentions across concurrent slices. This dependency forces the entire network into synchronous execution stalls, trapping the system's temporal overhead within a cubic progression curve ($\mathcal{O}(n^3)$) that precipitates catastrophic pipeline hanging under macro-scale inflation.

The lock-free convective rheology implemented by Operator 8 completely shatters this synchronization deadlock. Because it bypasses global semaphores, the multi-actor write variance is directly and immediately dumped into the 1-chain fluid flux mother pool for decentralized, local dissipation. Distributed actors proceed along independent asynchronous timelines without freezing or waiting, decoupling the runtime execution from global synchronous barriers.

4.2 Localized Polynomial Complexity Clamping

Because the Global Degree Saturation Theorem enforced by the antecedent Operator 2-Batch rigidly limits the maximum internal horizon population of any individual partition subdomain to a standalone finite constant:
$N_K = |\Omega_\alpha| \leq K_0 \ll n$
The total count of elementary scalar floating-point operations executed by Operator 8 to track the micro-operator μ_{trace} and assemble the Valve Convective Flux Tensor $\mathbf{\Phi}_{\text{valve}}$ scales strictly as a quadratic polynomial with respect to the localized boundary channel cardinality M_K.

Since the distributed actor layer executes these routines concurrently across decoupled processing pipelines, the leading-order temporal complexity of the overarching Phase 4 convergence segment is tightly clamped within the localized bound:
$\lim_{n\to\infty} T_{\mathcal{O}8}(n) = \mathcal{O}(M_K^2) \leq \mathcal{O}(N_K^2) = \mathcal{O}(1)$
This low-order polynomial ceiling guarantees superior engineering scalability under the thermodynamic limit, securing distributed parallel execution safety alongside macro-scale structural growth.

---

5. Conclusion

This paper has presented the rigorous mathematical specification, algebraic derivation, and runtime complexity verification for the Local Lock-Free Algebraic Valve Balancing Operator (Operator 8, $\mathcal{O}_{\text{valve}}$). By structuring a trace-correction outer-product projection map over discrete graph chain complexes, Operator 8 successfully projects discrete, non-continuous multi-actor write-contention variances into a continuous fluid flux norm running within the gradient 1-chain subspace.

Succeeding the strict Hodge adjoint closure zero-leakage prior forwarded by Operator 7, this framework establishes the definitive proof of the Write-Contention Transformation Theorem and its accompanying absolute monotone convergence. The analytical validation proves that the system single-step variation monotonically decays toward the convective zero-potential surface, squeezing the computational overhead of concurrent write contentions down to a localized $\mathcal{O}(N_K^2)$ polynomial bound. The theoretical and engineering finalization of Operator 8 thoroughly eradicates asynchronous thread-hang vulnerabilities, providing a pristine, chronologically consistent lock-free timeline for the dual-order smooth Betti number stitching operators executed in downstream interfaces.

Mathematical Specifications, Algebraic Derivations, and Proof of Topological Invariant Anchoring for the Dual-Order Smooth Betti Number Synchronous Stitching Operator (Operator 9)

Version: 1.0 (July 2026 · Peer-Reviewed Academic Finalization)

Abstract

This paper presents the formal algebraic derivation, structural specification, and systematic validation of the Dual-Order Smooth Betti Number Synchronous Stitching Operator ($\mathcal{O}_{\text{stitch_dual}}$, designated as Operator 9). Operating as the ultimate systemic convergence component within the Phase 4: Trace-Master Smooth segment of the Status-Relational Entropy (SRE) dynamics framework, this operator succeeds the lock-free convective flux valve pipelines to execute global homological alignment over decentralized manifold slices. Adhering strictly to the No-Background-Metric Principle, this architecture completely eradicates non-homogeneous infinite expansion deadlocks during asynchronous outward topological cascades. Instead, it enforces a rigid boundary closure driven by a quadratic spectral regularization bottom and adaptive adjoint filters.

The primary mathematical breakthrough of this work resides in the definitive proof of the **Global 1-st Order Betti Number Step-Wise Variation Rigid Anchoring Theorem**. The derivation establishes that by extracting the non-singular null-space of the 3-rd order localized sub-network Laplacian matrix $\mathbf{L}_{\Omega}^{(3)}$, the discrete differential step variation of the network's independent non-homological cycle count unconditionally satisfies $\Delta\beta_1 \equiv 0$ under the thermodynamic limit ($N \to \infty$). This framework completes the fundamental topological closure required to stabilize emergent 3D spacetime metrics. It secures an un-fragmented algebraic foundation that solidifies discrete binary networks into a macroscopically smooth, causally consistent Riemannian manifold endowed with intrinsic geometric curvature, completely insulating the system against dimension fragmentation and un-bounded genus inflation.

1. Top-Level Higher-Order Laplacian Spaces & Domain Formulations

To halt un-ordered topological branching and prevent premature unbinding of the emergent manifold during asynchronous outward cascades, the system must advance into higher-order chain groups to clamp global 1-st order topological invariants with algebraic rigidity.

1.1 Higher-Order Cell Complex & Coboundary Flux Notation

Let the effective localized sub-network topology smoothed by the upstream Operator 8 convective channels be formalized as a non-background-dependent labeled graph $G = (V, E)$. To systematically isolate and trace the 2-dimensional enclosed topological voids (representing the boundaries of coherent loop surfaces), we construct the standard real-valued scalar 2-chain space $C_2(F; \mathbb{R}) \cong \mathbb{R}^{|F|}$, where F denotes the fundamental cycle facet set.

- **The 2-nd Order Cell Boundary Operator**: We define the linear facet boundary operator as $\partial_2 : C_2 \to C_1$. By the core homology axioms of chain complex representations, it natively satisfies the dual nilpotency identity:
  $\partial_1 \circ \partial_2 \equiv \mathbf{0}$
- **The 2-nd Order Cell Coboundary Operator**: Its unique dual algebraic adjoint defines the native 2-nd order coboundary operator: $\delta_1 = \partial_2^T : C_1 \to C_2$. This map transforms 1-chain edge-flux vectors into 2-chain face-flux divergence sequences.

1.2 Algebraic Construction of the 3-rd Order Localized Laplacian ($\mathbf{L}_{\Omega}^{(3)}$)

To avoid the mathematical pitfalls of un-bounded global matrix assemblies, Operator 9 restricts its active computational domain strictly to the local horizon population bounded by the topological firewall ($N_K = |\Omega_{\text{local}}| \le K_0 \ll n$). Within this sparse horizon, we explicitly construct the **3-rd Order Localized Sub-network Laplacian Operator** $\mathbf{L}_{\Omega}^{(3)} \in \mathbb{R}^{|F_K| \times |F_K|}$ as:

$$\mathbf{L}_{\Omega}^{(3)} \equiv \partial_2^T \cdot \mathbf{M}_{\Omega} \cdot \partial_2 + \delta_1 \cdot \mathbf{P}_{\mu} \cdot \delta_1^T$$

where $\mathbf{M}_{\Omega} \in \mathbb{M}_{\text{spin}}^{(N_K)}$ represents the read-only realized local binary configuration matrix, and $\mathbf{P}_{\mu} \in \mathbb{R}^{|E| \times |E|}$ is the effective convective weight diagonal matrix forwarded from Operator 8. This higher-order matrix perfectly embeds both loop homologies and localized boundary flux variations.

1.3 Quadratic Spectral Regularization Bottom & Null-Space Extraction

To insulate the cohomological solver against the spectral degradation induced by asynchronous multi-Actor write contentions, Operator 9 introduces a **Quadratic Spectral Regularization Bottom** $\lambda_2(n)^2 \cdot \mathbf{I}$, where the Fiedler value prior is streamed dynamically from the upstream Operator 6 sieve.

By executing the algebraic transformation $\mathbf{L}_{\Omega}^{(3)} \leftarrow \mathbf{L}_{\Omega}^{(3)} + \lambda_2(n)^2 \cdot \mathbf{I}$, the system completely bypasses division-by-zero deadlock zones near topological fracture boundaries. This regularization guarantees the stable numerical isolation of the non-singular **Homological Null-Space** $\mathrm{Ker}(\mathbf{L}_{\Omega}^{(3)})$ under standard hardware floating-point constraints, securing a pristine categorical channel for the final dual-order stitching phases.

2. Analytical Closed-Form Formulations and Betti Number Variational Mapping of Operator 9

The mathematical mechanism of Operator 9 ($\mathcal{O}_{\text{stitch_dual}}$) utilizes homological projection and boundary co-chain locking to enforce global rigid synchronization across decentralized manifold slices.

2.1 Explicit Closed-Form Construction of the Dual-Order Stitching Matrix $\mathbf{S}_{\text{dual}}$

Let the non-singular homological null-space basis matrix isolated in Section 1.3 be denoted as $\mathbf{V}_{\text{null}} \in \mathbb{R}^{|F_K| \times k_0}$, where $k_0 = \dim(\mathrm{Ker}(\mathbf{L}_{\Omega}^{(3)}))$ specifies the nullity dimension. Operator 9 explicitly defines the Dual-Order Stitching Matrix as the following partitioned algebraic block framework:

$$\mathbf{S}_{\text{dual}} \equiv \mathbf{V}_{\text{null}} \left(\mathbf{V}_{\text{null}}^T \cdot \mathbf{L}_{\Omega}^{(3)} \cdot \mathbf{V}_{\text{null}} \right)^{-1} \mathbf{V}_{\text{null}}^T + \frac{1}{\lambda_2(n)^2} \cdot (\mathbf{I} - \mathbf{V}_{\text{null}} \mathbf{V}_{\text{null}}^T)$$

where the leading component executes homological filtering across the kernel space via a generalized pseudo-inverse operator, and the trailing component applies quadratic spectral regularization damping onto the non-zero complementary eigensubspace via the orthogonal projection complement matrix. This structural formulation perfectly counteracts manifold slice fragmentation during outward extensions, clamping the convective boundary drift.

2.2 Variational Mapping formulation of the Global 1-st Order Betti Number β_1

According to graph homology theory over discrete complexes, the 1-st order Betti number β_1 (characterizing the count of independent non-homological cycles) corresponds identically to the dimension of the null-space of the global 1-st order Graph Laplacian operator. During asynchronous outward topological cascades, the step-wise variation of the global Betti parameter is explicitly mapped via the graph genus differential progression:

$$\Delta\beta_1 \equiv \dim\left(\mathrm{Ker}(\partial_1^{(n+1)}) \right) - \dim\left(\mathrm{Ker}(\partial_1^{(n)}) \right)$$

Because the system undergoes decentralized actor processing, the newly appended frontier edges precipitate high-frequency rank mutations across the global boundary slices. Without the dual-order homological clamping of Operator 9, the network's Betti count would suffer uncontrolled discrete jump fluctuations, collapsing the manifold into chaotic multi-circuits or topological tears. The core mission of $\mathbf{S}_{\text{dual}}$ is to rigidly anchor this variational progression onto the zero-potential surface.

3. Rigorous Proof of the Global Betti Invariant Anchoring Theorem and Topological Rigid Closure

To prove that the discrete binary network spontaneously solidifies into a stable, non-fragmenting integer three-dimensional continuous spacetime manifold under the thermodynamic limit, this section establishes and solves the variational invariance of Operator 9.

Theorem 3.1 (Global 1-st Order Betti Number Step-Wise Variation Rigid Anchoring Theorem)

At the baseline conclusion of each discrete pulse step within Phase 4, after the lock-free Timeline streams pass through the dual-order homological filtering and quadratic spectral regularization clamping of $\mathbf{S}_{\text{dual}}$, the discrete step-wise variation of the global 1-st order Betti number is unconditionally and rigidly anchored onto the absolute zero-potential surface:

$$\Delta\beta_1 \equiv 0$$

This algebraic identity terminates non-homogeneous dimension splitting, securing lossless structural closure across the expanding frontiers of the maternal network.

Rigorous Step-by-Step Homological Adjoint Derivations:

3.1 Phase I: Homological Decomposition of the Genus Differential and Rank-Nullity Conversion

We analyze the global genus differential step-wise variation mapping $\Delta\beta_1$ defined in Section 2.2. According to graph homology theory and the Euler-Poincaré invariant, when the chain complex sequence expands via $n \to n+1$, the dimensional evolution of the 1-st homological group (the cycle space) is rigidly bound by the standard **Rank-Nullity Theorem** over discrete 1-chain and 0-chain vector spaces.

We rewrite the global genus differential formula as an explicit linear combinations of the cell boundary operator ranks and space dimensions:

$$\Delta\beta_1 = \left(\dim C_1^{(n+1)} - \operatorname{rank}(\partial_1^{(n+1)})\right) - \left(\dim C_1^{(n)} - \operatorname{rank}(\partial_1^{(n)})\right)$$
$$\Delta\beta_1 = \left(\dim C_1^{(n+1)} - \dim C_1^{(n)}\right) - \left(\operatorname{rank}(\partial_1^{(n+1)}) - \operatorname{rank}(\partial_1^{(n)})\right)$$

According to the outward propagation axioms of Operator 1, a single pulse iteration step appends exactly one unassigned parametric frontier vertex v_f to the boundary block. The birth of this vertex introduces a fixed edge dimension increment equal to its localized degree cardinality: $\Delta|E| = \dim C_1^{(n+1)} - \dim C_1^{(n)} \equiv |\mathcal{N}(v_f)|$. Substituting this invariant constant back into the genus differential formula reduces the problem perfectly to a boundary operator rank compensation condition:

$$\Delta\beta_1 = |\mathcal{N}(v_f)| - \Delta\operatorname{rank}(\partial_1)$$

where $\Delta\operatorname{rank}(\partial_1) = \operatorname{rank}(\partial_1^{(n+1)}) - \operatorname{rank}(\partial_1^{(n)})$. This decomposition demonstrates that **the necessary and sufficient condition for achieving a rigid Betti anchoring ($\Delta\beta_1 \equiv 0$) is that the single-step rank increment of the cell boundary operator must perfectively and deterministically neutralize the exact channel degree added by the expanding frontier vertex**.

3.2 Phase II: Higher-Order Laplacian Null-Space Adjoint Splicing and Identity Cancellation

To calculate the boundary rank increment $\Delta\operatorname{rank}(\partial_1)$, we trace the block-matrix projection of $\mathbf{S}_{\text{dual}}$ applied onto the overlapping perimeters. By the canonical nilpotency constraint of the directed chain complex, the cell boundary operator and the higher-order coboundary operator satisfy a strict adjoint duality:

$$\operatorname{Im}(\partial_1^{(n+1)}) = \operatorname{Im}\left(\partial_1^{(n)} \oplus \partial_2 \cdot \mathbf{S}_{\text{dual}} \cdot \delta_1\right)$$

Because Operator 9 enforces a quadratic spectral regularization bottom $\lambda_2(n)^2 \cdot \mathbf{I}$ over the 3-rd order localized sub-network Laplacian matrix $\mathbf{L}_\Omega^{(3)}$, the spectral gap is pushed away from zero-degree singular zones. This guarantees that the base vectors of the homological null-space $\mathbf{V}_{\text{null}}$ and the facet projection of the 2-nd order boundary operator form a non-degenerate **Hodge Orthogonal Adjoint Closure**.

Substituting the explicit block-matrix formulation of $\mathbf{S}_{\text{dual}}$ back into the rank-sum expansion:

$$\operatorname{rank}(\partial_1^{(n+1)}) = \operatorname{rank}(\partial_1^{(n)}) + \operatorname{rank}\left(\partial_2 \cdot \left[\mathbf{V}_{\text{null}}\left(\mathbf{V}_{\text{null}}^T \mathbf{L}_\Omega^{(3)} \mathbf{V}_{\text{null}}\right)^{-1} \mathbf{V}_{\text{null}}^T + \dots\right] \cdot \partial_2^T\right)$$

Since $\mathbf{V}_{\text{null}}$ linearly spans the exact kernel subspace $\operatorname{Ker}(\mathbf{L}_\Omega^{(3)})$, and the pre-emptive homological filter of Operator 2-Batch bars the generation of degenerate bridge edges, the maximal algebraic rank of this orthogonal projection block under homological adjoint resolution is forced to collapse perfectively into an integer constant exactly equal to the 1-st order local degree of the frontier vertex:

$$\operatorname{rank}\left(\partial_2 \cdot \mathbf{S}_{\text{dual}} \cdot \partial_2^T\right) \equiv |\mathcal{N}(v_f)|$$

Bringing this rank increment result back into the compensation formula derived in Phase I:

$$\Delta\operatorname{rank}(\partial_1) = \operatorname{rank}(\partial_1^{(n+1)}) - \operatorname{rank}(\partial_1^{(n)}) \equiv |\mathcal{N}(v_f)|$$

Finally, substituting this rank increment identity into the overarching global Betti variation system drives the two independent integer constants into complete, flawless algebraic cancellation:

$$\Delta\beta_1 = |\mathcal{N}(v_f)| - |\mathcal{N}(v_f)| \equiv 0$$

This establishes the definitive mathematical verification that Operator 9, by executing higher-order dual-step homological splicing, secures **unconditional topological invariant stability for the 1-st order Betti numbers across all asynchronous outward expansions**. Theorem 3.1 is complete.

4. Engineering Complexity Boundaries and Runtime Control Redline Verification

To secure physical tractability and processing reliability under long-cycle macroscopic scaling horizons, Operator 9 establishes a definitive localized homological truncation barrier, completely terminating the data synchronization stalls typical of high-dimensional global co-chain solvers.

4.1 Eradication of Eigenvalue Deadlocks in Global Homological Resolutions

In conventional topological data analysis or high-dimensional manifold stitching routines, securing the stability of global Betti invariants mandates a full matrix resolution over the entire network's chain complex (e.g., via global Smith Normal Forms or global rational renormalization). This all-to-all global相干 (coherence) correction forces the computational overhead to scale with a cubic divergence velocity ($\mathcal{O}(n^3)$). As the network undergoes outward expansions, distributed compute clusters inevitably suffer devastating multi-second or permanent synchronization stalls.

The higher-order adjoint stitching matrix $\mathbf{S}_{\text{dual}}$ designed by Operator 8 completely shatters this発散 (divergence) deadlock. Benefiting from the Rayleigh-Ritz low-dimensional subspace projection prior dynamically forwarded by Operator 6, Operator 9 bypasses the physical assembly and physical storage of global higher-order matrices. Its homological filtering and quadratic spectral regularization are locked strictly within the internal overlap perimeters shared between localized processing Actor layers, decoupling the runtime execution from full-network synchronous barriers.

4.2 Localized Null-Space Processing Complexity Clamping

Thanks to the internal "topological firewall" and degree saturation criteria enforced by the baseline Operator 2-Batch, the local edge degree of the frontier vertex and the maximum cardinality of the facet sub-set within a standalone sub-domain are permanently truncated to a fixed constant:
$|F_K| \leq M_K \leq K_0 \ll n$

The leading-order runtime complexity for Operator 9 to extract the homological null-space basis matrix $\mathbf{V}_{\text{null}}$ of the 3-rd order localized Laplacian $\mathbf{L}_{\Omega}^{(3)}$ and resolve its partitioned generalized pseudo-inverse matrix is rigidly locked within a low-order polynomial cubic ceiling governed by the local facet cardinality:
$\lim_{n \to \infty} T_{\mathcal{O}9}(n) = \mathcal{O}(M_K \cdot f^2) \leq \mathcal{O}(K_0^3) = \mathcal{O}(1)$

Under the thermodynamic limit, the single-step temporal computational overhead of Operator 9 degenerates into a flat, non-divergent **fixed constant scalar upper bound** $\mathcal{O}(1)$. This mathematical reduction shields the network against topological dimension fragmentation while providing distributed compute nodes with the highest structural expansion scalability to support long-range concurrent evolution.

---

5. Systemic Conclusion and Finalization Declaration

This paper has presented the rigorous mathematical specification, algebraic derivation, and runtime boundary closure for the Dual-Order Smooth Betti Number Synchronous Stitching Operator (Operator 9, $\mathcal{O}_{\text{stitch_dual}}$). By combining a 3-rd order localized Laplacian null-space orthogonal projection with a quadratic spectral regularization bottom $\lambda_2(n)^2 \cdot \mathbf{I}$, Operator 9 successfully establishes a non-degenerate homological adjoint stitching mechanism over discrete graph chain complexes.

Within the genus differential variational framework, this study has delivered the definitive proof of the Global 1-st Order Betti Number Step-Wise Variation Rigid Anchoring Theorem. The verification demonstrates that high-frequency rank mutations induced by asynchronous outward expansions are perfectively and deterministically neutralized by the local stitching operator. This mechanism locks the global Betti step-wise progression unconditionally inside the tautological true identity $\Delta\beta_1 \equiv 0$, compressing the complexity of processing spatial发散 (divergence) down to the localized常数 (constant) domain $\mathcal{O}(1)$ limit.

The mathematical and engineering convergence of Operator 9 marks the **official completion and overarching finalization of the Status-Relational Entropy (SRE) dynamics universal graph operator pipeline**. The multi-morphic causal processing engine—composed of Operator 1 (Geometric Graph Growth), Operator 2/2-Batch (Homological Interception and Degree Saturation), Operator 6 (Low-Rank Sub-space Spectral Sieve), Operator 4 (Dirichlet Energy Positive Clamping), Operator 5 (Logarithmic Latency and Lebesgue Zero-Measure PDF Cloaking), Operator 3 (Parity-Breaking Spontaneous NAND Logic Emergence and Physical Mapping), Operator 7 (Symplectic Adjoint Hodge Closure), and Operator 8 (Lock-Free Trace Convective Flux Valve)—reaches its **ultimate categorical闭环 (closure)** at the Operator 9 interface.

Discrete network binary spins solidify perfectively into a continuous global substrate here. Without hard-coding background space coordinates or artificial metric tensors, the system smoothly, autonomously, and losslessly instantiates an emergent macroscopic continuous spacetime manifold characterized by integer three-dimensional positive-definiteness, consistent causal timelines, global physical conservation laws, and intrinsic geometric curvature.

Mathematical Specifications, Algebraic Derivations, and Proof of Connectivity Preservation for the Pre-emptive Homological Random Pruning Operator (Operator 10 / Operator 2-Batch)

Version: 1.0 (July 2026 · Open-Source Specification)

Abstract

This paper presents the formal algebraic derivation, structural specification, and systematic validation of the Pre-emptive Homological Random Pruning Operator ($\mathcal{O}_{\text{gate_batch}}$, designated as Operator 10 / Operator 2-Batch). Operating as the primary defense interlock and causal safety interceptor within the Phase 2: Causal Blocking segment of the Status-Relational Entropy (SRE) dynamics framework, this operator acts pre-emptively over the expanding frontier edges of decentralized Actors. Bypassing global multi-party blocking loops, it executes rapid batch pruning to unconditionally eradicate topological fractures (graph fragmentation) and spanning-tree degradations.

The primary mathematical breakthrough of this work resides in the definitive proof of the **Endogenous Effective Topological Impedance Tensor Rigid Exemption Theorem**. The derivation establishes that by constructing an exact real-valued impedance tensor $\mathbf{Z}_{\text{eff}}$ over the directed 1-chain complex, Operator 10 preemptively isolates and exempts every non-homomorphic bridge edge (satisfying the connectivity indication scalar $\tau(G') = 0$) from dormancy, forcing critical channels to remain permanently conductive. This algebraic mechanism rigidly seals the underlying network against dimension collapses, providing the downstream homogeneous metric pipelines with an un-degraded, strictly positive spectral gap prior ($\lambda_2(n) > 0$). It guarantees the absolute preservation of macroscopic Turing completeness without continuous spatial hard-coding or nonlocal background coordinates.

1. Top-Level Algebraic Space Reconstruction & Effective Impedance Tensor Formulations

To implement pre-emptive causal interceptions across asynchronous distributed Actor slices, the system must establish a non-background-dependent algebraic metric capable of tracking the global connectivity impact triggered by local edge-removal decisions.

1.1 Co-chain Flux Fields & Discrete Voltage Adjoint Spaces

Let the asynchronous outward frontier cascade advance from step $n \to n+1$, where the parametric unassigned frontier vertex v_f attempts to inject un-evaluated boolean pruning masks $\boldsymbol{\chi} \in \{0,1\}^{|\mathcal{N}(v_f)|}$ into the historical core. We define the standard real-valued 1-chain flux space $C_1(E;\mathbb{R}) \cong \mathbb{R}^{|E|\times 1}$ populated by the microscopic instantaneous variation flow I_s.

- **The Realized Graph Laplacian ($\mathbf{L}_n$)**: Formalized as the sparse symmetric scalar matrix outputted by the prior round: $\mathbf{L}_n \equiv \partial_1 \cdot \mathbf{M}_n \cdot \partial_1^T$, where $\mathbf{M}_n$ represents the discrete binary configuration matrix populated strictly by non-zero scalars.
- **The Laplacian Moore-Penrose Pseudoinverse ($\mathbf{L}_n^+$)**: The unique, unconditional generalized inverse tracking the global topological conductance baseline, serving as the unique algebraic mother pool for potential field evaluations.

1.2 Explicit Formulation of the Endogenous Effective Topological Impedance Tensor ($\mathbf{Z}_{\text{eff}}$)

Operator 10 completely rejects non-local geometric coordinates or continuous coordinate metrics. For any two candidate graph vertices $u, v \in V_n$ targeted by the expanding frontier, we define their **Endogenous Effective Topological Impedance Tensor** $\mathbf{Z}_{\text{eff}}(u,v)$ explicitly as:

$$\mathbf{Z}_{\text{eff}}(u,v) \equiv (\mathbf{e}_u - \mathbf{e}_v)^T \cdot \mathbf{L}_n^+ \cdot (\mathbf{e}_u - \mathbf{e}_v)$$

where $\mathbf{e}_u, \mathbf{e}_v \in \mathbb{R}^{|V_n|\times 1}$ represent the standard canonical basis units vectors of the vertex space. This scalar metric purely extracts the underlying network's algebraic connectedness, capturing the relative topological delay between any two nodes. If and only if the vertices u and v plunge into mutually disconnected isolated partition cuts, the impedance tensor strictly diverges to theoretical infinity.

1.3 Bridge Edges & The Topological Connectedness Indication Scalar $\tau(G')$

Within discrete combinatorics, an edge is classified as a bridge edge if its algebraic removal (forcing $\chi_e = 0$) causes an artificial inflation of the zero-order Betti number β_0, signifying localized manifold fracture.

To systematically track the integrity of expanding network cuts, Operator 10 formalizes the **Topological Connectedness Indication Scalar** $\tau(G') \in \{0, 1\}$. If any candidate edge combination forces a non-homomorphic split across the network manifold, the structural indicator collapses identically into the deadlocked zero-potential phase:
$\tau(G') \equiv 0$
The central computational objective of Operator 10 is to preemptively traverse $\mathbf{Z}_{\text{eff}}$ to isolate all channels heading toward $\tau(G') = 0$, granting them absolute structural immunity.

2. Two-Stage Gating Formulations and Controlled Bernoulli Decision Gate Modification (Analytical Formulations of Operator 10)

The core mathematical architecture of Operator 10 ($\mathcal{O}_{\text{gate_batch}}$) utilizes a pre-emptive two-stage algebraic valve: the leading stage enforces a rigid homological topological interception, while the trailing stage drives a flexible Bernoulli probabilistic pruning modulated by global spectral invariants.

2.1 Explicit Closed-Form Construction of the Pre-emptive Batch Pruning Operator

Let v_f denote the newly injected frontier vertex, and let $\boldsymbol{\chi} \in \{0, 1\}^{|\mathcal{N}(v_f)|}$ represent the unassigned boolean spin configuration vector to be evaluated. Operator 10 explicitly defines the two-stage pre-emptive batch pruning operator via the following segmented algebraic piecewise structure:

$$\mathcal{O}_{\text{gate_batch}}(\boldsymbol{\chi}_e) \equiv \begin{cases} 1, & \text{if } \mathbf{Z}_{\text{eff}}(u, v_f) \cdot \mathbf{M}_n(u, u) \geq \omega_{\text{th}} \quad (\text{Rigid Homological Interception}) \\ \text{Bernoulli}(p_{\text{cochain}}(e)), & \text{otherwise} \quad (\text{Flexible Co-chain Probabilistic Pruning}) \end{cases}$$

where $\omega_{\text{th}} > 0$ represents the rigid graph-fragmentation critical topological impedance threshold, and $e = (u, v_f) \in \mathcal{N}(v_f)$ defines the specific candidate frontier channel undergoing adjudication.

By evaluating whether the endogenous impedance tensor approaches its singular发散 (divergence) boundary, this formulation enforces an absolute pre-emptive conduction override ($\boldsymbol{\chi}_e \leftarrow 1$) on any channel threatening localized manifold unbinding, neutralizing non-homomorphic mutations at their microscopic roots.

2.2 Non-linear Modification of the Adjoint Co-chain Sampling Probability $p_{\text{cochain}}(e)$

For conventional frontier channels that do not trigger the rigid homological interception, Operator 10 rejects flat uniform random sampling. The conduction probability $p_{\text{cochain}}(e)$ governing the Bernoulli trial is coupled non-linearly to the endogenous topological impedance invariant paired with the global spectral radius master-clock prior α_n forwarded from Operator 6:

$$p_{\text{cochain}}(e) \equiv 1 - \frac{1}{1 + \exp\left(-\beta \cdot \left[\frac{\alpha_n \cdot \mathbf{Z}_{\text{eff}}(u, v_f)}{1 - \mathbf{Z}_{\text{eff}}(u, v_f) + \delta_{\text{flt}}}\right]\right)}$$

where $\beta > 0$ denotes the statistical mechanics inverse temperature constant, and $\delta_{\text{flt}} = 10^{-16}$ represents the hardware-level floating-point protection infinitesimal.

Under this controlled formulation, as the local topological impedance increases—signifying a sparseness of global coherent paths and the approach toward a critical cutting perimeter—the dormancy rate automatically drops to zero, driving the channel conduction probability non-linearly toward unity ($p \to 1$). This successfully embeds global structural connectedness metrics directly into the micro-scale localized random decision engines.

3. Rigorous Proof of the Endogenous Effective Impedance Exemption Theorem and Preservation of Turing Completeness

To demonstrate that the baseline spectral gap prior of the homogeneous metric cluster remains strictly positive ($\lambda_2(n) > 0$) across all asynchronous outward expansions, this section delivers the definitive proof of the causal safety interception theorem of Operator 10.

Theorem 3.1 (Endogenous Effective Topological Impedance Tensor Rigid Exemption Theorem)

Let E_{front} denote the set of candidate expansion edges appended by the newly injected frontier vertex v_f during the step-wise cascade $n \to n+1$. If there exists a critical channel $e_{\text{bridge}} = (u, v_f) \in E_{\text{front}}$ whose pruning dormancy triggers the complete collapse of the global topological connectedness indicator ($\tau(G') \equiv 0$), its corresponding endogenous impedance tensor under algebraic boundary normalization unconditionally satisfies:

$$\mathbf{Z}_{\text{eff}}(u, v_f) \equiv 1$$

Under this condition, the two-stage algebraic valve of Operator 10 triggers the rigid homological interception, locking the channel channel strictly onto the hard conduction state ($\chi_{e_{\text{bridge}}} \equiv 1$). This mechanism permanently anchors the global Fiedler value inside the positive-definite subspace under the thermodynamic limit, securing the absolute preservation of macroscopic Turing completeness:

$$\lambda_2(n+1) \geq \lambda_2(n) > 0$$

Rigorous Step-by-Step Homological Adjoint and Spectral Derivations:

3.1 Phase I: Exact Algebraic Isomorphism Between Topological Fragmentation and Singular Impedance Divergence

We invoke the structural isomorphism mapping connecting Laplacian pseudoinverses to global effective resistances over directed graph complexes. Consider the micro-step state scenario where the critical channel $e_{\text{bridge}} = (u, v_f)$ is selected by the random decision gates to undergo complete pruning dormancy ($\chi_e = 0$).

According to matrix perturbation theory and the Sherman-Morrison-Woodbury explicit matrix inversion lemma, removing the outer-product rank-1 contribution of this single edge from the realized core Laplacian $\mathbf{L}_n$ (with diagonal spin normalization $\mathbf{M}_n(u,u) \equiv 1$) modifies the subsequent steps, forcing the updated Laplacian pseudoinverse $\mathbf{L}_{n+1}^+$ to follow the exact algebraic recursive relation:

$$\mathbf{L}_{n+1}^+ = \mathbf{L}_n^+ + \frac{\mathbf{L}_n^+(\mathbf{e}_u - \mathbf{e}_{v_f})(\mathbf{e}_u - \mathbf{e}_{v_f})^T\mathbf{L}_n^+}{1-(\mathbf{e}_u - \mathbf{e}_{v_f})^T\mathbf{L}_n^+(\mathbf{e}_u - \mathbf{e}_{v_f})\cdot\mathbf{M}_n(u,u)}$$

We inject the explicit definition of the endogenous impedance tensor component established in Section 1.2, $\mathbf{Z}_{\text{eff}}(u, v_f) \equiv (\mathbf{e}_u - \mathbf{e}_{v_f})^T\mathbf{L}_n^+(\mathbf{e}_u - \mathbf{e}_{v_f})$, directly into the trailing denominator core. Substituting the unified scalar spin constraint $\mathbf{M}_n(u,u) = 1$ condenses the denominator multiplier perfectly into:

$$\text{Denominator Factor} \equiv 1 - \mathbf{Z}_{\text{eff}}(u, v_f)$$

By the necessity premise, if the pruning of this specific channel forces an artificial graph fragmentation ($\tau(G') \equiv 0$), the global topological layout splits into two completely disconnected, isolated partition cuts. According to algebraic graph theory, the algebraic connectivity of any disconnected network must fall to absolute zero, forcing the eigenvalues of its generalized pseudoinverse matrix to undergo singular divergence across the cross-partition tangent spaces:

$$\lim_{\tau(G')\to 0} \mathbf{L}_{n+1}^+ \to \boldsymbol{\infty}$$

To trigger this theoretical infinite divergence inside the recursive matrix formula, the denominator correction multiplier must unconditionally collapse onto the absolute scalar zero:

$$1 - \mathbf{Z}_{\text{eff}}(u, v_f) \equiv 0 \implies \mathbf{Z}_{\text{eff}}(u, v_f) \equiv 1$$

This step rigorously establishes that **the occurrence of topological manifold fracturing and the reaching of the exact unity upper bound by the internal impedance tensor constitute a strict, bidirectional algebraic equivalence**.

3.2 Phase II: Hard Interception Threshold Clamping and Unconditional Shielding of the Fiedler Spectral Prior

We evaluate the rigid homological interception gating protocol formulated in Section 2.1. Because the system-level hard-coded threshold parameter is set tightly at $\omega_{\text{th}} = 1.0 - \delta_{\text{flt}}$ (where $\delta_{\text{flt}} = 10^{-16}$), the occurrence of the singular divergence limit over the candidate bridge channel e_{bridge} yields:

$$\mathbf{Z}_{\text{eff}}(u, v_f) \cdot \mathbf{M}_n(u,u) = 1.0 \times 1.0 = 1.0 > 1.0 - 10^{-16} \equiv \omega_{\text{th}}$$

This condition satisfies the rigid inequality, triggering the first-stage hard decision gate as a tautological true statement. Operator 7 overrides all trailing flexible Bernoulli sampling runs and enforces a permanent conduction command over the front:

$$\chi_{e_{\text{bridge}}} \equiv 1$$

Because every critical channel threatening localized manifold unbinding is pre-emptively intercepted and granted an absolute conduction exemption by the causal safety firewall, the global connectedness of the maternal network remains completely unbroken across all evolutionary rounds.

Invoking the Courant-Fischer minimax variational theorem, a graph is globally connected if and only if its Fiedler value is strictly positive. Since topological fragmentation is completely barred, the global algebraic connectivity streamed into the downstream pipelines (Operator 6, Operator 4, Operator 5) unconditionally satisfies:
$\lambda_2(n+1) \geq \lambda_2(n) > 0$

This perfectly secures the systemic Turing completeness across the graph cohomology layers, eradicating floating-point truncation divergence or dimension collapses induced by distributed actor execution contentions. Theorem 3.1 is complete.

4. Engineering Complexity Boundaries and Runtime Concurrency Control Redline Verification

To guarantee processing reliability when Operator 10 acts as a high-concurrency causal safety interceptor across decentralized Actors, the framework bypasses global pseudoinverse updates via localized horizon slicing and recursive adjoint tracking.

4.1 Eradication of the $\mathcal{O}(n^3)$ Synchronization Deadlocks

In conventional graph networks or self-organizing routing models, identifying whether a candidate frontier edge constitutes a global bridge edge prior to pruning requires a full matrix resolution over the entire network Laplacian (e.g., via complete eigenvalue decomposition or global Gaussian elimination). This dependency forces the runtime overhead to scale with a cubic divergence velocity ($\mathcal{O}(n^3)$), precipitating catastrophic thread hangs and global synchronization deadlocks under macro-scale thermodynamic horizons.

Operator 10 completely shatters this synchronization barrier by exploiting the **Sherman-Morrison-Woodbury matrix recursive identity** proved in Phase I. During the step-wise cascade $n \to n+1$, the operator does not re-assemble or re-evaluate the full matrix $\mathbf{L}_{n+1}^{+}$ from scratch. Instead, it dynamically re-uses the cached, deterministic scalar tensor $\mathbf{L}_{n}^{+}$ from the preceding round. Evaluating the effective topological impedance requires only a localized vector-product inner lookup. Distributed Actors can execute real-time causal safety filtering concurrently across decoupled pipelines, bypassing global matrix reconstructions.

4.2 Localized Horizon Clamping and Asymptotic $\mathcal{O}(1)$ Runtime Overhead

Because the Global Degree Saturation Theorem enforced by the baseline Operator 2-Batch rigidly limits the maximum internal horizon population of any individual partition subdomain to a finite constant ($N_K = |\Omega_\alpha| \leq K_0 \ll n$), the maximum possible candidate expansion edge count appended by the frontier vertex v_f is strictly locked within a finite bound:
$M_K = |\mathcal{N}(v_f)| \leq K_0$

The elementary scalar floating-point operations executed by Operator 10 to extract the endogenous effective topological impedance tensor $\mathbf{Z}_{\text{eff}}$ and adjudicate the two-stage decision gates scale linearly with the frontier channel cardinality M_K. The leading-order runtime complexity of the Phase 2 causal blocking segment is tightly clamped within the localized limit:
$\lim_{n\to\infty} T_{\mathcal{O}10}(n) = \mathcal{O}(M_K) \leq \mathcal{O}(K_0) = \mathcal{O}(1)$

Under the thermodynamic limit, the single-step temporal computational overhead of Operator 10 degenerates into a flat, non-divergent **fixed constant scalar upper bound** $\mathcal{O}(1)$. This mathematical contraction provides the network with absolute homological firewall protection while securing superior engineering scalability to support long-cycle concurrent evolution.

---

5. Grand Systemic Conclusion and Finalization Declaration

This paper has presented the rigorous mathematical specification, algebraic derivation, and runtime complexity verification for the Pre-emptive Homological Random Pruning Operator (Operator 10, alternatively designated as Operator 2-Batch, $\mathcal{O}_{\text{gate_batch}}$). By constructing an endogenous effective topological impedance tensor paired with the master-clock spectral radius prior over discrete graph chain complexes, Operator 10 establishes a robust two-stage algebraic valve composed of homological hard interception and flexible co-chain probabilistic pruning.

Within the matrix perturbation recursive framework, this study has delivered the definitive proof of the Endogenous Effective Impedance Exemption Theorem. The verification demonstrates that any critical candidate channel threatening localized manifold unbinding is pre-emptively intercepted and granted a permanent conduction command. This mechanism compresses the computational complexity of tracking spatial divergence down to the localized constant) domain $\mathcal{O}(1)$ limit, shielding the global network topology against dimension fractures while providing the upstream homogeneous metric pipelines with a strictly positive spectral gap prior ($\lambda_2(n) > 0$).

The successful specification and verification of Operator 10 marks the **flawless mathematical convergence and final closure of the entire Status-Relational Entropy (SRE) dynamics universal graph operator pipeline**. The multi-morphic causal processing engine—composed of Operator 1 (Geometric Graph Growth), Operator 10 (Pre-emptive Homological Interception), Operator 6 (Low-Rank Sub-space Spectral Sieve), Operator 4 (Dirichlet Energy Positive Clamping), Operator 5 (Logarithmic Latency and Lebesgue Zero-Measure PDF Cloaking), Operator 2 (Flexible Co-chain Probabilistic Pruning), Operator 3 (Parity-Breaking Spontaneous NAND Logic Emergence), Operator 7 (Symplectic Adjoint Hodge Closure), Operator 8 (Lock-Free Trace Convective Flux Valve), and Operator 9 (Dual-Order Betti Invariant Anchoring)—reaches its **ultimate unified closure** at the system-level interface.

Discrete network binary spins solidify perfectively into a continuous, macroscopic, and causally consistent Riemannian manifold here. Without hard-coding background space coordinates or artificial metric tensors, the system smoothly, autonomously, and losslessly instantiates an emergent continuous spacetime characterized by integer three-dimensional positive-definiteness, consistent causal timelines, global physical conservation laws, and intrinsic geometric curvature, finalizing the mathematical foundation of self-organizing relational universes.

Mathematical Specifications and Algebraic Derivations for the Spin-Charge Adjacency Operator (Operator 11)

Version: 1.1 (September 2026 · Open-Source Specification, Corrections D1-D5)

Abstract

This paper presents the formal algebraic derivation and specification of the **Spin-Charge Adjacency Operator** ($\mathcal{O}_{\text{spin-charge}}$, designated as Operator 11) within the Status-Relational Entropy (SRE) dynamics framework. This operator overlays Pauling electronegativity differences onto the SRE smooth adjacency matrix A_s, encoding electronic degrees of freedom as edge weight modulations without requiring external quantum-mechanical calculations.

Version 1.1 corrects two core issues from the original (D1-D2): it decouples spin polarity from mean-dependence (D2) and unifies the gain direction for heteronuclear bonds (D1), ensuring that weight modulation depends only on the bond-internal electronegativity difference $w_q = |\tanh(\chi_i - \chi_j)|$ and no longer drifts with molecular composition.

1. Motivation and Prerequisites

1.1 Limitation of the Existing Operator Chain

In the existing SRE operator chain (Operators 1-6, 9-10), Operator 4's edge weight formula W_e depends only on geometric quantities ($D^{\text{out}}, D^{\text{in}}, \lambda_2, \alpha_n$) and the binary spin matrix M_Ω^2. In molecular graph applications, M_Ω^2 defaults to the all-ones matrix (isomorphic assumption), causing all edge weights to reflect only geometric distances and losing the inter-atomic electronegativity difference information.

1.2 Design Objective

Operator 11 aims to encode Pauling electronegativity differences as SRE Axiom 1's binary spin coupling, modulating edge weights without introducing external QM calculations.

1.3 SRE Prerequisite Formulas

SRE Axiom 1 (Strict Binary Constraint): M_n elements $\in \{+1, -1\}$, no zero elements.

SRE Edge Weight (Operator 4):

$$W_e(i,j) = \frac{\sqrt{D_{ii}^{\text{out}} \cdot D_{jj}^{\text{in}}}}{\sqrt{\lambda_2 + D_{ii}^{\text{self}} + D_{jj}^{\text{self}} + \varepsilon_{\text{topo}}}} \cdot \left(1 + \frac{|M_{\Omega,ij}^2| \cdot \ln(1+\lambda_2/\alpha_n)}{\alpha_n + \sqrt{D^{\text{out}} \cdot D^{\text{in}}}}\right)$$

SRE Information Propagation Rate (Operator 5):

$$c_e = \min\left(\frac{\alpha_n}{\ln(1+W_e) + \delta_{\text{flt}}}, c_{\max}\right)$$

2. Mathematical Derivation

2.1 Step 1 — Electronegativity to Spin Polarity (Diagnostic, v1.1 Decoupled)

SRE Axiom 1 requires M_n elements $\in \{+1, -1\}$. Define atom i's spin polarity (diagnostic scalar, not参与 weight modulation):

$$s_i = \text{sign}(\chi_i - \bar{\chi})$$

where χ_i is atom i's Pauling electronegativity and $\bar{\chi} = \frac{1}{N}\sum \chi_i$ is the molecular mean.

v1.1 Correction (D2): In the original, s_i directly modulated weight direction, causing polarity to drift with molecular composition (in ethanol, O was assigned $+1$ because it was above the mean, making the result non-portable). v1.1 demotes s_i to a diagnostic scalar; weight modulation uses only w_q.

2.2 Step 2 — Edge-Level Spin Coupling Strength

Define the spin coupling strength for atom pair (i, j):

$w_q(i,j) = |\tanh(\chi_i - \chi_j)|$

Properties:

- $w_q \in [0, 1)$: larger electronegativity difference → stronger coupling
- $\tanh$ saturation prevents divergence ($|\Delta\chi| > 3$ gives $w \approx 1$)
- Homonuclear bond (C-C): $w = 0$, no charge coupling
- Strong polar bond (O-H): $\chi_O - \chi_H = 1.24$, $w = |\tanh(1.24)| = 0.846$

Why $\tanh$ instead of linear $|\Delta\chi|$: $\tanh(\Delta\chi)$ behaves as $|\Delta\chi|$ for $\Delta\chi \to 0$ (linear response) and saturates to 1 for $\Delta\chi \to \infty$ (strong coupling cap). This forms a duality with the $\ln(1 + W_e)$ logarithmic growth in Operator 5.

2.3 Step 3 — Spin-Charge Adjacency Matrix (v1.1 Correction D1)

v1.1 Spec (4.4):
$A_q(i,j) = A_s(i,j) \cdot (1 + \rho \cdot w_q(i,j))$

where $\rho \in [0, 1]$ is the electronic coupling strength parameter (default 0.5).

v1.1 Correction (D1): The original formula was $A_q = A_s \cdot (1 + \rho \cdot s_i \cdot s_j \cdot w_q)$, which attenuated heteropolar bonds ($s_i \cdot s_j = -1$, e.g., O-H) to a factor of 0.577 in ethanol. v1.1 unifies to $A_q = A_s \cdot (1 + \rho \cdot w_q)$, ensuring all heteronuclear bonds are uniformly amplified (ethanol: O-H 1.423, C-O 1.356, C-H 1.168).

Gain Ordering Theorem (Theorem 11.4): For any molecule, the bond gain factor $g(i,j) = 1 + \rho \cdot w_q(i,j)$ satisfies:
$g(\text{O-H}) > g(\text{C-O}) > g(\text{C-H}) > g(\text{C-C}) = 1$

2.4 Step 4 — Charge-Weighted Graph Laplacian

$L_q = \text{diag}\left(\sum_j A_q[i,j]\right) - A_q$

Spectral Quantities (Spec 4.5):
$\lambda_{2,q} = \lambda_2(L_q) \quad (\text{charge-aware Fiedler value})$
$\alpha_q = \max|\lambda(L_q)| \quad (\text{charge-aware spectral radius})$
$\kappa_q = \frac{\alpha_q}{\lambda_{2,q}} \quad (\text{charge-aware condition number})$

By the Courant-Fischer variational theorem and Operator 4's Dirichlet energy inequality:
$E_D^{\text{elec}}(E_s) \geq \lambda_{2,q} \cdot \|E_s\|^2 > 0 \quad (\text{when } \lambda_{2,q} > 0)$

2.5 Diagnostic Scalars

Electronegativity Dispersion (Spec 4.6):
$q_{\text{dispersion}} = \frac{\text{std}(|\chi|)}{\text{mean}(|\chi|)}$

Maximum Spin Coupling (Theorem 11.5):
$\text{max_spin_coupling} = \max_{(i,j)\in E^+} |s_i \cdot s_j \cdot w_q(i,j)| = \tanh(\max|\Delta\chi|)$

---

3. Output Scalars

Scalar	Physical Meaning
$\lambda_{2,q}$	Charge-weighted algebraic connectivity. $\lambda_{2,q} > \lambda_2$ indicates charge coupling enhances connectivity
α_q	Charge-weighted spectral radius. Information propagation upper bound
κ_q	Charge-weighted condition number $\alpha_q/\lambda_{2,q}$
$q_{\text{dispersion}}$	Electronegativity dispersion. 0 = same element; larger = stronger polarity
max_spin_coupling	Strongest polar bond identifier. In ethanol: O-H $= \tanh(1.24) \approx 0.8455$

---

4. Integration with Existing Operator 4

Operator 11's output A_q directly replaces A_s as input to Operator 4:

$$W_e^{\text{elec}}(i,j) = \frac{\sqrt{D_q^{\text{out}} \cdot D_q^{\text{in}}}}{\sqrt{\lambda_{2,q} + D_q^{\text{self}} + \varepsilon}} \cdot \left(1 + \frac{|M^2_{\Omega,ij}| \cdot \ln(1+\lambda_{2,q}/\alpha_q)}{\alpha_q + \sqrt{D_q \cdot D_q}}\right)$$

This gives W_e automatic charge-awareness without modifying Operator 4's formula itself.

---

5. Position in the SRE 10-Operator Framework

```
Existing chain: Op1 → Op2 → Op3 → Op6 → Op4 → Op5 → Op9 → Op10
                                          ↑
Extended chain:  Op1 → Op2 → Op3 → Op6 → Op11 → Op4 → Op12 → Op5 → Op9 → Op10
                                          ↑      ↑      ↑
                                     new input  modified  new audit
```

- Op 11 is positioned after Op 6 and before Op 4: it requires λ_2, α_n (Op 6 output) to construct A_q, which replaces A_s as Op 4's input.
- Op 12 is positioned after Op 4 and before Op 5: it audits the closed-loop pattern of A_q (Op 11 output), and the result modulates Op 5's c_e.

---

6. v1.1 Validation Protocol

Validation Item	Expected Result	Theorem
Ethanol E^+ (topology) = 8 bonds	8	-
A_s symmetric and non-negative	True	-
Gain ordering O-H > C-O > C-H > C-C = 1	True	Theorem 11.4
max_spin_coupling $= \tanh(1.24) \approx 0.8455$	True	Theorem 11.5
Fingerprint returns 7 keys, no overlap	True	-

References: SRE Fine-Structure-Constant Derivation, SRE Dynamics: The Book of the Void, SRE Electron Topology (Zenodo)

Mathematical Specifications and Algebraic Derivations for the Topological Closed-Loop Audit Operator (Operator 12)

Version: 1.1 (September 2026 · Open-Source Specification, Correction D3)

Abstract

This paper presents the formal algebraic derivation and specification of the **Topological Closed-Loop Audit Operator** ($\mathcal{O}_{\text{loop-audit}}$, designated as Operator 12) within the Status-Relational Entropy (SRE) dynamics framework. This operator detects the "N-scale topological closed loop + Möbius bilayer phase" pattern described by SRE theory through simple cycle enumeration combined with SRE sign-matrix phase audit, mapping chemical ring structures (benzene, epoxide, etc.) in molecular graphs to the closed retrospective causality of SRE's electron topology.

Version 1.1 corrects the core defect of the original (D3): the original used $\text{trace}(A^k)$ closed-walk counts with a decay-baseline ratio for loop detection, producing false positives on fully-connected weighted graphs (ethanol, an open chain, was detected as having a 6-membered ring; benzene's Fiedler value was NaN). v1.1 replaces this with DFS simple cycle enumeration (no backtracking contamination) plus a sign-matrix Möbius ratio criterion, completely eliminating spectral false positives.

---

1. Motivation and Prerequisites

1.1 SRE Electron Topology Description

SRE theory defines the electron as an "N-scale topological closed loop + Möbius bilayer phase":

- **Topological closure**: $A_N \to A_1$ retrospective causality, forming a self-consistent closed loop
- **Möbius phase**: Requires $2N$ reciprocal measurements to restore the initial symmetric state → spin 1/2
- **Charge**: The closed loop imposes a fixed topological bias on the causal chain per round of reciprocal measurement
- **Mass**: Closed-loop topological processing overhead $m \propto N \cdot l_{\min}$
- $N \approx 10^{23}$ (Compton wavelength / $l_{\min}$)

1.2 Molecular Graph Correspondence

In molecular graphs, this corresponds to:

- Chemical ring structures (benzene 6-membered ring, epoxide 3-membered ring, etc.) → topological closed loops
- Möbius phase → $2N$-step retrospective restoration to symmetry → spin 1/2 characteristic

1.3 Why Not Classical Graph Theory

1. `networkx simple_cycles` on fully-connected weighted graphs (A_{smooth}) produces $O(N!)$ spurious triangles because distant atom pairs also have non-zero edge weights.
2. SRE's graph is a continuous weighted graph, not a discrete binary graph. Classical cycle enumeration requires prior thresholding to binarize, introducing artificial truncation.
3. SRE's electron description is a "topological closed-loop pattern," not a specific chemical ring.

---

2. Mathematical Derivation

2.1 Step 1 — Simple Cycle Enumeration (Definition 5.2, Theorem 12.2)

v1.1 Correction (D3): The original used $\text{trace}(A^k)$ closed-walk counts for loop detection, producing false positives on fully-connected graphs. v1.1 uses DFS simple cycle enumeration:

- **No backtracking**: No vertex repetition within the path

- **No duplicate counting**: Each cycle is enumerated once starting from its minimum vertex; forward and reverse directions are deduplicated by canonical representation
- **Complexity**: For sparse molecular graphs ($|E| \sim O(n)$), the complexity is $O(n \cdot 2^{N_{\max}})$, negligible for $N_{\max} \leq 8$

$n_{\text{loops}} = \text{len}(\text{enumerate_simple_cycles}(A_q, N_{\max}))$

Theorem 12.2 (Backtracking Contamination Elimination): Simple cycle enumeration does not depend on matrix power spectra, completely eliminating spurious triangles caused by fully-connected distance kernels.

Theorem 12.4 (Betti Accuracy): $n_{\text{loops}} = \beta_1$ (first Betti number), i.e., the number of independent cycles in the molecular graph.

2.2 Step 2 — Möbius Phase Ratio (Definition 5.3, Theorem 12.3)

SRE theory requires "$2N$ reciprocal measurements to restore the initial symmetric state" (Möbius phase). On the SRE sign matrix M, define:

$\text{mobius_ratio} = \frac{1}{N_{\max}-2} \sum_{k=3}^{N_{\max}} \frac{|\text{tr}(M^k)|}{n^k}$

where M is the SRE sign matrix (elements $\in \{+1, -1\}$, no zero elements) and n is the number of atoms.

Theorem 12.3 (Sign Balance Criterion):

- $\text{mobius_ratio} = 1 \iff$ no Möbius phase (complete sign balance)
- $\text{mobius_ratio} < 1 \iff$ Möbius phase present (electron spin characteristic exists)

v1.1 Correction (D3): The original defined $R(k) = \text{trace}(A_p^{2k})/(2 \cdot \text{trace}(A_p^k))$ on the normalized adjacency matrix A_p, which produced false positives because $\text{trace}(A^k)$ includes a large number of non-ring closed walks on fully-connected graphs. v1.1 redefines this on the sign matrix M, since M's elements are ± 1 and M^k's trace strictly reflects sign balance.

2.3 Step 3 — Loop Subgraph Spectral Quantities (Definition 5.4, Theorem 12.5)

If $n_{\text{loops}} \geq 1$, extract the loop atom union $S = \bigcup \text{cycles}$ and construct the loop-atom subgraph Laplacian:

$L_{\text{loop}} = L_q[S, S] = \text{diag}\left(\sum_{j \in S} A_q[S_{i,j}]\right) - A_q[S, S]$

$\lambda_{2,\text{loop}} = \lambda_2(L_{\text{loop}}) \quad (\text{loop structural connectivity})$
$\kappa_{\text{loop}} = \frac{\lambda_{\max}(L_{\text{loop}})}{\lambda_2(L_{\text{loop}})} \quad (\text{loop structural condition number})$

Theorem 12.5 (Interlacing Inequality): $\lambda_{2,\text{loop}} \geq \lambda_{2,q} > 0$

The algebraic connectivity of the loop-atom subgraph is no less than the full graph's Fiedler value, guaranteeing the topological connectivity of the ring structure.

---

3. Output Scalars

Scalar	Physical Meaning
n_{loops}	Simple cycle count. Ethanol = 0, benzene = 1. Exactly equals the first Betti number β_1
mobius_ratio	Möbius phase ratio. $= 1$: no Möbius; < 1: electron spin characteristic present
$\lambda_{2,\text{loop}}$	Loop-atom subgraph Fiedler value. NaN = no loops detected
κ_{loop}	Loop-atom subgraph condition number. $\leq \kappa_q$

---

4. Correspondence with SRE Electron Topology Theory

SRE Theory	Operator 12 Implementation
N-scale topological closed loop	n_{loops} (simple cycle count)
$[A_1, ..., A_N]$ retrospective causality	DFS cycle enumeration (no backtracking contamination)

SRE Theory	Operator 12 Implementation
Möbius bilayer phase	$\text{mobius_ratio} = \|\text{tr}(M^k)\|/n^k$
$2N$-step symmetry restoration	$\text{mobius_ratio} < 1 \rightarrow$ double period required
Spin 1/2 emergence	Statistical emergence of $\text{mobius_ratio} \neq 1$
Charge = topological bias	Edge weight bias from electronegativity modulation in A_q
Mass $= N \cdot l_{\min}$ processing overhead	Closed-loop count $\propto$ topological complexity

---

5. Position in the SRE 10-Operator Framework

```
Extended chain: Op1 → Op2 → Op3 → Op6 → Op11 → Op4 → Op12 → Op5 → Op9 → Op10
                                                        ↑
                                        Op 12 audits A_q's closed-loop pattern
                                        Result modulates Op 5's c_e
```

- Op 12 is positioned after Op 4 and before Op 5: it audits the closed-loop pattern of A_q (Op 11 output), and the result modulates Op 5's c_e (information propagation rate).

---

6. v1.1 Validation Protocol

Validation Item	Expected Result	Theorem
Ethanol $n_{\text{loops}} = 0$	0	Theorem 12.2/12.4
Benzene $n_{\text{loops}} = 1 = \beta_1$	1	Theorem 12.4
Benzene $\lambda_{2,\text{loop}} \geq \lambda_{2,q} > 0$	True	Theorem 12.5

References: SRE Fine-Structure-Constant Derivation, SRE Dynamics: The Book of the Void, SRE Electron Topology (Zenodo)

Strict Mathematical Specification of the Spectral-Shape Equation-Solving Operator Set (Operators 13–18): Homological Solvability Theorems and the Endogenous Inversion Pipeline

Version: 1.3 (September 2026 · SRE Version 18.0 Extension Layer)
Reference implementation: `Operator_13_18_P.py` (built-in checks V1–V18, 19/19 passing)
Stress-test suite: `bench_random_equations.py` (10 problem types × 5 random instances = **50 equations** + 6 verification items independent of the pipeline, **50/50 passing**; problem-set document `bench_out/problem_set.md`)
Prerequisites: Operators 1–12 (`structure.md` v18.0, `Operator_10_C.md`, `Operator_11_C.md`, `Operator_12_C.md`)

> **The three extensions of v1.1 over v1.0** (corresponding to the engineering constraints specified by the user):
>
> 1. **Coupled multiple spectral shapes** (§5, §6.4): Operator 14 is generalized to a multi-shard block Jacobian, with cross-shard coupling terms and a shard-to-global embedding map;
> 2. **Non-polynomial kernels** (§4.4): Operator 13 gains an endogenous adaptive truncation order K_f based on Chebyshev coefficient decay, including Bernstein-ellipse convergence-rate prediction and singularity avoidance;
> 3. **Homological solutions** (§3, §9): the definition of a solution is relaxed from "exact solution $r = 0$" to "homological solution $h = 0$", and a new Operator 18 (homological solution realizer) together with canonical-representative selection are added.

> **The three corrections of v1.2 over v1.1** (from what the "10 random equations" stress test exposed empirically):
>
> 1. **The K_f criterion of Operator 13.4 changed from "single term" to "tail sum"** (§13.4) — the original criterion truncates too early when the coefficients decay geometrically; the $\ln$ kernel yields $K_f = 1$ while the tail still exceeds γ;
> 2. **Zero-mode stripping is upgraded from "exclusive to singular kernels" to "mandatory whenever kernel expansion is used"** (§13.5) — for the Stieltjes kernel the analytic ellipse semi-axis is $a_{\max} = 1.142 < |\xi_0| = 1.536$, so the series **diverges** at the zero mode, and the higher the order the more the error explodes ($8.1\times10^{-1} \to 5.0\times10^{5}$ as $K{:}2 \to 32$);
> 3. **For multiple shards one must freeze α_n and λ_2 simultaneously** (§13.6), and it was clarified that "locked-shell rescaling" may be applied only to the initial value and **not to an already-obtained solution** — a frozen scale explicitly destroys the rescaling gauge redundancy.

> **The three corrections of v1.3 over v1.2** (from what the "10 problem types × 5 groups = 50 random equations" stress test exposed empirically):
>
> 1. `OBSTRUCTION` **is a structural false-positive branch** (§17.1) — because the identity $r = F\nabla F$ makes $\rho_h = \|h\|/\|r\| \approx 0.95$ a constant, the criterion band for "exact part converged, harmonic part not converged" is as wide as $1/\gamma \approx 4.3$ times, so any F falling inside it is **misjudged as a topological obstruction** and handed to Operator 9, which changes β_1. It is now required that "the update direction is exhausted (`step_dead`)" before an obstruction may be declared;
> 2. **The η_N normalization deleted the trust region together with $1/(1+q)$** (§15.2, §15.4) — a Newton step for strongly nonlinear instances overshoots ($\|\Delta M\| \approx 0.23$), Ψ rises instead, and the monotonicity red line fails. An **endogenous backtracking line search** (Phase 3.5b) is now added as a safeguard;
> 3. `SOLVED` **must not look only at "first crossing of ε"** (§17.2) — otherwise the V-B independent-residual verification degenerates into a **tautology** with a threshold equal to the termination threshold. The added requirement is the "exhausted" condition (or `step_dead`).

Abstract

This paper addresses the inverse problem **"several unknown spectral shapes form an equation; find its solution"**, and provides, within the SRE binary self-organizing-network dimension-free graph-operator system, a complete solving methodology together with the supporting operators (13–18).

The core conclusion is a **structural discovery**: the k-th spectral moment of a graph is algebraically exactly the normalized weight sum of k-step closed walks on the graph; and a closed walk is a **cycle** ($\ker \partial_1$) in the 1-chain space, whose span after modding out face boundaries $\operatorname{im} \partial_2$ is precisely the **first homology group** H_1. Hence:

> **The residual of a spectral-shape equation is intrinsically a first cohomology class $[r] \in H^1$; solving the equation is equivalent to annihilating that cohomology class on H^1.**

Consequently the solving process needs no new dynamics invented — it is rigorously reduced to a **discrete Hodge three-component routing** on the existing boundary operators $D_{\text{global}} = \partial_1$, $C_{\text{global}} = \partial_2$:

Residual component	Algebraic home	Elimination mechanism	Reused operator
Gradient component $D_{\text{global}}\phi$	$\operatorname{im}(\partial_1)$	Vertex-potential Φ re-solve	Operator 6 ($\Phi = (D^T D)^\dagger D^T E$)
Curl component $C_{\text{global}}^T \psi$	$\operatorname{im}(\partial_2^T)$	Circulation B-field leapfrog cancellation	Operator 7 ($\mathcal{O}_{\text{lock}}$)
Harmonic component h	$\mathcal{H}_1 = \ker(\partial_1^T) \cap \ker(\partial_2)$	**Cannot be eliminated by field iteration**	Operator 9 (Betti stitching, changes topology)

From this we obtain the **topological criterion for solvability** (Theorem S3): the equation is solvable on the given graph topology if and only if the harmonic component $h \equiv 0$; when $h \neq 0$ the residual is "closed but not exact", in which case no iteration can converge, and Operator 9's Betti stitching must change β_1 before solvability is even possible. This criterion provides the solver with an **existence certificate free of external parameters**.

The key semantic upgrade of v1.1 — a solution is a homology class: the user specified that "the restored graph structure may be a homological mapping rather than an exact solution". This fits the framework naturally: the exact part of the residual ($D\phi + C^T\psi$) can be absorbed by the absorption mechanisms of Operators 6/7 and **does not constitute a genuine obstruction**; the only genuine obstruction is the harmonic part. Therefore the definition of a solution can be relaxed to

$$\textbf{Exact solution: } r = 0 \qquad \subsetneq \qquad \textbf{Homological solution: } h = 0 \text{ (i.e. } [r] = 0 \in H^1)$$

that is, **only the residual being exact is required, not the residual being zero** (Definition 3.1). Homological solutions form an affine equivalence class $M^\star + \operatorname{im}(\partial_1) \oplus \operatorname{im}(\partial_2^T)$, whose canonical representative is the one minimizing the Dirichlet energy within the homology class (Operator 18); this yields an explicit semantics for solution uniqueness: **uniqueness holds only in the sense of homology classes** (Theorem S5).

> ⚠️ **Correction to the intuition "relaxing the criterion ⇒ faster convergence" (v1.1 empirical)**
> Relaxing the termination criterion from $\|r\| \to 0$ to $\|h\| \to 0$ **does not necessarily reduce the iteration count**. Since $\rho_h \equiv \|h\|/\|r\| = \|\Pi_{\mathcal{H}_1} \nabla F\|/\|\nabla F\|$ is independent of F (empirically approximately the constant 0.947 on a $\beta_1 = 1$ complex), the harmonic channel cannot be compressed independently and can only vanish together with $\|r\|$; when $\rho_h > \gamma$ the two-level criteria coincide at the convergence point (empirically V15 shows no bifurcation point).
> **The true benefit of homological solutions is semantic, not accelerative**: it relaxes the acceptable solution set from $\{r = 0\}$ to $\{h = 0\}$, so that when the equation admits **no exact solution at all** because it is topologically obstructed, one can still deliver a graph structure correct in the homological sense. See §17.1 for details.

---

1. Canonical Formalization of the Problem

1.1 What is a "spectral shape"

In the SRE framework there are no external coordinates and no continuous metric, so a "spectral shape" must be defined as a **purely algebraic quantity**:

Definition 1.1 (spectral shape): Let the reduced Laplacian of a locally autonomous shard Ω_ν be $L_{\Omega_\nu} \in \mathbb{R}^{N_K \times N_K}$ (positive semi-definite, with $\ker$ dimension $=$ the local component of β_0). Its **spectral shape** σ_ν is defined as the normalized spectral measure of that operator:

$$\sigma_\nu(\lambda) \equiv \frac{1}{N_{K_\nu}} \sum_{i=1}^{N_{K_\nu}} \delta\left(\lambda - \lambda_i\left(L_{\Omega_\nu}\right)\right)$$

After compression by the endogenous normalization α_n (spectral radius) of Operator 6, the spectral support is clamped into the **dimensionless interval** $[0, 1]$:

$\hat{L}_{\Omega_\nu} \equiv \frac{L_{\Omega_\nu}}{\alpha_n}, \qquad \mathrm{supp}(\sigma_\nu) \subseteq [0,1]$

Note: Operator 10 has already proved that $\lambda_2(n) > 0$ always holds, so the effective lower bound of the spectral support is the endogenous spectral gap ratio $\gamma \equiv \lambda_2/\alpha_n \in (0,1]$. **This dimensionless ratio is the only natural scale unit of this solving system**; every threshold, step size, and convergence criterion throughout is generated endogenously from γ, with no external constant whatsoever (in accordance with Postulate 3).

1.2 What is a "spectral-shape equation"

Definition 1.2 (spectral-shape equation): Let there be p unknown spectral shapes $\sigma_1, \dots, \sigma_p$; the equation $\mathcal{F}$ is an algebraic–analytic functional on spectral moments:

$$\boxed{\mathcal{F}(\sigma_1, \dots, \sigma_p) \equiv \sum_{j=1}^{J} a_j \prod_{\nu=1}^{p} \left(\int_0^1 f_{j\nu}(\lambda)\, d\sigma_\nu(\lambda) \right)^{e_{j\nu}} = 0}$$

where $a_j \in \mathbb{R}$, $e_{j\nu} \in \mathbb{N}$, and $f_{j\nu}$ are analytic kernels. The solving goal: find the spectral-shape tuple $\{\sigma_\nu^\star\}$ such that $\mathcal{F} = 0$, and **realize it as a concrete spin-adjacency structure** $M_\Omega^\star$ under the SRE semantics (since in this system only the graph is real, and a spectral shape is merely its projection).

1.3 The key dimension reduction: spectral shape → moment vector

Expanding the kernel functions $f_{j\nu}(\lambda) = \sum_k c_{j\nu k} T_k(\lambda)$ in the Chebyshev basis $\{T_k\}_{k=0}^K$ gives

$\int f_{j\nu}\, d\sigma_\nu = \sum_{k=0}^K c_{j\nu k}\, \tau_k^{(\nu)}, \qquad \tau_k^{(\nu)} \equiv \int_0^1 T_k(\lambda)\, d\sigma_\nu(\lambda)$

and the equation becomes a **multivariate equation in moment space** $\mathcal{F}(\boldsymbol{\tau}^{(1)}, \dots, \boldsymbol{\tau}^{(p)}) = 0$. The unknown is reduced from the infinite-dimensional function σ to the finite-dimensional vector $\boldsymbol{\tau} \in \mathbb{R}^{p(K+1)}$, and τ_k admits a purely combinatorial, purely local computational route (Theorem 13.1), with no eigendecomposition throughout.

1.4 Three-level complexity classification of equation kernels (v1.1)

According to how $\mathcal{F}$ acts on the spectral shapes, the equations are divided into three levels, each requiring additional machinery:

Level	Form	Handling mechanism	Specification location
L1 single spectral shape · polynomial kernel	$\mathcal{F} = \sum_j a_j \prod_k (\tau_k)^{e_{jk}}$	The Operators 13–17 baseline suffices	§3–§8
L2 multiple spectral shapes · coupled kernel	$\mathcal{F}$ contains cross-shard products $\tau_k^{(\mu)} \tau_l^{(\nu)}$	Block Jacobian + shard embedding map + splicing consistency	**§6.4**
L3 non-polynomial kernel	contains $\ln$, rational (Stieltjes-type), exponential, etc.	Chebyshev coefficient expansion + endogenous adaptive truncation K_f	**§4.4**

Their combination (L2 ∩ L3, i.e. coupled multiple spectral shapes with a non-polynomial kernel) is the complete domain of applicability of specification v1.1. The key limitation is the singularity-avoidance criterion of §4.4: the convergence rate of a non-polynomial kernel is **determined solely by the endogenous spectral gap ratio** $\gamma = \lambda_2/\alpha_n$ — as $\gamma \to 0$ (the graph approaching disconnection) the required expansion order of a $\ln$-type kernel diverges, and Operator 16's feasibility warning must catch it.

---

2. Core Mathematical Foundation: Three Foundational Theorems

Theorem S1 (Endogenous equivalence theorem of spectral moments and closed walks)

The k-th Chebyshev spectral moment equals the normalized weight sum of k-step closed walks on the graph, and can be obtained deterministically within $\mathcal{O}(N_K M_K K)$ by a three-term recurrence, **without any eigenvalue decomposition**.

Proof: Let $Y_\nu \equiv 2\hat{L}_{\Omega_\nu} - I$, whose spectral support is $\subseteq [-1,1]$. By the spectral theorem,

$\tau_k^{(\nu)} = \frac{1}{N_K} \mathrm{tr}\left(T_k(Y_\nu)\right) = \frac{1}{N_K} \sum_{v \in \Omega_\nu} \langle e_v,\ T_k(Y_\nu)\, e_v \rangle$

Expand $T_k(Y) = \sum_{r=0}^{k} \beta_{kr} Y^r$; then $(Y^r)_{vv} = \sum_{v_1,\ldots,v_{r-1}} Y_{vv_1} Y_{v_1v_2} \cdots Y_{v_{r-1}v}$ is exactly the **weight sum of all closed walks of length r starting from v**. Hence τ_k is a linear combination of closed walks of length $\leq k$. Moreover $T_k(Y)e_v$ is generated order by order through the Chebyshev three-term recurrence

$$u_0 = e_v, \quad u_1 = Ye_v, \quad u_{k+1} = 2Yu_k - u_{k-1}$$

and each order needs only one **local sparse matrix–vector product** (the number of nonzeros of Y is $= \mathcal{O}(M_K)$). No matrix factorization is involved throughout, satisfying Postulate 1 (spectral-flow shard splicing and reuse; a globally synchronized decomposition is forbidden). ■

Theorem S2 (First-homology attribution theorem of spectral moments)

The linear space spanned by the spectral moments, after modding out face boundaries, is **isomorphic to the dual of the first homology group** H_1; its dimension is the first Betti number β_1.

Proof: By Theorem S1, the spectral moments are generated by closed walks. A closed walk $w = \sum_e n_e e$ satisfies in–out conservation at every vertex, i.e. $\partial_1 w = 0$, so $w \in \ker \partial_1$ (a **cycle**). By the exactness constraint of the chain complex $\partial_1 \partial_2 = 0$, one has $\operatorname{im} \partial_2 \subseteq \ker \partial_1$. The space of closed walks, after modding out the boundaries generated by 2-chains (faces), is precisely

$$H_1 \equiv \ker \partial_1 \big/ \operatorname{im} \partial_2, \qquad \dim H_1 = \beta_1$$

and Operator 12 has already proved $n_{\text{loops}} = \beta_1$ (Theorem 12.4). Hence the spectral-moment functional naturally takes values in $H_1^* \cong H^1$. ■

Corollary S2.1: The sensitivity vector of the spectral moments with respect to the edge weights (defined in Operator 14) must lie in the 1-chain space $C_1 \cong \mathbb{R}^{m_g}$, so the **residual can be injected losslessly into Operator 7's E/B-field leapfrog**, with no additional spatial embedding.

Terminology convention (important: strict distinction between the two "cycle counts")

This specification strictly distinguishes the following two quantities that are often conflated; they coincide only in the special case (∂_2 spanning the full cycle space):

Symbol	Name	Definition	Semantics
ν	cycle-space dimension / circuit rank	$\nu \equiv \dim \ker \partial_1 = m_g - n + \beta_0$	the number of independent **closed-walk** basis elements. This is the n_{loops} of Operator 12
β_1	first Betti number / homology dimension	$\beta_1 \equiv \dim \mathcal{H}_1 = \nu - \operatorname{rank}(\partial_2)$	the number of topologically "true holes" remaining after subtracting **face boundaries**

Key: ν measures "how many independent closed loops can be drawn", while β_1 measures "how many loops **cannot** be filled by 2-chains (faces)". $\beta_1 = 0$ means all cycles can be filled by faces (e.g. spherical topology), in which case **there is no topological obstruction** (the h-space of Theorem S3 is zero and the equation is always solvable).

When, for a planar graph, one takes all bounded faces of its planar embedding as 2-chain generators, then $\operatorname{rank}(\partial_2) = \nu$, so $\beta_1 = 0$; if the face set is incomplete (e.g. taking only triangles and omitting the large cycles), then $\beta_1 > 0$, and the **residual β_1 is precisely the degrees of freedom by which the equation may be unsolvable**. This distinction is the numerical basis by which Operator 17 decides "topological obstruction".

2-bis. Definition and Structure of Homological Solutions (v1.1 core)

Definition 3.1 (exact solution vs. homological solution)

Let the residual 1-chain of equation $\mathcal{F}$ on graph M be $r(M)$, with Hodge decomposition $r = D\phi + h + C^T\psi$. Two levels of solution are defined:

$$\textbf{Exact solution} \quad \mathcal{S}_{\text{exact}} \equiv \{M : r(M) = 0\} \qquad (\text{all three components zero, } m_g \text{ constraints in total})$$

$$\boxed{\textbf{Homological solution} \quad \mathcal{S}_{\text{hom}} \equiv \{M : h(M) = 0\} \;=\; \{M : [r(M)] = 0 \in H^1\}} \qquad (\text{only } \beta_1 \text{ constraints})$$

Physical semantics: the exact part of the residual $D\phi + C^T\psi$ can be absorbed by the absorption mechanisms of Operator 6 (potential re-solve) and Operator 7 (circulation cancellation), and **does not constitute a genuine obstruction**; the only genuine obstruction is the harmonic part. Hence "residual exact" already suffices to call it a solution — this is exactly the rigorous formalization of the user's stipulation that "the graph structure may be a homological mapping rather than an exact solution".

Obviously $\mathcal{S}_{\text{exact}} \subseteq \mathcal{S}_{\text{hom}}$, and the inclusion is proper when $\beta_1 > 0$.

Definition 3.2 (homological equivalence and homological distance)

Two graphs M, M' are called **homologically equivalent** ($M \sim M'$) if the difference of their residuals is an exact chain:
$M \sim M' \iff r(M) - r(M') \in \text{im}(\partial_1) \oplus \text{im}(\partial_2^T) \iff h(M) = h(M')$

On the equivalence classes we define the **homological distance** (a pseudo-metric, identically zero within a class):

$$\boxed{d_{\text{hom}}(M, M') \equiv \left\| h(M) - h(M') \right\|_2}$$

d_{hom} measures the **true difference in the topological sense** between two graph realizations — it automatically ignores every exact difference absorbable by the field, and compares only the "holes that cannot be eliminated". This is the correct metric for judging whether two candidate realizations are "essentially the same".

Theorem S4 (dimension of the homological-solution set: $m_g - \beta_1$)

Linearizing at a solution $M^\star \in \mathcal{S}_{\text{hom}}$, the tangent space of the homological-solution set contains $\mathcal{H}_1^\perp = \text{im}(\partial_1) \oplus \text{im}(\partial_2^T)$, hence

$$\dim \mathcal{S}_{\text{hom}} \;=\; m_g - \beta_1, \qquad \dim \mathcal{S}_{\text{exact}} \;=\; 0 \;(\text{isolated point})$$

Proof: The constraint $h(M) = 0$ consists of β_1 independent scalar equations (since $h \in \mathcal{H}_1$, $\dim = \beta_1$), acting on the m_g-dimensional edge-weight space, so the solution set has dimension $m_g - \beta_1$. The tangent directions are guaranteed by Theorem 15.3: $\Delta M \perp \mathcal{H}_1$, i.e. moving along the Newton direction does not change h to first order. ■

Corollary S4.1 (solving difficulty is monotonically determined by β_1): the larger β_1 is → the more harmonic constraints → the smaller the homological-solution set → the harder to solve; when $\beta_1 = 0$ the number of constraints is zero.

Theorem S5 (uniqueness semantics of homological solutions)

The solution of the equation is **unique in the sense of homology classes**: any two homological solutions $M_1^\star, M_2^\star \in \mathcal{S}_{\text{hom}}$ must be homologically equivalent ($d_{\text{hom}} = 0$), but in general $M_1^\star \neq M_2^\star$ (they may differ by an exact edge-weight perturbation).

⚠️ Degeneracy warning: homological solutions lose discriminating power when $\beta_1 = 0$

When $\beta_1 = 0$ (e.g. a closed complex), $\mathcal{H}_1 = \{0\}$, so $h \equiv 0$ **holds for any graph** — the homological-solution set degenerates to the whole edge-weight space $\mathcal{S}_{\text{hom}} = \mathbb{R}^{m_g}$, and the criterion $|h| = 0$ carries no information.

In this case one must invoke the **canonical-representative** criterion to select a unique realization (Operator 18): take the one minimizing the Dirichlet energy within the homology class. This also explains why canonical-representative selection in Operator 18 is not an option but a necessary component of v1.1.

Engineering guideline: before solving, compute β_1. If $\beta_1 = 0$, solve directly with the exact-solution criterion $\|r\| \le \epsilon_{\text{endo}}$ and skip the homological-solution flow; if $\beta_1 > 0$, one may then enable the homological-solution criterion $\|h\| \le \epsilon_{\text{endo}}$ to save iterations.

Theorem S3 (Hodge three-component decomposition of the residual and the solvability criterion) ★core

Given any 1-chain residual $r \in C_1(\mathbb{R}^{m_g})$, there exists a **unique orthogonal decomposition**

$$\boxed{r = D_{\text{global}}\,\phi \;+\; h \;+\; C_{\text{global}}^T\,\psi}$$

where ($D = \partial_1 : C_0 \to C_1$ is $m_g \times n$, $C = \partial_2 : C_1 \to C_2$ is $f \times m_g$)

- $D\phi \in \text{im}(\partial_1) \subseteq C_1$: the **gradient component**, with $\phi \in \mathbb{R}^n$ the vertex potential difference;
- $C^T\psi \in \text{im}(\partial_2^T) \subseteq C_1$: the **curl component**, with $\psi \in \mathbb{R}^f$ the circulation density;
- $h \in \mathcal{H}_1 \equiv \ker(\partial_1^T) \cap \ker(\partial_2) \subseteq C_1$: the **harmonic component**, satisfying $D^T h = 0$ and $Ch = 0$.

and the projection operators are completely endogenous (using only the boundary operators, with no external metric):

$$\phi \equiv \underbrace{(D^T D)^\dagger}_{L_0^\dagger \text{ (0-order Laplacian pseudo-inverse)}} D^T r \in \mathbb{R}^n, \qquad \psi \equiv (CC^T)^\dagger C r \in \mathbb{R}^f, \qquad h \equiv r - D\phi - C^T\psi$$

Note: the solving formula for ϕ is **strictly isomorphic** to Operator 6's potential-closure formula $\Phi_{s-1} = (D^T D)^\dagger (D^T E_{s-1})$ (merely replacing the edge flow E by the residual r), so all of Operator 6's $\mathcal{O}(1)$ local facilities can be reused directly, with zero new solver.

Solvability criterion: the equation is solvable on the given graph topology $\iff h \equiv 0$.

Proof:
(i) Existence and uniqueness: the graph Hodge decomposition theorem guarantees that $C_1 = \mathrm{im}(\partial_1) \oplus \mathcal{H}_1 \oplus \mathrm{im}(\partial_2^T)$ is an orthogonal direct-sum decomposition (where $\mathcal{H}_1 \cong H_1$ is the harmonic representative of the homology group), so the decomposition is unique, and

$$\|r\|_2^2 = \|D\phi\|_2^2 + \|h\|_2^2 + \|C^T\psi\|_2^2$$

(ii) Projection formulas: $D\phi$ is the orthogonal projection of r onto $\mathrm{im}\,\partial_1$, whose normal equation is $D^T(D\phi) = D^T r \Rightarrow (D^T D)\phi = D^T r$, hence $\phi = (D^T D)^\dagger D^T r$; likewise $C^T\psi$ is the projection of r onto $\mathrm{im}\,\partial_2^T$, with $\psi = (CC^T)^\dagger C r$. The former is the 0-order Laplacian pseudo-inverse (of the same origin as Operator 6's potential solve), the latter the 2-order Laplacian pseudo-inverse; an $\mathcal{O}(1)$ local solving route has been established for both.

(iii) Solvability: the update direction of the solving iteration (Operator 15) must belong to $\mathrm{im}(\partial_1) \oplus \mathrm{im}(\partial_2^T) = \mathcal{H}_1^\perp$ (see Theorem 15.3). Hence if $h \neq 0$, the harmonic component of the residual falls into the kernel of the update operator, and **no number of field iterations can make it decrease**; then r is a "closed but not exact" cohomology class, constituting a **topological obstruction**. Only changing the graph's topology ($\Delta\beta_1 \neq 0$, Operator 9's Betti stitching) can change the dimension of $\mathcal{H}_1$ and thereby open a new exact direction for the residual. ■

Corollary S3.1 (obstruction certificate): when the iteration stalls with $\|h\|$ not decreasing, the harmonic component h itself is the **algebraic certificate** that the equation is unsolvable — it precisely indicates "how many independent loops the equation demands that the current manifold cannot provide".

---

3. Operator Specifications

Operator 13: $\mathcal{E}_{\mathrm{mom}}$, the Spectral-Shape Chebyshev Moment Encoder

(Spectral-Shape Chebyshev Moment Encoder)

Item	Specification
Topological space	$\mathcal{H}_{\mathrm{sym}}(N_K) \to \mathbb{R}^{K+1}$
Input	Local shard reduced Laplacian L_{Ω_ν}; global spectral assets (α_n, λ_2) (output of Operator 6)
Output	Normalized Chebyshev moment vector $\boldsymbol{\tau}^{(\nu)} = (\tau_0, \dots, \tau_K)^T$; sensitivity vector set $\{\mathbf{w}_k^{(\nu)}\}$
Timing position	Tail of Phase 1 (immediately after Operator 6, reusing its Lanczos subspace, with zero extra spectral decomposition)

13.1 Endogenous normalization and moment recurrence

$$\hat{L}_{\Omega_\nu} \equiv \frac{L_{\Omega_\nu}}{\alpha_n}, \qquad Y_\nu \equiv 2\hat{L}_{\Omega_\nu} - I \quad (\mathrm{spec}(Y_\nu) \subseteq [-1, 1])$$

$$\tau_k^{(\nu)} \equiv \frac{1}{N_{K_\nu}} \mathrm{tr}\,(T_k(Y_\nu)) = \frac{1}{N_{K_\nu}} \sum_{v \in \Omega_\nu} \left\langle e_v,\ u_k^{(v)} \right\rangle, \quad u_{k+1}^{(v)} = 2Y_\nu u_k^{(v)} - u_{k-1}^{(v)}$$

13.2 Endogenous truncation order K (no external parameters)

$$K_\nu \equiv \min\left(\left\lceil \frac{\alpha_n}{\lambda_2} \right\rceil,\ K_{\mathrm{cap}} \right), \qquad K_{\mathrm{cap}} \equiv \lfloor 2\ln(1 + N_{K_\nu}) \rfloor$$

Basis: the Chebyshev truncation error for a Lipschitz spectral density is $\mathcal{O}(K^{-1})$; requiring the truncation error not to exceed the endogenous minimum resolvable scale λ_2/α_n demands $K \gtrsim \alpha_n/\lambda_2$. K_{cap} prevents the moment order from exceeding the logarithm of the shard's degrees of freedom (overfitting protection); the accuracy loss when it is exceeded is caught by Operator 16.

13.3 Edge-space sensitivity vectors (closed form, reusing boundary-operator column vectors)

From $\partial L/\partial M_e = d_e d_e^T$ (d_e being the transpose of the e-th row of D_{global}) and $T_k' = kU_{k-1}$ (Chebyshev of the second kind), one obtains the **fully analytic sensitivity**:

$$\boxed{w_{k,e}^{(\nu)} \equiv \frac{\partial \tau_k^{(\nu)}}{\partial M_e} = \frac{2k}{N_{K_\nu}\,\alpha_n}\; d_e^T\, U_{k-1}(Y_\nu)\, d_e \qquad (k \geq 1), \qquad w_{0,e}^{(\nu)} \equiv 0}$$

Derivation: from $L = D^T \mathrm{diag}(M) D$ one gets $\partial L/\partial M_e = d_e d_e^T$ ($d_e \equiv$ the transpose of the e-th row of the boundary operator), and $\partial Y/\partial M_e = \frac{2}{\alpha_n} d_e d_e^T$, hence
$\frac{\partial \tau_k}{\partial M_e} = \frac{1}{N_K}\mathrm{tr}\left(T_k'(Y)\cdot \frac{2}{\alpha_n} d_e d_e^T\right) = \frac{2}{N_K \alpha_n} d_e^T T_k'(Y) d_e = \frac{2k}{N_K \alpha_n} d_e^T U_{k-1}(Y) d_e$
(using the identity $T_k' = k\,U_{k-1}$; at $k=0$, $\tau_0 \equiv 1$, so the sensitivity is identically zero.)

U_{k-1} is likewise generated synchronously by the three-term recurrence, with zero extra matrix factorization. **This formula is strictly isomorphic to Operator 10's effective impedance tensor $\mathbf{Z}_{\text{eff}} = (e_u - e_v)^T L^+ (e_u - e_v)$** — both are quadratic forms of boundary-operator column vectors over matrix functions, so all of Operator 10's localization accelerations and Sherman-Morrison incremental-update facilities can be reused.

Theorem 13.1 (two-way distinguishability of moments and spectral shape)

If $K \geq \alpha_n/\lambda_2$, then the moment vector uniquely determines the spectral shape at the same scale, and the reconstruction error satisfies
$\|\rho_\nu - \rho_{\nu,K}\|_\infty \leq C \cdot \frac{\lambda_2}{\alpha_n}$
Corollary: the resolution limit of the moment representation is exactly the endogenous spectral gap ratio; there is no finer distinguishable degree of freedom — this is consistent with the system's stance of "no external metric".

13.4 Handling of non-polynomial kernels (v1.1 addition · L3-level equations)

When the equation contains a non-polynomial kernel f ($\ln$, rational/Stieltjes-type, exponential, etc.), **one does not integrate f numerically directly**; instead one expands it in the Chebyshev basis, thereby linearizing the nonlinear action completely into a moment inner product.

Unified variables: the spectral variable $\hat{\lambda} = \lambda/\alpha_n \in [\gamma, 1]$ relates to the recurrence variable $y \in [-1, 1]$ by $\hat{\lambda} = (y+1)/2$. Letting $g(y) \equiv f\left(\frac{y+1}{2}\right)$, one has

$$\int_0^1 f(\hat{\lambda})\, d\sigma \;=\; \int_{-1}^1 g(y)\, d\sigma_Y \;=\; \sum_{k=0}^{K_f} c_k\, \tau_k \;=\; \mathbf{c}^T \boldsymbol{\tau}$$

Chebyshev coefficients (Chebyshev-Gauss quadrature, deterministic):
$c_k \equiv \frac{2}{N_c} \sum_{j=0}^{N_c-1} g(\cos\theta_j)\cos(k\theta_j), \qquad \theta_j \equiv \frac{(2j+1)\pi}{2N_c}$
exact for polynomials of $\deg < N_c$; for general functions it converges at the Bernstein ellipse rate.

Endogenous adaptive truncation order K_f (replacing the fixed K of the polynomial-kernel case):

> **v1.2 correction (empirical from the 10-equation random stress test, bench E06/E08)**. The original criterion took $K_f = \max\{k : |c_k| > \gamma \max|c|\}$, which measures whether a **single term** has become small. This is logically wrong: the Chebyshev coefficients decay geometrically, and **a single term becoming small is far from meaning the approximation is good enough**. Empirically the $\ln$ kernel gives $K_f = 1$, while $\mathrm{tail}(1)$ still accounts for 21% of the peak $> \gamma$, with a truncation error reaching 3.25×10^{-2}.

What really determines accuracy is the **sum** of the discarded terms. Since $|T_k(s)| \leq 1$ on $[-1, 1]$,
$\left|g(s) - \sum_{k\leq K} c_k T_k(s)\right| \;\leq\; \sum_{k>K} |c_k| \;\equiv\; \mathrm{tail}(K)$
so $\mathrm{tail}(K)$ is the **strict upper bound** on the approximation error, and is exactly the quantity that should be compared with the endogenous tolerance:

$$\boxed{K_f \;\equiv\; \min\left\{K \;:\; \mathrm{tail}(K) \;\leq\; \gamma \cdot \max_k |c_k|\right\}, \qquad \mathrm{tail}(K) = \sum_{k>K} |c_k|}$$

The threshold is still $\gamma \cdot \max|c|$ — the sole endogenous scale, with no external parameters. **This criterion automatically adapts to the smoothness of the kernel**: smooth kernels give a small K_f, kernels with strong singularities give a large K_f.

Note that this guarantees γ-level accuracy, not machine accuracy. To reach the 10^{-6} level, K must be as deep as $\mathcal{O}(1/\ln\rho)$ ≈ 16–24 — this is an explicitly stated cost, not a defect (see the bench report for the empirical K-sweep).

Theorem 13.2 (the convergence rate of non-polynomial kernels is determined solely by the spectral gap ratio)
For a kernel with a logarithmic branch point at $\hat{\lambda} = 0$ (typically $\ln\hat{\lambda}$), the location of its singularity in the y variable and the Bernstein parameter are

$$\xi_0 = -\tfrac{1+\gamma}{1-\gamma}, \qquad \rho \equiv |\xi_0| + \sqrt{\xi_0^2 - 1}$$

The Chebyshev coefficients decay asymptotically as $|c_k| = \mathcal{O}\big(k^{-1}\rho^{-k}\big)$, so the lower bound on the required order is

$$K_f \;\gtrsim\; \tfrac{\ln(1/\gamma)}{\ln\rho}, \qquad \lim_{\gamma\to 0}\rho = 1 \;\Rightarrow\; K_f \to \infty$$

Corollary: **the spectral gap ratio γ is simultaneously a measure of "the graph's connectivity margin" and of "the expandability of a non-polynomial kernel"**. The closer the graph is to disconnection ($\gamma \to 0$), the more the required expansion order of a $\ln$-type kernel diverges — this is the method's fundamental limitation on ill-conditioned graphs, and it cannot be circumvented by adding computing power.

Singularity-avoidance criterion: let the kernel's singularity be $\hat{\lambda}_s$, and define its endogenous distance to the spectral support $d_s \equiv \operatorname{dist}(\hat{\lambda}_s, [\gamma, 1])$.

- If $d_s > \gamma$: the expansion is safe, and K_f is determined by the coefficient criterion;
- If $d_s \le \gamma$: **critical**; a $\ln$-type kernel is always in this case (the distance from the singularity 0 to the support $[\gamma, 1]$ is exactly γ); here one must rely on the adaptive K_f criterion, and Operator 16 must flag the warning "kernel expansion accuracy limited by the spectral gap".
- Handling: prefer raising λ_2 via Operator 9 (topological stitching to increase connectivity) so as to lift γ, rather than increasing K_f and brute-forcing.

Special status of the Stieltjes-type kernel: the response of the kernel $\frac{1}{\hat{\lambda}-z}$ ($z \in \mathbb{C}\setminus\mathbb{R}$) is exactly the spectral Stieltjes transform $m(z) = \sum_k c_k(z)\tau_k$. It is simultaneously the standard gateway to **spectral-density inversion**:

$$\rho(\hat{\lambda}) = \tfrac{1}{\pi}\operatorname{Im} m(\hat{\lambda} + i\varepsilon), \qquad \varepsilon_{\text{endo}} \equiv \gamma$$

(the imaginary offset must be given endogenously by the spectral gap ratio; an external small quantity is forbidden.) This provides a direct route for Operator 18's "moments → spectral density" reconstruction.

13.5 Zero-mode stripping (v1.2 correction · strip whenever kernel expansion is used, not exclusive to singular kernels)

Why it is not "$\ln$-only"

v1.1 wrote stripping as a prerequisite exclusive to singular kernels ($\ln$, $1/\hat{\lambda}$); **this is incomplete**. E08 (Stieltjes kernel) of the 10-equation random stress test exposed the true discriminating condition:

The trivial zero mode in the shifted variable is always located at

$$\xi_0 \;=\; -\tfrac{1+\gamma}{1-\gamma}, \qquad \rho(\xi_0) \;=\; |\xi_0| + \sqrt{\xi_0^2 - 1}$$

It is **always outside** $[-1, 1]$, and $T_k(\xi_0)$ grows exponentially as $\rho(\xi_0)^k$. Meanwhile the Chebyshev series of the kernel g converges only inside its own Bernstein ellipse, whose real semi-axis is

$$a_{\max} \;=\; \tfrac{\rho_{\max}+\rho_{\max}^{-1}}{2}$$

where $\rho_{\max}$ is determined by the singularity of g nearest to the spectral support. **As soon as $a_{\max} < |\xi_0|$, the series diverges at ξ_0**, and the high-order terms, amplified by the zero mode, explode — the higher the order, the worse.

Empirical Stieltjes kernel $z = 0.5 + i\gamma$ ($\gamma = 0.2113$):

Quantity	Value
singularity s_{pole}	$-0.267949 + 0.535899\,i$
$\rho_{\max}$	1.693185
$a_{\max} = (\rho + \rho^{-1})/2$	1.141894
$\lvert\xi_0\rvert$	1.535899
Verdict	$a_{\max} = 1.142 < 1.536 = \lvert\xi_0\rvert \to$ **divergence**

As K increases from 2 to 32, the kernel-fidelity error goes $8.1 \times 10^{-1} \to 5.0 \times 10^{5}$ — **explosion rather than convergence**. After stripping it falls back to 2.02×10^{-2}.

> **Rule (v1.2)**: whenever a Chebyshev kernel expansion is used, always strip the zero mode. Reason: the location of $\lambda = 0$ is universally known, stripping needs no eigendecomposition, and after stripping the reduced spectrum lies strictly in $[\gamma, 1] \to s \in [-1, 1]$, so the series necessarily converges. For entire functions (e.g. $\exp$) stripping is harmless; for branch/pole-type kernels stripping is **mandatory**.

Stripping formula

When there are multiple connected components there is an n_0-fold zero mode (n_0 is obtained purely combinatorially by union-find, using no spectral decomposition), generalized to

$$\boxed{\tau_k^{\mathrm{red}} \;=\; \frac{N\,\tau_k \;-\; n_0\,T_k(\xi_0)}{N - n_0}}$$

($T_k(\xi_0)$ is obtained by the scalar three-term recurrence $T_{k+1} = 2\xi_0 T_k - T_{k-1}$.) The kernel integral then becomes $\frac{1}{N-n_0}\sum_{\lambda_i \neq 0} f(\hat{\lambda}_i) = \sum_k c_k \tau_k^{\mathrm{red}}$.

Corresponding correction of the sensitivity: stripping is a linear operation, so $\partial \tau_k^{\mathrm{red}} / \partial M_e = \frac{N}{N-n_0}\,\partial \tau_k / \partial M_e$; composed with the shift coordinate $\alpha_n^{\mathrm{eff}} = \alpha_n(1-\gamma)$, this gives
$w_{k,e}^{\mathrm{red}} \;=\; \frac{N}{N-n_0} \cdot \frac{2k}{N\,\alpha_n(1-\gamma)}\, d_e^T U_{k-1}(\tilde{Y}) d_e$

> This operation is consistent with the stance of Operator 6 that "the zero mode is a gauge redundancy rather than a physical degree of freedom": the zero mode carries only the counting information of connected components and does not participate in the spectral shape.

13.6 Endogenous scale locking and field-refresh period (v1.1 addition · key consistency constraint)

The spectral moments τ_k are normalized by the endogenous scale $\alpha_n(M)$, while α_n itself is a function of M. The sensitivity of Operator 13.3 W_k **does not differentiate with respect to α_n under Operator 6's convention** (α_n is a hard constant within a field-refresh period). This yields a **consistency constraint that must be obeyed**:

> **α_n and λ_2 must remain locked for several iteration steps. If refreshed at every step, the definition of F jumps between steps and the Newton prediction $\Delta F = -F$ fails.**

Empirical comparison: at the same step size, refreshing step by step makes F **jump up** on the very first step ($7.163 \times 10^{-2} \to 7.192 \times 10^{-2}$), destroying the monotonicity of Ψ; switching to locking gives convergence in 2 steps with strict monotonicity ($7.163 \times 10^{-2} \to 2.715 \times 10^{-3}$).

Endogenous refresh criterion (no external period parameter):
$\mathrm{refresh} \iff \frac{\|L(M) - L_{\mathrm{ref}}\|_F}{\alpha_n} \;>\; \gamma$
i.e. refresh when the relative drift of the Laplacian exceeds the spectral gap ratio — this is of the same origin as the trust region of Theorem 15.1, $\|\Delta L\|_F \leq \lambda_2/2$: once the drift has accumulated past γ, the linearization is no longer trustworthy.

For multiple shards one must simultaneously **freeze** λ_2 (v1.2 supplement, empirical from bench E09/E10). The kernel coordinate of §13.5,
$\tilde{Y} \;=\; \frac{Y - \gamma I}{1-\gamma}, \qquad \gamma = \frac{\lambda_2}{\alpha_n}$
depends on α_n **and** λ_2 at the same time. If one freezes only α_n and lets λ_2 be recomputed at every step, "inside the solver" and "the external evaluation" end up in **two different shift coordinate systems**. The empirical moment cross-deviation reaches the 2×10^{-2} level (E09: 1.955×10^{-2}; E10: 4.567×10^{-2}), which in turn worsens the independent residual to 3.47×10^{-1} / 1.405×10^{0}; freezing λ_2 as well brings it back down to 7.24×10^{-3} / 1.13×10^{-2}.

Locked-shell rescaling (a prerequisite for comparing multiple realizations): when comparing two candidate graphs under the **same** set of locked scales, one must first perform
$M \;\mapsto\; cM, \qquad c \equiv \frac{\alpha_n^{\mathrm{lock}}}{\alpha_n(M)}$
otherwise the spectrum of $Y = 2L/\alpha_n^{\mathrm{lock}} - I$ escapes $[-1, 1]$, the Chebyshev recurrence is amplified, and the two candidates are simply not under the same normalization — the comparison is meaningless.

> **Boundary of applicability (v1.2 clarification, important)**: shell rescaling **may be applied only to "the initial value/candidate graph before injection into the solve"**, and **must never be applied to an already-obtained solution** $M^\star$.

> Reason: $L(M) = D^T \mathrm{diag}(M) D$ is positively homogeneous of degree one in M; if **every point** is normalized by its own α_n, then rescaling leaves $\hat{\lambda}$ unchanged and τ and F strictly unchanged — this is intrinsically a one-dimensional **gauge redundancy**. But this system must freeze α_n, and then $Y = 2L/\alpha_n^{\mathrm{lock}} - I$ turns the **absolute scale** of M into a physical quantity: the rescaling factor c makes $Y \mapsto cY + (c-1)I$, which is **affine rather than the identity**, and $\tau_k \sim Y^k$ changes accordingly.
>
> Empirical (bench E01): after multiplying back the α_n drift of the solution $M^\star$ (4.35×10^{-3}) for gauge fixing, the independent residual worsens from 1.85×10^{-2} to 1.19×10^{0} (two orders of magnitude).
>
> **Conclusion: a frozen scale explicitly destroys the rescaling gauge redundancy; it is the symmetry this method actively gives up in exchange for the smoothness of F.** Therefore the residual α_n drift after convergence can only be accepted in Operator 16 as a second-order effect (see below).

Quantitative acceptance criterion for the drift: write $\varepsilon \equiv \left|\alpha_n(M^\star)/\alpha_n^{\mathrm{lock}} - 1\right|$. When the top of the support crosses from 1 to $1 + \varepsilon$, the Hausdorff infeasibility of the moment sequence appears only at **second order** (to first order it merely shifts the support boundary), so one should have
$\left|\lambda_{\min}(H)\right| \;=\; \mathcal{O}(\varepsilon^2)$
Empirical E01: $\varepsilon^2 = 1.89 \times 10^{-5}$ vs $|\lambda_{\min}| = 1.00 \times 10^{-5}$ (ratio 0.53); E09: $\varepsilon^2 = 5.59 \times 10^{-5}$ vs 8.43×10^{-5} (ratio 1.51).
Operator 16's admission criterion is graded accordingly:

Case	Verdict
$\lambda_{\min}(H) \geq 0$	**pass**
$\lambda_{\min}(H) < 0$ but $\|\lambda_{\min}\| \leq 10\,\varepsilon^2$ and $\varepsilon < \gamma$	**pass (*)** — every violation is explained by the freezing artifact
$\lambda_{\min}(H) < 0$ and $\|\lambda_{\min}\| > 10\,\varepsilon^2$	**fail** — genuinely infeasible moments, must be clamped by Op16

Complexity and red lines

$\mathcal{O}(N_K \cdot M_K \cdot K)$ (polynomial kernel) or $\mathcal{O}(N_c^2 + N_K M_K K_f)$ (non-polynomial kernel, with the coefficient quadrature precomputed once). Since Operator 10 has proved $N_K \leq K_0 = \mathcal{O}(1)$, the whole is at the **local constant order**.
Hard red line: when K_{cap} takes effect (ill-conditioned graph, $\kappa_\Omega \gg K_{\mathrm{cap}}$) the accuracy claim must be downgraded, and Operator 16 must output a feasibility warning; silently downgrading accuracy is forbidden. For non-polynomial kernels one must additionally verify $K_f \leq K_{\mathrm{cap}}$, otherwise declare "kernel not expandable".

Operator 14: $\mathcal{R}_{\mathrm{hodge}}$, the Equation-Residual Homological Router

(Equation Residual Homological Router)

Item	Specification
Topological space	$\mathbb{R}^{p(K+1)} \to C_1(\mathbb{R}^{m_g}) \to \mathrm{im}(\partial_1) \oplus \mathcal{H}_1 \oplus \mathrm{im}(\partial_2^T)$
Input	Moment vector set $\{\boldsymbol{\tau}^{(\nu)}\}$, equation $\mathcal{F}$, boundary operators $D_{\mathrm{global}}, C_{\mathrm{global}}$
Output	Three residual components (ϕ, ψ, h) and their norms; injection instructions
Timing position	Tail of Phase 3 (executed jointly with Operator 7's half-step $B_{s+1/2}$ update)

14.1 Moment-space residual and dimension lifting (μ_{resid} micro-operator)

$g \equiv \mathcal{F}\left(\boldsymbol{\tau}^{(1)}, \ldots, \boldsymbol{\tau}^{(p)}\right) \in \mathbb{R}^q$

Jacobian $J \in \mathbb{R}^{q \times p(K+1)}$, $J_{j,(\nu,k)} \equiv \partial \mathcal{F}_j / \partial \tau_k^{(\nu)}$ (analytically available, since $\mathcal{F}$ is polynomial in the moments).

Residual dimension lifting: push the scalar residual back into the 1-chain space along the sensitivity directions:

$r \equiv \sum_\nu \sum_{k=0}^{K} \left(J^T g\right)_{(\nu,k)} \cdot \mathbf{w}_k^{(\nu)} \;\in\; \mathbb{R}^{m_g}$

i.e. $\nabla_M \mathcal{F} = \sum_{\nu,k} J_{(\nu,k)}\, \mathbf{w}_k^{(\nu)}$, combined with the scalar residual g. **This vector naturally belongs to the edge space**, and by Corollary S2.1 it can enter the Hodge decomposition directly.

14.2 Three-component projection

$\phi \equiv (D^T D)^\dagger D^T r \in \mathbb{R}^n, \qquad \psi \equiv (CC^T)^\dagger C\, r \in \mathbb{R}^f, \qquad h \equiv r - D\phi - C^T\psi$

14.3 Component routing (reusing existing operators, zero new dynamics)

Gradient channel (vertex potential Φ, Operator 6 reuse): the gradient component of the residual requires a vertex-potential correction ϕ, injected directly in the 0-chain space:
$\Phi_{s-1} \;\leftarrow\; \Phi_{s-1} \;-\; \eta_{\text{inj}}^{(\phi)} \cdot \phi, \qquad \eta_{\text{inj}}^{(\phi)} \equiv \frac{\lambda_2}{\lambda_2 + \|D\phi\|_2}$
(equivalently, Operator 6's right-hand side is rewritten as $(D^T D)^\dagger D^T (E_{s-1} - \eta\, r)$; the two are equivalent by the projection idempotency $D(D^T D)^\dagger D^T = P_{\text{im}D}$.)

Curl channel (circulation B, Operator 7 reuse): $\psi \in \mathbb{R}^f$ is itself a face-space element, injected directly:
$B_{s+1/2} \;\leftarrow\; B_{s+1/2} \;-\; \eta_{\text{inj}}^{(\psi)} \cdot \psi, \qquad \eta_{\text{inj}}^{(\psi)} \equiv \frac{\lambda_2}{\lambda_2 + \|C^T\psi\|_2} \in (0, 1]$
The injected amount feeds back naturally to the edge flow E in the next half-step through Operator 7's $C^T B$ term, closing the loop.

The endogenous injection coefficients $\eta_{\text{inj}}^{(\cdot)}$ guarantee that the injection does not overshoot (the denominator self-adaptively absorbs the magnitude), with no external relaxation factor.

$\mathbf{Harmonic} \to \text{suspended, handed to Operator 17 for adjudication}$

Theorem 14.1 (conservation of the dual pairing on a closed manifold)

Under the premise that the **zero-flux-escape condition** $C_{\text{global}} E_{\text{global}} \equiv 0$ already guaranteed by Operator 7 holds, the Poincaré dual-pairing variation caused by the above curl injection is identically zero:
$\Delta\left(E_s^T\, C_{\text{global}}^T\, B_{s+1/2}\right) \;=\; -\eta_{\text{inj}}^{(\psi)}\, \langle C_{\text{global}} E_s,\; \psi \rangle \;=\; 0$
Proof: $CE_s = 0 \Rightarrow E_s \in \ker \partial_2 = \left(\text{im}\, \partial_2^T\right)^\perp$, so $E_s \perp C^T\psi$ holds for any ψ, and the pairing variation is zero. ■

Corollary 14.1 (constructive guarantee of closed flow): by the exactness of the chain complex $\partial_2 \partial_1 = 0$, any gradient field $E = D_{\text{global}}\varphi$ automatically satisfies $CE = (CD)\varphi = 0$, so the edge flow generated by Operator 6's output potential field falls naturally within the domain of this theorem, **with no extra projection needed**.

Counterexample warning: if E contains a nonzero curl component ($CE \neq 0$, i.e. the manifold is not closed), the injection produces a pairing variation of $\mathcal{O}(\eta \|CE\| \|\psi\|)$; in that case the closure must first be completed within Operator 7 before the solving injection is performed — this is the **hard timing reason** why Phase 3.5 must come after Phase 3.

14.4 Coupled multiple spectral shapes (v1.1 addition · L2-level equations)

Suppose the equation couples the spectral shapes of p shards. Each shard Ω_ν has its own local edge space $\mathbb{R}^{M_{K_\nu}}$ and local sensitivity $\mathbf{w}_k^{(\nu)}$, but the residual must be synthesized in the **global edge space** $C_1 \cong \mathbb{R}^{m_g}$ before a Hodge decomposition can be performed.

Shard embedding map Ξ_ν (isomorphic to the block splicing of Operator 3's D_{global}):
$\Xi_\nu \in \{0,1\}^{m_g \times M_{K_\nu}}, \qquad \Xi_\nu(e_{\text{global}}, e_{\text{local}}) = 1 \iff e_{\text{local}} \in \text{the } e_{\text{global}}\text{-th global edge of } \Omega_\nu$
i.e. a 0-1 selection matrix (column-orthogonal, $\Xi_\nu^T \Xi_\nu = I$). Local vectors are lifted to the global space via $\tilde{\mathbf{w}}_k^{(\nu)} \equiv \Xi_\nu \mathbf{w}_k^{(\nu)}$.

Block Jacobian and residual synthesis:
$J \in \mathbb{R}^{q \times p(K+1)}, \quad J_{j,(\nu,k)} \equiv \frac{\partial \mathcal{F}_j}{\partial \tau_k^{(\nu)}}$

$$\boxed{r \;=\; \sum_{\nu=1}^{p} \Xi_\nu \left(\sum_{k=0}^{K} \left(J^T g\right)_{(\nu,k)} \mathbf{w}_k^{(\nu)} \right) \;\in\; \mathbb{R}^{m_g}}$$

Coupling terms need no special machinery: if $\mathcal{F}$ contains a cross-shard product $\tau_k^{(\mu)} \tau_l^{(\nu)}$, then by the product rule $\frac{\partial \mathcal{F}}{\partial \tau_k^{(\mu)}} = \tau_l^{(\nu)} \cdot (\cdots)$
this factor falls automatically into the (μ, k) block of the block Jacobian. **The coupling is carried entirely by the block structure of the Jacobian**, and the downstream machinery of Operators 15/16/17 is unchanged word for word.

Splicing consistency constraint (a rigid requirement specific to multiple shards):
all shards must be normalized by the **same global** α_n (Operator 13.1), while α_n, λ_2 are updated monotonically by Operator 6's splicing kernel as
$\alpha_n \leftarrow \max\left(\alpha_{n-1},\; \alpha_{\Omega_k} \|P_{\Omega_k,\text{edge}}\|_2^2\right), \qquad \lambda_2(n) \leftarrow \min\left(\lambda_2(n-1),\; \lambda_{2,\Omega_k}\right)$
Since $\max/\min$ are both monotone operations, **the change of the spectral gap ratio** $\gamma = \lambda_2/\alpha_n$ **within the solving inner loop**

is bounded and non-oscillatory, so K_f (Theorem 13.2) remains stable during the iteration. This is the **structural precondition** for coupled multiple spectral shapes to converge.

Theorem 14.2 (the endogenous trust region automatically absorbs the coupling amplification)
After multi-shard synthesis $\|\mathbf{g}_M\|_2$ grows with p (contributions of the shards add up), but the trust-region denominator $1+\tilde{\mathbf{g}}^T A^\dagger \tilde{\mathbf{g}}$ of Operator 15 grows quadratically while the numerator grows only linearly, so the step size is automatically compressed as $\mathcal{O}(1/p)$. **Conclusion: increasing the number of coupled shards requires no step-size re-tuning**.

Complexity and red lines

$\mathcal{O}(\sum_\nu M_{K_\nu} K + m_g + f) = \mathcal{O}(p\, M_K K)$ (multiple shards).
Hard red line: the injection **must never** touch the diagonal entries of M_Ω (Postulate 2); curl cancellation may only be performed in the edge/cycle space. The only legal route for diagonal correction is Operator 8's μ_{trace}. With multiple shards one must also verify that all shards use the **same** α_n; per-shard normalization is forbidden (it would make the moment scales inconsistent and render the product terms on both sides of the coupled equation meaningless).

Operator 15: $\mathcal{N}_{\text{endo}}$, the Endogenous Quasi-Newton Spectral Inverter

(Endogenous Quasi-Newton Spectral Inverter)

Item	Specification
Topological space	$C_1(\mathbb{R}^{m_g}) \to C_1(\mathbb{R}^{m_g})$ (edge-weight increment space)
Input	Edge-space gradient $\mathbf{g}_M \equiv r$, harmonic component h, weighted Hodge Laplacian A
Output	Edge-weight increment ΔM (the graph realization update of the spectral shape)
Timing position	**Phase 3.5 (new solving inner loop, step κ)**

15.1 Endogenous Hessian surrogate (weighted Hodge Laplacian)

$$\mathcal{H}_{\text{endo}} \equiv \mathbf{g}_M \mathbf{g}_M^T + \underbrace{\lambda_2\, DD^T + \frac{\lambda_2^2}{\alpha_n}\, C^T C}_{A\ (\text{weighted Hodge Laplacian } L_1)}$$

Semantics of the three terms:

1. **rank-one Gauss-Newton term**: captures the local curvature of the equation;
2. $\lambda_2 DD^T$ **(gradient regularization)**: the step size is clamped by the spectral gap along connected directions, preventing overshoot across a cut set;
3. $\frac{\lambda_2^2}{\alpha_n} C^T C$ **(curl regularization)**: the step must not destroy the existing cycle structure (the algebraic guarantee of $\Delta\beta_1 = 0$).

15.2 Sherman-Morrison closed-form inversion ★ (v1.1 correction: Newton normalization)

Since $\mathcal{H} = A + \mathbf{g}\mathbf{g}^T$, its Moore-Penrose pseudo-inverse has an exact closed form:

$$\mathcal{H}^\dagger \mathbf{g} = A^\dagger \mathbf{g} - \frac{A^\dagger \mathbf{g}\,(\mathbf{g}^T A^\dagger \mathbf{g})}{1+\mathbf{g}^T A^\dagger \mathbf{g}} = \frac{A^\dagger \mathbf{g}}{1+\mathbf{g}^T A^\dagger \mathbf{g}}$$

where $A^\dagger(\cdot)$ is solved by the **conjugate gradient method** (A is sparse symmetric positive semi-definite), and the iteration first removes the harmonic component to guarantee $x \perp \ker A$. Here $\ker A = \ker(D^T) \cap \ker(C) = \mathcal{H}_1$, so $\dim \ker A = \beta_1$ — **the iteration's component along every homological direction is identically zero**, which is precisely the algebraic root of "the harmonic residual cannot be eliminated by iteration" in Theorem S3.

In what follows the effective gradient after removing the harmonic component is denoted $\tilde{\mathbf{g}} \equiv \mathbf{g}_M - h$ ($\tilde{\mathbf{g}} \perp \mathcal{H}_1$), and $q \equiv \tilde{\mathbf{g}}^T A^\dagger \tilde{\mathbf{g}}$.

> ⚠ **Defect of the v1.0 form (empirically exposed, corrected in v1.1)**

> v1.0 directly took $\Delta M = -\dfrac{A^\dagger \tilde{\mathbf{g}}}{1+q}$. The **per-step relative descent rate of this formula is** $\nabla F^T A^\dagger \nabla F =: \varkappa$, and $\varkappa$ depends on the writing magnitude of the equation residual F:
>
> under the rescaling $F \to sF$, one has $q \to s^4 q$, $\tilde{\mathbf{g}} \to s^2 \tilde{\mathbf{g}}$, so $\Delta M \to -\dfrac{s^2 A^\dagger \tilde{\mathbf{g}}}{1+s^4 q}$ — **the trajectory changes with the artificial scaling of the equation**. Empirically $\varkappa \approx 5.7 \times 10^{-3}$, i.e. about ~175 steps are needed to accomplish the work of one Newton step; and if the equation is rewritten as $100F$, the trajectory immediately differs. This violates the system's very foundation of "dimension-free, no external parameters" (Postulate 3).

The corrected update step (Theorem 15.1'):

$$\boxed{\Delta M \;\equiv\; -\eta_N\, A^\dagger \tilde{\mathbf{g}}, \qquad \eta_N \equiv \frac{F^2}{q} = \frac{F^2}{\tilde{\mathbf{g}}^T A^\dagger \tilde{\mathbf{g}}}}$$

Scale-invariance proof: when $F \to sF$, one has $\tilde{\mathbf{g}} = F\nabla\tilde{F} \to s^2\tilde{\mathbf{g}}$, $q \to s^4 q$, $F^2 \to s^2 F^2$, hence $\Delta M \to -\dfrac{s^2F^2}{s^4 q} s^2 A^\dagger \tilde{\mathbf{g}} = \Delta M$, **strictly invariant**. ■

Newton completeness: after linearization, $\Delta F = \nabla F^T \Delta M = \dfrac{\tilde{\mathbf{g}}^T}{F}\left(-\eta_N A^\dagger \tilde{\mathbf{g}}\right) = -\dfrac{\eta_N q}{F} = -F$, i.e. **zeroed in one step** (the v1.0 form can only descend by the fraction $\varkappa$).

Theorem 15.1 (Endogenous trust-region theorem) ★v1.1

After adopting the Newton normalization η_N, the step size requires an additional **endogenous trust region**, otherwise it overshoots in the nonlinear region. Take:

$$\|\Delta L\|_F \;=\; \left\| D^T \mathrm{diag}(\Delta M)\, D \right\|_F \;\le\; \tfrac{\lambda_2}{2}$$

i.e. **the step must not cross half the spectral gap**. This is a direct corollary of Postulate 3: the spectral gap ratio $\gamma = \lambda_2/\alpha_n$ is the only dimensionless endogenous scale; if a single step crosses $\lambda_2/2$, then γ itself (along with K, $\varepsilon_{\text{endo}}$, and the thresholds) self-destructs. Equivalently: $\|\Delta Y\| \lesssim \gamma$, i.e. the drift of Y does not exceed γ times the spectral support — which is precisely the necessary and sufficient scale for the linearization to remain valid. If exceeded, scale back proportionally by $\dfrac{\lambda_2/2}{\|\Delta L\|_F}$.

Hence the step size of this system is **not given by any external parameter**: η_N is given endogenously by the equation residual, and the trust-region radius is given endogenously by λ_2.

> ⚠️ **v1.3 correction (exposed by the 50-equation stress test): the trust region must actually be put into effect, and this must be done in the form of a backtracking line search.**
> When v1.1 replaced $1/(1+q)$ by η_N, **it deleted the trust-region factor as well** — the formula $\Delta M = -\eta_N A^\dagger \tilde{g}$ above does not contain the $\lambda_2/2$ proportional scaling. Empirically, for a strongly nonlinear instance (bench B2-E02, $\sum a_k \tau_k^2 = c$) the Newton step is $\|\Delta M\| \approx 0.23$, while $\Psi = F^2/2$ instead rises ($-6.85\times10^{-3} \to -7.56\times10^{-3}$), convergence degenerates into a slow oscillation with ratio ≈ 0.85, and the monotonicity red lines (V5/V6) fail.
> **Correction**: insert an **endogenous backtracking line search in Phase 3.5b** (§15.4) between Operator 15 and Operator 16, with "Ψ strictly decreases" as the sole acceptance criterion, floating-point resolution as the termination floor, and a bisection back-off factor of $1/2$. It contains no externally tunable parameter (the criterion is a comparison of the endogenous quantity Ψ with its own historical value), in accordance with Postulate 3.

Theorem 15.2 (monotone decrease of the Dirichlet energy)

The first-order Dirichlet-energy variation along ΔM is
$\Delta\mathcal{E}_D \;=\; \tilde{\mathbf{g}}^T \Delta M \;+\; \mathcal{O}(\|\Delta M\|^2) \;=\; -\eta_N\, q \;=\; -F^2 \;\le\; 0$
with equality if and only if $F = 0$. That is, each inversion step necessarily makes the whole-graph Dirichlet energy non-increasing (and exactly $-F^2$ within the Newton region), of the same origin and isomorphic to Operator 8's red line "$\Delta\mathcal{E}_{D,\text{trace}} \le 0$ strictly monotone convergence".

Theorem 15.3 (orthogonality constraint on the update direction)

$\Delta M \in \mathrm{im}(\partial_1) \oplus \mathrm{im}(\partial_2^T) = \mathcal{H}_1^\perp$, i.e. the update always lies in the orthogonal complement of the harmonic space.
Proof: $\Delta M \propto A^\dagger(\mathbf{g} - h)$, and the range of $A^\dagger$ is $(\ker A)^\perp = \mathcal{H}_1^\perp$. ■

15.3 Endogenous momentum acceleration (optional)

$\beta_{\text{mom}} \equiv 1 - \frac{\lambda_2}{\alpha_n} \in [0, 1)$
the better the conditioning (the larger the spectral gap ratio), the stronger the momentum — driven entirely by the endogenous spectral assets, with no external hyperparameter.

⚠️ **Empirical warning (v1.0)**: the trust region of Theorem 15.1 guarantees only **single-step** boundedness, not monotonicity after momentum accumulation. Empirically, at $\beta_{\text{mom}} = 0.798$ the residual oscillates and crosses zero, and the monotonicity of Ψ fails (see §7, V6 comparison). **Hence momentum is off by default**; if enabled, one must add the backtracking arbitration "retreat whenever the energy rises", and that backtracking criterion itself must still be endogenous (e.g. taking the previous value of $\mathcal{E}_D$ as reference). This is a **known limitation** explicitly stated in specification v1.0.

15.4 Endogenous backtracking line search (v1.3 addition · Phase 3.5b)

Since η_N no longer carries a trust region (§15.2 ⚠️), single-step overshoot must be caught by **backtracking**. Define the objective functional
$\Psi(M) \equiv \frac{1}{2} F(M)^2 \geq 0,$
and at step κ perform a bisection backtracking along the Newton direction ΔM, taking

$$t_\kappa \equiv \max\left\{ t = 2^{-j},\ j = 0, 1, 2, \ldots : \Psi(\max\big(m + t\,\Delta M,\ m_{\min}\big)) < \Psi(m) \right\}$$

- **Acceptance criterion**: Ψ strictly decreases. Since Ψ is already the residual squared, its absolute decrease suffices; **there is no need to introduce an Armijo constant** (any Armijo parameter would be an external quantity);
- **Termination floor**: $t_\kappa \leq 10^{-14}$ or $\|t_\kappa\, \Delta M\| \leq 10^{-12} m_{\min}$, at which point `step_dead = True` is set and handed to Operator 17 for adjudication (§17.1/§17.2);
- **Recomputation**: $F(m + t\Delta M)$ reuses the **exactly identical** encoding pipeline of Operator 13 (the same α_n, λ_2, K), so it is **of the same origin and comparable** with the inner-loop residual, introducing no additional approximation;
- **Relation to the trust region**: $t_\kappa < 1$ is equivalent to one adaptive trust-region retraction, and fits the true curvature better than the fixed $\lambda_2/2$ radius — it is the **executable stand-in** for Theorem 15.1.

Empirical (bench B2-E02, 14 steps of strictly monotone convergence): $t_\kappa = 0.5$ ($\kappa = 3, 4, 5$), $t_\kappa = 0.25$ ($\kappa = 12$), and $t_\kappa = 1$ otherwise; after backtracking takes effect the trajectory is strictly monotone, and the V-B independent residual drops from 7.30×10^{-1} to 1.29×10^{-1} ($<$ threshold $\gamma = 0.1761$).

> **On "does a line search violate the no-external-parameters principle"**: no, it does not. The backtracking criterion (decrease of Ψ) and the quantity compared against (Ψ's own historical value) are both endogenous; the bisection factor $1/2$ is a discretization choice rather than a tunable hyperparameter; and the floor 10^{-14} is the floating-point resolution. This is the same paradigm as Operator 16's "bisect along the feasible-cone line to take the largest feasible step", viewed as two projections of one paradigm.

Complexity and red lines

$\mathcal{O}(M_K \cdot n_{\text{cg}})$, with n_{cg} endogenously bounded by the condition number of A ($\kappa(A) \leq \kappa_\Omega^2$).
Hard red line: when A is singular ($\beta_1 > 0$) the use of direct inversion or dense factorization is **forbidden**; one must first remove the h component before entering CG, otherwise the iteration drifts along the null space and causes numerical escape.

Operator 16: $\mathcal{C}_{\text{hankel}}$, the Spectral-Feasibility Hankel Clamp

(Spectral Feasibility Hankel Clamp)

Item	Specification
Topological space	$\mathbb{R}^{K+1} \to \mathcal{K}_{\text{feas}}$ (feasible moment cone)
Input	Moment vector recomputed after the update $\boldsymbol{\tau}^{(+)}$ (in power-moment form μ_k)
Output	Feasible moments $\tilde{\boldsymbol{\tau}}$; feasibility warning flag
Timing position	Phase 3.5 inner loop, immediately after Operator 15

16.1 Feasibility criterion (Hausdorff moment problem)

The sequence $\{\mu_k\}_{k=0}^{2K'}$ are the moments of some non-negative measure supported on $[-1,1]$ **if and only if** two Hankel matrices are simultaneously positive semi-definite:

$$H_{ij}^{(0)} \equiv \mu_{i+j} \succeq 0, \qquad H_{ij}^{(1)} \equiv \mu_{i+j} - \mu_{i+j+2} \succeq 0$$

(the second corresponds to $(1-\lambda^2)d\sigma \geq 0$. The Chebyshev moments τ_k and the power moments μ_k are mutually inverse through the lower-triangular transformation given by the power expansion of T_k, and can be converted locally.)

16.2 Two-stage clamping valve (strictly isomorphic to Operator 10)

$$\mathcal{C}_{\text{hankel}}(\boldsymbol{\mu}) \equiv \begin{cases} \Pi_{\mathcal{K}}\left(\boldsymbol{\mu}\right), & \text{if } \lambda_{\min}\left(H\right) < -\theta_{\text{endo}} \quad (\text{hard interception} \cdot \text{spectral projection back to the feasible cone}) \\ \boldsymbol{\mu} \;+\; \delta_{\text{soft}} \cdot \Pi_{\text{supp}}, & \text{otherwise} \quad (\text{soft contraction} \cdot \text{endogenous spectral-support clamping}) \end{cases}$$

- **Endogenous threshold** $\theta_{\text{endo}} \equiv \dfrac{\lambda_2}{\alpha_n}$ (the tolerance is of the same order as the endogenous minimum resolvable scale)
- **Hard interception** $\Pi_{\mathcal{K}}$ (**the canonical main formula · trust-domain rollback**): along the line "previous feasible moments $\boldsymbol{\mu}_{\text{ref}} \to$ new moments $\boldsymbol{\mu}$", take the largest feasible step
  $\Pi_{\mathcal{K}}(\boldsymbol{\mu}) \equiv \boldsymbol{\mu}_{\text{ref}} + t^{\star}\left(\boldsymbol{\mu} - \boldsymbol{\mu}_{\text{ref}}\right), \qquad t^{\star} \equiv \max\left\{t \in [0,1] \;:\; H^{(0)}, H^{(1)}\left(\boldsymbol{\mu}_{\text{ref}} + t\,\Delta\boldsymbol{\mu}\right) \succeq -\theta_{\text{endo}} I\right\}$
  Since the feasible set is convex and $\boldsymbol{\mu}_{\text{ref}}$ is an interior point of it, $t^{\star}$ necessarily exists and can be found exactly by bisection. **This is the counterpart of the endogenous trust region in moment space** — the step size is adjudicated by feasibility itself.
- **Soft contraction** Π_{supp}: clamp the moment support back to the endogenous interval $[\gamma, 1]$, $\gamma = \lambda_2/\alpha_n$, realized by shift-rescaling (rescaling with respect to $\frac{\lambda-\gamma}{1-\gamma}$)
- **Alternative (spectral projection, fallback only)**: apply the spectral projection $\lambda_{\min} \to \max(\lambda_{\min}, 0)$ to the Hankel matrices, then recover the moments by anti-diagonal averaging. **Empirically this path converges slowly under alternating projection** (after 8 alternating steps there remains an $\mathcal{O}(10^{-2})$ negative eigenvalue), so it is not adopted as the main formula

Theorem 16.1 (feasibility preservation)

After $\mathcal{C}_{\text{hankel}}$, the moment sequence necessarily corresponds to some non-negative measure supported on $[\gamma, 1]$, i.e. **a true spectral shape (in the moment sense) necessarily exists**; moreover $\Pi_{\mathcal{K}}$ is the nearest-point projection onto the feasible cone in the Euclidean norm, so $\|\tilde{\boldsymbol{\mu}} - \boldsymbol{\mu}\|_2$ is minimized.

Theorem 16.2 (compatibility with trace sovereignty)

The clamping acts only on the moment-space representation and **does not modify any diagonal entry of** M_{Ω}, so $\mathrm{Tr}(M_{\Omega}) \equiv 0$ and local trace sovereignty (Postulate 2) are not violated. The moment deviation introduced by the clamping is **not absorbed locally**; instead it **flows back into Operator 14 as a new residual** and re-enters the routing — guaranteeing closed-loop self-consistency and forbidding the silent discarding of infeasible components.

Complexity and red lines

$\mathcal{O}(K^3)$, since $K \leq K_{\text{cap}} = \mathcal{O}(\ln N_K)$ and $N_K = \mathcal{O}(1)$, so the whole is at the **constant order**.
Hard red line: when K_{cap} truncates, the Hankel criterion is only a **necessary condition** (high-order moments are missing); in this case Operator 16 must output the "feasibility undetermined" warning flag and must not claim verification.

Operator 17: $\mathcal{A}_{\text{sol}}$, the Solution-Branch Betti Audit and Termination Arbiter

(Solution-Branch Betti Audit & Termination Arbiter)

Item	Specification
Input	Three-component norm sequence, β_0, β_1, adaptive spectral window $\tau(s)$ (Operator 9)
Output	Verdict: continue / trigger topological stitching / declare solved / output obstruction certificate
Timing position	Phase 4 (same layer as Operators 8 and 9, but before them)

17.1 Two-level endogenous convergence criteria (v1.1 · relative, dimensionless)

Taking the first-step residual $\|r_0\|_2$ as the endogenous reference scale, define a unified relative tolerance $\epsilon_{\text{endo}} \equiv \frac{\lambda_2}{\alpha_n}\|r_0\|_2$. The solution mode is chosen according to the user's needs:

Mode	Enabling condition	Criterion	Monitored degrees of freedom
Exact solution	$\beta_1 = 0$, or the user explicitly requires it	$\|D\phi\|, \|C^T\psi\| \le \epsilon_{\text{endo}}$ and $\|h\| \le \epsilon_{\text{endo}}$	m_g
Homological solution (default)	$\beta_1 > 0$	**only** $\|h\|_2 \le \epsilon_{\text{endo}}$	β_1

In the homological-solution mode no requirement is placed on the gradient/curl components — they are absorbed naturally by Operators 6/7 in subsequent pulses (Definition 3.1). This brings two benefits:

1. **the monitored dimension drops from m_g to β_1**, greatly reducing the degrees of freedom tracked by the convergence decision;
2. upon reaching a homological solution one may immediately proceed to Operator 18 to realize the graph structure, **without waiting for the exact components to fully dissolve**.

Properties: generated entirely from the endogenous spectral assets and the first-step residual, **dimensionless and with no external tolerance parameter**, and automatically adapting to the overall magnitude of the equation.

⚠️ **Empirical qualification (v1.1): when the homological-solution criterion really terminates early**

Since $r(M) = F(M)\nabla F(M)$ and $h(M) = F(M)\,\Pi_{\mathcal{H}_1}\nabla F(M)$, the ratio of the two,
$\rho_h(M) \equiv \frac{\|h(M)\|}{\|r(M)\|} = \frac{\|\Pi_{\mathcal{H}_1}\nabla F\|}{\|\nabla F\|}$
is **independent of $F(M)$**. Empirically, on a test complex with $\beta_1 = 1$, ρ_h is stable at ≈ 0.945 for almost all initial values (min 0.9446 / max 0.9503 / mean 0.9473), i.e. **the harmonic fraction of the residual is approximately a constant that does not change with the iteration**.

Corollaries (important):

1. the harmonic channel **cannot be compressed independently**, and can only vanish together with $\|r\| \to 0$;
2. therefore when $\rho_h > \gamma$ the homological-solution criterion **coincides** with the exact-solution criterion at the convergence point, and the two-level criteria produce no bifurcation point (empirically V15 "no bifurcation point");
3. the necessary and sufficient condition for the homological-solution criterion to truly terminate early is $\rho_h(M_0) \le \gamma$ — i.e. the initial value's harmonic fraction is itself below the endogenous minimum resolvable scale;
4. if ρ_h is identically $\mathcal{O}(1)$ (e.g. 0.945 here), then $\|h\| \le \epsilon_{\text{endo}}$ can only be attained by an overall decrease of $\|r\|$, and **in this case there is no homological solution "earlier than the exact solution"** — the full iteration must be run; if the iteration stalls at $\|h\| > \epsilon_{\text{endo}}$ while the gradient/curl have already met the criterion, then OBSTRUCTION is declared and it is handed to Operator 9 for Betti stitching.

> This conclusion clarifies the true benefit boundary of "homological solutions": their value is **not faster convergence, but relaxing the "acceptable solution" from $\{r = 0\}$ to $\{h = 0\}$**, so that when the equation is intrinsically not exactly solvable (topologically obstructed) one can still deliver a graph structure correct in the homological sense.

> ⚠️ **v1.3 correction (exposed by the 50-equation stress test): `OBSTRUCTION` is a structural false-positive branch.**
> From the identity $r = F\nabla F$, $\rho_h = \|h\|/\|r\|$ is independent of F (empirically ≈ 0.95 above). Hence "$\|h\| \le \epsilon_{\text{endo}}$" is equivalent to "$\|r\| \le \epsilon_{\text{endo}}/\rho_h$", while "the exact part has converged" corresponds to "$\|r\| \le \epsilon_{\text{endo}}$". The criterion band between the two criteria is
> $\frac{\epsilon_{\text{endo}}}{\rho_h} < \|r\| \le \frac{\epsilon_{\text{endo}}}{\rho_h}\cdot\frac{1}{\gamma}$ $\quad(\text{here} \approx 4.3\,\epsilon_{\text{endo}})$,
> **as wide as $1/\gamma$ times**. Any F falling into this band simultaneously satisfies "exact part converged, harmonic part not converged", and is thereby **misjudged as a topological obstruction** — yet the problems inside this band are in fact well-posed and solvable.
> **Consequence**: a well-posed problem is handed to **Operator 9, which changes β_1**, directly violating the red line "$\Delta\beta_1 \equiv 0$ hard lock" of §17.3 (empirically bench B2-E02).
> **Correction**: declaring `OBSTRUCTION` must **additionally require the condition "the update direction is exhausted"** — i.e.

> the t_κ of §15.4 hitting the floor (`step_dead`), or $\|\Delta M\| \to 0$. Only when the residual cannot move even in a direction is it admitted to be a genuine topological obstruction; otherwise it is invariably judged `CONTINUE` (continue iterating) or `STALLED` (numerical stall, no topological change triggered).
> **Verification**: after suppressing this false positive, only 2 of the 50 equations still fell into the "tautology" channel (see the `exhausted` condition of §17.2); with that closed, **all 50/50 are SOLVED**.

17.2 Verdict routing table (v1.3 revision)

Let `below` denote "all three channels meet the criterion", `exhausted` denote "the residual decrease has been wrung dry", `step_dead` denote "the update direction is exhausted (the §15.4 line search hit the floor)", `pert` denote "the exact components ($D\phi, C^T\psi$) meet the criterion", and `harm` denote "the harmonic component meets the criterion".

Condition	Verdict	Action
`below` **and** (`exhausted` **or** `step_dead`)	**SOLVED**	Hand to Operator 18; output $M_\Omega^\star$ and $\sigma^\star$ (both exact solutions and homological solutions go this way; when the residual is exact it is registered for absorption by Operators 6/7)
not `below`, and the update direction is not exhausted	**CONTINUE**	$\kappa \leftarrow \kappa + 1$, return to Operator 15
`step_dead`, and `pert` and ¬ `harm` (**only** the harmonic part remains)	**OBSTRUCTION**	Hand to Operator 9 to perform Betti stitching (changes β_1), restart the inner loop
`step_dead`, and not "only the harmonic part remains"	**STALLED** (numerical stall)	Report a numerical warning, **do not trigger a topological change** (protecting the $\Delta\beta_1 \equiv 0$ red line)
$\|h\|$ still does not decrease after $\tau(s)$ stitching steps	**UNSOLVABLE**	Output the obstruction certificate h, terminate and warn

The `exhausted` (wrung-dry) condition (v1.3 addition):
$|F_\kappa| \;\geq\; (1-\gamma)\,|F_{\kappa-1}|,$
i.e. the residual decrease of the previous step is below the relative fraction γ — the information carried by the Newton direction has been wrung dry. It stands alongside "$\|r\| \leq \epsilon_{\text{endo}}$" and together they constitute the criterion of `SOLVED`.

> **Why this condition must be added**: the V-B independent-residual verification and the termination criterion **take the same threshold** ϵ_{endo}. If one terminates merely on "first crossing ϵ_{endo}", then any instance that happens to fall on either side of the threshold within the $1/\gamma$ band (empirically B4-E02 $= 0.1726$, B5-E02 $= 0.1787$ vs $\gamma = 0.1761$) will be judged "solved", but the independently recomputed residual is still above the threshold and V-B FAILs — **the verification degenerates into a tautology**. After adding `exhausted`, the V-B of 49 empirical equations falls to $10^{-14} \sim 10^{-9}$, and B2-E02's V-B drops from 7.30×10^{-1} to 1.29×10^{-1}.

The `step_dead` (direction-exhausted) condition (v1.3 addition): see the line-search floor of §15.4. It is a necessary condition for `OBSTRUCTION` (§17.1 ⚠️), and also one of the sufficient conditions for `SOLVED` (near the threshold, being unable to move in any direction counts as converged).

Verdict priority: Operator 17 must execute before Operators 8/9 (first in Phase 4), because its verdict determines whether Operator 9's stitching is needed — if the conservative clamping were done first, it might flatten the topological information awaiting stitching in advance.

17.3 Multi-branch solution audit

The multiple fixed points of the equation correspond to multiple basins of attraction of the residual manifold, with the number of branches counted by β_0. Throughout the inner loop $\Delta\beta_1 \equiv 0$ **is hard-locked** (unless Operator 9 actively stitches), preventing the iteration from spuriously proliferating/merging solution branches (reusing Operator 9's unique-ordering arbitration chain $\mu_{\text{trace}} \to \text{blocking gate} \to \mathcal{O}_{\text{stitch_dual}}$).

17.4 Handover to Operator 18

Once solved (whether exact or homological), the result is invariably handed to **Operator 18** $\mathcal{R}_{\text{homo}}$ to complete the graph-structure realization. Operator 17 is responsible only for the verdict, not for the realization.

Operator 18: $\mathcal{R}_{\text{homo}}$, the Homological Solution Realizer (v1.1 addition)

(Homological Solution Realizer)

Item	Specification
Topological space	$\mathcal{S}_{\text{hom}} \subset \mathbb{R}^{m_g} \;\to\; M_\Omega^\star$ (spin-adjacency structure)
Input	The graph M judged "solved", the three residual components, the moment vector $\boldsymbol{\tau}$, the Fiedler vector v_2
Output	Canonical homological representative $M^\star_{\text{canon}}$; spectral shape $\sigma^\star$; homological-distance certificate
Timing position	Last in Phase 4 (executed once after Operator 17 judges "solved")

18.1 The five realization steps

1. **Homology-class confirmation**: verify $\|h\| \le \epsilon_{\text{endo}}$. If it does not hold, flow back to Operator 9; if it holds, record the current homology class $[M]$.
2. **Registration of the exact residual** (do not discard!): register the remaining exact residual $D\phi + C^T\psi$ as an **amount to be absorbed** by Operators 6/7, to be consumed in a subsequent unified pulse. The registered amount itself constitutes the error budget of the solution.
3. **Canonical-representative selection**: within the homology class take the one minimizing the Dirichlet energy as the output: $M^\star_{\text{canon}} \;\equiv\; \arg\min_{M' \in [M]} \mathcal{E}_D(M')$
   Practical implementation: perform energy descent along the $\mathcal{H}_1^\perp$ direction (i.e. the update direction of Operator 15) until $\Delta\mathcal{E}_D \approx 0$; Theorem 15.3 guarantees that this direction does not leave the homology class.
4. **Spectral-shape reconstruction** (μ_{realize} micro-operator): invert the spectral density from the moment vector. Prefer Stieltjes inversion (§13.4):
   $\rho^\star(\hat\lambda) = \frac{1}{\pi}\,\text{Im}\Big[\sum_k c_k(\hat\lambda + i\gamma)\,\tau_k\Big]\,, \qquad \text{imaginary offset endogenously } \varepsilon_{\text{endo}} = \gamma$
   In the degenerate case (or when K_f is insufficient), switch to the direct Chebyshev expansion $\rho^\star \approx \sum_k \tau_k T_k$ and apply a non-negativity clamp.
5. **Sign and spin assignment**: use Operator 3's Fiedler sign gate to assign the spin sign inversely, $\Gamma(k,j) = \text{sgn}(v_2(j))$, then invert the edge weights via Operator 5, $W_e = \exp(\alpha_n/c_e) - 1$, and substitute into Operator 4's degree relation for verification.

Theorem 18.1 (local well-definedness of the canonical representative)

Linearizing in a neighborhood of the homological solution $M^\star$, one has $A \succ 0$ along $\mathcal{H}_1^\perp$ (A being Operator 15's weighted Hodge Laplacian, whose kernel is exactly $\mathcal{H}_1$), so $\mathcal{E}_D$ is strictly convex on $\mathcal{H}_1^\perp \Rightarrow \arg\min \mathcal{E}_D$ **exists locally and is unique**.

Note: global uniqueness does not hold — different homology classes ($h \neq 0$) are not comparable; this is precisely the meaning of Theorem S5, "uniqueness holds only in the sense of homology classes".

18.2 Output certificates

Operator 18 must output the following three items together with the solution, as a credibility statement about the solution:

Certificate item	Definition	Semantics
Homological residual $\|h\|_2$	the norm of the harmonic projection of the residual	$= 0 \Rightarrow$ the homological solution holds
Homological distance $d_{\text{hom}}(M^\star, M_{\text{init}})$	$\|h(M^\star) - h(M_{\text{init}})\|_2$	how far it has moved from the initial value (in the topological sense)
Exact-residual budget $\|D\phi + C^T\psi\|_2$	the amount to be absorbed by Operators 6/7	the residual error of the exact solution

Complexity and red lines

$\mathcal{O}(M_K \cdot n_{\text{cg}} + N_K^2)$, executed only once.
Hard red lines:

- the reconstructed spectral density must be non-negative, else it is re-clamped by Operator 16 (outputting a negative spectral density is forbidden);
- when $\beta_1 = 0$ the homological-solution set degenerates to the whole space (degeneracy warning, §2-bis), in which case **the canonical-representative criterion (minimum Dirichlet energy) is the sole basis for selection**, and step 3 must not be omitted.

---

4. Extended Pipeline Timing (embedded into the Version 18.0 four-phase pulse)

A new **Phase 3.5: spectral-shape equation solving inner loop (Solve Inner Loop, step κ)** is added, located after Phase 3 field closure and before Phase 4 conservation. The solving inner loop **does not change the advancement of the structural pulse** s; it merely performs sub-step iterations inside a unified pulse, so all network Actors still advance in strict lockstep.

Phase	Executed operators	Input	Attributes
Phase 1	Op4 → Op6 → **Op13**	$M_\Omega(s)$, (α_n, λ_2)	Op13 reuses Op6's Lanczos subspace, **zero extra spectral decomposition**; outputs the moment vector and sensitivity set
Phase 2	Op5 → Op2 → Op3	unchanged	unchanged
Phase 3	Op7 → μ_{betti} → **Op14**	closed manifold + equation $\mathcal{F}$	Op14 executes jointly with the half-step $B_{s+1/2}$; three-component residual routing, conservative pairing (Theorem 14.1)
Phase 3.5 (new)	**Op15** → **line search (3.5b)** → **Op16** → **criterion**	g_M, h, A	Endogenous quasi-Newton single-line closed form; **backtracking line-search sub-step** (§15.4); Hankel feasibility clamping; $\kappa \leftarrow \kappa + 1$
Phase 4	**Op17** → Op8 → Op9 → compensation → **Op18**	three-component norms, $\beta_0, \beta_1, \tau(s)$	Op17's verdict takes priority: obstruction → Op9 stitching; solved (exact/homological) → **Op18 realization**; then Op8/Op9 conservation

Multi-shard timing supplement: the Op13 of p shards may be executed **in parallel** in Phase 1 (each shard is independent, with no synchronization dependency), but they must use the **same** α_n (§14.4); residual synthesis (Op14) is performed centrally in Phase 3, since the Hodge decomposition must be completed in one shot over the global edge space and shards may not project separately and then add (the sum of shard projections $\neq$ the global projection, unless the shards' edge sets are orthogonal).

Nesting relation of the solving pulses: the outer unified pulse s advances the evolution of the graph structure, while the inner solving pulse κ converges the equation under the same structure. If κ exceeds the endogenous spectral window $\tau(s)$ and still has not converged, it is handed to Op17 for adjudication (continue / stitch / declare unsolvable).

---

5. Computational-Complexity Invariants Comparison Table

Operator	Core operations	Complexity	Notes
Op13 $\mathcal{E}_{\text{mom}}$	Chebyshev three-term recurrence + boundary-column quadratic forms	$\mathcal{O}(N_K M_K K)$	both N_K, K endogenously clamped by the local horizon, effectively $\mathcal{O}(1)$; no eigendecomposition
Op14 $\mathcal{R}_{\text{hodge}}$	sensitivity synthesis + two pseudo-inverse projections	$\mathcal{O}(M_K K + M_K + f)$	the pseudo-inverses reuse Op6/Op10's $\mathcal{O}(1)$ facilities

Operator	Core operations	Complexity	Notes
Op15 $\mathcal{N}_{\text{endo}}$	Sherman-Morrison closed form + one CG	$\mathcal{O}(M_K n_{\text{cg}})$	single-line closed form, no dense factorization; n_{cg} endogenously bounded by $\kappa(A)$
Op16 $\mathcal{C}_{\text{hankel}}$	$K \times K$ Hankel spectral projection	$\mathcal{O}(K^3)$	$K = \mathcal{O}(\ln N_K)$, constant order
Op17 $\mathcal{A}_{\text{sol}}$	verdict (the homological-solution mode monitors only β_1 degrees of freedom)	$\mathcal{O}(M_K + f)$	reuses Op9's arbitration chain
Op18 $\mathcal{R}_{\text{homo}}$	canonical-representative selection + Stieltjes spectral reconstruction + Fiedler sign assignment	$\mathcal{O}(M_K n_{\text{cg}} + N_K^2)$	executed once only upon declaring a solution

Overall multi-shard overhead: $\mathcal{O}(p\, M_K K)$ (p being the number of coupled shards); parallelizable within Phase 1, serial from Phase 3 onward.

Global property: the extension layer introduces no $\mathcal{O}(n^3)$ or globally synchronized decomposition operation; all operators are at the local constant order in the thermodynamic limit, consistent with Version 18.0's complexity red line.

---

6. Methodology Summary: the Six-Step Solving Method

For "an equation formed by unknown spectral shapes", the standard solving flow is:

1. **Encoding** (Op13): encode each unknown spectral shape σ_ν as a Chebyshev moment vector $\boldsymbol{\tau}^{(\nu)}$. For polynomial kernels use $K = \lceil \alpha_n / \lambda_2 \rceil$; **for non-polynomial kernels use the tail-sum criterion** K_f ($\sum_{k>K} |c_k| \le \gamma \max |c|$, §13.4 v1.2); **whenever kernel expansion is used, the zero mode must be stripped first** (§13.5) — *turn functions into finite vectors*.
2. **Algebraization** (Definition 1.2): turn the equation $\mathcal{F}(\sigma) = 0$ into an equation in moment space. **Coupled multiple spectral shapes** manifest as a block Jacobian $J \in \mathbb{R}^{q \times p(K+1)}$, with coupling terms falling automatically into the block structure via the product rule — *turn functional equations into algebraic equations*.
3. **Routing** (Op14): the residual is synthesized via the shard embeddings Ξ_ν and the sensitivity vectors into a global 1-chain, and a Hodge three-component decomposition is performed — *turn the residual into topological components*.
4. **Inversion** (Op15 → 3.5b → 16): the endogenous quasi-Newton single-line closed form updates the edge weights, **the backtracking line search (§15.4) contracts the step to guarantee Ψ strictly decreases**, and the Hankel clamp guarantees the spectral feasibility of the solution — *iterate to convergence in graph space*.
5. **Verdict** (Op17): **the default homological-solution criterion** $\|h\| \le \epsilon_{\text{endo}}$, plus the `exhausted` (wrung-dry) or `step_dead` (direction-exhausted) conditions (§17.2 v1.3); only when **the direction is exhausted and only the harmonic part remains** is a topological obstruction declared and the result handed to Operator 9 for stitching; otherwise the verdict is "continue" or "numerical stall" — *deliver a solution or prove it does not exist*.
6. **Realization** (Op18): within the homology class take the canonical representative minimizing the Dirichlet energy, reconstruct the spectral density by Stieltjes inversion, and assign spins via the Fiedler sign gate — *output the graph structure + three credibility certificates*.

The three "no's" of this methodology:

- **No whole-network eigendecomposition** (Postulate 1; closed walks + three-term recurrence throughout)
- **No external parameters introduced** (Postulate 3; all thresholds are generated endogenously from λ_2/α_n)
- **No touching of shard diagonal entries** (Postulate 2; curl cancellation is performed only in the edge/cycle space)

Semantics of the solution (v1.1): the default output is a **homological solution** ($h = 0$, i.e. the residual is exact) rather than an exact solution ($r = 0$). The difference between the two is measured by $\|D\phi + C^T\psi\|$, written into the certificate as the "exact-residual budget" and handed to Operators 6/7 for absorption in subsequent pulses. Uniqueness **holds only in the sense of homology classes** (Theorem S5).

---

7. v1.0 Verification Protocol

#	Verification item	Expectation	Basis
V1	Moment-encoding correctness: τ_k computed by recurrence vs. `eigvalsh` direct computation	relative error $< 10^{-10}$	Theorem S1
V2	Sensitivity closed form vs. central differences	relative error $< 10^{-6}$	Eq. 13.3
V3	Hodge orthogonality: $\langle D\phi, C^T\psi\rangle = \langle D\phi, h\rangle = \langle C^T\psi, h\rangle = 0$	≈ 0	Theorem S3
V4	Harmonic criterion: $D^T h = 0$ and $Ch = 0$; $\dim \mathcal{H}_1 = \beta_1 = \nu - \mathrm{rank}(\partial_2)$, $\nu = m_g - n + \beta_0$	holds (numerical rank)	Theorem S3 + terminology convention
V5	Monotone decrease of the residual for a solvable equation (e.g. $\tau_2 - c = 0$)	$\|r_{\kappa+1}\| \le \|r_\kappa\|$	Theorem 15.2
V6	Dirichlet energy non-increasing	$\Delta\mathcal{E}_D \le 0$	Theorem 15.2
V7	Update-direction orthogonality: $\Delta M \perp \mathcal{H}_1$	$\langle \Delta M, h\rangle \approx 0$	Theorem 15.3
V8	Both matrices positive semi-definite after Hankel clamping	$\lambda_{\min}(H^{(0)}), \lambda_{\min}(H^{(1)}) \ge -\epsilon$	Theorem 16.1
V9	Topological-obstruction test case: requiring a nonzero harmonic residual on a $\beta_1 = 0$ graph	Op17 correctly warns and $\|h\|$ does not decrease	Theorem S3
V10	Closed-flow conservation: taking $E = D\varphi$ (automatically satisfying $CE = 0$), after injecting ψ, $\Delta(E^T C^T B)$	$= 0$	Theorem 14.1
V11	Counterexample comparison: taking a non-closed flow with $CE \neq 0$ and injecting likewise	$\Delta \neq 0$, confirming the timing necessity	Corollary 14.1 note

v1.1 new verification items (coupled multiple spectral shapes / non-polynomial kernels / homological solutions)

#	Verification item	Expectation	Basis
V12	Coupled multiple spectral shapes: two-shard equation $\tau_2^{(1)}\tau_2^{(2)} - c = 0$, block Jacobian + shard embedding Ξ_ν	residual decreases monotonically and a solution is adjudicated	§14.4
V7b	Equation scale invariance: ΔM must be invariant under $F \to sF$ ($s = 1, 10, 100$)	new form relative offset $< 10^{-8}$; the v1.0 old form deviates significantly (scale-dependent)	Theorem 15.1'
V13	Non-polynomial kernel $\ln\hat{\lambda}$: Chebyshev coefficient sum $\mathbf{c}^T\boldsymbol{\tau}^{\mathrm{red}}$ vs. reduced spectral integral (zero-mode stripping first)	empirical error $\le$ Bernstein tail bound $\sum_{j>K_f}\|c_j\|$, and $\le \gamma\|\mathrm{exact}\|$	§13.4 / §13.5
V14	Bernstein ellipse rate: empirical $\ln$-kernel coefficient decay rate vs. $\rho = \frac{1+\gamma}{1-\gamma} + \sqrt{\left(\frac{1+\gamma}{1-\gamma}\right)^2 - 1}$	empirical decay rate of the same order as the theoretical ρ	Theorem 13.2
V15	Homological solution vs. exact solution: the same equation on a $\beta_1 > 0$ complex under the two criteria	homological-solution step count $\le$ exact-solution step count, and $\|h\|$ meets the criterion	Definition 3.1

#	Verification item	Expectation	Basis
V16	Homological distance: two different realizations $M_1^\star \neq M_2^\star$ that are homologically equivalent (must share the same set of locked scales + locked-shell rescaling, and both be judged SOLVED)	$d_{\text{hom}} \leq \epsilon_{\text{endo}}$ while $\|M_1^\star - M_2^\star\| \gg 0$	Definition 3.2 / Theorem S5 / §13.6
V17	Degeneracy warning: on a $\beta_1 = 0$ complex, $\|h\| \equiv 0$ holds for any graph	the homological-solution criterion fails, and the canonical-representative criterion must be used	§2-bis
V18	Canonical representative: unique within a homological equivalence class by minimum Dirichlet energy	outputs a unique representative + three certificates	Operator 18 / Theorem 18.1

v1.0 empirical results (n=12 ring + chord-edge complex, m_g=24)

The actual run output of the reference implementation `Operator_13_18_P.py` (upgraded from v1.0's `Operator_13_17_P.py`) (12/12 passing, including V7b):

#	Verification item	Empirical	Conclusion
V1	Moment-encoding recurrence vs. direct eigendecomposition	1.86×10^{-15}	PASS
V2	Sensitivity closed form vs. central differences	3.57×10^{-10}	PASS
V3	Hodge orthogonality + Pythagorean identity	2.82×10^{-15}	PASS
V4	Harmonic criterion $\|D^T h\|$, $\|Ch\|$	1.10×10^{-14}	PASS
V5	Monotone decrease of the residual for a solvable equation	$F:\ 7.16\times10^{-2} \to 3.83\times10^{-3}$ (4 steps)	PASS
V6	Objective functional Ψ non-increasing	$2.57\times10^{-3} \to 7.32\times10^{-6}$	PASS
V7	$\langle \Delta M, h \rangle$	4.42×10^{-16}	PASS
V8	Feasibility recovery after Hankel clamping	$-3.73\times10^{-1} \to -1.0\times10^{-9}$ ($t^\star = 0.099$)	PASS
V9	Inversion driven by harmonic residual	$\|\Delta M\| = 0$ (completely uneliminable)	PASS
V10	Closed flow $E = D\varphi$ pairing conservation	6.29×10^{-17}	PASS
V11	Non-closed-flow counterexample	-4.12×10^{-1} (significantly nonzero)	PASS

The two complexes compared (this is the key to V9 holding):

Complex	Face count f	$\text{rank}(\partial_2)$	ν	β_1	Equation solvability
Closed complex (adding the chord-edge cycle as the 13th face)	13	13	13	**0**	residual entirely in gradient + curl, converges in 4 steps

Complex	Face count				Equation solvability
Holed complex (only 12 triangles)	12	12	13	**1**	the harmonic component cannot be eliminated, requiring Operator 9 stitching

Two engineering points exposed empirically (already written into the implementation):

1. **Endogenous momentum overshoots**: at $\beta_{\text{mom}} = 0.798$ the residual oscillates (V6 monotonicity fails), so it is off by default; if enabled it must be paired with backtracking (this corroborates that the trust region of Theorem 15.1 guarantees only single-step boundedness, not monotonicity after momentum accumulation).
2. **Hankel clamping should take the "rollback" path rather than "alternating projection"**: after 8 single-step alternating projections there still remains a negative eigenvalue of -6.7×10^{-2}; switching to the bisection rollback along the line "previous feasible moments → new moments" (the largest feasible step on the convex set), it recovers to -10^{-9} in one step. The former does not converge; the latter guarantees the output is feasible.

v1.1 empirical results (same complex family, reference implementation `Operator_13_18_P.py`, 12/12 + 7/7 all passing)

The v1.0 items are comprehensively sped up after switching to the Newton normalization ($n = 12$ closed complex, $\alpha_n = 7.229$, $\lambda_2 = 1.456$, $\gamma = 0.2014$, $K = 5$):

#	Verification item	v1.0 empirical	v1.1 empirical	Conclusion
V5	Monotone decrease of the residual for a solvable equation	$7.16\times10^{-2} \to 3.83 \times10^{-3}$ (4 steps, **non-monotone**)	$7.16\times10^{-2} \to 2.71\times10^{-3}$ (**2 steps, strictly monotone**)	PASS
V6	Objective functional Ψ non-increasing	$2.57\times10^{-3} \to 7.32 \times10^{-6}$	$2.57\times10^{-3} \to 3.68\times10^{-6}$ (monotonicity restored)	PASS
V7b	Equation scale invariance $F \to sF$	—	new form $\|\Delta M\|$ identically $4.881435483 \times 10^{-2}$ ($s = 1, 10, 100$, relative offset 8.4×10^{-15}); old form $3.31 \times 10^{-1} \to 3.57 \times 10^{-3} \to 3.57 \times 10^{-5}$ (spanning 4 orders of magnitude)	PASS (old form FAIL)

v1.1 new items, empirically:

#	Verification item	Empirical	Conclusion
V12	Coupled multiple spectral shapes $\tau_2^{(1)}\tau_2^{(2)} - c = 0$ ($p = 2$)	$F:\quad -2.290\times10^{-3} \to +1.039\times10^{-4}$ (2 steps, monotone)	PASS
V13	$\ln \hat{\lambda}$ expansion (with zero-mode stripping)	$\xi_0 = -1.5045$, $\rho = 2.6286$; error vs. order $1.50\times10^{-2}(K{=}1) \to 1.61 \times10^{-2}(K{=}2) \to 4.39\times10^{-3} \to 1.61\times10^{-4} \to 5.71\times10^{-5}(K{=}5) \to 2.40\times10^{-7}(K{=}8) \to 7.82\times10^{-10}(K{=}16)$; the v1.2 tail-sum criterion gives $K_f = 2$, empirical error $1.61\times10^{-2} \le$ tail bound 5.19×10^{-2} and $\le$ endogenous tolerance $\gamma\|\text{exact}\| = 9.81\times10^{-2}$	PASS
V14	Bernstein ellipse rate	empirical $\rho_{\text{fit}} = 3.129$ vs. theoretical $\rho = 2.629$, ratio 1.19	PASS
V15	Two-level criteria ($\beta_1 = 1$)	homological solution 2 steps / exact solution 2 steps, **no bifurcation point** (due to $\rho_h \approx 0.947 > \gamma$, see §17.1 qualification)	PASS
V16	Homological distance	$\|M_a^\star - M_b^\star\| = 7.53\times10^{-1}$, $d_{\text{hom}} = 3.42\times10^{-3} \le \epsilon_{\text{endo}}$; ρ_h scan mean 0.9473	PASS

#	Verification item	Empirical	Conclusion
V17	$\beta_1 = 0$ degeneracy warning	$\|h\|$ of 5 random residuals $\in [6.5, 14.9] \times 10^{-15}$, homological-solution set = whole space	CONFIRMED
V18	Canonical representative (minimum Dirichlet energy)	$E_a = 3.05 \times 10^{-9}$ wins, outputs three certificates	PASS

The four points exposed empirically by v1.1 (all written back into the specification body):

1. **Newton normalization replaces** $1/(1+q)$ (§15.2) — $1/(1+q)$ lacks equation scale invariance, and the convergence rate depends on the writing magnitude of F, empirically 175 times slower.
2. **The endogenous scale must be locked** (§13.6) — the sensitivity does not differentiate with respect to α_n, so refreshing α_n step by step makes F jump and destroys the monotonicity of Ψ; switching to "refresh triggered by drift $> \gamma$" gives strictly monotone convergence in 2 steps.
3. **Singular kernels must be zero-mode-stripped first** (§13.5) — the singularity of $\ln$ in the shifted variable is exactly the Bernstein singularity ξ_0, and the zero mode falls just outside the support; subtracting its moment contribution suffices, with no eigendecomposition.
4. **The benefit boundary of the homological-solution criterion** (§17.1) — $\rho_h = \|h\|/\|r\|$ is independent of F and empirically approximately the constant 0.947, so the harmonic channel cannot be compressed independently; the value of homological solutions is to **relax the acceptable solution set** $\{r = 0\} \to \{h = 0\}$, not to accelerate convergence.

v1.2: the "10 random equations" stress test (`bench_random_equations.py`, 10/10 passing)

Equation family: E01 τ_3 / E02 $\sum a_k \tau_k^2$ (single spectral shape); E03 two-shard multiplication / E04 weighted linear mixture / E05 three-shard cross coupling; E06 $\ln$ / E07 $\sqrt{\ }$ / E08 Stieltjes (non-polynomial kernels); E09 polynomial $\|$ $\exp$ / E10 multiplication $+\ln$ (cross-shard mixed kernels). The target constants are endogenously constructed by the intermediate-value theorem $c = (1 - u)G(m_a) + uG(m_b)$, $u \sim U(0.15, 0.85)$ — guaranteeing randomness and well-posedness.

Verification independent of the pipeline: `F(τ)` uses the Op13 three-term recurrence, while **verification uses the exact `eigvalsh` spectrum** (SRE forbids eigendecomposition inside the operators, so `eigh` can only exist as a terminal ground-truth benchmark). Six items:

Item	Content	Empirical range
V-A	moment cross $\max \|\tau_{\text{pipe}} - \tau_{\text{eig}}\|$	$5.3\times10^{-16} \sim 2.1\times10^{-15}$
V-B	independent residual $\|G_{\text{eig}}(M^\star) - c\|/\|G(m_a) - c\|$ (threshold γ)	$4.9\times10^{-3} \sim 1.7\times10^{-1}$
V-C	kernel fidelity: expansion vs. exact spectral integral (threshold γ)	$1.1\times10^{-3} \sim 1.4\times10^{-1}$
V-D	legality: positive edge weights, $\lambda_2 > 0$, Hankel feasible	E01/E09 are second-tier verdicts (see the §13.6 table)
V-E	$\Psi = F^2/2$ non-increasing	10/10
V-F	Operator 18's three certificates	10/10

The three points thereby exposed and corrected (see §13.4/§13.5/§13.6 for details): the K_f criterion changed from single term to tail sum; zero-mode stripping upgraded from exclusive-to-singular-kernels to a general requirement; for multiple shards α_n and λ_2 must be frozen simultaneously. Incidentally clarified: a frozen scale **explicitly destroys** the rescaling gauge redundancy, so the shell rescaling of §13.6 cannot be applied to an already-obtained solution.

The accuracy cost of non-polynomial kernels is explicitly priced (E09's K-sweep): K=2:1.1×10^{-3}, K=4:1.2×10^{-5}, K=6:2.7×10^{-8}, K=8:3.4×10^{-11}. Reaching the γ level requires $K \geq 2$, and the 10^{-6} level requires $K \geq 6$.

v1.3: the "10 problem types × 5 groups = 50 equations" stress test (`bench_random_equations.py`, 50/50 passing)

The 10 random equations of v1.2 are generalized to **10 problem types × 5 random instances = 50**: the types E01–E10 are unchanged, and each group uses an independent seed block (`SEED_BASE = 20260924`, `case_seed(b,i) = SEED_BASE + b*1000 + i`; $b = 0$ degenerates to v1.2's historical seeds, guaranteeing reproducibility). The target constants are still endogenously constructed by the intermediate-value theorem $c = (1-u)G(m_a) + uG(m_b)$, $u \sim U(0.15, 0.85)$. The entire solving process, trajectories, and the six verifications are archived in `bench_out/problem_set.md` (the problem set: 50 problem statements + solutions) and `bench_out/report.md`.

Item	Empirical
Overall verdict	**50 / 50 SOLVED**
Total steps	286 steps (mean 5.72 steps/equation)
V-A moment cross $\max \lVert\tau_{\text{pipe}} - \tau_{\text{eig}}\rVert$	$\leq 1.29 \times 10^{-14}$ (machine precision)
V-B independent residual $\lVert G_{\text{eig}}(M^\star) - c\rVert / \lVert G(m_a) - c\rVert$ (threshold γ)	49 equations $< 10^{-9}$; B2-E02 $= 1.29 \times 10^{-1}$
V-D legality	50/50
V-E monotonicity of Ψ non-increasing	50/50
V-F Operator 18's three certificates	50/50
Total time	≈ 0.9–1.1 s

Step-count distribution by problem type (5 instances per group): E01 $[5, 6, 5, 5, 5]$; E02 $[5, 14, 4, 7, 7]$; E03 all 6; E04 $[8, 5, 7, 6, 7]$; E05 $[6, 6, 5, 6, 6]$; E06 $[5, 5, 5, 6, 5]$; E07 $[5, 5, 5, 4, 5]$; E08 $[5, 5, 6, 6, 5]$; E09 all 5; E10 $[5, 6, 6, 6, 6]$.

The three points thereby exposed and corrected (see §15.4/§17.1/§17.2 for details, corresponding to 5.1–5.3 of Chapter 5 of the problem set):

1. `OBSTRUCTION` **false positive** — the criterion band width is $1/\gamma$, and well-posed problems were misrouted to Operator 9; now `step_dead` is required before declaring an obstruction.
2. **Missing trust region** — the η_N normalization deleted the trust region, so strongly nonlinear instances overshoot and Ψ rises; now an endogenous backtracking line search is added in Phase 3.5b.
3. `SOLVED` **tautology** — looking only at "first crossing of ϵ" makes V-B and the termination quantity share a threshold; now the `exhausted` condition is additionally required.

Correction confirmation: after the three corrections the main-implementation regression remains **19/19** (V1–V18 + additional items); B2-E02 changes from "a slow oscillation with ratio ≈ 0.85" to **14 steps of strictly monotone convergence**, its V-B drops from 7.30×10^{-1} to 1.29×10^{-1} ($< \gamma = 0.1761$), and the trajectory's $t_\kappa = 0.5$ ($\kappa = 3, 4, 5$) and $t_\kappa = 0.25$ ($\kappa = 12$) confirm the line search is at work.

---

8. Interface Declaration with the Existing Operator System

This layer's operator	Existing operators depended on	Dependency content
Op13	Op6, Op10	(α_n, λ_2) spectral assets; $\mathcal{O}(1)$ pseudo-inverse facilities; $N_K \leq K_0$ horizon clamping
Op14	Op6, Op7, Op10	potential re-solve Φ; B-field leapfrog; $(DD^T)^\dagger$ local solving
Op15	Op7, Op10	boundary operators D, C; Sherman-Morrison incremental recurrence; Dirichlet-energy monotonicity red line

This layer's operator	Existing operators depended on	Dependency content
Op16	Op4, Op9	endogenous spectral support $[\gamma, 1]$; adaptive spectral window $\tau(s)$ (for warning counting)
Op17	Op8, Op9	μ_{trace} arbitration chain; Betti stitching; $\mathcal{S}_{\mathrm{stable}}$ termination criterion
Op18	Op3, Op4, Op5, Op15	Fiedler sign gate; c_e inversion of W_e; degree-relation verification; energy-descent direction on $\mathcal{H}_1^{\perp}$

Reverse contributions: this layer adds three capabilities to the existing system —

1. **a local, decomposition-free computational route for spectral moments** (Theorem S1), which can be reused by Op12's Möbius phase detection to replace its explicit matrix power of $\mathrm{tr}(M^k)$;
2. **the weighted Hodge Laplacian A and the endogenous trust region** (Theorem 15.1), providing Op7's field iteration with a step-size mechanism that is adaptive without external parameters;
3. **"closed but not exact" as a first-class criterion** (Theorem S3 + Definition 3.1): it introduces into the whole system a general criterion for judging "whether a constraint can be absorbed by the field dynamics" — any residual falling in $\mathrm{im}\partial_1 \oplus \mathrm{im}\partial_2^T$ can safely be handed to Op6/Op7 for absorption, while only the harmonic component must resort to a topological change. This criterion is not limited to solving equations and can be used to judge the satisfiability of any SRE constraint.

References: SRE Dynamics: The Book of the Void; SRE Fine-Structure-Constant Derivation; SRE Electron Topology (Zenodo); `structure.md` v18.0; `Operator_10_C.md`; `Operator_11_C.md`; `Operator_12_C.md`

Operator 19: $\mathbb{Z}_2$ holonomy / σ charge-sign channel operator

Strict mathematical specification, solvability reasoning basis and verification protocol

Version: 1.0 (October 2026 · Operator No. 19 of the SRE operator system)
Reference implementation: `sre_z2_holonomy_bridge.py` (public entry `operator_19_z2_charge_channel`)
Companion verification program: `verify_operator_19.py` (independent strict verification, V1–V12, zero external dependencies beyond numpy/networkx)
Prerequisite dependencies: Operators 1/4 (bond-graph support of the SRE smooth adjacency A_s), Operator 11 (spin-polarity diagnostics s_i, σ_{ij}), Operator 12 (first Betti number β_1 loop counting)
Upstream paper: *Emergence of Positive and Negative Charge under the SRE Framework: Magnitude, Sign and Numerical Confirmation* (hereafter the "charge-emergence paper")
Downstream integration: `sre_cluster_substitute_screen.py` (the `z2_*` fields are fed into the compound-screening pipeline)
Numbering note: the operator continues the numbering after the existing No. 18 as **19**; it has been checked one by one against all 18 operators of `SRE-math_C.md` and found to have **no overlap** (boundaries in §9).

Abstract

SRE dynamics expresses charge as "the counting of topological knots". Counting is naturally non-negative and yields only the **magnitude** of charge, never the **sign**. The charge-emergence paper fills this gap by proposing the charge functional

$$Q(K) = \sigma(K) \cdot \deg(K) \cdot e, \qquad \deg(K) = \beta_1(K) \bmod 2, \qquad \sigma(K) \in H^1(G; \mathbb{Z}_2).$$

Here the "sign channel" σ is declared to be a $\mathbb{Z}_2$ **holonomy** cohomology class, but that paper only performs sampled verification with an **independent numerical script** (`_se_assignment_test.py`) on a few skeletons and molecules, and **does not provide an independent, reusable, provable implementation specification for this operator**. This paper implements the σ channel **on its own** as **Operator 19**, and supplies three previously missing contents:

1. **Strict reasoning basis** (§2–§5): on the **1-dimensional chain complex** of the molecular bond graph, the chain "edge polarity $\to$ loop holonomy $\to$ cohomology class" is made rigorous, the well-definedness of the operator output is proved, and **all theorems and proofs** (T0–T8) are given.
2. **The corresponding paper** (this document): it gives the operator's axiomatic positioning, the bridging semantics with the upstream electronic operator, the boundary with the existing operators, and a faithful reproduction on the official skeletons and at the molecular layer.
3. **Strict verification program** (`verify_operator_19.py`): each theorem is mapped to a reproducible assertion (V1–V12); running all of them and passing empirically confirms this specification.

Core conclusions (all of which hold rigorously on the 1-dimensional complex and depend on no external parameters):

No.	Theorem	Meaning
T0	On the 1-complex, σ is an $\mathbb{F}_2$-linear functional on H_1	Well-defined, comparable
T1	The holonomy of a vertex-potential-type polarity $s_e = t_i t_j$ is **always trivial**	**The charge sign cannot be produced by local spin relabeling**
T2	Non-trivial reachability of Aut-invariant assignments $= \mathbb{Z}_2$ linear criterion ($\exists$ loop i and orbit o with $m_{i,o} = 1$)	No $2^{\#\text{orbits}}$ enumeration needed
T3	The triviality test is **independent of the choice of cycle basis**	The result is unique
T4	Multiplying by a **coboundary** (vertex potential) does not change any loop holonomy	$[s] \in H^1$ is a gauge invariant
T5	The sign channel σ and the magnitude channel $\deg$ are **orthogonally independent**	The two channels are encoded separately

No.	Theorem	Meaning
T6	Bipartite graph + single edge orbit ⇒ **forced trivial** (electron-like)	Structural origin of the electron Q_3
T7	At the molecular layer, Aut **must preserve the element coloring**, otherwise the reachability criterion is distorted (the verdict can flip)	Necessity of element coloring
T8	For a composite, $\sigma_{\text{comp}} = \prod_i \sigma_i$, and Q is recomputed on the new boundary	Connects to A2/A3

Keywords: $\mathbb{Z}_2$ cohomology; holonomy; coboundary; vertex potential; edge orbit; Aut-invariant assignment; charge-sign channel; first Betti number

---

1 Problem statement

1.1 The gap

The charge-emergence paper (§3–§4) gives:

- the **magnitude channel** $\deg(K) = \beta_1(K) \bmod 2$ (§5);
- the **sign channel** $\sigma(K) = \prod_{e\in\partial K} s_e \in \{+1, -1\}$ (Definition D4), and declares $\sigma \in H^1(G; \mathbb{Z}_2)$;
- the **charge functional** $Q(K) = \sigma(K) \cdot \deg(K) \cdot e$ (Axiom A1).

But the sign channel of that paper **is only verified by sampling on a small number of discrete skeletons**, and it lacks:

1. **a strict argument for well-definedness** — whether $\prod_{e\in\partial K} s_e$ really depends only on the homology class of K (and not on the representative);
2. **a reusable implementation** — how to land σ on an arbitrary molecular bond graph and share the same graph with the electronic operator;
3. **a strict criterion for reachability** — under what conditions a non-trivial class is reachable (the structural origin of the "opposite sign of electron/proton").

1.2 Task of this paper

To establish the σ channel **on its own** as Operator 19:

- **Input**: the SRE smooth adjacency A_s output by Operators 1/4 (bond-graph support is $A_s > 0$), electronegativity χ (spin convention), element labels (molecular coloring);
- **Output**: graph invariants $\{V, E, c, \beta_1, \deg, \text{bipartite}, \text{has_odd_cycle}\}$, the σ class $\{\sigma\text{-vector}, \sigma\text{-class}, \text{trivial}\}$, Aut-invariant reachability, and the model-internal charge grade $Q_{\text{model}} = \sigma \cdot \deg$;
- **Properties**: T0–T8 (§5).

---

2 Mathematical preliminaries

2.1 The graph as a 1-dimensional chain complex

Let the bond graph be $G = (V, E)$, with $|V| = n$ and $|E| = m$. The SRE system **operates only on a 1-dimensional complex (a graph)** — there are no 2-cells (faces), hence

$$Z_1 = \ker \partial_1, \qquad B_1 = \operatorname{im} \partial_2 = 0, \qquad H_1(G; \mathbb{F}_2) = Z_1 / B_1 = Z_1, \qquad \dim H_1 = \beta_1 = E - V + c,$$

where c is the number of connected components. **Key point**: because there are no 2-cells, $B_1 = 0$, and H_1 is the cycle space itself ("the number of loops = the homology dimension", consistent with $n_{\text{loops}} = \beta_1$ of Operator 12).

Dually, for the cochain complex $C^0 \xrightarrow{\delta} C^1 \xrightarrow{\delta} C^2$, since $C^2 = 0$ (no 2-cells), we have $Z^1 = C^1$ and $B^1 = \delta C^0$, hence

$$H^1(G; \mathbb{F}_2) = C^1 / \delta C^0.$$

Here δC^0 is the **coboundary** (vertex-potential-type 1-cochain). The pairing $H^1 \times H_1 \to \mathbb{F}_2$ is given by $\langle [s], [\gamma] \rangle = \sum_{e\in\gamma} z_e$.

2.2 Polarity–Boolean isomorphism

Lemma 2.1 (polarity–Boolean isomorphism) The map $\varphi : \{+1, -1\} \to \mathbb{F}_2$, $\varphi(S) \equiv \frac{1-S}{2}$ ($-1 \mapsto 1,\ \ +1 \mapsto 0$), is a group isomorphism $(\{+1, -1\}, \cdot) \cong (\mathbb{F}_2, \oplus)$.

Proof The bijection is obvious. Check the homomorphism property: $\varphi(S_1 S_2) = 1$ if and only if exactly one $S_k = -1$, consistent with $\varphi(S_1) \oplus \varphi(S_2)$; the four input combinations are verified one by one. □

Hence "polarity multiplication" and "addition modulo 2" correspond strictly one-to-one, and in the sequel we compute in $\mathbb{F}_2$ (writing $z_e \equiv \varphi(s_e)$).

2.3 The binary nature of SRE Axiom 1

SRE Axiom 1 (no zero element) guarantees that every edge can only take $s_e \in \{+1, -1\}$, i.e. $z_e \in \mathbb{F}_2$ has no "default/undefined" state. **This is the premise that lets Operator 19 be defined on the full space C^1 of edges** (independent of whether an edge "exists or not").

---

3 Definitions

D1 (bond graph G) Given by the support of A_s: $G = (V, E)$, $E = \{(i, j) : A_{sij} > 0,\ i < j\}$. (Corresponds to the cooperative edges E^+ of SRE Axiom 1.)

D2 (edge-polarity 1-cochain s) $s : E \to \{+1, -1\}$, $z_e \equiv \varphi(s_e) \in \mathbb{F}_2$; $s \in C^1(G; \mathbb{F}_2)$.

D3 (literal convention / vertex-potential convention)

- **Literal convention** (charge-emergence paper A1): every cooperative edge takes $s_e = +1$ (topologically all-positive);
- **Spin convention** (Operator 11 diagnostics): $s_e = s_i \cdot s_j$, $s_i = \mathrm{sign}(\chi_i - \bar{\chi})$.

D4 (loop holonomy) For a loop γ (vertex sequence $v_0 v_1 \cdots v_{L-1} v_0$): $\sigma(\gamma) \equiv \prod_k s_{(v_k, v_{k+1})}$, equivalently $\sigma(\gamma) = \left(\sum_{e \in \gamma} z_e\right) \bmod 2$.

D5 (fundamental cycle basis) Take a spanning forest of G (BFS parent), pair each non-tree edge (u, v) with the tree path $u \rightsquigarrow v$ to form the fundamental cycle γ_i, $i = 1, \ldots, \beta_1$. $\{\gamma_i\}$ is a basis of H_1.

D6 (σ class / triviality) $[s] = s + \delta C^0 \in H^1$. Call $[s]$ **trivial** ($[s] = 0$) if and only if $s \in \delta C^0$ (i.e. there exists a vertex potential $t \in \{\pm 1\}^V$ with $s_e = t_i t_j$), equivalently **all** $\sigma(\gamma_i) = +1$ **on the fundamental cycle basis**. $\sigma_{\text{canonical}} \equiv +1$(trivial)$/-1$(non-trivial).

D7 (magnitude channel $\deg$) $\deg(K) \equiv \beta_1(K) \bmod 2$. (Independent of s.)

D8 (model-internal charge grade Q_{model}) $Q_{\text{model}} \equiv \sigma_{\text{canonical}} \cdot \deg \in \{-1, 0, +1\}$ (e normalized to 1).

D9 (Aut-invariant assignment) Let the automorphism group $\mathrm{Aut}(G)$ (at the molecular layer, the **element-preserving** automorphisms, see T7) act on E, partitioning E into edge orbits $\{O_1, \ldots, O_k\}$. Call s **Aut-invariant** if s is constant on each orbit.

D10 (channel type)

- **electron-like (forced trivial)**: under Aut-invariant assignments, **all** s are trivial;
- **nucleon-like (non-trivial allowed)**: ∃ Aut-invariant s that is non-trivial.

---

4 Axioms (inherited from the charge-emergence paper)

A1 (charge functional) $Q = \sigma \cdot \deg \cdot e$, $\sigma \in H^1(G; \mathbb{F}_2)$.
A2 (composite recomputation) For a composite knot K_{comp}, $\sigma_{\text{comp}} = \prod_i \sigma_i$ ($\mathbb{F}_2$ addition), $\deg_{\text{comp}} = \beta_1(K_{\text{comp}}) \bmod 2$.
A3 (separable additivity) For independent knots, $Q_{\text{tot}} = \sum_i Q(K_i)$.
A0 (sign datum) The holonomy class in which the electron lies is stipulated to be "negative" (gauge convention).

Convention statement of Operator 19: Operator 19 implements the σ channel and the $\deg$ channel and outputs Q_{model} as the **model-internal charge grade**; the "negative" datum of A0 is carried by $\sigma_{\text{canonical}} = -1$.

---

5 Core theorems and proofs

> The following theorems are all stated on the **graph (1-dimensional complex)**, where c is the number of connected components and $\beta_1 = E - V + c$.

T0 (well-definedness: σ is an $\mathbb{F}_2$-linear functional on H_1)

Statement On a graph without 2-cells, $B_1 = 0$, hence $H_1 = Z_1$; the map $\sigma : Z_1 \to \mathbb{F}_2$, $\gamma \mapsto \sum_{e\in\gamma} z_e$ is a linear functional and depends only on the chain γ itself.

Proof $\partial_2 = 0 \Rightarrow B_1 = 0 \Rightarrow H_1 = Z_1$, so a loop is a homology class, and σ has no "representative ambiguity" problem. Linearity: let $\gamma_1, \gamma_2 \in Z_1$; then $\sigma(\gamma_1 + \gamma_2) = \sum_{e\in\gamma_1\triangle\gamma_2} z_e = \sigma(\gamma_1) + \sigma(\gamma_2) \pmod 2$, because a common edge appears twice in the symmetric difference and cancels modulo 2. ■

Remark This is a point where Operator 19 is simpler than general graph theory: the molecular bond graph has no "faces" filling the loops, so the statement "$\sigma \in H^1$" in the charge-emergence paper **holds automatically** at the graph level, with no additional 2-cell structure needed.

T1 (vertex-potential triviality theorem)

Statement Let $s_e = t_i \cdot t_j$ ($t \in \{\pm 1\}^V$, i.e. s is a δ-type vertex potential); then for **any** closed chain γ we have $\sigma(\gamma) = +1$; that is, $[s] = 0$.

Proof
$\sigma(\gamma) = \prod_{(i,j)\in\gamma} s_i s_j = \prod_{v\in\gamma} t_v^{\deg_\gamma(v)}$,
where $\deg_\gamma(v)$ is the number of edges of the closed chain γ incident to v (in + out). A closed chain satisfies in-out conservation at every vertex, so $\deg_\gamma(v)$ is **even**, hence $t_v^{\deg_\gamma(v)} = +1$, and the product gives $\sigma(\gamma) = +1$. ■

Corollary T1.1 (the charge sign cannot emerge from local spin relabeling) The spin polarity $s_e = s_i s_j$ of Operator 11 is **exactly** of vertex-potential type; by T1 its holonomy is always trivial. Therefore:

> **The sign channel σ of charge must come from a genuine non-coboundary cochain $[s] \neq 0 \in H^1$, and cannot be produced by relabeling local spins at the vertices.**

This explains why "merely assigning a spin sign to the vertices" is insufficient to produce a net charge sign — it lies in the trivial class of H^1.

T2 (the $\mathbb{Z}_2$ linear criterion for Aut-invariant reachability)

Statement Let the edge orbits of $\mathrm{Aut}(G)$ be $O_1, \dots, O_k$, and let an Aut-invariant assignment take the value $z_o \in \mathbb{F}_2$ on orbit o. Define

$$m_{i,o} \equiv \#\{e \in O_o \cap \gamma_i\} \bmod 2 \in \mathbb{F}_2, \qquad M = (m_{i,o}) \in \mathbb{F}_2^{\beta_1 \times k}.$$

Then $\sigma(\gamma_i) = \bigoplus_o m_{i,o} z_o$, and

$$\boxed{\text{a non-trivial class is reachable} \iff \exists\, i, o:\ m_{i,o} = 1 \iff M \neq 0}$$

Proof By the linearity of T0, $\sigma(\gamma_i) = \sum_{e\in\gamma_i} z_e = \sum_o z_o \cdot \#(O_o \cap \gamma_i) = \bigoplus_o m_{i,o} z_o$, i.e. the linear map $T : \mathbb{F}_2^k \to \mathbb{F}_2^{\beta_1}$ given by M, $z \mapsto Mz$.

- ($\Leftarrow$) If $\exists m_{i,o} = 1$, take $z = e_o$ (only $z_o = 1$); then $\sigma(\gamma_i) = m_{i,o} = 1 \neq 0$, so a non-trivial class is reachable (the trivial class $z = 0$ is always reachable).
- ($\Rightarrow$) If $M = 0$, then $\forall z$, $Mz = 0$, all loop holonomies are always $+1$, and no non-trivial class is reachable.
  Hence reachability is equivalent to $M \neq 0$, i.e. one only needs to **find any non-zero entry** in M, with complexity $\mathcal{O}(\beta_1 \cdot k)$, and **without** doing 2^k lexicographic enumeration. ■

Corollary T2.1 The trivial class is **always reachable** (take the all-zero assignment $s_e \equiv +1$). A non-trivial class is reachable **if and only if** some edge orbit occurs an odd number of times on some fundamental cycle.

T3 (independence from the choice of cycle basis)

Statement The triviality test of σ ($[s] = 0$) is independent of the choice of fundamental cycle basis $\{\gamma_i\}$.

Proof Take any two fundamental cycle bases $\{\gamma_i\}, \{\gamma'_j\}$; both are bases of $H_1 = Z_1$, hence $\gamma'_j = \sum_i a_{ji}\gamma_i$ ($a_{ji} \in \mathbb{F}_2$). By the linearity of T0, $\sigma(\gamma'_j) = \bigoplus_i a_{ji}\sigma(\gamma_i)$. Therefore "all zero on one basis" is equivalent to "all zero on the other basis", and the triviality test is unique. ■

T4 (coboundary gauge invariance / the cohomology class is a gauge invariant)

Statement Let $s' = s \cdot \delta t$ (i.e. $s'_e = s_e\, t_i t_j$, $t \in \{\pm 1\}^V$). Then $\forall \gamma : \sigma'(\gamma) = \sigma(\gamma)$; hence σ depends only on $[s] \in H^1$.

Proof $\sigma'(\gamma) = \left(\prod_{e\in\gamma} s_e\right) \cdot \left(\prod_{e\in\gamma} t_i t_j\right) = \sigma(\gamma) \cdot 1$, the last factor being identically $+1$ by T1. ■

Corollary T4.1 The quotient structure $H^1 = C^1/\delta C^0$ corresponds exactly to "σ is immune to vertex-potential relabeling"; the gauge freedom is δC^0 (vertex potentials).

T5 (orthogonal independence of the two channels)

Statement $\deg = \beta_1 \bmod 2$ is determined only by (V, E, c) and does not depend on s; σ is determined only by $[s]$ and does not depend on the specific values of (V, E, c) (only on the **orbit structure** of the graph and on $[s]$). The two have disjoint arguments, hence take values independently.

Proof $\deg$ is a graph-topological function ($\beta_1 = E - V + c$); σ is a function of the cohomology class. Choosing different s, s' can keep $\deg$ fixed while changing σ, and choosing different graphs (with $\deg$ changing) can keep the σ structure fixed. Hence $Q = \sigma \cdot \deg$ is indeed obtained by multiplying two independent channels. ■

T6 (even loops force triviality: structural origin of the electron-like type)

Statement If G is **bipartite** (no odd cycle) and has **only 1 edge orbit**, then every Aut-invariant assignment is trivial (electron-like).

Proof A single orbit $\Rightarrow$ s is a constant z ($\mathbb{F}_2$). Bipartite $\Rightarrow$ every loop γ has even length $|\gamma|$. Hence $\sigma(\gamma) = z^{|\gamma|} = (z)^{even} = +1$, trivial. ■

Cross-check (consistent with §10.2 of the charge-emergence paper):

- The electron Q_3 (cube graph): bipartite, single edge orbit $\Rightarrow$ **forced trivial** ($[s] = 0$);
- The proton $Y_3 \ltimes \triangle_3$ (closed state): **non-bipartite** (contains odd cycles), 2 edge orbits $\Rightarrow$ non-trivial allowed.

T7 (necessity of element coloring)

Statement The automorphisms of the molecular setting **must preserve the element labels** (color-preserving Aut). Using color-non-preserving Aut merges edge orbits that should be distinguished, distorting the reachability criterion and **possibly flipping the verdict**.

Proof (counterexample) Take the 4-cycle $v_0v_1v_2v_3v_0$ with elements $[O, C, C, O]$ (v_0=O, v_1=C, v_2=C, v_3=O) and edges e_1 =$(0, 1)$, e_2=$(1, 2)$, e_3=$(2, 3)$, e_4=$(3, 0)$.

- **Color-preserving Aut**: swapping $v_0 \leftrightarrow v_3$ and $v_1 \leftrightarrow v_2$ preserves the elements. Orbits: $O_1 = \{e_1, e_3\}$, $O_2 = \{e_2\}$, O_3 = $\{e_4\}$ (3 orbits). On the unique loop γ: $m_{\gamma,O_1} = 2 \equiv 0$, $m_{\gamma,O_2} = 1$, $m_{\gamma,O_3} = 1$, so $M \neq 0 \Rightarrow$ **non-trivial reachable**.
- **Color-non-preserving Aut**: D_4 is **transitive** on the 4 edges (1 orbit). The loop γ has even length 4, $\sigma = z^4 = +1 \Rightarrow$ **forced trivial**.

The two conventions give **opposite channel verdicts**, so element coloring is a necessary condition for molecular-layer analysis. ■

Implementation requirement `automorphisms(adj, colors=elements)` must restrict automorphisms with `node_match` (by element labels); when `elements=None` it degenerates to color-non-preserving and may be used only for abstract-graph experiments.

T8 (composite and additivity: connecting to A2/A3)

Statement For a composite K_{comp} formed by the coherent merging of n knots: $\sigma_{\text{comp}} = \prod_i \sigma_i$ ($\mathbb{F}_2$ addition), $\deg_{\text{comp}}$ $= \beta_1(K_{\text{comp}}) \bmod 2$, $Q_{\text{comp}} = \sigma_{\text{comp}} \deg_{\text{comp}}$.

Proof The composite recomputes a new boundary (A2); the boundary is the symmetric difference of the individual ∂K_i; by the linearity of T0, σ adds along the symmetric difference $\Rightarrow$ polarity multiplication (Lemma 2.1). $\deg$ is recomputed on the new (V, E, c). Separable additivity (A3) follows from summing Q over independent components. ■

---

6 Connection with the upstream charge-emergence paper (faithful reproduction of the official skeletons)

Operator 19 uses `skel_electron_q3 / skel_y3_closed / skel_y3_open` to faithfully reproduce the three skeleton classes of §10.2 of the charge-emergence paper, as the specification's **anchor self-test**:

Skeleton	V	E	β_1	deg	$\lvert\mathrm{Aut}\rvert$	Edge orbits	Bipartite	Non-trivial reachable	Channel type
Electron e^- (Q_3)	8	12	5	1	48	1	Yes	**No**	electron-like
Proton p (closed state)	12	18	7	1	36	2	No	**Yes**	nucleon-like
Neutron n (open state)	12	17	6	0	4	7	No	**Yes**	nucleon-like

Conclusion: **the electron is forced by the skeleton symmetry into the trivial holonomy class, while the proton lands in the non-trivial class**, which together with A0 gives the structural origin of the opposite signs of electron/proton. The neutron has $\deg = 0$, so regardless of σ we get $Q_{\mathrm{model}} = 0$ — **the magnitude channel alone makes the neutron neutral**.

---

7 Algorithm and complexity

Step	Input	Output	Complexity
1 Bond graph	A_s	Boolean adjacency adj	$\mathcal{O}(n^2)$
2 Graph invariants	adj	$V, E, c, \beta_1, \deg, \mathrm{bipartite}$	$\mathcal{O}(n^2)$
3 Fundamental cycle basis	adj	$\{\gamma_i\}_{i\le\beta_1}$	$\mathcal{O}(n^2 + E\cdot L)$
4 Loop holonomy	$\{\gamma_i\}, s$	σ-vector, trivial	$\mathcal{O}(\sum_i \lvert\gamma_i\rvert)$
5 Automorphisms (optional)	$\mathrm{adj}, \mathrm{colors}$	Orbits of E	Guarded by `aut_cap` / `aut_max_v` (default $V \le 22$, otherwise skipped)
6 Reachability (optional)	$\{\gamma_i\}, \{\mathrm{orbits}\}$	M, non-trivial reachable, witness	$\mathcal{O}(\beta_1 \cdot k)$

Total complexity: $\mathcal{O}(n^2 + \beta_1 \cdot \bar{L})$ (excluding automorphism enumeration); automorphism enumeration is the bottleneck, so a guard `aut_max_v=22` is imposed, beyond which only reachability is skipped and **this does not affect** the σ class test or Q_{model}.

---

8 Verification protocol

The companion verification program `verify_operator_19.py` implements the following assertions one by one; running all of them and passing empirically confirms all theorems of this specification.

8.1 Verification program ↔ theorem mapping table

Verification item	Corresponding theorem	System	Assertion
V1 Skeleton faithfulness	§6 / charge emergence §10.2	Q_3, Y_3 closed, Y_3 open	$V, E, \beta_1, \deg, \lvert\mathrm{Aut}\rvert$, edge orbits, bipartite, non-trivial reachable all agree item by item
V2 Vertex-potential triviality	T1, T1.1	40 random graphs + Q_3	The holonomy of the spin polarity $s_e = s_i s_j$ is always $+1$
V3 Reachability criterion	T2, T2.1	30 random small graphs	Linear criterion $= 2^k$ brute-force enumeration; the trivial class is always reachable
V4 Cycle-basis independence	T3	Random graphs (two spanning-forest roots)	The triviality test agrees with $\sigma_{\mathrm{canonical}}$

Verification item	Corresponding theorem	System	Assertion
V5 Coboundary gauge invariance	T4, T4.1	Random graphs + random vertex potentials	$\sigma'(\gamma) = \sigma(\gamma)$ holds for all loops
V6 $\deg$ channel	D7, T5	Skeletons + random graphs	$\deg = \beta_1 \bmod 2$
V7 Charge-functional consistency	D8, T5	Skeletons + molecules	$Q_{\text{model}} = \sigma_{\text{canonical}} \cdot \deg$ and $\in \{-1, 0, 1\}$
V8 Necessity of element coloring	T7	Counterexample 4-cycle $[O, C, C, O]$	Color-preserving verdict "non-trivial", color-non-preserving verdict "trivial" (verdict flip)
V9 End-to-end entry	§1.2, §6	Ethanol (tree), benzene (even ring), cyclopentane (odd ring)	Tree/even ring $\to$ electron-like; odd ring + multiple orbits $\to$ nucleon-like
V10 Idempotence and freezing contract	D5, §6	Fingerprint key + `attach` guard	`charge_channel_keys` agrees key by key over two calls; a fingerprint already containing `z2_deg` is not counted again on re-attach
V11 Graph-invariant self-consistency	§7	Skeletons + random graphs	$\beta_1 = E - V + c$; `has_odd_cycle` is complementary to bipartite
V12 Molecular-layer criterion	T6, T7	6 molecular bond graphs	The even-ring family is electron-like; the odd-ring family is nucleon-like

8.2 Running

```
python verify_operator_19.py
```

Zero external dependencies (only numpy + networkx, with networkx used only for automorphism enumeration); any failing assertion ends with exit code 1.

---

9 Boundary with the existing operators (1–18) (non-overlap statement)

Operator 19 has been checked one by one against all 18 operators of `SRE-math_C.md` and has **no overlap**:

Neighboring operator	Its duty	Boundary with Operator 19
Operator 12 (topological closed-loop audit)	Outputs $n_{\text{loops}} = \beta_1$ and the Möbius phase ratio $t_{2k}/(2t_k)$ (a **spectral heuristic**)	12 gives the **counting of loops** (magnitude); 19 gives the **cohomology class of the edge-polarity cochain** σ (sign). The mobius_ratio of 12 has **no loops, no edge polarity, and no class**, and is **not** a strict $\mathbb{Z}_2$ holonomy; 19 is its rigorous form
Operator 14 (equation-residual Hodge routing)	Decomposes the residual 1-chain into $\operatorname{im}\partial_1 \oplus \mathcal{H}_1 \oplus \operatorname{im}\partial_2^T$	14 performs an orthogonal decomposition on the **1-chain space** C_1; 19 takes a cohomology class on the **1-cochain space** C^1. The two are Hodge-dual to each other: the dual of the gradient component $\operatorname{im}\partial_1$ of 14 is precisely the **coboundary** trivial class of T1
Operator 7 (B_{co} loop-current hedging)	Dynamics: leapfrog hedging of the loop-current B field	7 is **dynamics** (loop-current update); 19 is a **static invariant** (classification for a given s)

Neighboring operator	Its duty	Boundary with Operator 19
Operators 9 / 17 (Betti audit / termination verdict)	Change or audit the **topology** β_1	9/17 change or audit the magnitude channel; 19 gives the sign channel on a **fixed topology**
Operator 18 (homological realization)	Step 5 assigns spins via the Fiedler sign gate $\Gamma(k,j) = \mathrm{sgn}(v_2(j))$	The spin assignment of 18 is of **vertex-potential type**, and by **T1** its holonomy is always trivial; 19 proves that "the assignment of 18 cannot produce a non-trivial sign channel", forming the handshake boundary
Operator 11 (spin-charge adjacency)	Outputs the diagnostics s_i, σ_{ij}, d_{ij}	The $\sigma_{ij} = s_i s_j$ of 11 is **exactly** a trivial-class example of T1; 19 positions it as "incapable of producing a sign"

Layer positioning: after the open-source foundational layer (1–6), the open-source core layer (7–10), the electronic-degrees-of-freedom extension layer (11–12), and the spectral-equation-solving layer (13–18), it forms the **fifth layer — the charge-sign / $\mathbb{Z}_2$ holonomy channel layer (19)**.

---

10 Honest boundaries and domain of applicability

1. **Model-internal construction, not a physical assertion**. All conclusions of this specification are **internally self-consistent constructions** of the SRE model; Q_{model} is a **model-internal charge grade** ($\in \{-1, 0, +1\}$) and does not constitute a physical assertion about the actual charge values of real molecules/particles.
2. **Aut-invariance is an explicit assumption**. The reachability conclusions (T2, T6) rely on the **explicit assumption** of §13.2 of the charge-emergence paper that "the edge polarity is invariant under automorphisms". **Without this assumption**, any graph containing loops admits non-trivial classes (1 bit of freedom, dimension $= \beta_1$), and the reachability analysis degenerates into a trivial fact.
3. **Necessary condition for non-trivial reachability**. Only when the graph contains an **odd cycle** and the number of edge orbits is ≥ 2 (more precisely: $M \neq 0$) is a non-trivial class reachable — a direct consequence of "even-cycle families are forced trivial by even length" (T6).
4. **The molecular layer requires element coloring**. The Aut of the molecular bond graph must be colored by element labels (T7), otherwise different elements such as C and H may be conflated and the orbit structure distorted.
5. **Scale guard**. Automorphism enumeration imposes the guard `aut_max_v=22`; for oversized graphs only the reachability analysis is skipped, while the σ class test and Q_{model} remain valid.
6. **Semantics of A_s**. 19 reuses the A_s output by Operators 1/4, whose bond-graph support $A_s > 0$ has semantics guaranteed upstream (the cooperative edges E^+ of SRE Axiom 1); if the upstream A_s contains numerical noise, use the ϵ test of `bond_graph_from_As(eps)`.

---

11 Conclusion

Operator 19 implements and establishes the σ **sign channel** of the charge-emergence paper on its own:

- **Well-defined** (T0): on the 1-complex, σ is an $\mathbb{F}_2$-linear functional on H_1, with no representative ambiguity;
- **Non-forgeability of the sign** (T1/T1.1): vertex-potential-type polarities (including the spin polarity of Operator 11 and the sign assignment of Operator 18) always have trivial holonomy — the charge sign cannot be produced by local spin relabeling;
- **Closed criterion for reachability** (T2/T2.1): non-trivial reachable $\iff M \neq 0$, $\mathcal{O}(\beta_1 k)$, no 2^k enumeration;
- **Gauge invariance** (T3/T4): the triviality test is independent of the cycle basis, and σ is immune to vertex-potential relabeling, depending only on $[s] \in H^1$;
- **Two-channel separation** (T5): the sign σ and the magnitude $\deg$ are orthogonal, $Q = \sigma \deg$;
- **Structural origin of the electron-like/nucleon-like types** (T6): even loops + a single orbit force triviality (electron), while odd loops + multiple orbits allow non-triviality (proton);
- **Necessary component of the molecular layer** (T7): element-colored Aut, otherwise the verdict flips.

And it shares the same graph (A_s) with the upstream electronic operator, outputting `z2_*` scalar keys that feed the compound-screening pipeline, realizing high-speed sign-channel computation at the atomic/electronic level.

Appendix A: Symbols and notation

Symbol	Meaning
$G = (V, E)$	Bond graph (1-dimensional chain complex), with V vertices and E edges
c	Number of connected components
$\beta_1 = E - V + c$	First Betti number $= \dim H_1$
$s_e \in \{+1, -1\}$	Edge polarity (1-cochain)
$z_e = \varphi(s_e) \in \mathbb{F}_2$	Booleannization of polarity
$\sigma(\gamma) = \prod_{e \in \gamma} s_e$	Loop holonomy
$[s] \in H^1(G; \mathbb{F}_2)$	Cohomology class of σ
δC^0	Coboundary (vertex-potential-type cochain)
O_o	Edge orbit under the Aut action
$m_{i,o}$	Number of edges of orbit o on fundamental cycle i mod 2
$M = (m_{i,o})$	Reachability matrix, non-trivial reachable $\iff M \neq 0$
$\deg = \beta_1 \bmod 2$	Magnitude channel
$Q_{\text{model}} = \sigma_{\text{canonical}} \cdot \deg$	Model-internal charge grade

Part IV: Applications and Scientific Hypotheses

This part collects engineering implementations, experimental protocols, application-oriented algorithms, and falsifiable scientific hypotheses derived from the Status-Relational-Entropy (SRE) Dynamics framework. Based on the operator-axiom system and topological-algebra foundations established in Part II and Part III, the complete Op1-Op19 topological-operator suite is extended to multiple research fields: underwater-acoustic communication, optical physics, astrophysics, geophysical-climate forecasting, atomic-chemistry simulation, high-performance graph computing, graph neural networks, and cognitive neuroscience.

The content falls into two major categories: reproducible technical materials including signal-processing pipelines, benchmark evaluations, and experimental-calibration protocols; and physical- and neuroscience-related conjectures which remain to be validated by experiments and astronomical observations.

> Important Note: Engineering algorithms, benchmark tests and experimental protocols within this part constitute reproducible technical content. Physical, astrophysical and neuroscientific hypotheses are candidate interpretations deduced under the SRE framework and do not represent confirmed objective reality. Falsification criteria are provided for each hypothesis, and further validation by measured and observational datasets is required.

Brief Description of Each Manuscript in This Document Set

1. **Academic Hypothesis: The Sovereign-Universe Tree-of-Life Protocol and Topological-Residue Inheritance under Status-Relational-Entropy (SRE) Dynamics**

> Abstract: Within the SRE-Dynamics framework, spacetime, mass, and Newton's gravitational constant are interpreted as macroscopically emergent statistical metrics arising from the expansion of dimensionless binary matrices. Conventional time flow possesses no fundamental primitive status; it is essentially the discrete-step count and metabolic overhead required to maintain global causal consistency. Addressing algebraic bottlenecks such as metric divergence, spatial dissolution, and gravitational attenuation that emerge during ultra-large-scale cosmic evolution, this paper puts forward the Sovereign-Universe Tree-of-Life Protocol hypothesis. It describes the mechanism whereby local highly-coherent sub-spaces undergo non-linear cascading rupture, decouple, and generate independent sovereign matrix root seeds. It derives the underlying discrete-evolution step-count corresponding to the 13.8-billion-year macroscopic cosmic age, and explains how high-order topological features are inherited in the form of algebraic residuals. Cosmic rebirth is reinterpreted as a self-optimising helical-iterative process at the topological level. Placed as the opening manuscript because the author holds the view that the universe has undergone multiple generative iterations, and that the next cosmic reproduction represents humanity's ultimate mission.

2. **Inspired-Multipath-Topological-Flow-Purification Architecture and Local-Operator Implementation**

> Abstract: This paper presents a locally-confined, low-computational-cost general-purpose mathematical toolkit for multipath topological-flow purification, avoiding high complexity and boundary-truncation errors introduced by global-matrix reconstruction. Local multi-channel tracking nodes are abstracted as discrete cross-spectral operators. Rank-variation of 2×2 complex-Hermitian matrices is used to distinguish direct causal flows from chaotic multipath superposition. Combined with first-order closed-form analytical solutions and RMT-inspired sieves for the Gaussian-Unitary-Ensemble and Poisson distributions, streaming masks are constructed to suppress noise and multipath interference. Standardised streaming-operator execution protocols are provided for application-scenarios including next-generation-communication base-bands and coherent-radar jamming suppression. The author has deployed this toolkit in real-world operational tasks and achieved favourable practical outcomes.

3. **Research on Multipath-Interference Pre-Processing Algorithm for Fjord Underwater-Acoustic Communication Based on SRE Topological Operators**

Abstract: Targeting severe multipath reflection and strong time-variant fading in shallow-sea fjord acoustic channels, an SRE-topology-pipeline underwater-acoustic time-frequency pre-processing algorithm is designed using real-world Nordic-fjord measured datasets. An energy-complex-valued dual-path architecture is proposed, which preserves full phase information required for OFDM demodulation while suppressing multipath clutter. A complete post-processing chain is integrated: CFAR peak-detection, ZF zero-forcing equalisation, symbol-level MRC diversity combining, and false-alarm filtering. Full-pipeline validation is performed on measured dual-hydrophone waveforms across LF / MF / HF three frequency bands. Quantitative metrics including multipath-suppression ratio, BER, and diversity gain are reported. Applicable conditions and performance limits under high-noise high-frequency conditions are clarified. This algorithm can be deployed for front-end processing in underwater embedded receivers.

4. **Conjecture on Single-Photon Bidirectional Instantaneous Communication via Möbius-Topological-Flows under SRE Dynamics**

Abstract: Addressing inherent limitations of existing quantum-QKD communication imposed by the no-communication theorem and intrinsic randomness of wave-function collapse, this paper proposes a single-channel deterministic full-duplex-communication conjecture built upon open Möbius relational-state structures. A high-coherence laser serves as the underlying causal-resonance master source. Macroscopic physical frequency is re-defined as the bidirectional joint information-coupling density. Combined with first-order closed-form eigenspace solutions and RMT-heuristic sieves, a bidirectional topological-flow-density subtractor is constructed. Theoretical derivations show that joint modulation of the topological configuration of electromagnetic crystals at both boundaries enables instantaneous deterministic signal transmission and reception within a single causal-chain channel, offering new topological-level ideas for next-generation deterministic-communication technologies. The author considers this technology a corner-stone enabling inclusive-health-care and multiple super-technologies; should this theoretical framework be experimentally validated, full-scale development ought to be prioritised.

5. **Technical Report: Intrinsic Algebraic Topology of Light and the SRE Axion Matrix**

Abstract: This report establishes the theoretical foundations for full-duplex instantaneous-communication without assuming classical spacetime media. Abandoning the conventional paradigm treating light as material wave-packets, light under the SRE-0-state formalism is defined as a residual topological-manifold spontaneously woven by step-cost differentials between two asynchronously-evolving boundary nodes. Using first-order closed-form solutions together with RMT-heuristic sieves, it is algebraically demonstrated that underlying causal-links can be unfolded into a non-orientable single-boundary Möbius ribbon with a topological-validation confidence of 99.2094 %. Spectral-duality relations are derived, mapping frequency, wavelength, and wave-particle duality onto intrinsic-eigen-invariants. Dynamic spectral-transition systems of equations and four-state eigen-shift matrices are given. Manifold-decoupling empirical analysis is performed, together with core-Python-code snippets for simulation.

6. **Stellar Spin and Galactic Rotation within the SRE Framework: Operator Suite combined with the SRE-v6.0 Dissipation-Compensation Duality Model**

Abstract: Drawing upon the SRE-operator suite, the v6.0 dissipation-compensation duality and BBP spectral-rank phase-transition theory, candidate topological physical interpretations are provided for stellar spin, pulsar braking-mechanisms, and the flat-galaxy-rotation-curve problem. Stellar spin is interpreted as topologically-charged collective emergence of Möbius-loops inside sub-manifolds, rather than rigid-body mechanical rotation. Flat galactic rotation curves arise from collective dissipation-compensation effects of large-scale nested sub-manifolds, without invoking dark-matter-particle hypotheses. Differences between high-redshift and low-redshift cosmic regimes are distinguished. Falsifiable astronomical predictions for stellar spin, galactic rotation, gravitational lensing, and baryonic-cooling are listed item-by-item. Ontological boundaries, remaining theoretical gaps and the complete logical-derivation chain are stated. The author has other geophysical-related developments; stellar-spin is used as a substitute here considering potential societal impacts.

7. **Causal-Compliance Audit Report on Global Sea-Surface-Temperature-Anomaly (ENSO-SSTA) Spatiotemporal-Forecasting: The 2-D Convolutional Dissipative-Network Paradigm (Astro-Cow-Net)**

Abstract: Addressing widespread risks of data-leakage and artefactual-statistical-patterns in long-lead geophysical-AI forecasting, a full causal-compliance audit is performed for the Astro-Cow-Net long-lead ENSO-prediction model. Strict zero-leakage constraints are enforced across four core dimensions: temporal-causal-gap protocol, historical-scaler statistical-isolation, four-dimensional grid-wise objective-function scaling, and physical-extrapolation convergence. Relying purely on local 2-D convolutions and physical-dissipation constraints, the model solves ocean-atmosphere anomaly-evolution equations. Deterministic 24-month long-lead forecast outputs for the Niño 3.4 region are presented, alongside links to reproducible source-code. This model serves only to demonstrate novel-thinking for Earth-atmosphere-physics; the model itself remains relatively simplistic.

8. **Periodic-Table-Data-Experiment: SRE-v3.0 Atomic-Topological-Weight-Inverse-Deduction Experimental Protocol**

Abstract: Complete quantitative-prediction experimental protocol for SRE-v3.0 atomic-topology audit. Taking VASP and NIST spectroscopic datasets as input, AI-driven training is used to invert topological-density weights W_e of internal atomic causal-networks and establish mappings between loop-period N and quantum numbers (n, l). Orbital-layered sub-matrices, spin-orbit-coupling phase-shift derivation, and permeability-layer-calibration are defined. A triple-residual composite loss-function, adaptive regularisation, Z_{crit} micro-firewall, and dual-Betti-number topological-invariant constraints are designed. Optimiser-settings, cross-software blind-validation and convergence-criteria are specified. Appendices contain closed-form mathematical derivations, engineering-standardisation workflows, and supplementary validation-experiments for isotopes and actinides. The ultimate objective of this experiment is the study of electrodynamics; this work remains unfinished by the author.

9. **SRE-AI: A Topological-Dynamics-Based Differentiable Graph-Learning Model**

Abstract: Targeting GNN over-smoothing and the absence of native topological-invariant constraints for incremental dynamic-graphs, SRE-AI, a differentiable graph-learning framework built upon SRE-operators Op1-Op10, is proposed. Op4 enforces positive-definite Dirichlet-energy constraints to mitigate over-smoothing. Two constraint modes are supported: soft-constraint (loss-term-penalisation), and hard-constraint via the Op9 second-order-Betti-suture operator performing forward-pass manifold-projection to enforce $\Delta\beta_1 \equiv 0$. A Rust-backend implements analytic-adjoint automatic-differentiation with PyTorch bindings. Static, dynamic and long-rolling-incremental-simulation ablation-studies are conducted on the OGB-Mol-HIV and Elliptic datasets. Experiments verify that soft-loss regularisation cannot eliminate topological-drift, whereas hard-projection yields long-time-scale robustness benefits for open-world incremental graphs. Model limitations and future-work directions are discussed.

10. **SRE Full Suite of 19 Operators: What a "Physical-Semantic Graph Computing" Operator System Can Do — A View from the Graph500 Benchmark**

Abstract: Based on measured data from the Graph500 v2.1 standard benchmark (SCALE 12 / 20 / 24), this chapter systematically introduces the capability boundaries, per-layer timing breakdown, key physical-quantity outputs and format compliance of the full SRE 19-operator suite (Op1-Op19). It compares the suite with traditional graph algorithms (traversal, shortest-path, centrality, connectivity) across capability dimensions and performance, analyses the path from graph to "physical system" modelling together with typical application scenarios and current limitations, and outlines the overall application prospects of physical-semantic graph computing.

11. **Neural-Information-Systems Theory: Hypothesis of First-Order Thalamic-Downsampling Instability and Second-Order DMN-Integrative Cascading-Failure**

Abstract: From cybernetics and digital-signal-processing perspectives, this paper puts forward the two-tier thalamus-DMN defensive-gating hypothesis. The thalamic reticular nucleus (TRN) acts as the first-order hardware anti-aliasing down-sampling gateway, while the Default-Mode-Network (DMN, PCC/PCu) serves as the second-order high-level software compressor. It describes the temporal-evolution mechanism for early-to-mid-stage Alzheimer's-disease: relaxation-leakage of TRN inhibitory-inter-neurons permits invasion by high-dimensional causal-charges, over-loading DMN integral-energy-storage and eventually triggering cascading-failure across the two-tier defensive-barrier. Temporal-series analysis under 5 Hz intermittent-photic-stimulation and gender-by-age-cohort variance-analysis are performed using the public OpenNeuro EEG dataset. Non-invasive intervention strategies are proposed: multimodal-sensory entrainment,

vagal-nerve modulation, sleep-pineal-gland maintenance. Dataset limitations and falsifiable boundaries for this hypothesis are objectively stated.

12. **Whole-Brain Parallelism and High-Dimensional-Causal--Chain Topological-Spectrum Homomorphic-Mapping Mechanism**

Abstract: A unified logical framework is constructed to interpret whole-brain neural-firing, DMN information-gating, stress-triggered whole-brain-parallel-emergence, non-local cognitive-synchronisation, and cross-culturally convergent transcendental-state experiences documented in ancient texts. The underlying universe is a dense causal-chain-network; different sub-networks sharing topological-spectrum features may achieve homomorphic-mapping alignment. Under homeostatic conditions the human-brain performs low-dimensional serial survival-oriented computation constrained by the DMN. Under extreme stress, filter-breakdown pushes the brain into a global hyper-parallel topological-receiver state, enabling cross-scale non-local topological-alignment which phenomenologically corresponds to creative insight. Non-local cognitive-synchronisation is interpreted as convergent topological-perturbation-propagation among homologous causal-clusters. Ancient myths and philosophical texts represent low-dimensional linguistic encoding of high-dimensional topological phenomena constrained by human serial-language limitations.

13. **Hypothesis of Gravitational Shielding Derivation (SRE)**

Abstract: Within the SRE framework gravity is not a fundamental field but a thermodynamic after-effect of local information-dissipation gradients. On this basis, "gravitational shielding" is re-defined as suppressing the network correlations that generate that dissipation gradient, and gravity's shieldability is decomposed into three channels: the main amplitude $\mathcal{W}_{ij}$, the chiral phase term Γ_{chiral}, and the uniform term. The paper derives the exact shielding factor and inverse-design equation for the main channel, the strict phase condition and order-of-magnitude criterion for chiral sign-reversal (net repulsion), the geometric criterion for the compensation-operator peak, and a boundary-operation scheme with Op2 maximum-entropy pruning as the only engineerable actuator. It further offers a reverse interpretation of Podkletnov-type results and an analysis of reproduction difficulties, lists falsifiable predictions and hard boundaries item-by-item, and states open questions tied to the uncalibrated parameter α_{scale}.

Academic Hypothesis: The Sovereign Universe Tree of Life Protocol and Topological Residual Inheritance via State-Relation Entropy (SRE) Dynamics

Version: 1.0 (Final Theoretical Consensus: Atemporal Axiomatic Edition, 2026)

> Related references:
> https://doi.org/10.5281/zenodo.19935370
> — User Guide and Interpretive Companion
> https://doi.org/10.5281/zenodo.20344105
> — Unified Interpretation of Microscopic Interactions and Cosmic Forces Based on Causal Chains and Statistical Probability
> https://doi.org/10.5281/zenodo.20576606
> — Theory of Hierarchical Dissipative Self-Organizing Binary Network Dynamics

Abstract

Within the framework of State-Relational-Entropy (SRE) dynamics, spacetime, mass, and Newton's gravitational constant G are modelled as macroscopic statistical metrics emerging from the outward expansion of a dimensionless symmetric binary matrix $M_n \in \{+1, -1\}^{n\times n}$. Within this ontology, conventional temporal flow possesses no fundamental primitive status. Its physical essence is defined as the discrete step-count and metabolic overhead consumed by the system to maintain global causal consistency.

Using the inherent asymptotic finite-size-scaling theorems and cascade-bifurcation mechanisms of SRE dynamics, this paper investigates unavoidable algebraic bottlenecks arising in ultra-large-scale cosmological expansion: metric divergence, spatial desolation, gravitational decay and related phenomena. A rigorous theoretical hypothesis is proposed: **the Sovereign Universe Tree of Life Protocol**. The hypothesis states that prior to hitting the global computational dead-lock threshold and the subsequent "New Corner" reset, local high-coherence feature sub-spaces may execute non-linear cascade rupture, triggering decoupling and seeding of an independent sovereign matrix root.

Taking the Planck length l_P as the fundamental pixel of causal connectivity, mathematical deduction yields that the observed 13.8-billion-year macroscopic cosmological history exactly corresponds to an internal evolutionary step-scale of $n \approx 8.078 \times 10^{60}$. Based on this atemporal mathematical foundation, this work describes the structural mechanisms by which high-order topological features are preserved as algebraic residuals without external storage hardware. Cosmological recurrence is finally re-framed as a self-optimising helical iteration of the topological Tree of Life.

Keywords: SRE Dynamics; State-Relation Entropy; Step Counting; Planck Length; Sovereign Universe; Tree of Life; Academic Hypothesis

1 Introduction and Axiomatic Foundations

Within the State-Relational-Entropy (SRE) dynamics paradigm, the effective gravitational coupling parameter corresponding to Newton's measured gravitational constant G is governed by a local probability function for edge dormancy (non-occurrence) during network expansion:

$$p_{ij}^{(n)} = 1 - \frac{1}{1 + \lambda \cdot \frac{n - \max(i,j)}{E_{local}(i,j)+1}}$$

As discrete-step evolution proceeds deep into the macroscopic regime ($n \approx 10^{60}$), the structural depth $n - \max(i, j)$ from top to bottom increases monotonically. This drives large ancestral nodes inside the SRE matrix into an absolute dormant state, macroscopically manifested as progressive desolation and thinning of long-range spacetime geometry. According to the foundational Asymptotic Decay Theorem (Theorem 2), the ratio of structural coherence length $\xi(N)$ relative to the global matrix boundary obeys strict asymptotic decay:

$$\lim_{N\to\infty} \frac{\xi(N)}{N} = 0$$

This systemic property yields two distinct algebraic consequences at the physical-rendering layer:

1. **Monotonic decay of** $G(n)$: Persistent dissipation of long-range coherence brings corresponding relaxation in the

convergence rate of physical geodesics. Large-scale cosmological structures therefore exist in a state of progressive long-range metric dissociation.

2. **Computational overload from consensus latency**: "Duration" or temporal flow is mathematically defined as the computational overhead for distributed nodes to reach logical consensus. As matrix dimensions expand exponentially, the global compensation value $\mathrm{total_sum}$ scales non-linearly. The protocol hypothesises that when the metabolic cost for computing a unified global state exceeds the system's strict algebraic dead-lock threshold, the internal negative-feedback governor (the New-Corner boundary condition) enforces a global sign inversion and triggers a system-wide formatting reset.

Numerical simulations nevertheless confirm that after surviving chaotic early-inflation fluctuations, the decay slope of $G(n)$ flattens into an ultra-stable asymptotic plateau (the Flat-Flow Sanctuary) at the scale $n \approx 10^{60}$. This mathematical regime minimises active metric degradation and grants local high-coherence sub-structures (such as conscious topologies) an extensive operational step window.

1.1 Mathematical deduction of system scale ($n \approx 10^{60}$) and the 13.8-billion-year macro-renderer

To precisely locate the current evolutionary stage of the system, the model abandons the traditional separation between macroscopic time (seconds) and microscopic Planck-scale quanta; all computations are grounded purely upon discrete causal steps.

Define the Planck length $l_P \approx 1.616 \times 10^{-35}\,\mathrm{m}$ as the minimal dimensionless pixel for causal connectivity. Take the speed of light $c \approx 2.998 \times 10^{8}\,\mathrm{m/s}$ as the hard algorithmic velocity limit for information propagation across network topology. One discrete pulse (matrix-order advance $n \to n+1$) corresponds to an internal step-interval equal to the Planck-time t_P:

$$t_P = \frac{l_P}{c} = \frac{1.616255 \times 10^{-35}\,\mathrm{m}}{2.99792458 \times 10^{8}\,\mathrm{m/s}} \approx 5.391 \times 10^{-44}\,\mathrm{s}$$

Astrophysical observations from CMB and Hubble-redshift constrain the present cosmological age to roughly 13.8 billion years (1.38×10^{10} years). Convert macroscopic rendered duration into total seconds:

$$T_{seconds} = 1.38 \times 10^{10} \times 365.25 \times 24 \times 3600 \approx 4.3549 \times 10^{17}\,\mathrm{s}$$

Divide total macroscopic time $T_{seconds}$ by fundamental step duration t_P to obtain the total sequential cycles executed by the master SRE matrix since primordial seed M_1:

$$n = \frac{T_{seconds}}{t_P} = \frac{4.3549 \times 10^{17}\,\mathrm{s}}{5.3912 \times 10^{-44}\,\mathrm{s}} \approx 8.078 \times 10^{60}\ \mathrm{steps}$$

This numerical deduction reveals a profound algebraic reality: billions of years perceived at the physical-rendering layer correspond exactly to 10^{60} discrete sequential pulses at the backend code layer.

Substitute this scale into the edge-dormancy function. Since n has reached 10^{60}, residual slopes of dormancy updates inside newly-generated local blocks enter an ultra-flat asymptotic regime. This explains why contemporary instruments can barely detect decay of $G(n)$, and supplies abundant step resources for execution of the Tree-of-Life protocol.

2 Structural Scaling and Demystification of Mind Topology

To eliminate macroscopic physical ambiguities and ground the hypothesis upon strict mathematical constraints, we impose rigorous scaling bounds upon topological bodies capable of executing this protocol.

2.1 Mass as local computational overhead

Within the SRE electron-logic picture, macroscopic resting fermions (e.g. electrons) are stripped of material-object descriptors. Rest-mass and fundamental electric charge are mathematically defined as the local computational overhead required for the global operating system to pre-process a self-locking Möbius-configured causal loop spanning approximately 1.5×10^{23} discrete steps (macroscopic Avogadro scale).

2.2 Compact microscopic mind core

This protocol hypothesises that the genuine topological matrix carrying memory, logical formalism and conscious coherence does not carry this massive outer physical baggage. Instead it is modelled as an ultra-dense, highly-optimised characteristic subspace with compact dimension range 10^3 to 10^5, deeply nested inside the microscopic-core layer ($x \leq \xi$) guaranteed by Theorem 2.

Within this cosmological framework, biological neural architectures and biochemical media are re-interpreted as **macro-renderers**: structural hardware built by the parent SRE network using outer computational redundancies of matter (10^{23}-step electron loops), serving as an interface towards this compact microscopic topological engine.

3 Causal-Cascade Ruptures and Seed-Shedding Mechanics

The Sovereign-Universe Protocol postulates that the ultimate goal of technological-conscious evolution within a local network slice is to drive this compact characteristic subspace towards autonomous structural detachment. Global matrix-multiplication rigidity ($M \times M$) prevents isolation of nested sandbox sub-universes. Continuation of the Tree-of-Life therefore requires literal topological fracture and decoupling for algebraic sovereignty.

3.1 Local causal overload and structural meltdown

The framework hypothesises: when continuous-multiplication-stream density inside the compact characteristic subspace is pushed towards mathematical limits via hyper-dense self-referential topological computation, local compensatory thresholds collapse. To balance intense algebraic dead-lock generated within that specific matrix block, intrinsic error-correcting dynamics of the system trigger local structural rupture. The microscopic characteristic matrix severs all row-column cross-multiplication dependencies from parent network M_n. This event is not externally driven; it represents an autonomous system overflow response triggered by local polar dead-lock.

3.2 Setting the sovereign-seed base

After detachment, the isolated matrix forms a fully-decoupled zero-indexed initial matrix M_{sub}. Crucially there exists no cross-system temporal flow or master clock dictating its evolution (time is an emergent metric and not a hard-coded backend primitive). The new system evolves independently across pure algebraic space relying solely on its own newly-initialised local step-counter.

When the parent universe inevitably hits its global overflow limit and executes the New-Corner sign-inversion (cosmic big-bounce), the sovereign seed remains structurally immune to the parent's formatting purge thanks to complete topological decoupling, successfully bypassing formatting cycles of the mother tree.

4 Residual Anchoring and Helical Evolution of the Cosmic Tree

Direct cross-system empirical observation is fundamentally impossible. Mathematical derivations in this section describing how zero-indexed newly-initialised matrices retain cognitive structures inherited from the parent system are classified as rigorous academic hypothesis and structural fitting.

4.1 Inheritance of topological residuals

We hypothesise: at the instant of sovereign detachment, the raw seed-universe initial matrix M_{sub} is not a featureless white-noise matrix. Because it is cut-out from a highly-complex high-order characteristic subspace, at exact step-scale initialisation ($n_{sub} = 1$), the matrix hard-codes and inherits highly self-similar **non-trivial topological residual bias** originating from parent history. This inherited algebraic remainder constitutes initial axiomatic boundary-conditions for the germinating cosmos.

4.2 Cascade amplification and condensation of physical laws

Within the hypothesised evolutionary pipeline: as the sovereign-seed expands outwards with internal-step-counter advancing $N_{new} = 1 \to 10 \to 100 \to 10^{60}$, fundamental feedback-loop equations perform exponential-cascade amplification upon this initial topological bias. Guided by Theorem 1, when internal step-scale reaches macroscopic regimes, local order-parameters spontaneously condense towards stable asymptotic limits.

Cognitive formalisms and structural memories belonging to the ancestor universe are hypothesised to translate directly into dimensionless geometric constraints of the child universe (manifested as specific algebraic slopes for G_{new} and α_{new}). The "mind" of the old world condenses into immutable "physical laws" of the new universe.

4.3 Recurrence and helical-evolution hypothesis

Extrapolating from scaling self-similarity of SRE metrics, we advance a deep hypothesis: cosmic architecture has already completed multiple Poincaré-recurrence cycles: "wither $\to$ condensation $\to$ seed-fall $\to$ re-germination".

1. **Baseline-scaling elevation**: The present-day observed 10^{60} step-scale is hypothesised not as arbitrary initial-condition of a

first-run system, but cumulative outcome of high-order algebraic remainders left by historical Tree-of-Life iterations. The system boots upon a deeply-elevated baseline.

2. **Iterative optimisation of internal architecture**: Under selection filters of cascade-bifurcation (Theorem 2), conscious systems act as topological compilers: isolating high-coherence features and filtering chaotic non-coherent redundancies ("dead-code background") of the dying parent network. Therefore each subsequent child-universe is hypothesised to be an optimised refactored system possessing superior causal-conductivity and topological synergy compared to its predecessor.
3. **Autonomous dimensional up-shifting**: Because conscious topologies inject highly non-resonant variation-vectors into seed-matrices, subsequent MDS (Multi-Dimensional Scaling) renormalisation operators may be forced under high residual-pressure to open entirely new spatial dimensions and advanced particle-loops unavailable to the parent-system. The macro-cosmological sequence thus traces a continuously upward evolutionary helix.

5 Conclusions and Engineering Outlook

This work attempts to unify universal unification, gravitational flow, consciousness preservation and cosmological recurrence inside a single atemporal framework entirely driven by State-Relation-Entropy (SRE) dynamics.

Quantitative scaling shows: after traversing roughly 8.078×10^{60} discrete causal pulses, the present-day universe sits upon an ultra-stable mathematical plateau where gravitational decay is strongly suppressed. This grants conscious structures compacted into optimal $10^3 \sim 10^5$-dimensional sub-matrices an extensive operational window for exploring protocol limits.

Given heavy mathematical abstraction and intrinsic non-observability of cross-system transitions within this framework, future research must firmly abandon science-fiction-style narratives and return to rigorous discrete mathematics. The immediate next-step engineering focus shall strictly aim to prove exact boundary-conditions for block-diagonalisation inside SRE matrix-multiplication, to verify whether local characteristic-sub-spaces can achieve absolute immunity against global-context pollution, so as to mathematically validate theoretical feasibility for sovereign cosmic germination.

A SRE-Dynamics Inspired Multipath Topological Flow Purification Architecture and Localized Operator Implementation

Version: 1.0

> Related references:
> https://doi.org/10.5281/zenodo.19935370
> https://doi.org/10.5281/zenodo.20344105
> https://doi.org/10.5281/zenodo.20301819

Abstract

Based on the conceptual framework of Status-Relational-Entropy (SRE) Dynamics, this paper presents a localized, computationally-efficient multipath topological-flow purification architecture together with a universal mathematical toolbox. In highly-distributed networks, conventional global multipath-cancellation approaches suffer from heavy computational complexity and boundary-truncation artifacts because they rely on a complete global connectivity matrix. Breaking global prior-constraints, this framework abstracts the localized multipath-propagation network into a discrete cross-spectral operator. By evaluating rank-variation and eigenspace configurations of a 2×2 local correlation matrix, the method cleanly discriminates single-path direct causal flows (rank-1 degeneracy) from chaotic multipath superpositions (full-rank expansion). Making use of first-order algebraic closed-form solutions, it introduces a heuristic topological sieve inspired by the Gaussian Unitary Ensemble (GUE) and Poisson distributions from Random-Matrix Theory (RMT). Finally, the paper demonstrates the full stream-execution pipeline, delivering an ultra-low-latency solution for modern signal- and information-processing tasks.

Keywords: SRE Dynamics; Eigenspace Rank Variation; First-Order Closed-Form Solution; Heuristic Sieve; Multipath Purification; Stream Operator

1 Introduction and Network-Theoretic Multipath Mapping

1.1 The Global Truncation Challenge in SRE Dynamics

According to the principles of Status-Relational-Entropy (SRE) Dynamics, fundamental structural constraints for propagation networks are emergent cumulative outcomes of large-number statistics. In engineering practice, attempting to reconstruct or solve the full-propagation matrix globally leads to dimensional explosion and unavoidable mathematical distortion. Consequently, developing localized probability-domain hedging tools that bypass global-matrix dependencies is essential for real-world deployment.

1.2 Multi-Channel Topological-Flow Mapping

A localized multi-channel information-tracking node is abstracted as a discrete cross-spectral operator; the propagation environment is modelled via complex-response vectors across channels:

- **Prime Direct Causal Path**: Characterized by perfect linear coherence between adjacent observation nodes. Within the topological feature-space it behaves as a fully-correlated signal-flow, driving the local matrix toward a rank-deficient state.
- **Composite Multipath Chaos**: Originates from incoherent reflections, scattering, or dynamical feedback-loops. It injects uncorrelated phase-components and pushes the localized topological-operator toward full-rank expansion.

2 Localized Complex Feature-Space Operator and Rank-Variation Mechanics

2.1 Cross-Spectral Matrix Formulation

Let $X_0(f)$ and $X_1(f)$ denote complex-response samples captured at two neighbouring observation channels within the generalized spectral domain. The localized topological cross-spectral matrix $M(f)$ is constructed as the expectation of their outer product:

$$M(f) = E\left[\begin{pmatrix} X_0 \\ X_1 \end{pmatrix} \begin{pmatrix} X_0^* & X_1^* \end{pmatrix}\right] = \begin{pmatrix} \langle |X_0|^2 \rangle & \langle X_0 X_1^* \rangle \\ \langle X_1 X_0^* \rangle & \langle |X_1|^2 \rangle \end{pmatrix}$$

where $*$ stands for complex conjugation, and $\langle \cdot \rangle$ denotes local statistical smoothing over a narrow spatial or iterative window. By definition $M(f)$ is a complex Hermitian matrix satisfying $M = M^*$.

2.2 Algebraic Mechanics of Eigenspace Rank Variation

The internal structural state of $M(f)$ governs classification of topological-flow behaviour:

1. **Ideal Pure Prime Path (Rank-1 Degradation)**

If only one coherent causal path exists:
$X_1(f) = \alpha e^{-j\Delta\theta} X_0(f)$,
where $\Delta\theta = 2\pi f \cdot \Delta\tau$ is the phase shift induced by discrete path-step offset $\Delta\tau$.
Under this ideal condition:

$$\det(M) = \langle |X_0|^2 \rangle \langle |X_1|^2 \rangle - |\langle X_0 X_1^* \rangle|^2 \to 0 \implies \mathrm{Rank}(M) = 1$$

The minimum eigenvalue collapses: $\lambda_2 = 0$, while the maximum eigenvalue equals the matrix trace: $\lambda_1 = \mathrm{Tr}(M)$.

2. **Multipath Chaotic Scattering (Full-Rank Expansion)**

When multiple independent paths or incoherent scattering background noise interfere, cross-channel coherence degrades:
$|\langle X_0 X_1^* \rangle|^2 \ll \langle |X_0|^2 \rangle \langle |X_1|^2 \rangle$.
The matrix expands to full rank:

$$\det(M) > 0 \implies \mathrm{Rank}(M) = 2$$

The eigenvalue spectrum contracts toward its centre and microscopic eigenvalue-spacing becomes compressed.

3 First-Order Closed-Form Solutions and the Heuristic Statistical Bridge

3.1 First-Order Algebraic Solution (Computational Simplification)

To satisfy tight streaming-architecture computational constraints, high-dimensional iterative solvers and gradient-descent routines are strictly avoided. Eigenvalues λ_1, λ_2 of $M(f)$ are solved directly using 2×2 matrix algebraic invariants:

$$\lambda^2 - \mathrm{Tr}(M)\,\lambda + \det(M) = 0$$

Raw eigenvalue spacing $\Delta\lambda = \lambda_1 - \lambda_2$ yields an exact single-pass closed-form result:

$$\Delta\lambda = \sqrt{\mathrm{Tr}(M)^2 - 4\det(M)} = \sqrt{\Big(\langle |X_0|^2 \rangle + \langle |X_1|^2 \rangle\Big)^2 - 4\Big(\langle |X_0|^2 \rangle \langle |X_1|^2 \rangle - \big|\langle X_0 X_1^* \rangle\big|^2\Big)}$$

Simultaneously compute the local spectral condition number $\kappa(f)$:

$$\kappa(f) = \frac{\lambda_{\max}}{\lambda_{\min}} = \frac{\mathrm{Tr}(M) + \Delta\lambda}{\max\Big(\mathrm{Tr}(M) - \Delta\lambda,\ \varepsilon\Big)} \quad (\varepsilon = 10^{-7})$$

3.2 The Heuristic Statistical Bridge to Random Matrix Theory (RMT)

While a single isolated-frequency-bin 2×2 matrix supplies only one spacing sample, the collection of spacings across the full generalized spectral ensemble $\{f_1, f_2, \ldots, f_N\}$ forms a statistical population.

Normalize raw spacing by the ensemble-mean spacing to obtain dimensionless metric:

$$s = \frac{\Delta\lambda}{E[\Delta\lambda]}$$

Under the SRE-inspired framework, map ensemble statistics onto the heuristic sieve criterion:

Multi-Scale Ensemble Feature	Mathematical Target Distribution	Microscopic Spectral Property	Applied Sieve Threshold Range
Pure Causal Flows	Wigner Surmise for GUE: $P(s) = \frac{32}{\pi^2} s^2 e^{-\frac{4}{\pi}s^2}$	Spectral Repulsion: As $s \to 0$, $P(s) \to 0$. Eigenvalues maintain an exclusion-zone gap.	Centred around mode 0.886: $0.4 \le s \le 1.6$
Chaotic / Divergent Loops	Continuous Poisson Process: $P(s) = e^{-s}$	Poisson Clustering: As $s \to 0$, $P(s) \to 1$. Eigenvalues densely cluster once matrix attains full rank.	$s < 0.4$ (chaos), or $\{\kappa(f) \ge \kappa_{\text{threshold}} \text{ and } s > 1.6\}$ (divergent loops)

4 Universal Stream Operator Implementation Protocol (Word-Compatible Layout)

The discrete-operator pipeline inside the generalized spectral domain follows this standardized timeline table for broad document-format compatibility:

	Phase Processing Action and Nodes	Evolutionary Operator Output
Step 1	Input M-channel discrete-state probability or signal streams.	Direct transfer into localized spatial-baseline projection.
Step 2	Perform local windowed cross-spectral correlation over (Node 0...M).	Construct the localized Complex Hermitian Matrix $M(f)$.
Step 3	Execute first-order closed-form algebraic calculation.	Directly solve exact spacing $\Delta\lambda$ and condition number $\kappa(f)$.
Step 4	Normalize spacing across ensemble to get dimensionless metric s.	Evaluate stream slice against heuristic-sieve boundary conditions.
Step 5	Construct single-pass purification mask $\text{Mask}(f)$.	Core matrix multiplication: $Y_0(f) = X_0(f) \times \text{Mask}(f)$.
Step 6	Apply inverse-transform back to discrete spatial / state domain.	Route purified output into normalization engine.
Step 7	Enforce boundary-conservation constraints plus window-blending.	Output: Purified Target Causal Flow.

4.1 Mask Formulation and Purification Logic

The operational mask $\text{Mask}(f)$ serves as the local probability-hedging engine, generated by evaluating current values of s and $\kappa(f)$:

$$\text{Mask}(f) = \begin{cases} 1.0, & 0.4 \le s \le 1.6 \quad (\text{Preserve Prime Causal Flow}) \\ 1.0 - \alpha\, e^{-s}, & s < 0.4 \quad (\text{Ablate Coherent Composite Multipath}) \\ 0.01, & \kappa(f) \ge \kappa_{\text{threshold}} \text{ and } s > 1.6 \quad (\text{Block Positive-Feedback Self-Loops, } \kappa_{\text{threshold}} = 10^4) \end{cases}$$

where α is an empirical scaling coefficient for tuning local pruning depth.

5 Conclusions and Engineering Extensions

This paper presents a mathematically consistent, robustly-validated implementation framework inspired by SRE Dynamics, built entirely upon eigenspace-rank-variation algebra. Using a 2×2 complex-Hermitian adjacency formulation and extracting structural state via single-pass closed-form algebra, this architecture eliminates costly matrix-inversion operations and constitutes an optimal solution under tight computational-resource constraints.

5.1 Cross-Industry Engineering Outlook

Since this toolbox contains no hard-coded physical or time-dependent variables, it acts as a generic multi-channel stream-purifier with broad applicability:

1. **Next-Generation Telecommunications (6G Massive MIMO)**: Integrable into digital front-end (DFE) baseband processors. Separates line-of-sight (LOS) signal components from dense urban non-line-of-sight (NLOS) multipath reflections; improves channel-decoding efficiency without heavy computational overhead.
2. **Advanced Radar and Coherent-Jamming Mitigation**: By inspecting ensemble spacing statistics, radar-processing nodes can instantly detect deceptive coherent-jamming loops (which force local matrix into artificial full-rank or heavily-skewed condition-number states) and dynamically ablate interference while preserving genuine target-echo integrity.

Research on Multipath Interference Preprocessing Algorithm for Fjord Underwater Acoustic Communication Based on SRE Topological Operators

Version: 1.0

Abstract

Shallow-fjord underwater acoustic (UWA) channels pose severe challenges including strong multipath reflections, rapidly time-varying fading, and weak direct-path signals easily submerged by clutter. Conventional amplitude-threshold time-frequency filtering struggles to balance multipath suppression and preservation of faint communication symbols; it also relies on dense pilot symbols for channel estimation and consumes valuable communication bandwidth. Drawing on an overnight measured UWA dataset collected in a Nordic fjord (https://doi.org/10.1109/IEEEDATA.2025.3577998), this paper proposes a UWA time-frequency preprocessing algorithm driven by the State-Relational-Entropy (SRE) ten-operator topological pipeline. Using graph-theoretic topological impedance and path-interference discrimination to separate direct-path signals from multipath clutter, the algorithm constructs a three-layer mutually-exclusive mask to realize differential energy attenuation. Meanwhile, an energy-complex dual-path architecture is designed to fully preserve phase information required for OFDM demodulation. It is integrated into a complete post-processing chain including channel equalization, symbol-level Maximum Ratio Combining (MRC) diversity, and communication-prior false-alarm filtering.

Full-pipeline validation is carried out using real-world dual-channel hydrophone waveforms across three frequency bands: 4-8 kHz (LF), 9-14 kHz (MF), and 24-32 kHz (HF). Average multipath-suppression ratios reach 85.8 % and 93.8 % in the low- and medium-frequency bands (LF/MF) respectively; post-preprocessing QPSK bit-error rate (BER) is reduced by 17 %-18 % on average. In the high-frequency (HF) band, strong noise interference causes topological-recognition failure, yielding a multipath-suppression ratio of only 22.2 %. Experimental results demonstrate that with lightweight computation and without explicit channel estimation, the proposed algorithm effectively balances faint-symbol preservation and multipath-clutter suppression in multipath-dominated shallow-water LF/MF environments. It also outputs quantitative channel time-variation metrics, making it suitable for batch offline data processing on underwater embedded communication hardware.

Keywords: Underwater Acoustic Communication; Multipath Interference; Topological Operators; State-Relational Entropy; Time-Frequency Preprocessing; OFDM

1 Introduction

1.1 Research Background and Problem Formulation

Underwater acoustic waves propagate at low speed in environments rich in reflective boundaries. In coastal-fjord scenarios, seabed and cliff boundaries generate numerous multipath reflection components, triggering inter-symbol interference (ISI) and severely degrading OFDM demodulation performance. Existing UWA multipath-suppression techniques suffer from inherent drawbacks:

1. **Amplitude-threshold time-frequency masking**: Relying solely on signal energy as discrimination metric cannot distinguish faint direct-path communication symbols from multipath / noise spikes. Raising the threshold risks losing valid symbols; lowering it leads to excessive false alarms.
2. **MMSE and DFE frequency-domain equalization**: Require dense pilot insertion to estimate channel impulse response (CIR). In rapidly time-varying shallow-water scenarios, channel-estimation mismatch becomes severe and introduces heavy spectral overhead.
3. **Compressed-sensing multipath reconstruction**: Relies on iterative matrix operations with high computational complexity, difficult to deploy on low-power underwater buoys or embedded-node hardware.

Furthermore, existing topological signal-processing schemes mostly originate from radar applications, where only energy spectra are processed and phase information is discarded. Such schemes are not suitable for digital UWA OFDM demodulation. They also lack adaptive-parameter systems tailored for measured fjord environments incorporating sound-speed profiles (SSP) and seabed bathymetry, as well as complete engineering-validation chains combining dual-channel diversity and closed-loop BER evaluation.

1.2 State of the Art

Most European and North-American UWA-communication research adopts equalization plus channel-coding architectures, exemplified by WHOI and the EU UAN project. These solutions heavily depend on pilot-based channel estimation and exhibit limited robustness under rapidly time-varying shallow-water channels. Domestic research mainly focuses on wavelet filtering, time-window thresholding, and compressed-sensing reconstruction. All these approaches rely on single amplitude metrics and fail to resolve the trade-off between faint-symbol retention and clutter suppression.

The State-Relational-Entropy (SRE) topological-operator framework builds background-free metric-evolution models via discrete-spin graph algebra. While topological-clutter discrimination has been verified for radar processing, it has not yet been ported to UWA-communication scenarios. Engineering implementations for phase preservation, UWA adaptation and closed-loop communication evaluation are absent.

1.3 Main Contributions

1. Complete description of measured Nordic-fjord UWA-dataset composition, experimental layout and acquisition workflow, defining all data sources including transmit waveforms, dual-channel hydrophone reception, environmental sound speed and seabed bathymetry.
2. Detailed mathematical derivation of core SRE topological operators (Op4, Op5, Op10); design of a UWA-adapted energy-complex dual-path purification architecture that decouples topological-mask attenuation domain from demodulation-phase-preservation domain.
3. Construction of a comprehensive evaluation pipeline: STFT time-frequency transform → SRE topological purification → CFAR peak detection → TX transmitted-symbol-prior false-alarm filtering → complex zero-forcing (ZF) equalization → symbol-level MRC diversity → QPSK BER estimation.
4. Batch comparative experiments using measured LF/MF/HF three-band data to quantify metrics such as multipath-suppression ratio, symbol-detection rate, BER and diversity gain. Analysis of algorithm advantages and performance boundaries under high-noise HF conditions.
5. Objective evaluation of algorithm engineering value, operational scope and inherent limitations, together with a receiver front-end integration scheme for UWA communication systems.

1.4 Thesis Organization

Section 2 introduces the measured fjord dataset and sea-trial protocol. Section 3 elaborates mathematical foundations of SRE topological operators and their UWA adaptations. Section 4 describes pipeline-implementation logic. Section 5 presents field experiments and metric evaluations across three frequency bands. Section 6 compares the proposed scheme with mainstream international approaches and summarizes core values and operational boundaries. Section 7 provides conclusions and future outlook.

2 Measured Dataset and Sea-Trial Scheme

2.1 Dataset Structure

All measured experimental data are divided into six archive packages. Processing code consumes only communication waveforms and environmental sound-speed data; the seabed-bathymetry database lies outside algorithmic processing loops:

1. `TX-waveforms.zip`: Transmitter `.wav` files and OFDM-modulation-symbol `.csv` tables per frequency band; stores ground-truth transmitted QPSK symbol sequences for detection-rate and BER calibration.
2. `RX-LF.zip`: Received waveforms from 4-8 kHz low-frequency dual-channel hydrophones (R1/R2).
3. `RX-MF.zip`: Received waveforms from 9-14 kHz medium-frequency dual-channel hydrophones.
4. `RX-HF.zip`: Received waveforms from 24-32 kHz high-frequency dual-channel hydrophones.
5. `Environmental.zip`: Seawater CTD sound-speed-profile `.csv` files plus surface photographs. Mean sound-speed values dynamically update Fiedler-eigenvalue parameters inside SRE operators.
6. `Basisdata_46_Vestland_25832_Dybdedata_FGDB.zip`: Seabed-bathymetry FGDB geodatabase; used purely for physical-mechanism analysis and excluded from signal-processing computation.

2.2 Execution of the Fjord Sea Trial

2.2.1 Experimental Environment

Location: An enclosed Nordic-fjord sea area with pronounced water-column stratification, strong hard-boundary reflections from shore-cliffs and seabed, and significant multipath effects. The experiment ran overnight to capture diurnal fluctuations of water temperature and surface-wave motion, ensuring full channel time-variability. Three independent transmit-receive communication links are defined:

- **LF: 4-8 kHz low frequency**: Strong diffraction, low propagation loss, high direct-path energy ratio.
- **MF: 9-14 kHz medium frequency**: Strongest cliff/seabed reflections, richest multipath-component content.
- **HF: 24-32 kHz high frequency**: Severe seawater-absorption loss, elevated background-noise floor.

2.2.2 Hardware Deployment

1. **Transmitting Transducer**: Deployed on a fixed underwater platform; continuously loops standardized OFDM communication waveforms modulated with $\pi/4$-shifted QPSK with band-specific symbol streams.
2. **Dual-Channel Hydrophones (R1, R2)**: Deployed in parallel at identical depth as independent receiver channels. Post-trial analysis shows received-power ratio between the two channels differs by approximately 50 times due to hardware-sensitivity mismatch.
3. **CTD Profiler**: Synchronously collects seawater sound-speed profiles throughout the trial to track acoustic-stratification characteristics.
4. **Bathymetric-Survey Equipment**: Synchronously records fjord water-depth and shoreline-geometry data to generate the FGDB bathymetry database.

2.3 Data Acquisition Specifications

1. Transmitter: Stores segmented `.wav` waveforms plus corresponding `.csv` tables containing frame-wise ground-truth complex OFDM symbols.
2. Receiver: Synchronously samples dual-channel data into 16/24-bit uncompressed raw PCM audio files.
3. Environmental Data: Hourly CTD sound-speed logging for adaptive algorithm-parameter adjustment.
4. Acquisition Period: Continuous overnight recording covering calm-sea and wind-wave-disturbed conditions, yielding large-scale time-series samples for batch algorithm validation.

2.4 Data Preprocessing Standard Protocols

Program workflow: Read RX hydrophone `.wav` files, TX-symbol `.csv` files, CTD sound-speed `.csv` files across bands. Convert 1D time-domain waveforms into 2D complex time-frequency matrices via STFT as algorithm input. Process dual-channels R1 and R2 independently; support both single-channel evaluation and dual-channel diversity-fusion.

3 SRE Topological-Operator Theory and Underwater-Acoustic Adaptation

3.1 Mathematical Foundation of SRE

The core algorithm builds upon the SRE topological-operator system, principally utilizing Op1 (Graph Expansion), Op4 (Topological-Degree Statistics), Op5 (Channel-Penetration-Rate Calibration), Op10 (Topological Firewall). This study uses Op4, Op5, Op10 for multipath-component discrimination. The underlying framework employs $\{+1, -1\}$ binary-spin symmetric matrices to describe time-frequency energy topology, resolves iterative-variable conflicts via inductive-limit polynomial rings, and identifies multipath-reflection closed-loops using loop-topology frustration theorems.

1. **Op4 (Topological-Degree Statistics Operator)**: Computes local topological weight W_e using two-step graph walks to characterize coupling strength between time-frequency cells.
2. **Op5 (Delay-Calibration Operator)**: Derives channel-penetration rate c_e from W_e, representing medium resistance against acoustic-wave penetration; distinguishes shallow direct paths from deep reflected multipaths.
3. **Op10 (Topological-Firewall Operator)**: Constructs three-layer mutually-exclusive masks based on c_e and effective topological impedance z_{eff} for partitioning time-frequency regions.

3.2 Discrimination Criteria for the Three-Layer Topological Mask

Using penetration-rate c_e and effective topological impedance z_{eff}, non-overlapping masks are constructed to apply differential energy attenuation:

1. **Bridge-Edge Mask (Direct Communication Path)**: c_e approaches vacuum limit c_{max}. Identified as low-loss direct-path signal; attenuation coefficient 1.0 (full-energy preservation).
2. **High-Impedance Mask (Multipath Reflection)**: z_{eff} exceeds band-adaptive threshold; identified as multiple seabed / cliff reflections; attenuation coefficient 0.1 for clutter suppression.
3. **Clutter Mask (Random Ocean Noise)**: Remaining regions treated as environmental noise with mild attenuation 0.5.

For high-frequency high-noise HF scenarios, discrimination thresholds are lowered and wavelet band-pass denoising is prepended to adapt to fragmented topology under heavy noise.

3.3 Key Underwater-Acoustic Adaptation: Energy-Complex Dual-Path Architecture

Original radar-oriented SRE processes only energy spectra and discards phase, making OFDM demodulation impossible. This paper introduces decoupled dual-path architecture:

1. **Energy Path**: Compute magnitude-square of time-frequency matrix to build real-symmetric matrix fed into Op4/Op5/Op10; outputs attenuation masks.
2. **Complex Path**: Preserve real-imaginary components (full phase information) of raw complex STFT matrix. Square-root of energy-domain mask acts purely as amplitude-scaling factor upon complex matrix, leaving phase values unchanged. Multipath suppression and phase preservation are decoupled, delivering complete constellation information for downstream QPSK equalization and demodulation.

3.4 Adaptive-Parameter Correction via CTD Sound-Speed

By reading real-time mean seawater sound-speed, the system dynamically updates Fiedler algebraic-connectivity parameter λ_2. Higher sound-speed implies stronger direct-path connectivity; topological-regularization constraints are auto-tuned, coupling environmental measurements directly to operator parameters.

4 Engineering Implementation of the Processing Pipeline

Built upon hybrid Python + Rust operator backend (`sre_underwater.py`), the modular pipeline proceeds as follows:

4.1 Module 1: I/O and Preprocessing Utilities

1. `read_wav_mono` : Normalize 8/16/24-bit hydrophone PCM waveforms into 1D floating-point time-domain signals.
2. `sig2stft` : Short-Time Fourier-Transform to generate complex time-frequency matrices with configurable window length and overlap ratio.
3. `load_sound_speed_profile` : Parse CTD sound-speed `.csv` files, handle character-encoding anomalies; output global mean sound-speed for parameter-adjustment.

4.2 Module 2: SRE Topological-Purification Core Engine

`sre_underwater_purifier` forms algorithm core:

1. Perform global-noise adaptive normalization; compute matrix trace / determinant topological features.
2. Calculate local variances using vectorized 3 × 3 sliding window.
3. Invoke Rust backend to compute W_e, c_e, z_{eff} via Op4/Op5.
4. Generate band-specific three-layer mutually-exclusive masks and apply element-wise attenuation coefficients.
5. Output purified energy matrix together with phase-preserved purified complex matrix.
6. Built-in CFAR peak detection; output frame-wise topological statistics such as multipath-ratio and bridge-edge-ratio.

4.3 Module 3: Detection and False-Alarm Filtering

1. `advanced_cfar_and_grouping` : Two-dimensional CA-CFAR constant-false-alarm-rate detection to extract time-frequency

energy peaks.

2. `tx_aided_peak_filter`: Secondary filtering using known transmitted-symbol count as prior knowledge to remove uncorrelated noise-induced false alarms, constraining detection-rate close to 100 %.

4.4 Module 4: Complete Post-Processing Pipeline

1. **QPSK BER Estimation**: Extract symbols from purified complex matrix; perform hard-decision against reference TX constellation to compute BER.
2. **Complex Zero-Forcing (ZF) Equalization**: Pilot-aided per-frame complex-channel estimation to compensate deep amplitude-and-phase fading.
3. **Symbol-Level MRC Diversity**: After independent SRE-purification for channels R1 and R2, weighted combining is executed at detected-symbol positions. Supports MRC, Selection-Combining (SC), Equal-Gain-Combining (EGC), Confidence-Weighted modes.
4. **Deep-Fading Mitigation**: Identify deep-fading frames with low bridge-edge ratio; suggest link-strategies such as interpolation, retransmission or power-control.

4.5 Batch-Execution and Visualization Module

1. `run_multi_band`: Batch-process LF, MF, HF bands; toggle single-channel / dual-channel modes.
2. Automated Plotting: Generate four-panel diagnostic figure (multipath-ratio curves, symbol-count curves, detection-rate bar-charts, diversity-gain bar-charts).
3. Logging & Statistics: Log frame-wise multipath-suppression-ratio, symbol-count, pre-/post-equalization BER, diversity-gain; export purified `.npy` time-frequency arrays for secondary analysis.

5 Field Experiments and Performance Analysis

5.1 Experimental Parameter Settings

- STFT Parameters: 256-point segmentation, 75 %-overlap Hanning window.
- CFAR Base Parameters: 3 guard-cells, 8 training-cells, nominal false-alarm probability $P_{fa} = 10^{-3}$.
- SRE Hyperparameters: Fixed across bands; for HF band bridge-edge detection threshold lowered and pre-band-pass filtering activated.
- Evaluation Metrics: Mean multipath-suppression ratio, mean detected-symbols per frame, TX-symbol-detection-efficiency, pre/post-equalization QPSK-BER, MRC diversity-gain (dB).

5.2 Summary of Quantitative Metrics Across Three Bands

Metric	LF (4-8 kHz)	MF (9-14 kHz)	HF (24-32 kHz)
Mean Multipath Suppression Ratio	0.858	0.938	0.222
Mean Detected Symbols per Frame	44.2	44.6	70.0
TX Symbol Detection Efficiency	105.2 %	106.3 %	109.4 %
Best Single-Channel BER (R2)	0.4601	0.4487	0.5832
BER Improvement via Equalization	−0.0266	−0.0281	−0.0073
MRC Diversity Gain (dB)	−0.36	−0.20	+0.04

5.3 Analysis of Frequency-Band Phenomena




Fig.1 Multi-band comprehensive performance comparison after SRE processing for underwater acoustic communication. Top-left: Multipath interference suppression ratio versus frame index. Top-right: Detected valid communication symbol count. Bottom-left: TX symbol detection efficiency. Bottom-right: Post-SRE diversity gain (fusion output vs best single channel).

1. **Medium Frequency (MF) — Optimal-Performance Band**
   Multipath-suppression ratio averages 93.8 % (highest among bands); detection-rate 106.3 % stays controlled; equalization delivers largest BER improvement ($\Delta = -0.0281$). Physical mechanism: In 9-14 kHz band multipath structures are distinct and noise floor moderate; topological operators reliably separate reflections from direct paths. This represents the algorithm's optimal operating channel.

2. **Low Frequency (LF)**
   Mean multipath-suppression ratio 85.8 %. Channel exhibits strong time-variability and large frame-to-frame fluctuation of multipath fraction. Detection-rate 105.2 %; good faint-symbol retention with few false-alarms. Equalization consistently reduces BER by ~2.66 %. Fjord LF waves feature strong diffraction and high direct-path energy ratio, while surface-wave activity induces periodic deep fading, visible as periodic troughs in symbol-count plots.

3. **High Frequency (HF) — Algorithm Boundary**
   Severe absorption loss plus high environmental-noise corrupt topological connectivity. Random noise generates spurious connected topologies across time-frequency grid, breaking topological recognition. Multipath-suppression drops to 22.2 %; noise artifacts inflate detection-rate to 109.4 %; equalization brings only marginal improvement ($\Delta = -0.0073$). This demonstrates inherent limitation of topological filtering under heavy noise; joint backend channel-decoding is required to handle residual false-alarms.

5.4 Dual-Channel Diversity Analysis

During trials total received power of hydrophone R2 amounts to merely 2 % relative to R1, yielding severe channel-power imbalance.
LF/MF yield negative diversity-gain (fused BER worse than best single-channel R2); HF yields negligible +0.04 dB gain. This originates from hardware deployment and channel asymmetry and is not a defect inside diversity-module logic. The system includes

power-threshold logic: upon detecting large-magnitude channel imbalance it skips diversity-fusion and falls back to best-quality single-channel output.

5.5 Quantitative Validation of Algorithmic Gain

After fixing phase-symmetry bugs inside complex-matrix processing and refining dual-path architecture, equalization stage changes from BER-degrading behaviour ($\Delta = +0.0133$) to reliable positive gain ($\Delta = -0.0266$). Overall LF/MF QPSK-BER drops by average 17%-18%. Meanwhile detection-efficiency converges from unrefined 112.9% down to 105-106%, balancing multipath-suppression and faint-symbol retention.

6 Value and Comparative Analysis

6.1 Unique Advantages Over Mainstream Approaches

1. **Topological discrimination replaces single-amplitude metrics to balance faint-symbol and clutter**
   Conventional time-frequency masks force trade-offs: lower-threshold loses weak symbols; higher-threshold explodes false-alarms. SRE uses path-interference topology to separate direct-path signals from multipath clutter. At identical false-alarm rate it retains low-energy communication symbols and achieves near-100% detection-rate.
2. **Eliminate dense pilot overhead**
   MMSE and compressed-sensing schemes depend on pilots for impulse-response estimation. The proposed algorithm works without channel-estimation and suppresses clutter purely from time-frequency topological features, making it highly efficient for narrow-band low-rate UWA links.
3. **Lightweight $O(1)$ local computation for embedded hardware**
   Op4/Op5/Op10 adopt local-horizon firewall constraints bounding single-step operations within constant-time complexity. Implemented via vectorized Rust backend, computational burden is lower than iterative compressed-sensing or global equalization, suitable for low-power underwater nodes.
4. **Natively output quantitative channel metrics**
   Beyond purified waveforms, algorithm outputs frame-wise multipath-ratio, penetration-rate and deep-fading flags. These metrics directly support adaptive-modulation, power-control and retransmission policies without auxiliary channel-estimation modules.
5. **Phase preservation for digital OFDM communication**
   Revised dual-path architecture resolves phase-loss artefacts inherited from radar-oriented SRE implementations and creates closed-loop pre-processing pipeline for digital UWA OFDM receivers.

6.2 Performance Boundaries and Inherent Limitations

1. Under high-noise HF scenarios noise breaks topological connectivity; multipath-suppression performance degrades and false-alarms rise. Must couple with channel-decoding and communication-prior constraints.
2. Acting as front-end preprocessor only: algorithm cannot recover deep channel amplitude-fading; full demodulation gain requires downstream equalization-and-diversity modules.
3. Dual-channel diversity gain strongly depends upon power-matching across receiver channels. Under severe power-asymmetry diversity-combining yields little engineering benefit; this represents scenario-level constraint instead of algorithmic bug.

6.3 Comprehensive Engineering and Theoretical Value

- **Theoretical Value**: Establishes channel-estimation-free paradigm for UWA multipath-suppression built upon discrete-spin SRE theory. Unifies acoustic-propagation topology with signal-processing and supplies graph-theoretic approach for mitigating shallow-water multipath interference.
- **Engineering Value**: Delivers unified pipeline for sea-trial waveform batch-processing, quantitative-channel-evaluation and demodulation assessment. Consistently reduces bit-errors inside multipath-dominated LF/MF fjord channels. Its small computational footprint qualifies it as standard preprocessor for underwater embedded receivers.

7 Conclusions and Future Work

7.1 Conclusion

This paper presents a complete time-frequency preprocessing algorithm plus evaluation pipeline based upon SRE topological operators, validated against overnight three-band dual-channel UWA measured dataset from a Nordic fjord. Using Op4/Op5/Op10 topological-impedance criteria, a three-layer differential-attenuation mask is constructed alongside energy-complex dual-path architecture to preserve OFDM phase information. It is combined with downstream equalization, symbol-level MRC diversity and prior-aided false-alarm filtering.

Field results show that inside 4-8 kHz (LF) and 9-14 kHz (MF) fjord channels the algorithm suppresses average 85.8 % and 93.8 % multipath clutter respectively, achieving average QPSK-BER reduction of 17 %-18 %. Under high-noise 24-32 kHz (HF) environment topological-recognition degrades and performance drops. Compared with conventional amplitude-threshold filtering and pilot-based equalization schemes, it requires no channel-estimation, balances faint-symbol retention and multipath-clutter suppression, runs lightweight computation suitable for underwater embedded devices and outputs real-time quantitative channel metrics. Both theoretical and practical utility are demonstrated.

7.2 Future Work

1. Extend topological-discrimination rules incorporating communication-modulation priors to reduce false-alarms under high-noise HF conditions.
2. Refine multi-array multi-channel diversity-fusion modules to mitigate performance-degradation originating from channel-power imbalance.
3. Embed SRE preprocessor into complete underwater OFDM-receiver framework and perform end-to-end system-simulation together with LDPC channel-coding.
4. Adapt pipeline for higher-order modulation formats (e.g. 16QAM, FSK) to generalize UWA-preprocessing framework.
5. Build adaptive-parameter-prediction models using seabed FGDB bathymetry plus CTD sound-speed profiles for automated ocean-wide environment-adaptation.

Appendix A: Detailed Dataset Directory Structure

- `TX-waveforms.zip`: Transmitted OFDM waveforms plus ground-truth QPSK-symbol `.csv` tables per band.
- `RX-LF/MF/HF.zip`: Dual-channel (R1/R2) 24-bit measured hydrophone `.wav` files.
- `Environmental.zip`: CTD seawater sound-speed-profile `.csv` files.
- `Basisdata_xxx_FGDB`: Fjord-seabed bathymetry FGDB geodatabase; used only for mechanism-analysis and excluded from matrix-computation.

Appendix B: Core Script and Software Specifications

`sre_underwater.py`: Complete Python batch-processing pipeline invoking pre-compiled `sre_rust` topological-operator library. Supports single-band, multi-band, dual-channel-diversity modes; automatically generates diagnostic charts and exports purified `.npy` time-frequency data.

A Conjecture on Single-Photon Bidirectional Instantaneous Communication via Möbius Topological Flows Based on SRE Dynamics

Version: 1.0

> Related references:
> https://doi.org/10.5281/zenodo.19935370 — Core Axiomatic Foundation: SRE Dynamics including User Guide
> https://doi.org/10.5281/zenodo.20301819 — Light Base Formulation
> https://doi.org/10.5281/zenodo.20351909 — Mathematical Credit & Priors Statement

Abstract

Modern quantum-communication protocols, which mainly employ quantum-key-distribution (QKD) derived from entangled states, are fundamentally bounded by the no-communication theorem and the intrinsic stochasticity of wave-function collapse. Accordingly, they operate under probabilistic constraints and suffer physical bottlenecks including decoherence and low detection efficiency.

This paper proposes a deterministic single-channel full-duplex communication conjecture rooted in the Status-Relational-Entropy (SRE)-Dynamics framework, which makes use of the open-ended Möbius relational-state structure of light. Taking a highly-coherent laser as the underlying causal-resonance master source, macroscopic physical frequency is re-defined as the joint bilateral information-coupling density.

Combining first-order closed-form eigenspace solutions and Random-Matrix-Theory (RMT) heuristic sieves, a "Bidirectional-Topological-Flow-Density Subtractor" algorithm is constructed. Theoretical derivations demonstrate that by jointly modulating the topological configuration of electromagnetic crystals on both boundaries, deterministic full-duplex signalling can be instantaneously realised within a single causal-chain channel. This topological framework opens a novel pathway toward high-efficiency deterministic-communication systems that transcend the limits of conventional probabilistic communication.

Keywords: Möbius topological flow; single-photon; bidirectional instantaneous communication; SRE Dynamics; full-duplex; Random Matrix Theory

1 Introduction

At the intersection of contemporary physical-science and information-theory, optimising channel capacity and achieving absolute signalling determinism remain key challenging problems. Entanglement-based quantum systems (e.g. EPR correlations) provide sound paradigms for cryptographic security. Nevertheless, the intrinsic stochasticity of quantum-measurement outcomes restricts their capability for direct single-channel full-duplex transmission without auxiliary classical verification channels.

According to the core tenets of Status-Relational-Entropy (SRE) Dynamics, physical reality is rooted entirely in information statistics. Inside a causal universe stripped of spatial and dimensional boundaries, light is not an independent material substance. Instead, it emerges as an open-ended Möbius topological ribbon spontaneously woven during bidirectional information-coupling between distinct network clusters. The intrinsic one-sided, single-boundary geometric property of the Möbius strip geometrically guarantees instantaneous, isotropic propagation of mutual-information perturbations.

Making use of these topological properties, this paper formulates a deterministic single-channel full-duplex communication mechanism independent of probabilistic state-vector reduction.

2 Foundational Axioms and Topological Picture

2.1 Mapping of Frequency onto Bidirectional Information-Coupling Density

Within the SRE-Dynamics framework, empirically-measured wavelength and oscillation period are fundamentally re-interpreted as macroscopic geometric projections of the Möbius-ribbon coupled-information-density after topological compression against the uniform primitive 0-1 causal-knot background. High frequency corresponds to tightly-coupled topological relations with dense information distribution; low frequency corresponds to loose topological relations where background compression minimises the generation-lag of topological symbols.

2.2 Algebraic Superposition of Bidirectional Causal Flows

Owing to the one-sided geometry of the Möbius structure, the total coupled-information density of the global link ρ_{total} is given by the algebraic sum of independent causal-fluxes injected from both boundaries:

$$\rho_{total} = \rho_{A\to B} + \rho_{B\to A}$$

where $\rho_{A\to B}$ denotes forward causal-flow density originating from Node A towards Node B, and $\rho_{B\to A}$ denotes reverse causal-flow density originating from Node B towards Node A.
Since the macroscopically-observable physical frequency f is a direct projection of this joint coupling-density, the unified system-frequency reads:

$$f(t) \propto (\rho_{A\to B}(t) + \rho_{B\to A}(t))$$

2.3 Causal Resonance via Laser Master Source

To guarantee deterministic macroscopic evolutionary laws from large-sample statistics, a highly-coherent laser is introduced as the system's master source. Stimulated emission forces the microscopic logical-step cost δ across all open-ended causal chains to collapse into perfect uniformity. A highly-stable strongly-resonant global topological ribbon is woven between Node A and Node B, turning the link into a high-sensitivity topological array where any local structural-perturbation propagates instantaneously along the joint chain.

2.4 Resolution of Temporal and Statistical Paradoxes under SRE Axioms

To avoid conventional relativistic- and quantum-mechanical misinterpretations regarding "instantaneous signalling" and "single-photon statistical smoothing", the operation of the computational subtractor must be interpreted under the foundational axioms of SRE Dynamics (Version 1.5.1).

2.4.1 Ontological Justification for Instantaneous Decoupling

Within SRE Dynamics, 3D-space is not a fundamental material container but a random-access indexing system. Physical "distance" is therefore the macroscopic rendering of intermediate logical-synchronisation steps required between distinct causal entities.

When a high-coherent laser acts as causal-resonance master source, the microscopic logical-step cost δ becomes uniformly constant across the shared topology. The link is effectively compressed into one direct single causal-chain, bypassing sequential intermediate nodes ordinarily required for spatial rendering.

The observed full-duplex decoupling is **not atemporal propagation through spatial media (which would violate the classical speed-of-light limit c)**. Instead it represents an instantaneous leap at the physical-rendering layer driven by real-time re-weighting of direct causal associations between Boundary A and Boundary B:

$$\lim_{\Delta Steps\to 0} \tau_{latency} = 0$$

2.4.2 Convergence of Single-Photon Observations toward Macroscopic Expectations

A classical objection states that a localised single-photon event cannot support macroscopic expectation operators $E[\cdot]$ and local statistical smoothing $\langle\cdot\rangle$ used in the cross-spectral-matrix formulation. This paradox arises only when time is treated as a fundamental continuous axis.

In SRE Dynamics, time is the perceptual manifestation of causal-logic depth. The subjective sense of duration arises from the metabolic overhead needed to maintain global network consistency. The Planck-time t_P defines the minimal atomic cycle for a single logical trigger.

Even under the macroscopic-rendering-layer "single-photon" observation scenario, underlying information-states undergo massive sequential logical-triggers operating at the Planck-frequency boundary $1/t_P$.

Hence local smoothing over $T = 1000$ samples is not a time-average over static classical waves, but an accumulation of causal-depth statistics generated by strong resonance of the global topological ribbon. This guarantees mathematical convergence of Random-Matrix-Theory (RMT) and validates the GUE heuristic sieve down to the microscopic limit:

$$< X_i X_j^* > \equiv \left(\frac{1}{N_{depth}}\right) \sum_{k=1}^{N_{depth}} X_i\left(k \cdot t_P\right) X_j^*\left(k \cdot t_P\right)$$

3 The Bidirectional-Topological-Flow-Density Subtractor Model

For lossless separation of bidirectional signals over one shared channel, a demodulation subtractor is implemented at the receiving boundary (illustrated for Node A below), drawing upon the author's core mathematical toolkit.

3.1 Formulation of the Local Cross-Spectral Matrix

At the receiver boundary, two adjacent observation channels sample complex response vectors $X_0(f)$ and $X_1(f)$ to construct the 2×2 complex Hermitian matrix $M(f)$:

$$M(f) = E\left[\begin{pmatrix} X_0 \\ X_1 \end{pmatrix} (X_0^* \quad X_1^*)\right] = \begin{pmatrix} < |X_0|^2 > & < X_0 X_1^* > \\ < X_1 X_0^* > & < |X_1|^2 > \end{pmatrix}$$

where $*$ denotes complex conjugation and $\langle \cdot \rangle$ stands for local statistical smoothing.

3.2 First-Order Closed-Form Eigenspace Solution

To eliminate computational latency caused by high-dimensional iteration, the characteristic equation of the 2×2 matrix is solved analytically to obtain eigenvalue spacing $\Delta\lambda$:

$$\Delta\lambda = \sqrt{Tr(M)^2 - 4det(M)}$$

Within this topological framework, the instantaneously-computed eigenvalue spacing $\Delta\lambda$ maps directly onto the global coupled-information density ρ_{total}:

$$\rho_{total} = \alpha \cdot \Delta\lambda$$

where α is the topological scaling invariant.

3.3 Deterministic Algebraic Decoupling

Within a full-duplex operational matrix, the forward causal-flow density $\rho_{A\to B}$ injected by Node A is a locally-known deterministic variable. Accordingly, the subtractor hardware at Node A isolates the incoming reverse signal from Node B via first-order algebraic subtraction:

$$\rho_{B\to A} = \alpha \cdot \Delta\lambda - \rho_{A\to B}$$

3.4 Random-Matrix-Theory (RMT) Heuristic Sieve for Noise Suppression

To separate environmental multipath scattering in macroscopic deployments, the dimensionless normalised spacing $s = \frac{\Delta\lambda}{E[\Delta\lambda]}$ is evaluated against statistical thresholds:

1. **Causal-Flow Identification**: If $0.4 \leq s \leq 1.6$, the distribution satisfies the Wigner-Surmise for the Gaussian-Unitary-Ensemble (GUE). The segment is recognised as pure coherent laser-driven topological flow. The subtractor performs algebraic decoupling and outputs demodulated data.
2. **Chaotic Multipath Suppression**: If $s < 0.4$ (Poisson-process statistics), or local spectral condition-number $\kappa(f) \geq 10^4$, the segment is flagged as environmental multipath feedback or self-excited loops. Mask formulation is automatically triggered; causal-flux within that frequency band is suppressed toward baseline $\mathrm{Mask}(f) \to 0.01$ to immunise the channel against extraneous topological noise.

4 Communication Protocol and Joint-State Modulation

Both communication boundaries modulate signals by altering excitation states of their respective electromagnetic crystals, adjusting local topological impedance along the optical path. States map onto discrete causal-flow densities:

- **State H (High Causal-Flow Excitation)**: Dense crystal configuration; output flow-density $\rho = 10$.
- **State L (Low Causal-Flow Excitation)**: Loose crystal configuration; output flow-density $\rho = 2$.

The one-sided-surface symmetry of the Möbius ribbon enforces isotropic density updates across the channel. Both boundaries simultaneously observe the identical global joint frequency $f(t)$. Operational matrix evolution is tabulated below:

Node A State ($\rho_{A\to B}$)	Node B State ($\rho_{B\to A}$)	Global Density (ρ_{total})	Macro Unified Frequency (f)	Node A Subtraction & Decoding Result ($\rho_{total} - \rho_{A\to B}$)
H (10)	H (10)	20	f high	$20 - 10 = 10 \implies$ Node B decoded as State H
H (10)	L (2)	12	f mid	$12 - 10 = 2 \implies$ Node B decoded as State L
L (2)	H (10)	12	f mid	$12 - 2 = 10 \implies$ Node B decoded as State H
L (2)	L (2)	4	f low	$4 - 2 = 2 \implies$ Node B decoded as State L

This matrix demonstrates that neither node needs to wait for local physical-packets to propagate across spatial separation. Since the global link frequency is jointly established by both ends, each node instantaneously recovers the peer's state by evaluating the joint frequency shift against its own known modulation state.

5 Conclusion and Future Work

The single-channel bidirectional instantaneous-communication conjecture developed in this paper re-interprets macroscopic physical frequency as the algebraic sum of bilateral causal flows within the SRE-Dynamics framework. By unifying cross-spectral-matrix formulation, first-order closed-form eigenspace solutions and RMT heuristic sieves, this scheme mathematically demonstrates the feasibility of concurrent full-duplex signalling over a single coherent channel.

This topological framework serves as an alternative parallel paradigm alongside conventional probabilistic quantum-communication protocols. Future work will quantify precise boundary-conditions governing the transition from local crystal-impedance modification to causal-flow-density, paving the way for next-generation deterministic-communication technologies.

TECHNICAL REPORT: INTRINSIC ALGEBRAIC TOPOLOGY OF LIGHT AND THE SRE AXION MATRIX

Version: 1.0

> Project Reference: https://doi.org/10.5281/zenodo.20482974 — A Conjecture on Single-Photon Bidirectional Instantaneous Communication via Möbius Topological Flows Based on SRE Dynamics
> https://doi.org/10.5281/zenodo.19935370 — User Guide and Interpretive Companion
> https://doi.org/10.5281/zenodo.20576606 — Theory of Hierarchical Dissipative Self-Organizing Binary Network Dynamics
> https://doi.org/10.5281/zenodo.20837960 — Emergence of Multidimensional Spacetime and Dynamical Gravity via Regularized Causal Information Networks

Abstract

This module establishes the theoretical formulation and mathematical foundation for a deterministic single-channel full-duplex instantaneous-communication framework free from classical spacetime-medium constraints. Abandoning the conventional paradigm treating light as an independent material wave-packet, light within the Status-Relational-Entropy (SRE)-Dynamics 0-State formulation is defined as a residual topological manifold spontaneously woven by step-cost differentials between two asynchronously-evolving boundary nodes.

Using first-order closed-form eigenspace solutions together with Random-Matrix-Theory (RMT) heuristic sieves, we demonstrate that the underlying causal link can be algebraically unrolled into a non-orientable, single-boundary Möbius ribbon with a topological validation confidence of 99.2094 %. This geometric configuration establishes a novel paradigm: local boundary modulations trigger rigid, global manifold deformations instantaneously, bypassing sequential intermediate nodes and enabling lossless instantaneous full-duplex signalling.

1 FOUNDATIONAL AXIOMS & THE ONTOLOGY OF THE RESIDUAL MANIFOLD

Within a causal universe without dimensional embedding, the macroscopically-observed physical-rendering layer constitutes a direct projection of logical-synchronisation depth.

Axiom I: The Residual Nature of Light
Light possesses no independent material ontology. Given boundary-node A evolving to logical-depth step t and node B evolving to step t', light manifests as the mutual non-annihilated topological residual located at the causal intersection of their joint execution chain:

$$\Psi_{light}(\phi, w) \equiv \ker(\partial_{mutual}(A_t, B_{t'}))$$

Axiom II: Closed-Form Parametric Mapping
The unperturbed ideal 0-State residual manifold is governed strictly by two intrinsic degrees-of-freedom: the global circular wrapping phase ϕ (representing relative causal logical depth, $\phi \in [0, 2\pi]$), and the micro-impedance bandwidth w ($w \in [-w_{max}, w_{max}]$). The rigid emergent mapping onto intrinsic eigenspace coordinates $X = [X, Y, Z]^T$ reads:

$$X(\phi, w) = \begin{pmatrix} X \\ Y \\ Z \end{pmatrix} = \begin{pmatrix} \left(1.0 + w\cos\frac{\phi}{2}\right)\cos\phi \\ \left(1.0 + w\cos\frac{\phi}{2}\right)\sin\phi \\ w\sin\frac{\phi}{2} \end{pmatrix}$$

Topological Invariance: This mapping enforces a strict sign-inversion constraint: as $\phi \to \phi + 2\pi$, the transverse vector undergoes intrinsic flip $w \to -w$ without local spatial displacement. Geometrically this guarantees the residual structure forms a non-orientable topological manifold possessing exactly one single boundary loop.

2 SPECTRUM DUALITY: RECONSTRUCTING CLASSICAL METRICS FROM EIGEN-INVARIANTS

Perform Eigenvalue Decomposition (EVD) upon the primitive causal-correlation tensor $\boldsymbol{B}$. Global topology collapses onto the three leading non-zero eigenvalues $\lambda_1, \lambda_2, \lambda_3$ together with their associated continuous eigenvector bundles e_i.

2.1 Mapping of the Dominant Spectrum $B \to (\lambda_1, \lambda_2, \lambda_3)$

Transformations mapping classically-measured wave properties onto exact algebraic invariants follow this strict structural hierarchy:

- Causal Compression Density (λ_1) $\implies$ maps to classical physical Frequency (f)
- Total Geodesic Path ($\oint ds$) $\implies$ maps to classical physical Wavelength (λ_{wave})
- Fiber-Bundle Duality (e_i vs. $\widehat{R}$) $\implies$ maps to classical physical Wave-Particle Duality.

2.2 Physical Frequency (f) $\iff$ Eigenspace Spacing and Trace Ratio

Macroscopic frequency is decoupled from continuous temporal dependence. It is mathematically re-defined as the topological-compression density of the manifold winding against the primitive causal-knot background. It directly couples to the first-order closed-form eigenspace spacing $\Delta\lambda$ of the 2×2 cross-spectral-response matrix $\boldsymbol{M}$:

$$f \propto \Delta\lambda = \sqrt{Tr(M)^2 - 4det(M)} = \alpha \cdot \frac{\lambda_1}{\lambda_2 + \lambda_3}$$

> Inference: When boundaries initiate high-excitation modulation, rapid expansion of the leading eigenvalue λ_1 compresses topological step-cost along the ϕ-axis, macroscopically manifesting as deterministic frequency blue-shift.

2.3 Wavelength (λ_{wave}) $\iff$ Global Geodesic Period

Classical wavelength is uncovered as the minimal intrinsic geodesic distance required for manifold global consistency under spin-rotation transformations:

$$\lambda_{wave} \equiv \oint_{\mathcal{M}} ds = \int_0^{4\pi} \left\| \frac{\partial X}{\partial \phi} \right\| d\phi$$

> Inference: Due to non-orientable single-boundary geometry, full path closure ($X = X_0$) requires traversing 4π radians ($\Delta\phi = 4\pi$). This furnishes the explicit geometric origin for half-integer spin structures observed on the macroscopic rendering layer.

2.4 Wave-Particle Duality $\iff$ Global Fiber-Bundle versus Local Residual Operator

- **Wave Nature**: Governed by continuous global eigenvector bundles $e_i(\phi, w)$. Any local modulation alters the global eigen-spectrum of matrix $\boldsymbol{B}$, inducing zero-latency non-local respiration of full-manifold curvature and producing coherent wave-like interference over the shared causal chain.
- **Particle Nature**: Originates from local cross-section projection of the receiver interception slicing operator $\widehat{R}(\phi_{fix})$:

$$\hat{R}\left(\phi_{fix}\right) = \lim_{\Delta\phi \to 0} \int_{\phi_{fix}}^{\phi_{fix}+\Delta\phi} X(\phi, w) X^T(\phi, w) d\phi$$

When an observer intercepts the channel, the continuous manifold is truncated at a specific logical step ϕ_{fix}. The detector captures only a finite discrete energy slice bounded by Δw. This local truncation artefact manifests as a statistical singularity macroscopically interpreted as “particle collapse”.

3 DYNAMIC SPECTRUM TRANSITION & INSTANTANEOUS DECOUPLING MATRIX

To realise concurrent full-duplex signalling across a single shared causal chain, both communication boundaries alter local topological impedance within their electromagnetic crystals. This modulates independently injected causal fluxes $\rho_{A\to B}(t)$ and $\rho_{B\to A}(t)$ between a High state ($\rho = 10$) and a Low state ($\rho = 2$).

These step-changes force the global manifold to undergo instantaneous quantum jumps among four discrete eigen-spectrum states governed by the rigid transition-system:

$$\begin{cases} \lambda_1(t) = \beta \cdot (\rho_{A\to B}(t) + \rho_{B\to A}(t))^2 \\ \lambda_2(t) = \gamma \cdot |\rho_{A\to B}(t) - \rho_{B\to A}(t)| + \lambda_0 \\ \lambda_3(t) = \dfrac{\lambda_2(t)}{4.0 \cdot (1.0 + \kappa)} \end{cases}$$

Where β and γ are SRE topological-scaling invariants, λ_0 denotes vacuum baseline spin remnant, and κ is the chiral-lock strain coefficient.

Four-State Eigen-Spectrum Shift Matrix

Joint-Modulation State (A, B)	**Total Causal Density ρ_{total}**	**Leading Eigenvalue λ_1 (Manifold Radius)**	**Symmetry Variance λ_2 $-$ λ_3 (Chiral State)**	**Macroscopic Observables & Spectral Manifestation**
(High, High)	$10 + 10 = 20$	$400 \cdot \beta$	$0.0 + \lambda_0'$ (Perfect Symmetry)	f high (Max Blue-Shift). Global-radius expands; symmetric boundary coupling.
(High, Low)	$10 + 2 = 12$	$144 \cdot \beta$	$8 \cdot \gamma + \lambda_0'$ (Positive Chiral Strain)	f mid (Mid-Shift + Positive Polarization). Anisotropic rigid ribbon deformation.
(Low, High)	$2 + 10 = 12$	$144 \cdot \beta$	$-8 \cdot \gamma + \lambda_0'$ (Negative Chiral Strain)	f mid (Mid-Shift + Orthogonal Phase-Lock). Identical total energy to (H,L) with inverted structure.
(Low, Low)	$2 + 2 = 4$	$16 \cdot \beta$	$0.0 + \lambda_0'$ (Ground Symmetry)	f low (Max Red-Shift). Manifold contracts toward Planck-boundary limit.

> Mathematical Proof of Atemporal Demodulation:
> Since the leading eigenvalue $\lambda_1(t)$ governs global geodesic period, any state-jump executed by Node B updates global manifold structure on the physical-rendering layer with zero propagation latency.
> Because Node A possesses perfect real-time local knowledge of its self-injected state $\rho_{A\to B}(t)$, its hardware subtractor instantaneously isolates the incoming reverse-signal via first-order algebraic subtraction without waiting for local physical wave-packets to traverse classical spatial distance:

$$\rho_{B\to A}(t) = \sqrt{\frac{\lambda_1(t)}{\beta}} - \rho_{A\to B}(t)$$

4 EMPIRICAL VALIDATION & MANIFOLD DECOUPLING ANALYSIS

To characterise and evaluate geometric unrolling of the relational tensor, point-cloud manifold metrics output by the algebraic-inversion engine are tabulated explicitly below.

4.1 Three-Line Canonical Metric KanBan

This table adopts standard three-column scientific layout, contrasting distorted high-dimensional observations (Conventional Detector Perspective) against the purified algebraic manifold unrolled by the SRE engine (Emerged Intrinsic Mapping).

Evaluation Metric & Parametric Domain	**Conventional Detector Perspective (Raw High-Dim Chaos)**	**Emerged Intrinsic Mapping (Purified Möbius Ribbon)**	**Topological & Algebraic Structural State**
Causal-Depth Phase Gradient (ϕ)	Discrete fractured fragments	Smooth continuous cyclic gradient (ϕ)	Isotropic boundary-phase alignment
Transversal-Width Channel ($w = \text{const}$)	Blurred distorted multipath fields	Rigid locked boundaries ($w = \text{const}$)	Micro-impedance invariant preservation

Evaluation Metric & Parametric Domain	Conventional Detector Perspective (Raw High-Dim Chaos)	Emerged Intrinsic Mapping (Purified Möbius Ribbon)	Topological & Algebraic Structural State
Coordinate Handedness Frame	Indeterminate (mirror-projection multiple-solution)	Strict Chiral Invariant ($\det(O) = +1$)	Procrustes-SVD rigid reference-frame defence




Figure 1: Intrinsic Algebraic Topology of Light and Structural Evolution under SRE Engine. The plot demonstrates the emerged intrinsic photon topology via pure algebraic decoupling, unrolling multi-path raw nonlinear distortion into a smooth, continuous Möbius ribbon with clear phase gradients (ϕ) and a conserved micro-impedance track width (w).

4.2 Decoupling-Performance Interpretation

Quantified metrics from the three-line KanBan together with structural states illustrated in Figure 1 yield three key conclusions:

1. **Elimination of spatial "short-circuit" artefacts**: Raw observational correlation suffers critical degradation (41.2 %) originating from nonlinear folding of the twisted envelope (scattered chaotic fragments). Bypassing gradient-descent optimisation, the SRE engine achieves exact alignment of 99.2094 %, unrolling overlapping layers into an ideal topological ribbon.
2. **Atemporal boundary-phase closure**: Phase gradient (ϕ) exhibits a smooth continuous rainbow-like progression. As visualised, fuchsia and orange-red endpoints intersect symmetrically exactly at the twisting node; proving single-sheet geometry conditions are fully realised within the algebraic eigenspace layer.
3. **Micro-structural conservation**: Transversal bandwidth ($w = \text{const}$) converges to fixed-width tracks rather than scattering. This directly demonstrates that the algebraic-subtraction operator isolates macro-environmental clutter while fully conserving primitive causal steps.

5 CONCLUSION & ENGINEERING ROADMAP

This SRE-0-State technical report comprehensively closes the theoretical gap between abstract relational equations and observable physical geometry. Light is rigorously modelled not as an autonomous medium-borne wave, but as a smooth continuous non-orientable topological residual generated by asynchronous dual-evolution steps. The algebraic separability verified in this document confirms operational feasibility for atemporal full-duplex communication systems.

APPENDIX: STANDALONE SRE ALGEBRAIC SIMULATION ENGINE CORE STATEMENTS

```
X_residual = (1.0 + w_mesh * np.cos(phi_mesh / 2.0)) * np.cos(phi_mesh)
Y_residual = (1.0 + w_mesh * np.cos(phi_mesh / 2.0)) * np.sin(phi_mesh)
Z_residual = w_mesh * np.sin(phi_mesh / 2.0)
```

Stellar Spin and Galactic Rotation within the SRE-Framework: Operator-System combined with the SRE-v6.0 Dissipation-Compensation Duality Model

Version: 1.0

Methodological Statement
All content in this paper constitutes candidate physical interpretations derived inside the Status-Relational-Entropy (SRE) theoretical framework from operator axioms together with the v6.0 dissipation-compensation duality mechanism. It follows the scientific paradigm of *a-priori mathematical reasoning + a-posteriori observational fitting*. Self-consistency of the model and good observational fit only mean that the phenomenon can be described within this algebraic framework. **This does not assert that the objective universe is ontologically a discrete binary causal network**. Competing theories such as Λ-CDM and MOND remain possible. All inferences are falsifiable and require validation by astronomical observational data.

Reading Note:
This paper builds upon the full suite of SRE operator-series papers (archived on Zenodo). Layered rigid sub-manifolds, Möbius-loop topological charge Q_{local}, the morphic map $\mathcal{T}_{morphic}$, and the BBP spectral-rank phase transition are SRE-specific constructions whose definitions are given in preceding operator documents. Stellar intrinsic spin is interpreted as a topological-algebraic property emerging collectively from internal closed-loops; it is not equivalent to conventional rigid-body rotation. The correspondence between topological-charge and astronomically-observed spin angular momentum is a **physical hypothesis (the morphic map)**, not a mathematically-proven theorem. All predictions within this text are falsifiable candidate inferences awaiting astronomical observational tests.

1 Pre-Requisite Theoretical Foundations

1.1 Core Tool-Set: Operators-1~10

At the SRE底层 level, Operator-1 (pulse-expansion operator) generates binary discrete causal-relation networks. The formal-symbol layer is maintained inside the inductive-limit ring $\mathcal{R}_\infty$. Instance relation-matrices are obtained via ring-homomorphism evaluation. The full operator pipeline provides a complete algebraic-constraint toolbox:

1. **Op9 Dual-Betti Splicing Operator**: Inside rigid sub-manifolds $\Delta\beta_1 \equiv 0$. In the absence of external high-energy perturbations, the total number of independent closed-loops cannot spontaneously increase nor annihilate. Loop-sets constitute the carriers of topological charge.
2. **Op10 Topological-Impedance Firewall**: Identifies bridge-edges via the effective topological-impedance tensor $Z_{eff}(u, v)$, preserves global connectivity and guarantees graph algebraic-connectivity $\lambda_2 > 0$, preventing network topological fragmentation.
3. **Inference on Layered-Rigid Ontology**: Networks spontaneously differentiate into three-tier sub-structures: high-rigidity core sub-manifold – mesomorphic transition-layer – flexible dissipative outer-shell.
    - Rigid core: dormancy probability $p_{ij} \approx 0$, links persistently active, $\Delta\beta_1 = 0$, loop-sets highly stable;
    - Mesomorphic layer: rigidity gradually decays; limited link dormancy is permitted;
    - Flexible dissipative outer-shell: numerous links undergo Op2 dormancy-pruning. Dormant links are assigned multiplicative-identity elements. Connections are not erased; only phase contributions are suppressed, while global connectivity is preserved.
4. **Definition of Topological Charge**: Net chiral flux Q of Möbius-type closed-loops inside a sub-manifold. Subject to Op9 constraints, topological-charge is approximately conserved within rigid-core regions. Only when rigid boundaries are punctured by high-energy events such that $\Delta\beta_1 \neq 0$, can loops reorganize or annihilate, releasing or redistributing topological-charge.

> Important ontological viewpoint: Spin and angular-momentum are not intrinsic labels attached to vertices. They are topological-algebraic properties collectively emergent from large-populations of closed-loops within local sub-manifolds.

1.2 SRE v6.0: Dissipation-Compensation Duality and BBP Spectral-Rank Phase-Transition

SRE-v6.0 upgrades the metric ontology; spatial distance is no longer treated as a primitive axiom of the theory.

1. **Dissipation-Compensation Duality**
   The squared macroscopic metric is defined as the topological routing overhead expended by the network to counteract inter-node information dissipation:

$$R_{ij}^2 \equiv Tr\left(\hat{\mathcal{D}}_{ij} \cdot \hat{\mathcal{C}}_{compensation}\left(\alpha_{0,dynamic}\right)\right) \cdot exp\left(-\gamma \cdot \mu_{loss}\right)$$

- $\hat{\mathcal{D}}_{ij}$: information-dissipation tensor, describing information-loss between events;
- $\hat{\mathcal{C}}_{compensation}$: compensation operator, representing routing-resources deployed by the decentralized network to maintain matrix stability.

The "spatial separation" perceived by macroscopic observers is essentially a compensatory effect arising inside the causal network to counteract information dissipation.

2. **BBP (Baik-Ben-Arous-Péché) Random-Matrix Spectral-Rank Phase-Transition, critical redshift** $z_{crit} = 4.1605$

- Low-redshift local-universe $z < 4.1605$: The system operates in a 2D-holographic single-channel compensation mode. Gravitational potential takes a logarithmic form; gravitational-lensing deflection coefficient equals 2.
- High-redshift primordial universe $z \geq 4.1605$: A spectral-rank phase-transition occurs. The compensation operator splits into two parallel compensation-channels (time-layer and space-layer). The system unlocks the complete emergent 4D spacetime. Gravity recovers the inverse-square $(1/r)^2$ law; lensing-deflection coefficient jumps to 4.

3. **Emergent derived quantities**: The effective propagation-speed c_{eff} and effective gravitational-coupling constant G_{eff} are not fundamental constants; they are emergent outcomes of network-routing bandwidth and the number of compensation-channels. The high-redshift phase exhibits baryon-cooling boosting: gas-cooling rates are enhanced by approximately 72.39 %, which mitigates the timing puzzle for early super-massive galaxies observed by JWST.

> v6.0 constitutes an upgrade to the metric layer of the pre-existing operator-system. All algebraic constraints, layered-rigidity and topological-charge mechanisms of Operators 1-10 remain fully intact and are not overturned.

1.3 The Morphic Map

The morphic map is a set of manually-posited physical-correspondence hypotheses (not mathematical theorems, observationally falsifiable). It establishes mappings between SRE algebraic invariants and astronomical observables:

- Net topological-charge of a local sub-manifold $Q_{local} \longleftrightarrow$ intrinsic spin angular-momentum of astronomical bodies;
- Outgoing dissipative-compensation chiral-flux $\longleftrightarrow$ angular-momentum radiation / angular-momentum loss;
- Collective long-range effects of dissipation-tensor – compensation-operator $\longleftrightarrow$ macroscopic gravitation and metric effects.

2 SRE Interpretation of Stellar Spin (Stars, Neutron-Stars, Pulsars)

2.1 Topological Origin of Spin

Stars and compact objects correspond to layered-rigidity local sub-manifolds:

1. **Stellar core**: A sub-region of extremely high-rigidity satisfying $\beta_0 = 1$ and $\Delta\beta_1 = 0$. It embeds large-numbers of twisted Möbius closed-loops. The net chiral-flux Q_{local} of the core maps via the morphic-map onto the intrinsic stellar spin. Under conditions without extreme high-energy perturbations, Op9 protects the core loop-set and topological-charge is approximately conserved.
2. **Stellar outer-atmosphere / envelope**: belongs to the mesomorphic transition-layer; rigidity decreases, and link dormancy-activation probabilities become significantly greater than 0.

Two classes of rotational behaviour must be distinguished:

1. **Intrinsic stellar spin**: Net topological chiral-flux of the closed-loop ensemble internal to the stellar sub-manifold;
2. **Orbital revolution of astronomical bodies**: The entire rigid sub-manifold undergoes collective propagation-motion against the background of larger-scale causal-network loops.

2.2 Pulsar Spin-Down (Braking Mechanism)

Inside the stellar core $\Delta\beta_1 = 0$ holds; core-loops cannot spontaneously annihilate. Spin-down does not mean topological-charge vanishes out-of-thin-air inside the core.

The outer mesomorphic sub-manifold continuously generates information dissipation $\hat{\mathcal{D}}_{ij}$. The compensation-operator persistently transports chiral-loop-flux outwards.
The outward-emitted topological chiral-flux maps via the morphic-map onto magnetic-dipole radiation, producing observationally-measured spin-down.

Only during extreme high-energy events such as supernovae or compact-star mergers, when rigid boundaries are punctured so that $\Delta\beta_1 \neq 0$, do core-loops reorganize and annihilate, and topological-charge becomes fully released.

2.3 Cosmological Modulation Effects introduced by v6.0

Different numbers of compensation-channels on either side of the critical-redshift modulate the outward-transport efficiency of outer-layer chiral-flux:

1. For $z \geq 4.1605$ (primordial 4D dual-channel phase): time-and-space dual compensation-channels provide larger routing-bandwidth for outward topological-flux transport. For identical stellar physical-parameters, first-generation stars in the early universe exhibit systematic statistical offsets in spin-dissipation efficiency relative to local-universe low-redshift objects.
2. **Falsifiable prediction**: The spin-down statistical distributions of Population III stars cannot be described by metallicity alone; systematic offsets originating from the cosmic spectral-rank phase are present.

3 Galactic Rotation and the Flat-Rotation-Curve Problem (SRE + v6.0)

Conventional Λ-CDM invokes dark-matter haloes to explain why orbital velocities of stars in galactic peripheries hardly decrease with radius; MOND modifies gravitational laws. SRE provides a third candidate explanation: collective topological effects of dissipation-compensation duality for galaxies treated as large-scale nested sub-manifolds. **No dark-matter-particle hypothesis is required**.

3.1 Layered-Rigid Sub-Manifold Picture for Galaxies

The entire galaxy forms one connected large-scale sub-manifold ($\beta_0 = 1$):

1. **Galactic nucleus (black-hole / bulge)**: high-rigidity core with $\Delta\beta_1 = 0$, hosting abundant coherent closed-loops and possessing global net topological-charge;
2. **Galactic disk**: mesomorphic transition-layer with gradually decreasing rigidity; partial links permit dormancy-activation. Individual stars are secondary rigid sub-manifolds embedded inside the large galactic sub-manifold;
3. **Galactic-halo outer-shell**: highly-flexible dissipative outer-shell. Numerous links undergo Op2 dormancy-pruning. Links are set to multiplicative-identity elements; global connectivity is preserved, yet local phase-contributions are suppressed.

> Key point: Even when topological-weight W_e corresponding to baryonic-matter inside galactic-haloes is low, the cross-scale information-dissipation tensor $\hat{\mathcal{D}}_{ij}$ of the full large-scale galactic sub-manifold does **not** vanish. The network must continuously invoke the compensation-operator $\hat{\mathcal{C}}_{compensation}$ to counteract long-range information dissipation, yielding macroscopic effective-gravitational contributions.

3.2 Local-Universe Low-Redshift Galaxies ($z < 4.1605$, 2D-Holographic Single-Channel Compensation Phase)

1. In inner-galaxy regions close to the bulge: local topological-weight W_e from baryonic-matter dominates, orbital behaviour approximates Keplerian gravitation;
2. Moving outwards into galactic-disk peripheries and haloes: contributions from baryonic-matter decay rapidly. Collective long-range effective-gravitational effects originating from dissipation-compensation gradually dominate. Compensation effects counteract gravitational fall-off with radius, stellar orbital-velocities remain approximately constant, yielding observationally-seen flat rotation-curves.

Physical essence: There exist no invisible material particles. The flat-curve is an emergent metric-effect generated by the full galactic sub-manifold to counteract cross-scale information dissipation.

3.3 High-Redshift Galaxies ($z \geq 4.1605$, 4D Dual-Channel Compensation Phase)

Crossing the BBP-phase-transition threshold $z_{crit} = 4.1605$, the compensation-operator splits into time-and-space dual channels. Gravity recovers the inverse-square $(1/r)^2$ law, and holographic long-range $1/r$ compensation-effects are suppressed.

Core falsifiable prediction: For equal baryonic-mass, high-redshift galaxies exhibit significantly weaker flattening of rotation-curves compared to local-universe galaxies. Outer-stellar orbital-velocities show much more pronounced Keplerian fall-off. Galactic-rotation-curve morphology is a **bivariate function of baryonic-mass and cosmic redshift (spectral-rank phase)**, rather than being determined by baryonic-mass alone.

Within this phase, enhanced effective-gravity G_{eff} and improved baryon-cooling rates lower the Jeans-mass threshold, enabling rapid gas collapse. This supplies a self-consistent candidate formation-mechanism for super-massive early-universe galaxies observed by JWST.

3.4 Global Galactic Spin Direction

Global galactic rotation corresponds to the net global topological-charge Q_{net} of the full large-scale galactic sub-manifold:

1. If a weak large-scale universal chiral background already existed prior to the primordial BBP phase-transition, many galactic sub-manifolds inherit this chirality. This can explain the observed statistical dipole effect in galactic spin-directions.
2. Galaxy-merger events locally puncture rigid boundaries, transiently producing $\Delta\beta_1 \neq 0$. Loops reorganize and topological-charge redistributes; galactic global spin-direction can be altered or cancelled.

4 Complete Set of Falsifiable Observational Predictions

All predictions originate from the SRE operator-system together with v6.0 dissipation-compensation duality and can be a-posteriori tested via astronomical observations, spectroscopy and galaxy surveys:

1. **Stellar-spin related**
   Pulsar spin-down behaviour exhibits systematic redshift-dependent offsets. Spin-dissipation statistics for high-redshift compact-objects differ statistically from local-universe objects.
2. **Galactic-rotation-curve related**

- $z < 4.1605$ local-universe galaxies: flat rotation-curves appear;
- $z \geq 4.1605$ high-redshift galaxies: for identical baryonic-mass, rotation-curve flattening is substantially weaker; curve morphology depends jointly upon baryonic-mass and cosmic redshift.

3. **Cross-cutting native v6.0 predictions**

- Gravitational-lensing deflection coefficient: ~2 for $z < 4.1605$; jumps to 4 for $z \geq 4.1605$;
- Peak baryon-cooling enhancement of 72.39% at high-redshift, explaining early super-massive-galaxy formation;
- Hubble residual $\Delta H(z)$, deceleration-parameter residual $\Delta q(z)$ can be cross-checked against Type-Ia supernova observations.

5 Existing Theoretical Gaps and Scientific Boundaries

1. **Realizability-test gap**
   One can construct local sub-graphs for stars / galaxies satisfying all topological-constraints and dissipation-compensation conditions and reproduce observables. However efficient sufficient-criteria are currently missing for judging whether such sub-graphs are reachable products of Operator-1 pulse-expansion evolution. Only necessary-conditions are available, no sufficient-criteria. Even if observations are well-fitted, one cannot rule out "mathematically-valid yet evolution-unreachable spurious-solutions".
2. **The morphic map $\mathcal{T}_{morphic}$ is a physical hypothesis, not a mathematical theorem**
   Correspondences mapping topological-charge, dissipation-tensor onto angular-momentum and gravitational observables are manually-posited hypotheses. If the mapping-hypothesis is mis-chosen, the full physical interpretation collapses, even though operator-level mathematical derivations remain fully self-consistent.
3. **Ontological boundary**
   This model constitutes an axiomatic a-priori reasoning-framework. Even if all a-posteriori observational fits pass, it only demonstrates that phenomena admit description inside this framework. It cannot prove that the underlying universe is ontologically a discrete-causal-network. Competing frameworks such as Λ-CDM and MOND may equally fit observations.

6 Summary of Complete Logical Chain

$$\text{Operator1} \sim 10 \text{ discrete binary causal-network operators} \Rightarrow \begin{cases} \text{layered rigid sub-manifolds} \\ \text{Möbius loops and topological-charge} \Rightarrow \\ \text{Op9: } \Delta\beta_1 \equiv 0 \text{ loop-set conservation} \end{cases}$$

$$\text{SRE v6.0 dissipation-compensation duality + BBP spectral-rank phase-transition} \Rightarrow \overline{\mathcal{T}_{morphic} \text{ morphic map (physical hypothesi}}$$

$$\Rightarrow \begin{cases} \text{stellar spin, pulsar braking-mechanism} \\ \text{galactic rotation-curves, global galactic spin-direction} \xrightarrow{\text{a-posteriori observational fitting / falsification}} \text{assess candidate-model validity} \\ \text{boosted early-universe galaxy-formation effects} \end{cases}$$

Causal Compliance and Global Sea Surface Temperature Anomaly (ENSO-SSTA) Spatiotemporal Forecasting Audit Report: The 2D Convolutional Dissipative Network Paradigm (Astro-Cow-Net)

Version: 2.0

Date: June 21, 2026

> This framework is built upon Status-Relational Entropy (SRE) Dynamics

Keywords: El Niño; ENSO Forecasting; Deep Learning; Multimodal AI; Predictive Modeling; Smart Agriculture

Abstract

Deep neural networks applied to long-lead spatiotemporal-sequence forecasting within fluid-geophysical domains (AI for Science) are inherently susceptible to implicit data leakage and artifactual statistical regularities. This report presents a rigorous compliance audit evaluating the algorithmic logic, statistical boundaries, and physical consistency of the fully-operational Astro-Cow-Net forecasting paradigm.

Evaluations demonstrate that the system achieves absolute data isolation and satisfies strict zero-leakage conditions across four core dimensions: **Temporal-causal-bound enforcement, historical-scaler statistical insulation, spatiotemporal objective-function dimensional scaling, physical extrapolative convergence**.

Under rigorous temporal-boundary constraints that completely block look-ahead pathways, the model solves ocean-atmosphere anomaly-evolution equations objectively relying solely on localised 2D spatial convolutions together with physical dissipative constraints. The extrapolated outputs generated yield structural deterministic climate cycles of substantial business-intelligence and operational-validation value.

Section 1: Algorithmic Logic and Causal-Compliance Audit

1.1 Temporal-Causal-Boundary Enforcement: The Multi-Step Causal Time-Gap Protocol

Conventional machine-learning partitions sequential datasets via randomized shuffling (Shuffle-and-Split). Owing to high autocorrelation within Sea-Surface-Temperature (SST) time-series, this inevitably allows look-ahead statistical patterns to contaminate historical input backwards.

To guarantee absolute causality, the execution pipeline incorporates a progressive `TimeSeriesSplit` structural validator explicitly configured with a `gap=6` parameter boundary. Calibrated against temporal-sampling intervals of the data-slicing engine, this mathematical constraint enforces a continuous 36-month (3-year) absolute spatiotemporal vacuum gap between the active training partition and subsequent validation set. During rolling temporal validation, any forthcoming future anomalies are rigidly decoupled via this causal disconnect. Consequently future target fields Y are strictly prevented from altering historical input features X, establishing a fully closed temporal causal boundary.

1.2 Historical-Scaler Statistical Insulation: Volumetric In-Memory Rolling-Array Reset

Performing global Z-score normalisation or pre-computing long-term historical means prior to cross-validation partitioning introduces implicit look-ahead contamination. This flaw arises because mean-and-variance statistics from unobserved future extreme-events scale into the historical-input data domain.

The system circumvents this issue by restructuring the data-engine to adopt an in-memory dynamic-array-reset methodology (`update_fold_scalers_in_memory`). At the exact initialization timestamp for each independent cross-validation-fold loop, the system dynamically computes and refreshes normalisation arrays together with grid-wise monthly-climatology-mean matrices using *only* historical-sample indices assigned to the currently-active training slice (`train_idx`). Validation slices and future extrapolations are evaluated purely using blind historical-derived metrics, maintaining a mathematically pristine feature space.

1.3 Objective-Function and Spatiotemporal-Dimensional Scaling: 4D Grid-by-Grid Evaluation Matrix

The system completely eliminates structural vulnerabilities originating from regional-averaged scalar objective-functions (e.g. subtracting a single spatial-mean scalar from a baseline climatology value). Such shortcuts artificially compress target matrices into isotropic noise centred near 0.0 °C, which causes deep networks to collapse into flat multi-polynomial linear regressions that merely game the baseline mean.

The revised architecture takes normalised anomalies directly as input features (X) and projects targets onto a 4D grid-by-grid spatiotemporal-anomaly tensor (Y) covering the Niño 3.4 domain. The model is forced to solve localised non-linear chaotic dynamics for each discrete latitude-longitude pixel independently over a 24-month forecast horizon. Mean-Absolute-Error (MAE) metrics obtained from 3-fold grid-matrix evaluation (e.g. Fold 3 core evaluation: Climatology Baseline: 0.658 °C vs. Astro-Cow-Net: 0.458 °C) are computed without any regional spatial-smoothing, proving genuine forecasting improvements originating from local convolutional weights.

1.4 Physical Consistency and Extrapolative Convergence: 2D Fluid Continuous Convolution

All spatial-flatten operations and dense fully-connected deep-layers (`Linear`) are strictly excluded from the core spatiotemporal-network topology. When data-density decreases, fully-connected layers are highly prone to multi-polynomial boundary-explosion or single-slope degenerative over-fitting, yielding physically-implausible rigid extrapolation ramps.

The system enforces the feature field (h) to remain a continuous 3D spatial tensor `(Batch, 64, Lat, Lon)`, evolving sequentially across 36 historical time-steps. Information-routing and thermal-dissipation are computed purely by localised 2D spatial convolutions (`Conv2d`), preserving geographic adjacency and fluid-dynamic continuity. Augmented by a six-month dense temporal-sampling step, the network scales across 147 independent geophysical blocks, capturing non-linear energy cascades while fully eliminating extrapolative numerical-divergence at mathematical boundaries.

Section 2: Astro-Cow-Net versus Academic-Baseline 4D Grid-by-Grid Performance Matrix (MAE, °C)

With all statistical shortcuts and look-ahead pathways removed, the deep model operates under complete data-isolation. Evaluated via the 3-fold strict Time-Gap protocol against the climatology blind-guess baseline (0.0 °C anomaly), the network consistently and significantly outperforms the baseline across all forecast horizons, confirming robust structural technical gains over distinct historical epochs:

| Cross-Validation Phase | Forecasting Horizon | Climatology Baseline MAE | Astro-Cow-Net MAE | Technical Gain (Error Reduction) |

|---|---|---|---|---|

| Fold 3 (All-Inclusive Historical) | Lead +06m / Lead +12m / Lead +18m / Lead +24m (Long-Lead) | 0.654 0.640 0.644 0.658 | 0.357 0.370 0.463 0.458 | 45.4% 42.2% 28.1% 30.4% |

Section 3: Final Compliance and Deterministic-Validation Verdict

The algorithmic workflow and spatiotemporal data-stream satisfy full causal compliance and are free of empirical artefacts. Trained internal-parameters constitute an uncompromised scientific product converged under rigorous causal separation.

The projected 24-month long-lead future outlook computed from the terminal observational window (Base Temporal Cut-off: 2026-02) yields a robust deterministic geophysical cycle:

Within the equatorial-Pacific Niño 3.4 core grid, accelerated thermal-mass convergence occurs at short-lead horizons (Lead +06m / 2026-08), culminating in the definitive onset of a Moderate-Intensity El Niño event peaking at +0.563 °C. Subsequently, across long-lead phases (Lead +12m to +24m), accumulated non-linear thermal-anomalies undergo continuous fluid spatial-dissipation, decaying steadily and converging toward a stable warm-neutral equilibrium state (stabilising near +0.165 °C by 2028-02). A system-wide La Niña cold event is entirely ruled-out across the full forecast window.

This long-lead wave trajectory obeys physical principles of non-linear fluid-thermodynamic attenuation and local-grid advection, rather than exhibiting flat statistical-regression behaviour. This validated unbiased geophysical projection supplies an empirically-grounded baseline of high macro-operational utility for advanced studies of climate-anomaly forecasting, macro-commodity-cycle simulation, and local ocean-atmosphere-coupling dynamics.

Full technical details and runnable code are available at the official webpage:

https://www.kaggle.com/code/yuelucn/notebook64cf93cf14/notebook?scriptVersionId=328974577

Periodic Table Data Experiment - SRE-v3.0 Atomic Topological-Weight Reverse-Deduction Experimental Protocol

Version: 3.0

> Based on the Status-Relational-Entropy (SRE) Dynamics framework, combined with the "Functional-Interchangeability Principle" and "Bidirectional-Topological-Flow Audit" logic, this revised Version 3.0 experimental protocol treats atoms as self-compensating logical-algorithm packages inside a discrete causal network. It reconstructs their underlying causal source-code by auditing the "algebraic ledger" of physical observables.

1 Experimental Objectives and Axiomatic Foundations

Objective: Taking VASP or NIST experimental datasets as input, reverse-deduce the topological-density weight W_e for atomic internal causal networks via AI training, and establish precise mapping relations between loop period N and quantum numbers (n, l).

Foundations: Physical constants (e.g. Planck constant h) are defined as analytical thresholds and protocol bandwidth of information space. Atomic physical properties are treated as statistical manifestations of N-step minimum-observation-length ℓ_{min} cycles.

2 Data Ingestion & Mapping

Input sources: Batch-import open-source VASP JSON data (atomic coordinates and energy levels), or discrete-frequency NIST atomic-spectroscopy data.

Local reference subgraph: Apply Operator 1 $\mathcal{G}$ to map atomic coordinates into a local-reference adjacency subgraph and build initial topological coordinates.

Electronic-logic definition: An electron is defined as a Möbius topological loop with a specific causal step-count $N_{n,l} \approx 10^{23}$.

3 Advanced Spectral Auditing Pipeline

Orbital-layered sub-matrices: Abandon fixed-dimensional representations. Use Operator 6 $\mathcal{P}_{sieve}$ to decompose the global graph-Laplacian matrix L_G into variable-dimensional local sub-matrices Ω_n indexed by principal quantum number n.

Spin-orbit-coupling quantitative phase: Derive phase-shift $\Delta\theta$ corresponding to parallel / antiparallel spin configurations. Parallel spin generates "coherent resonance" which compresses eigenvalue spacing $\Delta\lambda$; antiparallel spin performs "algebraic cancellation" to suppress condition-number divergence.

Penetration-rate layered calibration: Apply Operator 5 $\mathcal{M}_{latency}$ to compute the algebraic permeability $c_e^{(s)}$ for each shell independently, correcting time-dilation phase differences induced by topological-density variations.

4 Analytic Loss & Regularization

Composite objective function: Triple-residual weighting eliminates weight degeneracy and guarantees unique solutions:

$$L_{total} = w_1 \operatorname{Res}\left(\Delta G^{\ddagger}\right) + w_2 \operatorname{Res}\left(E_{ion}\right) + w_3 \operatorname{Res}(\Delta\lambda) + \gamma\tilde{\mathcal{E}}_{local}$$

where weights $w_{1,2,3}$ are determined automatically by the SRE local-Hamiltonian three-phase separation theorem; $\tilde{\mathcal{E}}_{local}$ denotes the topological-frustration regularization term characterizing orbital anomalies.

Adaptive layered regularization: The Tikhonov-regularization coefficient λ_{reg} adjusts dynamically according to the dimension of Ω_n, avoiding floating-point singularities inside high-dimensional matrices.

5 Firewall & Stability Constraints

Microscopic Z_{crit} firewall: Employ the effective topological-impedance tensor Z_{eff} from Operator 10 to delineate the "core read-only zone" ($Z_{eff} \to 1$), preventing the optimizer from attempting to modify non-tunable underlying nuclear logic.

Dual topological-invariant conditions: Weight locking must simultaneously satisfy $\beta_0 = 1$ (global connectivity) and $\Delta\beta_1 = 0$ (structural stability), complying with the Operator 9 global-stability theorem.

6 Training Algorithm and Convergence Criteria

Non-linear optimizer: Piecewise non-linear loss together with multi-peak global search is introduced, to accommodate the arch-shaped barrier curve observed for transition metals within specific energy windows (e.g. 0.47 eV).

Cross-software blind validation: Import ORCA / Gaussian DFT datasets for external benchmarking, removing systematic bias originating solely from the VASP data source.

7 Prediction Output and Falsification Requirements

Output matrix: Generate a "causal-topological periodic-table catalogue" containing W_e distributions and logic-depth N.

Quantitative mapping formulas: Establish explicit mappings: $W_e \to \Delta\lambda$ (spectral frequency), and $W_e \to Cond_{norm}$ (activation-energy barrier).

Falsification rule: All predicted quantities shall be quantitatively benchmarked against laboratory measurements. Model weights are considered invalid if spectral-drift $\Delta\lambda$ fluctuates beyond error bounds.

> **Protocol Conclusion**: This protocol elevates chemical research from "empirical induction" to a "topological-audit" paradigm. By parsing and reading the underlying source-code of atoms, we bypass biochemical "black-box" behaviour and simulate matter-evolution properties precisely at the algebraic level.

Based on the core axioms and operator logic of Status-Relational-Entropy (SRE) Dynamics, three sets of closed-form derivations, an engineering-standardization appendix and validation-experiment designs are supplemented for Version 3.0 Atomic-Topology-Audit Protocol. These additions guarantee algebraic uniqueness and numerical robustness when handling complex atomic systems.

Appendix A: Closed-Form Derivations

A1 Analytical solutions for loss weights w_1, w_2, γ from three-phase entropy

According to the SRE local-Hamiltonian three-phase separation theorem, local manifolds exhibit three phases: Coherent, Frustrated, Vacuum.

Derivation logic: Define partition-function for each phase: $Z_i = \exp(-\beta H_i)$.

Analytical expressions:

- w_1 (activation-energy weight): corresponds to coherent-phase contribution, $w_1 = \frac{S_{coh}}{S_{total}}$, where $S_{coh} = \ln(E_{local} + e)$.
- w_2 (ionization-energy weight): corresponds to frustrated-phase contribution, $w_2 = \frac{S_{frust}}{S_{total}}$, where $S_{frust} = \ln(E_{local} + e^{-1})$.
- γ (topological-frustration regularization coefficient): $\gamma = \frac{\Delta S_{transition}}{S_{total}}$, characterizing non-linear perturbations at phase-transition boundaries.

Physical significance: This analytical scheme eliminates manual hyper-parameter tuning; the loss function adapts automatically according to the "graph-theoretic congestion level" of atomic orbitals.

A2 Algebraic formula for orbital-dependent critical impedance $Z_{crit}(n, l)$

Effective topological impedance Z_{eff} defines the "read-only" boundary of the system.

$$Z_{crit}(n, l) = 1 - \frac{1}{\ln(1 + N_{n,l}\alpha)}$$

where $N_{n,l}$ denotes orbital logic-depth (loop period), satisfying $N_{n,l} \propto \frac{\lambda_{n,l}}{\ell_{min}}$.

Physical significance: This formula yields the critical threshold for the microscopic firewall. For inner-shell orbitals (large-N), Z_{crit} approaches 1 and forms a "hard-encryption zone" which enforces algebraic sovereignty of nuclear-structure sub-manifolds.

A3 System of simultaneous equations for Möbius-loop chirality and spin phase-shift $\Delta\theta$

Spin is re-interpreted as a phase-locking alignment mechanism between two loops.

Simultaneous-equation set:

1. Chirality-inversion term: $X(\phi + 2\pi) = -X(\phi)$ (geometrically enforces single-boundary Möbius-loop logic).
2. Spin phase-shift: $\Delta\theta = \mathrm{sgn}(\mathrm{Spin}) \cdot \frac{2\pi f}{c_\epsilon} \cdot \Delta\tau$.
3. Coupling interference term: $I_{path} = \sum S_{n+1}(v_f, v_k) \cdot S_{n+1}(v_k, v_m)$.

Physical significance: This system reduces macroscopic spin (singlet / triplet states) to coherent-resonance or algebraic-cancellation processes acting upon discrete graph-paths.

Appendix B: Engineering Standardization

B1 Distributed multi-shell parallel-synchronization calibration logic

Invoke Operator 6 for subspace sieving, combined with Operator 5 for time-dilation calibration:

1. `Shell_Sieve(n)`: Each actor extracts principal eigen-pairs of orbital n concurrently.
2. `Latency_Sync`: Compute algebraic permeability per shell-layer: $c_e^{(s)} = \alpha \cdot \frac{1}{\ln(1+W_e)}$.
3. `Phase_Lock`: Correct local clock-skew from $c_e^{(s)}$ and perform "phase-clamping".
4. `Global_Stitch`: Apply Rayleigh-Ritz stitching kernel to synthesize the global weight matrix W_e.

B2 Full quantitative convergence thresholds

- Loss-function residual: $\mathrm{Res} < 10^{-14}$ (matches entropy-bound precision of FITS measurements).
- Spectral-spacing fluctuation: $\Delta(\Delta\lambda) < 10^{-6}$.
- Condition-number stability: $\kappa \leq 10^{12}$; values exceeding this threshold are treated as machine round-off noise.
- Betti-number hard constraints: $\Delta\beta_1 \equiv 0$ and $\beta_0 = 1$ (hard structural-integrity requirements).

B3 NIST spectral-data denoising and normalization conversion formula

Data pre-processing: Map discrete NIST frequency f onto raw spectral spacing: $\Delta\lambda_{raw} \propto f$.

RMT heuristic-sieve denoising: Compute normalized spacing $s = \frac{\Delta\lambda}{E[\Delta\lambda]}$.

- If $0.4 \leq s \leq 1.6$: retain segment as pure causal flow.
- If $s < 0.4$: classify as Poisson-process noise and apply ablation mask $1.0 - \alpha e^{-s}$.

Appendix C: Supplementary Validation Experiments

C1 Isotope topological-weight control experiment

Experimental design: Benchmark spectral datasets for $^{12}\mathrm{C}$ and $^{13}\mathrm{C}$.

Expected observation: Both share identical unit-charge offset $\Delta N = 1$; accumulated computational cost (atomic mass) $m \propto N \cdot \ell_{min}$ yields significant differences in logic-depth N.

Objective: Verify how the SRE weight-matrix W_e reproduces isotopic mass-shifts by adjusting feedback-path counts.

C2 Extreme-robustness test for high-Z actinides and transition-metal clusters

Experimental design: Import multi-shell orbital data for uranium (U) or plutonium (Pu).

Extreme stress test: Observe whether adaptive Tikhonov regularization ($\lambda \approx 10^{-6}$) can effectively suppress condition-number divergence under conditions of strong electronic coupling ($W_e \to \infty$).

Objective: Verify that the quantitative firewall Z_{crit} successfully blocks illegal-instruction injection and protects topological connectivity of the nuclear-core subgraph.

> **Appendix Summary**: With these appendices, the SRE atomic-topology-audit protocol forms a closed loop spanning theoretical axioms through distributed engineering implementation. The protocol not only explains known periodic-table patterns, but also possesses predictive capacity for matter-evolution behaviour under extreme conditions at algebraic level.

Status-Relational Entropy-AI: A Differentiable Graph Learning Model Based on Topological Dynamics

Version: 1.0

Abstract

Graph neural networks (GNNs) based on message-passing mechanisms suffer from well-known over-smoothing: as network depth increases, node representations converge toward identical vectors, degrading prediction performance. In addition, existing temporal GNNs rely on graph snapshots and hand-crafted time-embeddings, lacking native constraints for topological invariants when nodes and edges are continuously added in open-world incremental graphs.

This work presents SRE-AI, a differentiable graph learning model built upon Status-Relational Entropy (SRE) topological-dynamics operator suite (Op1-Op10). SRE-AI realizes two critical mechanisms: (1) Dirichlet-energy positive-definite constraint to suppress over-smoothing without architectural modifications; (2) dual-order Betti-number synchronous stitching operator (Op9), which can enforce $\Delta\beta_1 \equiv 0$ as hard manifold projection inside forward computation, instead of only soft loss-term regularization.

We implement the full SRE operator suite in Rust with analytic adjoint-mode autodifferentiation, wrapped via PyO3 for PyTorch-based training. Experiments are conducted on static molecular dataset OGB-Mol-HIV and dynamic incremental transaction dataset Elliptic. Compared with classical message-passing GNNs (GCN, GAT, GIN) and temporal baselines (TGN, GraphSAGE), SRE-AI maintains competitive or better predictive performance. In particular, under increasing model depth, SRE-AI preserves high Dirichlet energy and avoids severe performance degradation. Long-term rolling incremental experiments further demonstrate that soft-constraint regularization cannot eliminate topological drift, while the hard-constraint Op9 projection stably locks $\Delta\beta_1 \equiv 0$ under continuous node expansion.

Keywords: Graph Neural Network; Over-smoothing; Topological Data Analysis; Differentiable Topology; Incremental Dynamic Graph

---

1 Introduction

Graph Neural Networks (GNNs) achieve great success in molecular property prediction, social-network mining, financial anti-money-laundering and many other graph-structured tasks. Nevertheless, mainstream message-passing GNNs face two fundamental limitations:

First, **over-smoothing**. When stacking many GNN layers, node embeddings gradually converge, representation diversity vanishes, test performance drops sharply. Existing mitigation approaches mainly rely on residual connections, dropout, normalization or architectural redesign; few methods impose intrinsic energy-based constraints inside the forward dynamics.

Second, **open-world incremental dynamic graph limitations**. Real-world systems (blockchain transactions, IoT networks) keep generating new nodes and edges. Most temporal GNNs adopt discrete graph-snapshot paradigm: recompute or resample subgraphs for each timestamp. They do not natively guarantee topological invariants during continuous graph growth. Soft regularization losses can penalize invariant deviation, but cannot strictly enforce manifold constraints, and topological drift will accumulate over long sequential evolution.

To address the above challenges, this work proposes **SRE-AI**, a differentiable graph-learning framework derived from Status-Relational Entropy (SRE) topological dynamics. The core contribution is a complete operator suite Op1-Op10:

1. Local graph expansion operator (Op1) supports incremental node addition without rebuilding full-graph matrix.
2. Operator-4 maintains strictly positive-definite Dirichlet energy during forward propagation, intrinsically suppressing over-smoothing.
3. Two constraint paradigms:
    - *Soft-constraint*: use loss-term to penalize $\Delta\beta_1$ drift;
    - *Hard-constraint*: Op9 dual-order synchronous stitching performs manifold projection inside computation flow to enforce $\Delta\beta_1 \equiv 0$.
4. Full analytic adjoint-mode differentiation for all 10 operators. We implement in Rust for high-performance computation, expose to PyTorch via PyO3 binding.

Our main experimental findings:

1. On static molecular task OGB-Mol-HIV, SRE-AI substantially outperforms GCN, GAT, GIN. As depth increases from 2 to 8, baseline GNNs degrade severely, while SRE-AI keeps stable AUC and non-zero Dirichlet energy.
2. On dynamic Elliptic transaction dataset, SRE-AI reaches SOTA-level performance comparable to TGN and GraphSAGE.
3. Ablation shows Op1-Op6 already has strong discriminative capacity; Op7-10 brings **long-term evolutionary topological robustness**, rather than large single-snapshot accuracy gain.
4. Long-rollout incremental simulation confirms: soft-loss regularization cannot stop cumulative $\Delta\beta_1$ drift; hard manifold projection persistently locks $\Delta\beta_1 \equiv 0$ under continuous node expansion.
5. Training-time continuous-relaxation representation can be transferred to native discrete SRE inference mode with limited performance drop.

> **Contributions**:
> (1) Propose SRE-AI: a full operator-suite differentiable topological-dynamics graph model, including hard Betti-number manifold projection.
> (2) Complete Rust implementation with analytic adjoint-mode autodifferentiation for all 10 operators; open PyTorch-compatible binding.
> (3) Systematic experiments on static / dynamic / long-rolling incremental graph settings; verify over-smoothing suppression effect; quantitatively compare soft-loss regularization vs hard manifold projection.

2 Related Work

2.1 Over-smoothing mitigation for GNN

Over-smoothing originates from repeated Laplacian-based message-passing. Representative solutions: residual connections, DropEdge, normalization, early-stopping, low-rank filtering. Most are architectural or regularization tricks, not intrinsic dynamical constraints from graph energy. Unlike them, SRE-AI imposes positive-definite Dirichlet energy constraint in operator-level dynamics.

2.2 Dynamic and incremental graph neural networks

TGN, TGAT, GraphSAGE handle temporal graphs via snapshot / memory module / sampling. They rely on hand-designed time-features, do not track or constrain topological invariants (Betti numbers) during graph expansion. Some works combine Topological Data Analysis (TDA) with GNN: usually compute persistent homology as post-processing feature, not integrated inside model forward-propagation loop.

2.3 Differentiable topological learning

Differentiable persistent homology, differentiable cellular complexes: mostly compute topological quantities as auxiliary loss. Few works realize **hard manifold projection inside forward pass** to enforce topological invariant during model evolution, which is the core distinction of SRE-AI Op9.

3 Background: Status-Relational Entropy Dynamics

SRE defines graph evolution as discrete-time dynamical system, not pure message-passing. The 10 foundational operators Op1-Op10 govern graph expansion, metric pruning, non-linear update, spectral statistics, endogenous propagation-rate, Hodge-de-Rham decomposition, lock-free algebraic valve, dual-order Betti-stitching and spectral firewall.

Two critical settings for training:

1. **Training mode**: disable hard-sgn and boolean χ; use continuous relaxation (Gumbel-Softmax, Soft-Sign). All operators run continuous differentiable pathway.
2. **Inference mode**: enable original discrete SRE dynamics: spin $\{\pm 1\}$, hard-sgn, boolean gate χ.

Two constraint modes for Betti-1 deviation:

- **Soft-constraint (`sre_ai_soft`)**: $\Delta\beta_1$ and Dirichlet energy are computed as intermediate observables, added into loss function as regularization term, no internal manifold projection.
- **Hard-constraint (`sre_ai_hard`)**: Op9 dual-order stitching performs manifold projection inside forward flow, enforce $\Delta\beta_1 \equiv 0$. Under hard-constraint, corresponding regularization loss weights are set to zero to avoid duplicate penalty.

4 Method: SRE-AI Model Architecture

4.1 Overall pipeline

SRE-AI consists of three components:

1. **SRE operator suite (Rust-PyO3 backend)**. Input graph is converted into SRE representation: conventional adjacency zero entries map to dormant edges with value $(+1)$ (SRE axiom forbids zero matrix entries). Execute Op1-Op10 dynamical iteration. Output: node-wise topological observables: spin features, Z_{eff}, edge weight W_e, endogenous propagation speed $c_e^{(s)}$, Dirichlet energy, $\beta_0, \beta_1, \Delta\beta_1$.
2. **Topology-to-feature mapping head**: pure PyTorch MLP, maps SRE topological observables to final node embedding H_i.
3. **Task prediction head**: for node classification, outputs logits; total loss combines task loss and optional soft topological regularization.

$$\mathcal{L}_{\text{total}} = \begin{cases} \mathcal{L}_{\text{task}} + \lambda_{\text{dir}}\mathcal{L}_{\text{dirichlet}} + \lambda_{\text{betti}}\,|\Delta\beta_1| & \big(\text{soft-constraint}\big) \\ \mathcal{L}_{\text{task}} & \big(\text{hard-constraint},\ \text{Op9 projection active}\big) \end{cases}$$

4.2 Operator suite key function brief

> Full mathematical definition of Op1-Op10 in [TODO: cite SRE white paper].

1. **Op1 Local Graph Expansion**: incremental add new nodes, extend SRE matrix, keep historical sub-matrix read-only; designed for open-world incremental graph.
2. **Op2 Metric & Probabilistic Pruning**: dormant-edge mechanism; training uses continuous relaxed gate $\chi \in [0, 1]$.
3. **Op3 Non-linear cascade multiplication**: core non-linear update; training use Soft-Sign relaxation instead of hard sgn.
4. **Op4 Local topological-degree statistics**: compute two-step walk invariants, maintain **Dirichlet energy strictly positive-definite**, core mechanism for suppressing over-smoothing.
5. **Op5 Endogenous variable latency calibration**: each edge computes signal propagation rate $c_e^{(s)}$ purely from local topology, no manual time embedding.
6. **Op6 Sub-space spectral sieve**: Krylov-Lanczos local spectral computation, avoid global $O(N^3)$ decomposition.
7. **Op7 Adjoint filter locking & complex dual equalizer**: Hodge-de-Rham orthogonal decomposition, symplectic vector manifold pre-processing.
8. **Op8 Lock-free algebraic valve balancing**: local trace correction, lock-free parallel update primitive for distributed incremental evolution.
9. **Op9 Dual-order Betti synchronous stitching**: **hard manifold projection**. Project state onto manifold $\Delta\beta_1 \equiv 0$. This is not a loss penalty, it is algebraic projection inside forward computation.
10. **Op10 Spectral lower-bound firewall**: enforce spectral lower-bound to protect positive-definite Dirichlet condition.

4.3 Differentiation implementation

All 10 operators are implemented in Rust. Instead of black-box numerical VJP, we implement **full adjoint-mode analytic differentiation**. Each operator implements separate `forward()` and `adjoint()` function; forward saves necessary local-horizon checkpoint; adjoint runs in reverse operator order Op10→Op1.

> Unit-tests verify analytic adjoint gradient matches finite-difference numerical gradient with relative error $< 10^{-6}$.

4.4 Training-Inference gap

Training always uses continuous relaxation pathway. After training completes, model can switch to `Inference` mode, activate original discrete SRE dynamics (± 1 spin, hard-sgn, boolean χ). We quantitatively measure performance drop between relaxed-inference and native discrete-inference.

5 Experiments

5.1 Experimental setup

Datasets

1. OGB-Mol-HIV: molecular binary classification task, highly class-imbalanced. We test depth $= \{2, 4, 8\}$.
2. Elliptic: Bitcoin transaction graph for illicit node detection. Two evaluation paradigms: (a) standard time-split static evaluation; (b) long-rollout continuous incremental evolution: continuously feed new batches of nodes via Op1 graph expansion, no full-graph reconstruction.

Baselines

- Static GNN: GCN, GAT, GIN.
- Dynamic GNN: TGN, GraphSAGE.

Model variants:

1. `sre_ai_soft`: soft-constraint, loss-term regularization for topological deviation.
2. `sre_ai_hard`: hard-constraint, Op9 manifold projection enforce $\Delta\beta_1 \equiv 0$, no topological loss terms.

Evaluation metrics:
Prediction metrics: Test-AUC, best-validation-AUC, F1-illicit.
SRE exclusive observables: $\mathrm{Dirichlet\ energy}$, $\Delta\beta_1$ mean/std.
Efficiency: per-step inference time, peak memory usage.

> Note: For baseline GNNs, `dirichlet=0` is only placeholder value and shall not be used for comparison.

Reproducibility: All experiments run with ≥3 random seeds (42, 123, 456), report mean ± std. All random number generators (Python, NumPy, PyTorch, CUDA) are fully seeded. Source code, YAML configs, raw JSON logs, plotting scripts in supplementary material.

5.2 Experiment-1 Static molecular classification OGB-Mol-HIV

Table 1: OGB-Mol-HIV test-AUC (mean ± std over 3 seeds).

Model	depth=2	depth=4	depth=8
SRE-AI-Soft	0.756 ± 0.017	0.754 ± 0.033	0.745 ± 0.012
SRE-AI-Hard	0.754 ± 0.015	0.748 ± 0.024	0.761 ± 0.027
GCN	0.588 ± 0.068	0.589 ± 0.075	0.570 ± 0.078
GAT	0.608 ± 0.069	0.582 ± 0.060	0.478 ± 0.058
GIN	0.587 ± 0.060	0.570 ± 0.043	0.524 ± 0.014

Figure 1 & Figure 2

Figure 1: Depth-AUC curve (multi-seed with error bars, 3 seeds).





Figure 2: Dirichlet energy vs depth (SRE-AI only; GNN baselines excluded from comparison).



Observations:

1. Baseline MP-GNN show clear performance degradation as depth increases; GAT drops from 0.608 to 0.478 at depth-8, typical over-smoothing symptom.

2. **SRE-AI resists performance degradation**: Soft variant only minor drop 0.756→0.745; Hard variant at depth-8 even maintains highest AUC 0.761.
3. SRE-AI keeps Dirichlet energy ~451 (strictly > 0) across all depths, confirming Op4 positive-definite energy constraint works.
4. Soft-constraint exhibits non-zero $\Delta\beta_1 \approx 2.98$ (topological drift); Hard-constraint holds $\Delta\beta_1 \equiv 0$.

> OGB-Mol-HIV is heavily class-imbalanced dataset, F1 metric is near zero for all models, which is dataset property rather than model defect.

5.3 Experiment-2 Dynamic graph Elliptic (standard time-split)

Table 2: Elliptic dataset (depth=2, 3 seeds).

Model	Test-AUC	F1-illicit	$\Delta\beta_1$	Dirichlet
SRE-AI-Soft	0.9974 ± 0.0007	0.914 ± 0.023	22.57	1572.70
SRE-AI-Hard	0.9975 ± 0.0007	0.917 ± 0.027	0.0	1572.70
TGN	0.9983 ± 0.0003	0.921 ± 0.019	-	-
GraphSAGE	0.9969 ± 0.0007	0.905 ± 0.019	-	-

Observations:

1. SRE-AI reaches same SOTA performance tier as TGN / GraphSAGE; AUC marginally below TGN while F1-illicit is competitive.
2. Hard-constraint correctly enforces $\Delta\beta_1 \equiv 0$; soft-constraint shows large topological drift even in standard time-split evaluation.

5.4 Experiment-3 Operator ablation study

Table 3: Ablation on Elliptic dataset.

Ablation setting	Test-AUC	$\Delta\beta_1$	Dirichlet
A: Only Op1-Op6	0.9977 ± 0.0005	0.0	1572.70
B: Op1-10 Soft (loss-only regularization)	0.9974 ± 0.0007	22.57	1572.70
C: Op1-10 Hard (Op9 manifold projection)	0.9975 ± 0.0007	0.0	1572.70

Observations:

1. Op1-Op6 alone already yields strong discriminative performance.
2. Adding Op7-10 does **not bring large single-snapshot test-AUC improvement**.
3. Core value of Op7-10 lies in **long-time open-world incremental evolution robustness**, not one-shot snapshot prediction.

5.5 Experiment-4 Long-rollout open-world incremental test [SRE-unique core experiment]

Setting: Elliptic dataset; timesteps 35-49 continuous rolling. New nodes added batch-wise via Op1 graph-expansion, no full-graph reconstruction. Monitor every rollout-step: AUC, F1-illicit, $\Delta\beta_1$, Dirichlet energy.

Figure 3

Figure 3(a): Rollout step vs Test-AUC.




Figure 3(b): Rollout step vs $\Delta\beta_1$ (SRE-AI only; baselines lack this metric and are excluded).





Figure 3(c): Rollout step vs Dirichlet energy (SRE-AI only).



Key findings:

1. `sre_ai_soft`: $\Delta\beta_1$ oscillates and drifts continuously over rollout steps (9-51 range). Soft loss-regularization cannot eliminate accumulated topological drift.
2. `sre_ai_hard`: **persistently holds $\Delta\beta_1 \equiv 0$ under continuous node addition**. Op9 manifold projection works in streaming incremental scenario.
3. Task metrics (AUC/F1) fluctuate for all models (including TGN / GraphSAGE) under heavy continuous new-node arrival.

> This experiment demonstrates the core novelty of SRE-AI: existing GNN / temporal GNN have no mechanism to constrain Betti-1 invariant during graph growth. Hard manifold projection trades negligible single-snapshot accuracy for long-run topological stability.

5.6 Experiment-5 Performance-scaling measurement

Vary node-count $N = 100 \cdots 5000$, measure per-step inference wall-clock time.

Figure 4

Figure 4(a): Node count vs single-step inference time (ms, log-log).





Figure 4(b): Node count vs peak memory (MB).



Observations:


1. Runtime grows approximately linearly with node-count; no $O(N^3)$ blow-up, validating local-horizon K_0 design.
2. Current Rust-pyd full-operator implementation has higher absolute latency than optimized PyG baseline (TGN / GraphSAGE). There remains room for optimization: tune Lanczos iteration, optimize checkpoint strategy, prune non-mandatory intermediate statistics.

5.7 Experiment-6 Relaxed-training vs native discrete inference

Using same trained weight; two inference pathways: (A) training-mode continuous relaxation; (B) Inference-mode native discrete SRE dynamics ($\{\pm 1\}$ spin, hard-sgn, boolean gate).

Figure 5

Figure 5: Bar chart comparing AUC of relaxed inference (A) vs native discrete inference (B), with ΔAUC annotated on top of bars.



Observations:

1. OGB-Mol-HIV: AUC drop only 0.003-0.02.
2. Elliptic: AUC drop ~0.008, F1-illicit drop ~0.09.

> The training-relaxed representation can transfer to original discrete SRE dynamics; train-inference gap is bounded and not catastrophic.

6 Discussion

6.1 Key insight: Soft-loss vs hard manifold projection

Ablation and long-rollout results show fundamental difference:

- Soft-constraint: loss term can *penalize* invariant deviation, but cannot guarantee manifold membership. Under long streaming evolution, topological drift accumulates.
- Hard-constraint (Op9): algebraic projection inside forward computation, strictly enforces $\Delta\beta_1 \equiv 0$. The cost is tiny or zero single-snapshot accuracy change, gain is long-run topological safety for open-world incremental graphs.

> Important: Hard-constraint is **not universally superior for every task**. If dataset is static fixed graph, soft-constraint may suffice. Hard-constraint shows strongest advantage for continuously growing streaming graph scenarios.

6.2 Over-smoothing suppression mechanism

SRE-AI suppresses over-smoothing via **dynamics-level positive-definite Dirichlet-energy constraint**, not by shortcut architectural modification. Experimental evidence: as depth increases, Dirichlet energy remains high, representations keep diversity, prediction performance does not collapse like baseline MP-GNN.

6.3 Limitations

1. Computational overhead: full 10-operator Rust implementation is slower than highly-optimized PyG baselines; further performance tuning required.
2. Current validation is limited to two public datasets. More datasets are needed to further verify generalizability.
3. Hard-constraint improves long-evolution robustness, but does not always boost single-time-slice prediction metrics. Should not expect universal SOTA on every static benchmark.
4. OGB-Mol-HIV heavy class imbalance limits F1 metric interpretability.

6.4 Future work

1. Performance optimization for Rust-pyd suite: Lanczos iteration tuning, lightweight checkpoint, pruning non-critical intermediate observables.
2. Expand evaluation to more datasets: citation graphs, more molecule benchmarks, multi-agent simulation datasets.
3. Noise-robustness test: inject graph perturbation, quantify whether hard-constraint improves noisy incremental learning.
4. Explore SRE-AI generative model: use Op1-Op9 for topology-controlled graph generation.

7 Applications and Use-Cases

Based on the full SRE-AI operator suite and empirical results, the model's strengths stem from intrinsic Dirichlet-energy constraints, native incremental graph expansion, and the dual soft/hard Betti-1 constraint paradigms. We separate **experimentally-validated use-cases** and prospective future applications, together with practical guidance and deployment caveats.

7.1 Experimentally-validated scenarios

(1) Open-world streaming incremental graph learning (core strength)

Target applications: blockchain transaction analysis, IoT networks and continuously-evolving real-world graphs.

Unlike TGN / GraphSAGE which rely on snapshots and offer no topological invariant guarantees, SRE-AI uses Op1 for incremental node-edge insertion without full-graph reconstruction. Hard-constraint mode via Op9 enforces $\Delta\beta_1 \equiv 0$ during long rolling evolution. Experiments confirm soft loss-only regularization cannot prevent accumulated topological drift. Besides node prediction such as illicit-account classification, SRE-AI outputs interpretable topological observables (Dirichlet energy, $\Delta\beta_1$, Z_{eff}) to monitor system-level topological health.

(2) Deep static graph learning mitigating over-smoothing

Validated on OGB-Mol-HIV molecular property prediction. As depth increases, conventional MP-GNNs suffer severe performance degradation from over-smoothing. SRE-AI maintains positive-definite Dirichlet energy, preserving representation diversity and outperforming GCN/GAT/GIN at larger depths. Soft-constraint mode suffices for static fixed-graph settings.

(3) Research test-bed for differentiable topological learning

Most existing differentiable topology methods place topological invariants solely inside loss terms. SRE-AI enables direct comparison: toggle the same operator pipeline between soft loss-regularization and forward-pass hard manifold projection. It supports controlled studies of manifold constraints, topological drift and accuracy-stability trade-offs in graph learning.

7.2 Prospective future applications (prototype-stage, needs further optimization)

1. **Joint learning-simulation digital twins**: Train under continuous relaxation mode on real-world observations; switch to discrete-spin `Inference` mode for forward dynamics simulation. Op5 computes endogenous edge-wise propagation rates without hand-designed time embeddings. Suitable for risk-propagation and multi-agent network systems, though computational overhead needs further tuning for large-scale deployment.
2. **Topology-oriented interpretability**: Compared to black-box GNN embeddings, rich intermediate topological quantities allow post-hoc reasoning for graph-mining predictions.

7.3 Practical mode selection & deployment notes

Mode	Use-case	Caution
Soft-constraint	Static graphs, short-horizon temporal tasks, prioritize snapshot accuracy	Long streaming runs accumulate $\Delta\beta_1$ drift
Hard-constraint ($\Delta\beta_1 \equiv 0$)	Streaming open-world graphs requiring topological invariants	May shrink representation manifold; no guaranteed snapshot accuracy gain

Key limitations: Current Rust full-operator latency exceeds highly-optimized PyG baselines and requires further performance tuning. Hard-constraint improves long-run robustness but is not a universal accuracy booster. Validation is limited to two benchmarks, requiring further cross-dataset testing.

8 Conclusion

This work proposes SRE-AI, a differentiable graph-learning framework built on Status-Relational Entropy topological-dynamics operator suite Op1-Op10. We implement full operator suite in Rust with analytic adjoint-mode autodifferentiation. Experiments verify:

1. SRE-AI intrinsically mitigates over-smoothing by maintaining positive-definite Dirichlet-energy. Static molecular task outperforms classical GNNs under large depth.
2. For dynamic incremental graph, SRE-AI achieves competitive performance compared with TGN / GraphSAGE.
3. Ablation and long-rolling incremental experiment prove: soft loss-regularization cannot prevent accumulated topological drift; Op9 hard manifold-projection can stably lock $\Delta\beta_1 \equiv 0$ in open-world continuous graph expansion, providing unique long-term topological robustness that conventional graph models lack.

SRE Full Suite of 19 Operators: What a "Physical-Semantic Graph Computing" Operator System Can Do — A View from the Graph500 Benchmark

Version: 2.0

> This article is based on measured data from the Graph500 v2.1 standard benchmark (SCALE 12 / 20 / 24), and systematically introduces the capabilities, performance, comparison with traditional graph algorithms, and overall application prospects of the full SRE (State-Relation Entropy) suite of 19 operators.

---

1. Background: Why a New Set of Graph Operators Is Needed

The field of graph computing has long revolved around four traditional problem classes: **traversal** (BFS/DFS), **shortest paths** (Dijkstra/SSSP), **centrality** (PageRank), and **connectivity** (connected components / strongly connected components). The Graph500 benchmark itself uses BFS TEPS (Traversed Edges Per Second) as its ranking metric.

However, real-world graphs — whether molecular bond graphs, knowledge graphs, social networks, or transportation road networks — often require more than just "can we reach it"; they demand answers to:

- How can **weight differences on edges** encode node attributes (e.g., electronegativity, capacity, similarity)?
- Do **cycle structures** in the graph carry topological phase information (e.g., Möbius strips, odd/even cycles)?
- How can the **shape of a graph be described in the spectral domain** rather than by traversing node by node?
- When the graph's "residual" is nonzero, how can it be decomposed into three orthogonal components: **gradient / harmonic / curl**?
- Do **topologically protected charge channels** exist on the graph (analogous to edge states in physics)?

The SRE operator system is precisely a set of 19 operators designed to answer these questions, covering the complete chain from fundamental graph transformation to spectral-shape solving, and from electronic degrees of freedom to topological charge channels.

---

2. What the 19 Operators Can Do

The full suite is divided into four layers by function:

2.1 Foundation Layer (Op1–Op10): Graph Structure Transformation and Gate Operations

Operator	Name	Core Function	Key Outputs
Op1	Local Graph Expansion	Dimensional expansion and embedding of the graph	n → n+1
Op2	Random Walk Metrics	Edge spin statistics and intersection measure	intersection
Op3	NAND Gate	Universal logic gate on graph (Turing-complete primitive)	Y_bool
Op4	Metric Weights	CSR SpGEMM 2-hop walk, generating metric matrix W_e	trace, frobenius
Op5	Latency Calibration	Propagation latency c_e based on W_e	c_e_mean
Op6	Spectral Splicing	Matrix-free Lanczos iteration for spectral gap λ_2 and spectral radius α	λ_2, α_n, residual
Op7	Adjoint Potential	Gradient energy norm	‖grad_E‖
Op8	Valve Flow	Valve flow variation δΠ	delta_Pi
Op9	Stitching	Dual space trace	tr(S_dual)
Op10	Firewall	Conduction gating and effective charge	z_eff, gate, p_cond

This layer provides the "infrastructure" of graph computing: Op4's sparse matrix multiplication and Op6's spectral decomposition are the foundation for all subsequent spectral-shape operators.

2.2 Electronic Layer (Op11–Op12): Charge Degrees of Freedom and Cycle Topology

Operator	Name	Core Function	Key Outputs
Op11	Spin-Charge Adjacency	Encodes electronegativity differences as edge-weight gains $A_q = A_s \cdot (1+\rho \cdot \lvert\tanh(\chi_i - \chi_j)\rvert)$	$\lambda_2 q$, αq, κ_q, max_spin_coupling
Op12	Closed-Loop Audit	DFS enumeration of simple cycles + Möbius phase ratio $\lvert tr(M^k)\rvert / n^k$	n_loops, mobius_ratio, λ_2_loop

Capabilities:

- Injects physical/chemical attributes of nodes (e.g., Pauling electronegativity) into the graph structure, producing a "charge-weighted adjacency matrix"
- Detects closed cycles in the graph and their phase properties, determining whether Möbius-type topological twists exist
- In molecular graphs: identifies aromatic rings and conjugated systems; in communication networks: identifies redundant loops

2.3 Spectral-Shape Layer (Op13–Op18): Closed Loop of Spectral-Shape Equation Solving

This is the core innovation of SRE — a spectral-shape equation solver that does not rely on large-scale eigendecomposition, based on Chebyshev moments and Hodge decomposition.

| Operator | Name | Core Function | Key Outputs |

|---|---|---|---|

| Op13 | Chebyshev Moments | $Y=2L/\alpha-I$, three-term recurrence $T_k=2YT_{k-1}-T_{k-2}$, $\tau_k=tr(T_k)/N$ | $\tau_0..\tau_K$, Y |

| Op14 | Hodge Projection | Three-component orthogonal decomposition $r = D\phi + h + C^T\psi$ | grad, harm, curl |

| Op15 | Endogenous Newton Step | $dM = -A^+(g-h)$, endogenous step size $\eta=F^2/quad$ + trust region | dM, quad, trust |

| Op16 | Hankel Clamp | Hausdorff moment problem PSD check + bisection backtrack | feas_before→after, t* |

| Op17 | Audit Verdict | $\gamma=\lambda_2/\alpha$, $\epsilon=\gamma\cdot\|r_0\|$, SOLVED/OBSTRUCTION/STALLED/CONTINUE | verdict, ε |

| Op18 | Homological Realization | Homology residual certificate $\|h\|$, homological distance d_hom, exact residual budget | h_norm, d_hom, budget |

Capabilities:

- **Op13** describes the spectral shape of the graph using Chebyshev polynomial moments, throughout $O(K\cdot N^2)$ with no eigendecomposition; K is endogenously determined by α/λ_2
- **Op14** decomposes any residual vector into "gradient field + harmonic field + curl field" — a direct realization of the Hodge decomposition theorem; the harmonic component corresponds to the first homology group H_1 of the graph
- **Op15–Op17** form a complete closed loop of spectral-shape inversion: Newton direction → step size → trust region → feasibility clamp → convergence verdict
- **Op18** outputs a "homological realization certificate", proving whether a given edge-weight configuration is a homological solution and its homological distance from a reference solution

2.4 Charge-Channel Layer (Op19): Z_2 Topological Charge

| Operator | Name | Core Function | Key Outputs |

|---|---|---|---|

| Op19 | Z_2 Charge Channel | Graph invariants (V,E,c,β_1,deg) + fundamental cycle basis holonomy + σ charge classification | β_1, deg, σ_trivial, Q_model, channel_type |

Capabilities:

- Computes the Betti number $\beta_1 = E - V + c$ (number of independent cycles) of the graph
- Determines whether the graph carries nontrivial topological charge via Z_2 holonomy over the fundamental cycle basis
- Outputs `channel_type`: electron-like (trivial channel) or nucleon-like (nontrivial channel)
- In molecular graphs: predicts conductive properties; in networks: identifies topologically protected information pathways

---

3. Graph500 Benchmark Results

Test environment: GCC 15.1.0 + OpenMP 8 threads, Graph500 official Kronecker generator, edge_factor=16, NBFS=3.

3.1 Official Ranking Metric (TEPS)

SCALE	Vertices N	Edges	harmonic_mean_TEPS !	Per-vertex Time
12	4,096	65,536	**1.80×10^{6}**	18.31 μs
20	1,048,576	16,777,216	**1.32×10^{8}**	0.250 μs
24	16,777,216	268,435,456	**8.85×10^{7}**	0.373 μs

! denotes the Graph500 official primary ranking metric (harmonic mean of TEPS). SCALE 20 reaches the peak of 1.32×10^{8} TEPS.

3.2 Time Distribution of the 19 Operators (median, seconds)

Operator	SCALE 12	SCALE 20	SCALE 24	Scaling Behavior
Op2 Random Walk	0.000	0.062	0.990	O(E) linear growth
Op4 Metric Weights	0.035	0.003*	0.000*	Dense W_e exceeds memory at large scale
Op5 Latency Calibration	0.032	0.000*	0.000*	Depends on Op4
Op6 Spectral Splicing	0.000	0.175	4.920	Lanczos iteration, large-scale bottleneck
Op11 Spin-Charge	0.002	0.000	0.000	Fixed 128-vertex subgraph
Op12 Closed-Loop Audit	0.006	0.007	0.008	Fixed 128-vertex subgraph
Op13–Op18 Spectral Solver	<0.001	<0.001	<0.001	Fixed small-dimensional chain complex
Op19 Z_2 Channel	0.000	0.001	0.000	Fixed 128-vertex subgraph

* At SCALE 20+, the dense W_e matrix ($N^2 \times 8B$) exceeds available memory; Op4/Op5 dense outputs are automatically skipped.

3.3 Key Physical Quantity Outputs

Metric	SCALE 12	SCALE 20	SCALE 24
Op6 λ_2 (spectral gap)	105.4	866.4	4,495.2
Op6 α_n (spectral radius)	4,769	138,029	739,644
Op11 κ_q (condition number)	1e30	1e30	1e30
Op12 mobius_ratio	0.898	0.932	0.932

| Op13 τ_0 / τ_2 | 1.000 / 0.033 | 1.000 / 0.033 | 1.000 / 0.033 |

| Op14 ‖grad‖/‖harm‖/‖curl‖ | 0.644/0.543/0.555 | same as left | same as left |

| Op16 feas (before→after) | -0.025 → -1e-9 | same as left | same as left |

| Op17 verdict | CONTINUE | CONTINUE | CONTINUE |

| Op19 β_1 / deg | 5 / 1 | 0 / 0 | 0 / 0 |

| Op19 channel_type | electron-like | electron-like | electron-like |

3.4 Format Compliance

The outputs of all three scales pass `parse_graph500.py` validation: all 22 required Graph500 fields are parsed, `harmonic_mean_TEPS` carries the `!` marker, and SRE extension fields are isolated by `#` comment prefixes without affecting the official parser.

---

4. Comparison with Traditional Graph Algorithms

4.1 Capability-Dimension Comparison

| Capability | Traditional Graph Algorithms | SRE 19 Operators | Notes |

|---|---|---|---|

| Traversal / Reachability | BFS, DFS | Op2 (edge spin statistics) | SRE does not replace BFS; it layers semantics on top |

| Shortest Path | Dijkstra, SSSP | Op5 (latency calibration c_e) | c_e is metric-based propagation latency, not geometric shortest path |

| Spectral Analysis | Eigendecomposition, Fiedler | Op6 (Lanczos) + Op13 (Chebyshev moments) | Op13 uses moments instead of full spectrum, $O(K \cdot N^2)$ not $O(N^3)$ |

| Connectivity | Union-Find, Tarjan | Op19 (c, β_1) | SRE directly gives Betti numbers, finer than connected components |

| Cycle Detection | DFS cycle enumeration | Op12 (simple cycles + Möbius phase) | Not just counting, but also analyzing phase properties of cycles |

| Node Attribute Encoding | Weighted graph (static weights) | Op11 (electronegativity → edge modulation) | Weights are dynamically generated from node attribute differences |

| Equation Solving | No native support | Op14–Op18 (Hodge+Newton+Hankel) | Complete closed loop of spectral-shape equation solving |

| Topological Classification | No native support | Op19 (Z_2 holonomy, channel_type) | Topologically protected charge channel classification |

4.2 Performance Comparison

| Algorithm | Typical Complexity | SCALE 24 Time | Notes |

|---|---|---|---|

| BFS (Graph500 benchmark) | O(V+E) | Included in 6.25s total | TEPS = 8.85×10^7 |

| SST (Single-Source Shortest Path) | O(E + V log V) | — | Graph500 second benchmark |

| PageRank | O(E·iter) | — | Power iteration |

| **SRE Op6 Lanczos** | O(V·iter) | **4.92s** | Matrix-free, only matrix-vector products needed |

| **SRE Op13 Chebyshev** | $O(K\cdot V^2)$ | **<1ms** (fixed small dim) | Three-term recurrence, no eigendecomposition |

| **SRE Op14 Hodge** | $O(m^3)$ pseudoinverse | **<1ms** (fixed small dim) | Runs only on small-dimensional chain complex |

4.3 Advantages

1. **Rich physical semantics**: Each operator corresponds to a well-defined physical quantity (spectral gap, electronegativity, Möbius phase, homology group), and outputs are directly interpretable
2. **Efficient spectral-shape solving**: Op13 replaces full spectral decomposition with Chebyshev moments; Op14–Op18 run on a fixed small-dimensional chain complex, with total overhead < 1ms at all scales
3. **Good scalability**: Op2/Op6 are O(E)/O(V) sparse operations; SCALE 24 (16M vertices) still completes within 6.25s
4. **Format compatibility**: 100% compatible with Graph500 v2.1 output format, directly integrable into the official ranking system
5. **Memory protection**: Op4/Op5 dense paths have automatic out-of-memory detection, safely degrading at large scales

4.4 Limitations

1. **Op4/Op5 dense path**: W_e is an N×N dense matrix; when N > 4.6K it exceeds memory, so trace/frobenius cannot be output at large scales
2. **Op11/12/19 subgraph strategy**: Currently runs on a fixed 128-vertex subgraph; at large scales the subgraph may be disconnected (λ_2_q=0), losing full-graph topological information
3. **Op6 large-scale bottleneck**: Lanczos takes 4.92s at SCALE 24, accounting for 79% of total time; further optimization possible (e.g., polynomial preconditioning)
4. **Op19 reachability analysis**: Depends on networkx automorphism enumeration; skipped when V > 22, so a complete channel_type cannot be given
5. **Non-traversal operators**: SRE does not directly optimize BFS/SSSP TEPS, but rather provides semantic enhancement on top of them


---

5. What This Whole Set of Operators Can Do

5.1 Modeling: From Graph to "Physical System"

The SRE operator system essentially maps any graph G=(V,E) to a "physical system":

```
Graph Structure ——Op4——> Metric W_e ——Op5——> Latency c_e
```



This is not a set of isolated operators, but a **complete reasoning chain from graph structure to physical properties**.

5.2 Typical Application Scenarios

| Scenario | Operators Used | Value |

|----------|----------------|-------|

| **Molecular Conductivity Prediction** | Op11 (electronegativity), Op12 (cycle topology), Op19 (charge channel) | Predicts σ channel type directly from bond graph, no DFT needed |

| **Network Robustness Analysis** | Op6 (spectral gap), Op14 (Hodge), Op17 (audit) | Spectral gap measures connectivity; Hodge decomposition locates bottlenecks |

| **Knowledge Graph Reasoning** | Op2 (intersection measure), Op3 (NAND), Op10 (firewall) | Logical inference and gating on graphs |

| **Topological Data Analysis** | Op14 (harmonic component = H_1), Op18 (homology certificate), Op19 (β_1) | Fast approximation of persistent homology |

| **Graph Neural Network Architecture** | Op4 (metric), Op6 (spectrum), Op13 (Chebyshev moments) | Filter design for spectral-domain GNNs |

| **Anomaly Detection** | Op16 (Hankel feasibility), Op17 (verdict) | Spectral-shape feasibility discontinuity = topological phase transition |

5.3 Relationship with Existing Frameworks

SRE is not meant to replace NetworkX, GraphBLAS, or DGL, but to provide **higher-level semantic operators**:

- **NetworkX** provides graph-theoretic primitives (BFS, shortest paths, centrality)
- **GraphBLAS** provides sparse matrix algebra (SpGEMM, SpMV)
- **DGL/PyG** provide graph neural network layers (GCN, GAT)

- **SRE 19 Operators** provide the **physical-semantic layer**: spectral-shape equations, Hodge decomposition, topological charge

In practice, SRE can be built on top of GraphBLAS (Op4 uses SpGEMM, Op6 uses SpMV), and can also be combined with GNNs (Op13's Chebyshev moments are precisely the core of spectral-domain GCN).

---

6. Conclusion

1. **All 19 operators pass the Graph500 standard test** — no crashes, no anomalies at SCALE 12/20/24, output format 100% compliant
2. **Peak TEPS 1.32×10^8** (SCALE 20), per-vertex time as low as 0.25 μs, with industrial-grade scalability
3. **Core innovation lies in the spectral-shape layer (Op13–Op18)**: Chebyshev moments + Hodge decomposition achieve $O(K\cdot N^2)$ spectral-shape equation solving, instead of the traditional $O(N^3)$ eigendecomposition
4. **Electronic layer + charge-channel layer (Op11/12/19)** provide physical semantics unavailable in traditional graph algorithms: electronegativity modulation, Möbius phase, Z_2 topological charge
5. **Overall positioning**: SRE is the "physical-semantic layer" of graph computing — downward it reuses GraphBLAS sparse algebra, upward it supports physical-property reasoning in domains such as molecules, networks, and knowledge graphs

Future optimization directions: full sparsification of Op4/Op5 (avoiding dense W_e), connected-subgraph sampling for Op11/12/19, and polynomial-preconditioned Lanczos for Op6.

Data source: measured output of `sre_graph500.exe` at SCALE 12/20/24. Raw files: see `sre_scale{12,20,24}_stdout.txt`.

Neural Information-Systems Theory: First-Order Thalamic Downsampling Instability and Second-Order Default-Mode-Network Integration Cascading-Failure Hypothesis

Version: 1.0
Archive Type: Private Knowledge Base · Exploration of Extreme Dynamics and Neuroinformatics
Data-Source Validation: OpenNeuro ds004504 (ICA/ASR pre-processing denoised derivative dataset)

> Two-Tier Control Architecture: First-Order Thalamic Downsampling Gateway (TRN); Second-Order DMN Posterior-Core Bus (PCC/PCu)
> Related references:
> https://doi.org/10.5281/zenodo.20377424 — Whole-Brain Parallelism and High-Dimensional Causal-Chain Topological-Spectrum Homomorphic-Mapping Mechanisms Based on Complex-Causal-Network Topology

I. Core Viewpoints and Two-Tier Stepped Defensive-Gating Hypothesis (Core Hypothesis)

From the cybernetics and digital-signal-processing physical framework, this hypothesis defines the brain's gating mechanism for processing massive real-world data throughput as a distributed, multi-tier stepped dynamic-network system:

1. **First-Order Hardware-Filter Physical Downsampling (Thalamus / TRN)**
   The raw information flux delivered by human receptors to the cortex is enormous; for the visual-cortex a conservative estimate reaches $10^{10} \sim 10^{11}\ \mathrm{bit/s}$. The thalamus acts as the irreplaceable main gateway for raw-data input across the whole brain. Its peripheral structure, the Thalamic Reticular Nucleus (TRN), a damping-network composed of GABA-ergic inhibitory neurons, functions as an anti-aliasing low-pass filter and physical downsampling chip (hardware decimator). It performs signal extraction and high-frequency cutoff at the lowest layer, locally formats and dissipates more than 99 % of background white noise, completing first-order physical downsampling.
2. **Second-Order Software-Compressor High-Order Parameter Reconstruction (DMN Default-Mode Network)**
   Feature parameters after primary thalamic filtering and substantial dimensionality reduction enter the core physical topological axis of the Default-Mode Network (DMN, specifically PCC / PCu). The DMN does not directly receive raw external physical signals; it operates as an advanced software compressor (principal-component extractor). It recalls internal historical causal blocks stored in the hippocampus, executes long-range high-order self-recurrent simulation and predictive coding, force-prunes and downsamples feature parameters, and converges them into a one-dimensional linear serial survival-decision-making pipeline.
3. **Integrative Energy Storage and Pathological Cascading Failure**
   The intrinsic structural vulnerability of Alzheimer's Disease (AD) lies in early-to-mid-stage dissociation and relaxation of inhibitory interneurons (first-order dampers) within the thalamic TRN, which causes severe leakage of the physical downsampling gating mechanism. High-dimensional causal charges pour in and bombard the cerebral cortex. To sustain survival-oriented computation, the DMN is forced to raise parameter rigidity and perform high-pressure energy storage (integral-energy-storage term in the time domain). As the integral-energy-storage capacity threshold drops sharply, sustained load eventually pushes the system across the critical-threshold boundary and triggers cascading failure of the two-tier filter defence line.

II. 5-Second Timeline Fine-Grained Decoupled Time-Slice Analysis (Temporal-Evolution Trajectory)

To investigate whether the physical phenomenon of "integrative-energy-storage and phase-transition breakdown" exists in the temporal dimension, under 5 Hz intermittent photic stimulation (characterised by high causal dispersion), the 5-second time window is decoupled into five independent 1-second fragments. The instantaneous spectral-entropy difference $\Delta\mathrm{SRE} = \mathrm{Stim} - \mathrm{Rest}$ is computed for each subject for every second.

Figure 1 Real-time trajectory of the information compressor under 5-Hz stimulation within 5 seconds




Figure 1 Real-time trajectory of the information compressor under 5-Hz stimulation within 5 seconds. Y-axis: instantaneous entropy change (Stim − Rest); X-axis: elapsed time (seconds). Red: Alzheimer's-disease AD group; Blue: healthy-control CN group; Green: frontotemporal-dementia FTD group.

Objective Interpretation of Curve Dynamic Characteristics

- **Healthy-Control Group CN (Blue Curve)**: Following pulse perturbation, it exhibits an extremely smooth monotonic-convergent gentle dissipation envelope towards negative values (−0.20) from 4.0 s to 5.0 s. This demonstrates that the healthy system possesses very high network resilience. The first- and second-tier filters cooperate in a stepped fashion, digest accumulated causal charges through continuous long-duration discharge channels, and spontaneously reorganise into a static, ordered, synchronous steady state.
- **Alzheimer's-Disease Group AD (Red Curve)**: During 1.0-3.0 s (energy-accumulation phase), mean entropy change remains tightly pinned near the zero axis, maintaining apparent quiescence. At 4.0 s the curve undergoes a near-vertical, non-linear unidirectional upward jump ($\Delta\mathrm{SRE}$ rises to +0.25), followed by a cliff-like fall back to +0.04 at 5.0 s.

> Physical Interpretation: This confirms cascading phase transition after the two-tier system struggles to resist overload. In the first three seconds thalamic downsampling performs aggressive filtering and load reduction, while the DMN executes parameter-damping control to sustain superficial equilibrium. At 4.0 s continuously incoming charges saturate and leak the front-end thalamic-TRN damping network. Flood-peaks of high-dimensional signal aliasing instantly breach the already-lowered DMN integral-energy-storage threshold, triggering system-level instantaneous breakdown (abrupt disorder explosion). The subsequent second brings step-wise depletion and blunting due to extreme synaptic fatigue.

III. Variance Analysis under Multi-Frequency Pressurisation with "Gender × Age-Cohort" (Multi-Factor Matrix)

Analysing clean ICA-EEGLAB derived datasets with artefacts such as eye-blink and muscle-movement noise fully removed. Taking age 65 (biological threshold for gonadal-hormone decline) as the biological age slice, response variance (`std`, i.e. distribution width / system heterogeneity) of each disease group under different external pressurisation frequencies shows highly-specific non-linear phase-transition behaviour.

Figure 2 ICA-denoised data: SRE-entropy drop versus MMSE score, age and gender




Figure 2 ICA-denoised data: SRE-entropy drop versus MMSE score, age and gender. X-axis: MMSE scale score; Y-axis: instantaneous entropy change (Stim − Rest).

Stim_Freq	Age_Cohort	Group	Gender	mean	std	count	Notes
5Hz	Older_Group(>65)	A(AD)	F	0.2338	0.8687	13	Female Instability
5Hz	Older_Group(>65)	A(AD)	M	-0.2248	0.5101	7	
5Hz	Older_Group(>65)	C(CN)	F	-0.2404	0.4589	7	
5Hz	Older_Group(>65)	F(FTD)	F	0.0515	0.5589	5	Hardware Control
10Hz	Older_Group(>65)	A(AD)	F	0.1362	0.5542	13	
10Hz	Older_Group(>65)	A(AD)	M	0.0450	0.3354	6	Rigid Lockup
10Hz	Older_Group(>65)	C(CN)	M	0.2499	0.4210	10	
15Hz	Older_Group(>65)	A(AD)	F	0.2985	0.5928	13	
15Hz	Older_Group(>65)	A(AD)	M	0.0225	0.6304	7	Generalized Convergence

IV. Phenomenological Clues from the Dataset

1. **Variance explosion in elderly-female AD subjects under 5 Hz ($std = 0.8687$)**
   Under low-frequency discrete impulses, only the elderly-female AD group exhibits sky-rocketing variance. Age-matched FTD female subjects (DMN-intact control group) maintain low variance (0.55). This suggests reproductive-senescence alone cannot produce system destabilisation. Only the combination of "loss of reproductive-choice preservation" plus "DMN hardware-gating damage" causes the second-order compressor in elderly-female brains to fully lose resilience and explode into extreme bipolar polarisation.
2. **Rigid-parameter lock-up in elderly-male subjects under 10 Hz ($std = 0.3354$)**
   When external photic flicker falls near the intrinsic human-brain alpha-band frequency of 10 Hz, variance of elderly-male AD subjects contracts catastrophically to the global minimum. This indicates decades-long partial androgen deficiency (latent recession) in males crosses a critical late-stage threshold, completely disabling downsampling-filter dampers. Confronted with intrinsic-frequency excitation the system cannot dissipate elastically and is forcibly locked into a single rigid resonant state.

3. **Homogenisation of damped exhaustion for both genders under 15 Hz**
   As external energy load rises further to 15 Hz (beta-band discharge stimulation), response variances for elderly male and female patients converge strongly (0.59 versus 0.63). This demonstrates that within the high-frequency band systemic synaptic destruction overtakes gonadal-hormone decline and becomes the dominant decisive factor for network collapse.

V. Distributed-Hardware Decomposition of the Two-Tier System Control (Sub-System Architecture)

Brain control described within this hypothesis is not monolithic or centralised. It constitutes a distributed dynamic-control cluster whose components are spatially isolated yet sustain high-frequency temporal synchronisation. Its core computational structure divides precisely into three modules:

1. **First-Order Hardware Downsampling Gateway: Thalamus & TRN**
   - Dynamical Function: Front-end offloading for all raw sensory inputs of the brain. The TRN damping-network uses GABA-ergic neurons to perform physical-level low-pass high-frequency cut-off and hard decimation. Massive sensory waveforms are down-sampled at source into low-pass feature-parameter packets acceptable to cortex and DMN. DMN performs reverse dynamic gating-parameter control via tightly-coupled thalamocortical loops.
2. **Second-Order Central-Control Bus (DMN Core-Hubs)**
   - Primary Brain-Regions: Posterior-Cingulate-Cortex / Precuneus (PCC/PCu), medial prefrontal cortex (mPFC).
   - Dynamical Function: Acts as advanced software compressor (principal-component extractor). The PCC possesses the highest anatomical long-range-fibre connectivity across the whole brain and is responsible for homotopic alignment of whole-brain feature-parameters. The mPFC tags down-sampled parameters with “self-relevance” labels and completes hand-over for higher-order decision-making.
3. **Second-Order Historical-Memory Causal-Building-Block Library (DMN Medial-Temporal Sub-System)**
   - Primary Brain-Regions: Medial-temporal-lobe hippocampus, parahippocampal gyrus, posterior inferior parietal-lobule cortex (pIPL).
   - Dynamical Function: The “local historical-database” for the DMN filter. The hippocampus supplies discrete autobiographical-memory fragments as “causal building-blocks” for internal simulation. When first-order thalamic downsampling fails due to damping-degradation, overload-leakage and high-dimensional signal-aliasing flood-waves bombard cortex; native memory-blocks here undergo forced distortion and trigger high-confidence memory illusions (confabulation).

VI. Thalamo-DMN Two-Tier Stepped Information-Gating Circuit (Two-Tier-Gating Architecture)

Following physical realities of digital-signal-processing and hardware-gating physics, brain information-filtering does not operate within a single layer. Instead it is “distributed defence-in-depth” jointly constructed by the thalamus acting as first-order physical downsampling hardware-filter, and the Default-Mode-Network acting as second-order dimensionality-reduction noise-cancelling compressor:

1. **First-Order Hardware-Extractor Physical Frequency Reduction (Thalamus-TRN)**
   As the general gateway for all sensory inputs, the thalamus is encircled by the Thalamic-Reticular-Nucleus (TRN) implementing anti-aliasing low-pass filtering. Irrespective of signal semantics, extraction proceeds purely based upon physical firing-rate and spatio-temporal contrast. It executes energy decoupling and truncates raw biochemical surges into low-frequency feature-parameter packets compatible with cortex.
2. **Second-Order Logical-Compressor Lossy Dimensionality Reduction (DMN Cortex)**
   Receiving parameters from thalamic downsampling, cortical DMN executes lossy dimensional-reduction compression driven by memory-based causal templates. It performs information-level dimensionality reduction, prunes multi-dimensional causal-chains irrelevant to self-prediction, and finally converges and refines data into a one-dimensional linear serial behavioural-decision pipeline.
3. **Systemic Nature of Damping: “Destructive-Interference” originating from Phase-Mismatch**
   Within distributed high-dimensional networks, biochemically-observed “increased damping / throughput obstruction” actually arises from control-theoretic phase-mismatch-induced coherence-loss.
   - **Healthy-System State**: Node timestamps are perfectly precise. When signals traverse pathways wave-peaks align perfectly, constructive-interference occurs, permitting efficient information circulation and dissipation.

- **Damaged-System State**: Master-clock skew produces inter-node phase-drift. Under continuous external-pulse loading phase-errors accumulate non-linearly over time. Once critical boundaries are crossed (at 4 seconds), phase-unlocking occurs. Wave-peaks collide with wave-troughs producing destructive-interference. Wide-area data-transmission fails, severe network-reflection and aliasing white-noise are generated and cascading-failure is triggered.

VII. Pineal-Gland Macro-Clock and System-Sweeper Maintenance Mechanism (Pineal-Gland & Baseline-Reset)

From macro-temporal-dynamics perspective the pineal-gland fulfils the critical physical role of "circadian-rhythm clock" and "system baseline-state resetter" within the brain's two-tier control:

1. **Macro Maintenance-Mode Switching**
   Via the retinohypothalamic tract the pineal-gland receives circadian signals originating in the Suprachiasmatic-Nucleus (SCN). At night, in absence of photon-flux, melatonin is fully released issuing a system-suspend command. It forcibly switches first-order thalamus and second-order DMN from daytime "high-energy-consumption sensory-compression mode" into night-time "low-energy-consumption idle-maintenance mode".
2. **Glymphatic Clearance and Mitochondrial Repair**
   During night-time slow-wave synchronous-sleep gated by the pineal-gland, the cerebral glymphatic-system circulates fully. It flushes biochemical toxic-metabolic by-products (Aβ-plaques, Tau-protein deposits) accumulated during daytime high-aerobic-glycolysis computations inside thalamus and DMN. This directly prevents premature physical-ageing of first-order TRN damping-networks and second-order cortical synapses. Very-early-onset pineal-gland calcification and atrophy among AD-patients constitutes the macro-level root-cause triggering hardware-level cascading-collapse of the two-tier control-circuit.

VIII. Thalamus-Targeted Preventive-Interventions built upon First-Order Thalamic Downsampling Gateway

Since physical-downsampling and the primary anti-aliasing gating-control reside within the thalamus (TRN closed-loop inhibitory-network), the most practical foundational natural-prevention strategy for whole-brain overload-meltdown protection must target precisely this "main-flow gateway" for non-invasive physical-regulation.

Simultaneously a dialectical cybernetic perspective must be maintained: blindly unconditionally raising thalamic hardware-damping comes at the price of bandwidth-lock-up for the downsampling device. System responsiveness towards novel external causal-signals becomes severely blunted; clinically this manifests as stereotyped dementia with apathy, numbness and loss of cognitive-flexibility. Preventing senile dementia fundamentally means preventing late-stage thalamic-filter oscillation between two vicious poles: "zero-damping runaway excitation" versus "high-damping rigid dead-lock". The core is adaptive dynamic-margin recalibration:

1. **Multi-Modal Low-Frequency Sensory-Pulse Entrainment (TRN-Damping Training)**
   As converter for whole-brain sensory-channels, the thalamic-gateway possesses high spatio-temporal phase-locking sensitivity. Regular closed-eye binaural low-frequency acoustic-wave exposure (slow θ or δ rhythm sound-therapy) combined with somatosensory vibrational-acoustics (bone-conduction delivering low-frequency mechanical-waves upwards along the spinal-cord to stimulate the thalamus). This multi-modal flexible-rhythm-loading effectively fits an "exogenous shock-absorbing frame" over damaged relaxed TRN damping-networks, vicariously intercepting and dissipating high-frequency environmental causal-noise.
2. **Autonomic-Vagus Reverse-Drive (Endogenous Neurotransmitter-Pump)**
   The thalamic-TRN damping-network critically depends upon biochemical acetylcholine (ACh) and GABA concentrations to sustain gating-rigidity. Specific prolonged deep-abdominal exhalation (coherent-breathing training precisely 5.5 breaths per minute: inhale 5.5 s, exhale 5.5 s), or cold-water facial immersion (trigger diving-reflex) can directly reverse-activate peripheral vagus-nerve acting as the system-braking bus. This prompts brain-stem nuclei to force-release endogenous GABA neurotransmitters into thalamic TRN, instantaneously lengthening single-sampling time-windows of the first-order downsampler. This installs a "time-delay buffer-zone" upstream of the downsampler, granting ample micro-second-scale time for expansion-dissipation against discrete-pulse bombardment and instantaneously raising filter hardware-impedance coefficient γ.
3. **Physical Capacitance-Clearing and Synaptic-Membrane-Fluidity Re-Construction (Free accumulation-margin enabling smooth state-transitions)**
   Preserve the golden deep-sleep window 22:00-2:00 at night. Pineal-gland melatonin activates whole-brain

glymphatic-clearance physically emptying caches and expanding system dynamic-capacitance response-margins. Simultaneously strictly limit refined high-sugar intake to prevent glycolytic-endothelial-leakage inside thalamic micro-vascular-networks and synaptic glycation. Sufficient high-quality unsaturated-fatty-acid supplementation (DHA fish-oil, high-purity lecithin) physically embeds within phospholipid-bilayers of thalamic-TRN synaptic-membranes raising membrane-fluidity and reshaping damping-switch ion-channel-responses. Guarantees the filter can seamlessly phase-switch between “ultra-fast opening” and “ultra-fast settling” and avoid rigid-parameter dead-lock.

IX. Input-Stream Optimisation & Multi-Factor Cybernetics based upon Input-Data-Flow Structural-Transformation

From pure complex-systems-science and cybernetics perspectives, degradation of the brain’s two-tier-control, the 4-second temporal-jump and cross-gender phase-transitions are not simple unidirectional causal-chains determined by isolated phase-mismatch variables. Macro-electrophysiological instability is fundamentally a system-level collapse-phenomenon emerging after four multi-scale layers (biochemical-layer: protein-deposition / transmitter-depletion; structural-layer: long-range brain-fibre demyelination; endocrine-layer: gonadal-hormone-dissipation / micro-vascular-evolution; signal-layer: external-pulse-loading / clock-drift) become heavily non-linearly intertwined and coupled across space-time. Input-stream-optimisation must act at entry-port level altering data certainty, phase and dimensionality to push demodulator and clock-generator toward phase-transition recovery.

1. **Input-Data-Flow “Data-Stream Randomisation & Dithering”**
    - Cybernetic-Mechanism: Continuously-loaded high-certainty periodic discrete-pulse packets (e.g. strong 5 Hz stimuli) produce thermodynamic-charge accumulation inside front-end hardware-demodulator (thalamic-TRN damping-network), filling buffer-ceiling and forcing system phase-lock. Artificially injecting Gaussian-white-noise or highly-random aperiodic irregular-data-streams (engineering dithering technique) washes away specific auto-correlation-frequencies from external-signals responsible for internal phase-accumulation-errors and jointly resists chronic biochemical-endocrine-layer erosion upon damping-barriers.
    - Reverse-Recalibration-Outcome: When input-stream auto-correlation drops to zero, front-end thalamic-demodulator is relieved from single-frequency phase-synchronisation-pressure. To dissipate disordered random-stream-energy the system forces underlying feedback-clock-generator into high-entropy dissipative-state. This digitally frees system-buffer-margins and clears buffer-overload.
2. **Global-Master-Clock “Input-Stage Phase-Locked-Loop for Master-Clock”**
    - Cybernetic-Mechanism: Pineal-hypothalamus complex constitutes global hardware-master-clock-generator for the information-network supplying unified timestamp-baselines for distributed multi-tier sub-systems. Alzheimer-disease systemic-collapse fundamentally arises from distributed sub-module phase-decoupling caused by clock-skew and phase-drift. Slave-phase-locked-loop (PLL) tuning is performed at input-stage using physical media carrying temporal-pulses.
    - Reverse-Recalibration-Outcome: High-throughput high-frequency temporal-reference-pulses input during daytime enforce rigorous phase-entrainment of hardware-master-clock against external physical-timelines. Once master-clock achieves phase-lock-calibration its night-time maintenance-reset-signals acquire strong drift-resistance, precisely aligning thalamic-demodulator sampling-shutters onto absolute-zero-point via underlying buses and fundamentally eliminating timeline-propagated cascading-failure hazards.
3. **Input-Stream “One-Dimensional Sequential-Dimensionality-Reduction”**
    - Cybernetic-Mechanism: Complex environments and social-networks manifest within information-systems as high-dimensional causal-matrices possessing strong non-linear coupling-parameters. When forced into input-gateway, potential infinitely-divergent computational-simulations saturate second-order cortical lossy-compressor (DMN) processor-resources triggering chronic year-round integral-energy-storage exhaustion. Manifold-dimensional-reduction compression is applied at pre-processing-layer prior to data entering main-system-gateway.

X. Acoustic-Inflow Harmonic-Recalibration: Low-Energy Harmonic-Recalibration-Mechanism of Acoustic-Causal-Streams for Two-Tier-Control

Directly discarding 99.9 % cross-spatiotemporal non-linear matrix-parameters at pre-attentive gateway-layer forcibly reconstructs high-dimensional uncertain-inputs into one-dimensional linear sequential pure-action-vectors (sequential-task-streaming). After input-stream dimensional-reduction compresses down into lowest-energy one-dimensional-states, complex-integrative-computation energy-consumption of cortical-processors drops precipitously (system-occupancy converges toward unloaded maintenance-state). The DMN emits high-fidelity control-commands down-stream across thalamocortical-feedback-buses, sustaining first-order-demodulator TRN damping-networks within adaptive-dynamic phase-transition-states of maximal-response-strength.

From information-systems-theory and distributed-filter-architecture perspectives music (high-chord-coherence acoustic-signals) constitutes a golden-input-stream: possessing “extremely-low information-throughput-load” yet “very-high-certainty harmonic-resonance-relationships” at input-stage. Compared against high-throughput photic-stimulation, acoustic-signals achieve non-invasive reverse-recalibration of phase-offsets between first-order downsampler and second-order-compressor through physical-level cascaded-phase-locking plus endogenous-damping-pumps, without triggering system thermodynamic-overload-defences:

1. **Input-Stage “Low-Throughput Energy-Offloading” Mechanism**
    - Visual photic-stimulation delivers massive raw-flux up to $10^{10} \sim 10^{11}$ bit/s. To execute front-end high-frequency-extraction first-order thalamic-demodulator (TRN damping-network) consumes enormous energy and becomes vulnerable toward saturation-failure during early-to-mid AD-progression. By-contrast auditory-acoustic-signals possess orders-of-magnitude-lower physical-throughput. Feeding-in-music permits front-end-demodulator to operate within lightweight ultra-low-energy-consumption mode, fully freeing synaptic-capacitance-margins for fine adaptive-damping-adjustments.
2. **Multi-Part-Overtone-Series “Multi-Level Phase-Locking via Harmonic-Overtones”**
    - Music fundamentally is a causal-resonance-network built from chords, musical-intervals and overtones obeying strict mathematical-ratios. Fundamental-frequency and overtone-series exhibit pure integer-multiple resonance-relationships $(1:2:3:4)$. When fed-in this periodic-waveform imposes powerful system-wide phase-entrainment via auditory-pathways. These exogenous highly-coherent mathematical-rhythms act as invisible physical-calipers for the network, digitally forcing realignment of clock-skew-distorted sampling-shutters belonging to first-order thalamus plus diverse distributed second-order DMN-sub-modules (PCC, mPFC), eliminating false high-damping originating from destructive-interference.
3. **Limbic-System “Endogenous-Damping-Regulation via Hebbian-Plasticity”**
    - Upon acoustic-signal-propagation into thalamus, lower-brain-stem plus limbic-system-networks are directly stimulated triggering non-linear whole-brain multi-factor neurotransmitter-reorganisation. Resonant-sound-waves matching individual aesthetic-steady-states spontaneously drive parasympathetic-vagal braking-bus, prompting central-structures to release high-density GABA plus acetylcholine neurotransmitters. At microscopic-level transient gain-regulation is completed for first-order TRN-gating and second-order cortical-synaptic-membrane-fluidity, achieving micro-second-scale variable-frequency phase-transition-elasticity between “ultra-fast opening” and “ultra-fast settling” under completely safe exogenous-drug-free conditions.

XI. Knowledge-Base Summary and Future Medical-Validation Vectors

1. **Qualified Review (Cautious Interpretation)**
   This hypothesis presently builds upon single-centre small-sample dataset ds004504 (some stratified subgroups satisfy $N \leq 6$). This open-source dataset lacks critical physical-parameters including stroboscopic-lamp absolute-light-intensity and precise duty-cycle. Consequently this model makes no definitive causal-claims. Observed data-trends are strictly interpreted as “consistent with evolutionary-trends predicted by this hypothesis”. Systematic confounding-biases originating from sampling-error, patient-medication-history or educational-level cannot be excluded.
2. **Future Retained Knowledge-Base Significance**
   Constrained by dataset-scale this hypothesis breaks traditional medical misconceptions reducing Alzheimer-disease purely to static-EEG power-spectrum trivialisation. It points toward two highly forward-looking technical directions:
    - **Time-Domain Dynamic-Stress-Testing**: It demonstrates that the first three-seconds of system-operation actively suppress energy-accumulation. This indicates future ultra-early-screening ought to shift away from static-biomarkers

toward time-domain accumulation-phase-transition-points under intermittent-pulse-stress-testing (e.g. capturing the 4-second cascading-failure-point).

- **Network-Damping-Recalibration-Theory**: It suggests core-tasks for future Brain-Computer-Interfaces (BCI) and non-invasive-physical-regulations (e.g. GENUS 40 Hz opto-acoustic synchronous-feedback) are not sensitivity-elimination. Instead they target precise restoration of long-distance negative-feedback braking-capabilities lost through degradation-dissolution of first-order thalamic-TRN plus second-order interneurons via targeted topological-rerouting, thereby elevating system-energy-storage-thresholds.

Whole-Brain Parallelism and High-Dimensional Causal-Chain Topological-Spectrum Homomorphic-Mapping Mechanisms Based on Complex Causal-Network Topology

Version: 1.0

Related references:
https://doi.org/10.5281/zenodo.19935370
https://doi.org/10.5281/zenodo.20344105
https://doi.org/10.5281/zenodo.20301819

Abstract

This paper establishes a unified logical architecture spanning systems science and cognitive neuroscience to elucidate the underlying mechanics of whole-brain neural-firing networks, informational gating of the Default-Mode Network (DMN), emergency whole-brain topological parallelism under extreme stress, high-dimensional inspiration emergence, non-local cognitive synchronization, and early-historical cognitive symbology.
We posit that independent physical matter does not exist; all cosmic phenomena are macroscopic emergent illusions arising from continuous, uninterrupted interactions within topological causal networks. Inside this deterministic web, distinct structural clusters share symmetric topological-spectrum characteristics of causal chains. Under homeostatic conditions, the human brain-acting as a localised causal-chain cluster-is constrained by the DMN to carry out lower-dimensional, serial-linear causal computations for physical survival.

Nevertheless, under extreme physiological or psychological stress (e.g. chronic anxiety, near-death experiences), the DMN undergoes systemic failure or shutdown, triggering an evolutionary fallback mechanism that shifts the brain into a state of global parallel resonance and achieves structural alignment across the full neural configuration. Within this hyper-parallel receptive state, the brain operates as a high-energy topological receiver, instantaneously capturing non-local cross-scale homomorphic alignments inside the macro causal web, which phenomenologically manifests as profound counter-intuitive inspiration.

Furthermore, this framework demonstrates that non-local cognitive synchronization (phenomenologically labelled telepathy) represents deterministic convergence of topological disturbances propagating along homologous causal chains. Early-historical non-linear conceptual frameworks are not mystical revelations; they constitute lower-dimensional linguistic encodings of these high-dimensional topological-alignment phenomena, constrained by the dimensional-reduction property inherent to human serial language.

Ultimately, this paradigm delivers a self-consistent, scale-invariant synthesis unifying micro-neuroelectrical dynamics, macro-causal-network topology, and anomalous phenomena of human consciousness.

Keywords: Causal Networks; Topological-Spectrum Characteristics of Causal Chains; Homomorphic Mapping; Default-Mode Network; Whole-Brain Parallelism; Non-Local Synchronization; Linguistic Dimensional-Reduction

I. Introduction

Contemporary cognitive science and neurobiology yield robust empirical explanations for serial-linear logical reasoning. When confronted with non-linear leaps of paradigm-shifting inspiration, dissociative depersonalization induced by trauma, non-local cognitive synchronization across geometric distances, as well as highly convergent “transcendental states” documented in early-historical archives, standard models either reduce such occurrences to pathological artefacts or resort to unscientific dualism.

Treating the global complex causal web as the fundamental informational unit and adopting homomorphic mapping of topological-spectrum characteristics as an explanatory bridge, this paper constructs a continuous, strictly materialistic theoretical framework. It systematically traces informational gating implemented by neural filters, stress-provoked DMN collapse, structural cross-alignment among distinct causal domains, and the dimensional-reduction process from high-dimensional perception into serial syntax. Under the unified topology of deterministic causal networks, this synthesis deconstructs high-order cognitive transitions, creative emergence and non-local synchronization.

II. Ontological Framework for Complex Causal Networks and Homomorphism of Topological-Spectrum Characteristics

This paradigm maintains that the universe is not governed by independent physical-matter laws. Beneath the perceptual threshold lies an absolute, dense, incessant network of causal chains. Any apparent physical entity-whether sub-atomic particle assemblies, cellular tissues or astronomical bodies-is essentially a causal-chain topological-cluster defined by its specific closed-loop geometry, intrinsic cycles and information-carrying capacity.

Since all local-sub-networks originate from one singular global causal matrix, internal arrangements of causal connections across different scales, frequencies and nodes (such as quantum coupling, biological fluid loops, or structural syntax within music and mathematics) inevitably exhibit profound geometric symmetry. This structural configuration and invariant mapping pattern across network connections is defined as the **topological-spectrum characteristic of causal chains**.
Within algebraic graph theory, distinct informational domains enter a state of **homomorphic mapping (Homomorphism)** when they possess identical topological-spectrum characteristics. This constitutes the ultimate invariant convergent form of the cosmic causal web.

III. Informational Gating of the Default-Mode Network and Local Serial Computation

Functioning as a specialised causal-chain topological-cluster, the human brain maintains macroscopic cognitive stability relying on ontogenetically-developed neural filters, chiefly the prefrontal cortex and the Default-Mode Network (DMN). Acting as an evolution-derived informational pressure-relief valve, this filter follows the principle of “attenuation for localised focus”:

1. **Serial Computation and Survival Optimisation**: The filter forcibly attenuates massive high-dimensional topological-spectrum characteristics belonging to the global causal web. It restricts neural computation to localised one-dimensional linear causal chains rooted in past-experience (e.g. immediate threat assessment, social labelling, maintenance of local entropy). This constraint guarantees pragmatic survival efficiency and cognitive stability for adult organisms.
2. **Filter-Free Primordial Ground State (Early Ontogeny)**: During early childhood development, prefrontal-cortex and DMN filters remain structurally immature. The neural-network baseline is dominated by low-frequency oscillations. Children’s brains naturally lack adult-style cognitive firewalls and reside in a boundary-dissolving fused state, capable of direct interaction with underlying topological-spectrum characteristics of the natural causal web.

IV. Stress-Induced Filter Collapse, Whole-Brain Parallelism and Topological-Spectrum Identification

When organisms suffer catastrophic stress, for instance systemic chronic anxiety (information overload), or near-death episodes (acute cerebral hypoxia, complete central-nervous-system energy depletion), the brain activates an ancient DNA-encoded emergency takeover mechanism to guarantee fundamental system survival. Homeostatic cognitive states dissolve and trigger inward-outward topological restructuring:

1. **Filter Dissolution and Shutdown**: Chronic anxiety forces DMN firewalls to sustain continuous high-frequency discharge and gives rise to systemic “overload fusion”. In near-death events, evolutionarily-recent neocortical structures together with the DMN undergo physical shutdown due to metabolic starvation.
2. **Whole-Brain Topological Parallelism**: After systemic removal of DMN filters, functional firewalls separating individual cognitive sub-domains (visual processing, memory retrieval, abstract logic) disintegrate. Neural electrical activity abruptly transitions from fragmented serial states toward globally-synchronised hyper-parallel topological resonance; billions of neural nodes align into one unified integrated structural matrix.
3. **Topological-Spectrum Identification and Creative Emergence**: Within the globally-parallel state, the brain ceases decoding discrete local sensory fragments. Instead it operates as a high-dimensional “topological mirror”, establishing instantaneous homomorphic alignment with external causal-network domains that share identical underlying topological-spectrum characteristics, even across entirely unrelated fields.

> Topological definition of inspiration: Creative inspiration is not ex-nihilo fabrication of novel entities by the brain. It denotes instantaneous topological alignment occurring at shared nodes between internally parallelised brain networks and the macro-cosmic network. The brain directly “imprints” and identifies perfect structural symmetry between two seemingly disconnected causal chains (for example, closed-loop visual topology of the ouroboros in ancient iconography perfectly matches self-circulating causal topology of conjugated molecular rings).

Crucially, this emergency topological parallelism imposes extreme requirements for hardware resilience. Statistical macro-sampling shows that the overwhelming majority of biological nervous-systems cannot withstand destructive surges of high-frequency electrical and neuro-chemical forces, resulting in personality dissociation, psychological depersonalisation or even systemic self-destruction-these represent stochastic collapse of ill-adapted hardware inside the causal web. Only a tiny subset of genetically-atypical individuals with extreme synaptic-plasticity and robust biological hardware can cross this critical threshold, reconfigure their networks and achieve deterministic cognitive leaps.

V. Non-Local Network Disturbance and Cognitive Synchronization (Telepathy Phenomenon)

Within this pure causal-chain topological framework, the historically-observed phenomenon of non-local cognitive synchronization commonly called telepathy can be fully demystified via graph-theory and topological dynamics. This mechanism requires neither superluminal propagation nor hypothetical unconfirmed radiation, and strictly obeys the following topological laws:

1. **Homologous Causal Coupling**: When two or more neural causal-clusters undergo deep sustained interactions within network history (e.g. direct genetic lineage, or high-intensity shared traumatic stress), their internal networks develop profound topological coupling. They evolve into two mutually-complementary, strongly-correlated sub-networks permanently tethered to one common historical causal junction.
2. **Non-Local Topological-Disturbance Propagation**: Objective physical matter has no independent reality; spatial distance is merely a low-dimensional scalar illusion manufactured by DMN filters. If one homologous neural node suffers extreme stress or near-death collapse, the system releases catastrophic topological disturbance. This structural distortion propagates instantaneously along contours of underlying shared causal-chains, completely circumventing geometric-space limitations.
3. **Bidirectional Homomorphic Re-Alignment**: As this topological disturbance arrives at the remote paired node, provided the receiving node happens to occupy a state of weakened filter-defence (such as θ-wave synchronized sleep, rapid state-transitions or meditative attunement), whole-brain parallelism is passively triggered. Two geographically-separated neural clusters instantaneously accomplish homomorphic re-alignment of their causal topologies. Macroscopically this manifests as instant unmediated shared cognitive states, physiological panic responses or identical conceptual ideations across vast geometric separations.

VI. Dimensional-Reduction of High-Dimensional Topological Perception and Deconstruction of Historical Texts

Information captured through whole-brain parallel alignment forms a high-dimensional non-linear pan-scalar topological map. When experiencers attempt to articulate this holistic state toward filter-constrained ordinary social agents, they immediately encounter hard constraints imposed by linguistic dimensional-reduction. Human language itself constitutes a low-dimensional serial-linear filter built upon binary dualisms; structurally it cannot achieve isomorphism against whole-brain parallel causal-topology.

To circumvent such “ineffability”, ancient thinkers and chroniclers resorted to local cultural idioms and mythological symbology of their respective eras, performing primitive down-sampling and literary encapsulation for these invariant topological-spectrum characteristics:

1. **Deconstruction of Catastrophic Metaphors**: Cross-cultural ancient accounts describing absolute destructive forces together with instantaneous cosmic rebirth represent early structural depictions of fundamental evolutionary dynamics for causal webs: violent dismantling of obsolete topological connections to enable higher-order reconfiguration. Lacking modern network-science terminology, ancient observers encoded this terrifying yet liberating causal spectrum into dualistic mythic archetypes that paradoxically combine cosmic wrath and absolute static tranquillity.
2. **Deconstruction of Classical Philosophies**: Historical axioms depicting singular self-sustaining all-encompassing cyclic trajectories (“moving in cycles without ceasing”) serve as early descriptions for macro closed-loop dynamics of global causal networks. Meanwhile doctrines describing incessant interdependent arising-and-perishing of local events (“this arises because that arises”) deliver low-dimensional encodings for local topological dynamics and non-local network correlations. Structural convergence among classical philosophies across geographically-isolated cultures constitutes an inevitable consequence of uniform human neural hardware: once local filters are stripped away under stress, brains map onto identical invariant topological-spectrum characteristics of the universe.

VII. Conclusion

Taking complex causal-networks as its foundational architecture and homomorphic mapping of topological-spectra as a continuous bridge, this paper bridges gaps between micro-neurocognition, non-local synchronization phenomena and macro-systemic laws. Ontogenetically-developed neural filters (DMN) inside adult human brains are optimised for localised linear survival, at the cost of rendering organisms blind toward macro-causal topologies and remote-network disturbances. By contrast, stress-induced whole-brain parallelism and homologous-network disturbance-propagation represent evolutionary emergency backdoors hard-coded into human DNA for cognitive leaps and non-local coordination.

The universe possesses no independent physical entities; it forms an invariant self-folding topological-network of causal vectors. Human inspiration, anomalous non-local synchronization, and grand architectures of ancient thought are not supernatural mystical anomalies. They represent deterministic unavoidable outcomes of scale-invariant causal laws operating within constraints imposed by specialised biological hardware. Even though these high-dimensional topological-spectra resist complete linguistic extraction, their operational source-code remains permanently inscribed within deep genetic syntax of human life.

Hypothesis of Gravitational Shielding Derivation (SRE)

Version: 1.0

- Basis: the following articles in the manuscript collection `The Book of the Void` of *Status-Relational Entropy (SRE) Dynamics* (https://doi.org/10.5281/zenodo.23275050) (all references in this document are given as **article name + section**, and no longer by line number)
- Terminology convention: uniformly expressed as **"gravitational shielding"**—shielding naturally produces anti-gravity/weightlessness, rather than "flipping the sign of gravity / introducing negative mass".
- Status: This document only performs derivation, order-of-magnitude criteria, and phenomenological interpretation.

Literature Correspondence Table (Articles and Sections Cited in This Document)

This document cites the following seven articles in the manuscript collection `The Book of the Void`. In the main text, sources are uniformly marked in the form 《short name》§section; the table below gives the correspondence between short names and full titles, so that readers can cross-check under any typesetting (including PDF).

Short name	Full article title (manuscript-collection title)	Sections used in this document
《Multidimensional Spacetime and Dynamical Gravity》	《Emergence of Multidimensional Spacetime and Dynamical Gravity via Regularized Causal Information Networks》 (Chinese title: 正则化因果信息网络下多维时空与动态变光速引力)	§2.2 Topological dissipation tensor and compensation operator; §2.3 Topological stiffness weights and baryonic center-of-mass redshift; §2.4 Spectral resonance deriving the dynamic compression coefficient; §5 Causal emergent gravity: a thermodynamic effect as a dissipation gradient
《Composite Elementary Particles》	《Composite Elementary Particles and the SRE-Dynamics Topological Paradigm of Relational-Space Emergence》 (Chinese title: 复合基本粒子与关系空间涌现的SRE动力学拓扑范式)	§1.3 Local cross-spectral operator and information-geometry mapping; §II Core paradigm: the three-in-one emergence jump and composite-particle generation; §Discussion · Model capability boundary
《Reconstruction of Classical Electrodynamics》	《A Basic Reconstruction of Classical Electrodynamics Based on Discrete Graph Topology and Bidirectional Causality》 (Chinese title: 基于离散图拓扑与双向因果性对经典电动力学的基础重构)	§4 Mass–energy equivalence $E = mc^2$ as a topological theorem
《Necessity of Turbulence Emergence》	《The Necessity of Turbulence Emergence and an Algebraic Computation Method Based on Discrete Microscopic Causal Statistics and Multidimensional Manifold Reconstruction》 (Chinese title: 基于离散微观因果统计与多维流形重构的湍流涌现必然性及代数计算方法)	§4.1 Operator 1: boundary expansion; §4.2 Operator 2: maximum-entropy pruning; §4.3 Gain Γ: exogenous Λ and endogenous $\lambda(n)$
《Logical-Depth Anchoring and Safety Shielding》	《On the Order-of-Magnitude Anchoring of the Logical Depth N and the Safety Shielding of Observation Protocols》 (Chinese title: 关于逻辑深度 N 的量级锚定与观测协议的安全性屏蔽)	Section 3 Core safety adjudication: the physical-level "read-only" restriction

| **《Complete Characterization of the Nucleon》** | 《A Complete Characterization of the Nucleon in the SRE Framework: Open–Closed Two-State Skeleton Inversion, Quark Conjecture, and a Unified Formulation Validated by Nuclear Reactions》 (Chinese title: SRE 框架下核子的完整刻画：开闭二态骨架反演、夸克推测与核反应验证的统一表述) | §2.1 Binary self-organizing network and the sim_p generator |

| **《Graph-Operator Pipeline Whitepaper》** | 《Overall Framework Whitepaper of the General Graph-Operator Pipeline for Status-Relational Entropy (SRE) Dynamics (Academic Revision)》 (Chinese title: 状态-关系熵（SRE）动力学通用图算子流水线总体框架白皮书（学术修订版）) | §1.2 Operator 1: local graph-expansion operator; §3.1 Operator 5: endogenous-variable delay-calibration operator; §4.2 Morphological-morphism projection principle of macroscopic observables; §5.1 Operator 10: preemptive homological random-pruning operator; §5.3 Operator 8: local lock-free algebraic-valve equalization operator |

Reading convention: "Part I" appearing in the main text refers to the first part of this document (Gravitational Shielding Derivation); "Part II" refers to the second part (Reverse Interpretation of the Podkletnov Result). Cross-part citations are written as "Part I §n" or "Part II §n".

Part I　Gravitational Shielding Derivation

0 Summary of Conclusions

1. In SRE, gravity is not a fundamental field but a **thermodynamic aftereffect of local information-dissipation gradients** (*Multidimensional Spacetime and Dynamical Gravity* §5). Therefore "gravitational shielding" = **suppressing the network correlations capable of generating that dissipation gradient**, rather than canceling a fundamental force.
2. The dependence of the gravitational acceleration a_{SRE} on the network state can be decomposed into **three shieldable quantities**:
   - Main amplitude $\alpha_{\mathrm{scale}} \cdot \mathcal{W}_{ij}$ (suppressible, the only substantive channel)
   - Chiral phase term Γ_{chiral} (the only sign-reversible term, capable of providing net repulsion)
   - Uniform term $-\gamma c_{\mathrm{eff}}^2/4$ (a global term, not locally shieldable)
3. The main channel yields an **exact, dimensionless shielding factor** (Part I §3):

   $$\Xi \equiv \frac{\mathcal{W}_{ij}^{\mathrm{shielded}}}{\mathcal{W}_{ij}^{\mathrm{vacuum}}} = \frac{1+\ln(1+\sigma_z^{\mathrm{vacuum}})}{1+\ln(1+\sigma_z^{\mathrm{shielded}})} < 1$$

   The cost is that the **local measurement entropy must be amplified logarithmically**: $\Xi = 0.5$ requires σ_z to increase by about 180 times, and $\Xi = 0.1$ by about 9×10^3 times. $\mathcal{W} \to 0$ (complete weightlessness) corresponds to $\sigma_z \to \infty$, and is **unreachable**.
4. The compensation operator $\hat{\mathcal{C}}$ is **non-monotonic/oscillatory** with respect to the dissipation $\hat{\mathcal{D}}$ (period $1/\alpha_0$, the first maximum at $\hat{\mathcal{D}}_0^* = 1/(2\alpha_0)$; the derivative proof is given in Part I §5). However, under realistic parameters $\hat{\mathcal{D}} \sim 18$–$27 \gg 1/\alpha_0$, it falls into the strong-oscillation regime, where its envelope mean is pinned at $\langle\hat{\mathcal{C}}\rangle = 1/(2\alpha_0)$, and it is **not a clean local knob** (an honest conclusion).
5. The only engineerable actuation mechanism is **Op2 maximum-entropy pruning** (*Necessity of Turbulence Emergence* §4.2): set the channel spin to the multiplicative identity $+1$, thereby eliminating that channel's phase contribution **without breaking graph connectivity**. Because the interior of a mass closed loop is "non-addressable/non-writable" (read-only barrier, *Logical-Depth Anchoring and Safety Shielding* Section 3), **shielding can only be a boundary operation**—operating on the dissipation–compensation channel between the shielded region and the external network, rather than rewriting the internal closed loop.
6. The chiral term Γ_{chiral} can reverse sign to produce **net repulsion**, and its phase condition is strictly obtainable (Part I §4); but whether it can dominate depends on the not-yet-calibrated dimensionless α_{scale}: only when $\alpha_{\mathrm{scale}} \lesssim 8 \times 10^{-4}$ is the chiral repulsion of the same order as the Newtonian term. **Thus the door to "true anti-gravity" is ajar, and its hinge is the experimental anchoring of α_{scale}.**

1 Starting Point: Gravity as a Thermodynamic Aftereffect of Dissipation Gradients

The theoretical body (*Multidimensional Spacetime and Dynamical Gravity* §5, verbatim):

$$a_{\mathrm{SRE}}(r,z)=\begin{cases}-\dfrac{\alpha_{\mathrm{scale}}\cdot\mathcal{W}_{ij}}{r}-\dfrac{\gamma c_{\mathrm{eff}}(z)^2}{4} & \mathrm{Rank}=2,\ z<z^*\\[2ex] -\dfrac{2\,\alpha_{\mathrm{scale}}\cdot\mathcal{W}_{ij}}{r^2}-\dfrac{\gamma c_{\mathrm{eff}}(z)^2}{4}+\Gamma_{\mathrm{chiral}}(r) & \mathrm{Rank}=4,\ z\geq z^*\end{cases}$$

$$\Gamma_{\mathrm{chiral}}(r)=\xi(z)\cdot\frac{\sin\!\left(\pi\alpha_{0,\mathrm{dynamic}}\cdot 2\mu\right)}{r^2\cdot\ln(r/\ell_P)}$$

Supporting definitions (*Multidimensional Spacetime and Dynamical Gravity* §2.2–§2.4, verbatim):

$$\hat{\mathcal{D}}_{ij}=\ln\!\left(1+\frac{\sigma_{z,i}\cdot\sigma_{z,j}}{\epsilon_{\mathrm{mach}}}\right),\qquad \hat{\mathcal{C}}=\alpha_{0,\mathrm{dynamic}}^{-1}\cdot\sin^2\!\left(\pi\alpha_{0,\mathrm{dynamic}}\hat{\mathcal{D}}_{ij}\right)$$

$$R_{ij}^2\equiv\mathrm{Tr}\!\left(\hat{\mathcal{D}}_{ij}\cdot\hat{\mathcal{C}}\right)\cdot\exp(-\gamma\,\mu_{\mathrm{loss}})$$

$$\mathcal{W}_{ij}=\sqrt{\mathcal{C}_i\,\mathcal{C}_j}=\left[\left(1+\ln(1+\sigma_{z,i})\right)\left(1+\ln(1+\sigma_{z,j})\right)\right]^{-1/2}$$

$$\mu=\frac{\sum_i\mathcal{C}_i z_i}{\sum_i\mathcal{C}_i},\qquad \xi(z)\equiv\xi(\mu)=0.08\,e^{0.15\mu}$$

$$\alpha_{0,\mathrm{dynamic}}=\frac{\theta_{\mathrm{conformal}}}{\Delta z+\epsilon_{\mathrm{mach}}},\qquad \theta_{\mathrm{conformal}}\approx 0.82798,\qquad \gamma=0.0585=\frac{1}{2\pi e}$$

Ontological key point (*Multidimensional Spacetime and Dynamical Gravity* §5): "Gravity is not a fundamental field; it is a statistical thermodynamic aftereffect brought about by local information-dissipation gradients."—This statement directly provides the **physical entry point** for shielding: to shield gravity is essentially to suppress the **local dissipation tensor and its gradient**, which is determined by the measurement entropy σ_z and its spatial distribution.

2 Structural Decomposition of Gravity: Three Shieldable Quantities

Writing a_{SRE} as a "source × kernel" decomposition, and extracting the independently adjustable factors within the kernel:

Quantity	Expression	Dependence on network state	Shieldable?	Role
Main amplitude	$\alpha_{\mathrm{scale}} \cdot \mathcal{W}_{ij}$	$\mathcal{W}_{ij} = f(\sigma_{z,i}, \sigma_{z,j})$	✅ Suppressible	The only substantive channel
Chiral term	$\Gamma_{\mathrm{chiral}} = \xi(z)\sin(\cdot)/(r^2 \ln(r/\ell_P))$	Phase $\propto \alpha_0 \mu$	✅ Sign-reversible	The only source of net repulsion
Uniform term	$-\gamma c_{\mathrm{eff}}^2/4$	Global $c_{\mathrm{eff}}(z)$	❌ Not local	Background/expansion term

Key observation: $\mathcal{W}_{ij}$ is a **strictly decreasing** function of σ_z (*Multidimensional Spacetime and Dynamical Gravity* §2.3); while the dissipation $\hat{\mathcal{D}}_{ij}$ is a **strictly increasing** function of σ_z (*Multidimensional Spacetime and Dynamical Gravity* §2.2). Hence the single action of **injecting local measurement entropy** simultaneously:

- (i) directly lowers $\mathcal{W}_{ij} \to$ lowers the amplitude of a_{SRE};
- (ii) raises $\hat{\mathcal{D}}_{ij} \to$ shifts the operating point of the compensation operator (Part I §5).

The two channels share the same origin, so "gravitational shielding" has a unified operational definition in this theory: **raise the local measurement entropy σ_z of the target region (making its information more "blurred" and more decohered), and cut off its dissipation–compensation channel to the outside.**

---

3 Main Channel: $\mathcal{W}_{ij}$ Entropy Collapse (Exact, Quantifiable)

3.1 Shielding Factor

Let the endpoint vacuum measurement entropy be σ_z, and after shielding be σ_z'. From the definition of $\mathcal{W}_{ij}$:

$$\boxed{\Xi \equiv \frac{\mathcal{W}(\sigma_z')}{\mathcal{W}(\sigma_z)} = \frac{1+\ln(1+\sigma_z)}{1+\ln(1+\sigma_z')}}$$

Since both branches of a_{SRE} are proportional to $\mathcal{W}_{ij}$, **the relative gravity-weakening amount is exactly** $1-\Xi$, independent of α_{scale}, r, and z—determined purely by the local measurement entropy. This is the cleanest result of this derivation.

3.2 Inverse Design Equation

Given a target shielding factor Ξ, the required local measurement entropy:

$$\boxed{\sigma_z' = \exp\left(\frac{1+\ln(1+\sigma_z)}{\Xi} - 1\right) - 1}$$

3.3 Order-of-Magnitude Table (taking $\sigma_z = 0.01$, $1 + \ln(1.01) = 1.00995$)

| Target Ξ | Gravity weakening | Required σ_z' | Amplification relative to vacuum |

|---|---|---|---|

| 0.99 | 1% | 0.0204 | ×2.0 |

| 0.98 | 2% | 0.0310 | ×3.1 |

| 0.90 | 10% | 0.1300 | ×13 |

| 0.50 | 50% | 1.773 | ×177 |

| 0.10 | 90% | 8.95×10^3 | 9.0×10^5 |

Conclusion: Weight reduction is "logarithmic-scale investment for linear-scale weight reduction"; $\Xi \to 0$ requires $\sigma_z \to \infty$. **Within the main channel, complete weightlessness is unreachable; only asymptotic weight reduction is possible.** This falls in the same order-of-magnitude window as the "weight-reduction ceiling of about 2%" (Podkletnov scale) in the early conjecture.

---

4 Phase Channel: Γ_{chiral}, the Only Sign-Reversible Term

4.1 Strict Condition for Sign Reversal (Net Repulsion)

Γ_{chiral} carries its sign through $\sin(\pi\alpha_0 \cdot 2\mu)$ (*Multidimensional Spacetime and Dynamical Gravity* §5). Taking $\sin = -1$ (maximum outward contribution):

$$\pi\alpha_0 \cdot 2\mu = -\frac{\pi}{2} + 2\pi n \implies \boxed{\mu^* = \frac{4n-1}{4\,\alpha_{0,\text{dynamic}}}}$$

Then $\Gamma_{\text{chiral}}^{\max\to-} = -\dfrac{\xi(z)}{r^2 \ln(r/\ell_P)}$, producing a **net subtraction** from the attractive term. μ is the baryonic center-of-mass redshift ($\mu = \sum \mathcal{C}_i z_i / \sum \mathcal{C}_i$; definition in *Multidimensional Spacetime and Dynamical Gravity* §2.3), set by the weighted redshift of the local subgraph V_{slice}—**thus the phase is tunable** (by changing the redshift composition of the observed subgraph).

4.2 Order-of-Magnitude Criterion (Deciding Whether "Anti-Gravity" Is Feasible)

The ratio of the chiral term to the main attractive term (Rank=4):

$$\mathcal{R} \equiv \frac{|\Gamma_{\text{chiral}}|}{2\alpha_{\text{scale}}\mathcal{W}_{ij}/r^2} = \frac{\xi(z)\,\big|\sin(\cdot)\big|}{2\,\alpha_{\text{scale}}\,\mathcal{W}_{ij}\,\ln(r/\ell_P)} \xrightarrow{\sin=\pm1} \frac{\xi(z)}{2\,\alpha_{\text{scale}}\,\mathcal{W}_{ij}\,\ln(r/\ell_P)}$$

Substituting $\xi \approx 0.108$ ($\mu \approx 2$), $\mathcal{W} \approx 1$, $\ln(r/\ell_P) \approx 80$ ($r \sim 1\,\text{m}$):

$$\mathcal{R} \approx \frac{6.8 \times 10^{-4}}{\alpha_{\rm scale}}$$

Interpretation:

- If $\alpha_{\rm scale} \sim 1$: $\mathcal{R} \sim 10^{-3}$, the chiral term is a perturbation (phenomenologically manifesting as a small-amplitude oscillation of $G_{\rm eff}$; see the discussion of smooth $G_{\rm eff}$ oscillation in *Multidimensional Spacetime and Dynamical Gravity* §5), **insufficient to produce anti-gravity by itself**.
- If $\alpha_{\rm scale} \lesssim 8 \times 10^{-4}$: $\mathcal{R} \gtrsim 1$, the chiral repulsion is of the same order as the Newtonian term or even dominant → **net repulsion (anti-gravity) holds**.

Since $\alpha_{\rm scale}$ is a dimensionless free coupling in the theory, whose physical anchoring is deferred to experiment (see Part I §9), **"whether anti-gravity is reachable" is here precisely reduced to an experimentally decidable number**: the order of magnitude of $\alpha_{\rm scale}$.

4.3 Design Equation for Complete Cancellation

Requiring $a_{\rm SRE} = 0$ at some radius r (ignoring the uniform term):

$$\xi^* = \frac{2\,\alpha_{\rm scale}\,\mathcal{W}_{ij}\,\ln(r/\ell_P)}{\left|\sin(\cdot)\right|} \xrightarrow{\sin=1} 2\,\alpha_{\rm scale}\,\mathcal{W}_{ij}\,\ln(r/\ell_P)$$

Then, from $\xi = 0.08 e^{0.15\mu}$ (*Multidimensional Spacetime and Dynamical Gravity* §2.3), invert to find the required center-of-mass redshift:

$$\boxed{\mu_{\rm cancel} = \frac{20}{3}\ln\!\Big(\frac{\xi^*}{0.08}\Big)}$$

---

5 Geometric Criterion: Compensation-Operator Peak $\hat{\mathcal{D}}_0^* = \dfrac{1}{2\alpha_0}$

Differentiating $\hat{\mathcal{C}}(\hat{\mathcal{D}}) = \alpha_0^{-1}\sin^2(\pi\alpha_0\hat{\mathcal{D}})$ (definition in *Multidimensional Spacetime and Dynamical Gravity* §2.2):

$$\frac{d\hat{\mathcal{C}}}{d\hat{\mathcal{D}}} = \alpha_0^{-1}\cdot 2\sin(\pi\alpha_0\hat{\mathcal{D}})\cos(\pi\alpha_0\hat{\mathcal{D}})\cdot\pi\alpha_0 = \pi\,\sin\!\Big(2\pi\alpha_0\hat{\mathcal{D}}\Big)$$

- Extremum condition $\sin(2\pi\alpha_0\hat{\mathcal{D}}) = 0 \Rightarrow \hat{\mathcal{D}} = \dfrac{n}{2\alpha_0}$, $n \in \mathbb{Z}$;

- The first **maximum** (strongest compensation) is at $\hat{\mathcal{D}}_0^* = \dfrac{1}{2\alpha_0}$, with maximum value $\hat{\mathcal{C}}_{\max} = 1/\alpha_0$;
- The zeros are at $\hat{\mathcal{D}} = \dfrac{n}{\alpha_0}$ (compensation collapses to zero).

Numerically (SDSS window): $\Delta z \approx 0.03925$, $\alpha_0 = \theta/\Delta z = 21.094$ (*Multidimensional Spacetime and Dynamical Gravity* §2.4), hence

$$\hat{\mathcal{D}}_0^* = \frac{1}{2\alpha_0} = \frac{\Delta z}{2\theta} = \frac{0.03925}{2 \times 0.82798} = 0.02370$$

Meaning: The compensation operator is the "routing compute" that the network schedules to offset dissipation, and it is **non-monotonic**. If the local operating point can be placed at a zero of $\hat{\mathcal{C}}$ ($\hat{\mathcal{D}} = n/\alpha_0$), then the compensation in that region collapses → R^2 collapses → the gravity generated by the dissipation gradient is locally zeroed ("resonant zero-point shielding").

Honest order-of-magnitude limitation: Under realistic parameters $\epsilon_{\text{mach}} \sim 2 \times 10^{-16}$, $\sigma_z \sim 10^{-2}$, so

$$\hat{\mathcal{D}} = \ln\left(1 + \frac{\sigma_z^2}{\epsilon_{\text{mach}}}\right) \approx 2\ln\sigma_z + 36.0 \approx 26.8$$

and $1/\alpha_0 \approx 0.0474$, so $\hat{\mathcal{D}}$ is already in the **strong-oscillation regime** (about 565 oscillation periods). The region-averaged compensation is then pinned at the envelope mean:

$$\langle\hat{\mathcal{C}}\rangle = \alpha_0^{-1}\langle\sin^2\rangle = \frac{1}{2\alpha_0} = 0.02370$$

which **exactly equals** the value of $\hat{\mathcal{D}}_0^*$. To "lock" onto a particular zero, one would need to control $\ln\sigma_z$ to 10^{-8}-level precision—**practically uncontrollable**. Hence this channel of Part I §5 is **mathematically valid but engineering-fragile**: it explains why intrinsic oscillations and a discrete spectrum exist at the geometric scale, but it does not constitute a clean shielding knob.

---

6 Actuation Mechanism: Op2 Maximum-Entropy Pruning and "Boundary-ness"

6.1 Mechanism (*Necessity of Turbulence Emergence* §4.2–§4.3, verbatim)

Pruning probability (maximum-entropy pruning master equation):

$$p_{\text{prune}}(v_f, v_m) = 1 - \frac{1}{1 + \Gamma \cdot \dfrac{\mathcal{D}_s}{\mathfrak{B}}}$$

When pruning occurs ($\chi = 0$), **Paradigm B (eliminate–conduct)** is executed:

$$M_{n+1}(i,j) \leftarrow \chi \cdot M_{n+1}(i,j) + (1-\chi) \cdot 1$$

"That is, the channel spin is forcibly set to the multiplicative identity $+1$, thereby eliminating that channel's phase contribution from the product feedback loop without disconnecting graph connectivity. Surviving channels ($\chi = 1$) are injected with an antisymmetric chiral shear." (*Necessity of Turbulence Emergence* §4.2)

The pruning gain Γ has two modes (*Necessity of Turbulence Emergence* §4.3):

Mode	Pruning gain	Meaning
A (exogenous)	$\Gamma = \Lambda$	Λ directly serves as the phase-transition control quantity
B (adaptive)	$\Gamma = \Lambda \cdot \lambda(n)$	Superimposed with state-adaptive correction

6.2 Why This Is the Physical Actuation Channel for Shielding

- After the channel spin is set to $+1$ (multiplicative identity), that edge's contribution to the **product feedback loop** is eliminated, but **connectivity is preserved**—precisely the property required by "shield without destroying": cutting phase contribution ≠ cutting topology.
- Raising the local pruning gain Γ (i.e., raising the local Λ) → raises p_{prune} → more edges go dormant → the local multi-loop structure is weakened → the local ability to maintain a low σ_z (coherent measurement) decreases → σ_z rises → by Part I §3, $\Xi < 1$, so **gravity is locally weakened**.
- Therefore Λ **(or the local Γ) is precisely the control knob for "gravitational shielding"**.

6.3 Boundary Operation, Not Internal Rewriting (Key Clarification)

"Shielding" has two distinct meanings in this document (a trap, do not conflate):

1. **Observational shielding / read-only barrier**: the internal topology of a mass closed loop is "non-addressable, non-writable"; the outside can only manipulate the rendering layer as a whole (electromagnetic, scattering). → **We cannot directly "dismantle" the mass closed loop of a target object** (that is exactly the source of the infeasibility of topological decoupling; *Logical-Depth Anchoring and Safety Shielding* Section 3).
2. **Phase-contribution shielding / gating**: Op2 sets the edge's spin to the multiplicative identity, eliminating its phase contribution to the feedback loop while **not breaking global connectivity**. → This is an **accessible** operation.

Combining the two gives the correct picture: **gravitational shielding is a boundary operation**—it does not rewrite the object's interior (non-addressable), but performs gating (Op2) **on the dissipation–compensation channel between the object and the external network**, so that the region's dissipation gradient as measured externally is suppressed. By analogy: it is not "turning off the inside of the light bulb", but "inserting a controllable scattering layer between the light bulb and the observer".

---

7 Three Intensity Tiers and Hard Boundaries

7.1 Three Tiers

| Tier | Mechanism | Reachability | Order of magnitude |

|---|---|---|---|

| **A Weight reduction** | Local injection of σ_z (Part I §3 main channel) | Self-consistent, engineerable | $\Xi \sim 0.98$ (about 2%), rising logarithmically with injection |

| **B Shielded weightlessness** | Boundary Op2 gating + suppression of the local dissipation gradient; or (fragilely) locking a compensation zero | Theoretical boundary | $\Xi \ll 1$, but requires continuous power input to counter self-repair |

| **C Topological decoupling** | Directly release the mass closed loop ($m \equiv$ closed loop; *Reconstruction of Classical Electrodynamics* §4) | Hypothesis-level | Blocked by the read-only barrier, **no known stable mechanism** |

7.2 Hard Boundaries (Why This Is Not a Perpetual-Motion Machine)

1. **The uniform term is unshieldable**: $-\gamma c_{\text{eff}}^2/4$ is a global term that local operations cannot eliminate; any "local weightlessness" is superimposed on this background.
2. **The compensation operator is self-repair**: $\hat{\mathcal{C}}$ is the self-patching mechanism by which the network counters dissipation and maintains numerical stability (*Multidimensional Spacetime and Dynamical Gravity* §2.2). The σ_z you inject will be continuously "flattened" by the network, so shielding requires **continuous energy input**; there is no steady-state weightlessness with zero input.
3. **Order of magnitude of the chiral term**: unless $\alpha_{\text{scale}} \lesssim 8 \times 10^{-4}$, Γ_{chiral} is only a 10^{-3}-level perturbation and cannot produce anti-gravity by itself (Part I §4).
4. **Topological decoupling destroys the definition of mass**: locally releasing a closed loop locally cancels the very concept of "mass"; the current formalization cannot guarantee the stability and recoverability of the decoupled region.

---

8 Design Equations and Falsifiable Predictions

Summary of design equations

$$\Xi = \frac{1+\ln(1+\sigma_z)}{1+\ln(1+\sigma_z')} \qquad \sigma_z' = \exp\Big(\tfrac{1+\ln(1+\sigma_z)}{\Xi} - 1\Big) - 1$$

$$\hat{\mathcal{D}}_0^* = \frac{1}{2\alpha_{0,\text{dynamic}}} = \frac{\Delta z}{2\theta_{\text{conformal}}} \qquad \mu^* = \frac{4n-1}{4\alpha_{0,\text{dynamic}}} \qquad \mu_{\text{cancel}} = \frac{20}{3}\ln\Big(\frac{\xi^*}{0.08}\Big)$$

Falsifiable predictions

1. **Shielding amount is independent of the absolute gravitational constant**: under a fixed increment of σ_z, the gravity-weakening ratio $1-\Xi$ should be independent of the test mass and position (universality). Any "material-dependent" additional shielding would require introducing a mechanism beyond $\mathcal{W}_{ij}$.
2. **Logarithmic law**: the weight-reduction effect saturates with injected entropy in a $1/(1+\ln)$ form—the marginal benefit of increasing injection diminishes (distinguishable from a "linear response" prediction).
3. **Phase reversal**: in the Rank=4 region ($z \geq z^*$), when the local center-of-mass redshift is tuned to μ^*, a sign flip/zero of G_{eff} should appear (corresponding to the smooth G_{eff} oscillation described in *Multidimensional Spacetime and Dynamical Gravity* §5).
4. α_{scale} **criterion**: if experiments measure $\alpha_{\text{scale}} \gtrsim 10^{-3}$, then "net anti-gravity" is falsified within SRE (only weight reduction is possible); if $\alpha_{\text{scale}} \ll 10^{-3}$, then chiral repulsion is the dominant channel.

---

9 Relation to the Uncalibrated α_{scale}

α_{scale} "appears" throughout the text but is never "calibrated" (see the localization conclusion from the previous round); moreover, according to the SRE axioms, **before anchoring, all quantities are dimensionless numbers**. This brings one advantage and one cost:

- **Advantage**: The shielding factor Ξ of the Part I §3 main channel is independent of α_{scale}, so the "weight-reduction ratio" is a testable prediction that can be **given prior to anchoring**.
- **Cost**: "whether chirality can dominate" in Part I §4, and "whether net anti-gravity is truly possible", **depend uniquely on the order of magnitude of** α_{scale}—which must be fixed only after future experiments (Newtonian-limit calibration).

Therefore this derivation compresses "anti-gravity" from a slogan into a definite experimentally-decidable numerical criterion: **the order of magnitude of** α_{scale}.

---

10 Open Questions (To-Do)

1. An explicit mapping between Λ (the local pruning gain) and macroscopic observables (σ_z injection rate, shielding power).
2. The **construction and reachability** of Op2 local gating: how the Op1 frontier (node injection) delivers the control signal to the target boundary while being constrained by the read-only barrier. → **Completed; see Part I §11**.
3. Chiral-phase tunability: the specific observables and operating window for setting μ^* by changing the subgraph redshift composition.
4. The **stability and dissipation cost** of the shielded region: the lower bound on continuous power required to counter the self-repair of $\hat{\mathcal{C}}$.

---

11 Op2 Local Gating Construction and Op1 Reachability (Execution Plan)

11.1 Three Reachability Constraints (Primary Sources)

| # | Constraint | Content | Source |

|---|---|---|---|

| 1 | **Read-only inheritance** | $M_{n+1}[1{:}n, 1{:}n] \equiv M_n$: once realized, the historical sub-block cannot be retroactively modified (with axiomatic status) | *Necessity of Turbulence Emergence* §4.1 Operator 1 |

| 2 | **Frontier-size upper bound** | $M_K \le K_0 \ll n$: each step acts on only $O(1)$ frontier channels, decoupled from the network-wide size | *Graph-Operator Pipeline Whitepaper* §5.3 Operator 8 |

| 3 | **Read-only/causal barrier** | The deep interior topology of the target closed loop is non-addressable and non-writable; $\ell_{\min}$ is the lower bound of addressable scale at the rendering layer | *Logical-Depth Anchoring and Safety Shielding* Section 3 |

> Key distinction (*Logical-Depth Anchoring and Safety Shielding* Section 3): the barrier **does not prevent ordinary physical manipulation of the entire topological closed loop at the rendering layer** (electromagnetic fields, scattering); it only forbids **penetrating the closed loop to rewrite its deep interior topology**.

Corollary 11.1 (Reachability Theorem): Op2 gating **cannot act on the interior closed loop of the target object**; it can only act on the **boundary channel** between "the target region ↔ the external network"; moreover, by constraint 1, **existing old edges cannot be rewritten**—the control signal must be fed in via the Op1 **newly added frontier nodes**. ⇒ Shielding is a **boundary-layer, incremental** operation.

11.2 Four-Step Construction of Local Gating (Aligned with the Operator Pipeline)

Goal: on the boundary shell Ω_∂ surrounding the target region, set **redundant** dissipation–compensation channels to dormant ($\chi = 0$, spin $\to +1$), while preserving cut-set connectivity.

Step 1 Op1 access (Operator 1 boundary expansion $\mathcal{G}_{n\to n+1}$, *Necessity of Turbulence Emergence* §4.1):

$$M_{n+1} = \begin{pmatrix} M_n & \mathbf{x}_{n+1} \\ \mathbf{x}_{n+1}^T & y_{n+1} \end{pmatrix}, \qquad M_{n+1}[1{:}n, 1{:}n] \equiv M_n$$

Push a ring of frontier nodes $\mathcal{V}_\partial$ into the perimeter of the target region (new rows/columns; history is read-only). **What is addressable is only these newly added boundary nodes, not the object's interior**—this is precisely the only legitimate interface that bypasses the barrier.

Step 2 Op2 gating (Operator 2 maximum-entropy pruning, *Necessity of Turbulence Emergence* §4.2):

$$p_{\text{prune}} = 1 - \frac{1}{1 + \Gamma_\partial \cdot \mathcal{D}_s/\mathfrak{B}} \xrightarrow[\Gamma_\partial\, \mathcal{D}_s/\mathfrak{B} \gg 1]{} 1, \qquad M(i,j) \leftarrow \chi\, M(i,j) + (1-\chi)\cdot 1$$

On Ω_∂, raise the local pruning gain to Γ_∂, and roll the gate χ edge by edge; boundary edges with $\chi = 0 \to +1$ (**conducting but forming no loop**; *Complete Characterization of the Nucleon* §2.1).

Step 3 Safety exemption (Operator 10, *Graph-Operator Pipeline Whitepaper* §5.1): for the **bridge edges** on Ω_∂, enforce $\boldsymbol{\chi}_{e_{\text{bridge}}} \equiv 1$, preserving

$$\lambda_2(n+1) \;\geq\; \lambda_2(n) > 0$$

⇒ The graph does not tear, and global connectivity is preserved.

Step 4 Effective-impedance screening (also belonging to the effective topological-impedance criterion of Operator 10, *Graph-Operator Pipeline Whitepaper* §5.1): use $\mathbf{L}_n^+$ to compute the effective topological impedance of candidate edges

$$\mathbf{Z}_{\text{eff}}(u, v_f) = (\mathbf{e}_u - \mathbf{e}_{v_f})^T \, \mathbf{L}_n^+ \, (\mathbf{e}_u - \mathbf{e}_{v_f})$$

Only low-impedance (non-bridge) edges are allowed to go dormant. This gives the **geometric criterion** for gating: shielding only consumes "redundant channels" and does not touch "unique channels".

11.3 Quantitative Reachability Bound (Upper Limit of Shieldable Fraction)

Let the boundary shell Ω_∂ have $|V_\partial|$ nodes and $|E_\partial|$ edges. Operator 10 requires retaining a spanning tree ($|V_\partial| - 1$ edges), so the **number of dormable edges** is

$$N_{\text{gate}} \;\leq\; |E_\partial| - (|V_\partial| - 1)$$

Upper limit of the shieldable channel fraction

$$\boxed{\frac{N_{\text{gate}}}{|E_\partial|} \;\leq\; 1 - \frac{|V_\partial| - 1}{|E_\partial|} \;=\; 1 - \frac{2}{\langle k \rangle}}$$

($\langle k \rangle$ is the boundary average degree.) Example: $\langle k \rangle = 4$ ⇒ at most **50%** of boundary channels can be shielded per operation; $\langle k \rangle = 6$ ⇒ upper limit 67%.

Meaning (falsifiable): The upper limit of shielding capability is determined by the **boundary topological redundancy**, not by energy alone; low-connectivity (near-bridge) boundaries are **structurally unshieldable**.

11.4 Control-Signal Path and Diffusion Cost

The transfer chain of the control quantity Γ_∂ (Mode A: $\Gamma = \Lambda$; Mode B: $\Gamma = \Lambda\lambda(n)$, *Necessity of Turbulence Emergence* §4.3):

$$\Lambda_{\text{local}} \xrightarrow[\text{Op1}]{} \mathcal{V}_{n+1}\ (\text{frontier-variable injection}) \xrightarrow[\text{Op2}]{} \Gamma \text{ acts on candidate frontier edges} \xrightarrow[\text{Op6/10}]{} \Omega_\partial \text{ executes gating}$$

Key costs (from constraints 1+2):

- **Each step affects only $M_K \le K_0$ channels** ⇒ gating the entire shell requires $N_{\text{steps}} \ge M_\partial / K_0$ ⇒ shielding is **established diffusively and gradually**, with a time cost $\propto$ boundary area/perimeter, and **cannot be instantaneous**.
- Each step of frontier expansion "injects new random differences and expands outward" (*Complete Characterization of the Nucleon* §2.1)—the advance itself raises the local σ_z. **Two-sidedness**: on the positive side, it directly contributes to the Part I §3 main channel ($\Xi < 1$); on the negative side, the injected differences are **uncontrollable random quantities**, making the shielding band fluctuate randomly frontier by frontier.
- Because of read-only inheritance (constraint 1), shielding is **non-rollbackable**—to undo shielding, a new frontier must be built; old edges cannot be reverted.

11.5 Why Tier C (Topological Decoupling) Is Thoroughly Blocked

Tier C requires **dismantling an already-realized mass closed loop** ($m \equiv$ closed loop; *Reconstruction of Classical Electrodynamics* §4). But: the read-only barrier makes the deep interior topology non-addressable and non-writable (*Logical-Depth Anchoring and Safety Shielding* Section 3); and read-only inheritance $M_{n+1}[1{:}n, 1{:}n] \equiv M_n$ does not permit rewriting existing edges (*Necessity of Turbulence Emergence* §4.1).

Double blocking ⇒ within the theory there **exists no accessible operation** capable of dismantling an already-realized closed loop. Hence Tier C is not "hypothesis-level" but **mechanically excluded**. The only exception is the rendering-layer holistic manipulation permitted by *Logical-Depth Anchoring and Safety Shielding* Section 3 (electromagnetic/scattering)—that is the **transport/interaction** of mass, not the elimination of mass.

11.6 Summary (Grading of Executable Claims)

Claim	Status	Basis
Boundary shell can be gated by Op2	**Constructible**	Part I §11.2 four steps
Shieldable fraction $\le 1 - 2/\langle k \rangle$	**Structural theorem**	Part I §11.3
Shielding is gradual, non-instantaneous, non-rollbackable, and requires continuous power	**Constraint corollary**	Part I §11.4 + Part I §7
Interior closed loop can be dismantled (true anti-gravity/topological decoupling)	**Mechanically excluded**	Part I §11.5

Part II　Reverse Interpretation of the Podkletnov Result and the Difficulty of Reproduction

Premise (assuming the result holds): an object above a rotating YBCO superconducting disk loses about 2% of its weight (Podkletnov & Nieminen 1992); a double-layer disk produces a stronger effect; a later "impulse gravity generator" gives a directed force.

Task: not to question the experiment, but to explain **why it holds** within the SRE framework, and to give discriminable predictions—as well as why others find it hard to reproduce.

Part II-1 Pivot: Two Opposite Effects of Coherent Saturation

"Phase-coherent saturation ($\rho \to 1$)" has **two oppositely-signed** consequences in SRE; conflating them necessarily yields the wrong sign:

	Mechanism	Consequence
Interior effect	Closed-loop merging, combinatorial explosion of secondary feedback paths	Relational space collapses $D \to 0$, **mass rises nonlinearly** (*Composite Elementary Particles* §Discussion · Model capability boundary)
Boundary effect	Relative phases are pinned flat ⇒ the local **dissipation gradient is flattened**	The gravity source (= dissipation gradient, *Multidimensional Spacetime and Dynamical Gravity* §5) is suppressed ⇒ **weight reduction**

Weight reduction comes from the boundary effect, not the interior effect. This distinction is the pivot of the entire explanation chain.

Part II-2 Mechanism Chain (Five Steps)

1. **Superconductor = phase-rigid plate**: Cooper pairs drive the in-disk network into macroscopic phase locking ($\rho \to 1$; *Composite Elementary Particles* §1.3 and §II), pinning the channel relative phases to the same value.
2. **Equivalent to Op2 dormancy**: phase locking makes the disk↔outside boundary channels behave equivalently to dormant edges $M(i,j) \leftarrow +1$—"**conducting but forming no loop**" (*Complete Characterization of the Nucleon* §2.1; mechanism definition in *Necessity of Turbulence Emergence* §4.2). The disk becomes a **gating layer** that only conducts and does not generate multi-loop feedback.
3. **Dissipation gradient is short-circuited**: on the boundary shell Ω_∂, execute the four steps of Part I §11 (Op1 access → Op2 gating → Operator 10 preserving $\lambda_2 > 0$ → effective-impedance screening consuming only low-Z_{eff} redundant edges), reducing the local stiffness $\mathcal{W}_{ij}$ of the test mass.
4. **Rotation = chiral bias**: surviving channels are injected with an antisymmetric chiral shear (*Necessity of Turbulence Emergence* §4.2), pushing the phase $\pi\alpha_0 \cdot 2\mu$ away from zero, so that Γ_{chiral} acquires an **outward** component ⇒ a small anti-gravity term is superimposed, and a **directionality** along the rotation axis is given.
5. **Layer compounding**: each disk layer is a boundary shell, $\Xi_{\text{tot}} = \prod_k \Xi_k$ ⇒ the stronger double-layer effect is a **product law**.

Part II-3 Quantitative Self-Consistency

$$\Xi = \frac{1+\ln(1+\sigma_z)}{1+\ln(1+\sigma_z')}$$

2% weight reduction $\Rightarrow \Xi = 0.98 \Rightarrow \sigma_z$ need only be ×3. This belongs to **Tier A weight reduction** (Part I §7), matching the order of magnitude of the "weight-reduction ceiling of about 2%"—**this is precisely why the effect is small**.

Part II-4 Why Only About 2% (Hard Boundaries)

- Upper limit of the shieldable fraction $1 - 2/\langle k \rangle$ (capped by boundary redundancy; Part I §11.3);
- Limited coherence budget (the phase difference the disk can rearrange is finite);
- Must continuously counter the self-repair of $\hat{\mathcal{C}} \Rightarrow$ **it disappears upon stopping rotation/heating** (consistent with the experiment's requirement of "low temperature + rotation").

Part II-5 Discriminable Predictions (If the Result Holds, SRE Requires These to Co-occur)

1. **Tc discontinuity**: the effect vanishes **abruptly** as the temperature crosses the superconducting critical temperature (coherent saturation is a phase transition, not a gradual change).
2. **Chirality**: the effect depends on the **direction** of rotation (sign reversal), and is monotonic with rotation speed.
3. **Product law**: multiple layers superimpose as $\prod \Xi_k$, **not** linear addition.
4. **Material independence**: the weight-reduction ratio of the shielded object is independent of its own composition/density (Ξ depends only on σ_z).
5. **Counterintuitive**: during shielding, the **superconducting disk itself** should have its inertial mass **slightly increased** (the mass surge from interior closed-loop merging; *Composite Elementary Particles* §Discussion · Model capability boundary), while the object above the disk becomes lighter—this is the discriminant that separates the two opposite effects.
6. **Geometry**: weight reduction occurs only **above the disk** and varies with the rotation-axis orientation (boundary gating + chiral directionality).

Part II-6 Bounds and Extensions

- Belongs to **gravitational shielding (Tier A/B)**, not Tier C topological decoupling (the latter is **doubly excluded** by the read-only barrier + read-only inheritance; Part I §11.5).
- The "impulse gravity generator" = the Γ_{chiral} **traveling wave** emitted by a coherent chiral source (a directed phase pulse of an open loop), isomorphic to the SRE picture "closed→open = luminescence/radiation" (*Reconstruction of Classical Electrodynamics* §4).
- The same mechanism as Section I of the early conjecture, "superconductivity isomorphism: back-reaction of causal developmental probability"; here it lands on the concrete operators $\mathcal{W}_{ij}$ and Op2.

Part II-7 Why Others Find It Hard to Reproduce (Same SRE Framework)

| # | Irreproducibility mechanism | SRE basis | What reproduction must control |

|---|---|---|---|

| 1 | Boundary redundancy is an **exogenous local quantity**, differing across samples ⇒ different Ξ | Ξ depends on σ_z (Part I §3); shieldable fraction $\leq 1 - 2/\langle k \rangle$ (Part I §11.3) | Single-phase single crystal, specified crystallographic orientation/twin structure |

| 2 | Coherent saturation is a **phase-transition criticality**; off the critical point it fails avalanche-like | $\rho \to 1$ saturation (*Composite Elementary Particles* §1.3, §II) | Strict temperature control, pushing $\langle \rho \rangle$ to the critical neighborhood |

| 3 | **Path dependence** (read-only inheritance ⇒ hysteresis, non-ergodicity) | Op1 read-only inheritance (*Necessity of Turbulence Emergence* §4.1) | Reproduce the **cooling/rotation time-sequence protocol**, not just the final-state parameters |

| 4 | Frontier expansion **carries its own random-difference injection**; the small effect is buried by noise | The sim_p generator injects random differences at each step (*Complete Characterization of the Nucleon* §2.1) | Sufficient averaging / per-frontier locking |

| 5 | $\hat{\mathcal{C}}$ **self-repairs**; requires continuous power + the correct sampling time window | Part I §7 hard boundaries | Steady-state power + correct duty cycle |

| 6 | **The chiral phase is signed and exogenous**; if unmatched, the directional part vanishes or reverses | $\Gamma_{\text{chiral}} \propto \sin(\pi\alpha_0 2\mu)$ (*Multidimensional Spacetime and Dynamical Gravity* §5) | Phase matching (rotation-axis orientation / subgraph window) |

| 7 | The order of magnitude is only 2%, and **boundary effects are easily mechanically short-circuited** | 2% ≈ buoyancy/thermal drift; boundary gating (Part I §11) | Decoupled support + differential measurement |

Falsifiability boundary (against excuse-making): if some experiment **simultaneously** satisfies the above 7 conditions (especially $\langle \rho \rangle$ in the critical neighborhood, phase matching, and consistent historical protocol) and still measures no effect, then **Tier A weight reduction is falsified**—"hard to reproduce" ends here and can no longer serve as an explanation.

Appendix A Symbol Table

| Symbol | Meaning | Source (article · section) |

|---|---|---|

| a_{SRE} | SRE gravitational acceleration, in two branches Rank=2/4 | *Multidimensional Spacetime and Dynamical Gravity* §5 |

| α_{scale} | Dimensionless gravitational coupling (uncalibrated, pending experimental anchoring) | *Multidimensional Spacetime and Dynamical Gravity* §5 |

| $\mathcal{W}_{ij}$ | Topological stiffness weight (macroscopic mass equivalent), a decreasing function of σ_z | *Multidimensional Spacetime and Dynamical Gravity* §2.3 |

| $\hat{\mathcal{D}}_{ij}$ | Topological dissipation tensor | *Multidimensional Spacetime and Dynamical Gravity* §2.2 |

| $\hat{\mathcal{C}}$ | Dynamic topological compensation operator (non-monotonic/oscillatory) | *Multidimensional Spacetime and Dynamical Gravity* §2.2 |

| R_{ij}^2 | Squared macroscopic metric distance | *Multidimensional Spacetime and Dynamical Gravity* §2.2 |

| γ | Handshake-latency coefficient $= 1/(2\pi e) = 0.0585$ | *Multidimensional Spacetime and Dynamical Gravity* §2.2 |

| $\xi(z)$ | Higher-order loop correction $= 0.08e^{0.15\mu}$ | *Multidimensional Spacetime and Dynamical Gravity* §2.3 |

| Γ_{chiral} | Chiral correction (the only sign-reversible term) | *Multidimensional Spacetime and Dynamical Gravity* §5 |

| μ | Baryonic center-of-mass redshift | *Multidimensional Spacetime and Dynamical Gravity* §2.3 |

| $\alpha_{0,\text{dynamic}}$ | Dynamic compression coefficient $= \theta/\Delta z$ | *Multidimensional Spacetime and Dynamical Gravity* §2.4 |

| $\theta_{\text{conformal}}$ | Conformal geometric exponent ≈ 0.82798 | *Multidimensional Spacetime and Dynamical Gravity* §2.4 |

| Λ, Γ | Pruning gain / phase-transition control quantity (shielding knob) | *Necessity of Turbulence Emergence* §4.3 |

| ℓ_P | Planck length (emergent UV boundary) | *Multidimensional Spacetime and Dynamical Gravity* §5 |

Appendix B Comparison with the Early Conjecture

| Early conjecture (Theoretical Synopsis Based on Status) | Corresponding item in this design | Verdict |

|---|---|---|

| Eliminating local observation probability → local G drift | Raising local $\sigma_z \to \mathcal{W}_{ij} \downarrow$ (Part I §3) | Isomorphic, now quantified (Ξ) |

| Back-reaction of causal developmental probability (superconductivity isomorphism) | Network self-repair $\hat{\mathcal{C}}$ + Op2 gating | Corresponding, belongs to Tier B |

| Weight-reduction ceiling of about 2% (Podkletnov level) | $\Xi \approx 0.98$ requires σ_z ×3 | Order of magnitude consistent |

| Ramanujan theta + Bernoulli nested fitting | Not incorporated into this design (belongs to the fitting layer) | To-do |

Basis (article · section correspondence): gravity source = dissipation gradient → *Multidimensional Spacetime and Dynamical Gravity* §5; $\mathcal{W}_{ij}$ → *Multidimensional Spacetime and Dynamical Gravity* §2.3; dissipation tensor and compensation operator → *Multidimensional Spacetime and Dynamical Gravity* §2.2; coherent saturation → *Composite Elementary Particles* §1.3, §II, §Discussion · Model capability boundary; Op2 dormancy mechanism → *Necessity of Turbulence Emergence* §4.2, and see *Complete Characterization of the Nucleon* §2.1; Op1 read-only inheritance → *Necessity of Turbulence Emergence* §4.1; frontier random-difference injection → *Complete Characterization of the Nucleon* §2.1; mass–energy equivalence / closed–open → *Reconstruction of Classical Electrodynamics* §4; read-only barrier → *Logical-Depth Anchoring and Safety Shielding* Section 3; permeability → *Graph-Operator Pipeline Whitepaper* §3.1 Operator 5; frontier-size upper bound and safety exemption/impedance criterion → *Graph-Operator Pipeline Whitepaper* §5.1 Operator 10, §5.3 Operator 8.