



Bispecific T-cell engagers From receptor to systems' level modeling to rationalize dose-effect relationships


No need to take
pictures: get the slides



Alexander Kulesza^{1,2}, Wilbert de Witte¹,
Luis David Jimenez Franco¹, Venetia
Karamitsou¹, Stephan Schaller¹

¹ ESQlabs, Saterland, Germany

² University Namur, Belgium



Bispecific T-cell engagers

From receptor to systems' level modeling to rationalize dose-effect relationships

Alexander Kulesza^{1,2}, Wilbert de Witte¹,
Luis David Jimenez Franco¹, Venetia Karamitsou¹, Stephan Schaller¹

¹ ESQlabs, Saterland, Germany

² University Namur, Belgium

Agenda

SLIDES



Disclosures:

AK is employee of ESQlabs GmbH, holds shares of CreativeQuantum GmbH and Ablatom SAS

3

01 BISPECIFIC T-CELL ENGAGERS? *Introduction, challenges and available models*

02 OPPORTUNITIES FOR MECHANISTIC MODELING *Literature, published models, current work*

- a** Cellular and systems level
A whole-body T-cell PBPK approach
- b** Receptor level and cellular level
Immune synapse modeling
- c** Cellular and systems level
Towards modeling of CRS incidence, severity and mitigation

WHAT ARE BISPECIFIC T-CELL ENGAGERS?

A very short introduction

Recombination of a Mixture of Univalent Antibody Fragments of Different Specificity¹

A. NISONOFF
M. M. RIVERS

*Department of Microbiology
University of Illinois
Urbana, Illinois*

Received March 27, 1961

oxidized, the recombined product contains a considerable amount of bivalent 5S "hybrid antibody" (6, 7).¹ The recombination of fragments of two different specificities appears to be essentially random, as shown by quantitative analysis of the

sults indicate that a reconstituted 5S antibody can induce passive hemagglutination, and that reconstituted 5S hybrid antibody preparations produce passive mixed agglutination of two visually distinguishable red cell types, each coated with a different antigen. With an artificial mixture of 5S antibodies (not hybridized), mixed agglutination does not occur; the two cell types are aggluti-


SEROLOGIC DEMONSTRATION OF DUAL SPECIFICITY OF RABBIT BIVALENT HYBRID ANTIBODY*

By H. H. FUDENBERG, M.D., GENEVIEVE DREWS, M.D., AND A. NISONOFF, Ph.D.

(From the Department of Medicine, University of California School of Medicine, San Francisco, and the Department of Microbiology, University of Illinois, Urbana)

PLATE 5


(Received for publication, September 4, 1963)



Obtained by peptic digestion (pepsin plus mild reduction of purified antibody) and then recombined by oxidation

WHAT ARE BISPECIFIC T-CELL ENGAGERS?

A very short introduction




A bispecific T-cell engager is a protein that simultaneously binds through a target antigen on a tumor cell and CD3 on a T-cell to form a TCR-independent artificial immune synapse and circumvent HLA restriction

Vafa (2020) 10.3389/fonc.2020.00446

WHAT ARE BISPECIFIC T-CELL ENGAGERS?

A very short introduction


6



Modified from Table 1 of
Ball et al. (2023) 10.1080/19420862.2023.2181016
See also Qi (2023) 10.1016/j.tips.2023.09.009

The bsTCE modeling landscape

7



The bsTCE modeling landscape

8




COMPARTMENTS	○	○	+	+	-	++	-	○	+
RECEPTORS	++	+	○	-	○	+	++	○	+
T-CELLS	-	○	-	++	++	○	-	-	++
CYTOKINES	-	-	○	○	-	○	-	++	-



A bsTCE modeling challenge: Cytokine release syndrome

9

- How can we better design the therapeutic window?
 - Early prediction of cytokine release profiles from affinities
- How do we proceed more efficiently from the starting dose towards the optimal doses?
 - Simulation of cytokine release under step-up dosing, escalation, CRS mitigation, with clinically relevant regimen
- How do we determine which have a low benefit/risk and how can we address that?
 - Explore between patient variability in CRS, support biomarker programs and use virtual twins



EMERGENT TOPICS

T-CELL
BIODISTRIBUTION

IMMUNE SYNAPSE


CRS INCIDENCE &
MITIGATION

Lowering CD3
affinity to reduce
CRS

Consequence
for T-cell
activation ??



Consequence for
tumor
penetration?

See Qin (2024)
10.1016/j.apsb.2024.03.027
Scheme adopted from:
Gibbs et al. (2020) 10.1002/jcph.1706



A WB-PBPK T-cell / bsTCE model in PK-Sim / MoBi

11



1

Implement T cell in existing mouse mAb PBPK model in PKSim

Literature based T cell PBPK model from Khot *et al.* 2019

Available data is exogenous active T cell

Optimization of T cell distribution properties in PKSim PBPK model backbone

2

Capture the distribution of antibody with CD3 binding only

Distribution of gD/CD3 TDB

Distribution data from Mandikian *et al.* 2018

Reproduce how CD3 affinity alters TDB distribution to T cell rich tissues

3

Capture the distribution of TDB with CD3 and tumor target binding

Distribution of HER2/CD3 TDB

Distribution data from Mandikian *et al.* 2018

Reproduce how CD3 affinity and target binding alters TDB distribution

Measurement and Quantitative Characterization of Whole-Body Pharmacokinetics of Exogenously Administered T Cells in Mice

Antari Khot, Satoko Matsueda, Veena A. Thomas, Richard C. Koya, and Dhaval K. Shah
Journal of Pharmacology and Experimental Therapeutics March 2019, 368 (3) 503-513; DOI: <https://doi.org/10.1124/jpet.118.252858>

Susilo (2022) www.page-meeting.org/?abstract=10024

LARGE MOLECULE THERAPEUTICS | APRIL 01 2018

Relative Target Affinities of T-Cell-Dependent Bispecific Antibodies Determine Biodistribution in a Solid Tumor Mouse Model **FREE**

Danielle Mandikian; Nene Takahashi; Amy A. Lo; Ji Li; Jeffrey Eastham-Anderson; Dionysos Slaga; Jason Ho; Maria Hristopoulos; Robyn Clark; Klara Totpal; Kedan Lin; Sean B. Joseph; Mark S. Dennis; Saileta Prabhu; Teemu T. Junttila; C. Andrew Boswell

ACoP15, Phoenix, AZ Nov. 10 - 13, 2024

EMERGENT TOPICS

T-CELL
BIODISTRIBUTION

IMMUNE SYNAPSE

CRS INCIDENCE &
MITIGATION

A WB-PBPK T-cell model in PK-Sim / MoBi

Implemented T cell PBPK model from Khot et al. into PKSim protein PBPK model backbone

- PKSim protein PBPK model has been validated across different species and different proteins (from small peptides to mAbs)

Single location-agnostic lymph node compartment

Two organ-specific parameters to calibrate:

- T cell transmigration rate: T cell migration from vascular to extravascular space
- T cell retention factor: accounting for exogenous T cell retention in the organ


Major limitations / assumptions:

- T cell elimination only in the lung vascular space
- No T cell proliferation (short time scale)

Niederalt (2018) 10.1007/s10928-017-9559-4

Khot (2019) 10.1124/jpet.118.252858

Susilo (2022) www.page-meeting.org/?abstract=10024



EMERGENT TOPICS

T-CELL BIODISTRIBUTION

IMMUNE SYNAPSE

CRS INCIDENCE & MITIGATION


A WB-PBPK T-cell model in PK-Sim / MoBi

Model simulation captures the distribution of exogenously administered active T cells

13

→ Base understanding on T cell distribution

Animal 1 2 3 Data digitized from Khot et al.



LN model parameters not calibrated using this data because

- Model is lumped location agnostic LN
- Data from Khot et al. 2019 is measurement from only the inguinal LN

EMERGENT TOPICS

T-CELL BIODISTRIBUTION

IMMUNE SYNAPSE

CRS INCIDENCE & MITIGATION

A WB-PBPK T-cell / bsTCE model in PK-Sim / MoBi


14

→ How affinities affect in vivo biodistribution

Higher CD3 affinity:

- Reduces plasma PK
- Increases HER2/CD3 TDB distribution to T-cell rich tissues
- Reduces HER2/CD3 TDB concentration in HER2+ tumor

Addition of HER2 target in the TDB reduces plasma and spleen concentration



Khot (2019) 10.1124/jpet.118.252858

Mandikian (2018) 10.1158/1535-7163.MCT-17-0657

Susilo (2022, 2024) www.page-meeting.org/?abstract=10024; to be submitted for publication



ACoP15, Phoenix, AZ Nov. 10 - 13, 2024

EMERGENT TOPICS

T-CELL
BIODISTRIBUTION

IMMUNE SYNAPSE

CRS INCIDENCE &
MITIGATION

Summary

15

- It is possible to use WB-PBPK platforms for describing T-cell and biomolecular distributions with minor modifications
- Kinetics / biodistribution of exogenous T-cells and labeled bsTCE impacted by T-cell biodistribution (as a function of affinities) can be described with that framework

Ongoing work

- Investigating synapse formation and TMDD in the WB-PBPK framework extending „affinity – effects“ to „affinity – on/off tumor effects“



M. Susilo
et al.

Genentech
A Member of the Roche Group

Next steps

- Fully modular bsTCE model in MoBi v12
- Integration with mechanistic and structured TGI models to describe indication-specific tumor penetration and modulation of the TME



EMERGENT TOPICS


T-CELL
BIODISTRIBUTION

IMMUNE SYNAPSE


CRS INCIDENCE &
MITIGATION

A basic CD3-Target receptor binding model

16



An often used model for the immune synapse is a simple version of the homogeneous, well stirred phase binding between the bispecific and freely diffusible receptors leading to ternary complex formation.



Models are
sensitive to

CD3/target affinity

Cell density

E:T ratio



EMERGENT TOPICS


T-CELL
BIODISTRIBUTION

IMMUNE SYNAPSE


CRS INCIDENCE &
MITIGATION

A basic CD3-Target receptor binding model

17



An often used model for the immune synapse is a simple version of the homogeneous, well stirred phase binding between the bispecific and freely diffusible receptors leading to ternary complex formation.



Models are
sensitive to

CD3/target affinity

Cell density

E:T ratio



EMERGENT TOPICS


T-CELL
BIODISTRIBUTION

IMMUNE SYNAPSE

CRS INCIDENCE &
MITIGATION

A basic CD3-Target receptor binding model

18



An often used model for the immune synapse is a simple version of the homogeneous, well stirred phase binding between the bispecific and freely diffusible receptors leading to ternary complex formation.

Then empirical „signals“ (often sigmoid functions) translate ternary complex into different downstream effects



EMERGENT TOPICS

T-CELL
BIODISTRIBUTION

IMMUNE SYNAPSE


CRS INCIDENCE &
MITIGATION

Physiology of the immune synapse

19

The picture of homogeneously distributed ternary complexes of freely diffusible species is not very biologically plausible as cellular rather than molecular kinetics govern the probability while complex cooperative multistep processes the strength of immune synapse formation.

There is quite some knowledge about the physiology of immune synapses



EMERGENT TOPICS


T-CELL
BIODISTRIBUTION

IMMUNE SYNAPSE

CRS INCIDENCE &
MITIGATION

Physiology of the immune synapse

20




CD3/target expression

Cell density

E:T ratio

*Integrated into an in vivo
physiological model it
could predict organ specific
responses, and optimal
regimen*

Liu et al (2023)
10.7554/eLife.83659



Optimizing Clinical Translation of Bispecific
T-cell Engagers through Context Unification
with a Quantitative Systems Pharmacology
Model

Xiaozhi Liao¹, Timothy Qi¹, Jiawei Zhou¹, Can Liu¹ and Yanguang Cao^{1,2,*}



ACoP15, Phoenix, AZ Nov. 10 - 13, 2024

EMERGENT TOPICS


T-CELL
BIODISTRIBUTION

IMMUNE SYNAPSE

CRS INCIDENCE &
MITIGATION

Physiology of the immune synapse

21



Optimizing Clinical Translation of Bispecific T-cell Engagers through Context Unification with a Quantitative Systems Pharmacology Model

Liu et al (2023)
10.7554/eLife.83659

Xiaozhi Liao¹, Timothy Qi¹, Jiawei Zhou¹, Can Liu¹ and Yanguang Cao^{1,2,*}



ACoP15, Phoenix, AZ Nov. 10 - 13, 2024

EMERGENT TOPICS


T-CELL
BIODISTRIBUTION

IMMUNE SYNAPSE

CRS INCIDENCE &
MITIGATION

Physiology of the immune synapse

22



The physiological IS has a „bullseye“ super structure“ with a distinct signaling and adhesion – dedicated regions

And the bsTCE is believed to be structurally very similar to this structure

Huppa (2013) 10.1016/B978-0-12-407707-2.00001-1
 Gao (2021) 10.3389/fimmu.2021.664329
 Robinson (2018) 10.1182/blood-2018-02-830992



EMERGENT TOPICS

T-CELL
BIODISTRIBUTION

IMMUNE SYNAPSE

CRS INCIDENCE &
MITIGATION

Summary


- The immune synapse needs to be modeled under consideration of receptor and cellular level mechanisms for optimal context unification
- Multi-step processes are already in place in different modeling approaches to take that into account

Next steps

- Kinetic proofreading explains specificity of antigen recognition that could apply to bsTCE synapses
- Testing of context unification model integrating a kinetic proofreading module for describing the adhesion kinetics once the synapse can be formed

25

Y. Cao lab



EMERGENT TOPICS

IMMUNE SYNAPSE

T-CELL
BIODISTRIBUTION

CRS INCIDENCE
& MITIGATION

Cytokine release syndrome is a condition that may occur after treatment with immunotherapy. It is caused by a large, rapid and systemic release of cytokines from immune cells affected by the immunotherapy.

Adopted from NCI Dictionary of Cancer Terms

26

Cytokine Release Syndrome Following Blinatumomab Therapy

[Pius E Ojemolon](#)^{1,✉}, [Sunaina Kalidindi](#)², [Taylor A Ahlborn](#)², [Osaigboka P Aihie](#)³, [Moyosoluwa I Awoyomi](#)¹

Editors: Alexander Muacevic, John R Adler

11-15% of patients who receive blinatumomab infusions develop CRS, and 2-5% have severe (grade ≥ 3) CRS


Blinatumomab-induced macrophage activating syndrome (MAS) in adult with B-cell acute lymphoblastic leukemia (B-ALL)

[Adam Braun](#)^{1,✉,#}, [Salman Otoukesh](#)^{1,#}, [Jose Tinajero](#)¹, [Guido Marcucci](#)¹, [Ibrahim Aldoss](#)¹

► [Author information](#) ► [Article notes](#) ► [Copyright and License information](#)

Other cells like (i.e. myeloid cells play a very important role)

Current concepts in the diagnosis and management of cytokine release syndrome

 Clinical Trials & Observations

Daniel W. Lee, Rebecca Gardner, David L. Porter, Chrystal U. Louis, Nabil Ahmed, Michael Jensen, Stephan A. Grupp, Crystal L. Mackall

Tocilizumab (anti IL6R/sIL6R) is used front line to treat CRS



EMERGENT TOPICS

IMMUNE SYNAPSE

T-CELL
BIODISTRIBUTION

CRS INCIDENCE
& MITIGATION


Dissecting the Mechanisms Underlying the Cytokine Release Syndrome (CRS) Mediated by T-Cell Bispecific Antibodies



Gabrielle Leclercq-Cohen¹, Nathalie Steinhoff¹, Lluïcia Albertí Servera², Sina Nassiri², Sabrina Danilin², Emily Piccione³, Emilio Yángüez¹, Tamara Hüsser¹, Sylvia Herter¹, Stephan Schmeing¹, Petra Gerber¹, Petra Schwalie², Johannes Sam¹, Stefanie Briner¹, Sylvia Jenni¹, Roberta Bianchi¹, Marlene Biehl¹, Floriana Cremasco¹, Katerina Apostolopoulou¹, Hélène Haegel², Christian Klein¹, Pablo Umaña¹, and Marina Bacac¹

IL-6 trans-signaling induces plasminogen activator inhibitor-1 from vascular endothelial cells in cytokine release syndrome

Sujin Kang¹, Toshio Tanaka², Hitomi Inoue¹, Chikako Ono³, Shoji Hashimoto⁴, Yoshiyuki Kioi¹, Hisatake Matsumoto⁵, Hiroshi Matsuura⁵, Tsunehiro Matsubara⁵, Kentaro Shimizu⁵, Hiroshi Ogura⁵, Yoshiharu Matsuura³, Tadamitsu Kishimoto⁶



Created with Biorender.com

Consensus

- involvement of T-cells as a major source of „activating stimulus“ (IFN γ and TNF α)
- Involvement of myeloid cells (macrophages and monocytes) producing IL6, IL10 and IL8

Hypotheses

- Wiring between positive and negative cytokine feedback loops
- Importance of between IL6 cis and trans signaling
- Involvement of endothelial cells and EC derived IL6




EMERGENT TOPICS

IMMUNE SYNAPSE

T-CELL
BIODISTRIBUTION

CRS INCIDENCE
& MITIGATION

Proposed ingredients of a model to test CRS hypotheses and drive data generation



Consensus

- Context dependent immune synapse formation driving T-cell activation and effector function
- involvement of T-cells as a major source of „activating stimulus“ (IFN γ and TNF α)
- Involvement of myeloid cells (macrophages and monocytes) producing IL6, IL10 and IL8

Hypotheses

- Wiring between positive and negative cytokine feedback loops
- Importance of between IL6 cis and trans signaling
- Involvement of endothelial cells and derived IL6

EMERGENT TOPICS


IMMUNE SYNAPSE

T-CELL
BIODISTRIBUTION

CRS INCIDENCE
& MITIGATION

A macrophage centric cytokine secretion model 31

We adopted and simplified a cellular-cytokine model that has good overlap with measured cytokine concentrations and added T-cell derived interferon as master macrophage activator



Complex, non-identifiable model as an example for this talk and to highlight workflow

In preparation for more detailed IL6 modeling, IL6 signaling has been dissociated from IL6 secretion.


EMERGENT TOPICS

IMMUNE SYNAPSE

T-CELL
BIODISTRIBUTION

CRS INCIDENCE
& MITIGATION

A macrophage centric cytokine secretion model 32




Remember?

11-15% of patients who receive blinatumomab infusions develop CRS, and 2-5% have severe (grade ≥ 3) CRS

cytokinesplasma_g_IL10IL6
cytokinesplasma_Khil_IL6IL6_ng
cytokinesplasma_g_IL6IL6
cytokinesplasma_kIL6byTNF_ng
cytokinesplasma_g_MIL10
cytokinesplasma_Khil_IL6TNF_ng
cytokinesplasma_g_IL6TNF
cytokinesplasma_kIL6_ng
cytokinesplasma_kIL10_ng
cytokinesplasma_Khil_MIL10_ng

We need a Vpop



EMERGENT TOPICS


IMMUNE SYNAPSE

T-CELL
BIODISTRIBUTION

CRS INCIDENCE
& MITIGATION

A macrophage centric cytokine secretion model 33

Cytokine Release: 5th-95th Percentile Range Across Virtual Patients




The model is too „well behaved“ to produce severe CRS, which is why we need another positive feedback loop leading to amplification in line with trans IL6 signaling

Remember?

11-15% of patients who receive blinatumomab infusions develop CRS, and 2-5% have severe (grade ≥ 3) CRS

cytokinesplasma_g_IL10IL6
cytokinesplasma_Khil_IL6IL6_ng
cytokinesplasma_g_IL6IL6
cytokinesplasma_kIL6byTNF_ng
cytokinesplasma_g_MIL10
cytokinesplasma_Khil_IL6TNF_ng
cytokinesplasma_g_IL6TNF
cytokinesplasma_kIL6_ng
cytokinesplasma_kIL10_ng
cytokinesplasma_Khil_MIL10_ng



EMERGENT TOPICS

IMMUNE SYNAPSE

T-CELL
BIODISTRIBUTION

CRS INCIDENCE
& MITIGATION



A detailed IL6 – (s)IL6R – (s)gp130 model

34

For proof of concept, we have integrated and extended the model by Rossi et al. (2022)
10.3389/fimmu.2022.919489

The balance of interleukin (IL)-6, IL-6-soluble IL-6 receptor (sIL-6R), and IL-6·sIL-6R·sgp130 complexes allows simultaneous classic and trans-signaling

[Paul Baran[‡]](#), [Selina Hansen[‡]](#), [Georg H Waetzig[§]](#), [Mohammad Akbarzadeh[‡]](#), [Larissa Lamertz[‡]](#), [Heinrich J Huber[¶]](#),
[M Reza Ahmadian[‡]](#), [Jens M Moll[‡]](#), [Jürgen Scheller^{‡,1}](#)



+ Additional between patient variability in sIL6 levels and sgp130 levels



EMERGENT TOPICS

IMMUNE SYNAPSE


T-CELL
BIODISTRIBUTION

CRS INCIDENCE
& MITIGATION


A detailed IL6 – (s)IL6R – (s)gp130 model

36

IL-6 Concentration in Virtual Population Stratified by Peak IL-6 Level



For a subset of virtual patients, IL6 crosses the threshold to activate endothelial cells which amplify IL6 secretion



Front-line tocilizumab disrupts excessive (i.e. trans) IL6 signaling

Could be also used to model risk-adapted preemptive tocilizumab

see Kadauke (2024) [10.1200/JCO.20.02477](https://doi.org/10.1200/JCO.20.02477)



EMERGENT TOPICS

IMMUNE SYNAPSE


T-CELL
BIODISTRIBUTION

CRS INCIDENCE
& MITIGATION

A detailed IL6 – (s)IL6R – (s)gp130 model

37

IL-6 Concentration in Virtual Population Stratified by Peak IL-6 Level




For a subset of virtual patients, IL6 crosses the threshold to activate

Functional *IL6R* 358Ala Allele Impairs Classical IL-6 Receptor Signaling and Influences Risk of Diverse Inflammatory Diseases

Ricardo C. Ferreira , Daniel F. Freitag , Antony J. Cutler, Joanna M. M. Howson, Daniel B. Rainbow, Deborah J. Smyth, Stephen Kaptoge, Pamela Clarke, Charlotte Boreham, Richard M. Coulson, Marcin L. Pekalski, Wei-Min Chen, Suna Onengut-Gumuscu, [...], John A. Todd [view all]

Published: April 4, 2013 • <https://doi.org/10.1371/journal.pgen.1003444>



Could be also used to model risk-adapted preemptive tocilizumab which is being tested in CART-trials

Kadauke (2024) [10.1200/JCO.20.02477](https://doi.org/10.1200/JCO.20.02477)



EMERGENT TOPICS

IMMUNE SYNAPSE

T-CELL
BIODISTRIBUTION

CRS INCIDENCE
& MITIGATION

Summary

38

- Expansion of T-cell-cytokine models can cover co-variability of a multitude of cytokines important for CRS
- Detailed IL6 modeling can help to explain between-patient differences in switching to an amplified / severe cytokine storm and action of anti-IL6 axis mitigation
- Looking beyond oncology / bsTCE models can be helpful to get inspiration for concrete models

Next steps

Repositories of immune digital twin literature, models and best practice

Systems immunogenicity of T-cell agonists



Anna
Niarakis


































UNIVERSITY of NEBRASKA-LINCOLN

Tomáš
Helikar

ACoP15, Phoenix, A



A fully modular approach PK-Sim / MoBi v12 (outlook)

Type	Module description	Building Blocks used in module
	bsAb ADME and binding <ul style="list-style-type: none"> ○ Target expression ○ Protein binding partner ○ FcRn mediated recycling 	       
	Extended physiology <ul style="list-style-type: none"> ○ Tumor ○ Lymph nodes ○ SC depots 	    
	Receptors <ul style="list-style-type: none"> ○ TMDD ○ Kinetic proofreading 	   
	T-cell module and IS <ul style="list-style-type: none"> ○ Transmigration ○ Retention factor ○ Cell encounter + Adhesion ○ Activation + Proliferation 	    
	Cytokine release <ul style="list-style-type: none"> ○ IL6 release 	   



PK-Sim/Base module



Extension module



https://github.com/Open-Systems-Pharmacology/Suite/releases/tag/v12.0_beta1



ACoP15, Phoenix, AZ Nov. 10 - 13, 2024

Take home messages

1. Quantitative systems pharmacology (QSP) models can derisk various aspects of bispecific T-cell engager design and development
2. Emerging topics for future QSP models are immune synapse modeling, extended cytokine network and effects modeling and T-cell/biTCE biodistribution modeling
3. Using modular workflows based on whole body PBPK and QSP extension modules allows for assembly of flexible „fit for purpose“ models streamlined to the question at hand

Acknowledgements:

- Susilo M, Li C, Chen C, Gadkar K, and team (Genentech)
- Jimenez L, Schaller S, deWitte W, Karamitsou V (ESQlabs)

Thank
you