


arXiv:2008.01610

# The joints problem for varieties

joint work with




Jonathan Tidor and Hung-Hsun Hans Yu


**Yufei Zhao**Massachusetts Institute  
of Technology

Joints problem What's the max # of joints  
that  $N$  lines in  $\mathbb{R}^3$  can make?


A joint is a point contained  
in 3 non-coplanar lines



Examples  
 $N$  lines  
 $\Theta(N^{3/2})$  joints



②  $k \sim \sqrt{2N}$  generic planes  
 $\leadsto$  pairwise form  
 $\binom{k}{2} \sim N$  lines  
& triplewise form  
 $\binom{k}{3} \sim \frac{\sqrt{2}}{3} N^{3/2}$  joints



Counting and cutting cycles of lines and rods in space\*

Bernard Chazelle

Dept. Comput. Sci., Princeton Univ., Princeton, NJ 08544, USA

Herbert Edelsbrunner

Dept. Comput. Sci., Univ. Illinois, Urbana, IL 61801, USA

Leonidas J. Guibas

Lab. Comput. Sci., MIT, Cambridge, MA 02139 and DEC SRC, Palo Alto, CA 94301, USA

Richard Pollack

Courant Inst. Math. Sci., New York Univ., New York, NY 10012, USA

Raimund Seidel

Dept. Comput. Sci. Electr. Engin., Univ. California, Berkeley, CA 94720, USA

Micha Sharir

Courant Inst. Math. Sci., New York Univ., New York, NY 10012, USA  
School Math. Sci., Tel Aviv Univ., Tel Aviv 69978, Israel

Jack Snoeyink

Dept. Comput. Sci., Stanford Univ., Stanford, CA 94305, USA

Communicated by Kurt Mehlhorn

Submitted 2 October 1990

Accepted 1 February 1991

Introduced the joints problem  
& proved  $O(N^{7/4})$  upper bd on #joints

... many subsequent improvements on until :

Algebraic methods in discrete analogs of the Kakeya problem

Larry Guth<sup>a</sup>, Nets Hawk Katz<sup>b,\*</sup>

**Theorem 1.1.** Any set of  $N$  lines in  $\mathbb{R}^3$  form at most  $O(N^{3/2})$  joints.

Recently Hans Yu & I improved the constant factor to optimal :  $\leq \frac{\sqrt{2}}{3} N^{3/2}$  joints

## Connections

- To **Kakeya** problem in analysis (Wolff)
- Finite field Kakeya problem (Dvir) ← polynomial method
- Erdős distinct distance problem (Guth-Katz)
- Multilinear Kakeya, "joints of tubes" (Bennett-Carbery-Tao, Guth)

Many extentions & generalizations of the joints problem

[Kaplan-Sharir-Shustin, Quilodrán] Simplification & extension to all dimensions

Ithm (joints of lines)  $N$  lines in  $\mathbb{F}^d$  have  $O_d(N^{\frac{d}{d-1}})$  joints.

arb. field  $\mathbb{F}$  tight const now known [Yu-Z]

Joints of flats : max # joints for  $N$  planes in  $\mathbb{F}^6$  ?

a point contained in a triple ↑  
of planes in spanning & indep directions

↑ 2-dim flats

Construction  $\Theta(N^{3/2})$  joints : generic 4-flats,  $\cap$  pair → planes,  $\cap$  triple → joints

Why I like this problem : 

- natural extension of joints
- a key step of pf of joints thm fails badly

# Incidence geometry for higher dimensional objects

[Solymosi - Tao]: nearly tight ( $\pm \text{oli}$  in exponent) bound for point-varietiy incidences in  $\mathbb{R}^d$  (in the spirit of Szemerédi-Trotter)

- Extension of the Guth-Katz polynomial partitioning method
  - ↑ Use bounded degree polynomials

[Walsh]: a different, algebraic, partitioning method

→ incidence among higher dim, higher degree varieties in  $\mathbb{F}^d$

Previous results:

[Yang]  $N$  planes in  $\mathbb{R}^6$  have  $N^{\frac{3}{2} + \text{oli}}$  joints

(technique: bounded deg partitioning, restrict to codim-1 variety)

Limitations ① Error term in exponent ② Only in  $\mathbb{R}$

[Yu-Z. / Carbery - Iliopoulos]  $N$  lines &  $M$  planes in  $\mathbb{F}^4$  make  $O(NM^{1/2})$  joints  
(plane-line<sup>2</sup>)

line-line-plane, in indep & spanning directions

Our results  $N$  planes in  $\mathbb{F}^6$  have  $O(N^{\frac{3}{2}})$  joints

[Tidor-Yu-Z.]  $N$  k-flats in  $\mathbb{F}^{mk}$  have  $O_{m,k}(N^{\frac{m}{m-1}})$  joints

A set of k-dim varieties in  $\mathbb{F}^{mk}$  of total degree  $N$  has  $O_{m,k}(N^{\frac{m}{m-1}})$  joints

A new way to apply polynomial method  
to higher dim objects

$p \in V_1, \dots, V_m$  regular point  
tangent spaces at  $p$  spanning & indep directions

Several sets of lines (& flats, varieties) [Conj by Carbery]

and count joints formed by taking one object from each set

Multijoints of lines (Iliopoulos  $\mathbb{R}^d$  &  $\mathbb{F}^3$ ; Zhang  $\mathbb{F}^d$ )

$L_1, L_2, \dots, L_d$  sets of lines in  $\mathbb{F}^d$

# joints formed by taking one line from each set is  $\lesssim_d (|L_1| \dots |L_d|)^{\frac{1}{d-1}}$ .

[Tidor-Yu-Z.] We extend from lines to varieties of arb dim

Joints of lines with multiplicities (Carbery conj; Iliopoulos  $\mathbb{R}^3$ ; Zhang  $\mathbb{F}^d$ )

$L_1, L_2, \dots, L_d$  sets of lines in  $\mathbb{F}^d$

$\sum_{\text{joints } p} \left( \#(l_1, \dots, l_d) \in L_1 \times \dots \times L_d \text{ making a joint at } p \right)^{\frac{1}{d-1}} \lesssim_d (|L_1| \dots |L_d|)^{\frac{1}{d-1}}$ .

[Tidor-Yu-Z.] Also ~~lines~~ varieties

Review of the proof of: [Kaplan-Sharir-Shustein, Quilodrán]

$N$  lines in  $\mathbb{R}^3$  have  $O(N^{3/2})$  joints

① Parameter counting:

using  $\dim \mathbb{R}[x_1, \dots, x_d] \leq_n = \binom{n+d}{d}$

deduce that  $\exists$  non-zero poly  $g$ ,  $\deg \leq C J^{1/3}$ , vanishing on joints


Take  $g$  with min deg.  $J = \# \text{joints}$

② Vanishing lemma: a single-variable polynomial cannot vanish more times than its degree

③ A joints-specific argument. If all lines have  $> C J^{1/3}$  joints, then vanishing lemma  $\Rightarrow g$  vanishes on all lines  $\Rightarrow \nabla g$  vanishes on all joints  $\Rightarrow$  one of  $\partial_x g, \partial_y g, \partial_z g$  is nonzero, lower deg & vanish on all joints

So some line has  $\leq C J^{1/3}$  joints. Remove this line & induction  $\square$

How to generalize vanishing lemma to 2-var polynomials?



Thm (Tidor-Yu-Z.)  $N$  planes in  $\mathbb{R}^6$  have  $O(N^{3/2})$  joints

Wishful thinking



only if we had something like ...

- every nonzero  $g(x,y)$  of  $\deg \leq n$  has  $\leq n^2$  zeros
- OR
- two polynomials, each  $\deg \leq N^{1/6}$ , vanishing at all joints and no common factors when restricted to each plane  
(related: inverse Bézout)

**Method of multiplicities**: ask a polynomial to vanish at each joint to some high order

By counting parameters, maybe hope for:



- Every nonzero  $g(x,y)$  of  $\deg \leq n$  vanish to order  $\geq s$  at  $\leq n^2$  pts.

Counterexample:  $g(x,y) = y^5$


Linear dependencies among vanishing conditions

$$\text{e.g. } g(p)=0, \partial_x g(p)=0, (\partial_{xx} - \partial_{xy})g(p)=0$$

**Key idea 1** Restricting to a plane for now

.	.	.
.	.	.
.	.	.
.	.	.


We will construct a set of  $\dim \mathbb{R}[x,y]_{\leq n} = \binom{n+2}{2}$  linearly indep vanishing conditions on  $\mathbb{R}[x,y]_{\leq n}$



Attached to each point  $p$  is a set of vanishing conditions for  $g \in \mathbb{R}[x,y]_{\leq n}$ :

$$g(p) = 0, \quad \partial_x g(p) = 0, \quad \partial_y g(p) = 0 \\ \partial_{xx} g(p) = 0, \quad \partial_{xy} g(p) = 0, \quad \partial_{yy} g(p) = 0, \quad \partial_{xxx} g(p) = 0, \dots$$

If we take all these conditions up to order  $n+1$  then any satisfying  $g \in \mathbb{R}[x,y]_{\leq n}$  must be zero.



The above vanishing conditions attached to several different points are lin. dep. as linear functionals on  $\mathbb{R}[x,y]_{\leq n}$

We will select a basis of linear functionals on  $\mathbb{R}[x,y]_{\leq n}$  via the following procedure.

### First attempt

Cycle through the points on the plane (say 100 pts)

$$P_1, P_2, P_3, \dots, P_1, P_2, P_3, \dots, P_1, P_2, P_3, \dots$$

$P_1$ : add vanishing condition  $g(p_1) = 0$

$P_2$ : add vanishing condition  $g(p_2) = 0$ , as long as it does not already follow from previously added vanishing cond.


$P_1$ : add a nonredundant subset of  $\partial_x g(p_1) = 0, \partial_y g(p_1) = 0$   
 ↑ none implied by other added + prev.added  
 ie. basis extension

$P_2$ : add a nonredundant subset of  $\partial_x g(p_2) = 0, \partial_y g(p_2) = 0$

The process assigns a total of  $\binom{n+2}{2}$  vanishing conditions each attached to a point

Can we control the # of vanishing attached to each pt?

Example



(This example also comes up for inverse Bézout; see Tao blog)

\*Key idea 2 Let some points get a head start

e.g.  $P_1, P_2, \dots, P_{50}, P_1, \dots, P_{50}, \dots, P_1, \dots, P_{50}, P_1, \dots, P_{100}, P_1, \dots, P_{100}, \dots$   
(100 pts)

Handicap  $\vec{\alpha}$  assigns an integer to each point

e.g.	points	a	b	c	d	e
handicap $\vec{\alpha}$	0	1	3	0	-1	

→ order: c c b c a b c d a b c d e a b c d e ..

Modify process of assigning vanishing conditions

c : add a nonredundant set of 0<sup>th</sup> order derivative vanishing @ c

c	1 <sup>st</sup>	c
b	0 <sup>th</sup>	b
c	2 <sup>nd</sup>	c
a	0 <sup>th</sup>	a
b	1 <sup>st</sup>	b
c	3 <sup>rd</sup>	c

Want a "good" choice of handicaps: treating all joints "fairly"

Hard to compute how # vanishing cond at each pt  $\vec{\alpha} \rightarrow$  <sup>distr of vanish cond</sup>

depends on the handicap.

- ① Monotonicity -  $\alpha_p \nearrow \Rightarrow$  # van. cond at  $p$  cannot  $\downarrow$
- ② Lipschitz continuity - small  $\Delta$  in handicap  $\rightsquigarrow$  small  $\Delta$  in # van cond
- ③ Bounded domain - suffices to consider handicaps with bounded values (else some pt gets 0 van. cond.)

\* Key idea 3 Decide on handicap choice later, implicitly (existence via compactness/smoothing)

Putting different planes together at joints of planes in  $\mathbb{R}^b$

Handicap  $\vec{\alpha} \in \mathbb{Z}^J$  assigns an integer to each joint

Separately for each plane  $F$ , apply above process to assign vanishing conditions restricted to  $F$

to joints on  $F$

assigning derivative operators

A new Vanishing lemma Given  $0 \neq g \in \mathbb{R}[x_1, \dots, x_b]_{\leq n}$ ,  
 $\exists$  joint  $p$ , contained in planes  $F_1, F_2, F_3$  (indep & spanning directions)  
& derivative operator  $D_i$  assigned to  $p$  on  $F_i$  (& likewise  $D_2, D_3$ )  
s.t.  $D_1 D_2 D_3 g(p) \neq 0$ .

By parameter counting,

$$\sum_{\text{joints } p} (\# \text{ choices of } D_1, D_2, D_3 \text{ at } p) \geq \dim \mathbb{R}[x_1, \dots, x_b]_{\leq n} = \binom{n+6}{6}$$

By compactness/smoothing,  $\exists$  handicap so that these terms are roughly all equal


Also recall that # van. cond. on each plane is exactly  $\binom{n+2}{2}$

Putting together + AM-GM  $\Rightarrow$  theorem □

## Joints of varieties

Flats: higher order directional directives along a flat

Varieties: derivatives in local coordinates

e.g.   $y = x^2 + y^2$  on the circle,

$$\begin{aligned} y &= x^2 + y^2 \\ &= x^2 + (x^2 + y^2)^2 \\ &= x^2 + (x^2 + (x^2 + y^2)^2)^2 = \dots \\ &= x^2 + x^4 + 2x^6 + \dots \end{aligned}$$

completion

Power series in local coord

2<sup>nd</sup> order derivative operator at the origin is  $\frac{1}{2} \frac{\partial^2}{\partial x^2} + \frac{\partial}{\partial y}$  (not  $\frac{\partial^2}{\partial x^2}$ )  
so that evaluations give linear functional on the  
space of regular functions

## Extension to arbitrary fields $\mathbb{F}$

When differentiating, we only care about coeff extraction

Hasse derivatives (formal algebraic derivatives)

Question Other applications of this variant  
of polynomial method for higher dim objects?