

ABOUT THE TEST FoundationOne®Liquid CDx is a next generation sequencing (NGS) assay that identifies clinically relevant genomic alterations in circulating cell-free DNA.

PATIENT	DISEASE Unknown primary carcinoma (NOS)	PHYSICIAN	ORDERING PHYSICIAN Yeh, Yi-Chen	SPECIMEN	SPECIMEN ID WLC 12/10/1971
	NAME Chien, Wen-Lin		MEDICAL FACILITY Taipei Veterans General Hospital		SPECIMEN TYPE Blood
	DATE OF BIRTH 10 December 1971		ADDITIONAL RECIPIENT None		DATE OF COLLECTION 01 December 2022
	SEX Male		MEDICAL FACILITY ID 205872		SPECIMEN RECEIVED 05 December 2022
	MEDICAL RECORD # 14285575		PATHOLOGIST Not Provided		

Biomarker Findings

Blood Tumor Mutational Burden - 4 Muts/Mb
Microsatellite status - MSI-High Not Detected
Tumor Fraction - Elevated Tumor Fraction Not Detected

Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

NF1I679fs*21

GNASR201H

TP53G245S

Report Highlights

- Targeted therapies with potential clinical benefit **approved in another tumor type**: Cobimetinib (p. [7](#)), Selumetinib (p. [7](#)), Trametinib (p. [8](#))
- Evidence-matched clinical trial options based on this patient's genomic findings: (p. [9](#))

BIOMARKER FINDINGS

Blood Tumor Mutational Burden -
4 Muts/Mb

Microsatellite status -
MSI-High Not Detected

Tumor Fraction -
Elevated Tumor Fraction Not Detected

THERAPY AND CLINICAL TRIAL IMPLICATIONS

No therapies or clinical trials. See Biomarker Findings section

MSI-High not detected. No evidence of microsatellite instability in this sample (see Appendix section).

Tumor fraction is considered elevated when ctDNA levels are high enough that aneuploidy can be detected. The fact that elevated tumor fraction was not detected in this specimen indicates the possibility of lower levels of ctDNA but does not compromise confidence in any reported alterations. However, in the setting of a negative liquid biopsy result, orthogonal testing of a tissue specimen should be considered if clinically indicated (see Biomarker Findings section).

GENOMIC FINDINGS

VAF%

NF1 - I679fs*21 0.99%

10 Trials [see p. 9](#)

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)

None

THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)

Cobimetinib

Selumetinib

Trametinib

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Nimesh Patel, M.D. | 12 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

GNAS - R201H **p. 5** **TP53 - G245S** **p. 6**

NOTE Genomic alterations detected may be associated with activity of certain approved therapies; however, the therapies listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and/or exhaustive. Neither the therapies nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies. Therapies contained in this report may have been approved by the US FDA or other national authorities; however, they might not have been approved in your respective country. In the appropriate clinical context, germline testing of *APC, ATM, BAP1, BRCA1, BRCA2, BRIP1, CHEK2, FH, FLCN, MEN1, MLH1, MSH2, MSH6, MUTYH, NF1, NF2, PALB2, PMS2, POLE, PTEN, RAD51C, RAD51D, RB1, RET, SDHA, SDHB, SDHC, SDHD, SMAD4, STK11, TGFBR2, TP53, TSC1, TSC2, VHL*, and *WT1* is recommended.

Variant Allele Frequency is not applicable for copy number alterations.


Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Nimesh Patel, M.D. | 12 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1517889-01



HISTORIC PATIENT FINDINGS

ORD-1517889-01
VAF%

Blood Tumor Mutational Burden

4 Muts/Mb

Microsatellite status

MSI-High Not Detected

Tumor Fraction

Elevated Tumor Fraction Not Detected

NF1	● I679fs*21	0.99%
GNAS	● R201H	0.25%
TP53	● G245S	0.55%

NOTE This comparison table refers only to genes and biomarkers assayed by prior FoundationOne®Liquid CDx or FoundationOne®CDx tests. Up to five previous tests may be shown.

For some genes in FoundationOne Liquid CDx, only select exons are assayed. Therefore, an alteration found by a previous test may not have been confirmed despite overlapping gene lists. Please refer to the Appendix for the complete list of genes and exons assayed. The gene and biomarker list will be updated periodically to reflect new knowledge about cancer biology.

As new scientific information becomes available, alterations that had previously been listed as Variants of Unknown Significance (VUS) may become reportable.

Tissue Tumor Mutational Burden (TMB) and blood TMB (bTMB) are estimated from the number of synonymous and non-synonymous single-nucleotide variants (SNVs) and insertions and deletions (indels) per area of coding genome sampled, after the removal of known and likely oncogenic driver events and germline SNPs. Tissue TMB is calculated based on variants with an allele frequency of $\geq 5\%$, and bTMB is calculated based on variants with an allele frequency of $\geq 0.5\%$.

Not Tested = not baited, not reported on test, or test preceded addition of biomarker or gene

Not Detected = baited but not detected on test

Detected = present (VAF% is not applicable)

VAF% = variant allele frequency percentage

Cannot Be Determined = Sample is not of sufficient data quality to confidently determine biomarker status

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Nimesh Patel, M.D. | 12 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1517889-01

BIOMARKER FINDINGS

BIOMARKER

Blood Tumor Mutational Burden

RESULT

4 Muts/Mb

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased blood tumor mutational burden (bTMB) may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1¹⁻³, anti-PD-1³⁻⁴, anti-PD-1/CTLA4 therapies⁵⁻⁶, anti-PD-L1/CTLA4 therapies⁷⁻¹⁰. A Phase 2 multi-solid-tumor trial showed that bTMB ≥ 16 Muts/Mb (as measured by this assay) was associated with improved survival from treatment with a PD-1 inhibitor alone or in combination with a CTLA-4 inhibitor⁵. In non-small cell lung cancer (NSCLC), multiple clinical trials have shown patients with higher bTMB derive clinical benefit from immune checkpoint inhibitors following single-agent or combination treatments with either CTLA4 inhibitors or chemotherapy, with reported high bTMB cutpoints ranging from 6 Muts/Mb-16 Muts/Mb^{1,8-10}. In head and neck squamous cell

carcinoma (HNSCC), a Phase 3 trial showed that bTMB ≥ 16 Muts/Mb (approximate equivalency ≥ 8 Muts/Mb as measured by this assay) was associated with improved survival from treatment with a PD-L1 inhibitor alone or in combination with a CTLA-4 inhibitor¹¹. In colorectal cancer (CRC), a Phase 2 study showed that bTMB ≥ 28 Muts/Mb (approximate equivalency ≥ 14 Muts/Mb as measured by this assay) was associated with improved OS from a PD-L1 inhibitor alone or in combination with a CTLA-4 inhibitor⁷.

FREQUENCY & PROGNOSIS

Average bTMB levels in solid tumors other than NSCLC have not been evaluated (PubMed, Mar 2022). Published data investigating the prognostic implications of TMB have mainly been investigated in the context of tissue TMB. In patients with NSCLC, increased TMB is associated with higher tumor grade and poor prognosis¹², as well as with a decreased frequency of known driver mutations in EGFR, ALK, ROS1, or MET (1% of high-TMB samples each), but not BRAF (10.3%) or KRAS (9.4%)¹³. Although some studies have reported a lack of association between smoking and increased TMB in NSCLC^{12,14}, several other large studies did find a strong link¹⁵⁻¹⁸. In CRC, elevated TMB is associated with a higher frequency of BRAF V600E driver mutations¹⁹⁻²⁰ and with

microsatellite instability (MSI)²⁰, which in turn has been reported to correlate with better prognosis²¹⁻²⁸. Although increased TMB is associated with increased tumor grade in endometrioid endometrial carcinoma²⁹⁻³² and bladder cancer³³, it is also linked with improved prognosis in patients with these tumor types³⁰.

FINDING SUMMARY

Blood tumor mutational burden (bTMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations from circulating tumor DNA in blood. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma³⁴⁻³⁵ and cigarette smoke in lung cancer³⁶⁻³⁷, treatment with temozolomide-based chemotherapy in glioma³⁸⁻³⁹, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes^{19,30,40-42}, and microsatellite instability (MSI)^{19,30,42}. High bTMB levels were not detected in this sample. It is unclear whether the bTMB levels in this sample would be predicted to be associated with sensitivity to PD-1- or PD-L1-targeting immune checkpoint inhibitors, alone or in combination with other agents¹⁻²⁴. Depending on the clinical context, TMB testing of an alternate sample or by another methodology could be considered.

BIOMARKER

Tumor Fraction

RESULT

Elevated Tumor Fraction Not Detected

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Specimens with elevated tumor fraction values have high circulating-tumor DNA (ctDNA) content, and thus high sensitivity for identifying genomic alterations. Such specimens are at low risk of false negative results. However, if elevated tumor fraction is not detected, it does not exclude the presence of disease burden or compromise the confidence of reported alterations. Tumor fraction levels currently have limited implications for diagnosis, surveillance, or therapy and should not

be overinterpreted or compared from one blood draw to another. There are currently no targeted approaches to address specific tumor fraction levels. In the research setting, changes in tumor fraction estimates have been associated with treatment duration and clinical response and may be a useful indicator for future cancer management⁴³⁻⁴⁸.

FREQUENCY & PROGNOSIS

Detectable ctDNA levels have been reported in a variety of tumor types, with higher tumor fraction levels reported for patients with metastatic (Stage 4) tumors compared with patients with localized disease (Stages 1 to 3)⁴⁹. Elevated tumor fraction levels have been reported to be associated with worse prognosis in a variety of cancer types, including pancreatic cancer⁵⁰, Ewing sarcoma and osteosarcoma⁵¹, prostate cancer⁴⁶, breast cancer⁵², leiomyosarcoma⁵³, esophageal cancer⁵⁴, colorectal

cancer⁵⁵, and gastrointestinal cancer⁵⁶.

FINDING SUMMARY

Tumor fraction provides an estimate of the percentage of ctDNA present in a cell-free DNA (cfDNA) sample. The tumor fraction estimate for this sample is based on the observed level of aneuploid instability. The tumor fraction algorithm utilized for FoundationOne Liquid CDx uses the allele frequencies of approximately 1,000 single-nucleotide polymorphism (SNP) sites across the genome. Unlike the maximum somatic allele frequency (MSAF) method of estimating ctDNA content⁵⁷, the tumor fraction metric does not take into account the allele frequency of individual variants but rather produces a more holistic estimate of ctDNA content using data from across the genome. The amount of ctDNA detected may correlate with disease burden and response to therapy⁵⁸⁻⁵⁹.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Nimesh Patel, M.D. | 12 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1517889-01

GENOMIC FINDINGS

GENE
NF1

ALTERATION

I679fs*21

TRANSCRIPT ID

NM_001042492.2

CODING SEQUENCE EFFECT

2033_2034insC

VARIANT CHROMOSOMAL POSITION

chr17:29553477

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence in neurofibromatosis Type 1-associated neurofibroma⁶⁰⁻⁶³, glioma or glioblastoma⁶³⁻⁶⁷, and non-small cell lung cancer⁶⁸, NF1 inactivation may predict sensitivity to MEK inhibitors such as cobimetinib, trametinib, binimetinib, and selumetinib. On the basis of limited clinical data⁶⁹⁻⁷¹ and preclinical data⁷²⁻⁷³, loss or inactivation of NF1 may predict sensitivity to mTOR inhibitors, including everolimus and temsirolimus. A preclinical study suggests that

combined mTOR and MEK inhibition is effective in a model of NF1-deficient malignant peripheral nerve sheath tumors (MPNST)⁷⁴.

FREQUENCY & PROGNOSIS

NF1 alterations are common in various solid tumors, with mutations being more frequent than gene loss, and are often seen in melanoma (18%), glioma (14%), gastrointestinal stromal tumor (GIST; 10%), and in salivary gland (6.1%), ovarian (5.4%), breast (4.8%), non-small cell lung (4.7%), non-melanoma skin (4.7%), and endometrial (3.7%) cancer⁷⁵. Published data investigating the prognostic implications of NF1 mutation in carcinomas are limited (PubMed, Jan 2022). NF1 loss was significantly associated with decreased overall and disease-specific survival in patients with grade II and III gliomas, but not in those with GBM⁷⁶. One study reported increased risk of death and poor overall survival in the NF1-mutated melanoma⁷⁷. Another study demonstrated that NF1 is an independent prognostic factor of survival in gastric carcinoma in univariate and multivariate analysis, with no or low NF1 expression significantly associated with shorter OS ($P < 0.001$)⁷⁸.

FINDING SUMMARY

NF1 encodes neurofibromin, a GTPase-activating protein (GAP) that is a key negative regulator of the RAS signaling pathway⁷⁹. Neurofibromin acts as a tumor suppressor by repressing RAS signaling⁸⁰. Alterations such as seen here may disrupt NF1 function or expression⁸⁰⁻⁸⁹.

POTENTIAL GERMLINE IMPLICATIONS

One or more of the NF1 variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters) associated with neurofibromatosis type 1 (ClinVar, Sep 2022)⁹⁰. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. Germline mutations in NF1 cause the autosomal dominant disorder neurofibromatosis type 1, which is characterized in part by increased risk of developing various tumors, including sarcoma, glioma, breast carcinoma, and neuroendocrine and hematological neoplasms⁹¹⁻⁹³. Estimates for the prevalence of the disorder in the general population range from 1:2,500 to 1:3,000⁹⁴⁻⁹⁵, and in the appropriate clinical context, germline testing of NF1 is recommended.

GENE
GNAS

ALTERATION

R201H

TRANSCRIPT ID

NM_000516.4

CODING SEQUENCE EFFECT

602G>A

VARIANT CHROMOSOMAL POSITION

chr20:57484421

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no therapies targeted to GNAS mutation in cancer. However, there is limited data indicating that a patient with appendiceal adenocarcinoma and a GNAS mutation (R201H) benefited from trametinib for 4 months⁹⁶. Additionally, a patient

with GNAS-mutated Erdheim-Chester disease exhibited a PR following treatment with single-agent trametinib⁹⁷.

FREQUENCY & PROGNOSIS

The highest incidences of GNAS mutations have been reported in intraductal papillary mucinous neoplasms (40-66%)⁹⁸⁻⁹⁹ and appendiceal mucinous neoplasms (50-72%)¹⁰⁰⁻¹⁰¹ as well as in tumors affecting the peritoneum (22%), pituitary gland (20%), bone (14%), pancreas (11%), and small intestine (11%)(COSMIC, 2022)¹⁰². Amplification of GNAS has been reported in ovarian epithelial carcinomas (12-30%)¹⁰³⁻¹⁰⁵, colorectal adenocarcinoma (9%)¹⁹, stomach adenocarcinoma (7%)¹⁰⁶, lung adenocarcinoma (6.5%)¹⁰⁷, breast invasive carcinoma (6.5%)¹⁰⁸, pancreatic adenocarcinoma (6%)¹⁰⁹, and sarcomas (5.8%)¹¹⁰. GNAS mutations are rare in hematological malignancies generally (COSMIC, 2022)^{102,111-112}. Activating GNAS mutations have been identified in gastrointestinal polyps in 75% (3/4) of patients

with McCune-Albright syndrome¹¹³. Amplification of GNAS has been associated with shorter progression-free survival in patients with ovarian cancer¹⁰⁴⁻¹⁰⁵, while activating GNAS mutations have been correlated with tumor progression and poor prognosis in patients with gastric cancer¹¹⁴.

FINDING SUMMARY

GNAS encodes the alpha subunit of the stimulatory G protein (Gs-alpha)¹¹⁵. Gs-alpha is a guanine-nucleotide binding protein (G protein) that is involved in hormonal regulation of adenylate cyclase¹¹⁵. GNAS has been reported to be amplified in cancer¹¹⁶ and may be biologically relevant in this context¹¹⁷⁻¹¹⁸. GNAS alterations that have been shown to result in constitutive activation of adenylyl cyclase and an increase in cellular cAMP concentration¹¹⁹⁻¹²⁴ are predicted to be activating. Mutations at R201 specifically are commonly associated with McCune-Albright syndrome, a disease that can co-occur with various cancers in patients with GNAS activating mutations¹²⁵⁻¹²⁷.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Nimesh Patel, M.D. | 12 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1517889-01

GENOMIC FINDINGS

GENE

TP53

ALTERATION

G245S

TRANSCRIPT ID

NM_000546.4

CODING SEQUENCE EFFECT

733G>A

VARIANT CHROMOSOMAL POSITION

chr17:7577548

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved therapies to address TP53 mutation or loss. However, tumors with TP53 loss of function alterations may be sensitive to the WEE1 inhibitor adavosertib¹²⁸⁻¹³¹ or p53 gene therapy such as SGT-53¹³²⁻¹³⁶. In a Phase 1 study, adavosertib in combination with gemcitabine, cisplatin, or carboplatin elicited PRs in 9.7% and SDs in 53% of patients with solid tumors; the response rate was 21% (4/19) for patients with TP53 mutations versus 12% (4/33) for patients who were TP53 wildtype¹³⁷. A Phase 2 trial of adavosertib in combination with chemotherapy (gemcitabine, carboplatin, paclitaxel, or doxorubicin) reported a 32% (30/94, 3 CR) ORR and a 73% (69/94) DCR for patients with platinum-refractory TP53-mutated ovarian, Fallopian tube, or peritoneal cancer¹³⁸. A smaller Phase 2 trial of adavosertib in combination with carboplatin achieved a 43% (9/21, 1 CR) ORR and a 76% (16/21) DCR for patients with TP53-mutated ovarian cancer¹³⁹. The combination of adavosertib with paclitaxel and carboplatin for patients with TP53-mutated ovarian cancer also significantly increased PFS compared with paclitaxel and carboplatin alone¹⁴⁰. In the Phase 2 VIKTORY trial, patients with TP53-mutated metastatic and/or recurrent gastric cancer experienced a 24% (6/25) ORR with adavosertib combined with paclitaxel¹⁴¹. A Phase 1 trial of neoadjuvant adavosertib in combination with cisplatin and docetaxel for head and neck squamous cell carcinoma (HNSCC) elicited a 71% (5/7) response rate for patients with TP53 alterations¹⁴². The Phase 2 FOCUS4-C trial for patients with TP53- and RAS-mutated colorectal

cancer reported improvement in PFS (3.61 vs. 1.87 months, HR=0.35, p=0.0022), but not OS (14.0 vs 12.8 months, p=0.93), following adavosertib treatment compared with active monitoring¹⁴³. In a Phase 1b clinical trial of SGT-53 in combination with docetaxel for patients with solid tumors, 75% (9/12) of evaluable patients experienced clinical benefit, including 2 confirmed and 1 unconfirmed PRs and 2 instances of SD with significant tumor shrinkage¹³⁶. Missense mutations leading to TP53 inactivation may be sensitive to therapies that reactivate mutated p53 such as eprenetapopt. In a Phase 1b trial for patients with p53-positive high-grade serous ovarian cancer, eprenetapopt combined with carboplatin and pegylated liposomal doxorubicin achieved a 52% (11/21) response rate and 100% DCR¹⁴⁴. A Phase 1 trial of eprenetapopt with pembrolizumab for patients with solid tumors reported an ORR of 10% (3/29)¹⁴⁵.

FREQUENCY & PROGNOSIS

Pan-cancer analysis of the TCGA datasets across 12 cancer types identified TP53 as the most frequently mutated gene, with 42% of more than 3,000 tumors harboring a TP53 mutation; in this study TP53 mutation occurred most frequently in ovarian serous carcinoma (95%), lung squamous cell carcinoma (SCC) (79%), head and neck SCC (70%), colorectal adenocarcinoma (59%), lung adenocarcinoma (52%), and bladder urothelial carcinoma (50%)¹⁴⁶. TP53 loss of heterozygosity (LOH) is frequently seen in tumors and often occurs when one copy of TP53 harbors a mutation; in some tumors, LOH is correlated with progression¹⁴⁷⁻¹⁵⁰. While the prognostic significance of TP53 alteration or dysregulation varies according to tumor type, studies have shown an association with poor prognosis for patients with breast cancer¹⁵¹⁻¹⁵³, endometrial cancer¹⁵⁴⁻¹⁵⁵, HNSCC¹⁵⁶⁻¹⁵⁸, or urothelial cancer¹⁵⁹⁻¹⁶⁰. In one study of 55 patients with lung adenocarcinoma, TP53 alterations correlated with immunogenic features including PD-L1 expression, tumor mutation burden and neoantigen presentation; likely as a consequence of this association TP53 mutations correlated with improved clinical outcomes to PD-1 inhibitors pembrolizumab and nivolumab in this study¹⁶¹. TP53 mutation has not been consistently demonstrated to be a significant independent prognostic marker in the context of

CRC¹⁶².

FINDING SUMMARY

Functional loss of the tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers¹⁶³. Alterations such as seen here may disrupt TP53 function or expression¹⁶⁴⁻¹⁶⁸.

POTENTIAL GERMLINE IMPLICATIONS

One or more of the TP53 variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters) associated with Li-Fraumeni syndrome (ClinVar, Sep 2022)⁹⁰. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. Germline mutations in TP53 are associated with the very rare autosomal dominant disorder Li-Fraumeni syndrome and the early onset of many cancers¹⁶⁹⁻¹⁷¹, including sarcomas¹⁷²⁻¹⁷³. Estimates for the prevalence of germline TP53 mutations in the general population range from 1:5,000¹⁷⁴ to 1:20,000¹⁷³. For pathogenic TP53 mutations identified during tumor sequencing, the rate of germline mutations was 1% in the overall population and 6% in tumors arising before age 30¹⁷⁵. In the appropriate clinical context, germline testing of TP53 is recommended.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion¹⁷⁶⁻¹⁸¹. CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy¹⁷⁶⁻¹⁷⁷. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease¹⁸². Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH^{180,183-184}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Nimesh Patel, M.D. | 12 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1517889-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Cobimetinib

Assay findings association

NF1
I679fs*21

AREAS OF THERAPEUTIC USE

Cobimetinib is a MEK inhibitor that is FDA approved to treat patients with histiocytic neoplasms. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical evidence in neurofibromatosis type 1 (NF1)-associated neurofibroma^{60-63,185-189}, glioma^{63-67,190}, and non-small cell lung cancer⁶⁸, NF1 inactivation may predict sensitivity to MEK inhibitors.

SUPPORTING DATA

Single-agent cobimetinib has shown clinical activity in the

context of histiocytic neoplasms, including Langerhans cell histiocytosis, Erdheim-Chester disease, Rosai-Dorfman disease, nodular histiocytosis, and mixed histiocytosis¹⁹¹⁻¹⁹⁸. A Phase 1 study of cobimetinib monotherapy in solid tumors reported CRs for 1.0% (1/97) of patients and PRs for 6.2% (6/97) of patients, all of whom had melanoma¹⁹⁹. Clinical benefit following treatment with cobimetinib has been seen for patients with non-small cell lung cancer (NSCLC)¹⁹⁹⁻²⁰⁰, adenoid cystic carcinoma¹⁹⁹, and anaplastic pleomorphic xanthoastrocytoma²⁰¹.

Selumetinib

Assay findings association

NF1
I679fs*21

AREAS OF THERAPEUTIC USE

Selumetinib is a MEK inhibitor that is FDA approved to treat pediatric patients with neurofibromatosis type 1 (NF1)-associated plexiform neurofibromas (PNs). Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical evidence in neurofibromatosis type 1 (NF1)-associated neurofibroma^{60-63,185-189}, glioma^{63-67,190}, and non-small cell lung cancer⁶⁸, NF1 inactivation may predict sensitivity to MEK inhibitors.

SUPPORTING DATA

Selumetinib has demonstrated efficacy in NF1-associated neurofibroma in Phase 2 studies^{61,185-186} and a Phase 1 study⁶⁰. Phase 2 studies reported clinical responses in low-grade glioma^{64,202}, melanoma²⁰³⁻²⁰⁷, and in

lung^{68,208-209} and endometrial cancer²¹⁰. A Phase 2 study of selumetinib for patients with activating alterations in the MAPK pathway reported a DCR of 15% (3/20), with no objective responses observed²¹¹. Phase 1 studies of selumetinib to treat patients with solid tumors reported 1/15 PR for a patient with colorectal cancer (CRC) and 5/15 SDs for patients with tonsil squamous cell carcinoma (SCC), non-small cell lung cancer (NSCLC), and CRC²¹²; 2/39 PRs (for patients with CRC) and 18/39 SDs were achieved when selumetinib was administered in combination with cyclosporin A²¹³. Multiple Phase 1 studies combining selumetinib with erlotinib or temsirolimus²¹⁴, docetaxel or dacarbazine²¹⁵, AKT inhibitors²¹⁶, or cixutumumab (an anti-IGF-1R antibody)²¹⁷ reported clinical responses for patients with advanced solid tumors including NSCLC, thyroid carcinoma, tongue SCC, and ovarian cancer.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Nimesh Patel, M.D. | 12 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1517889-01

THERAPIES WITH CLINICAL BENEFIT
IN OTHER TUMOR TYPE

Trametinib

Assay findings association
NF1
I679fs*21

AREAS OF THERAPEUTIC USE

Trametinib is a MEK inhibitor that is FDA approved as a monotherapy to treat patients with melanoma with BRAF V600E or V600K mutations. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical evidence in neurofibromatosis type 1 (NF1)-associated neurofibroma^{60-63,185-189}, glioma^{63-67,190}, and non-small cell lung cancer⁶⁸, NF1 inactivation may predict sensitivity to MEK inhibitors.

SUPPORTING DATA

A Phase 1 trial of trametinib in 206 patients with solid tumors reported 21 (10%) objective responses²¹⁸. Phase 1 monotherapy trials of RO4987655, another MEK inhibitor, have shown significant response rates in patients with melanoma, including those with BRAF and NRAS mutations, but very low response rates in patients with other solid tumors, including those with KRAS mutations²¹⁹⁻²²⁰. A Phase 1b trial of trametinib in

combination with gemcitabine in patients with solid tumors showed a complete response in a patient with breast cancer, as well as partial responses in pancreatic and salivary gland cancer²²¹. A Phase 1b trial of combination treatment with the MEK inhibitor binimetinib and the PI3K-alpha inhibitor alpelisib reported disease control (partial responses or stable disease) in 47% (21/45) of patients, including partial responses in 2 of 3 patients with KRAS-mutant ovarian cancer and 1 of 3 patients with NRAS-mutant melanoma; a 43% rate of stable disease was observed in patients with KRAS-mutant colorectal cancer, with responses independent of PIK3CA mutation status²²². Whereas frequent adverse events precluded a recommended Phase 2 dose and schedule for the combination of trametinib and everolimus in a Phase 1b trial for solid tumors²²³, a retrospective study for heavily pretreated patients with solid tumors reported tolerable regimens of the combination for 23/31 patients, with 16 patients treated >3 months and evaluable patients achieving a median PFS of 6.5 months²²⁴.

NOTE Genomic alterations detected may be associated with activity of certain US FDA or other specific country approved therapies; however, the therapies listed in this report may have varied evidence in the patient's tumor type. The listed therapies are not ranked in order of potential or predicted efficacy for this patient or in order of level of evidence for this patient's tumor type. The therapies listed in this report may not be complete and/or exhaustive. Furthermore, the listed therapies are limited to US FDA approved pharmaceutical drug products that are linked to a specific genomic alteration. There may also be US FDA approved pharmaceutical drug products that are not linked to a genomic alteration. Further there may also exist pharmaceutical drug products that are not approved by the US FDA or other national authorities. There may also be other treatment modalities available than pharmaceutical drug products.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Nimesh Patel, M.D. | 12 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1517889-01

CLINICAL TRIALS

IMPORTANT Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually updated and should be investigated by the physician or

research staff. This is not a comprehensive list of all available clinical trials. There may also be compassionate use or early access programs available, which are not listed in this report. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials are not ranked in order of potential or predicted efficacy for this patient or

in order of level of evidence for this patient's tumor type. Clinical trials listed here may have additional enrollment criteria that may require medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see clinicaltrials.gov. However, clinicaltrials.gov does not list all clinical trials that might be available.

GENE
NF1
ALTERATION
I679fs*21
RATIONALE

On the basis of clinical evidence and strong preclinical evidence, NF1 inactivation may predict sensitivity to MEK inhibitors. Limited clinical

data and strong preclinical data indicate that loss or inactivation of NF1 may also predict sensitivity to mTOR inhibitors.

NCT03239015
PHASE 2

Efficacy and Safety of Targeted Precision Therapy in Refractory Tumor With Druggable Molecular Event

TARGETS

EGFR, ERBB4, ERBB2, PARP, mTOR, MET, ROS1, RET, VEGFRs, BRAF, CDK4, CDK6

LOCATIONS: Shanghai (China)

NCT04337463
PHASE NULL

ATG-008 Combined With Toripalimab in Advanced Solid Tumors

TARGETS

mTORC1, mTORC2, PD-1

LOCATIONS: Chongqing (China), Chengdu (China)

NCT04985604
PHASE 1/2

DAY101 Monotherapy or in Combination With Other Therapies for Patients With Solid Tumors

TARGETS

BRAF, MEK

LOCATIONS: Busan (Korea, Republic of), Seoul (Korea, Republic of), Oregon, Barcelona (Spain), Madrid (Spain), California, Colorado, Toronto (Canada), Indiana

NCT04803318
PHASE 2

Trametinib Combined With Everolimus and Lenvatinib for Recurrent/Refractory Advanced Solid Tumors

TARGETS

mTOR, FGFRs, RET, PDGFRA, VEGFRs, KIT, MEK

LOCATIONS: Guangzhou (China)

NCT05125523
PHASE 1

A Study of Sirolimus for Injection (Albumin Bound) in Patients With Advanced Solid Tumors

TARGETS

mTOR

LOCATIONS: Tianjin (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Nimesh Patel, M.D. | 12 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1517889-01

CLINICAL TRIALS
NCT04801966
PHASE NULL

Safety and Oversight of the Individually Tailored Treatment Approach: A Novel Pilot Study

TARGETS

CDK4, CDK6, PI3K-alpha, PD-L1, MEK,
PARP, PD-1, BRAF

LOCATIONS: Melbourne (Australia)

NCT03498521
PHASE 2

A Phase II Randomized Study Comparing the Efficacy and Safety of Targeted Therapy or Cancer Immunotherapy Versus Platinum-Based Chemotherapy in Patients With Cancer of Unknown Primary Site

TARGETS

ALK, RET, SMO, AKTs, PARP, PD-L1,
EGFR, VEGFA, MEK, BRAF, ERBB2,
TRKB, TRKC, ROS1, TRKA

LOCATIONS: Kecskemet (Hungary), Monterrey (Mexico), Cdmx (Mexico), Temuco (Chile), Recoleta (Chile)

NCT03297606
PHASE 2

Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR)

TARGETS

VEGFRs, ABL, SRC, ALK, ROS1, AXL,
TRKA, MET, TRKC, DDR2, KIT, EGFR,
PD-1, CTLA-4, PARP, CDK4, CDK6,
FLT3, CSF1R, RET, mTOR, ERBB2, MEK,
BRAF, SMO

LOCATIONS: Vancouver (Canada), Edmonton (Canada), Saskatoon (Canada), Regina (Canada), Ottawa (Canada), Montreal (Canada), Toronto (Canada), Kingston (Canada), London (Canada)

NCT04720976
PHASE 1/2

JAB-3312 Activity in Adult Patients With Advanced Solid Tumors

TARGETS

MEK, SHP2, PD-1, EGFR, KRAS

LOCATIONS: Utah, California, Arizona, Minnesota, Illinois, Michigan, Oklahoma, Missouri, Indiana, Connecticut

NCT04965818
PHASE 1/2

Phase 1b/2 Study of Futibatinib in Combination With Binimetinib in Patients With Advanced KRAS Mutant Cancer

TARGETS

MEK, FGFRs

LOCATIONS: California, Indiana, Texas

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Nimesh Patel, M.D. | 12 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1517889-01

APPENDIX
Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

APC
V1125A

CARD11
S694L

CD22
S230P

CREBBP
S908I

EGFR
D800V

EPHA3
H61Q

FGF6
L3M

FGFR1
P469S

IRS2
A512T

MAP3K1
L78P

NTRK3
G547R

PALB2
S652N

PDCD1LG2 (PD-L2)
Y106C

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Nimesh Patel, M.D. | 12 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1517889-01

APPENDIX

Genes assayed in FoundationOne®Liquid CDx

FoundationOne Liquid CDx interrogates 324 genes, including 309 genes with complete exonic (coding) coverage and 15 genes with only select non-coding coverage (indicated with an *); 75 genes (indicated in bold) are captured with increased sensitivity and have complete exonic (coding) coverage unless otherwise noted.

ABL1 Exons 4-9	<i>ACVR1B</i>	AKT1 Exon 3	<i>AKT2</i>	<i>AKT3</i>	ALK Exons 20-29, Introns 18, 19	<i>ALOX12B</i>	<i>AMER1</i> (FAM123B or WTX)	APC
AR	ARAF Exons 4, 5, 7, 11, 13, 15, 16	<i>ARFRP1</i>	<i>ARID1A</i>	<i>ASXL1</i>	ATM	ATR	<i>ATRX</i>	<i>AURKA</i>
<i>AURKB</i>	<i>AXIN1</i>	<i>AXL</i>	<i>BAP1</i>	<i>BARD1</i>	<i>BCL2</i>	<i>BCL2L1</i>	<i>BCL2L2</i>	<i>BCL6</i>
<i>BCOR</i>	<i>BCORL1</i>	<i>BCR*</i> Introns 8, 13, 14	BRAF Exons 11-18, Introns 7-10	BRCA1 Introns 2, 7, 8, 12, 16, 19, 20	BRCA2 Intron 2	<i>BRD4</i>	<i>BRIP1</i>	<i>BTG1</i>
<i>BTG2</i>	BTB Exons 2, 15	<i>CALR</i>	<i>CARD11</i>	<i>CASP8</i>	<i>CBFB</i>	<i>CBL</i>	CCND1	<i>CCND2</i>
<i>CCND3</i>	<i>CCNE1</i>	<i>CD22</i>	<i>CD70</i>	<i>CD74*</i> Introns 6-8	<i>CD79A</i>	<i>CD79B</i>	CD274 (PD-L1)	<i>CDC73</i>
CDH1	CDK12	CDK4	CDK6	<i>CDK8</i>	<i>CDKN1A</i>	<i>CDKN1B</i>	CDKN2A	<i>CDKN2B</i>
<i>CDKN2C</i>	<i>CEBPA</i>	<i>CHEK1</i>	CHEK2	<i>CIC</i>	<i>CREBBP</i>	CRKL	<i>CSF1R</i>	<i>CSF3R</i>
<i>CTCF</i>	<i>CTNNA1</i>	CTNNB1 Exon 3	<i>CUL3</i>	<i>CUL4A</i>	<i>CXCR4</i>	<i>CYP17A1</i>	<i>DAXX</i>	<i>DDR1</i>
DDR2 Exons 5, 17, 18	<i>DIS3</i>	<i>DNMT3A</i>	<i>DOT1L</i>	<i>EED</i>	EGFR Introns 7, 15, 24-27	<i>EMSY</i> (C11orf30)	<i>EP300</i>	<i>EPHA3</i>
<i>EPHB1</i>	<i>EPHB4</i>	ERBB2	ERBB3 Exons 3, 6, 7, 8, 10, 12, 20, 21, 23, 24, 25	<i>ERBB4</i>	<i>ERCC4</i>	<i>ERG</i>	ERRF1	ESR1 Exons 4-8
<i>ETV4*</i> Intron 8	<i>ETV5*</i> Introns 6, 7	ETV6* Introns 5, 6	<i>EWSR1*</i> Introns 7-13	EZH2 Exons 4, 16, 17, 18	<i>EZR*</i> Introns 9-11	<i>FANCA</i>	<i>FANCC</i>	<i>FANCG</i>
<i>FANCL</i>	<i>FAS</i>	<i>FBXW7</i>	<i>FGF10</i>	<i>FGF12</i>	<i>FGF14</i>	<i>FGF19</i>	<i>FGF23</i>	<i>FGF3</i>
<i>FGF4</i>	<i>FGF6</i>	FGFR1 Introns 1, 5, Intron 17	FGFR2 Intron 1, Intron 17	FGFR3 Exons 7, 9 (alternative designation exon 10), 14, 18, Intron 17	<i>FGFR4</i>	<i>FH</i>	<i>FLCN</i>	<i>FLT1</i>
FLT3 Exons 14, 15, 20	FOXL2	<i>FUBP1</i>	<i>GABRA6</i>	<i>GATA3</i>	<i>GATA4</i>	<i>GATA6</i>	<i>GID4</i> (C17orf39)	GNA11 Exons 4, 5
<i>GNA13</i>	GNAQ Exons 4, 5	GNAS Exons 1, 8	<i>GRM3</i>	<i>GSK3B</i>	<i>H3-3A</i> (H3F3A)	<i>HDAC1</i>	<i>HGF</i>	<i>HNFI1A</i>
HRAS Exons 2, 3	<i>HSD3B1</i>	<i>ID3</i>	IDH1 Exon 4	IDH2 Exon 4	<i>IGF1R</i>	<i>IKBKE</i>	<i>IKZF1</i>	<i>INPP4B</i>
<i>IRF2</i>	<i>IRF4</i>	<i>IRS2</i>	<i>JAK1</i>	JAK2 Exon 14	JAK3 Exons 5, 11, 12, 13, 15, 16	<i>JUN</i>	<i>KDM5A</i>	<i>KDM5C</i>
<i>KDM6A</i>	<i>KDR</i>	<i>KEAP1</i>	<i>KEL</i>	KIT Exons 8, 9, 11, 12, 13, 17, Intron 16	<i>KLHL6</i>	<i>KMT2A</i> (MLL) Introns 6, 8-11, Intron 7	<i>KMT2D</i> (MLL2)	KRAS

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1517889-01

APPENDIX

Genes assayed in FoundationOne®Liquid CDx

FoundationOne Liquid CDx interrogates 324 genes, including 309 genes with complete exonic (coding) coverage and 15 genes with only select non-coding coverage (indicated with an *); 75 genes (indicated in bold) are captured with increased sensitivity and have complete exonic (coding) coverage unless otherwise noted.

LTK	LYN	MAF	MAP2K1 (MEK1) Exons 2, 3	MAP2K2 (MEK2) Exons 2-4, 6, 7	MAP2K4	MAP3K1	MAP3K13	MAPK1
MCL1	MDM2	MDM4	MED12	MEF2B	MEN1	MERTK	MET	MITF
MKNK1	MLH1	MPL Exon 10	MRE11 (MRE11A)	MSH2 Intron 5	MSH3	MSH6	MST1R	MTAP
MTOR Exons 19, 30, 39, 40, 43-45, 47, 48, 53, 56	MUTYH	MYB* Intron 14	MYC Intron 1	MYCL (MYCL1)	MYCN	MYD88 Exon 4	NBN	NF1
NF2	NFE2L2	NFKBIA	NKX2-1	NOTCH1	NOTCH2 Intron 26	NOTCH3	NPM1 Exons 4-6, 8, 10	NRAS Exons 2, 3
NSD2 (WHSC1 or MMSET)	NSD3 (WHSC1L1)	NT5C2	NTRK1 Exons 14, 15, Introns 8-11	NTRK2 Intron 12	NTRK3 Exons 16, 17	NUTM1* Intron 1	P2RY8	PALB2
PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)	PDGFRA Exons 12, 18, Introns 7, 9, 11	PDGFRB Exons 12-21, 23 9, 11
PDK1	PIK3C2B	PIK3C2G	PIK3CA Exons 2, 3, 5-8, 10, 14, 19, 21 (Coding Exons 1, 2, 4-7, 9, 13, 18, 20)	PIK3CB	PIK3R1	PIM1	PMS2	POLD1
POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI	PRKN (PARK2)	PTCH1
PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51	RAD51B	RAD51C
RAD51D	RAD52	RAD54L	RAF1 Exons 3, 4, 6, 7, 10, 14, 15, 17, Introns 4-8	RARA Intron 2	RB1	RBM10	REL	RET Introns 7, 8, Exons 11, 13-16, Introns 9-11
RICTOR	RNF43	ROS1 Exons 31, 36-38, 40, Introns 31-35	RPTOR	RSP02* Intron 1	SDC4* Intron 2	SDHA	SDHB	SDHC
SDHD	SETD2	SF3B1	SGK1	SLC34A2* Intron 4	SMAD2	SMAD4	SMARCA4	SMARCB1
SMO	SNCAIP	SOCS1	SOX2	SOX9	SPEN	SPOP	SRC	STAG2
STAT3	STK11	SUFU	SYK	TBX3	TEK	TENT5C (FAM46C)	TERC* ncRNA	TERT* Promoter
TET2	TGFBR2	TIPARP	TMPRSS2* Introns 1-3	TNFAIP3	TNFRSF14	TP53	TSC1	TSC2
TYRO3	U2AF1	VEGFA	VHL	WT1	XPO1	XRCC2	ZNF217	ZNF703

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS

Microsatellite (MS) status
Blood Tumor Mutational Burden (bTMB)
Tumor Fraction

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Nimesh Patel, M.D. | 12 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1517889-01

APPENDIX

About FoundationOne® Liquid CDx

FoundationOne Liquid CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Ciplastraat 3, 2440 Geel, Belgium. The CE-IVD regulatory status of FoundationOne Liquid CDx is applicable in countries that accept and/or recognize the CE mark.



ABOUT FOUNDATIONONE LIQUID CDx

FoundationOne Liquid CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne Liquid CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details.

INTENDED USE

FoundationOne Liquid CDx is a next generation sequencing based *in vitro* diagnostic device that analyzes 324 genes. Substitutions and insertion and deletion alterations (indels) are reported in 311 genes, copy number alterations (CNAs) are reported in 310 genes, and gene rearrangements are reported in 324 genes. The test also detects the genomic signatures blood tumor mutational burden (bTMB), microsatellite instability (MSI), and tumor fraction. FoundationOne Liquid CDx utilizes circulating cell-free DNA (cfDNA) isolated from plasma derived from the anti-coagulated peripheral whole blood of cancer patients. The test is intended to be used as a companion diagnostic to identify patients who may benefit from treatment with targeted therapies in accordance with the approved therapeutic product labeling. Additionally, FoundationOne Liquid CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with malignant neoplasms.

TEST PRINCIPLES

The FoundationOne Liquid CDx assay is performed exclusively as a laboratory service using circulating cell-free DNA (cfDNA) isolated from plasma derived from anti-coagulated peripheral whole blood from patients with solid malignant neoplasms. The assay employs a single DNA extraction method to obtain cfDNA from plasma from whole blood. Extracted

cfDNA undergoes whole-genome shotgun library construction and hybridization-based capture of 324 cancer-related genes including coding exons and select introns of 309 genes, as well as only select intronic regions or non-coding regions of 15 genes. Hybrid-capture selected libraries are sequenced with deep coverage using the NovaSeq® 6000 platform. Sequence data are processed using a customized analysis pipeline designed to accurately detect genomic alterations, including base substitutions, indels, select copy number variants, and select genomic rearrangements. Substitutions and insertion and deletion alterations (indels) are reported in 311 genes, copy number alterations (CNAs) are reported in 310 genes, and gene rearrangements are reported in 324 genes. The assay also reports tumor fraction, and genomic signatures including MSI and bTMB. A subset of targeted regions in 75 genes is baited for increased sensitivity.

THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. *Note:* A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

QUALIFIED ALTERATION CALLS (EQUIVOCAL)

All equivocal calls, regardless of alteration type, imply that there is adequate evidence to call the alteration with confidence. However, the repeatability of equivocal calls may be lower than non-equivocal calls.

RANKING OF THERAPIES AND CLINICAL TRIALS

Ranking of Therapies in Summary Table

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

Ranking of Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

LIMITATIONS

1. For *in vitro* diagnostic use.
2. For prescription use only. This test must be ordered by a qualified medical professional in accordance with clinical laboratory regulations.
3. A negative result does not rule out the presence of a mutation below the limits of detection of the assay. Patients for whom no companion diagnostic alterations are detected should be considered for confirmation with an appropriately validated tumor tissue test, if available.
4. The FoundationOne Liquid CDx assay does not detect heterozygous deletions.
5. The test is not intended to provide information on cancer predisposition.
6. Performance has not been validated for cfDNA input below the specified minimum input.
7. Tissue TMB and blood TMB (bTMB) are estimated from the number of synonymous and nonsynonymous single-nucleotide variants (SNVs) and insertions and deletions (indels) per area of coding genome sampled, after the removal of known and likely oncogenic driver events and germline SNPs. Tissue TMB is calculated based on variants with an allele frequency of $\geq 5\%$, and bTMB is calculated based on variants with an allele frequency of $\geq 0.5\%$.
8. Tumor fraction is the percentage of circulating tumor DNA (ctDNA) present in a cell-free DNA (cfDNA) sample. The tumor fraction estimate is computationally derived from the observed level of aneuploidy in the sample. Tumor fraction is considered elevated when ctDNA levels are high enough that aneuploidy can be detected and is significantly distinct from that typically found in non-tumor samples.
9. Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the tumor genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor. The MSI algorithm is based on genome wide analysis of 1765 microsatellite loci and not based on the 5 or 7 MSI loci described in current clinical practice guidelines for solid tissue testing.
10. Genomic findings from circulating cell-free DNA (cfDNA) may originate from circulating tumor DNA fragments, germline alterations, or non-tumor somatic alterations, such as clonal hematopoiesis of indeterminate potential (CHIP). Genes with alterations that may be derived from CHIP include, but are not limited to: *ASXL1*, *ATM*, *CBL*, *CHEK2*, *DNMT3A*, *JAK2*,

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1517889-01

APPENDIX

About FoundationOne® Liquid CDx

KMT2D (MLL2), MPL, MYD88, SF3B1, TET2, TP53, and U2AF1.

11. Alterations reported may include somatic (not inherited) or germline (inherited) alterations; however, the test does not distinguish between germline and somatic alterations. If a reported alteration is suspected to be germline, confirmatory testing should be considered in the appropriate clinical context.
12. The test is not intended to replace germline testing or to provide information about cancer predisposition.

REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic, nontargeted treatment, germline, or clonal hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >30%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM, BAP1, BRCA1, BRCA2, BRIP1, CHEK2, FH, FLCN, MLH1, MSH2, MSH6, MUTYH, PALB2, PMS2, POLE, RAD51C, RAD51D, RET, SDHA, SDHB, SDHC, SDHD, TSC2, and VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's

tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1, ATM, CBL, CHEK2, DNMT3A, IDH2, JAK2, KMT2D (MLL2), MPL, MYD88, SF3B1, TET2, and U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) (www.nccn.org). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2022. All rights reserved. To view the most recent and complete version of the guidelines, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

This report makes no promises or guarantees that a particular drug will be effective in the treatment of

disease in any patient. This report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne Liquid CDx.

TREATMENT DECISIONS ARE THE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this test or the information contained in this report.

Certain sample of variant characteristics may result in reduced sensitivity. These include: low sample quality, deletions and insertions >40bp, or repetitive/high homology sequences. FoundationOne Liquid CDx is performed using cell-free DNA, and as such germline events may not be reported.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Nimesh Patel, M.D. | 12 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1517889-01

APPENDIX
About FoundationOne®Liquid CDx
SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
Muts/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

REFERENCE SEQUENCE INFORMATION

Sequence data is mapped to the human genome, Genome Reference Consortium Human Build 37 (GRCh37), also known as hg19.

MR Suite Version (RG) 7.4.0

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Nimesh Patel, M.D. | 12 December 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1517889-01

APPENDIX **References**

1. Gandara DR, et al. *Nat. Med.* (2018) PMID: 30082870
2. Wang Z, et al. *JAMA Oncol* (2019) PMID: 30816954
3. Sturgill EG, et al. *Oncologist* (2022) PMID: 35274716
4. Aggarwal C, et al. *Clin. Cancer Res.* (2020) PMID: 32102950
5. Schenker et al., 2022; AACR Abstract CT022
6. Saori et al., 2021; ESMO Abstract 80P
7. Chen EX, et al. *JAMA Oncol* (2020) PMID: 32379280
8. Rizvi NA, et al. *JAMA Oncol* (2020) PMID: 32271377
9. Si H, et al. *Clin Cancer Res* (2021) PMID: 33355200
10. Leigh NB, et al. *J Thorac Oncol* (2022) PMID: 34800700
11. Li et al., 2020; ASCO Abstract 6511
12. Xiao D, et al. *Oncotarget* (2016) PMID: 27009843
13. Spigel et al., 2016; ASCO Abstract 9017
14. Shim HS, et al. *J Thorac Oncol* (2015) PMID: 26200269
15. Govindan R, et al. *Cell* (2012) PMID: 22980976
16. Ding L, et al. *Nature* (2008) PMID: 18948947
17. Imielinski M, et al. *Cell* (2012) PMID: 22980975
18. Kim Y, et al. *J. Clin. Oncol.* (2014) PMID: 24323028
19. *Nature* (2012) PMID: 22810696
20. Stadler ZK, et al. *J. Clin. Oncol.* (2016) PMID: 27022117
21. Samowitz WS, et al. *Cancer Epidemiol. Biomarkers Prev.* (2001) PMID: 11535541
22. Elsaleh H, et al. *Clin Colorectal Cancer* (2001) PMID: 12445368
23. Brueckl WM, et al. *Anticancer Res.* () PMID: 12820457
24. Guidoboni M, et al. *Am. J. Pathol.* (2001) PMID: 11438476
25. Gryfe R, et al. *N. Engl. J. Med.* (2000) PMID: 10631274
26. Sinicrope FA, et al. *Gastroenterology* (2006) PMID: 16952542
27. Guastadisegni C, et al. *Eur. J. Cancer* (2010) PMID: 20627535
28. Laghi L, et al. *Dig Dis* (2012) PMID: 22722556
29. Mehnert JM, et al. *J. Clin. Invest.* (2016) PMID: 27159395
30. Cancer Genome Atlas Research Network, et al. *Nature* (2013) PMID: 23636398
31. Hussein YR, et al. *Mod. Pathol.* (2015) PMID: 25394778
32. Church DN, et al. *Hum. Mol. Genet.* (2013) PMID: 23528559
33. Cazier JB, et al. *Nat Commun* (2014) PMID: 24777035
34. Pfeifer GP, et al. *Mutat. Res.* (2005) PMID: 15748635
35. Hill VK, et al. *Annu Rev Genomics Hum Genet* (2013) PMID: 23875803
36. Pfeifer GP, et al. *Oncogene* (2002) PMID: 12379884
37. Rizvi NA, et al. *Science* (2015) PMID: 25765070
38. Johnson BE, et al. *Science* (2014) PMID: 24336570
39. Choi S, et al. *Neuro-oncology* (2018) PMID: 29452419
40. Briggs S, et al. *J. Pathol.* (2013) PMID: 23447401
41. Heitzner E, et al. *Curr. Opin. Genet. Dev.* (2014) PMID: 24583393
42. Roberts SA, et al. *Nat. Rev. Cancer* (2014) PMID: 25568919
43. Bronkhorst AJ, et al. *Biomol Detect Quantif* (2019) PMID: 30923679
44. Raja R, et al. *Clin. Cancer Res.* (2018) PMID: 30093454
45. Hrebien S, et al. *Ann. Oncol.* (2019) PMID: 30860573
46. Choudhury AD, et al. *JCI Insight* (2018) PMID: 30385733
47. Goodall J, et al. *Cancer Discov* (2017) PMID: 28450425
48. Goldberg SB, et al. *Clin. Cancer Res.* (2018) PMID: 29330207
49. Bettgowda C, et al. *Sci Transl Med* (2014) PMID: 24553385
50. Lapin M, et al. *J Transl Med* (2018) PMID: 30400802
51. Shulman DS, et al. *Br. J. Cancer* (2018) PMID: 30131550
52. Stover DG, et al. *J. Clin. Oncol.* (2018) PMID: 29298117
53. Hemming ML, et al. *JCO Precis Oncol* (2019) PMID: 30793095
54. Egyud M, et al. *Ann. Thorac. Surg.* (2019) PMID: 31059681
55. Fan G, et al. *PLoS ONE* (2017) PMID: 28187169
56. Vu et al., 2020; DOI: 10.1200/PO.19.00204
57. Li G, et al. *J Gastrointest Oncol* (2019) PMID: 31602320
58. Zhang EW, et al. *Cancer* (2020) PMID: 32757294
59. Butler TM, et al. *Cold Spring Harb Mol Case Stud* (2019) PMID: 30833418
60. Dombi E, et al. *N. Engl. J. Med.* (2016) PMID: 28029918
61. Schalkwijk S, et al. *Cancer Chemother Pharmacol* (2021) PMID: 33903938
62. Toledano H, et al. *Childs Nerv Syst* (2021) PMID: 32780261
63. Ronsley R, et al. *Cancer Med* (2021) PMID: 33939292
64. Fangusaro J, et al. *Lancet Oncol.* (2019) PMID: 31151904
65. Manoharan N, et al. *J Neurooncol* (2020) PMID: 32780261
66. Kondyli M, et al. *J Neurooncol* (2018) PMID: 30097824
67. Awada G, et al. *Case Rep Oncol* () PMID: 33082744
68. Middleton G, et al. *Nature* (2020) PMID: 32669708
69. Lim SM, et al. *Oncotarget* (2016) PMID: 26859683
70. Weiss B, et al. *Neuro-oncology* (2015) PMID: 25314964
71. Janku F, et al. *Oncotarget* (2014) PMID: 24931142
72. Johannessen CM, et al. *Curr. Biol.* (2008) PMID: 18164202
73. Johannessen CM, et al. *Proc. Natl. Acad. Sci. U.S.A.* (2005) PMID: 15937108
74. Malone CF, et al. *Cancer Discov* (2014) PMID: 24913553
75. Zehir A, et al. *Nat. Med.* (2017) PMID: 28481359
76. Vizcaino MA, et al. *Hum. Pathol.* (2015) PMID: 26190195
77. Cirenajwis H, et al. *Mol Oncol* (2017) PMID: 28267273
78. Liu D, et al. *Oncotarget* (2017) PMID: 29137312
79. Hattori S, et al. *Biochem. Biophys. Res. Commun.* (1991) PMID: 1904223
80. Morcos P, et al. *Mol. Cell. Biol.* (1996) PMID: 8628317
81. Ballester R, et al. *Cell* (1990) PMID: 2121371
82. Xu GF, et al. *Cell* (1990) PMID: 2116237
83. Martin GA, et al. *Cell* (1990) PMID: 2121370
84. Thomas L, et al. *Hum. Mutat.* (2012) PMID: 22807134
85. Skuse GR, et al. *Hum. Mol. Genet.* (1997) PMID: 9300663
86. Messiaen LM, et al. *Genet. Med.* () PMID: 11258625
87. Ars E, et al. *Hum. Mol. Genet.* (2000) PMID: 10607834
88. Messiaen LM, et al. *J. Med. Genet.* (2005) PMID: 15863657
89. Poulet P, et al. *Mol. Cell. Biol.* (1994) PMID: 8264648
90. Landrum MJ, et al. *Nucleic Acids Res.* (2018) PMID: 29165669
91. Jett K, et al. *Genet. Med.* (2010) PMID: 20027112
92. Patil S, et al. *Oncologist* (2012) PMID: 22240541
93. Evans DG, et al. *Clin Sarcoma Res* (2012) PMID: 23036231
94. Upadhyaya M, et al. *J. Med. Genet.* (1995) PMID: 8544190
95. Williams VC, et al. *Pediatrics* (2009) PMID: 19117870
96. Ang C, et al. *Case Rep Oncol* () PMID: 28868010
97. Saunders IM, et al. *Oncologist* (2019) PMID: 31740567
98. Furukawa T, et al. *Sci Rep* (2011) PMID: 22355676
99. Wu J, et al. *Sci Transl Med* (2011) PMID: 21775669
100. Nishikawa G, et al. *Br. J. Cancer* (2013) PMID: 23403822
101. Singhi AD, et al. *Hum. Pathol.* (2014) PMID: 24925222
102. Tate JG, et al. *Nucleic Acids Res.* (2019) PMID: 30371878
103. *Nature* (2011) PMID: 21720365
104. Kan Z, et al. *Nature* (2010) PMID: 20668451
105. Tominaga E, et al. *Gynecol. Oncol.* (2010) PMID: 20537689
106. *Nature* (2014) PMID: 25079317
107. *Nature* (2014) PMID: 25079552
108. *Nature* (2012) PMID: 23000897
109. Witkiewicz AK, et al. *Nat Commun* (2015) PMID: 25855536
110. Barretina J, et al. *Nat. Genet.* (2010) PMID: 20601955
111. Lohr JG, et al. *Cancer Cell* (2014) PMID: 24434212
112. Chapman MA, et al. *Nature* (2011) PMID: 21430775
113. Zacharin M, et al. *J. Med. Genet.* (2011) PMID: 21357941
114. Alakus H, et al. *World J. Gastroenterol.* (2009) PMID: 20027678
115. Hayward BE, et al. *Proc. Natl. Acad. Sci. U.S.A.* (1998) PMID: 9860993
116. Gao J, et al. *Sci Signal* (2013) PMID: 23550210
117. Zack TI, et al. *Nat. Genet.* (2013) PMID: 24071852
118. Beroukhi R, et al. *Nature* (2010) PMID: 20164920
119. Masters SB, et al. *J. Biol. Chem.* (1989) PMID: 2549064
120. Graziano MP, et al. *J. Biol. Chem.* (1989) PMID: 2549065
121. Jang IS, et al. *Exp. Mol. Med.* (2001) PMID: 11322485
122. Landis CA, et al. *Nature* (1989) PMID: 2549426
123. Tobar-Rubin R, et al. *J. Mol. Endocrinol.* (2013) PMID: 23288949
124. Mariot V, et al. *Bone* (2011) PMID: 20887824
125. Weinstein LS, et al. *N. Engl. J. Med.* (1991) PMID: 1944469
126. Collins MT, et al. *J. Clin. Endocrinol. Metab.* (2003) PMID: 12970318
127. Nault JC, et al. *J. Hepatol.* (2012) PMID: 21835143
128. Hirai H, et al. *Cancer Biol. Ther.* (2010) PMID: 20107315
129. Bridges KA, et al. *Clin. Cancer Res.* (2011) PMID: 21799033
130. Rajeshkumar NV, et al. *Clin. Cancer Res.* (2011) PMID: 21389100
131. Osman AA, et al. *Mol. Cancer Ther.* (2015) PMID: 25504633
132. Xu L, et al. *Mol. Cancer Ther.* (2002) PMID: 12489850
133. Xu L, et al. *Mol. Med.* (2001) PMID: 11713371
134. Camp ER, et al. *Cancer Gene Ther.* (2013) PMID: 23470564
135. Kim SS, et al. *Nanomedicine* (2015) PMID: 25240597
136. Pirollo KF, et al. *Mol. Ther.* (2016) PMID: 27357628
137. Leijen S, et al. *J. Clin. Oncol.* (2016) PMID: 27601554
138. Moore et al., 2019; ASCO Abstract 5513
139. Leijen S, et al. *J. Clin. Oncol.* (2016) PMID: 27998224
140. Oza et al., 2015; ASCO Abstract 5506
141. Lee J, et al. *Cancer Discov* (2019) PMID: 31315834
142. Méndez E, et al. *Clin. Cancer Res.* (2018) PMID: 29535125
143. Seligmann JF, et al. *J Clin Oncol* (2021) PMID: 34538072
144. Gourley et al., 2016; ASCO Abstract 5571
145. Park H, et al. *ESMO Open* (2022) PMID: 36084396
146. Kandath C, et al. *Nature* (2013) PMID: 24132290
147. Wongsurawat VJ, et al. *Cancer Epidemiol. Biomarkers Prev.* (2006) PMID: 16537709
148. Brosh R, et al. *Nat. Rev. Cancer* (2009) PMID: 19693097
149. Baker SJ, et al. *Science* (1989) PMID: 2649981
150. Calcagno DQ, et al. *BMC Gastroenterol* (2013) PMID: 24053468
151. Alsnér J, et al. *Acta Oncol* (2008) PMID: 18465328
152. Olivier M, et al. *Clin. Cancer Res.* (2006) PMID: 16489069
153. Végran F, et al. *PLoS ONE* (2013) PMID: 23359294
154. Wild PJ, et al. *EMBO Mol Med* (2012) PMID: 22678923
155. Lee EJ, et al. *Gynecol. Oncol.* (2010) PMID: 20006376

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # **ORD-1517889-01**
APPENDIX
References

156. Ganci F, et al. Ann. Oncol. (2013) pmid: 24107801
157. Lindenberg-van der Plas M, et al. Clin. Cancer Res. (2011) pmid: 21467160
158. Peltonen JK, et al. Head Neck Oncol (2011) pmid: 21513535
159. Bringuier PP, et al. Int. J. Cancer (1998) pmid: 9761125
160. Feng C, et al. Sci Rep (2014) pmid: 24500328
161. Dong ZY, et al. Clin. Cancer Res. (2017) pmid: 28039262
162. Russo A, et al. J. Clin. Oncol. (2005) pmid: 16172461
163. Brown CJ, et al. Nat. Rev. Cancer (2009) pmid: 19935675
164. Joerger AC, et al. Annu. Rev. Biochem. (2008) pmid: 18410249
165. Kato S, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) pmid: 12826609
166. Kamada R, et al. J. Biol. Chem. (2011) pmid: 20978130
167. Zerdoumi Y, et al. Hum. Mol. Genet. (2017) pmid: 28472496
168. Yamada H, et al. Carcinogenesis (2007) pmid: 17690113
169. Bougeard G, et al. J. Clin. Oncol. (2015) pmid: 26014290
170. Sorrell AD, et al. Mol Diagn Ther (2013) pmid: 23355100
171. Nichols KE, et al. Cancer Epidemiol. Biomarkers Prev. (2001) pmid: 11219776
172. Kleihues P, et al. Am. J. Pathol. (1997) pmid: 9006316
173. Gonzalez KD, et al. J. Clin. Oncol. (2009) pmid: 19204208
174. Laloo F, et al. Lancet (2003) pmid: 12672316
175. Mandelker D, et al. Ann. Oncol. (2019) pmid: 31050713
176. Jaiswal S, et al. N. Engl. J. Med. (2014) pmid: 25426837
177. Genovese G, et al. N. Engl. J. Med. (2014) pmid: 25426838
178. Xie M, et al. Nat. Med. (2014) pmid: 25326804
179. Acuna-Hidalgo R, et al. Am. J. Hum. Genet. (2017) pmid: 28669404
180. Severson EA, et al. Blood (2018) pmid: 29678827
181. Fuster JJ, et al. Circ. Res. (2018) pmid: 29420212
182. Hematology Am Soc Hematol Educ Program (2018) pmid: 30504320
183. Chabon JJ, et al. Nature (2020) pmid: 32269342
184. Razavi P, et al. Nat. Med. (2019) pmid: 31768066
185. Glassberg et al., 2020; ASPHO Abstract 2015
186. Coyne et al., 2020; ASCO Abstract 3612
187. McCowage et al., 2018; ASCO Abstract 10504
188. Mueller et al., 2020; SNO Abstract NFB-17
189. Waldner et al., 2020; DOI: 10.1055/s-0040-1715638
190. Romo et al., 2019; SNO Abstract RARE-54
191. Diamond EL, et al. Cancer Discov (2016) pmid: 26566875
192. Diamond EL, et al. Nature (2019) pmid: 30867592
193. Jacobsen E, et al. N. Engl. J. Med. (2017) pmid: 29236635
194. Berce PC, et al. JAAD Case Rep (2022) pmid: 35372654
195. Giuffrè C, et al. Indian J Ophthalmol (2020) pmid: 32823478
196. Razanamahery J, et al. Clin Exp Rheumatol () pmid: 31376269
197. Moyon Q, et al. Chest (2020) pmid: 31669429
198. Gupta RK, et al. Neurol Int (2022) pmid: 36135991
199. Rosen LS, et al. Invest New Drugs (2016) pmid: 27424159
200. Cho et al., 2020; AACR Abstract CT201
201. Touat et al., 2019; DOI: 10.1200/PO.18.00298
202. Banerjee A, et al. Neuro-oncology (2017) pmid: 28339824
203. Gupta A, et al. Ann. Oncol. (2014) pmid: 24567366
204. Robert C, et al. Lancet Oncol. (2013) pmid: 23735514
205. Kirkwood JM, et al. Clin. Cancer Res. (2012) pmid: 22048237
206. Banerji U, et al. Clin. Cancer Res. (2010) pmid: 20179232
207. Boers-Sonderen MJ, et al. Anticancer Drugs (2012) pmid: 22293660
208. Lopez-Chavez A, et al. J. Clin. Oncol. (2015) pmid: 25667274
209. Hainsworth JD, et al. J Thorac Oncol (2010) pmid: 20802351
210. Coleman RL, et al. Gynecol. Oncol. (2015) pmid: 25887099
211. Eckstein OS, et al. J Clin Oncol (2022) pmid: 35363510
212. Deming DA, et al. Invest New Drugs (2016) pmid: 26666244
213. Krishnamurthy A, et al. Cancer Res. (2018) pmid: 30042150
214. Infante JR, et al. Invest New Drugs (2017) pmid: 28424891
215. LoRusso PM, et al. BMC Cancer (2017) pmid: 28264648
216. Tolcher AW, et al. Clin. Cancer Res. (2015) pmid: 25516890
217. Wilky BA, et al. Br. J. Cancer (2015) pmid: 25268371
218. Infante JR, et al. Lancet Oncol. (2012) pmid: 22805291
219. Leijen S, et al. Clin. Cancer Res. (2012) pmid: 22767668
220. Zimmer L, et al. Clin. Cancer Res. (2014) pmid: 24947927
221. Infante JR, et al. Eur. J. Cancer (2013) pmid: 23583440
222. Juric et al., 2014; ASCO Abstract 9051
223. Tolcher AW, et al. Ann. Oncol. (2015) pmid: 25344362
224. Patterson et al., 2018; AACR Abstract 3891

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Nimesh Patel, M.D. | 12 December 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531