

ABOUT THE TEST FoundationOne®Liquid CDx is a next generation sequencing (NGS) assay that identifies clinically relevant genomic alterations in circulating cell-free DNA.

PATIENT
DISEASE Gallbladder carcinoma
NAME Chang Yang, Hsueh-Hua
DATE OF BIRTH 17 July 1947
SEX Female
MEDICAL RECORD # 17527343

PHYSICIAN
ORDERING PHYSICIAN Yeh, Yi-Chen
MEDICAL FACILITY Taipei Veterans General Hospital
ADDITIONAL RECIPIENT None
MEDICAL FACILITY ID 205872
PATHOLOGIST Not Provided

SPECIMEN
SPECIMEN ID HHCY 07/17/1947
SPECIMEN TYPE Blood
DATE OF COLLECTION 06 December 2022
SPECIMEN RECEIVED 12 December 2022

Biomarker Findings

Blood Tumor Mutational Burden - 8 Muts/Mb
Microsatellite status - MSI-High Not Detected
Tumor Fraction - Elevated Tumor Fraction

Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

AKT2 amplification
ATM L1524fs*7
CCNE1 amplification
CASP8 G20fs*18
DNMT3A E463fs*188
RB1 K95fs*15
TP53 D259Y, I195T

Report Highlights

- Evidence-matched clinical trial options based on this patient's genomic findings: (p. [10](#))
- Variants that may represent clonal hematopoiesis and may originate from non-tumor sources: **ATM** L1524fs*7 (p. [6](#)), **DNMT3A** E463fs*188 (p. [8](#))

BIOMARKER FINDINGS

Blood Tumor Mutational Burden -
 8 Muts/Mb

Microsatellite status -
 MSI-High Not Detected

Tumor Fraction -
 Elevated Tumor Fraction

THERAPY AND CLINICAL TRIAL IMPLICATIONS

No therapies or clinical trials. See Biomarker Findings section

MSI-High not detected. No evidence of microsatellite instability in this sample (see Appendix section).

Tumor fraction is considered elevated when ctDNA levels are high enough that aneuploidy can be detected. There is higher sensitivity for identifying genomic alterations and a lower risk of false negative results in specimens with elevated tumor fraction; the positive percent agreement observed between liquid and tissue for defined short variants is $\geq 90\%$ (Li et al., 2021; AACR Abstract 2231) (see Biomarker Findings section).

GENOMIC FINDINGS

VAF%

AKT2 -	amplification	-
10 Trials see p. 10		
ATM -	L1524fs*7	0.09%
10 Trials see p. 12		
CCNE1 -	amplification	-
4 Trials see p. 14		

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)

THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)

None	None
None	None
None	None

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Julie Tse, M.D. | 22 December 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS (CH)

Genomic findings below may include nontumor somatic alterations, such as CH. The efficacy of targeting such nontumor somatic alterations is unknown. This content should be interpreted based on clinical context. Refer to appendix for additional information on CH.

ATM - L1524fs*7 p. [6](#) **DNMT3A - E463fs*188** p. [8](#)

GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

CASP8 - G20fs*18 p. [7](#) **RB1 - K95fs*15** p. [8](#)
DNMT3A - E463fs*188 p. [8](#) **TP53 - D259Y, I195T** p. [9](#)

NOTE Genomic alterations detected may be associated with activity of certain approved therapies; however, the therapies listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and/or exhaustive. Neither the therapies nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies. Therapies contained in this report may have been approved by the US FDA or other national authorities; however, they might not have been approved in your respective country. In the appropriate clinical context, germline testing of APC, ATM, BAP1, BRCA1, BRCA2, BRIP1, CHEK2, FH, FLCN, MEN1, MLH1, MSH2, MSH6, MUTYH, NF1, NF2, PALB2, PMS2, POLE, PTEN, RAD51C, RAD51D, RB1, RET, SDHA, SDHB, SDHC, SDHD, SMAD4, STK11, TGFB2, TP53, TSC1, TSC2, VHL, and WT1 is recommended.

Variant Allele Frequency is not applicable for copy number alterations.


Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Julie Tse, M.D. | 22 December 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1521387-01



FoundationOne®Liquid CDx
22 Dec 2022

HISTORIC PATIENT FINDINGS

ORD-1521387-01
VAF%

Blood Tumor Mutational Burden

8 Muts/Mb

Microsatellite status

MSI-High Not Detected

Tumor Fraction

26%

AKT2	amplification	Detected
ATM	● L1524fs*7	0.09%
CCNE1	amplification	Detected
CASP8	● G20fs*18	69.5%
DNMT3A	● E463fs*188	1.1%
RB1	● K95fs*15	23.8%
TP53	● I195T	0.48%
	● D259Y	20.7%

NOTE This comparison table refers only to genes and biomarkers assayed by prior FoundationOne®Liquid CDx or FoundationOne®CDx tests. Up to five previous tests may be shown.

For some genes in FoundationOne Liquid CDx, only select exons are assayed. Therefore, an alteration found by a previous test may not have been confirmed despite overlapping gene lists. Please refer to the Appendix for the complete list of genes and exons assayed. The gene and biomarker list will be updated periodically to reflect new knowledge about cancer biology.

As new scientific information becomes available, alterations that had previously been listed as Variants of Unknown Significance (VUS) may become reportable.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Julie Tse, M.D. | 22 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1521387-01

Tissue Tumor Mutational Burden (TMB) and blood TMB (bTMB) are estimated from the number of synonymous and non-synonymous single-nucleotide variants (SNVs) and insertions and deletions (indels) per area of coding genome sampled, after the removal of known and likely oncogenic driver events and germline SNPs. Tissue TMB is calculated based on variants with an allele frequency of $\geq 5\%$, and bTMB is calculated based on variants with an allele frequency of $\geq 0.5\%$.

Not Tested = not baited, not reported on test, or test preceded addition of biomarker or gene

Not Detected = baited but not detected on test

Detected = present (VAF% is not applicable)

VAF% = variant allele frequency percentage

Cannot Be Determined = Sample is not of sufficient data quality to confidently determine biomarker status

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Julie Tse, M.D. | 22 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1521387-01

BIOMARKER FINDINGS

BIOMARKER

Blood Tumor Mutational Burden

RESULT

8 Muts/Mb

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased blood tumor mutational burden (bTMB) may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1¹⁻³, anti-PD-1³⁻⁴, anti-PD-1/CTLA4 therapies⁵⁻⁶, anti-PD-L1/CTLA4 therapies⁷⁻¹⁰. A Phase 2 multi-solid-tumor trial showed that bTMB ≥ 16 Muts/Mb (as measured by this assay) was associated with improved survival from treatment with a PD-1 inhibitor alone or in combination with a CTLA-4 inhibitor⁵. In non-small cell lung cancer (NSCLC), multiple clinical trials have shown patients with higher bTMB derive clinical benefit from immune checkpoint inhibitors following single-agent or combination treatments with either CTLA4

inhibitors or chemotherapy, with reported high bTMB cutpoints ranging from 6 Muts/Mb-16 Muts/Mb¹⁸⁻¹⁰. In head and neck squamous cell carcinoma (HNSCC), a Phase 3 trial showed that bTMB ≥ 16 Muts/Mb (approximate equivalency ≥ 8 Muts/Mb as measured by this assay) was associated with improved survival from treatment with a PD-L1 inhibitor alone or in combination with a CTLA-4 inhibitor¹¹. In colorectal cancer (CRC), a Phase 2 study showed that bTMB ≥ 28 Muts/Mb (approximate equivalency ≥ 14 Muts/Mb as measured by this assay) was associated with improved OS from a PD-L1 inhibitor alone or in combination with a CTLA-4 inhibitor⁷.

FREQUENCY & PROGNOSIS

Average bTMB levels in solid tumors other than NSCLC have not been evaluated (PubMed, Mar 2022). Published data investigating the prognostic implications of bTMB levels in biliary tract cancer are limited (PubMed, Jul 2022). Although cases with hypermutated biliary tract cancer were enriched in a subgroup with poor prognosis in 1 study¹², TMB-high (≥ 10 mut/Mb) status in biliary adenocarcinoma not treated with immunotherapy was not significantly associated with OS in another

study, in which patients with TMB-high tumors experienced numerically longer OS compared with patients with TMB-low tumors (11.5 vs. 8.4 months, adjusted HR=0.65)¹³.

FINDING SUMMARY

Blood tumor mutational burden (bTMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations from circulating tumor DNA in blood. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma¹⁴⁻¹⁵ and cigarette smoke in lung cancer¹⁶⁻¹⁷, treatment with temozolomide-based chemotherapy in glioma¹⁸⁻¹⁹, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes²⁰⁻²⁴, and microsatellite instability (MSI)^{20,23-24}. High bTMB levels were not detected in this sample. It is unclear whether the bTMB levels in this sample would be predicted to be associated with sensitivity to PD-1- or PD-L1-targeting immune checkpoint inhibitors, alone or in combination with other agents¹⁻²⁴. Depending on the clinical context, TMB testing of an alternate sample or by another methodology could be considered.

BIOMARKER

Tumor Fraction

RESULT

Elevated Tumor Fraction

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Specimens with elevated tumor fraction have high circulating-tumor DNA (ctDNA) content, and thus high sensitivity for identifying genomic alterations. Such specimens are at low risk of false negative results. Tumor fraction levels currently have limited implications for diagnosis, surveillance, or therapy and should not be overinterpreted or compared from one blood draw to another. There are currently no targeted approaches to address

specific tumor fraction levels. In the research setting, changes in tumor fraction estimates have been associated with treatment duration and clinical response and may be a useful indicator for future cancer management²⁵⁻³⁰.

FREQUENCY & PROGNOSIS

Detectable ctDNA levels have been reported in a variety of tumor types, with higher tumor fraction levels reported for patients with metastatic (Stage 4) tumors compared with patients with localized disease (Stages 1 to 3)³¹. Elevated tumor fraction levels have been reported to be associated with worse prognosis in a variety of cancer types, including pancreatic cancer³², Ewing sarcoma and osteosarcoma³³, prostate cancer²⁸, breast cancer³⁴, leiomyosarcoma³⁵, esophageal cancer³⁶, colorectal cancer³⁷, and gastrointestinal cancer³⁸.

FINDING SUMMARY

Tumor fraction provides an estimate of the percentage of ctDNA present in a cell-free DNA (cfDNA) sample. The tumor fraction estimate for this sample is based on the observed level of aneuploid instability. The tumor fraction algorithm utilized for FoundationOne Liquid CDx uses the allele frequencies of approximately 1,000 single-nucleotide polymorphism (SNP) sites across the genome. Unlike the maximum somatic allele frequency (MSAF) method of estimating ctDNA content³⁹, the tumor fraction metric does not take into account the allele frequency of individual variants but rather produces a more holistic estimate of ctDNA content using data from across the genome. The amount of ctDNA detected may correlate with disease burden and response to therapy⁴⁰⁻⁴¹.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Julie Tse, M.D. | 22 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1521387-01

GENOMIC FINDINGS

GENE

AKT2

ALTERATION
amplification

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Amplification or activation of AKT2 may promote AKT-mTOR pathway activation and may predict sensitivity to inhibitors of the AKT and downstream mTOR pathways. Clinical benefit has been achieved in 1 patient with AKT2 amplification treated with an mTOR inhibitor⁴². In preclinical studies, the AKT inhibitor MK-2206 showed

evidence of enhancing anti-tumor activity of other chemotherapeutic agents in lung and ovarian tumor cells⁴³.

FREQUENCY & PROGNOSIS

AKT2 amplification has been reported most frequently in endometrial cancer (3.7%), ovarian cancer (3.6%), cancer of unknown primary (CUP) (2.2%), pancreatic cancer (1.8%), head and neck cancer (1.6%), and bladder cancer (1.4%)⁴⁴. One study found AKT2 protein expression to be more common in hilar cholangiocarcinomas (54%) than peripheral cholangiocarcinomas (27%)⁴⁵. AKT expression was associated with favorable prognosis in cholangiocarcinoma⁴⁶, while another study reported no prognosis relevance of pAKT in the intrahepatic cholangiocarcinoma⁴⁷.

FINDING SUMMARY

AKT2 encodes an intracellular serine/threonine kinase that is also known as PKB-beta. AKT2 is one of three members of the AKT gene family, and activation of AKT2 has been implicated in multiple malignancies⁴⁸⁻⁴⁹. AKT isoforms appear to have different roles in tumorigenesis; AKT1 appears to contribute to the initiation of tumors, whereas AKT2 promotes invasion and metastasis in breast tumors⁵⁰. Although AKT2 amplification has been reported to associate with AKT2 overexpression⁵¹⁻⁵³, studies in various cancers suggest that AKT2 phosphorylation may have greater clinical relevance than AKT2 amplification or mRNA overexpression⁵⁴⁻⁵⁵.

GENE

ATM

ALTERATION
L1524fs*7

TRANSCRIPT ID
NM_000051.3

CODING SEQUENCE EFFECT
4569_4570insA

VARIANT CHROMOSOMAL POSITION
chr11:108163477

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Loss of functional ATM results in a defective DNA damage response and homologous recombination-mediated DNA repair and may predict sensitivity to PARP inhibitors⁵⁶⁻⁵⁷. Clinical responses have been reported for patients with ATM-mutated prostate cancer treated with PARP inhibitors⁵⁸⁻⁶⁰ and PARP inhibitors have shown limited clinical benefit for patients with other ATM-mutated solid tumors including pancreatic cancer⁶¹⁻⁶², colorectal cancer⁶³, papillary renal cell carcinoma⁶⁴, ovarian cancer⁶⁵, small cell bowel cancer⁶², and biliary tract cancer⁶⁶. In Phase 1 trials of ATR inhibitors, a heavily pretreated patient with colorectal cancer who achieved a CR to berzosertib⁶⁷ and 4 out of 4 patients with diverse solid tumors who achieved PRs to BAY1895344⁶⁸ harbored ATM inactivation

or protein loss. In a Phase 2 study of a combination of the ATR inhibitor ceralasertib and durvalumab for patients with advanced gastric cancer, objective responses (ORs) were experienced by 50% (4/8) of patients with loss of ATM expression, compared with 14% (3/21) patients with intact ATM⁶⁹. Studies showing reduced cell viability and increased DNA damage in preclinical models of solid tumors⁷⁰⁻⁷² and hematologic malignancies^{70,73} also support the increased sensitivity of ATM-deficient cells to ATR inhibitors. Preclinical experiments also indicate that loss of ATM causes dependency on DNA-PKcs in cancer cells; DNA-PKcs inhibitors promoted apoptosis in ATM-deficient cells and were active in a lymphoma mouse model lacking ATM activity⁷⁴.

FREQUENCY & PROGNOSIS

ATM alterations have been observed in various solid tumors, with mutations being more frequent than gene loss and often seen in small bowel (7-9%), endometrial (7%), non-melanoma skin (3-6%), bladder (5%), hepatobiliary (4-5%), colorectal (4-5%), and lung (3%) cancer^{44,75}. Published data investigating the prognostic implications of ATM alterations in biliary tract carcinoma are limited (PubMed, Feb 2022).

FINDING SUMMARY

ATM encodes the protein ataxia telangiectasia mutated, which is a serine/threonine protein kinase that plays a key role in the DNA damage

response⁷⁶. Loss of functional ATM promotes tumorigenesis⁷⁷. Alterations such as seen here may disrupt ATM function or expression⁷⁸⁻⁸⁰.

POTENTIAL GERMLINE IMPLICATIONS

ATM mutation carriers have increased cancer risk, with female carriers displaying a 38% lifetime risk of breast cancer⁸¹. Biallelic mutations in ATM underlie the rare autosomal-recessive inherited disorder ataxia-telangiectasia (A-T), also referred to as genome instability or DNA damage response syndrome⁸². This disease is characterized by genomic instability, sensitivity to DNA-damaging agents, and increased risk of developing cancer^{76,82}. The prevalence of A-T is estimated at 1:40,000 to 1:100,000 worldwide⁸². In the appropriate clinical context, germline testing of ATM is recommended.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion⁸³⁻⁸⁸. Comprehensive genomic profiling of solid tumors may detect nontumor alterations that are due to CH^{87,89-90}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Julie Tse, M.D. | 22 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1521387-01

GENOMIC FINDINGS

GENE
CCNE1

ALTERATION
amplification

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved therapies that directly target CCNE1 alterations. Because amplification or overexpression of CCNE1 leads to increased genomic instability through the ATR-Chk1-WEE1 pathway⁹¹⁻⁹² and cyclin E1 promotes cell cycle progression in a complex with CDK2⁹³, clinical and preclinical studies have investigated inhibitors of Chk1, ATR, CDK2, and WEE1 as potential therapeutic approaches for tumors with CCNE1 activation. Clinical benefit has been reported for patients with recurrent high-grade serous ovarian carcinoma (HGSOC) with CCNE1 amplification or expression in response to treatment with the

Chk1 inhibitor prexasertib⁹⁴. Studies of the WEE1 inhibitor adavosertib observed PRs in patients with CCNE1-amplified HGSOC and ovarian cancer⁹⁵⁻⁹⁶. Similarly, in a Phase 2 study of patients with CCNE1-amplified solid tumors, adavosertib elicited an ORR of 26% with PRs reported for patients with ovarian cancer, urothelial carcinoma, or melanoma⁹⁷. Preclinical studies have demonstrated that cell lines with CCNE1 amplification or overexpression were sensitive to inhibitors of ATR⁹⁸⁻⁹⁹, CDK2¹⁰⁰, or WEE1^{92,101}. However, other studies have shown that sensitivity of various cell lines to CDK2 inhibitors, including SNS-032, dinaciclib, and seliciclib, at clinically achievable doses, is largely independent of CCNE1 copy number or expression¹⁰²⁻¹⁰⁵. One study has reported a reduction in tumor CCNE1 levels in 4/6 lung and esophageal cancer cases following treatment with the HDAC inhibitor vorinostat¹⁰⁶.

FREQUENCY & PROGNOSIS

In one study, CCNE1 amplification was observed 7% of in gallbladder samples¹⁰⁷. CCNE1

amplification has been reported in 1-2% of samples in published cholangiocarcinoma datasets¹⁰⁸⁻¹⁰⁹. Published data investigating the prognostic implications of CCNE1 alterations in biliary tract carcinoma are limited (PubMed, Mar 2022).

FINDING SUMMARY

CCNE1 encodes the protein cyclin E1, which plays a role in the regulated transition from the G1 to S phase by binding to and activating cyclin-dependent protein kinase 2 (CDK2). It also has a direct role in initiation of replication and the maintenance of genomic stability⁹³. Amplification of chromosomal region 19q12-q13 has been demonstrated in many types of cancer, and CCNE1 is a well-studied gene within this amplicon¹¹⁰⁻¹¹¹. Increased copy number of CCNE1 is highly associated with overexpression of the cyclin E1 protein¹¹²⁻¹¹³. Cyclin E1 overexpression can lead to cell transformation as a result of an increase in cyclin E1 activity^{93,114}.

GENE
CASP8

ALTERATION
G20fs*18

TRANSCRIPT ID
NM_001080125.1

CODING SEQUENCE EFFECT
56_57insAACTTCTTCT

VARIANT CHROMOSOMAL POSITION
chr2:202122999

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted approaches to directly address alterations in CASP8.

FREQUENCY & PROGNOSIS

CASP8 mutations have been observed across solid tumors⁴⁴, including 8.2% of head and neck squamous cell¹¹⁵, 3.2% of colorectal¹¹⁶, and 4% of cervical¹¹⁷ carcinoma cases. Reduced expression of CASP8 has been associated with poor prognosis in ovarian cancer¹¹⁸, prostate cancer¹¹⁹, and medulloblastoma¹²⁰.

FINDING SUMMARY

CASP8 encodes caspase-8, a multifunctional protein that mediates apoptosis¹²¹⁻¹²⁴, cell motility¹²⁵⁻¹²⁶, and cell signaling, including through the NFkB¹²⁷⁻¹²⁹ and MAPK¹³⁰⁻¹³¹ pathways. The role of CASP8 in cancer is complex and context-dependent, with diverse cancer types exhibiting either overexpression or loss of expression. CASP8 mutations found in the context of cancer tend to be truncating or missense mutations; most of the characterized mutations impair apoptosis¹³²⁻¹³⁶ and promote NFkB activation¹³⁷.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Julie Tse, M.D. | 22 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1521387-01

GENOMIC FINDINGS

GENE

DNMT3A

ALTERATION

E463fs*188

TRANSCRIPT ID

NM_022552.3

CODING SEQUENCE EFFECT

1388delA

VARIANT CHROMOSOMAL POSITION

chr2:25469069-25469070

FREQUENCY & PROGNOSIS

DNMT3A alterations have been reported at relatively low frequencies in solid tumors and are more prevalent in hematological malignancies (cBioPortal, Feb 2022)¹³⁸⁻¹³⁹. Published data investigating the prognostic implications of DNMT3A alterations in solid tumors are limited (PubMed, Feb 2022).

FINDING SUMMARY

The DNMT3A gene encodes the protein DNA methyltransferase 3A, an enzyme that is involved in the methylation of newly synthesized DNA, a function critical for gene regulation¹⁴⁰⁻¹⁴¹. The role of DNMT3A in cancer is uncertain, as some reports describe increased expression and contribution to tumor growth, whereas others propose a role for DNMT3A as a tumor suppressor¹⁴²⁻¹⁴⁷. Alterations such as seen here may disrupt DNMT3A function or expression¹⁴⁸⁻¹⁵¹.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion⁸³⁻⁸⁸. CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy⁸³⁻⁸⁴. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease¹⁵². Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH^{87,89-90}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted therapies available to address genomic alterations in DNMT3A in solid tumors.

GENE

RB1

ALTERATION

K95fs*15

TRANSCRIPT ID

NM_000321.2

CODING SEQUENCE EFFECT

284_287delAAAA

VARIANT CHROMOSOMAL POSITION

chr13:48916752-48916756

kinase A, particularly in small cell lung cancer (SCLC). A clinical study evaluating the Aurora kinase A inhibitor alisertib for patients with prostate cancer did not find an association between RB1 deletion and clinical benefit¹⁵⁸. Other approaches to target RB1 inactivation under investigation in preclinical studies include inhibitors of BCL-2 family members¹⁵⁹ and activation of the NOTCH pathway¹⁶⁰.

FREQUENCY & PROGNOSIS

RB1 mutations have been reported across solid tumors including small cell lung cancer (60%), bladder (13%), uterine sarcoma (10%), non-melanoma skin cancer (9.5%), and gastrointestinal neuroendocrine tumors (9.3%)⁷⁵. The prognostic significance of RB1 mutation in gallbladder carcinoma has not been extensively studied (PubMed, Jul 2022).

FINDING SUMMARY

RB1 encodes the retinoblastoma protein (Rb), a tumor suppressor and negative regulator of the cell cycle¹⁶¹⁻¹⁶². Alterations such as seen here may disrupt RB1 function or expression¹⁶³⁻¹⁶⁹.

POTENTIAL GERMLINE IMPLICATIONS

Mutations in RB1 underlie the development of retinoblastoma (RB), a rare tumor that arises at a rate of approximately 1:20,000 live births, with nearly 5,000 new cases worldwide per year¹⁷⁰. Germline mutations in RB1 account for approximately 40% of RB tumors¹⁷¹ and are associated with an increased risk of developing secondary malignancies that include soft tissue and bone sarcoma and malignant melanoma¹⁷²⁻¹⁷³. In the appropriate clinical context, germline testing of RB1 is recommended.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of limited clinical data¹⁵³ and strong preclinical data¹⁵⁴⁻¹⁵⁷, RB1 inactivation may be associated with sensitivity to inhibitors of Aurora

ORDERED TEST # ORD-1521387-01

GENOMIC FINDINGS

GENE

TP53

ALTERATION

D259Y, I195T

TRANSCRIPT ID

NM_000546.4, NM_000546.4

CODING SEQUENCE EFFECT

775G>T, 584T>C

VARIANT CHROMOSOMAL POSITION

chr17:7577506, chr17:7578265

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved therapies to address TP53 mutation or loss. However, tumors with TP53 loss of function alterations may be sensitive to the WEE1 inhibitor adavosertib¹⁷⁴⁻¹⁷⁷ or p53 gene therapy such as SGT53¹⁷⁸⁻¹⁸². In a Phase 1 study, adavosertib in combination with gemcitabine, cisplatin, or carboplatin elicited PRs in 9.7% and SDs in 53% of patients with solid tumors; the response rate was 21% (4/19) for patients with TP53 mutations versus 12% (4/33) for patients who were TP53 wildtype¹⁸³. A Phase 2 trial of adavosertib in combination with chemotherapy (gemcitabine, carboplatin, paclitaxel, or doxorubicin) reported a 32% (30/94, 3 CR) ORR and a 73% (69/94) DCR for patients with platinum-refractory TP53-mutated ovarian, Fallopian tube, or peritoneal cancer¹⁸⁴. A smaller Phase 2 trial of adavosertib in combination with carboplatin achieved a 43% (9/21, 1 CR) ORR and a 76% (16/21) DCR for patients with platinum-refractory TP53-mutated ovarian cancer¹⁸⁵. The combination of adavosertib with paclitaxel and carboplatin for patients with TP53-mutated ovarian cancer also significantly increased PFS compared with paclitaxel and carboplatin alone¹⁸⁶. In the Phase 2 VIKTORY trial, patients with TP53-mutated metastatic and/or recurrent gastric cancer experienced a 24% (6/25) ORR with adavosertib combined with paclitaxel¹⁸⁷. A Phase 1 trial of neoadjuvant adavosertib in combination with cisplatin and docetaxel for head and neck squamous cell carcinoma (HNSCC) elicited a 71%

(5/7) response rate for patients with TP53 alterations¹⁸⁸. The Phase 2 FOCUS4-C trial for patients with TP53- and RAS-mutated colorectal cancer reported improvement in PFS (3.61 vs. 1.87 months, HR=0.35, p=0.0022), but not OS (14.0 vs 12.8 months, p=0.93), following adavosertib treatment compared with active monitoring¹⁸⁹. In a Phase 1b clinical trial of SGT-53 in combination with docetaxel for patients with solid tumors, 75% (9/12) of evaluable patients experienced clinical benefit, including 2 confirmed and 1 unconfirmed PRs and 2 instances of SD with significant tumor shrinkage¹⁸². Missense mutations leading to TP53 inactivation may be sensitive to therapies that reactivate mutated p53 such as eprenetapopt. In a Phase 1b trial for patients with p53-positive high-grade serous ovarian cancer, eprenetapopt combined with carboplatin and pegylated liposomal doxorubicin achieved a 52% (11/21) response rate and 100% DCR¹⁹⁰. A Phase 1 trial of eprenetapopt with pembrolizumab for patients with solid tumors reported an ORR of 10% (3/29)¹⁹¹.

FREQUENCY & PROGNOSIS

Inactivation of p53, through mutation, deletion, or loss of heterozygosity (LOH), has been observed in 25-63% of gallbladder carcinomas and 10-61% of cholangiocarcinomas^{12,192-200}. TP53 mutations occur more frequently in tumors caused by liver fluke (*O. viverrini*) infection (40%) than in cholangiocarcinoma cases not related to infection (9%)¹⁹⁵. TP53 alteration and inactivation were shown to be early and common events in gallbladder carcinogenesis²⁰¹. Aberrant TP53 expression, which is indicative of TP53 dysregulation, has been observed in 20-62% of gallbladder carcinomas and 25% (5/20) of cholangiocarcinomas²⁰²⁻²⁰⁴. Data regarding the prognostic significance of TP53 mutation in cholangiocarcinoma are conflicting²⁰⁵⁻²¹³. Overexpression of p53 protein has been associated with reduced patient survival in poorly differentiated gallbladder adenocarcinomas and biliary tract cancers²¹⁴⁻²¹⁵; however, another study did not find such a correlation²⁰⁷.

FINDING SUMMARY

Functional loss of the tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers²¹⁶. Alterations such as seen here may disrupt TP53 function or expression²¹⁷⁻²²¹.

POTENTIAL GERMLINE IMPLICATIONS

One or more of the TP53 variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters) associated with Li-Fraumeni syndrome (ClinVar, Sep 2022)²²². Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. Germline mutations in TP53 are associated with the very rare autosomal dominant disorder Li-Fraumeni syndrome and the early onset of many cancers²²³⁻²²⁵, including sarcomas²²⁶⁻²²⁷. Estimates for the prevalence of germline TP53 mutations in the general population range from 1:5,000²²⁸ to 1:20,000²²⁷. For pathogenic TP53 mutations identified during tumor sequencing, the rate of germline mutations was 1% in the overall population and 6% in tumors arising before age 30²²⁹. In the appropriate clinical context, germline testing of TP53 is recommended.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion⁸³⁻⁸⁸. CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy⁸³⁻⁸⁴. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease¹⁵². Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH^{87,89-90}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Julie Tse, M.D. | 22 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1521387-01

CLINICAL TRIALS

IMPORTANT Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually updated and should be investigated by the physician or

research staff. This is not a comprehensive list of all available clinical trials. There may also be compassionate use or early access programs available, which are not listed in this report. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials are not ranked in order of potential or predicted efficacy for this patient or

in order of level of evidence for this patient's tumor type. Clinical trials listed here may have additional enrollment criteria that may require medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see clinicaltrials.gov. However, clinicaltrials.gov does not list all clinical trials that might be available.

GENE
AKT2
RATIONALE
 AKT2 amplification or mutation may lead to AKT-mTOR pathway activation and may predict

sensitivity to inhibitors of this pathway.

ALTERATION
 amplification

NCT04341259
PHASE 1

A Study Of The Pharmacokinetics And Safety Of Ipatasertib In Chinese Participants With Locally Advanced Or Metastatic Solid Tumors.

TARGETS
 AKTs

LOCATIONS: Shanghai City (China)

NCT03239015
PHASE 2

Efficacy and Safety of Targeted Precision Therapy in Refractory Tumor With Druggable Molecular Event

TARGETS
 EGFR, ERBB4, ERBB2, PARP, mTOR, MET, ROS1, RET, VEGFRs, BRAF, CDK4, CDK6

LOCATIONS: Shanghai (China)

NCT04803318
PHASE 2

Trametinib Combined With Everolimus and Lenvatinib for Recurrent/Refractory Advanced Solid Tumors

TARGETS
 mTOR, FGFRs, RET, PDGFRA, VEGFRs, KIT, MEK

LOCATIONS: Guangzhou (China)

NCT05125523
PHASE 1

A Study of Sirolimus for Injection (Albumin Bound) in Patients With Advanced Solid Tumors

TARGETS
 mTOR

LOCATIONS: Tianjin (China)

NCT03772561
PHASE 1

Phase I Study of AZD5363 + Olaparib + Durvalumab in Patients With Advanced or Metastatic Solid Tumor Malignancies

TARGETS
 PARP, AKTs, PD-L1

LOCATIONS: Singapore (Singapore)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Julie Tse, M.D. | 22 December 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1521387-01

CLINICAL TRIALS
NCT03297606
PHASE 2

Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR)

TARGETS

VEGFRs, ABL, SRC, ALK, ROS1, AXL, TRKA, MET, TRKC, DDR2, KIT, EGFR, PD-1, CTLA-4, PARP, CDK4, CDK6, FLT3, CSF1R, RET, mTOR, ERBB2, MEK, BRAF, SMO

LOCATIONS: Vancouver (Canada), Edmonton (Canada), Saskatoon (Canada), Regina (Canada), Ottawa (Canada), Montreal (Canada), Toronto (Canada), Kingston (Canada), London (Canada)

NCT04551521
PHASE 2

CRAFT: The NCT-PMO-1602 Phase II Trial

TARGETS

PD-L1, AKTs, MEK, BRAF, ALK, RET, ERBB2

LOCATIONS: Würzburg (Germany), Mainz (Germany), Heidelberg (Germany), Tübingen (Germany)

NCT03673787
PHASE 1/2

A Trial of Ipatasertib in Combination With Atezolizumab

TARGETS

AKTs, PD-L1

LOCATIONS: Sutton (United Kingdom)

NCT01582191
PHASE 1

A Phase 1 Trial of Vandetanib (a Multi-kinase Inhibitor of EGFR, VEGFR and RET Inhibitor) in Combination With Everolimus (an mTOR Inhibitor) in Advanced Cancer

TARGETS

mTOR, EGFR, SRC, RET, VEGFRs

LOCATIONS: Texas

NCT03203525
PHASE 1

Combination Chemotherapy and Bevacizumab With the NovoTTF-100L(P) System in Treating Participants With Advanced, Recurrent, or Refractory Hepatic Metastatic Cancer

TARGETS

VEGFA, mTOR

LOCATIONS: Texas

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Julie Tse, M.D. | 22 December 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1521387-01

CLINICAL TRIALS
GENE
ATM
RATIONALE
 Loss or inactivation of ATM may increase sensitivity to PARP inhibitors, ATR inhibitors, or

DNA-PKcs inhibitors.

ALTERATION
 L1524fs*7

NCT04644068
PHASE 1/2

Study of AZD5305 as Monotherapy and in Combination With Anti-cancer Agents in Patients With Advanced Solid Malignancies

TARGETS
 ERBB2, TROP2, PARP

LOCATIONS: Shanghai (China), Guangzhou (China), Seoul (Korea, Republic of), Chongqing (China), Chengdu (China), Chuo-ku (Japan), Koto-ku (Japan), Melbourne (Australia), Lublin (Poland), Warszawa (Poland)

NCT04123366
PHASE 2

Study of Olaparib (MK-7339) in Combination With Pembrolizumab (MK-3475) in the Treatment of Homologous Recombination Repair Mutation (HRRm) and/or Homologous Recombination Deficiency (HRD)-Positive Advanced Cancer (MK-7339-007/KEYLYNK-007)

TARGETS
 PARP, PD-1

LOCATIONS: Fukuoka (Japan), Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Okayama (Japan), Nagoya (Japan), Tokyo (Japan), Kashiwa (Japan), Sapporo (Japan), Nedlands (Australia), Southport (Australia)

NCT03742895
PHASE 2

Efficacy and Safety of Olaparib (MK-7339) in Participants With Previously Treated, Homologous Recombination Repair Mutation (HRRm) or Homologous Recombination Deficiency (HRD) Positive Advanced Cancer (MK-7339-002 / LYNK-002)

TARGETS
 PARP

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Nedlands (Australia), Darlinghurst (Australia), Adana (Turkey), Jerusalem (Israel), Konya (Turkey), Ramat Gan (Israel), Beer-Sheva (Israel), Istanbul (Turkey)

NCT02264678
PHASE 1/2

Ascending Doses of AZD6738 in Combination With Chemotherapy and/or Novel Anti Cancer Agents

TARGETS
 ATR, PARP, PD-L1

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Goyang-si (Korea, Republic of), Cambridge (United Kingdom), Withington (United Kingdom), Manchester (United Kingdom), London (United Kingdom), Coventry (United Kingdom), Sutton (United Kingdom), Oxford (United Kingdom)

NCT04298008
PHASE 2

AZD6738 Plus Durvalumab in Biliary Tract Cancer

TARGETS
 ATR, PD-L1

LOCATIONS: Seoul (Korea, Republic of)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Julie Tse, M.D. | 22 December 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1521387-01

CLINICAL TRIALS
NCT04298021
PHASE 2

DDR-Umbrella Study of DDR Targeting Agents in Advanced Biliary Tract Cancer

TARGETS
 PD-L1, ATR, PARP

LOCATIONS: Seoul (Korea, Republic of)

NCT05035745
PHASE 1/2

Selinexor & Talazoparib in Advanced Refractory Solid Tumors; Advanced/Metastatic Triple Negative Breast Cancer (START)

TARGETS
 XPO1, PARP

LOCATIONS: Singapore (Singapore)

NCT03772561
PHASE 1

Phase I Study of AZD5363 + Olaparib + Durvalumab in Patients With Advanced or Metastatic Solid Tumor Malignancies

TARGETS
 PARP, AKTs, PD-L1

LOCATIONS: Singapore (Singapore)

NCT04801966
PHASE NULL

Safety and Oversight of the Individually Tailored Treatment Approach: A Novel Pilot Study

TARGETS
 CDK4, CDK6, PI3K-alpha, PD-L1, MEK, PARP, PD-1, BRAF

LOCATIONS: Melbourne (Australia)

NCT02693535
PHASE 2

TAPUR: Testing the Use of Food and Drug Administration (FDA) Approved Drugs That Target a Specific Abnormality in a Tumor Gene in People With Advanced Stage Cancer

TARGETS
 ALK, ROS1, AXL, TRKA, MET, TRKC, CDK4, CDK6, FLT3, VEGFRs, CSF1R, KIT, RET, mTOR, ERBB2, MEK, BRAF, PARP, PD-1, CTLA-4, EGFR, ERBB4

LOCATIONS: Hawaii, Washington, Oregon, California

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Julie Tse, M.D. | 22 December 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1521387-01

CLINICAL TRIALS
GENE
CCNE1
RATIONALE

Strong preclinical and clinical data suggest that CCNE1 amplification may predict sensitivity to

WEE1 inhibitors.

ALTERATION

amplification

NCT04768868
PHASE 1

The Safety and Pharmacokinetics Preliminary Efficacy of IMP7068 in Patients With Advanced Solid Tumors

TARGETS
WEE1
LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Shanghai (China), Wuhan (China), Beijing (China), Chengdu (China), Kansas, Texas

NCT05128825
PHASE 2

A Study of ZN-c3 in Subjects With Malignant Tumors

TARGETS
WEE1
LOCATIONS: Nevada, Colorado, Texas, Ohio, Pennsylvania, Maryland, Virginia

NCT03968653
PHASE 1

Study of Oral Debio 0123 in Combination With Carboplatin in Participants With Advanced Solid Tumors

TARGETS
WEE1
LOCATIONS: Groningen (Netherlands), Nijmegen (Netherlands), Leiden (Netherlands), Barcelona (Spain)

NCT05109975
PHASE 1

A Study to Evaluate Safety and Preliminary Anti-tumor Activity of Debio 0123 as Monotherapy in Adult Participants With Advanced Solid Tumors

TARGETS
WEE1
LOCATIONS: Bellinzona (Switzerland), Zürich (Switzerland), Michigan, Texas

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Julie Tse, M.D. | 22 December 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1521387-01

APPENDIX
Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

BRCA1

E1849Q

CD22

amplification

CEBPA

amplification

CSF3R

H54D

EP300

Q950P

GABRA6

T414I

HSD3B1

G90S

IDH2

M293T

NF1

M1461>IR

NF2

E106K

POLE

R260*

SMO

A729V

SOC1

A76G and P50L

STAG2

E276K

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Julie Tse, M.D. | 22 December 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1521387-01

APPENDIX

Genes assayed in FoundationOne®Liquid CDx

FoundationOne Liquid CDx interrogates 324 genes, including 309 genes with complete exonic (coding) coverage and 15 genes with only select non-coding coverage (indicated with an *); 75 genes (indicated in bold) are captured with increased sensitivity and have complete exonic (coding) coverage unless otherwise noted.

ABL1 Exons 4-9	ACVR1B	AKT1 Exon 3	AKT2	AKT3	ALK Exons 20-29, Introns 18, 19	ALOX12B	AMER1 (FAM123B or WTX)	APC
AR	ARAF Exons 4, 5, 7, 11, 13, 15, 16	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX	AURKA
AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2	BCL6
BCOR	BCORL1	BCR* Introns 8, 13, 14	BRAF Exons 11-18, Introns 7-10	BRCA1 Introns 2, 7, 8, 12, 16, 19, 20	BRCA2 Intron 2	BRD4	BRIP1	BTG1
BTG2	BTK Exons 2, 15	CALR	CARD11	CASP8	CBFB	CBL	CCND1	CCND2
CCND3	CCNE1	CD22	CD70	CD74* Introns 6-8	CD79A	CD79B	CD274 (PD-L1)	CDC73
CDH1	CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B
CDKN2C	CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R
CTCF	CTNNA1	CTNNB1 Exon 3	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1
DDR2 Exons 5, 17, 18	DIS3	DNMT3A	DOT1L	EED	EGFR Introns 7, 15, 24-27	EMSY (C11orf30)	EP300	EPHA3
EPHB1	EPHB4	ERBB2	ERBB3 Exons 3, 6, 7, 8, 10, 12, 20, 21, 23, 24, 25	ERBB4	ERCC4	ERG	ERRF1	ESR1 Exons 4-8
ETV4* Intron 8	ETV5* Introns 6, 7	ETV6* Introns 5, 6	EWSR1* Introns 7-13	EZH2 Exons 4, 16, 17, 18	EZR* Introns 9-11	FANCA	FANCC	FANCG
FANCL	FAS	FBXW7	FGF10	FGF12	FGF14	FGF19	FGF23	FGF3
FGF4	FGF6	FGFR1 Introns 1, 5, Intron 17	FGFR2 Intron 1, Intron 17	FGFR3 Exons 7, 9 (alternative designation exon 10), 14, 18, Intron 17	FGFR4	FH	FLCN	FLT1
FLT3 Exons 14, 15, 20	FOXL2	FUBP1	GABRA6	GATA3	GATA4	GATA6	GID4 (C17orf39)	GNA11 Exons 4, 5
GNA13	GNAQ Exons 4, 5	GNAS Exons 1, 8	GRM3	GSK3B	H3-3A (H3F3A)	HDAC1	HGF	HNFI1A
HRAS Exons 2, 3	HSD3B1	ID3	IDH1 Exon 4	IDH2 Exon 4	IGF1R	IKBKE	IKZF1	INPP4B
IRF2	IRF4	IRS2	JAK1	JAK2 Exon 14	JAK3 Exons 5, 11, 12, 13, 15, 16	JUN	KDM5A	KDM5C
KDM6A	KDR	KEAP1	KEL	KIT Exons 8, 9, 11, 12, 13, 17, Intron 16	KLHL6	KMT2A (MLL) Introns 6, 8-11, Intron 7	KMT2D (MLL2)	KRAS

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1521387-01

APPENDIX

Genes assayed in FoundationOne® Liquid CDx

FoundationOne Liquid CDx interrogates 324 genes, including 309 genes with complete exonic (coding) coverage and 15 genes with only select non-coding coverage (indicated with an *); 75 genes (indicated in bold) are captured with increased sensitivity and have complete exonic (coding) coverage unless otherwise noted.

LTK	LYN	MAF	MAP2K1 (MEK1) Exons 2, 3	MAP2K2 (MEK2) Exons 2-4, 6, 7	MAP2K4	MAP3K1	MAP3K13	MAPK1
MCL1	MDM2	MDM4	MED12	MEF2B	MEN1	MERTK	MET	MITF
MKNK1	MLH1	MPL Exon 10	MRE11 (MRE11A)	MSH2 Intron 5	MSH3	MSH6	MST1R	MTAP
MTOR Exons 19, 30, 39, 40, 43-45, 47, 48, 53, 56	MUTYH	MYB* Intron 14	MYC Intron 1	MYCL (MYCL1)	MYCN	MYD88 Exon 4	NBN	NF1
NF2	NFE2L2	NFKBIA	NKX2-1	NOTCH1	NOTCH2 Intron 26	NOTCH3	NPM1 Exons 4-6, 8, 10	NRAS Exons 2, 3
NSD2 (WHSC1 or MMSET)	NSD3 (WHSC1L1)	NT5C2	NTRK1 Exons 14, 15, Introns 8-11	NTRK2 Intron 12	NTRK3 Exons 16, 17	NUTM1* Intron 1	P2RY8	PALB2
PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)	PDGFRA Exons 12, 18, Introns 7, 9, 11	PDGFRB Exons 12-21, 23
PDK1	PIK3C2B	PIK3C2G	PIK3CA Exons 2, 3, 5-8, 10, 14, 19, 21 (Coding Exons 1, 2, 4-7, 9, 13, 18, 20)	PIK3CB	PIK3R1	PIM1	PMS2	POLD1
POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI	PRKN (PARK2)	PTCH1
PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51	RAD51B	RAD51C
RAD51D	RAD52	RAD54L	RAF1 Exons 3, 4, 6, 7, 10, 14, 15, 17, Introns 4-8	RARA Intron 2	RB1	RBM10	REL	RET Introns 7, 8, Exons 11, 13-16, Introns 9-11
RICTOR	RNF43	ROS1 Exons 31, 36-38, 40, Introns 31-35	RPTOR	RSP02* Intron 1	SDC4* Intron 2	SDHA	SDHB	SDHC
SDHD	SETD2	SF3B1	SGK1	SLC34A2* Intron 4	SMAD2	SMAD4	SMARCA4	SMARCB1
SMO	SNCAIP	SOCS1	SOX2	SOX9	SPEN	SPOP	SRC	STAG2
STAT3	STK11	SUFU	SYK	TBX3	TEK	TENT5C (FAM46C)	TERC* ncRNA	TERT* Promoter
TET2	TGFBR2	TIPARP	TMPRSS2* Introns 1-3	TNFAIP3	TNFRSF14	TP53	TSC1	TSC2
TYRO3	U2AF1	VEGFA	VHL	WT1	XPO1	XRCC2	ZNF217	ZNF703

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS

Microsatellite (MS) status
Blood Tumor Mutational Burden (bTMB)
Tumor Fraction

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Julie Tse, M.D. | 22 December 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1521387-01

APPENDIX

About FoundationOne® Liquid CDx

FoundationOne Liquid CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Ciplastraat 3, 2440 Geel, Belgium. The CE-IVD regulatory status of FoundationOne Liquid CDx is applicable in countries that accept and/or recognize the CE mark.



ABOUT FOUNDATIONONE LIQUID CDx

FoundationOne Liquid CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne Liquid CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details.

INTENDED USE

FoundationOne Liquid CDx is a next generation sequencing based *in vitro* diagnostic device that analyzes 324 genes. Substitutions and insertion and deletion alterations (indels) are reported in 311 genes, copy number alterations (CNAs) are reported in 310 genes, and gene rearrangements are reported in 324 genes. The test also detects the genomic signatures blood tumor mutational burden (bTMB), microsatellite instability (MSI), and tumor fraction. FoundationOne Liquid CDx utilizes circulating cell-free DNA (cfDNA) isolated from plasma derived from the anti-coagulated peripheral whole blood of cancer patients. The test is intended to be used as a companion diagnostic to identify patients who may benefit from treatment with targeted therapies in accordance with the approved therapeutic product labeling. Additionally, FoundationOne Liquid CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with malignant neoplasms.

TEST PRINCIPLES

The FoundationOne Liquid CDx assay is performed exclusively as a laboratory service using circulating cell-free DNA (cfDNA) isolated from plasma derived from anti-coagulated peripheral whole blood from patients with solid malignant neoplasms. The assay employs a single DNA extraction method to obtain cfDNA from plasma from whole blood. Extracted

cfDNA undergoes whole-genome shotgun library construction and hybridization-based capture of 324 cancer-related genes including coding exons and select introns of 309 genes, as well as only select intronic regions or non-coding regions of 15 genes. Hybrid-capture selected libraries are sequenced with deep coverage using the NovaSeq® 6000 platform. Sequence data are processed using a customized analysis pipeline designed to accurately detect genomic alterations, including base substitutions, indels, select copy number variants, and select genomic rearrangements. Substitutions and insertion and deletion alterations (indels) are reported in 311 genes, copy number alterations (CNAs) are reported in 310 genes, and gene rearrangements are reported in 324 genes. The assay also reports tumor fraction, and genomic signatures including MSI and bTMB. A subset of targeted regions in 75 genes is baited for increased sensitivity.

THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. *Note:* A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

QUALIFIED ALTERATION CALLS (EQUIVOCAL)

All equivocal calls, regardless of alteration type, imply that there is adequate evidence to call the alteration with confidence. However, the repeatability of equivocal calls may be lower than non-equivocal calls.

RANKING OF THERAPIES AND CLINICAL TRIALS

Ranking of Therapies in Summary Table

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

Ranking of Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

LIMITATIONS

1. For *in vitro* diagnostic use.
2. For prescription use only. This test must be ordered by a qualified medical professional in accordance with clinical laboratory regulations.
3. A negative result does not rule out the presence of a mutation below the limits of detection of the assay. Patients for whom no companion diagnostic alterations are detected should be considered for confirmation with an appropriately validated tumor tissue test, if available.
4. The FoundationOne Liquid CDx assay does not detect heterozygous deletions.
5. The test is not intended to provide information on cancer predisposition.
6. Performance has not been validated for cfDNA input below the specified minimum input.
7. Tissue TMB and blood TMB (bTMB) are estimated from the number of synonymous and nonsynonymous single-nucleotide variants (SNVs) and insertions and deletions (indels) per area of coding genome sampled, after the removal of known and likely oncogenic driver events and germline SNPs. Tissue TMB is calculated based on variants with an allele frequency of $\geq 5\%$, and bTMB is calculated based on variants with an allele frequency of $\geq 0.5\%$.
8. Tumor fraction is the percentage of circulating tumor DNA (ctDNA) present in a cell-free DNA (cfDNA) sample. The tumor fraction estimate is computationally derived from the observed level of aneuploidy in the sample. Tumor fraction is considered elevated when ctDNA levels are high enough that aneuploidy can be detected and is significantly distinct from that typically found in non-tumor samples.
9. Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the tumor genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor. The MSI algorithm is based on genome wide analysis of 1765 microsatellite loci and not based on the 5 or 7 MSI loci described in current clinical practice guidelines for solid tissue testing.
10. Genomic findings from circulating cell-free DNA (cfDNA) may originate from circulating tumor DNA fragments, germline alterations, or non-tumor somatic alterations, such as clonal hematopoiesis of indeterminate potential (CHIP). Genes with alterations that may be derived from CHIP include, but are not limited to: *ASXL1*, *ATM*, *CBL*, *CHEK2*, *DNMT3A*, *JAK2*,

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1521387-01

APPENDIX
About FoundationOne® Liquid CDx

KMT2D (MLL2), MPL, MYD88, SF3B1, TET2, TP53, and U2AF1.

11. Alterations reported may include somatic (not inherited) or germline (inherited) alterations; however, the test does not distinguish between germline and somatic alterations. If a reported alteration is suspected to be germline, confirmatory testing should be considered in the appropriate clinical context.
12. The test is not intended to replace germline testing or to provide information about cancer predisposition.

REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic, nontargeted treatment, germline, or clonal hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >30%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM, BAP1, BRCA1, BRCA2, BRIP1, CHEK2, FH, FLCN, MLH1, MSH2, MSH6, MUTYH, PALB2, PMS2, POLE, RAD51C, RAD51D, RET, SDHA, SDHB, SDHC, SDHD, TSC2, and VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's

tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1, ATM, CBL, CHEK2, DNMT3A, IDH2, JAK2, KMT2D (MLL2), MPL, MYD88, SF3B1, TET2, and U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) (www.nccn.org). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2022. All rights reserved. To view the most recent and complete version of the guidelines, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

This report makes no promises or guarantees that a particular drug will be effective in the treatment of

disease in any patient. This report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne Liquid CDx.

TREATMENT DECISIONS ARE THE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this test or the information contained in this report.

Certain sample of variant characteristics may result in reduced sensitivity. These include: low sample quality, deletions and insertions >40bp, or repetitive/high homology sequences. FoundationOne Liquid CDx is performed using cell-free DNA, and as such germline events may not be reported.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Julie Tse, M.D. | 22 December 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1521387-01

APPENDIX

About FoundationOne®Liquid CDx

SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
Muts/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

REFERENCE SEQUENCE INFORMATION

Sequence data is mapped to the human genome, Genome Reference Consortium Human Build 37 (GRCh37), also known as hg19.

MR Suite Version (RG) 7.4.0

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Julie Tse, M.D. | 22 December 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1521387-01

APPENDIX
References

1. Gandara DR, et al. *Nat. Med.* (2018) PMID: 30082870
2. Wang Z, et al. *JAMA Oncol* (2019) PMID: 30816954
3. Sturgill EG, et al. *Oncologist* (2022) PMID: 35274716
4. Aggarwal C, et al. *Clin. Cancer Res.* (2020) PMID: 32102950
5. Schenker et al., 2022; AACR Abstract CT022
6. Saori et al., 2021; ESMO Abstract 80P
7. Chen EX, et al. *JAMA Oncol* (2020) PMID: 32379280
8. Rizvi NA, et al. *JAMA Oncol* (2020) PMID: 32271377
9. Si H, et al. *Clin Cancer Res* (2021) PMID: 33355200
10. Leighl NB, et al. *J Thorac Oncol* (2022) PMID: 34800700
11. Li et al., 2020; ASCO Abstract 6511
12. Nakamura H, et al. *Nat. Genet.* (2015) PMID: 26258846
13. Shao C, et al. *JAMA Netw Open* (2020) PMID: 33119110
14. Pfeifer GP, et al. *Mutat. Res.* (2005) PMID: 15748635
15. Hill VK, et al. *Annu Rev Genomics Hum Genet* (2013) PMID: 23875803
16. Pfeifer GP, et al. *Oncogene* (2002) PMID: 12379884
17. Rizvi NA, et al. *Science* (2015) PMID: 25765070
18. Johnson BE, et al. *Science* (2014) PMID: 24336570
19. Choi S, et al. *Neuro-oncology* (2018) PMID: 29452419
20. Cancer Genome Atlas Research Network, et al. *Nature* (2013) PMID: 23636398
21. Briggs S, et al. *J. Pathol.* (2013) PMID: 23447401
22. Heitz E, et al. *Curr. Opin. Genet. Dev.* (2014) PMID: 24583393
23. *Nature* (2012) PMID: 22810696
24. Roberts SA, et al. *Nat. Rev. Cancer* (2014) PMID: 25568919
25. Bronkhorst AJ, et al. *Biomol Detect Quantif* (2019) PMID: 30923679
26. Raja R, et al. *Clin. Cancer Res.* (2018) PMID: 30093454
27. Hrebien S, et al. *Ann. Oncol.* (2019) PMID: 30860573
28. Choudhury AD, et al. *JCI Insight* (2018) PMID: 30385733
29. Goodall J, et al. *Cancer Discov* (2017) PMID: 28450425
30. Goldberg SB, et al. *Clin. Cancer Res.* (2018) PMID: 29330207
31. Bettgeowda C, et al. *Sci Transl Med* (2014) PMID: 24553385
32. Lapin M, et al. *J Transl Med* (2018) PMID: 30400802
33. Shulman DS, et al. *Br. J. Cancer* (2018) PMID: 30131550
34. Stover DG, et al. *J. Clin. Oncol.* (2018) PMID: 29298117
35. Hemming ML, et al. *JCO Precis Oncol* (2019) PMID: 30793095
36. Egyud M, et al. *Ann. Thorac. Surg.* (2019) PMID: 31059681
37. Fan G, et al. *PLoS ONE* (2017) PMID: 28187169
38. Vu et al., 2020; DOI: 10.1200/PO.19.00204
39. Li G, et al. *J Gastrointest Oncol* (2019) PMID: 31602320
40. Zhang EW, et al. *Cancer* (2020) PMID: 32757294
41. Butler TM, et al. *Cold Spring Harb Mol Case Stud* (2019) PMID: 30833418
42. Basho RK, et al. *JAMA Oncol* (2017) PMID: 27893038
43. Hirai H, et al. *Mol. Cancer Ther.* (2010) PMID: 20571069
44. Zehir A, et al. *Nat. Med.* (2017) PMID: 28481359
45. Guedj N, et al. *J. Hepatol.* (2009) PMID: 19446907
46. Javle MM, et al. *J. Gastroenterol. Hepatol.* (2006) PMID: 16984600
47. Schomitz et al., 2007, 18161916
48. Liu AX, et al. *Cancer Res.* (1998) PMID: 9679957
49. Vivanco I, et al. *Nat. Rev. Cancer* (2002) PMID: 12094235
50. Chin YR, et al. *Cell Adh Migr* () PMID: 21519185
51. Cheng JQ, et al. *Proc. Natl. Acad. Sci. U.S.A.* (1992) PMID: 1409633
52. Thompson FH, et al. *Cancer Genet. Cytogenet.* (1996) PMID: 8646743
53. Altomare DA, et al. *Oncogene* (2005) PMID: 16288292
54. Int. J. Biol. Markers () PMID: 18409144
55. Scrima M, et al. *PLoS ONE* (2012) PMID: 22363436
56. Michels J, et al. *Oncogene* (2014) PMID: 24037533
57. Bryant HE, et al. *Nucleic Acids Res.* (2006) PMID: 16556909
58. Mateo J, et al. *N. Engl. J. Med.* (2015) PMID: 26510020
59. Mateo J, et al. *Lancet Oncol.* (2019) PMID: 31806540
60. Abida W, et al. *Clin. Cancer Res.* (2020) PMID: 32086346
61. Ma et al., 2022; ASCO GI Abstract 563
62. Dhawan et al., 2020; ASCO Abstract 3513
63. Papageorgiou GI, et al. *Front Oncol* (2021) PMID: 35004311
64. Olson D, et al. *Clin Genitourin Cancer* (2016) PMID: 27079472
65. Swisher EM, et al. *Nat Commun* (2021) PMID: 33941784
66. Zhang W, et al. *Oncologist* (2020) PMID: 32045060
67. O'Carrigan et al., 2016; ASCO Abstract 2504
68. Yap TA, et al. *Cancer Discov* (2021) PMID: 32988960
69. Kwon M, et al. *J Immunother Cancer* (2022) PMID: 35790315
70. Menezes DL, et al. *Mol. Cancer Res.* (2015) PMID: 25232030
71. Vendetti FP, et al. *Oncotarget* (2015) PMID: 26517239
72. Min A, et al. *Mol. Cancer Ther.* (2017) PMID: 28138034
73. Kwok M, et al. *Blood* (2016) PMID: 26563132
74. Riabinska A, et al. *Sci Transl Med* (2013) PMID: 23761041
75. Nguyen B, et al. *Cell* (2022) PMID: 35120664
76. Shiloh Y, et al. *Nat. Rev. Mol. Cell Biol.* (2013) PMID: 23847781
77. Cremona CA, et al. *Oncogene* (2014) PMID: 23851492
78. Jiang X, et al. *J. Biol. Chem.* (2006) PMID: 16603769
79. Fernandes N, et al. *J. Biol. Chem.* (2005) PMID: 15713674
80. Scott SP, et al. *Proc. Natl. Acad. Sci. U.S.A.* (2002) PMID: 11805335
81. van Os NJ, et al. *Clin Genet* (2016) PMID: 26662178
82. Rothblum-Oviatt C, et al. *Orphanet J Rare Dis* (2016) PMID: 27884168
83. Jaiswal S, et al. *N. Engl. J. Med.* (2014) PMID: 25426837
84. Genovese G, et al. *N. Engl. J. Med.* (2014) PMID: 25426838
85. Xie M, et al. *Nat. Med.* (2014) PMID: 25326804
86. Acuna-Hidalgo R, et al. *Am. J. Hum. Genet.* (2017) PMID: 28669404
87. Severson EA, et al. *Blood* (2018) PMID: 29678827
88. Fuster JJ, et al. *Circ. Res.* (2018) PMID: 29420212
89. Chabon JJ, et al. *Nature* (2020) PMID: 32269342
90. Razavi P, et al. *Nat. Med.* (2019) PMID: 31768066
91. Lin AB, et al. *Clin. Cancer Res.* (2017) PMID: 28331049
92. Chen X, et al. *Clin Cancer Res* (2018) PMID: 30181387
93. Mörry T, et al. *Int. J. Biochem. Cell Biol.* (2004) PMID: 15147722
94. Lee JM, et al. *Lancet Oncol.* (2018) PMID: 29361470
95. Lheureux S, et al. *Lancet* (2021) PMID: 33485453
96. Oza AM, et al. *Clin Cancer Res* (2020) PMID: 32611648
97. Fu et al., 2021; AACR abstract 974
98. Toledo LI, et al. *Nat. Struct. Mol. Biol.* (2011) PMID: 21552262
99. Buisson R, et al. *Mol. Cell* (2015) PMID: 26365377
100. Yang L, et al. *Oncotarget* (2015) PMID: 26204491
101. Kok YP, et al. *Oncogenesis* (2020) PMID: 33028815
102. Taylor-Harding B, et al. *Oncotarget* (2015) PMID: 25557169
103. Etemadmoghadam D, et al. *Clin. Cancer Res.* (2013) PMID: 24004674
104. Scaltriti M, et al. *Proc. Natl. Acad. Sci. U.S.A.* (2011) PMID: 21321214
105. Nanos-Webb A, et al. *Breast Cancer Res. Treat.* (2012) PMID: 21695458
106. Ma T, et al. *Mol. Cancer Ther.* (2013) PMID: 23686769
107. Narayan RR, et al. *Cancer* (2019) PMID: 30427539
108. Lowery MA, et al. *Clin. Cancer Res.* (2018) PMID: 29848569
109. Sanchez-Vega F, et al. *Cell* (2018) PMID: 29625050
110. Leung SY, et al. *Mod. Pathol.* (2006) PMID: 16575401
111. Lin L, et al. *Cancer Res.* (2000) PMID: 11156406
112. Mayr D, et al. *Am. J. Clin. Pathol.* (2006) PMID: 16753589
113. Nakayama N, et al. *Cancer* (2010) PMID: 20336784
114. Stamatakis M, et al. *World J Surg Oncol* (2010) PMID: 21176227
115. *Nature* (2015) PMID: 25631445
116. Yaeger R, et al. *Cancer Cell* (2018) PMID: 29316426
117. Cancer Genome Atlas Research Network, et al. *Nature* (2017) PMID: 28112728
118. Hernandez L, et al. *Cell Death Discov* (2015) PMID: 28179987
119. Rodríguez-Berriguete G, et al. *BMC Cancer* (2015) PMID: 26507126
120. Pingoud-Meier C, et al. *Clin. Cancer Res.* (2003) PMID: 14695141
121. Galluzzi L, et al. *Immunity* (2016) PMID: 26885855
122. Tummers B, et al. *Immunol. Rev.* (2017) PMID: 28462525
123. Crowder RN, et al. *Exp. Oncol.* (2012) PMID: 23070000
124. Liu J, et al. *Mol. Cancer* (2011) PMID: 21801448
125. Graf RP, et al. *Curr. Mol. Med.* (2014) PMID: 24467204
126. Barbero S, et al. *Cancer Res.* (2009) PMID: 19383910
127. Zhang L, et al. *Cell. Signal.* (2015) PMID: 25446254
128. Su H, et al. *Science* (2005) PMID: 15746428
129. Henry CM, et al. *Mol. Cell* (2017) PMID: 28212752
130. Kober AM, et al. *Cell Death Dis* (2011) PMID: 21975294
131. Sun BK, et al. *Oncol. Rep.* (2011) PMID: 21152872
132. Soung YH, et al. *Cancer Res.* (2005) PMID: 15705878
133. Soung YH, et al. *Oncogene* (2005) PMID: 15531912
134. Mandruzzato S, et al. *J. Exp. Med.* (1997) PMID: 9271594
135. Kim HS, et al. *Gastroenterology* (2003) PMID: 12949717
136. Li C, et al. *Mol Oncol* (2014) PMID: 24816188
137. Ando M, et al. *Cancer Sci.* (2013) PMID: 23659359
138. Cerami E, et al. *Cancer Discov* (2012) PMID: 22588877
139. Gao J, et al. *Sci Signal* (2013) PMID: 23550210
140. Trowbridge JJ, et al. *Nat. Genet.* (2011) PMID: 22200773
141. Prog Mol Biol Transl Sci () PMID: 21507354
142. Yang J, et al. *Mol Med Rep* () PMID: 21887466
143. Vallböhmer D, et al. *Clin Lung Cancer* (2006) PMID: 16870044
144. Daskalos A, et al. *Cancer* (2011) PMID: 21351083
145. Fabbri M, et al. *Proc. Natl. Acad. Sci. U.S.A.* (2007) PMID: 17890317
146. Gao Q, et al. *Proc. Natl. Acad. Sci. U.S.A.* (2011) PMID: 22011581
147. Kim MS, et al. *APMIS* (2013) PMID: 23031157
148. Chen ZX, et al. *J. Cell. Biochem.* (2005) PMID: 15861382
149. Guo X, et al. *Nature* (2015) PMID: 25383530
150. Sandoval JE, et al. *J. Biol. Chem.* (2019) PMID: 30705090
151. Zhang ZM, et al. *Nature* (2018) PMID: 29414941
152. Hematology Am Soc Hematol Educ Program (2018) PMID: 30504320
153. Owonikoko et al., 2016; ESMO Abstract 14230
154. Hook KE, et al. *Mol. Cancer Ther.* (2012) PMID: 22222631

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # **ORD-1521387-01**
APPENDIX
References

155. Gong X, et al. Cancer Discov (2019) PMID: 30373917
156. Oser MG, et al. Cancer Discov (2019) PMID: 30373918
157. Yang W, et al. Kaohsiung J Med Sci (2022) PMID: 34741392
158. Beltran H, et al. Clin. Cancer Res. (2019) PMID: 30232224
159. Allaman-Pillet N, et al. Ophthalmic Genet. () PMID: 21955141
160. Viatour P, et al. J. Exp. Med. (2011) PMID: 21875955
161. Burkhardt DL, et al. Nat. Rev. Cancer (2008) PMID: 18650841
162. Knudsen ES, et al. Nat. Rev. Cancer (2008) PMID: 19143056
163. Berge EO, et al. Mol. Cancer (2010) PMID: 20594292
164. Giacinti C, et al. Oncogene (2006) PMID: 16936740
165. Otterson GA, et al. Proc. Natl. Acad. Sci. U.S.A. (1997) PMID: 9342358
166. Otterson GA, et al. Am. J. Hum. Genet. (1999) PMID: 10486322
167. Qin XQ, et al. Genes Dev. (1992) PMID: 1534305
168. Rubin SM, et al. Cell (2005) PMID: 16360038
169. Sun H, et al. Mol. Cell. Biol. (2006) PMID: 16449662
170. Chen Z, et al. Hum. Mutat. (2014) PMID: 24282159
171. Yun J, et al. Int J Ophthalmol (2011) PMID: 22553621
172. Houston SK, et al. Int Ophthalmol Clin (2011) PMID: 21139478
173. Ng AK, et al. Semin Radiat Oncol (2010) PMID: 19959033
174. Hirai H, et al. Cancer Biol. Ther. (2010) PMID: 20107315
175. Bridges KA, et al. Clin. Cancer Res. (2011) PMID: 21799033
176. Rajeshkumar NV, et al. Clin. Cancer Res. (2011) PMID: 21389100
177. Osman AA, et al. Mol. Cancer Ther. (2015) PMID: 25504633
178. Xu L, et al. Mol. Cancer Ther. (2002) PMID: 12489850
179. Xu L, et al. Mol. Med. (2001) PMID: 11713371
180. Camp ER, et al. Cancer Gene Ther. (2013) PMID: 23470564
181. Kim SS, et al. Nanomedicine (2015) PMID: 25240597
182. Pirollo KF, et al. Mol. Ther. (2016) PMID: 27357628
183. Leijen S, et al. J. Clin. Oncol. (2016) PMID: 27601554
184. Moore et al., 2019; ASCO Abstract 5513
185. Leijen S, et al. J. Clin. Oncol. (2016) PMID: 27998224
186. Oza et al., 2015; ASCO Abstract 5506
187. Lee J, et al. Cancer Discov (2019) PMID: 31315834
188. Méndez E, et al. Clin. Cancer Res. (2018) PMID: 29535125
189. Seligmann JF, et al. J Clin Oncol (2021) PMID: 34538072
190. Gourley et al., 2016; ASCO Abstract 5571
191. Park H, et al. ESMO Open (2022) PMID: 36084396
192. Li M, et al. Nat. Genet. (2014) PMID: 24997986
193. Ross JS, et al. Oncologist (2014) PMID: 24563076
194. Jiao Y, et al. Nat. Genet. (2013) PMID: 24185509
195. Chan-On W, et al. Nat. Genet. (2013) PMID: 24185513
196. Borger DR, et al. Oncologist (2012) PMID: 22180306
197. Suto T, et al. J Surg Oncol (2000) PMID: 10738270
198. Ong CK, et al. Nat. Genet. (2012) PMID: 22561520
199. Javle M, et al. Cancer (2016) PMID: 27622582
200. Nault JC, et al. Semin. Liver Dis. (2011) PMID: 21538283
201. Moreno M, et al. Ann Hepatol () PMID: 16177659
202. Won HS, et al. BMC Cancer (2010) PMID: 20955617
203. Chaube A, et al. BMC Cancer (2006) PMID: 16686942
204. Chuang SC, et al. World J Surg (2004) PMID: 15573254
205. Ruzzenente A, et al. Ann. Surg. Oncol. (2016) PMID: 26717940
206. Xiaofang L, et al. World J Surg Oncol (2012) PMID: 22230750
207. Ajiki T, et al. Hepatogastroenterology () PMID: 8799388
208. J Surg Oncol (2006) PMID: 16724348
209. Guo R, et al. Hum. Pathol. (2014) PMID: 24746206
210. Boerner T, et al. Hepatology (2021) PMID: 33765338
211. Conci S, et al. Updates Surg (2020) PMID: 32020551
212. Simbolo M, et al. Sci Rep (2018) PMID: 29740198
213. Churi CR, et al. PLoS ONE (2014) PMID: 25536104
214. Lee CS, et al. Pathology (1995) PMID: 7567135
215. Ahrendt SA, et al. J Hepatobiliary Pancreat Surg (2000) PMID: 11180865
216. Brown CJ, et al. Nat. Rev. Cancer (2009) PMID: 19935675
217. Joerger AC, et al. Annu. Rev. Biochem. (2008) PMID: 18410249
218. Kato S, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) PMID: 12826609
219. Kamada R, et al. J. Biol. Chem. (2011) PMID: 20978130
220. Zerdoumi Y, et al. Hum. Mol. Genet. (2017) PMID: 28472496
221. Yamada H, et al. Carcinogenesis (2007) PMID: 17690113
222. Landrum MJ, et al. Nucleic Acids Res. (2018) PMID: 29165669
223. Bougeard G, et al. J. Clin. Oncol. (2015) PMID: 26014290
224. Sorrell AD, et al. Mol Diagn Ther (2013) PMID: 23355100
225. Nichols KE, et al. Cancer Epidemiol. Biomarkers Prev. (2001) PMID: 11219776
226. Kleihues P, et al. Am. J. Pathol. (1997) PMID: 9006316
227. Gonzalez KD, et al. J. Clin. Oncol. (2009) PMID: 19204208
228. Lalloo F, et al. Lancet (2003) PMID: 12672316
229. Mandelker D, et al. Ann. Oncol. (2019) PMID: 31050713

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.