

ABOUT THE TEST FoundationOne®CDx is a next-generation sequencing (NGS) based assay that identifies genomic findings within hundreds of cancer-related genes.

PATIENT	DISEASE Brain glioblastoma (GBM)	PHYSICIAN	ORDERING PHYSICIAN Yeh, Yi-Chen	SPECIMEN	SPECIMEN SITE Brain
	NAME Fan, Kuang Hua		MEDICAL FACILITY Taipei Veterans General Hospital		SPECIMEN ID S111-09619A (PF22121)
	DATE OF BIRTH 11 August 1963		ADDITIONAL RECIPIENT None		SPECIMEN TYPE Slide Deck
	SEX Male		MEDICAL FACILITY ID 205872		DATE OF COLLECTION 09 March 2022
	MEDICAL RECORD # 48284577		PATHOLOGIST Not Provided		SPECIMEN RECEIVED 28 October 2022

Biomarker Findings

Microsatellite status - MS-Stable
Tumor Mutational Burden - 8 Muts/Mb

Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

KIT amplification

PDGFRA deletion exons 8-9, amplification, E229K - subclonal, R841K - subclonal[†]

MDM4 amplification

PIK3CA E545K

CDKN2A/B p16INK4a loss and p14ARF loss exons 2-3

CDKN2C L24fs*5

HGFE199K

PIK3C2B amplification

TERT promoter -146C>T

2 Disease relevant genes with no reportable alterations: **EGFR**, **IDH1**

[†] See About the Test in appendix for details.

Report Highlights

- Variants with **diagnostic implications** that may indicate a specific cancer type: **TERT** promoter -146C>T (p. 9)
- Targeted therapies with potential clinical benefit **approved in another tumor type**: Imatinib (p. 10), Nilotinib (p. 10), Sorafenib (p. 11), Sunitinib (p. 11)
- Evidence-matched **clinical trial options** based on this patient's genomic findings: (p. 12)
- Variants with **prognostic implications** for this tumor type that may impact treatment decisions: **TERT** promoter -146C>T (p. 9)

BIOMARKER FINDINGS

Microsatellite status - MS-Stable

Tumor Mutational Burden - 8 Muts/Mb

THERAPY AND CLINICAL TRIAL IMPLICATIONS

No therapies or clinical trials. See Biomarker Findings section

No therapies or clinical trials. See Biomarker Findings section

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 07 November 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

GENOMIC FINDINGS	THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)	THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)
KIT - amplification	none	Imatinib
		Nilotinib
		Sorafenib
		Sunitinib
10 Trials see p. 12		
PDGFRA - deletion exons 8-9, amplification, E229K - subclonal, R841K - subclonal	none	Imatinib
		Sorafenib
6 Trials see p. 15		
MDM4 - amplification	none	none
1 Trial see p. 14		
PIK3CA - E545K	none	none
10 Trials see p. 16		

GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

CDKN2A/B - p16INK4a loss and p14ARF loss	HGF - E199K	p. 8
exons 2-3	PIK3C2B - amplification	p. 9
CDKN2C - L24fs*5	TERT - promoter -146C>T	p. 9

NOTE Genomic alterations detected may be associated with activity of certain approved therapies; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and exhaustive. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies.

Therapies contained in this report may have been approved by the US FDA.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 07 November 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

BIOMARKER FINDINGS

BIOMARKER

Microsatellite status

RESULT

MS-Stable

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence, MSS tumors are significantly less likely than MSI-H tumors to respond to anti-PD-1 immune checkpoint inhibitors¹⁻³, including approved therapies nivolumab and pembrolizumab⁴. In a retrospective analysis of 361 patients with solid tumors treated with pembrolizumab, 3% were MSI-H and

experienced a significantly higher ORR compared with non-MSI-H cases (70% vs. 12%, $p=0.001$)⁵.

FREQUENCY & PROGNOSIS

Low-level MSI has been reported in 5-9% of glioblastoma (GBM) samples⁶⁻⁸. A large-scale study did not find high-level microsatellite instability (MSI-H) in any of 129 GBM samples⁶, although a small-scale study reported MSI-H in 4 of 15 pediatric GBMs and 1 of 12 adult GBMs⁹. The frequency of MSI has been reported to be increased in relapsed compared to primary GBM⁶, in GBMs with a previous lower grade astrocytoma⁷, and in giant cell GBM compared to classic GBM⁸.

FINDING SUMMARY

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor¹⁰. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2¹⁰⁻¹². This sample is microsatellite-stable (MSS), equivalent to the clinical definition of an MSS tumor: one with mutations in none of the tested microsatellite markers¹³⁻¹⁵. MSS status indicates MMR proficiency and typically correlates with intact expression of all MMR family proteins^{10,12,14-15}.

BIOMARKER

Tumor Mutational Burden

RESULT

8 Muts/Mb

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1¹⁶⁻¹⁸, anti-PD-1 therapies¹⁶⁻¹⁹, and combination nivolumab and ipilimumab²⁰⁻²⁵. In glioma, a lack of association between TMB and clinical benefit from immune checkpoint inhibitors has been reported^{16,26-27}. However, multiple case studies have reported that patients with ultramutated gliomas driven by POLE mutations

have benefited from treatment with anti-PD-1²⁸⁻²⁹ or anti-PD-L1³⁰ therapies. Therefore, although increased TMB alone may not be a strong biomarker for PD-1 or PD-L1 inhibitors in this cancer type, these agents may have efficacy for patients with glioma harboring both high TMB and POLE mutation.

FREQUENCY & PROGNOSIS

Glioblastoma (GBM) harbors a median TMB of 2.7 mutations per megabase (mut/Mb), and 4.2% of cases have high TMB (>20 muts/Mb)³¹. For pediatric patients, high TMB has been reported in a subset of high-grade gliomas, frequently in association with mutations in mismatch repair or proofreading genes and in TP53, whereas BRAF alterations or other oncogene fusions were observed more frequently in brain tumors harboring low TMB³²⁻³³. Increased TMB has been reported to correlate with higher tumor grade in glioma³⁴ and glioblastoma (GBM) tissue samples with biallelic mismatch repair deficiency

(bMMRD)²⁸, as well as with shorter OS of patients with diffuse glioma³⁵.

FINDING SUMMARY

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma³⁶⁻³⁷ and cigarette smoke in lung cancer³⁸⁻³⁹, treatment with temozolomide-based chemotherapy in glioma⁴⁰⁻⁴¹, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes⁴²⁻⁴⁶, and microsatellite instability (MSI)^{42,45-46}. This sample harbors a TMB below levels that would be predicted to be associated with sensitivity to PD-1- or PD-L1-targeting immune checkpoint inhibitors, alone or in combination with other agents^{16,26-30}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 07 November 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

GENOMIC FINDINGS

GENE
KIT

ALTERATION
amplification

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence, primarily in gastrointestinal stromal tumor (GIST), melanoma, AML, and systemic mastocytosis, KIT activating alterations are associated with sensitivity to TKIs including imatinib, sunitinib, sorafenib, dasatinib, nilotinib, pazopanib, regorafenib, ponatinib, midostaurin, apatinib, avapritinib, and ripretinib⁴⁷⁻⁵⁵. The use of mTOR inhibitors as an alternative therapeutic strategy has demonstrated limited success in KIT-mutated, imatinib-resistant melanoma, with 1 PR and 3 SD observed for 4 patients treated with everolimus⁵⁶. However, no

responses were observed for 10 patients with mastocytosis following everolimus monotherapy, with 8/10 patients harboring the KIT D816V mutation⁵⁷. The role of KIT amplification as a biomarker for response to mTOR inhibitors has not been investigated (PubMed, Mar 2022). Clinical benefit has been observed for patients with KIT amplified or overexpressing tumors following treatment with imatinib⁵⁸⁻⁶⁸, nilotinib⁶⁹, sorafenib⁷⁰⁻⁷³, and sunitinib⁷⁴⁻⁷⁵, suggesting that KIT amplification may be sensitive to these inhibitors. However, evidence demonstrating clinical benefit for regorafenib, dasatinib, pazopanib, or ponatinib in the context of KIT amplified or overexpressing tumors is limited.

FREQUENCY & PROGNOSIS

In the TCGA datasets, KIT amplification has been reported in 2.5% of lower grade gliomas (grades 2 and 3)⁷⁶ and 9.2% of glioblastomas (Grade 4 astrocytoma)⁷⁷. KIT amplification has been variously reported in 4-47% of glioblastomas in the

scientific literature⁷⁸⁻⁸⁰. Amplification of KIT has been strongly correlated with the presence of KDR and/or PDGFRA amplification in glioblastoma^{79,81-82}. One study found no correlation between KIT amplification and overall survival in patients with glioblastoma, while a separate study reported that overexpression of KIT was associated with tumor grade and shorter survival in patients with malignant glioma^{78,83}.

FINDING SUMMARY

KIT (also called c-KIT) encodes a cell surface tyrosine kinase receptor that, upon ligand binding and dimerization, activates the PI3K-AKT and RAS-MAPK signaling pathways⁸⁴. KIT aberrations, including point mutations, translocations, amplification, and overexpression, have been associated with various malignancies, and KIT is considered an oncoprotein⁸⁵. KIT has been reported to be amplified in cancer⁸⁶ and may be biologically relevant in this context⁸⁷⁻⁸⁸.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 07 November 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

GENOMIC FINDINGS

GENE
PDGFRA

ALTERATION

deletion exons 8-9, amplification, E229K - subclonal, R841K - subclonal

TRANSCRIPT ID

NM_006206.4, NM_006206.4

CODING SEQUENCE EFFECT

685G>A, 2522G>A

VARIANT CHROMOSOMAL POSITION

chr4:55131142, chr4:55152090

VARIANT ALLELE FREQUENCY (% VAF)

0.58%, 1.3%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of extensive clinical evidence in solid tumors and hematologic cancers, PDGFRA activating alterations are associated with sensitivity to imatinib⁸⁹⁻¹²⁶. Sorafenib has shown clinical and preclinical activity against the FIP1L1-PDGFRA fusion in chronic eosinophilic leukemia (CEL) and mutations associated with clinical resistance to imatinib and sunitinib in both CEL and gastrointestinal stromal tumor (GIST)¹²⁷⁻¹³². Complete responses to nilotinib have been reported in patients with CEL or hypereosinophilic syndrome with FIP1L1-PDGFRA or activating mutations^{105,133-134}; preclinical evidence has reported efficacy of nilotinib in the context of PDGFRA mutations associated with GIST¹³⁵⁻¹³⁶. Patients with GIST harboring PDGFRA activating mutations have been reported to derive clinical

benefit from treatment with sunitinib¹³⁷ or regorafenib¹³⁸⁻¹³⁹. Preclinical studies have reported sensitivity of activating PDGFRA mutations and FIP1L1-PDGFRA fusion to dasatinib^{129,135}. PDGFRA D842 mutations were reported to be sensitive to avapritinib in clinical¹⁴⁷ and preclinical¹⁴⁷ studies of GIST, and demonstrated sensitivity to ripretinib for 1 patient¹⁴⁰.

FREQUENCY & PROGNOSIS

PDGFRA amplification has been suggested to be more common in higher grade astrocytomas than in lower grade astrocytomas; studies have reported PDGFRA amplification in 16.3% (27/166) of Grade 2 astrocytomas and in 23.6% (91/386) of Grade 3 and 4 astrocytomas analyzed^{81,141-142}. PDGFRA amplification has been reported in 5.2-33% of glioblastoma cases^{77-79,141,143-144}. PDGFRA mutation has been identified in 5.6% of Grade 3 and 5.4% of Grade 4 astrocytomas, 2.4% of Grade 3 oligodendrogliomas, and 12% (3/25) of gliosarcomas analyzed in COSMIC (Feb 2022)¹⁴⁵. PDGFRA mutations have been reported in 0-5% of lower grade glioma and glioblastoma samples^{77,146-152}. Ceccarelli et al., 2016; 26824661, Cancer Genome Atlas Research Network., 2015; 26061751, cBio-Johnson et al., 2014; 24336570, cBio-Thomas et al., 2017; 28472509, cBio-Jones et al., 2013; 23817572). A retrospective analysis of TCGA glioma samples reported elevated expression of ERBB3 correlated with PDGFRA expression and co-expression of these genes was an indicator of poor prognosis in a GBM patient cohort¹⁵³. Amplification of PDGFRA has been associated with tumor grade and poor progression-free and overall survival in patients with

glioblastoma^{141,143-144}. In addition, PDGFRA amplification has been reported to occur in conjunction with IDH1 mutation in glioblastoma, and both alterations in the same tumor have been associated with poor patient prognosis¹⁴¹. Amplification of PDGFRA has also been strongly correlated with the presence of KDR and/or KIT amplification in glioblastomas, as well as with EGFR amplification^{79,81-82,154}.

FINDING SUMMARY

PDGFRA encodes platelet-derived growth factor receptor alpha (PDGFR-alpha), a tyrosine kinase receptor that, upon binding of cognate ligands (PDGFA or PDGFB), activates several signaling pathways, including PI3K and MAPK¹⁵⁵. PDGFR aberrations, including point mutations, translocations, amplification, and/or overexpression, have been associated with various malignancies⁸⁵. Amplification of PDGFRA, frequently occurring with amplification of the genes KDR and KIT, has been associated with increased PDGFRA expression^{80,156-158} and poor prognosis^{80,141,159-160} in some subtypes of glioma. The PDGFRA rearrangement in this tumor results in deletion of exons 8-9. The PDGFRA exon 8-9 deletion mutant was demonstrated to be active in the absence of ligand and has been characterized as transforming¹⁶¹⁻¹⁶². This mutant was sensitive to imatinib and the kinase inhibitor vatalanib¹⁶². Although alterations such as E229K and R841K seen here have not been fully characterized and are of unknown functional significance, similar alterations have been previously reported in the context of cancer, which may indicate biological relevance.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 07 November 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

GENOMIC FINDINGS

GENE

MDM4

ALTERATION
amplification

clinical development¹⁶⁷⁻¹⁷³ and have been shown to be effective against cancer cells in the presence of a wildtype p53 allele¹⁷⁴⁻¹⁷⁷. Additional therapeutic mechanisms that target MDM4 are also in preclinical development¹⁷²⁻¹⁷³.

MDM4 amplification in 4% (4/106) of GBMs and in 4% (1/27) of anaplastic oligodendrogliomas; MDM4 amplification was not detected in any of the 56 low-grade (Grade 1 or 2) gliomas investigated¹⁸³. The association of MDM4 amplification with tumor grade in astrocytomas is unclear¹⁷⁸⁻¹⁷⁹.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Small molecules targeting MDM4 or the MDM2-MDM4 complex, such as idasanutlin, have reported clinical benefit for patients with polycythemia vera¹⁶³⁻¹⁶⁴ and AML¹⁶⁵ and have shown anti-tumor activity in preclinical assays¹⁶⁶. Additional nutlins have been in preclinical and

FREQUENCY & PROGNOSIS

In the TCGA dataset, amplification of MDM4 was observed in 8.5% of glioblastoma cases⁷⁷. MDM4 amplification or amplification of the 1q32.1 chromosomal locus, which encompasses MDM4 and PIK3C2B, has been frequently reported, particularly in Grade 3 and 4 astrocytoma or glioblastoma multiforme (GBM) cases; one study reported MDM4 amplification in 27% (23/86) of astrocytoma samples¹⁷⁸⁻¹⁸³. A study also reported

FINDING SUMMARY

MDM4 acts as a negative regulator of p53, but a fraction of MDM4 localized to the mitochondria acts in concert with p53 to promote apoptosis¹⁸⁴. MDM4 has been reported to be amplified in cancer and therapies targeting MDM4 or MDM2 have been shown to increase levels of the tumor suppressor p53 in cancer cells^{86,177,185-186}.

GENE

PIK3CA

ALTERATION
E545K

TRANSCRIPT ID
NM_006218.2

CODING SEQUENCE EFFECT
1633G>A

VARIANT CHROMOSOMAL POSITION
chr3:178936091

VARIANT ALLELE FREQUENCY (% VAF)
34.0%

studies of copanlisib for patients with genomically unselected tumors reported 1 CR and 2 PRs (1 unconfirmed) among 16 total patients with PIK3CA-mutated solid tumors with or without PTEN alterations¹⁹¹⁻¹⁹². In the Phase 2 MATCH trial for patients with PIK3CA-mutated solid tumors, 28% (18/65) of patients experienced PFS lasting at least 6 months after treatment with taselisib; however, no ORs were observed in this study²⁰⁵. A separate Phase 1b study of taselisib in combination with the CDK4/6 inhibitor palbociclib for patients with PIK3CA-mutated solid tumors reported an ORR of 0% (n=12) and a DCR of 17% (2/12)²⁰⁶. In a Phase 1 trial of the dual PI3K/mTOR kinase inhibitor apitolisib, 79% (11/14) of patients with PIK3CA-mutated advanced solid tumors experienced disease control (3 PRs, 8 SDs)²⁰⁷. The PI3K inhibitor alpelisib is approved as a single agent for the treatment of patients with PIK3CA-related overgrowth spectrum (PROS)²⁰⁸, but has shown limited activity as monotherapy for PIK3CA-mutated solid tumors with a Phase 1a study reporting an ORR of 6.0% (8/134) and a DCR of 58% (78/134)²⁰⁹.

FREQUENCY & PROGNOSIS

PIK3CA mutations have been reported in 9% of glioblastoma (GBM) samples analyzed in the TCGA dataset⁷⁷, and other studies report the incidence of PIK3CA mutations in primary GBMs as 5-18%²¹⁰⁻²¹². One study detected PIK3CA mutation in 16% (36/232) of IDH-wildtype GBM samples analyzed²¹³. PIK3CA mutations have been reported

in 5-23% of high-grade gliomas (including glioblastomas, anaplastic astrocytomas, and anaplastic oligodendrogliomas)^{146,210-212,214}. While another study did not observe PIK3CA mutations in low-grade astrocytomas or in anaplastic astrocytomas, it did report high ERK and AKT activity²¹². One study found that PIK3CA mutation in glioblastoma (GBM) was associated with shorter median PFS in both a discovery cohort (6.9 vs. 12.4 months, HR=2.89, p=0.01) and in the TCGA cohort (6.1 vs. 9 months, p=0.008), but was not consistently associated with median OS²¹⁵. In a study of IDH-wildtype GBM, patients with alterations in PI3K class I genes (PIK3CA, PIK3R1, PIK3CG, and PIK3R2) had significantly longer OS (20.0 months altered vs. 16.9 months wildtype, HR=0.62, p=0.002) and PFS (11.0 months altered vs. 7.4 months wildtype, p=0.0043); patients with PIK3CA alterations experienced an improved OS but this association was not highly significant (20.0 months altered vs. 18.1 months wildtype, p=0.0407)²¹³.

FINDING SUMMARY

PIK3CA encodes p110-alpha, which is the catalytic subunit of phosphatidylinositol 3-kinase (PI3K). The PI3K pathway is involved in cell signaling that regulates a number of critical cellular functions, including cell growth, proliferation, differentiation, motility, and survival²¹⁶⁻²¹⁷. PIK3CA alterations that have been characterized as activating, such as observed here, are predicted to be oncogenic²¹⁸⁻²³⁹.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Clinical and preclinical data in various tumor types indicate that PIK3CA activating alterations may predict sensitivity to therapies targeting PI3K¹⁸⁷⁻¹⁹⁴, AKT¹⁹⁵⁻¹⁹⁶, or mTOR¹⁹⁷⁻²⁰⁴. The Phase 2 NCI-MATCH study of copanlisib for patients with refractory solid tumors harboring PIK3CA mutations with or without PTEN loss met its primary endpoint with an ORR of 16% (4/25 PRs); responses (PR or SD >6 months) were seen in patients with ameloblastoma, liposarcoma, and carcinomas of the endometrium, ovary, esophagus, lung, and prostate¹⁹⁴. However, the Phase 2 study of copanlisib for patients with endometrial carcinoma harboring PIK3CA hotspot mutations failed to report any objective responses (n=11)¹⁹³. Two other

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 07 November 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

GENOMIC FINDINGS

GENE

CDKN2A/B

ALTERATION

p16INK4a loss and p14ARF loss exons 2-3

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Preclinical data suggest that tumors with loss of p16INK4a function may be sensitive to CDK4/6 inhibitors, such as abemaciclib, ribociclib, and palbociclib²⁴⁰⁻²⁴³. Clinical data in mesothelioma, breast cancer, and uterine leiomyosarcoma indicate that CDKN2A loss may predict sensitivity to abemaciclib²⁴⁴ and palbociclib treatment²⁴⁵⁻²⁴⁶. However, multiple other clinical studies have shown no significant correlation between p16INK4a loss or inactivation and therapeutic benefit of these agents²⁴⁷⁻²⁵³; it is not known whether CDK4/6 inhibitors would be beneficial in this case. Although preclinical studies have suggested that loss of p14ARF function may be associated with reduced sensitivity to MDM2 inhibitors²⁵⁴⁻²⁵⁵, the clinical relevance of p14ARF as a predictive biomarker is not clear.

FREQUENCY & PROGNOSIS

Concurrent putative homozygous deletion of CDKN2A and CDKN2B has been reported in 35% of patients with gliomas²⁵⁶ and detected more

frequently in patients with glioblastoma multiforme (GBM; 58%)⁷⁷ than in those with lower grade gliomas (6%)²⁵⁷. In other studies, loss of CDKN2A/B by deletion has been reported in up to 78% of astrocytomas (including anaplastic astrocytomas and GBM)^{143,158,258}. A study found homozygous deletion of both p16INK4a and p14ARF in 26% (13/50) of glioblastomas (GBMs); 18% (9/50) of cases showed homozygous deletion of the p14ARF-encoding locus alone²⁵⁹. One study detected CDKN2A/B loss in 69% (161/232) and mutation in 2.6% (6/232) of IDH-wildtype GBM samples analyzed²¹³. Decreased p14ARF and p16INK4a expression levels were found to be tightly associated in a study of glioma samples²⁶⁰. Homozygous deletion of the genomic region including CDKN2A and CDKN2B has been found to be associated with poor prognosis in GBM and likely serves as an early event in GBM progression^{143,261}. In addition, expression of p16INK4a has been found to be lower in patients with high grade malignant gliomas compared to patients with low grade gliomas, and loss of p16INK4a expression has been associated with shorter overall survival in pilocytic astrocytomas²⁶²⁻²⁶³.

FINDING SUMMARY

CDKN2A encodes two different, unrelated tumor suppressor proteins, p16INK4a and p14ARF, whereas CDKN2B encodes the tumor suppressor p15INK4b²⁶⁴⁻²⁶⁵. Both p15INK4b and p16INK4a

bind to and inhibit CDK4 and CDK6, thereby maintaining the growth-suppressive activity of the Rb tumor suppressor; loss or inactivation of either p15INK4b or p16INK4a contributes to dysregulation of the CDK4/6-cyclin-Rb pathway and loss of cell cycle control²⁶⁶⁻²⁶⁷. The tumor suppressive functions of p14ARF involve stabilization and activation of p53, via a mechanism of MDM2 inhibition²⁶⁸⁻²⁶⁹. One or more alterations observed here are predicted to result in p16INK4a loss of function²⁷⁰⁻²⁹¹. One or more alterations seen here are predicted to result in p14ARF loss of function^{274,291-294}.

POTENTIAL GERMLINE IMPLICATIONS

Germline CDKN2A mutation is associated with melanoma-pancreatic cancer syndrome, a condition marked by increased risk of developing malignant melanoma and/or pancreatic cancer²⁹⁵. Mutation carriers within families may develop either or both types of cancer, and melanoma cases may be referred to as familial or hereditary melanoma²⁹⁶⁻²⁹⁷. CDKN2A is the most implicated gene in familial melanoma, with germline mutations present in 16% to 20% of familial melanoma cases²⁹⁸⁻³⁰⁰. CDKN2A alteration has also been implicated in familial melanoma-astrocytoma syndrome, an extremely rare tumor association characterized by dual predisposition to melanoma and nervous system tumors³⁰¹⁻³⁰³. In the appropriate clinical context, germline testing of CDKN2A is recommended.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 07 November 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

GENOMIC FINDINGS

GENE

CDKN2C

ALTERATION

L24fs*5

TRANSCRIPT ID

NM_001262.2

CODING SEQUENCE EFFECT

71_74delTGTT

VARIANT CHROMOSOMAL POSITION

chr1:51436107-51436111

VARIANT ALLELE FREQUENCY (% VAF)

45.6%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of preclinical studies, CDKN2C inactivation may promote sensitivity to CDK4/6 inhibitors^{242,304-306}, although this has not been demonstrated clinically.

FREQUENCY & PROGNOSIS

Although CDKN2C missense mutations are rare in the context of cancer, loss of p18INK4c expression, genomic loss, deletion, and promoter methylation have been frequently reported in Merkel cell carcinoma³⁰⁷⁻³⁰⁸, anaplastic meningioma³⁰⁹⁻³¹¹, uterine leiomyomata³¹², leiomyosarcoma³¹³, pituitary adenoma³¹⁴⁻³¹⁵, Hodgkin lymphoma³¹⁶, multiple myeloma³¹⁷⁻³¹⁹, hepatocellular carcinoma (HCC)³²⁰, adenoid cystic carcinoma of the salivary gland³²¹, secondary angiosarcoma³²², pancreatic carcinoma³²³⁻³²⁴, gastroenteropancreatic neuroendocrine tumors³²⁵, glioblastoma^{77,326-327}, Wilms tumors³²⁸, and mantle cell lymphoma (MCL)³²⁹⁻³³¹. CDKN2C alterations, including missense mutations, have also been reported in parathyroid adenomas³³²⁻³³⁴ and thyroid carcinomas (TCs)³³⁵⁻³³⁷; single nucleotide polymorphisms within the region encoding CDKN2C have been associated with increased risk of papillary TC³³⁸ and with increased tumor size in sporadic medullary TC³³⁹. CDKN2C alterations or loss of p18INK4c expression are negative

prognostic factors in multiple myeloma³¹⁷⁻³¹⁹, HCC³²⁰, MCL³²⁹, and acute myeloid leukemia (AML)³⁴⁰. However, p18INK4c has been reported to be overexpressed in mesothelioma³⁴¹, and nuclear p18INK4c expression has been reported to be a poor prognostic factor in oligodendroglioma³⁴².

FINDING SUMMARY

CDKN2C encodes p18, also known as INK4c, a member of a family of cyclin-dependent kinase 4 (CDK4) inhibitors. INK4 family members are commonly deleted in cancer, which results in unrestrained CDK4/6 activity and dysregulated cell cycle entry³⁴³. CDKN2C alterations that result in loss or disruption of the ankyrin repeats (amino acids 4-132), are predicted to be inactivating³⁴⁴⁻³⁴⁵, although some alterations seen in the context of cancer have not been directly functionally characterized.

GENE

HGF

ALTERATION

E199K

TRANSCRIPT ID

NM_000601.4

CODING SEQUENCE EFFECT

595G>A

VARIANT CHROMOSOMAL POSITION

chr7:81381466

VARIANT ALLELE FREQUENCY (% VAF)

32.1%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of several preclinical studies in different cancer types, high HGF gene expression may associate with sensitivity to MET-targeted therapies, such as the approved multikinase inhibitors crizotinib and cabozantinib³⁴⁶⁻³⁵⁰.

However, this hypothesis has not been extensively tested in clinical studies. Whereas patients with glioblastoma and high tumor HGF gene expression experienced longer survival and a higher objective response rate (5/14 vs. 0/16) on the MET-targeting antibody onartuzumab combined with the anti-VEGF antibody bevacizumab than with placebo plus bevacizumab³⁵¹, tumor HGF gene expression did not predict significant benefit from onartuzumab added to the EGFR-inhibitor erlotinib for patients with non-small cell lung cancer³⁵². Anti-HGF antibodies, such as ficlatuzumab, are also under clinical investigation³⁵³⁻³⁵⁴. It is unclear if missense mutations in HGF may lead to increased expression or activation, and therefore it is unknown whether these therapeutic approaches would be relevant.

FREQUENCY & PROGNOSIS

Characterized HGF mutations have rarely been reported in solid tumors³⁵⁵⁻³⁵⁶. HGF expression within tumor glioma cells is associated with high-

grade glioma and increased microvessels, and tumor-derived HGF expression has been shown to correlate with reduced survival time³⁵⁷. Elevated expression of HGF and MET mRNA have been reported in GBM³⁵⁸⁻³⁵⁹, and HGF expression in GBM models has been shown to be associated with responsiveness to MET inhibition³⁴⁶.

FINDING SUMMARY

HGF encodes hepatocyte growth factor, also known as scatter factor, an activating ligand of the receptor tyrosine kinase MET. Certain splice isoforms of HGF may also act as MET antagonists³⁶⁰⁻³⁶¹. HGF plays an important role in normal development, acting as a growth factor in a number of different tissues³⁶⁰⁻³⁶¹. HGF and its receptor, MET, have been implicated in growth, invasion, and metastasis of many solid tumors³⁶¹. Although alterations such as seen here have not been fully characterized and are of unknown functional significance, similar alterations have been previously reported in the context of cancer, which may indicate biological relevance.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 07 November 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

GENOMIC FINDINGS

GENE

PIK3C2B

ALTERATION
amplification

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no therapies known to effectively target mutations in PIK3C2B.

FREQUENCY & PROGNOSIS

Although PIK3C2B mutation has not been reported for many tumor types, microarray analysis of glioblastoma cell lines revealed that increased gene expression of PIK3C2B significantly correlated with insensitivity to erlotinib³⁶². In addition, 8% of 465 glioblastoma multiforme (GBM) tumor samples analyzed in one study demonstrated copy number amplification of PIK3C2B¹⁸¹. Although elevated expression of PIK3C2B has been observed in GBM, it is unclear if this plays any role in oncogenesis.

FINDING SUMMARY

PI3K signaling is implicated in the regulation of metabolic control, immunity, angiogenesis and cardiovascular homeostasis, and is one of the most frequently dysregulated pathways in cancer. In contrast to class I PI3Ks, including p110- α and p110- β , the functional roles of class II PI3Ks, encoded by PIK3C2A, PIK3C2B, and PIK3C2G, are not well understood³⁶³.

GENE

TERT

ALTERATION
promoter -146C>T

TRANSCRIPT ID
NM_198253.2

CODING SEQUENCE EFFECT
-146C>T

VARIANT CHROMOSOMAL POSITION
chr5:1295250

VARIANT ALLELE FREQUENCY (% VAF)
34.4%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Therapeutic options for targeting tumors with TERT mutations are limited, although a variety of approaches have been investigated, including immunotherapies using TERT as a tumor-associated antigen and antisense oligonucleotide- or peptide-based therapies. TERT peptide vaccines showed limited anticancer efficacy in clinical trials³⁶⁴; however, in one preclinical study, the combination of a TERT peptide vaccine and anti-CTLA-4 therapy suppressed tumor growth³⁶⁵. A Phase 2 study of the TERT inhibitor imetelstat for

patients with advanced non-small cell lung cancer reported no improvement in PFS or OS³⁶⁶.

FREQUENCY & PROGNOSIS

TERT promoter mutations have been reported in 51-59% of gliomas³⁶⁷⁻³⁶⁸, most frequently in glioblastoma (GBM, 54-84%), gliosarcoma (81%), oligodendroglioma (78%), and historically in lower grade astrocytomas (25-31%) but less frequently in ependymomas³⁶⁷⁻³⁷¹. In patients with glioblastoma (GBM), the prevalence of TERT promoter mutation is lower in pediatric primary GBM (11%) and adult secondary GBM (28%) compared with adult primary GBM (58-83%)^{367,369}. One study detected TERT promoter mutations in 78% (181/232) of IDH-wildtype GBM samples analyzed²¹³. TERT promoter mutation has been shown to be significantly associated with increased TERT gene expression in astrocytoma, oligodendroglioma, and GBM³⁷². TERT promoter mutations significantly associate with poor prognosis in patients with GBM, although this correlation may be due to the association with primary GBM as opposed to IDH-positive secondary GBM^{367,369,372-373}. In the context of IDH-wildtype glioma, TERT mutations are associated with reduced OS (NCCN CNS Cancers Guidelines, v1.2022).

FINDING SUMMARY

Telomerase reverse transcriptase (TERT, or hTERT) is a catalytic subunit of the telomerase complex, which is required to maintain appropriate chromosomal length³⁷⁴. Activation of TERT is a hallmark of cancer, being detected in up to 80-90% of malignancies and absent in quiescent cells³⁷⁵⁻³⁷⁷. Mutations within the promoter region of TERT that confer enhanced TERT promoter activity have been reported in two hotspots, located at -124 bp and -146 bp upstream of the transcriptional start site (also termed C228T and C250T, respectively)³⁷⁸⁻³⁸⁰, as well as tandem mutations at positions -124/-125 bp and -138/-139 bp³⁷⁸.

POTENTIAL DIAGNOSTIC IMPLICATIONS

TERT mutations are associated with 1p/19q co-deletion in oligodendrogliomas, and are highly recurrent in IDH/ATRX-wildtype glioblastoma (GBM) (NCCN CNS Cancers Guidelines, v1.2022)³⁸¹. The presence of EGFR gene amplification or TERT promoter mutations are indicative of diffuse astrocytic glioma with molecular features of glioblastoma, WHO grade 4 in IDH1/2-wildtype tumors (NCCN CNS Cancers Guidelines, v1.2022)³⁸².

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 07 November 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Imatinib

Assay findings association
KIT
amplification

PDGFRA
deletion exons 8-9, amplification,
E229K - subclonal, R841K -
subclonal

AREAS OF THERAPEUTIC USE

Imatinib targets the BCR-ABL fusion protein, PDGFR, and KIT. It is FDA approved for the treatment of KIT-positive gastrointestinal stromal tumors (GIST), Ph+ chronic myeloid leukemia (CML) and acute lymphoblastic leukemia (ALL), myelodysplastic syndrome/myeloproliferative syndrome (MDS/MPS), aggressive systemic mastocytosis without a D816V KIT mutation, hypereosinophilic syndrome and/or chronic eosinophilic leukemia, and dermatofibrosarcoma protuberans. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of strong clinical evidence, PDGFRA activating mutations^{91,96-97,101,126}, fusions^{90,94,100,102,110,113,115,119,122,383}, and expression⁹⁹ may predict sensitivity to imatinib. On the basis of clinical and preclinical data in KIT-mutated^{59-60,101,384}, KIT-amplified⁵⁸⁻⁶¹, or KIT-expressing tumors^{63-68,385-386}, KIT

activating alterations may confer sensitivity to imatinib. PDGFRA amplification may predict sensitivity to tyrosine kinase inhibitors such as imatinib; a patient with Merkel cell carcinoma expressing PDGFRA achieved a complete response to imatinib⁹⁹.

SUPPORTING DATA

In a clinical study where patients with recurrent glioblastoma were given imatinib, 2/24 patients achieved a PR, 10 patients reported SD, and median OS and PFS was observed to be 6.2 and 3 months, respectively³⁸⁷. However, other Phase 2 clinical trials of imatinib have reported no anti-tumor activity, with a study of 231 patients with glioblastoma reporting a radiographic response rate of only 3.4%^{68,388}. In another Phase 2 study, imatinib plus hydroxyurea was shown to be well tolerated among patients with recurrent/progressive low-grade glioma, but had negligible antitumor activity³⁸⁹.

Nilotinib

Assay findings association
KIT
amplification

AREAS OF THERAPEUTIC USE

Nilotinib targets tyrosine kinases such as ABL (including BCR-ABL), PDGFRs, KIT, CSF1R, DDR1, and DDR2. It is FDA approved to treat newly diagnosed pediatric or adult patients with Philadelphia chromosome-positive (Ph+) chronic myeloid leukemia (CML) in chronic phase, adults with Ph+ CML in chronic or accelerated phase with resistance or intolerance to prior therapy including imatinib, and pediatric patients with Ph+ CML in chronic phase with resistance or intolerance to prior tyrosine-kinase inhibitor therapy. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical and preclinical data in KIT-mutated^{69,390-393}, KIT-amplified⁶⁹, or KIT-expressing tumors³⁹⁴, KIT activating alterations may confer sensitivity to nilotinib.

SUPPORTING DATA

Clinical data on the efficacy of nilotinib for the treatment of CNS tumors are limited (PubMed, Jul 2022). Nilotinib

has been primarily investigated as a therapeutic option for the treatment of CML or gastrointestinal stromal tumors (GIST). In the context of CML, a Phase 3 clinical trial of Ph+ patients treated with imatinib or nilotinib (300 mg or 400 mg) reported progression-free survival (PFS) rates of 93% and 97-98% and overall survival (OS) rates of 93% and 94-97%, respectively, at 4 years³⁹⁵. For imatinib-resistant Japanese patients with CML, a Phase 2 trial reported a 47.8% major medical response rate to treatment with nilotinib at 12 months³⁹⁶. A Phase 3 clinical trial of single-agent nilotinib in 240 patients with advanced GIST who failed prior treatment with imatinib or sunitinib reported no significant difference in progression-free survival between nilotinib and the best supportive care, but did report increased overall survival for nilotinib-treated patients³⁹⁷. A Phase 2 trial has shown that nilotinib was well tolerated and suggested it may be particularly useful for treating patients with GIST harboring mutations in KIT exon 17³⁹⁸. Preclinical, cell-based assays have reported efficacy for nilotinib alone and in combination with additional therapies in the context of leiomyosarcoma and synovial sarcoma³⁹⁹.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 07 November 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Sorafenib

Assay findings association
KIT
amplification

PDGFRA
deletion exons 8-9, amplification,
E229K - subclonal, R841K -
subclonal

AREAS OF THERAPEUTIC USE

Sorafenib is a kinase inhibitor that targets the RAF kinases, KIT, FLT3, RET, VEGFRs, and PDGFRs. It is FDA approved for the treatment of unresectable hepatocellular carcinoma, advanced renal cell carcinoma, and recurrent or metastatic differentiated thyroid carcinoma. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical and preclinical data in KIT-mutated⁴⁰⁰⁻⁴⁰⁷ or KIT-expressing tumors⁷⁰⁻⁷³, KIT activating alterations may predict sensitivity to sorafenib. On the basis of clinical responses in patients with GIST, PDGFRA activating mutations may predict sensitivity to sorafenib^{131,408}.

SUPPORTING DATA

Phase 2 studies of sorafenib plus temozolomide report limited activity in patients with relapsed glioblastoma

multiforme (GBM)⁴⁰⁹. A Phase 1/2 trial of temsirolimus in combination with sorafenib in patients with glioblastoma was terminated at the Phase 2 interim analysis after patients failed to meet the primary endpoint of 6 month progression-free survival⁴¹⁰. A Phase 2 trial of sorafenib and erlotinib in glioblastoma also did not meet its primary endpoint, and erlotinib clearance was increased by the addition of sorafenib⁴¹¹. In a Phase 1 trial in patients with high-grade glioma, the combination of sorafenib with radiation therapy (RT) and temozolomide (TMZ) resulted in increased toxicity and did not result in significant improvement in clinical efficacy compared with RT and TMZ alone⁴¹². In a clinical study of sorafenib in pediatric patients with low-grade astrocytoma, one patient achieved a partial response (PR), one had stable disease (SD), and 9 patients had progressive disease; this study was terminated early due to unexpectedly high disease progression rates⁴¹³.

Sunitinib

Assay findings association
KIT
amplification

AREAS OF THERAPEUTIC USE

Sunitinib is a small-molecule tyrosine kinase inhibitor that targets PDGFRs, VEGFRs, KIT, FLT3, CSF-1R, and RET. It is FDA approved for the treatment of advanced or metastatic pancreatic neuroendocrine tumors, gastrointestinal stromal tumors (GISTs) in patients who have progressed on or are intolerant to imatinib, and advanced renal cell carcinoma (RCC) as well as for the adjuvant treatment of patients at high risk of recurrent RCC after nephrectomy. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical and preclinical data in KIT-mutated^{74,414-418} or KIT-expressing tumors⁷⁴⁻⁷⁵, KIT activating alterations may predict sensitivity to sunitinib.

SUPPORTING DATA

Phase 2 clinical trials of sunitinib in glioblastoma have reported no significant improvement in clinical outcome⁴¹⁹⁻⁴²⁰. A Phase 2 trial that examined sunitinib treatment followed by radiation therapy in patients with glioblastoma reported a median progression-free survival (PFS) of 7.7 weeks, and a median overall survival (OS) of 12.8 weeks; 83.3% (10/12) of patients experienced neurological deterioration prior to radiation therapy⁴²¹. Another Phase 2 study that examined daily sunitinib treatment in patients with glioblastoma reported no objective response in any of the 40 patients, with a median PFS of 2.2 months and a median OS of 9.2 months; five patients in the study had stable disease for more than six months⁴²².

NOTE Genomic alterations detected may be associated with activity of certain FDA approved drugs, however, the agents listed in this report may have varied evidence in the patient's tumor type.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 07 November 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

CLINICAL TRIALS

NOTE Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually

updated and should be investigated by the physician or research staff. This is not a comprehensive list of all available clinical trials. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials listed here may have additional enrollment criteria that

may require medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see [clinicaltrials.gov](https://www.clinicaltrials.gov). Or, visit <https://www.foundationmedicine.com/genomic-testing#support-services>.

GENE
KIT
ALTERATION
 amplification

RATIONALE

KIT amplification or activating mutations may predict sensitivity to small molecule tyrosine kinase inhibitors. Also, because KIT activation

leads to activation of the PI3K-AKT-mTOR pathway, PI3K and mTOR pathway inhibitors may be relevant in a tumor with KIT activation.

NCT05024214
PHASE 1/2

Phase Ib/II Trial of Envafolelimab Plus Lenvatinib for Subjects With Solid Tumors

TARGETS

PD-L1, FGFRs, RET, PDGFRA, VEGFRs, KIT, FLT3, CSF1R

LOCATIONS: Hangzhou (China), Shanghai (China), Dongguan (China), Guangzhou (China), Zhuhai (China), Benbu (China), Zhengzhou (China), Jinan (China), Dalian (China), Tianjin (China)

NCT05098847
PHASE 2

Cryoablation Combined With Sintilimab Plus Lenvatinib In Previously Treated Unresectable Liver Metastasis From Solid Tumors

TARGETS

FGFRs, RET, PDGFRA, VEGFRs, KIT, PD-1

LOCATIONS: Shanghai (China)

NCT04337463
PHASE NULL

ATG-008 Combined With Toripalimab in Advanced Solid Tumors

TARGETS

mTORC1, mTORC2, PD-1

LOCATIONS: Chongqing (China), Chengdu (China)

NCT04803318
PHASE 2

Trametinib Combined With Everolimus and Lenvatinib for Recurrent/Refractory Advanced Solid Tumors

TARGETS

mTOR, FGFRs, RET, PDGFRA, VEGFRs, KIT, MEK

LOCATIONS: Guangzhou (China)

NCT04977453
PHASE 1/2

GI-101 as a Single Agent or in Combination With Pembrolizumab, Lenvatinib or Local Radiotherapy in Advanced Solid Tumors

TARGETS

FGFRs, RET, PDGFRA, VEGFRs, KIT, PD-1, CTLA-4

LOCATIONS: Daejeon (Korea, Republic of), Suwon-si (Korea, Republic of), Seoul (Korea, Republic of)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 07 November 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

CLINICAL TRIALS
NCT03564691
PHASE 1

Study of MK-4830 as Monotherapy and in Combination With Pembrolizumab (MK-3475) in Participants With Advanced Solid Tumors (MK-4830-001)

TARGETS

ITL4, FGFRs, RET, PDGFRA, VEGFRs, KIT, PD-1

LOCATIONS: Seoul (Korea, Republic of), Liverpool (Australia), Petah Tikva (Israel), Ramat Gan (Israel), Tel Aviv (Israel), Haifa (Israel), Warszawa (Poland), Gdansk (Poland), Thessaloniki (Greece), Heraklion (Greece)

NCT04008797
PHASE 1

A Study of E7386 in Combination With Other Anticancer Drug in Participants With Solid Tumor

TARGETS

CBP, Beta-catenin, FGFRs, RET, PDGFRA, VEGFRs, KIT

LOCATIONS: Osakasayama (Japan), Chuo-Ku (Japan), Chiba (Japan), Kashiwa (Japan)

NCT03025893
PHASE 2/3

A Phase II/III Study of High-dose, Intermittent Sunitinib in Patients With Recurrent Glioblastoma Multiforme

TARGETS

FLT3, VEGFRs, CSF1R, KIT, RET

LOCATIONS: Groningen (Netherlands), Nijmegen (Netherlands), Amsterdam (Netherlands)

NCT04729348
PHASE 2

Pembrolizumab And Lenvatinib In Leptomeningeal Metastases

TARGETS

PD-1, KIT, VEGFRs, FGFRs, PDGFRA, RET

LOCATIONS: Massachusetts

NCT02379416
PHASE 1

Combination Nilotinib and Paclitaxel in Adults With Relapsed Solid Tumors

TARGETS

ABL, KIT

LOCATIONS: Maryland

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 07 November 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

CLINICAL TRIALS
GENE
MDM4
RATIONALE

Tumors with MDM4 amplification or overexpression and wild-type p53 may be

sensitive to inhibitors of MDM-p53 interactions.

ALTERATION

amplification

NCT03725436
PHASE 1

ALRN-6924 and Paclitaxel in Treating Patients With Advanced, Metastatic, or Unresectable Solid Tumors

TARGETS
 MDM2, MDM4

LOCATIONS: Texas

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 07 November 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

CLINICAL TRIALS
GENE
PDGFRA
RATIONALE

PDGFRA amplification may predict sensitivity to imatinib and to anti-PDGFRA antibodies.

PDGFRA activating mutations may predict sensitivity to certain PDGFRA-targeted therapies.

ALTERATION

 deletion exons 8-9, amplification, E229K
 - subclonal, R841K - subclonal

NCT03970447
PHASE 2/3

A Trial to Evaluate Multiple Regimens in Newly Diagnosed and Recurrent Glioblastoma

TARGETS

BRAF, VEGFRs, RET, KIT

LOCATIONS: Utah, Michigan, New York, Alabama

NCT03025893
PHASE 2/3

A Phase II/III Study of High-dose, Intermittent Sunitinib in Patients With Recurrent Glioblastoma Multiforme

TARGETS

FLT3, VEGFRs, CSF1R, KIT, RET

LOCATIONS: Groningen (Netherlands), Nijmegen (Netherlands), Amsterdam (Netherlands)

NCT05159245
PHASE 2

The Finnish National Study to Facilitate Patient Access to Targeted Anti-cancer Drugs

TARGETS

BRAF, VEGFRs, RET, KIT, ERBB2, TRKB, ALK, TRKC, ROS1, TRKA, SMO, PD-L1, MEK, CDK4, CDK6

LOCATIONS: Kuopio (Finland), Helsinki (Finland), Tampere (Finland), Turku (Finland)

NCT02379416
PHASE 1

Combination Nilotinib and Paclitaxel in Adults With Relapsed Solid Tumors

TARGETS

ABL, KIT

LOCATIONS: Maryland

NCT04771520
PHASE 2

Avapritinib for the Treatment of CKIT or PDGFRA Mutation-Positive Locally Advanced or Metastatic Malignant Solid Tumors

TARGETS

KIT, PDGFRA

LOCATIONS: Texas

NCT01738139
PHASE 1

Ipilimumab and Imatinib Mesylate in Advanced Cancer

TARGETS

KIT, ABL, CTLA-4

LOCATIONS: Texas

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 07 November 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

CLINICAL TRIALS
GENE
PIK3CA
ALTERATION
E545K
RATIONALE

PIK3CA activating mutations may lead to activation of the PI3K-AKT-mTOR pathway and may therefore indicate sensitivity to inhibitors of

this pathway. Strong clinical data support sensitivity of PIK3CA-mutated solid tumors to the PI3K-alpha inhibitor alpelisib.

NCT04589845
PHASE 2

Tumor-Agnostic Precision Immuno-Oncology and Somatic Targeting Rational for You (TAPISTRY) Platform Study

TARGETS

TRKB, ALK, TRKC, ROS1, TRKA, RET, PD-L1, AKTs, ERBB2, MDM2, PI3K-alpha, RAFs, NRAS

LOCATIONS: Taipei City (Taiwan), Taoyuan County (Taiwan), Tainan (Taiwan), Shanghai City (China), Shanghai (China), Shatin (Hong Kong), Hong Kong (Hong Kong), Seoul (Korea, Republic of), Xi'an (China), Tianjin (China)

NCT04341259
PHASE 1

A Study Of The Pharmacokinetics And Safety Of Ipatasertib In Chinese Participants With Locally Advanced Or Metastatic Solid Tumors.

TARGETS

AKTs

LOCATIONS: Shanghai City (China)

NCT03239015
PHASE 2

Efficacy and Safety of Targeted Precision Therapy in Refractory Tumor With Druggable Molecular Event

TARGETS

EGFR, ERBB4, ERBB2, PARP, mTOR, MET, ROS1, RET, VEGFRs, BRAF, CDK4, CDK6

LOCATIONS: Shanghai (China)

NCT04337463
PHASE NULL

ATG-008 Combined With Toripalimab in Advanced Solid Tumors

TARGETS

mTORC1, mTORC2, PD-1

LOCATIONS: Chongqing (China), Chengdu (China)

NCT04803318
PHASE 2

Trametinib Combined With Everolimus and Lenvatinib for Recurrent/Refractory Advanced Solid Tumors

TARGETS

mTOR, FGFRs, RET, PDGFRA, VEGFRs, KIT, MEK

LOCATIONS: Guangzhou (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 07 November 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

CLINICAL TRIALS
NCT04526470
PHASE 1/2

Alpelisib and Paclitaxel in PIK3CA-altered Gastric Cancer

TARGETS
 PI3K-alpha

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of)

NCT05125523
PHASE 1

A Study of Sirolimus for Injection (Albumin Bound) in Patients With Advanced Solid Tumors

TARGETS
 mTOR

LOCATIONS: Tianjin (China)

NCT03772561
PHASE 1

Phase I Study of AZD5363 + Olaparib + Durvalumab in Patients With Advanced or Metastatic Solid Tumor Malignancies

TARGETS
 PARP, AKTs, PD-L1

LOCATIONS: Singapore (Singapore)

NCT04801966
PHASE NULL

Safety and Oversight of the Individually Tailored Treatment Approach: A Novel Pilot Study

TARGETS
 CDK4, CDK6, PI3K-alpha, PD-L1, MEK, PARP, PD-1, BRAF

LOCATIONS: Melbourne (Australia)

NCT04317105
PHASE 1/2

Testing the Addition of an Anti-cancer Drug, Copanlisib, to the Usual Immunotherapy (Nivolumab With or Without Ipilimumab) in Patients With Advanced Solid Cancers That Have Changes in the Following Genes: PIK3CA and PTEN

TARGETS
 PD-1, CTLA-4, PI3K

LOCATIONS: Toronto (Canada), Texas, Virginia

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 07 November 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

APPENDIX
Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

ERBB4
D1179E

FGFR4
R234H

FOXL2
R349G

GSK3B
T403A

KDM5A
T269N

MLL2
S2215T

PDGFRA
C290Y, S348_E358del and
S348_I363del

PIK3C2G
S1238L

RAD52
G125C

ROS1
Y1783H

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 07 November 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

APPENDIX

Genes Assayed in FoundationOne®CDx

FoundationOne CDx is designed to include genes known to be somatically altered in human solid tumors that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay interrogates 324 genes as well as introns of 36 genes involved in rearrangements. The assay will be updated periodically to reflect new knowledge about cancer biology.

DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS

ABL1	ACVR1B	AKT1	AKT2	AKT3	ALK	ALOX12B	AMER1 (FAM123B or WTX)	
APC	AR	ARAF	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX
AURKA	AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2
BCL6	BCOR	BCORL1	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1
BTG2	BTK	CALR	CARD11	CASP8	CBFB	CBL	CCND1	CCND2
CCND3	CCNE1	CD22	CD274 (PD-L1)	CD70	CD79A	CD79B	CDC73	CDH1
CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B	CDKN2C
CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R	CTCF
CTNNA1	CTNNB1	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1	DDR2
DIS3	DNMT3A	DOT1L	EED	EGFR	EMSY (C11orf30)	EP300	EPHA3	EPHB1
EPHB4	ERBB2	ERBB3	ERBB4	ERCC4	ERG	ERRF1	ESR1	EZH2
FANCA	FANCC	FANCG	FANCL	FAS	FBXW7	FGF10	FGF12	FGF14
FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1	FGFR2	FGFR3	FGFR4
FH	FLCN	FLT1	FLT3	FOXL2	FUBP1	GABRA6	GATA3	GATA4
GATA6	GID4 (C17orf39)	GNA11	GNA13	GNAQ	GNAS	GRM3	GSK3B	H3-3A (H3F3A)
HDAC1	HGF	HNFI1A	HRAS	HSD3B1	ID3	IDH1	IDH2	IGF1R
IKBKE	IKZF1	INPP4B	IRF2	IRF4	IRS2	JAK1	JAK2	JAK3
JUN	KDM5A	KDM5C	KDM6A	KDR	KEAP1	KEL	KIT	KLHL6
KMT2A (MLL)	KMT2D (MLL2)	KRAS	LTK	LYN	MAF	MAP2K1 (MEK1)	MAP2K2 (MEK2)	MAP2K4
MAP3K1	MAP3K13	MAPK1	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1
MERTK	MET	MITF	MKNK1	MLH1	MPL	MRE11 (MRE11A)	MSH2	MSH3
MSH6	MST1R	MTAP	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN	MYD88
NBN	NF1	NF2	NFE2L2	NFKBIA	NKX2-1	NOTCH1	NOTCH2	NOTCH3
NPM1	NRAS	NSD2 (WHSC1 or MMSET)	NSD3 (WHSC1L1)	NT5C2	NTRK1	NTRK2	NTRK3	NTRK3
P2RY8	PALB2	PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)
PDGFRA	PDGFRB	PDK1	PIK3C2B	PIK3C2G	PIK3CA	PIK3CB	PIK3R1	PIM1
PMS2	POLD1	POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI
PRKN (PARK2)	PTCH1	PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51
RAD51B	RAD51C	RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10
REL	RET	RICTOR	RNF43	ROS1	RPTOR	SDHA	SDHB	SDHC
SDHD	SETD2	SF3B1	SGK1	SMAD2	SMAD4	SMARCA4	SMARCB1	SMO
SNCAIP	SOC1	SOX2	SOX9	SPEN	SPOP	SRC	STAG2	STAT3
STK11	SUFU	SYK	TBX3	TEK	TENT5C (FAM46C)	TET2	TET2	TGFB2
TIPARP	TNFAIP3	TNFRSF14	TP53	TSC1	TSC2	TYRO3	U2AF1	VEGFA
VHL	WT1	XPO1	XRCC2	ZNF217	ZNF703			

DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS

ALK	BCL2	BCR	BRAF	BRCA1	BRCA2	CD74	EGFR	ETV4
ETV5	ETV6	EWSR1	EZR	FGFR1	FGFR2	FGFR3	KIT	KMT2A (MLL)
MSH2	MYB	MYC	NOTCH2	NTRK1	NTRK2	NUTM1	PDGFRA	RAF1
RARA	RET	ROS1	RSP02	SDC4	SLC34A2	TERC*	TERT**	TPRSS2

*TERC is an NCRNA

**Promoter region of TERT is interrogated

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS

Homologous Recombination status
Loss of Heterozygosity (LOH) score
Microsatellite (MS) status
Tumor Mutational Burden (TMB)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.


Electronically signed by Erik Williams, M.D. | 07 November 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

APPENDIX

About FoundationOne®CDx

FoundationOne CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Ciplstraat 3, 2440 Geel, Belgium. 

ABOUT FOUNDATIONONE CDx

FoundationOne CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details:
www.rochefoundationmedicine.com/f1cdxtech.

INTENDED USE

FoundationOne®CDx (F1CDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI), tumor mutational burden (TMB), and for selected forms of ovarian cancer, loss of heterozygosity (LOH) score, using DNA isolated from formalin-fixed, paraffin-embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with therapies in accordance with approved therapeutic product labeling. Additionally, F1CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms.

TEST PRINCIPLE

FoundationOne CDx will be performed exclusively as a laboratory service using DNA extracted from formalin-fixed, paraffin-embedded (FFPE) tumor samples. The proposed assay will employ a single DNA extraction method from routine FFPE biopsy or surgical resection specimens, 50-1000 ng of which will undergo whole-genome shotgun library construction and hybridization-based capture of all coding exons from 309 cancer-related genes, one promoter region, one non-coding (ncRNA), and select intronic regions from 34 commonly rearranged genes, 21 of which also include the coding exons. The assay therefore includes

detection of alterations in a total of 324 genes.

Using an Illumina® HiSeq platform, hybrid capture-selected libraries will be sequenced to high uniform depth (targeting >500X median coverage with >99% of exons at coverage >100X). Sequence data will be processed using a customized analysis pipeline designed to accurately detect all classes of genomic alterations, including base substitutions, indels, focal copy number amplifications, homozygous gene deletions, and selected genomic rearrangements (e.g. gene fusions). Additionally, genomic signatures including loss of heterozygosity (LOH), microsatellite instability (MSI) and tumor mutational burden (TMB) will be reported.

THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. The F1CDx report may be used as an aid to inform molecular eligibility for clinical trials. Note: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

Diagnostic Significance

FoundationOne CDx identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

Qualified Alteration Calls (Equivocal and Subclonal)

An alteration denoted as "amplification – equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne CDx for identifying a copy number amplification is four (4) for *ERBB2* and six (6) for all other genes. Conversely, an alteration denoted as "loss – equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is one that the FoundationOne CDx analytical

methodology has identified as being present in <10% of the assayed tumor DNA.

Ranking of Therapies and Clinical Trials

Ranking of Therapies in Summary Table

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

Ranking of Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) (www.nccn.org). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2022. All rights reserved. To view the most recent and complete version of the guidelines, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

Limitations

1. In the fraction-based MSI algorithm, a tumor specimen will be categorized as MSI-H, MSS, or MS-Equivocal according to the fraction of microsatellite loci determined to be altered or unstable (i.e., the fraction unstable loci score). In the F1CDx assay, MSI is evaluated based on a genome-wide analysis across >2000 microsatellite loci. For a given microsatellite locus, non-somatic alleles are discarded, and the microsatellite is categorized as unstable if remaining alleles differ from the reference genome. The final fraction unstable loci score is calculated as the number of unstable microsatellite loci divided by the number of evaluable microsatellite loci. The MSI-H and MSS cut-off thresholds were determined by

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 07 November 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

APPENDIX

About FoundationOne®CDx

analytical concordance to a PCR comparator assay using a pan-tumor FFPE tissue sample set. Patients with results categorized as "MS-Stable" with median exon coverage <300X, "MS-Equivocal," or "Cannot Be Determined" should receive confirmatory testing using a validated orthogonal (alternative) method.

2. TMB by F1CDx is determined by counting all synonymous and non-synonymous variants present at 5% allele frequency or greater (after filtering) and the total number is reported as mutations per megabase (mut/Mb) unit. Observed TMB is dependent on characteristics of the specific tumor focus tested for a patient (e.g., primary vs. metastatic, tumor content) and the testing platform used for the detection; therefore, observed TMB results may vary between different specimens for the same patient and between detection methodologies employed on the same sample. The TMB calculation may differ from TMB calculations used by other assays depending on variables such as the amount of genome interrogated, percentage of tumor, assay limit of detection (LoD), filtering of alterations included in the score, and the read depth and other bioinformatic test specifications. Refer to the SSED for a detailed description of these variables in FMI's TMB calculation https://www.accessdata.fda.gov/cdrh_docs/pdf17/P170019B.pdf. The clinical validity of TMB defined by this panel has been established for TMB as a qualitative output for a cut-off of 10 mutations per megabase but has not been established for TMB as a quantitative score.
3. Homologous Recombination status may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas (Coleman et al., 2017; 28916367). Samples with deleterious *BRCA1/2* alteration and/or Loss of Heterozygosity (LOH) score $\geq 16\%$ will be reported as "HRD Positive" and samples with absence of these findings will be reported as "HRD Not Detected," agnostic of potential secondary *BRCA1/2* reversion alterations. Certain potentially deleterious missense or small in-frame deletions in *BRCA1/2* may not be classified as deleterious and, in the absence of an elevated LOH profile, samples with such mutations may be classified as "HRD Not Detected." A result of "HRD Not Detected" does not rule out the presence of a *BRCA1/2* alteration or an elevated LOH profile outside the assay performance characteristic limitations.
4. The LOH score is determined by analyzing SNPs spaced at 1Mb intervals across the genome on the FoundationOne CDx test and

extrapolating an LOH profile, excluding arm- and chromosome-wide LOH segments. Detection of LOH has been verified only for ovarian cancer patients, and the LOH score result may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas. The LOH score will be reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine LOH. Performance of the LOH classification has not been established for samples below 35% tumor content. There may be potential interference of ethanol with LOH detection. The interfering effects of xylene, hemoglobin, and triglycerides on the LOH score have not been demonstrated.

5. Alterations reported may include somatic (not inherited) or germline (inherited) alterations; however, the test does not distinguish between germline and somatic alterations. The test does not provide information about susceptibility.
6. Biopsy may pose a risk to the patient when archival tissue is not available for use with the assay. The patient's physician should determine whether the patient is a candidate for biopsy.
7. Reflex testing to an alternative FDA approved companion diagnostic should be performed for patients who have an *ERBB2* amplification result detected with copy number equal to 4 (baseline ploidy of tumor +2) for confirmatory testing. While this result is considered negative by FoundationOne®CDx (F1CDx), in a clinical concordance study with an FDA approved FISH test, 70% (7 out of 10 samples) were positive, and 30% (3 out of 10 samples) were negative by the FISH test with an average ratio of 2.3. The frequency of *ERBB2* copy number 4 in breast cancer is estimated to be approximately 2%. Multiple references listed in <https://www.mycancergenome.org/content/disease/breast-cancer/ERBB2/238/> report the frequency of HER2 overexpression as 20% in breast cancer. Based on the F1CDx HER2 CDx concordance study, approximately 10% of HER2 amplified samples had copy number 4. Thus, total frequency is conservatively estimated to be approximately 2%.

REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic, nontargeted treatment, germline, or clonal

hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

VARIANT ALLELE FREQUENCY

Variant Allele Frequency (VAF) represents the fraction of sequencing reads in which the variant is observed. This attribute is not taken into account for therapy inclusion, clinical trial matching, or interpretive content. Caution is recommended in interpreting VAF to indicate the potential germline or somatic origin of an alteration, recognizing that tumor fraction and tumor ploidy of samples may vary.

Precision of VAF for base substitutions and indels

BASE SUBSTITUTIONS	%CV*
Repeatability	5.11 - 10.40
Reproducibility	5.95 - 12.31
INDELS	%CV*
Repeatability	6.29 - 10.00
Reproducibility	7.33 - 11.71

*Interquartile Range = 1st Quartile to 3rd Quartile

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >10%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM*, *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *CHEK2*, *FH*, *FLCN*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *PALB2*, *PMS2*, *POLE*, *RAD51C*, *RAD51D*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *TSC2*, and *VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 07 November 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1488563-01

APPENDIX

About FoundationOne®CDx

tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1*, *CBL*, *DNMT3A*, *IDH2*, *JAK2*, *KMT2D (MLL2)*, *MPL*, *MYD88*, *SF3B1*, *TET2*, and *U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne CDx.

TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking

into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. FoundationOne CDx is performed using DNA derived from tumor, and as such germline events may not be reported.

SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
mut/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

REFERENCE SEQUENCE INFORMATION

Sequence data is mapped to the human genome, Genome Reference Consortium Human Build 37 (GRCh37), also known as hg19.

MR Suite Version (RG) 7.3.0

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of the data. The median exon coverage for this sample is 734x. The suitability of use.

ORDERED TEST # **ORD-1488563-01**
APPENDIX **References**

1. Gatalica Z, et al. Cancer Epidemiol. Biomarkers Prev. (2014) PMID: 25392179
2. Kroemer G, et al. Oncoimmunology (2015) PMID: 26140250
3. Lal N, et al. Oncoimmunology (2015) PMID: 25949894
4. Le DT, et al. N. Engl. J. Med. (2015) PMID: 26028255
5. Ayers et al., 2016; ASCO-SITC Abstract P60
6. Martinez R, et al. Oncology (2004) PMID: 15331927
7. Martinez R, et al. J. Cancer Res. Clin. Oncol. (2005) PMID: 15672285
8. Martinez R, et al. Cancer Genet. Cytogenet. (2007) PMID: 17498554
9. Szybka M, et al. Clin. Neuropathol. (2012) PMID: 12908754
10. Kocarnik JM, et al. Gastroenterol Rep (Oxf) (2015) PMID: 26337942
11. You JF, et al. Br. J. Cancer (2010) PMID: 21081928
12. Bairwa NK, et al. Methods Mol. Biol. (2014) PMID: 24623249
13. Boland CR, et al. Cancer Res. (1998) PMID: 9823339
14. Pawlik TM, et al. Dis. Markers (2004) PMID: 15528785
15. Boland CR, et al. Gastroenterology (2010) PMID: 20420947
16. Samstein RM, et al. Nat. Genet. (2019) PMID: 30643254
17. Goodman AM, et al. Mol. Cancer Ther. (2017) PMID: 28835386
18. Goodman AM, et al. Cancer Immunol Res (2019) PMID: 31405947
19. Cristescu R, et al. Science (2018) PMID: 30309915
20. Ready N, et al. J. Clin. Oncol. (2019) PMID: 30785829
21. Hellmann MD, et al. N. Engl. J. Med. (2018) PMID: 29658845
22. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
23. Hellmann MD, et al. Cancer Cell (2018) PMID: 29731394
24. Rozeman EA, et al. Nat Med (2021) PMID: 33558721
25. Sharma P, et al. Cancer Cell (2020) PMID: 32916128
26. Zhao J, et al. Nat. Med. (2019) PMID: 30742119
27. Touat M, et al. Nature (2020) PMID: 32322066
28. Bouffett E, et al. J. Clin. Oncol. (2016) PMID: 27001570
29. Johanns TM, et al. Cancer Discov (2016) PMID: 27683556
30. Lukas RV, et al. J. Neurooncol. (2018) PMID: 30073642
31. Chalmers ZR, et al. Genome Med (2017) PMID: 28420421
32. Patel RR, et al. Pediatr Blood Cancer (2020) PMID: 32386112
33. Johnson A, et al. Oncologist (2017) PMID: 28912153
34. Draaisma K, et al. Acta Neuropathol Commun (2015) PMID: 26699864
35. Wang L, et al. BMC Cancer (2020) PMID: 32164609
36. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
37. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
38. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
39. Rizvi NA, et al. Science (2015) PMID: 25765070
40. Johnson BE, et al. Science (2014) PMID: 24336570
41. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
42. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
43. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
44. Heitzer E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
45. Nature (2012) PMID: 22810696
46. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
47. Evans EK, et al. Sci Transl Med (2017) PMID: 29093181
48. Abbaspour Babaei M, et al. Drug Des Devel Ther (2016) PMID: 27536065
49. Ramaswamy A, et al. J Gastrointest Oncol (2016) PMID: 27563456
50. Demetri GD, et al. Lancet (2013) PMID: 23177515
51. Gotlib J, et al. N. Engl. J. Med. (2016) PMID: 27355533
52. Jawhar M, et al. Blood (2017) PMID: 28424161
53. Xu X, et al. Int J Clin Exp Pathol (2014) PMID: 25031773
54. Gotlib J, et al. Blood (2005) PMID: 15972446
55. Luo C, et al. Onco Targets Ther (2017) PMID: 29066909
56. Si L, et al. J. Clin. Oncol. (2012) PMID: 22162580
57. Parikh SA, et al. Leuk Lymphoma (2010) PMID: 20038218
58. Wei X, et al. Oncol. Res. (2019) PMID: 30075827
59. Hodi FS, et al. J. Clin. Oncol. (2013) PMID: 23775962
60. Carvajal RD, et al. JAMA (2011) PMID: 21642685
61. Guo J, et al. J. Clin. Oncol. (2011) PMID: 21690468
62. Debiec-Rychter M, et al. Gastroenterology (2005) PMID: 15685537
63. Dematteo RP, et al. Lancet (2009) PMID: 19303137
64. Faivre S, et al. J. Clin. Oncol. (2005) PMID: 16135502
65. Hotte SJ, et al. J. Clin. Oncol. (2005) PMID: 15659505
66. Alcedo JC, et al. Head Neck (2004) PMID: 15350030
67. Brandwein JM, et al. Leukemia (2011) PMID: 21403650
68. Reardon DA, et al. Br. J. Cancer (2009) PMID: 19904263
69. Lee SJ, et al. Oncologist (2015) PMID: 26424760
70. Llovet JM, et al. Clin. Cancer Res. (2012) PMID: 22374331
71. Zhang HL, et al. Clin Genitourin Cancer (2013) PMID: 23058498
72. Seino S, et al. Gastroenterology (2014) PMID: 25450081
73. Li XF, et al. Med. Oncol. (2009) PMID: 18846437
74. Minor DR, et al. Clin. Cancer Res. (2012) PMID: 22261812
75. Mahipal A, et al. Melanoma Res. (2012) PMID: 23114504
76. Cancer Genome Atlas Research Network, et al. N. Engl. J. Med. (2015) PMID: 26061751
77. Brennan CW, et al. Cell (2013) PMID: 24120142
78. Nobusawa S, et al. Neuropathology (2011) PMID: 21382095
79. Joensuu H, et al. J. Pathol. (2005) PMID: 16021678
80. Burford A, et al. PLoS ONE (2013) PMID: 23990986
81. Holtkamp N, et al. Neuro-oncology (2007) PMID: 17504929
82. Puputti M, et al. Mol. Cancer Res. (2006) PMID: 17189383
83. Skardelly M, et al. Transl Oncol (2009) PMID: 19701495
84. Int. J. Biochem. Cell Biol. (1999) PMID: 10582339
85. Semin. Oncol. (2004) PMID: 15175998
86. Gao J, et al. Sci Signal (2013) PMID: 23550210
87. Zack TI, et al. Nat. Genet. (2013) PMID: 24071852
88. Beroukhim R, et al. Nature (2010) PMID: 20164920
89. Arefi M, et al. Int. J. Hematol. (2012) PMID: 22806436
90. Baccarani M, et al. Haematologica (2007) PMID: 17666373
91. Cassier PA, et al. Clin. Cancer Res. (2012) PMID: 22718859
92. Chalmers ZR, et al. Blood Cancer J (2015) PMID: 25658984
93. Cools J, et al. N. Engl. J. Med. (2003) PMID: 12660384
94. Curtis CE, et al. Br. J. Haematol. (2007) PMID: 17555450
95. Debiec-Rychter M, et al. Eur. J. Cancer (2004) PMID: 15010069
96. Dileo P, et al. Int. J. Cancer (2011) PMID: 20473908
97. Fanta PT, et al. J. Clin. Oncol. (2015) PMID: 24638008
98. Florian S, et al. Leuk. Res. (2006) PMID: 16406018
99. Frenard C, et al. JAAD Case Rep (2016) PMID: 27051816
100. Griffin JH, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) PMID: 12808148
101. Heinrich MC, et al. J. Clin. Oncol. (2003) PMID: 14645423
102. Helbig G, et al. Br. J. Haematol. (2009) PMID: 19120352
103. Helbig G, et al. Am. J. Hematol. (2014) PMID: 24009127
104. Hus M, et al. Leuk. Res. (2011) PMID: 21093052
105. Ikezoe T, et al. Leuk. Res. (2010) PMID: 20303172
106. Intermetz T, et al. Br. J. Haematol. (2009) PMID: 19735261
107. Jain N, et al. Leuk. Res. (2009) PMID: 19013640
108. Jovanovic JV, et al. Blood (2007) PMID: 17299092
109. Kang HJ, et al. Acta Oncol (2012) PMID: 22150077
110. Klion AD, et al. Blood (2004) PMID: 14504092
111. Kobayashi M, et al. Respirology (2009) PMID: 19192229
112. Kocáková I, et al. Klin Onkol (2014) PMID: 24635438
113. Metzgeroth G, et al. Br. J. Haematol. (2008) PMID: 18950453
114. Murayama Y, et al. World J Gastrointest Oncol (2012) PMID: 22645636
115. Ogbogu PU, et al. J. Allergy Clin. Immunol. (2009) PMID: 19910029
116. Ohnishi H, et al. Br. J. Haematol. (2006) PMID: 16856885
117. Pardanani A, et al. Blood (2003) PMID: 12842979
118. Pardanani A, et al. Blood (2004) PMID: 15284118
119. Qu SQ, et al. Oncotarget (2016) PMID: 27120808
120. Score J, et al. Leukemia (2006) PMID: 16498388
121. Shah S, et al. J Hematol Oncol (2014) PMID: 24669761
122. Sugimoto Y, et al. Cancer Genet (2015) PMID: 26319757
123. Volz HC, et al. Int. J. Cardiol. (2011) PMID: 20609486
124. von Bubnoff N, et al. Leukemia (2005) PMID: 15618966
125. Walz C, et al. Genes Chromosomes Cancer (2006) PMID: 16845659
126. Yoo C, et al. Cancer Res Treat (2016) PMID: 26130666
127. Al-Riyami AZ, et al. Leuk. Lymphoma (2013) PMID: 23157309
128. Lierman E, et al. Blood (2006) PMID: 16645167
129. Lierman E, et al. Leukemia (2009) PMID: 19212337
130. Metzgeroth G, et al. Leukemia (2012) PMID: 21818111
131. Roubaud G, et al. Ann. Oncol. (2012) PMID: 22294526
132. von Bubnoff N, et al. Oncogene (2011) PMID: 20972453
133. Hochhaus A, et al. J. Cancer Res. Clin. Oncol. (2013) PMID: 24057647
134. Tabouret E, et al. Leuk. Res. (2011) PMID: 20832858
135. Dewaele B, et al. Clin. Cancer Res. (2008) PMID: 18794084
136. Weisberg E, et al. Gastroenterology (2006) PMID: 17087936
137. Brohl AS, et al. Clin Sarcoma Res (2015) PMID: 26396737
138. Grellety T, et al. Future Sci OA (2015) PMID: 28031906
139. Kollár A, et al. Clin Sarcoma Res (2014) PMID: 25905001
140. Jaku et al., 2017; ASCO Abstract 2515
141. Phillips JJ, et al. Brain Pathol. (2013) PMID: 23438035
142. Motomura K, et al. J. Neuropathol. Exp. Neurol. (2013) PMID: 23242283
143. Sottoriva A, et al. Proc. Natl. Acad. Sci. U.S.A. (2013) PMID: 23412337
144. Alentorn A, et al. Neuro-oncology (2012) PMID: 23074200
145. Tate JG, et al. Nucleic Acids Res. (2019) PMID: 30371878
146. Nature (2008) PMID: 18772890
147. Hoadley KA, et al. Cell (2018) PMID: 29625048
148. Ellrott K, et al. Cell Syst (2018) PMID: 29596782
149. Taylor AM, et al. Cancer Cell (2018) PMID: 29622463
150. Gao Q, et al. Cell Rep (2018) PMID: 29617662
151. Liu J, et al. Cell (2018) PMID: 29625055
152. Sanchez-Vega F, et al. Cell (2018) PMID: 29625050

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # **ORD-1488563-01**
APPENDIX
References

153. Song K, et al. Am J Cancer Res (2018) PMID: 29888103
154. Szerlip NJ, et al. Proc. Natl. Acad. Sci. U.S.A. (2012) PMID: 22323597
155. Andrae J, et al. Genes Dev. (2008) PMID: 18483217
156. Flavahan WA, et al. Nature (2016) PMID: 26700815
157. Roszik J, et al. Sci Rep (2016) PMID: 26787600
158. Verhaak RG, et al. Cancer Cell (2010) PMID: 20129251
159. Koschmann C, et al. Oncotarget (2016) PMID: 27582545
160. Puget S, et al. PLoS ONE (2012) PMID: 22389665
161. Clarke ID, et al. Oncogene (2003) PMID: 12569364
162. Ozawa T, et al. Genes Dev. (2010) PMID: 20889717
163. Mascarenhas J, et al. Blood Adv (2022) PMID: 34933330
164. Mascarenhas J, et al. Blood (2019) PMID: 31167802
165. Reis et al 2016; 26869629
166. Chen L, et al. Int J Cancer (2019) PMID: 30536898
167. Vaseva AV, et al. Cell Death Dis (2011) PMID: 21562588
168. Wang H, et al. Neoplasia (2011) PMID: 21750655
169. Wang H, et al. Mol. Cancer Ther. (2011) PMID: 21075910
170. Reed D, et al. J. Biol. Chem. (2010) PMID: 20080970
171. Smalley KS, et al. Cancer Res. (2007) PMID: 17210701
172. Vogel SM, et al. Proc. Natl. Acad. Sci. U.S.A. (2012) PMID: 23035244
173. Mandke P, et al. PLoS ONE (2012) PMID: 22870278
174. Hirose M, et al. Oncoscience (2014) PMID: 25621298
175. Chang YS, et al. Proc. Natl. Acad. Sci. U.S.A. (2013) PMID: 23946421
176. Bernal F, et al. Cancer Cell (2010) PMID: 21075307
177. Wade M, et al. Nat. Rev. Cancer (2013) PMID: 23303139
178. Arjona D, et al. Diagn. Mol. Pathol. (2005) PMID: 16319692
179. Schiffman JD, et al. Cancer Res. (2010) PMID: 20068183
180. Nobusawa S, et al. Brain Pathol. (2010) PMID: 20406234
181. Rao SK, et al. J. Neurooncol. (2010) PMID: 19609742
182. Jin G, et al. Neuro-oncology (2010) PMID: 20472715
183. Riemenschneider MJ, et al. Cancer Res. (1999) PMID: 10626796
184. Mancini F, et al. EMBO J. (2009) PMID: 19521340
185. Danovi D, et al. Mol. Cell. Biol. (2004) PMID: 15199139
186. Biegling KT, et al. Nat. Rev. Cancer (2014) PMID: 24739573
187. Fritsch C, et al. Mol. Cancer Ther. (2014) PMID: 24608574
188. Juric D, et al. J. Clin. Oncol. (2018) PMID: 29401002
189. Gallant JN, et al. NPJ Precis Oncol (2019) PMID: 30793038
190. Delestre F, et al. Sci Transl Med (2021) PMID: 34613809
191. Morschhauser F, et al. Mol Cancer Ther (2020) PMID: 31619463
192. Patnaik A, et al. Ann. Oncol. (2016) PMID: 27672108
193. Santin AD, et al. Gynecol Oncol Rep (2020) PMID: 31934607
194. Damodaran S, et al. J Clin Oncol (2022) PMID: 35133871
195. André F, et al. N. Engl. J. Med. (2019) PMID: 31091374
196. Smyth LM, et al. NPJ Breast Cancer (2021) PMID: 33863913
197. Varnier R, et al. Eur J Cancer (2019) PMID: 31351267
198. Basse C, et al. JCO Precis Oncol (2018) PMID: 32914004
199. Sultova E, et al. Arch Gynecol Obstet (2021) PMID: 33277683
200. Mackay HJ, et al. Cancer (2014) PMID: 24166148
201. Myers AP, et al. Gynecol. Oncol. (2016) PMID: 27016228
202. Dhami J, et al. Cold Spring Harb Mol Case Stud (2018) PMID: 29588307
203. Harris EJ, et al. Front Oncol (2019) PMID: 30863722
204. Hanna GJ, et al. Clin Cancer Res (2018) PMID: 29301825
205. Krop et al., 2018; ASCO Abstract 101
206. Pascual J, et al. Cancer Discov (2021) PMID: 32958578
207. Dolly SO, et al. Clin. Cancer Res. (2016) PMID: 26787751
208. Canaud et al., 2021; ESMO Abstract LBA23
209. Aust Fam Physician (1986) PMID: 2941002
210. Gallia GL, et al. Mol. Cancer Res. (2006) PMID: 17050665
211. Broderick DK, et al. Cancer Res. (2004) PMID: 15289301
212. El-Habr EA, et al. Clin. Neuropathol. () PMID: 20569675
213. Yan et al. 2020; DOI:10.1200/PO.19.00385
214. Derakhshandeh-Peykar P, et al. J. Neurogenet. (2011) PMID: 22026810
215. Tanaka S, et al. Acta Neuropathol Commun (2019) PMID: 31036078
216. Samuels Y, et al. Cancer Cell (2005) PMID: 15950905
217. Nat. Rev. Cancer (2009) PMID: 19629070
218. Kang S, et al. Proc. Natl. Acad. Sci. U.S.A. (2005) PMID: 15647370
219. Ikenoue T, et al. Cancer Res. (2005) PMID: 15930273
220. Gymnopoulos M, et al. Proc. Natl. Acad. Sci. U.S.A. (2007) PMID: 17376864
221. Horn S, et al. Oncogene (2008) PMID: 18317450
222. Rudd ML, et al. Clin. Cancer Res. (2011) PMID: 21266528
223. Hon WC, et al. Oncogene (2012) PMID: 22120714
224. Burke JE, et al. Proc. Natl. Acad. Sci. U.S.A. (2012) PMID: 22949682
225. Wu H, et al. Proc. Natl. Acad. Sci. U.S.A. (2009) PMID: 19915146
226. Laurenti R, et al. Rev Saude Publica (1990) PMID: 2103068
227. Dan S, et al. Cancer Res. (2010) PMID: 20530683
228. Oda K, et al. Cancer Res. (2008) PMID: 18829572
229. Zhao L, et al. Oncogene (2008) PMID: 18794883
230. Lui VW, et al. Cancer Discov (2013) PMID: 23619167
231. Ross RL, et al. Oncogene (2013) PMID: 22430209
232. Riviere JB, et al. Nat. Genet. (2012) PMID: 22729224
233. Shibata T, et al. Cancer Lett. (2009) PMID: 19394761
234. Dogruluk T, et al. Cancer Res. (2015) PMID: 26627007
235. Croessmann S, et al. Clin. Cancer Res. (2018) PMID: 29284706
236. Ng PK, et al. Cancer Cell (2018) PMID: 29533785
237. Spangle JM, et al. (2020) PMID: 32929011
238. Chen L, et al. Nat Commun (2018) PMID: 29636477
239. Jin N, et al. J Clin Invest (2021) PMID: 34779417
240. Konecny GE, et al. Clin. Cancer Res. (2011) PMID: 21278246
241. Katsumi Y, et al. Biochem. Biophys. Res. Commun. (2011) PMID: 21871868
242. Cen L, et al. Neuro-oncology (2012) PMID: 22711607
243. Logan JE, et al. Anticancer Res. (2013) PMID: 23898052
244. Fennell DA, et al. Lancet Oncol (2022) PMID: 35157829
245. Elvin JA, et al. Oncologist (2017) PMID: 28283584
246. Gao J, et al. Curr Oncol (2015) PMID: 26715889
247. Gopalan et al., 2014; ASCO Abstract 8077
248. Peguero et al., 2016; ASCO Abstract 2528
249. Konecny et al., 2016; ASCO Abstract 5557
250. DeMichele A, et al. Clin. Cancer Res. (2015) PMID: 25501126
251. Finn RS, et al. Lancet Oncol. (2015) PMID: 25524798
252. Infante JR, et al. Clin. Cancer Res. (2016) PMID: 27542767
253. Johnson DB, et al. Oncologist (2014) PMID: 24797823
254. Van Maerken T, et al. Mol. Cancer Ther. (2011) PMID: 21460101
255. Gamble LD, et al. Oncogene (2012) PMID: 21725357
256. Ceccarelli M, et al. Cell (2016) PMID: 26824661
257. Jonsson P, et al. Clin. Cancer Res. (2019) PMID: 31263031
258. Weber RG, et al. Oncogene (2007) PMID: 16909113
259. Nakamura M, et al. Brain Pathol. (2001) PMID: 11303791
260. Chakravarti A, et al. Clin. Cancer Res. (2001) PMID: 11489817
261. Feng J, et al. Cancer (2012) PMID: 21713760
262. Raabe EH, et al. Clin. Cancer Res. (2011) PMID: 21636552
263. Liu W, et al. J. Exp. Clin. Cancer Res. (2011) PMID: 21843312
264. Quelle DE, et al. Cell (1995) PMID: 8521522
265. Mutat. Res. (2005) PMID: 15878778
266. Gazzeri S, et al. Oncogene (1998) PMID: 9484839
267. Oncogene (1999) PMID: 10498883
268. Sherr CJ, et al. Cold Spring Harb. Symp. Quant. Biol. (2005) PMID: 16869746
269. Ozenne P, et al. Int. J. Cancer (2010) PMID: 20549699
270. Ruas M, et al. Oncogene (1999) PMID: 10498896
271. Jones R, et al. Cancer Res. (2007) PMID: 17909018
272. Haferkamp S, et al. Aging Cell (2008) PMID: 18843795
273. Huot TJ, et al. Mol. Cell. Biol. (2002) PMID: 12417717
274. Rizos H, et al. J. Biol. Chem. (2001) PMID: 11518711
275. Gombart AF, et al. Leukemia (1997) PMID: 9324288
276. Yang R, et al. Cancer Res. (1995) PMID: 7780957
277. Parry D, et al. Mol. Cell. Biol. (1996) PMID: 8668202
278. Greenblatt MS, et al. Oncogene (2003) PMID: 12606942
279. Yarbrough WG, et al. J. Natl. Cancer Inst. (1999) PMID: 10491434
280. Poi MJ, et al. Mol. Carcinog. (2001) PMID: 11255261
281. Byeon IJ, et al. Mol. Cell (1998) PMID: 9660926
282. Kannengiesser C, et al. Hum. Mutat. (2009) PMID: 19260062
283. Lal G, et al. Genes Chromosomes Cancer (2000) PMID: 10719365
284. Koh J, et al. Nature (1995) PMID: 7777061
285. McKenzie HA, et al. Hum. Mutat. (2010) PMID: 20340136
286. Miller PJ, et al. Hum. Mutat. (2011) PMID: 21462282
287. Kutscher CL, et al. Physiol. Behav. (1977) PMID: 905385
288. Scaini MC, et al. Hum. Mutat. (2014) PMID: 24659262
289. Jenkins NC, et al. J. Invest. Dermatol. (2013) PMID: 23190892
290. Walker GJ, et al. Int. J. Cancer (1999) PMID: 10389768
291. Rutter JL, et al. Oncogene (2003) PMID: 12853981
292. Itahana K, et al. Cancer Cell (2008) PMID: 18538737
293. Zhang Y, et al. Mol. Cell (1999) PMID: 10360174
294. Zhang Y, et al. Cell (1998) PMID: 9529249
295. Whelan AJ, et al. N Engl J Med (1995) PMID: 7666917
296. Adv Exp Med Biol (2010) PMID: 20687502
297. Hogg D, et al. J Cutan Med Surg (1998) PMID: 9479083
298. De Unamuno B, et al. Melanoma Res (2018) PMID: 29543703
299. Soura E, et al. J Am Acad Dermatol (2016) PMID: 26892650
300. Huerta C, et al. Acta Derm Venereol (2018) PMID: 29405243
301. Kaufman DK, et al. Neurology (1993) PMID: 8414022
302. Bahau M, et al. Cancer Res (1998) PMID: 9622062
303. Chan AK, et al. Clin Neuropathol () PMID: 28699883
304. Wiedemeyer WR, et al. Proc. Natl. Acad. Sci. U.S.A. (2010) PMID: 20534551
305. Eguichi T, et al. Mol. Cancer Ther. (2009) PMID: 19509251
306. Jalili A, et al. J. Natl. Cancer Inst. (2012) PMID: 22997239
307. Leonard JH, et al. Cancer Detect. Prev. (2000) PMID: 11198276
308. Vortmeyer AO, et al. Am. J. Clin. Pathol. (1998) PMID: 9825252

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1488563-01

APPENDIX
References

- 9576574
309. Leuraud P, et al. J. Neurooncol. (2000) PMID: 11263500
310. Boström J, et al. Am. J. Pathol. (2001) PMID: 11485924
311. Santarius T, et al. Neuropathol. Appl. Neurobiol. (2000) PMID: 10736068
312. Christacos NC, et al. Genes Chromosomes Cancer (2006) PMID: 16320247
313. Williams EA, et al. JCO Precis Oncol (2020) PMID: 33015533
314. Hossain MG, et al. Endocr. Pathol. (2009) PMID: 19401813
315. Kirsch M, et al. Genes Chromosomes Cancer (2009) PMID: 18973139
316. Sánchez-Aguilera A, et al. Blood (2004) PMID: 14645011
317. Leone PE, et al. Clin. Cancer Res. (2008) PMID: 18829482
318. Boyd KD, et al. Clin. Cancer Res. (2011) PMID: 21994415
319. Walker BA, et al. Blood (2010) PMID: 20616218
320. Morishita A, et al. Hepatology (2004) PMID: 15349907
321. Daa T, et al. APMIS (2008) PMID: 18254776
322. Styring E, et al. Br. J. Cancer (2014) PMID: 24983371
323. Lindberg D, et al. Clin. Endocrinol. (Oxf) (2008) PMID: 17803708
324. Liang JW, et al. PLoS ONE (2014) PMID: 25502777
325. Kim HS, et al. Cancer Res Treat (2014) PMID: 25036575
326. Solomon DA, et al. Cancer Res. (2008) PMID: 18381405
327. Wiedemeyer R, et al. Cancer Cell (2008) PMID: 18394558
328. Arcellana-Panlilio MY, et al. Genes Chromosomes Cancer (2000) PMID: 10918395
329. Hartmann EM, et al. Blood (2010) PMID: 20421449
330. Flordal Thelander E, et al. Leuk. Res. (2007) PMID: 17161458
331. Beà S, et al. Blood (2009) PMID: 18984860
332. Tahara H, et al. J. Bone Miner. Res. (1997) PMID: 9286748
333. Costa-Guda J, et al. Horm Cancer (2013) PMID: 23715670
334. Gluick T, et al. Endocr. Relat. Cancer (2013) PMID: 24127162
335. van Veelen W, et al. Int. J. Cancer (2009) PMID: 18942719
336. Pita JM, et al. J. Clin. Endocrinol. Metab. (2014) PMID: 24423316
337. Kunstman JW, et al. Hum. Mol. Genet. (2015) PMID: 25576899
338. Neta G, et al. Carcinogenesis (2011) PMID: 21642358
339. Barbieri RB, et al. Eur. J. Endocrinol. (2014) PMID: 25565272
340. Hattori H, et al. Pediatr Hematol Oncol (2006) PMID: 16651240
341. Romagnoli S, et al. Am. J. Pathol. (2009) PMID: 19218339
342. Korshunov A, et al. Arch. Pathol. Lab. Med. (2002) PMID: 11800646
343. Ortega S, et al. Biochim. Biophys. Acta (2002) PMID: 11960696
344. Venkataramani R, et al. Nat. Struct. Biol. (1998) PMID: 9437433
345. Li J, et al. Biochemistry (1999) PMID: 10074345
346. Zhang Y, et al. Clin. Cancer Res. (2013) PMID: 23386689
347. Xie Q, et al. Proc. Natl. Acad. Sci. U.S.A. (2012) PMID: 22203985
348. Kentsis A, et al. Nat. Med. (2012) PMID: 22683780
349. Xie Q, et al. Genes Cancer (2013) PMID: 24167653
350. Wanjala J, et al. Mol. Cancer Ther. (2015) PMID: 25381262
351. Cloughesy et al., 2015; ASCO Abstract 2015
352. Koeppen H, et al. Clin. Cancer Res. (2014) PMID: 24687921
353. Patnaik A, et al. Br. J. Cancer (2014) PMID: 24901237
354. Iveson T, et al. Lancet Oncol. (2014) PMID: 24965569
355. Zehir A, et al. Nat. Med. (2017) PMID: 28481359
356. Nguyen B, et al. Cell (2022) PMID: 35120664
357. Guo YF, et al. World J Surg Oncol (2012) PMID: 22741575
358. Garnett J, et al. Neoplasia (2013) PMID: 23359207
359. Kunkel P, et al. Neuro-oncology (2001) PMID: 11296484
360. Stamos J, et al. EMBO J. (2004) PMID: 15167892
361. Tolbert WD, et al. Proc. Natl. Acad. Sci. U.S.A. (2010) PMID: 20624990
362. Löw S, et al. Anticancer Res. () PMID: 19189657
363. Biswas K, et al. J. Biol. Chem. (2013) PMID: 23192342
364. Nat Rev Clin Oncol (2017) PMID: 27245281
365. Duperret EK, et al. Mol Ther (2018) PMID: 29249395
366. Chiappori AA, et al. Ann Oncol (2015) PMID: 25467017
367. Killela PJ, et al. Proc. Natl. Acad. Sci. U.S.A. (2013) PMID: 23530248
368. Killela PJ, et al. Oncotarget (2014) PMID: 24722048
369. Nonoguchi N, et al. Acta Neuropathol. (2013) PMID: 23955565
370. Liu X, et al. Cell Cycle (2013) PMID: 23603989
371. Koelsche C, et al. Acta Neuropathol. (2013) PMID: 24154961
372. Arita H, et al. Acta Neuropathol. (2013) PMID: 23764841
373. Reitman ZJ, et al. Acta Neuropathol. (2013) PMID: 24217890
374. Shay JW, et al. Semin. Cancer Biol. (2011) PMID: 22015685
375. Shay JW, et al. Eur. J. Cancer (1997) PMID: 9282118
376. Kim NW, et al. Science (1994) PMID: 7605428
377. Hanahan D, et al. Cell (2000) PMID: 10647931
378. Horn S, et al. Science (2013) PMID: 23348503
379. Huang FW, et al. Science (2013) PMID: 23348506
380. Vinagre J, et al. Nat Commun (2013) PMID: 23887589
381. Weller M, et al. Nat Rev Clin Oncol (2021) PMID: 33293629
382. Louis DN, et al. Neuro Oncol (2021) PMID: 34185076
383. Metzgeroth G, et al. Leukemia (2007) PMID: 17377585
384. Debiec-Rychter M, et al. Eur. J. Cancer (2006) PMID: 16624552
385. Kamenz T, et al. World J. Gastroenterol. (2006) PMID: 16570351
386. Wang YY, et al. Proc. Natl. Acad. Sci. U.S.A. (2005) PMID: 15650049
387. Hassler MR, et al. Springerplus (2014) PMID: 25674429
388. Razis E, et al. Clin. Cancer Res. (2009) PMID: 19789313
389. Reardon DA, et al. Cancer (2012) PMID: 22371319
390. Carvajal RD, et al. Clin. Cancer Res. (2015) PMID: 25695690
391. Hochhaus A, et al. J. Cancer Res. Clin. Oncol. (2015) PMID: 26002753
392. Blay JY, et al. Lancet Oncol. (2015) PMID: 25882987
393. Kajimoto N, et al. Int J Clin Exp Pathol (2015) PMID: 26722383
394. Sako H, et al. PLoS ONE (2014) PMID: 25221952
395. Hughes TP, et al. Blood (2014) PMID: 24335106
396. Takahashi N, et al. Biomark Res (2014) PMID: 24650752
397. Reichardt P, et al. Ann. Oncol. (2012) PMID: 22357255
398. Cauchi C, et al. Cancer Chemother. Pharmacol. (2012) PMID: 22119758
399. Villar VH, et al. PLoS ONE (2012) PMID: 22662203
400. Quintás-Cardama A, et al. Nat Clin Pract Oncol (2008) PMID: 18936790
401. Bisagni G, et al. J Thorac Oncol (2009) PMID: 19461405
402. Handolias D, et al. Br. J. Cancer (2010) PMID: 20372153
403. Dişel U, et al. Lung Cancer (2011) PMID: 20970876
404. Park SH, et al. Invest New Drugs (2012) PMID: 22270258
405. Catania C, et al. Onco Targets Ther (2014) PMID: 24855380
406. Guo T, et al. Clin. Cancer Res. (2007) PMID: 17699867
407. Hu S, et al. Mol. Cancer Ther. (2008) PMID: 18483300
408. Fumagalli et al., 2012; ESMO Abstract 1491P
409. Zustovich et al., 2013; 23898124; Reardon et al.
410. Lee EQ, et al. Neuro-oncology (2012) PMID: 23099651
411. Peereboom DM, et al. Neuro-oncology (2013) PMID: 23328813
412. Hottinger AF, et al. Br. J. Cancer (2014) PMID: 24786603
413. Karajannis MA, et al. Neuro-oncology (2014) PMID: 24803676
414. Heinrich MC, et al. J. Clin. Oncol. (2008) PMID: 18955458
415. Buchbinder EI, et al. Cancer (2015) PMID: 26264378
416. Reichardt P, et al. BMC Cancer (2016) PMID: 26772734
417. Hirai F, et al. Mol Clin Oncol (2016) PMID: 27073655
418. Goemans BF, et al. Leuk. Res. (2010) PMID: 20435347
419. Pan E, et al. J. Neurooncol. (2012) PMID: 22832897
420. Kreisl TN, et al. J. Neurooncol. (2013) PMID: 23086433
421. Balaña C, et al. Target Oncol (2014) PMID: 24424564
422. Hutterer M, et al. Neuro-oncology (2014) PMID: 24311637

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 07 November 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531