



## Sample Information

**Patient Name:** 陸玟臻  
**Gender:** Female  
**ID No.:** F220100214  
**History No.:** 43240217  
**Age:** 50

**Ordering Doctor:** DOC1242E 劉峻宇  
**Ordering REQ.:** H3LF8KK  
**Signing in Date:** 2021/11/04

**Path No.:** S110-99947  
**MP No.:** F21093  
**Assay:** Oncomine Focus Assay  
**Sample Type:** FFPE  
**Block No.:** S110-35808A  
**Percentage of tumor cells:** 30%

**Reporting Doctor:** DOC5466K 葉奕成 (Pho

**Note:**

## Sample Cancer Type: Non-Small Cell Lung Cancer

Table of Contents	Page
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2
Biomarker Descriptions	2
Relevant Therapy Summary	4
Relevant Therapy Details	5
Clinical Trials Summary	23
Alert Details	24

**Report Highlights**  
2 Relevant Biomarkers  
16 Therapies Available  
15 Clinical Trials

## Relevant Non-Small Cell Lung Cancer Variants

Gene	Finding	Gene	Finding
ALK	None detected	NTRK1	None detected
BRAF	None detected	NTRK2	None detected
EGFR	<b>EGFR p.(L858R) c.2573T&gt;G</b>	NTRK3	None detected
ERBB2	None detected	RET	None detected
KRAS	None detected	ROS1	None detected
MET	None detected		

## Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	<b>EGFR p.(L858R) c.2573T&gt;G</b> epidermal growth factor receptor Allele Frequency: 25.93%  <b>Prognostic significance:</b> None <b>Diagnostic significance:</b> None	<b>afatinib</b> <sup>1,2</sup> <b>bevacizumab* + erlotinib</b> <sup>2</sup> <b>dacomitinib</b> <sup>1,2</sup> <b>erlotinib</b> <sup>1,2</sup> <b>erlotinib + ramucirumab</b> <sup>1,2</sup> <b>gefitinib</b> <sup>1,2</sup> <b>osimertinib</b> <sup>1,2</sup> afatinib + cetuximab atezolizumab + bevacizumab + chemotherapy bevacizumab + gefitinib gefitinib + chemotherapy osimertinib + chemotherapy	None	15
IIC	<b>AR amplification</b> androgen receptor  <b>Prognostic significance:</b> None <b>Diagnostic significance:</b> None	None	hormone therapy	0

Public data sources included in relevant therapies: FDA<sup>1</sup>, NCCN, EMA<sup>2</sup>, ESMO

Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

**Tier Reference:** Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

\* Includes biosimilars

## Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

### DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
EGFR	p.(L858R)	c.2573T>G	COSM6224	chr7:55259515	25.93%	NM_005228.5	missense	1994

### Copy Number Variations

Gene	Locus	Copy Number
AR	chrX:66776186	9.7

## Biomarker Descriptions

### AR (androgen receptor)

**Background:** The AR gene encodes the androgen receptor protein (AR), a ligand-activated transcription factor regulated by the binding of the hormones testosterone and dihydrotestosterone<sup>1,2</sup>. Hormone binding to AR results in receptor dimerization, nuclear translocation, and target gene transcription, thus activating the RAS/RAF/MEK/ERK and PI3K/AKT/MTOR signaling pathways, which promote cell proliferation and survival<sup>2,3,4</sup>.

**Alterations and prevalence:** Alterations in AR function can result from overexpression, gene amplification, or mutations. AR mutations, including L702H, W742C/L, H875Y, and T878A, are commonly observed in 10-30% of castration-resistant prostate cancer and result in decreased ligand specificity, allowing other nuclear hormones to activate AR<sup>5</sup>. Androgen receptor splice variants have been reported in castration resistant prostate cancer<sup>6,7</sup>. The androgen receptor splice variant 7 (AR-V7) is a result of aberrant mRNA splicing of AR exons 1-3 and a cryptic exon 3, resulting in the expression of a constitutively active protein<sup>7</sup>.

## Biomarker Descriptions (continued)

**Potential relevance:** The FDA has granted fast track designation (2016) to seviteronel for AR-positive triple-negative breast cancer (TNBC) patients<sup>8</sup>. Androgen deprivation therapy (ADT) such as abiraterone<sup>9</sup> (2011) and enzalutamide<sup>10</sup> (2011) are FDA approved for use in locally advanced and metastatic prostate cancers. Although many men initially respond to ADT, most will develop hormone resistance. Resistance to ADT is also associated with other aberrations of the AR gene including mutations within the ligand binding domain and gene amplification<sup>5,11,12</sup>. The androgen receptor splice variant, AR-V7, lacks the ligand binding domain, resulting in constitutive activation and is associated with resistance to androgen deprivation therapy (ADT) in advanced prostate cancer<sup>6</sup>.

### EGFR (epidermal growth factor receptor)

**Background:** The EGFR gene encodes the epidermal growth factor receptor (EGFR) tyrosine kinase, a member of the ERBB/human epidermal growth factor receptor (HER) family. In addition to EGFR/ERBB1/HER1, other members of the ERBB/HER family include ERBB2/HER2, ERBB3/HER3, and ERBB4/HER4<sup>13</sup>. EGFR ligand induced dimerization results in kinase activation and leads to stimulation of oncogenic signaling pathways including the PI3K/AKT/MTOR and RAS/RAF/MEK/ERK pathways. Activation of these pathways promote cell proliferation, differentiation, and survival<sup>14,15</sup>.

**Alterations and prevalence:** Recurrent somatic mutations in the tyrosine kinase domain (TKD) of EGFR are observed in approximately 10-20% of lung adenocarcinoma, and at higher frequencies in never-smoker, female, and Asian populations<sup>16,17,18,19</sup>. The most common mutations occur near the ATP-binding pocket of the TKD and include short in-frame deletions in exon 19 (EGFR exon 19 deletion) and the L858R amino acid substitution in exon 21<sup>20</sup>. These mutations constitutively activate EGFR resulting in downstream signaling, and represent 80% of the EGFR mutations observed in lung cancer. A second group of less prevalent activating mutations include E709K, G719X, S768I, L861Q, and short in-frame insertion mutations in exon 20<sup>21,22,23,24</sup>. EGFR activating mutations in lung cancer tend to be mutually exclusive to KRAS activating mutations<sup>25</sup>. In contrast, a different set of recurrent activating EGFR mutations in the extracellular domain include R108K, A289V and G598V and are primarily observed in glioblastoma<sup>20,26</sup>. Amplification of EGFR is observed in several cancer types including 30% of glioblastoma, 12% of esophageal cancer, 10% of head and neck cancer, 5% of bladder cancer, and 5% of lung squamous cell carcinoma<sup>17,18,19,26,27</sup>. Deletion of exons 2-7, encoding the extracellular domain of EGFR (EGFRvIII), results in overexpression of a ligand-independent constitutively active protein and is observed in approximately 30% of glioblastoma<sup>28,29,30</sup>.

**Potential relevance:** Approved first-generation EGFR tyrosine kinase inhibitors (TKIs) include erlotinib<sup>31</sup> (2004) and gefitinib<sup>32</sup> (2015), which block the activation of downstream signaling by reversible interaction with the ATP-binding site. Although initially approved for advanced lung cancer, the discovery that drug sensitivity was associated with exon 19 and exon 21 activating mutations allowed first-generation TKIs to become subsequently approved for front-line therapy in lung cancer tumors containing exon 19 or exon 21 activating mutations. Second-generation TKIs afatinib<sup>33</sup> (2013) and dacomitinib<sup>34</sup> (2018) bind EGFR and other ERBB/HER gene family members irreversibly and were subsequently approved. First- and second-generation TKIs afatinib, dacomitinib, erlotinib, and gefitinib are recommended for the treatment NSCLC harboring EGFR exon 19 insertions, exon 19 deletions, point mutations L861Q, L858R, S768I, and codon 719 mutations, whereas most EGFR exon 20 insertions, except p.A763\_Y764insFQEA, confer resistance to the same therapies<sup>35,36,37,38</sup>. In lung cancer containing EGFR exon 19 or 21 activating mutations, treatment with TKIs is eventually associated with the emergence of drug resistance<sup>39</sup>. The primary resistance mutation that emerges following treatment with first-generation TKI is T790M, accounting for 50-60% of resistant cases<sup>20</sup>. Third generation TKIs were developed to maintain sensitivity in the presence of T790M. Osimertinib<sup>40</sup> (2015) is an irreversible inhibitor indicated for metastatic EGFR T790M positive lung cancer and for the first-line treatment of metastatic NSCLC containing EGFR exon 19 deletions or exon 21 L858R mutations. Like first-generation TKIs, treatment with osimertinib is associated with acquired resistance. In this case, resistance is associated with the C797S mutation, and occurs in 22-44% of cases<sup>39</sup>. The T790M and C797S mutations may be each selected following sequential treatment with a first-generation TKI followed by a third-generation TKI or vice versa<sup>41</sup>. T790M and C797S can occur in either cis or trans allelic orientation<sup>41</sup>. If C797S is observed following progression after treatment with a third-generation TKI in the first-line setting, sensitivity may be retained to first-generation TKIs<sup>41</sup>. If C797S co-occurs in trans with T790M following sequential treatment with first- and third-generation TKIs, patients may exhibit sensitivity to combination first- and third-generation TKIs, but resistance to third-generation TKIs alone<sup>41,42</sup>. However, C797S occurring in cis conformation with T790M, confers resistance to first- and third-generation TKIs<sup>41</sup>. Fourth-generation TKIs are in development to overcome acquired C797S and T790M resistance mutations after osimertinib treatment. EGFR targeting antibodies including cetuximab (2004), panitumumab (2006), and necitumumab (2016) are under investigation in combination with EGFR-targeting TKIs for efficacy against EGFR mutations. The bispecific antibody, JNJ-61186372<sup>43</sup>, targeting EGFR and MET, and the TKI mobocertinib<sup>44</sup>, each received a breakthrough designation from the FDA (2020) for NSCLC tumors harboring EGFR exon 20 insertion mutations. The Oncoprex immunogene therapy CNVN-202<sup>45</sup> in combination with osimertinib received a fast track designation from the FDA (2020) for NSCLC tumors harboring EGFR mutations that progressed on osimertinib alone. BDTX-189<sup>46</sup> was granted a fast track designation (2020) for the treatment of solid tumors harboring an EGFR exon 20 insertion mutation.

## Relevant Therapy Summary

● In this cancer type

○ In other cancer type

● In this cancer type and other cancer types

✕ No evidence

### EGFR p.(L858R) c.2573T>G

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
osimertinib	●	●	●	●	● (III)
afatinib	●	●	●	●	✕
dacomitinib	●	●	●	●	✕
erlotinib	●	●	●	●	✕
erlotinib + ramucirumab	●	●	●	●	✕
gefitinib	●	●	●	●	✕
bevacizumab + erlotinib	✕	●	●	●	✕
afatinib + cetuximab	✕	●	✕	✕	✕
osimertinib + chemotherapy	✕	●	✕	✕	✕
osimertinib + chemotherapy + surgical intervention	✕	●	✕	✕	✕
bevacizumab (Allergan) + erlotinib	✕	✕	●	✕	✕
bevacizumab (Fujifilm Kyowa Kirin Biologics) + erlotinib	✕	✕	●	✕	✕
bevacizumab (Mabxience) + erlotinib	✕	✕	●	✕	✕
bevacizumab (Pfizer) + erlotinib	✕	✕	●	✕	✕
bevacizumab (Samsung Bioepis) + erlotinib	✕	✕	●	✕	✕
atezolizumab + bevacizumab + carboplatin + paclitaxel	✕	✕	✕	●	✕
bevacizumab + gefitinib	✕	✕	✕	●	✕
gefitinib + carboplatin + pemetrexed	✕	✕	✕	●	✕
amivantamab, lazertinib, osimertinib	✕	✕	✕	✕	● (III)
osimertinib, chemotherapy	✕	✕	✕	✕	● (III)
atezolizumab, bevacizumab, chemotherapy	✕	✕	✕	✕	● (II)
bevacizumab, erlotinib	✕	✕	✕	✕	● (II)
datopotamab deruxtecan	✕	✕	✕	✕	● (II)
durvalumab, tremelimumab, chemotherapy	✕	✕	✕	✕	● (II)
osimertinib, savolitinib	✕	✕	✕	✕	● (II)
patritumab deruxtecan	✕	✕	✕	✕	● (II)
DZD-9008	✕	✕	✕	✕	● (I/II)
amivantamab, lazertinib	✕	✕	✕	✕	● (I)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

**Disclaimer:** The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2021.09(004).

## Relevant Therapy Summary (continued)

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

### EGFR p.(L858R) c.2573T>G (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
lazertinib, amivantamab	×	×	×	×	● (I)
telisotuzumab vedotin, osimertinib	×	×	×	×	● (I)
TNO-155, nazartinib	×	×	×	×	● (I)

### AR amplification

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
androgen receptor therapy	×	○	×	×	×
bicalutamide	×	○	×	×	×
leuprorelin	×	○	×	×	×

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Details

### Current FDA Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

FDA information is current as of 2021-08-18. For the most up-to-date information, search [www.fda.gov](https://www.fda.gov).

### EGFR p.(L858R) c.2573T>G

#### ● afatinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2019-10-11

Variant class: EGFR L858R mutation

#### Indications and usage:

GILOTRIF® is a kinase inhibitor indicated for:

- First-line treatment of patients with metastatic non-small cell lung cancer (NSCLC) whose tumors have non-resistant epidermal growth factor receptor (EGFR) mutations as detected by an FDA-approved test.

Limitation of Use: Safety and efficacy of GILOTRIF® were not established in patients whose tumors have resistant EGFR mutations

- Treatment of patients with metastatic, squamous NSCLC progressing after platinum-based chemotherapy

#### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2019/201292s015lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/201292s015lbl.pdf)

## EGFR p.(L858R) c.2573T>G (continued)

### ● dacomitinib

**Cancer type:** Non-Small Cell Lung Cancer

**Label as of:** 2020-12-18

**Variant class:** EGFR L858R mutation

**Indications and usage:**

VIZIMPRO® is a kinase inhibitor indicated for the first-line treatment of patients with metastatic non-small cell lung cancer (NSCLC) with epidermal growth factor receptor (EGFR) exon 19 deletion or exon 21 L858R substitution mutations as detected by an FDA-approved test.

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/211288s003lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/211288s003lbl.pdf)

### ● erlotinib

**Cancer type:** Non-Small Cell Lung Cancer

**Label as of:** 2016-10-18

**Variant class:** EGFR L858R mutation

**Indications and usage:**

TARCEVA® is a kinase inhibitor indicated for:

- The treatment of patients with metastatic non-small cell lung cancer (NSCLC) whose tumors have epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by an FDA-approved test receiving first-line, maintenance, or second or greater line treatment after progression following at least one prior chemotherapy regimen.
- First-line treatment of patients with locally advanced, unresectable or metastatic pancreatic cancer, in combination with gemcitabine.

**Limitations of Use:**

- Safety and efficacy of TARCEVA® have not been established in patients with NSCLC whose tumors have other EGFR mutations.
- TARCEVA® is not recommended for use in combination with platinum-based chemotherapy.

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2016/021743s025lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2016/021743s025lbl.pdf)

### ● erlotinib + ramucirumab

**Cancer type:** Non-Small Cell Lung Cancer

**Label as of:** 2021-06-15

**Variant class:** EGFR L858R mutation

**Indications and usage:**

CYRAMZA® is a human vascular endothelial growth factor receptor 2 (VEGFR2) antagonist indicated:

- as a single agent or in combination with paclitaxel, for treatment of advanced or metastatic gastric or gastro-esophageal junction adenocarcinoma with disease progression on or after prior fluoropyrimidine- or platinum-containing chemotherapy.
- in combination with erlotinib, for first-line treatment of metastatic non-small cell lung cancer with epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) mutations.
- in combination with docetaxel, for treatment of metastatic non-small cell lung cancer with disease progression on or after platinum-based chemotherapy. Patients with EGFR or ALK genomic tumor aberrations should have disease progression on FDA-approved therapy for these aberrations prior to receiving CYRAMZA®.
- in combination with FOLFIRI, for the treatment of metastatic colorectal cancer with disease progression on or after prior therapy with bevacizumab, oxaliplatin, and a fluoropyrimidine.
- as a single agent, for the treatment of hepatocellular carcinoma in patients who have an alpha fetoprotein of  $\geq 400$  ng/mL and have been treated with sorafenib.

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/125477s039lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125477s039lbl.pdf)

**EGFR p.(L858R) c.2573T>G (continued)****● gefitinib****Cancer type:** Non-Small Cell Lung Cancer**Label as of:** 2021-05-05**Variant class:** EGFR L858R mutation**Indications and usage:**

IRESSA® is a tyrosine kinase inhibitor indicated for the first-line treatment of patients with metastatic non-small cell lung cancer (NSCLC) whose tumors have epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by an FDA-approved test.

Limitation of Use: Safety and efficacy of IRESSA® have not been established in patients whose tumors have EGFR mutations other than exon 19 deletions or exon 21 (L858R) substitution mutations.

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/206995s004lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/206995s004lbl.pdf)

**● osimertinib****Cancer type:** Non-Small Cell Lung Cancer**Label as of:** 2021-07-26**Variant class:** EGFR L858R mutation**Indications and usage:**

TAGRISSO® is a kinase inhibitor indicated for:

- as adjuvant therapy after tumor resection in adult patients with non-small cell lung cancer (NSCLC) whose tumors have epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test.
- the first-line treatment of adult patients with metastatic NSCLC whose tumors have EGFR exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test.
- the treatment of adult patients with metastatic EGFR T790M mutation positive NSCLC, as detected by an FDA-approved test, whose disease has progressed on or after EGFR TKI therapy.

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/208065s022lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/208065s022lbl.pdf)

## Current NCCN Information

- ☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

NCCN information is current as of 2021-08-02. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org).  
For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

### EGFR p.(L858R) c.2573T>G

#### ● afatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

#### ● dacomitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

#### ● erlotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

#### ● gefitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]



**EGFR p.(L858R) c.2573T>G (continued)****● osimertinib****Cancer type:** Non-Small Cell Lung Cancer**Variant class:** EGFR L858R mutation**NCCN Recommendation category:** 1**Population segment (Line of therapy):**

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Preferred intervention

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]**● afatinib****Cancer type:** Non-Small Cell Lung Cancer**Variant class:** EGFR L858R mutation**NCCN Recommendation category:** 2A**Population segment (Line of therapy):**

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (Subsequent therapy)

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]**● afatinib + cetuximab****Cancer type:** Non-Small Cell Lung Cancer**Variant class:** EGFR L858R mutation**NCCN Recommendation category:** 2A**Population segment (Line of therapy):**

- Progression (Subsequent therapy)

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]**● bevacizumab + erlotinib****Cancer type:** Non-Small Cell Lung Cancer**Variant class:** EGFR L858R mutation**NCCN Recommendation category:** 2A**Population segment (Line of therapy):**

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Other recommended intervention
- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (Subsequent therapy)

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

## EGFR p.(L858R) c.2573T>G (continued)

### ● dacomitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

### ● erlotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

### ● erlotinib + ramucirumab

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Other recommended intervention
- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

### ● gefitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

## EGFR p.(L858R) c.2573T>G (continued)

### ● osimertinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Brain Metastases, Leptomeningeal Metastases, Spine Metastases (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 1.2021]

### ● osimertinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (Subsequent therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

### ● osimertinib + chemotherapy

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Stage IIB, Stage IIIA, Stage IIIB (Adjuvant therapy)
- Stage IIIA; Resectable (Adjuvant therapy)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

### ● osimertinib + chemotherapy + surgical intervention

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Stage IIB (Adjuvant therapy)
- Stage IIIA; Resectable (Adjuvant therapy)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

## EGFR p.(L858R) c.2573T>G (continued)

### ● erlotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Leptomeningeal Metastases, Spine Metastases (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 1.2021]

### ● erlotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFRi sensitizing mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Brain Metastases (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 1.2021]

### ● afatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFRi sensitizing mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Brain Metastases (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 1.2021]

### ● gefitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFRi sensitizing mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Brain Metastases (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 1.2021]

## AR amplification

### ○ androgen receptor therapy

Cancer type: Head and Neck Cancer

Variant class: AR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Salivary Gland Neoplasm; Recurrent, Unresectable, Distant Metastases (Line of therapy not specified); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Head and Neck Cancers [Version 3.2021]

### ○ bicalutamide

Cancer type: Head and Neck Cancer

Variant class: AR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Salivary Gland Neoplasm; Recurrent, Unresectable, Distant Metastases (Line of therapy not specified); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Head and Neck Cancers [Version 3.2021]

### ○ leuprorelin

Cancer type: Head and Neck Cancer

Variant class: AR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Salivary Gland Neoplasm; Recurrent, Unresectable, Distant Metastases (Line of therapy not specified); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Head and Neck Cancers [Version 3.2021]

## Current EMA Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

EMA information is current as of 2021-08-18. For the most up-to-date information, search [www.ema.europa.eu/ema](https://www.ema.europa.eu/ema).

### EGFR p.(L858R) c.2573T>G

#### ● afatinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2021-04-21

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/giotrif-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/giotrif-epar-product-information_en.pdf)

#### ● bevacizumab (Allergan) + erlotinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2021-05-21

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/mvasi-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/mvasi-epar-product-information_en.pdf)

#### ● bevacizumab (Fujifilm Kyowa Kirin Biologics) + erlotinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2021-06-23

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/equidacent-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/equidacent-epar-product-information_en.pdf)

#### ● bevacizumab (Mabxience) + erlotinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2021-08-11

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/alymsys-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/alymsys-epar-product-information_en.pdf)

#### ● bevacizumab (Pfizer) + erlotinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2021-07-07

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/zirabev-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/zirabev-epar-product-information_en.pdf)

#### ● bevacizumab (Samsung Bioepis) + erlotinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2021-05-18

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/onbevzi-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/onbevzi-epar-product-information_en.pdf)

**EGFR p.(L858R) c.2573T>G (continued)****● bevacizumab (Samsung Bioepis) + erlotinib**

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2021-06-21

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/aybintio-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/aybintio-epar-product-information_en.pdf)**● bevacizumab + erlotinib**

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2021-01-28

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/avastin-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/avastin-epar-product-information_en.pdf)**● dacomitinib**

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2021-07-21

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/vizimpro-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/vizimpro-epar-product-information_en.pdf)**● erlotinib**

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2019-04-24

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/documents/product-information/tarceva-epar-product-information\\_en.pdf](https://www.ema.europa.eu/documents/product-information/tarceva-epar-product-information_en.pdf)**● erlotinib + ramucirumab**

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2020-07-02

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/cyramza-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/cyramza-epar-product-information_en.pdf)**● gefitinib**

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2021-03-05

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/iressa-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/iressa-epar-product-information_en.pdf)**● osimertinib**

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2021-07-01

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/tagrisso-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/tagrisso-epar-product-information_en.pdf)

## Current ESMO Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

ESMO information is current as of 2021-08-02. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

### EGFR p.(L858R) c.2573T>G

#### ● atezolizumab + bevacizumab + carboplatin + paclitaxel

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Non-squamous Cell; Metastatic (First-line therapy); ESMO-MCBS v1.1 score: 3
- Metastatic (Second-line therapy)

**Reference:** ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

#### ● afatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFRi sensitizing mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Advanced (First-line therapy)

**Reference:** ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

#### ● erlotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFRi sensitizing mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Advanced (First-line therapy)

**Reference:** ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]



## EGFR p.(L858R) c.2573T>G (continued)

### ● gefitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFRi sensitizing mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Advanced (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● osimertinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFRi sensitizing mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Advanced (First-line therapy); ESMO-MCBS v1.1 score: 4

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● dacomitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFRi sensitizing mutation

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Advanced (First-line therapy); ESMO-MCBS v1.1 score: 3

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● erlotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFRi sensitizing mutation

ESMO Level of Evidence/Grade of Recommendation: III / B

Population segment (Line of therapy):

- Non-squamous Cell (Maintenance therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

## EGFR p.(L858R) c.2573T>G (continued)

### ● afatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● bevacizumab + erlotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● bevacizumab + gefitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● dacomitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

## EGFR p.(L858R) c.2573T>G (continued)

### ● erlotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● erlotinib + ramucirumab

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● gefitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● gefitinib + carboplatin + pemetrexed

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

## EGFR p.(L858R) c.2573T>G (continued)

### ● bevacizumab + erlotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Stage IV (First-line therapy); ESMO-MCBS v1.1 score: 3

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● bevacizumab + gefitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Stage IV (First-line therapy); ESMO-MCBS v1.1 score: 3

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● erlotinib + ramucirumab

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Stage IV (First-line therapy); ESMO-MCBS v1.1 score: 3

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● gefitinib + carboplatin + pemetrexed

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Advanced (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

## EGFR p.(L858R) c.2573T>G (continued)

### ● afatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● bevacizumab + erlotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● bevacizumab + gefitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● dacomitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

## EGFR p.(L858R) c.2573T>G (continued)

### ● erlotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● erlotinib + ramucirumab

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● gefitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● gefitinib + carboplatin + pemetrexed

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

## Clinical Trials in Taiwan region:

### Clinical Trials Summary

#### EGFR p.(L858R) c.2573T>G

NCT ID	Title	Phase
NCT04487080	A Phase III, Randomized Study of Amivantamab and Lazertinib Combination Therapy Versus Osimertinib Versus Lazertinib as First-Line Treatment in Patients With EGFR-Mutated Locally Advanced or Metastatic Non-Small Cell Lung Cancer.	III
NCT03521154	A Phase III, Randomized, Double-blind, Placebo-controlled, Multicenter, International Study of Osimertinib as Maintenance Therapy in Patients With Locally Advanced, Unresectable EGFR Mutation-positive Non-Small Cell Lung Cancer (Stage III) Whose Disease Has Not Progressed Following Definitive Platinum-based Chemoradiation Therapy (LAURA)	III
NCT04035486	A Phase III, Open-label, Randomized Study of Osimertinib With or Without Platinum Plus Pemetrexed Chemo, as First-line Treatment in Patients With Epidermal Growth Factor Receptor (EGFR) Mutation Positive, Locally Advanced or Metastatic Non-small Cell Lung Cancer (FLAURA2)	III
NCT04351555	A Phase III, Randomised, Controlled, Multi-center, 3-Arm Study of Neoadjuvant Osimertinib as Monotherapy or in Combination With Chemotherapy Versus Standard of Care Chemotherapy Alone for the Treatment of Patients With Epidermal Growth Factor Receptor Mutation Positive, Resectable Non-small Cell Lung Cancer	III
NCT04147351	A Phase II Study of Atezolizumab in Combination With Bevacizumab, Carboplatin or Cisplatin, and Pemetrexed for EGFR-mutant Metastatic Non-small Cell Lung Cancer Patients After Failure of EGFR Tyrosine Kinase Inhibitors.	II
NCT02655536	A Phase II, Open Label, Multicenter Study of Bevacizumab in Combination With Erlotinib Versus Erlotinib Alone in Patients With EGFR Mutant Non-small Cell Lung Cancer Who Have Brain Metastases	II
NCT03994393	A Phase II Trial of Durvalumab (MEDI4736) and Tremelimumab With Chemotherapy in Metastatic EGFR Mutant Non-squamous Non-small Cell Lung Cancer (NSCLC) Following Progression on EGFR Tyrosine Kinase Inhibitors (TKIs)	II
NCT03778229	A Phase II, Single Arm Study Assessing Efficacy of Osimertinib With Savolitinib in Patients With EGFRm + MET+, Locally Advanced or Metastatic Non Small Cell Lung Cancer Who Have Progressed Following Osimertinib Treatment (SAVANNAH Study)	II
NCT04619004	HERTHENA-Lung01: A Phase II Randomized Open-Label Study of Patritumab Deruxtecan (U3-1402) in Subjects With Previously Treated Metastatic or Locally Advanced EGFR-mutated Non-Small Cell Lung Cancer (NSCLC)	II
NCT02609776	A Phase I, First-in-Human, Open-Label, Dose Escalation Study of JNJ-61186372, a Human Bispecific EGFR and cMet Antibody, in Subjects With Advanced Non-Small Cell Lung Cancer.	I
NCT04077463	An Open-label Phase I/Ib Study to Evaluate the Safety and Pharmacokinetics of JNJ-73841937 (Lazertinib), a Third Generation EGFR-TKI, as Monotherapy or in Combinations With JNJ-61186372, a Human Bispecific EGFR and cMet Antibody in Participants With Advanced Non-Small Cell Lung Cancer	I
NCT02099058	A Multicenter, Phase I/Ib, Open-Label, Dose-Escalation Study of ABBV-399, an Antibody Drug Conjugate, in Subjects With Advanced Solid Tumors	I
NCT04484142	Phase II, Single-arm, Open-label Study of DS-1062a in Advanced or Metastatic Non-small Cell Lung Cancer With Actionable Genomic Alterations and Progressed on or After Kinase Inhibitor Therapy and Platinum Based Chemotherapy (TROPION-Lung05)	II
NCT03114319	An Open-label, Multi-center, Phase I, Dose Finding Study of Oral TNO155 in Adult Patients With Advanced Solid Tumors.	I

## Clinical Trials Summary (continued)

### EGFR p.(L858R) c.2573T>G (continued)

NCT ID	Title	Phase
NCT03974022	A Phase I/II, Open-Label, Multicenter Study to Assess the Safety, Tolerability, Pharmacokinetics and Anti-tumor Efficacy of DZD9008 in Patients With Advanced Non-Small Cell Lung Cancer (NSCLC) With EGFR or HER2 Mutation	I/II

## Alerts Informed By Public Data Sources

### Current FDA Information

 Contraindicated
  Not recommended
  Resistance
  Breakthrough
  Fast Track

FDA information is current as of 2021-08-18. For the most up-to-date information, search [www.fda.gov](https://www.fda.gov).

### EGFR p.(L858R) c.2573T>G

#### osimertinib + quaratusugene ozeplasmid

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR mutation

##### Supporting Statement:

The FDA has granted Fast Track Designation to the immunogene therapy, quaratusugene ozeplasmid, in combination with EGFR inhibitor osimertinib for the treatment of non-small cell lung cancer (NSCLC) with EFGR mutations that progressed after treatment with osimertinib alone.

##### Reference:

<https://www.genprex.com/news/genprex-receives-u-s-fda-fast-track-designation-for-gene-therapy-that-targets-lung-cancer/>

### AR amplification

#### seviteronel

**Cancer type:** Triple Negative Breast Cancer

**Variant class:** AR positive

##### Supporting Statement:

The FDA has granted Fast Track Designation to the small-molecule CYP17 lyase-selective inhibitor, seviteronel, for:

- Androgen receptor (AR) positive advanced triple negative breast cancer (TNBC).
- Estrogen receptor (ER) positive advanced breast cancer.

##### Reference:

<https://www.businesswire.com/news/home/20160106006206/en/Innocrin-Pharmaceuticals-Granted-Fast-Track-Designation-FDA>



## Current NCCN Information

 Contraindicated
  Not recommended
  Resistance
  Breakthrough
  Fast Track

NCCN information is current as of 2021-08-02. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org).  
For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

### EGFR p.(L858R) c.2573T>G

#### alectinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFRi sensitizing mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Crizotinib, ceritinib, alectinib, brigatinib, or lorlatinib are not recommended as subsequent therapy for patients with sensitizing EGFR mutations who relapse on EGFR TKI therapy."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

#### brigatinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFRi sensitizing mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Crizotinib, ceritinib, alectinib, brigatinib, or lorlatinib are not recommended as subsequent therapy for patients with sensitizing EGFR mutations who relapse on EGFR TKI therapy."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

#### ceritinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFRi sensitizing mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Crizotinib, ceritinib, alectinib, brigatinib, or lorlatinib are not recommended as subsequent therapy for patients with sensitizing EGFR mutations who relapse on EGFR TKI therapy."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

#### crizotinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFRi sensitizing mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Crizotinib, ceritinib, alectinib, brigatinib, or lorlatinib are not recommended as subsequent therapy for patients with sensitizing EGFR mutations who relapse on EGFR TKI therapy."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

## EGFR p.(L858R) c.2573T>G (continued)

### – lorlatinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFRi sensitizing mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Crizotinib, ceritinib, alectinib, brigatinib, or lorlatinib are not recommended as subsequent therapy for patients with sensitizing EGFR mutations who relapse on EGFR TKI therapy."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

### – atezolizumab

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "subsequent therapy with pembrolizumab, nivolumab, or atezolizumab is not recommended in patients with EGFR mutations or ALK fusions."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

### – nivolumab

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "subsequent therapy with pembrolizumab, nivolumab, or atezolizumab is not recommended in patients with EGFR mutations or ALK fusions."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

### – pembrolizumab

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "subsequent therapy with pembrolizumab, nivolumab, or atezolizumab is not recommended in patients with EGFR mutations or ALK fusions."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

## Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

## References

1. Lu et al. International Union of Pharmacology. LXV. The pharmacology and classification of the nuclear receptor superfamily: glucocorticoid, mineralocorticoid, progesterone, and androgen receptors. *Pharmacol. Rev.* 2006 Dec;58(4):782-97. PMID: 17132855
2. Davey et al. Androgen Receptor Structure, Function and Biology: From Bench to Bedside. *Clin Biochem Rev.* 2016 Feb;37(1):3-15. PMID: 27057074
3. Crumbaker et al. AR Signaling and the PI3K Pathway in Prostate Cancer. *Cancers (Basel).* 2017 Apr 15;9(4). PMID: 28420128
4. Leung et al. Non-Genomic Actions of the Androgen Receptor in Prostate Cancer. *Front Endocrinol (Lausanne).* 2017 Jan 17;8:2. PMID: 28144231
5. Waltering et al. Androgen receptor (AR) aberrations in castration-resistant prostate cancer. *Mol. Cell. Endocrinol.* 2012 Sep 5;360(1-2):38-43. PMID: 22245783
6. Antonarakis et al. AR-V7 and resistance to enzalutamide and abiraterone in prostate cancer. *N. Engl. J. Med.* 2014 Sep 11;371(11):1028-38. PMID: 25184630
7. Zhu et al. Novel Junction-specific and Quantifiable In Situ Detection of AR-V7 and its Clinical Correlates in Metastatic Castration-resistant Prostate Cancer. *Eur. Urol.* 2018 May;73(5):727-735. PMID: 28866255
8. <https://www.businesswire.com/news/home/20160106006206/en/Innocrin-Pharmaceuticals-Granted-Fast-Track-Designation-FDA>
9. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/202379s031s033lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/202379s031s033lbl.pdf)
10. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/203415s016lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/203415s016lbl.pdf)
11. Lallous et al. Functional analysis of androgen receptor mutations that confer anti-androgen resistance identified in circulating cell-free DNA from prostate cancer patients. *Genome Biol.* 2016 Jan 26;17:10. PMID: 26813233
12. Robinson et al. Integrative clinical genomics of advanced prostate cancer. *Cell.* 2015 May 21;161(5):1215-1228. PMID: 26000489
13. King et al. Amplification of a novel v-erbB-related gene in a human mammary carcinoma. *Science.* 1985 Sep 6;229(4717):974-6. PMID: 2992089
14. Zhixiang. ErbB Receptors and Cancer. *Methods Mol. Biol.* 2017;1652:3-35. PMID: 28791631
15. Gutierrez et al. HER2: biology, detection, and clinical implications. *Arch. Pathol. Lab. Med.* 2011 Jan;135(1):55-62. PMID: 21204711
16. Pines et al. Oncogenic mutant forms of EGFR: lessons in signal transduction and targets for cancer therapy. *FEBS Lett.* 2010 Jun 18;584(12):2699-706. PMID: 20388509
17. Cancer Genome Atlas Research Network. Comprehensive molecular profiling of lung adenocarcinoma. *Nature.* 2014 Jul 31;511(7511):543-50. doi: 10.1038/nature13385. Epub 2014 Jul 9. PMID: 25079552
18. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
19. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
20. da et al. EGFR mutations and lung cancer. *Annu Rev Pathol.* 2011;6:49-69. doi: 10.1146/annurev-pathol-011110-130206. PMID: 20887192
21. Arcila et al. EGFR exon 20 insertion mutations in lung adenocarcinomas: prevalence, molecular heterogeneity, and clinicopathologic characteristics. *Mol. Cancer Ther.* 2013 Feb;12(2):220-9. PMID: 23371856
22. Kobayashi et al. EGFR Exon 18 Mutations in Lung Cancer: Molecular Predictors of Augmented Sensitivity to Afatinib or Neratinib as Compared with First- or Third-Generation TKIs. *Clin Cancer Res.* 2015 Dec 1;21(23):5305-13. doi: 10.1158/1078-0432.CCR-15-1046. Epub 2015 Jul 23. PMID: 26206867
23. Yasuda et al. Structural, biochemical, and clinical characterization of epidermal growth factor receptor (EGFR) exon 20 insertion mutations in lung cancer. *Sci Transl Med.* 2013 Dec 18;5(216):216ra177. PMID: 24353160
24. Chiu et al. Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitor Treatment Response in Advanced Lung Adenocarcinomas with G719X/L861Q/S768I Mutations. *J Thorac Oncol.* 2015 May;10(5):793-9. PMID: 25668120
25. Karachaliou et al. KRAS mutations in lung cancer. *Clin Lung Cancer.* 2013 May;14(3):205-14. PMID: 23122493
26. Brennan et al. The somatic genomic landscape of glioblastoma. *Cell.* 2013 Oct 10;155(2):462-77. PMID: 24120142
27. Cancer Genome Atlas Network. Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature.* 2015 Jan 29;517(7536):576-82. PMID: 25631445
28. Mitsudomi et al. Epidermal growth factor receptor in relation to tumor development: EGFR gene and cancer. *FEBS J.* 2010 Jan;277(2):301-8. PMID: 19922469

## References (continued)

29. Gazdar. Activating and resistance mutations of EGFR in non-small-cell lung cancer: role in clinical response to EGFR tyrosine kinase inhibitors. *Oncogene*. 2009 Aug;28 Suppl 1:S24-31. PMID: 19680293
30. Gan et al. The EGFRvIII variant in glioblastoma multiforme. *J Clin Neurosci*. 2009 Jun;16(6):748-54. PMID: 19324552
31. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2016/021743s025lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2016/021743s025lbl.pdf)
32. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/206995s004lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/206995s004lbl.pdf)
33. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2019/201292s015lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/201292s015lbl.pdf)
34. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/211288s003lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/211288s003lbl.pdf)
35. NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]
36. Naidoo et al. Epidermal growth factor receptor exon 20 insertions in advanced lung adenocarcinomas: Clinical outcomes and response to erlotinib. *Cancer*. 2015 Sep 15;121(18):3212-3220. PMID: 26096453
37. Vyse et al. Targeting EGFR exon 20 insertion mutations in non-small cell lung cancer. *Signal Transduct Target Ther*. 2019;4:5. PMID: 30854234
38. Yi et al. A comparison of epidermal growth factor receptor mutation testing methods in different tissue types in non-small cell lung cancer. *Int J Mol Med*. 2014 Aug;34(2):464-74. PMID: 24891042
39. Madic et al. EGFR C797S, EGFR T790M and EGFR sensitizing mutations in non-small cell lung cancer revealed by six-color crystal digital PCR. *Oncotarget*. 2018 Dec 21;9(100):37393-37406. PMID: 30647840
40. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/208065s022lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/208065s022lbl.pdf)
41. Niederst et al. The Allelic Context of the C797S Mutation Acquired upon Treatment with Third-Generation EGFR Inhibitors Impacts Sensitivity to Subsequent Treatment Strategies. *Clin. Cancer Res*. 2015 Sep 1;21(17):3924-33. PMID: 25964297
42. Wang et al. Lung Adenocarcinoma Harboring EGFR T790M and In Trans C797S Responds to Combination Therapy of First- and Third-Generation EGFR TKIs and Shifts Allelic Configuration at Resistance. *J Thorac Oncol*. 2017 Nov;12(11):1723-1727. PMID: 28662863
43. <https://www.jnj.com/janssen-announces-u-s-fda-breakthrough-therapy-designation-granted-for-jnj-6372-for-the-treatment-of-non-small-cell-lung-cancer>
44. <https://www.takeda.com/newsroom/newsreleases/2020/takeda-announces-u.s.-fda-breakthrough-therapy-designation-for-mobocertinib-tak-788-for-the-treatment-of-nscl-patients-with-egfr-exon-20-insertion-mutations/>
45. <https://www.genprex.com/news/genprex-receives-u-s-fda-fast-track-designation-for-gene-therapy-that-targets-lung-cancer/>
46. <https://investors.blackdiamondtherapeutics.com/news-releases/news-release-details/black-diamond-therapeutics-granted-fast-track-designation-fda>