



Sample Information

Patient Name: 王玉  
Gender: Female  
ID No.: F228804068  
History No.: 30487402  
Age: 84  
  
Ordering Doctor: DOC1654E 林庭安  
Ordering REQ.: H45DB31  
Signing in Date: 2023/01/18

Path No.: M112-00010  
MP No.: MY23005  
Assay: Oncomine Myeloid Assay  
Sample Type: Blood  
Date of blood drawing: 2023/01/13

Reporting Doctor: DOC5466K 葉奕成 (Phone: 8#5466)

Note:

Sample Cancer Type: Myelodysplastic Syndrome

Table of Contents	Page	Report Highlights
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2	2 Relevant Biomarkers
Biomarker Descriptions	3	15 Therapies Available
Relevant Therapy Summary	4	0 Clinical Trials
Relevant Therapy Details	5	
Prognostic Details	12	
Diagnostic Details	13	

Relevant Myelodysplastic Syndrome Variants

Gene	Finding	Gene	Finding
ASXL1	<b>ASXL1 p.(R404*) c.1210C&gt;T</b>	NPM1	None detected
BCOR	None detected	NRAS	None detected
CBL	None detected	NUP214	None detected
CREBBP	None detected	RUNX1	None detected
ETV6	None detected	SF3B1	<b>SF3B1 p.(K700E) c.2098A&gt;G</b>
EZH2	None detected	SRSF2	None detected
FLT3	None detected	STAG2	None detected
GATA2	None detected	TP53	None detected
IDH2	None detected	U2AF1	None detected
KIT	None detected	WT1	None detected
KMT2A	None detected	ZRSR2	None detected

## Relevant Myelodysplastic Syndrome Variants (continued)

Gene	Finding	Gene	Finding
MECOM	None detected		

## Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	<i>SF3B1</i> p.(K700E) c.2098A>G splicing factor 3b subunit 1 Allele Frequency: 26.01%  <b>Prognostic significance:</b> NCCN: Favorable <b>Diagnostic significance:</b> Myelodysplastic Syndrome	luspatercept	hypomethylating agent + lenalidomide lenalidomide luspatercept	0
IA	<i>ASXL1</i> p.(R404*) c.1210C>T ASXL transcriptional regulator 1 Allele Frequency: 5.60%  <b>Prognostic significance:</b> NCCN: Poor	None	allogeneic stem cells azacitidine cytarabine cytarabine + daunorubicin cytarabine + daunorubicin + etoposide cytarabine + etoposide + idarubicin cytarabine + fludarabine + idarubicin + filgrastim cytarabine + idarubicin cytarabine + mitoxantrone decitabine gemtuzumab ozogamicin + chemotherapy venetoclax + chemotherapy	0

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

## Prevalent cancer biomarkers without relevant evidence based on included data sources

*SETBP1* p.(G870S) c.2608G>A

## Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

### DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
SF3B1	p.(K700E)	c.2098A>G	COSM84677	chr2:198266834	26.01%	NM_012433.4	missense	1999
SETBP1	p.(G870S)	c.2608G>A	COSM1234973	chr18:42531913	27.25%	NM_015559.3	missense	2000
ASXL1	p.(R404*)	c.1210C>T	.	chr20:31021211	5.60%	NM_015338.6	nonsense	2000
DNMT3A	p.(M801V)	c.2401A>G	.	chr2:25462006	26.45%	NM_022552.4	missense	2000
EZH2	p.(D90Ifs*9)	c.268delG	.	chr7:148529820	51.04%	NM_004456.5	frameshift Deletion	1979

## Biomarker Descriptions

### ASXL1 (ASXL transcriptional regulator 1)

**Background:** The ASXL1 gene encodes the ASXL transcriptional regulator 1 protein, a ligand-dependent co-activator and epigenetic scaffolding protein involved in transcriptional regulation<sup>1,2</sup>. ASXL1 belongs to the ASXL gene family, which also includes ASXL2 and ASXL3<sup>2</sup>. ASXL proteins contain a conserved c-terminal plant homeodomain (PHD) which facilitates interaction with DNA and histones<sup>2,3</sup>. ASXL1 influences chromatin remodeling and transcription through interaction with BAP1 and polycomb repressive complex (PRC) proteins, as well as other transcriptional activators and repressors<sup>2,4</sup>. In cancer, ASXL1 is the target of somatic mutations which often result in a truncated ASXL1 protein and loss of its PHD<sup>5,6,7</sup>. Such mutations can lead to impaired protein function and consequent upregulation of HOXA gene expression, supporting a tumor suppressor role for ASXL1<sup>8</sup>.

**Alterations and prevalence:** Missense, nonsense, and frameshift mutations in ASXL1 are reported in 3-6% of de novo acute myeloid leukemia (AML), up to 36% of secondary AML, approximately 15% of myelodysplastic syndromes (MDS), up to 23% of myeloproliferative neoplasms (MPN), up to 30% of systemic mastocytosis (SM), and approximately 45% of chronic myelomonocytic leukemia (CMML)<sup>4,9,10,11,12,13,14,15,16</sup>. The ASXL1 G646Wfs\*12 mutation accounts for over 50% of ASXL1 mutated cases in myeloid malignancies<sup>6,11,17</sup>. This mutation results from a single nucleotide expansion that occurs within an eight base pair guanine repeat that extends from c.1927 to c.1934. It is proposed that the high prevalence of the G646Wfs\*12 variant is due to replication slippage which can occur in areas of repetitive sequence<sup>18</sup>. As a consequence, detection of G646Wfs\*12 may result as an artifact of PCR and/or sequencing<sup>19</sup>. However, multiple studies observe an increase in the frequency of G646Wfs\*12 in myeloid cancer relative to normal suggesting that G646Wfs\*12 is a bona fide somatic mutation<sup>9,18,20</sup>.

**Potential relevance:** The majority of frameshift and nonsense mutations in ASXL1 that result in protein truncation and removal of the PHD domain are considered pathogenic<sup>21</sup>. Mutations in ASXL1 confer poor/adverse risk in AML and are associated with poor outcomes and adverse risk<sup>16</sup>. Additionally, ASXL1 nonsense or frameshift mutations are independently associated with poor prognosis in MDS and CMML<sup>22</sup>. Moreover, ASXL1 mutations are independently associated with inferior overall survival (OS) in patients with MPN or SM<sup>23,24</sup>.

### SETBP1 (SET binding protein 1)

**Background:** The SETBP1 gene encodes the SET binding protein 1, a multi-functional protein which contributes to several cellular processes including transcriptional regulation, proliferation, differentiation, and transformation<sup>25</sup>. SETBP1 contains a SET binding domain, which enables SETBP1 to form complexes with SET domain containing proteins, including the nuclear SET oncoprotein, a potent inhibitor of protein phosphatase 2A (PP2A)<sup>25,26,27</sup>. SETBP1 binding stabilizes SET, leading to elevated SET expression and increased inhibition of PP2A<sup>25,28,29</sup>. SETBP1 mediated inhibition of PP2A facilitates leukemic transformation in hematological malignancies including acute myeloid leukemia (AML)<sup>29</sup>. SETBP1 also contains three AT-hook domains, three nuclear localization motifs, and a SKI-homologous region which can influence transcriptional regulation<sup>25</sup>. SETBP1 is the target of somatic mutations in both hematological malignancies as well as solid tumors<sup>12,30</sup>. SETBP1 mutations often result in a gain of function and can lead to HOX gene upregulation, suggesting an oncogenic role for SETBP1 in cancer<sup>30,31</sup>. Additionally, germline gain of function mutations in SETBP1 are found to be causal of Schinzel-Giedion syndrome (SGS), a rare developmental disorder characterized by multiple malformations, severe neurological alterations and increased risk of cancer<sup>32</sup>.

**Alterations and prevalence:** SETBP1 mutations are observed in up to 32% of atypical chronic myeloid leukemia (aCML), 24% of juvenile myelomonocytic leukemia (JMML), 18% of chronic myelomonocytic leukemia (CMML), 10% of myelodysplastic/myeloproliferative neoplasms (MDS/MPN), 1-3% of primary AML and up to 17% of secondary AML (sAML)<sup>12,30,33,34,35</sup>. Additionally, mutations in SETBP1 are reported in solid tumors including up to 12% of melanoma, 11% of lung adenocarcinoma, 9% of stomach and uterine cancer, as well as, 6% of esophageal and colorectal carcinoma<sup>12</sup>. SETBP1 mutations are predominantly missense, the most recurrent involving amino acid substitutions at D868, G870, and I871<sup>30,34,35</sup>. SETBP1 fusions have also been described in hematological malignancies. The t(11;18)(p15;q12)/NUP98-SETBP1 and t(12;18)(p13;q12)/ETV6-SETBP1 fusions have been reported in individual cases of T-cell acute lymphoblastic leukemia (T-ALL) and AML, respectively<sup>36,37</sup>.

**Potential relevance:** The presence of SETBP1 mutations is one of the diagnostic criteria for CMML as defined by the World Health Organization (WHO)<sup>38</sup>. Overexpression of SETBP1 is associated with accelerated leukemic transformation and poor prognosis in AML<sup>29,33</sup>. Additionally, mutations in SETBP1 are associated with poor prognosis in MDS/MPN, CMML, JMML, and aCML<sup>33,35,39,40</sup>.

### SF3B1 (splicing factor 3b subunit 1)

**Background:** The SF3B1 gene encodes the splicing factor 3b subunit 1 protein, a core component of the U2 small nuclear ribonucleoprotein (snRNP) complex of the spliceosome responsible for RNA splicing. SF3B1 is involved in recognition of the branch point sequence during selection of the 3' splice site. Recurrent somatic mutations in SF3B1 and other components of the splicing machinery including SRSF2, U2AF1, and ZRSR2, are common in myelodysplasia. These components experience mutations in a mutually exclusive manner suggesting a common impact on RNA splicing and the pathogenesis of myelodysplasia<sup>41</sup>. SF3B1 mutations

## Biomarker Descriptions (continued)

are believed to contribute to aberrant post-translational inactivation of the regulatory complex PPP2R5A of protein phosphatase 2A (PP2A), leading to the activation and stabilization of MYC activation and impairing apoptosis<sup>42</sup>.

**Alterations and prevalence:** SF3B1 mutations occur in the majority (70-80%) of myelodysplastic syndromes (MDS) with ring sideroblasts (RS) and at lower frequency in other myeloid neoplasms including MDS without RS (7%), chronic myelomonocytic leukemia (5-6%), therapy-related acute myeloid leukemia (AML) or AML with MDS features (5%), and de novo AML (3%)<sup>41,43,44</sup>. Recurrent somatic SF3B1 mutations are also common in certain solid cancers including uveal melanoma (20-30%) and breast cancer (2%) and at lower frequencies in diverse cancer types<sup>12,45,46,47,48,49,50,51</sup>. Cancer-associated recurrent missense mutations in SF3B1 occur within the HEAT repeat domains 5-9 at codon positions R625, K666, K700, G742, and D781<sup>52</sup>. The functional significance of recurrent SF3B1 mutations is to alter branch point selection thus inducing cryptic 3' splice site selection<sup>52,53,54</sup>.

**Potential relevance:** Currently, no therapies are approved for SF3B1 aberrations. SF3B1 mutations are associated with aggressive disease and shorter survival in patients diagnosed with chronic lymphocytic leukemia (CLL)<sup>55</sup>. Investigational inhibitors of the spliceosome are in early clinical development<sup>56,57</sup>.

## Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

### SF3B1 p.(K700E) c.2098A>G

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
luspatercept	×	○	×	●	×
hypomethylating agent + lenalidomide	×	○	×	×	×
lenalidomide	×	○	×	×	×

### ASXL1 p.(R404\*) c.1210C>T

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
Allogeneic hematopoietic stem cell transplantation	×	○	×	×	×
azacitidine	×	○	×	×	×
cytarabine	×	○	×	×	×
cytarabine + daunorubicin	×	○	×	×	×
cytarabine + daunorubicin + etoposide	×	○	×	×	×
cytarabine + etoposide + idarubicin	×	○	×	×	×
cytarabine + fludarabine + idarubicin + filgrastim	×	○	×	×	×
cytarabine + idarubicin	×	○	×	×	×
cytarabine + mitoxantrone	×	○	×	×	×
decitabine	×	○	×	×	×
gemtuzumab ozogamicin + cytarabine + daunorubicin	×	○	×	×	×
venetoclax + azacitidine	×	○	×	×	×

## Relevant Therapy Summary (continued)

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

### ASXL1 p.(R404\*) c.1210C>T (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
venetoclax + cytarabine	×	○	×	×	×
venetoclax + decitabine	×	○	×	×	×

## Relevant Therapy Details

### Current NCCN Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

NCCN information is current as of 2022-11-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org).  
For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

### SF3B1 p.(K700E) c.2098A>G

#### ☐ hypomethylating agent + lenalidomide

Cancer type: Myelodysplastic/Myeloproliferative Neoplasm Variant class: SF3B1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Line of therapy not specified); Consider

Reference: NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]

#### ☐ lenalidomide

Cancer type: Myelodysplastic/Myeloproliferative Neoplasm Variant class: SF3B1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Line of therapy not specified); Consider

Reference: NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]

**SF3B1 p.(K700E) c.2098A>G (continued)****○ luspatercept**

Cancer type: Myelodysplastic/Myeloproliferative Neoplasm      Variant class: SF3B1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Line of therapy not specified); Consider

Reference: NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]

**ASXL1 p.(R404\*) c.1210C>T****○ cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia      Variant class: ASXL1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**○ cytarabine + daunorubicin + etoposide**

Cancer type: Acute Myeloid Leukemia      Variant class: ASXL1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**○ cytarabine + etoposide + idarubicin**

Cancer type: Acute Myeloid Leukemia      Variant class: ASXL1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**ASXL1 p.(R404\*) c.1210C>T (continued)****○ cytarabine + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**○ Allogeneic hematopoietic stem cell transplantation**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**○ cytarabine**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**○ cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**○ cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**ASXL1 p.(R404\*) c.1210C>T (continued)****○ cytarabine + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**○ cytarabine + mitoxantrone**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**○ decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**○ gemtuzumab ozogamicin + cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**○ venetoclax + azacitidine**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]



**ASXL1 p.(R404\*) c.1210C>T (continued)****○ venetoclax + cytarabine**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**○ venetoclax + decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**○ azacitidine**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**○ cytarabine + daunorubicin + etoposide**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**○ cytarabine + etoposide + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**ASXL1 p.(R404\*) c.1210C>T (continued)****○ cytarabine + fludarabine + idarubicin + filgrastim**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

## Current ESMO Information

☒ In this cancer type    ☐ In other cancer type    ☐ In this cancer type and other cancer types

ESMO information is current as of 2022-11-01. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

### SF3B1 p.(K700E) c.2098A>G

#### ☒ luspatercept

Cancer type: Myelodysplastic Syndrome

Variant class: SF3B1 mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Refractory (Second-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Myelodysplastic Syndromes [Annals of Oncology (2020), <https://doi.org/10.1016/j.annonc.2020.11.002>]

## Prognostic Details

### Current NCCN Information

NCCN information is current as of 2022-11-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org).  
For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

#### SF3B1 p.(K700E) c.2098A>G

##### Prognostic significance: NCCN: Favorable

Cancer type: Myelodysplastic Syndrome

Variant class: SF3B1 K700E mutation

NCCN Recommendation category: 2A

##### Summary:

- NCCN Guidelines® independently associate the biomarker with favorable prognosis

Reference: NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]

#### ASXL1 p.(R404\*) c.1210C>T

##### Prognostic significance: NCCN: Poor

Cancer type: Myelodysplastic Syndrome

Variant class: ASXL1 truncating mutation

NCCN Recommendation category: 2A

##### Summary:

- NCCN Guidelines® independently associate the biomarker with poor prognosis in MDS and CMML

Reference: NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]

## Diagnostic Details

### Current NCCN Information

NCCN information is current as of 2022-11-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org).  
For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

#### SF3B1 p.(K700E) c.2098A>G

### Diagnostic significance: Myelodysplastic Syndrome

Variant class: SF3B1 mutation

NCCN Recommendation category: 2A

#### Diagnostic notes:

- 2022 WHO Classification of Myelodysplastic Neoplasms (MDS); MDS-SF3B1 (low blasts) with mutated SF3B1 variant allele frequency (VAF)  $\geq 10\%$ .  $\geq 15\%$  ring sideroblasts (RS) may substitute for SF3B1 mutation (MDS with low blasts and ring sideroblasts). Normal karyotype or any cytogenetic abnormality other than del(5q), monosomy 7, inv(3) or abnormal 3q26, complex ( $\geq 3$ ) or somatically mutated TP53, RUNX1, or EZH2 genes

Reference: NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]

## Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

## References

1. O'Leary et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res.* 2016 Jan 4;44(D1):D733-45. PMID: 26553804
2. Katoh. Functional and cancer genomics of ASXL family members. *Br. J. Cancer.* 2013 Jul 23;109(2):299-306. PMID: 23736028
3. Gelsi-Boyer et al. Mutations of polycomb-associated gene ASXL1 in myelodysplastic syndromes and chronic myelomonocytic leukaemia. *Br. J. Haematol.* 2009 Jun;145(6):788-800. PMID: 19388938
4. Gelsi-Boyer et al. Mutations in ASXL1 are associated with poor prognosis across the spectrum of malignant myeloid diseases. *J Hematol Oncol.* 2012 Mar 21;5:12. doi: 10.1186/1756-8722-5-12. PMID: 22436456
5. Larsson et al. The changing mutational landscape of acute myeloid leukemia and myelodysplastic syndrome. *Mol. Cancer Res.* 2013 Aug;11(8):815-27. PMID: 23645565
6. Alvarez et al. ASXL1 mutations in myeloid neoplasms: pathogenetic considerations, impact on clinical outcomes and survival. *Curr Med Res Opin.* 2018 May;34(5):757-763. PMID: 28027687
7. Yang et al. Gain of function of ASXL1 truncating protein in the pathogenesis of myeloid malignancies. *Blood.* 2018 Jan 18;131(3):328-341. PMID: 29113963
8. Abdel-Wahab et al. ASXL1 mutations promote myeloid transformation through loss of PRC2-mediated gene repression. *Cancer Cell.* 2012 Aug 14;22(2):180-93. PMID: 22897849
9. Alberti et al. Discriminating a common somatic ASXL1 mutation (c.1934dup; p.G646Wfs\*12) from artifact in myeloid malignancies using NGS. *Leukemia.* 2018 Aug;32(8):1874-1878. PMID: 29959414
10. Kakosaiou et al. ASXL1 mutations in AML are associated with specific clinical and cytogenetic characteristics. *Leuk. Lymphoma.* 2018 Oct;59(10):2439-2446. PMID: 29411666
11. Paschka et al. ASXL1 mutations in younger adult patients with acute myeloid leukemia: a study by the German-Austrian Acute Myeloid Leukemia Study Group. *Haematologica.* 2015 Mar;100(3):324-30. PMID: 25596267
12. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
13. Jawhar et al. The clinical and molecular diversity of mast cell leukemia with or without associated hematologic neoplasm. *Haematologica.* 2017 Jun;102(6):1035-1043. PMID: 28255023
14. Jawhar et al. KIT D816 mutated/CBF-negative acute myeloid leukemia: a poor-risk subtype associated with systemic mastocytosis. *Leukemia.* 2019 May;33(5):1124-1134. PMID: 30635631
15. Damaj et al. ASXL1 but not TET2 mutations adversely impact overall survival of patients suffering systemic mastocytosis with associated clonal hematologic non-mast-cell diseases. *PLoS ONE.* 2014;9(1):e85362. PMID: 24465546
16. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]
17. Boulton et al. Frequent mutation of the polycomb-associated gene ASXL1 in the myelodysplastic syndromes and in acute myeloid leukemia. *Leukemia.* 2010 May;24(5):1062-5. doi: 10.1038/leu.2010.20. Epub 2010 Feb 25. PMID: 20182461
18. Yannakou et al. ASXL1 c.1934dup;p.Gly646Trpfs\*12-a true somatic alteration requiring a new approach. *Blood Cancer J.* 2017 Dec 15;7(12):656. doi: 10.1038/s41408-017-0025-8. PMID: 29242575
19. Abdel-Wahab et al. The most commonly reported variant in ASXL1 (c.1934dupG;p.Gly646TrpfsX12) is not a somatic alteration. *Leukemia.* 2010 Sep;24(9):1656-7. doi: 10.1038/leu.2010.144. Epub 2010 Jul 1. PMID: 20596031
20. Montes-Moreno et al. Clinical molecular testing for ASXL1 c.1934dupG p.Gly646fs mutation in hematologic neoplasms in the NGS era. *PLoS ONE.* 2018;13(9):e0204218. PMID: 30222780
21. Landrum et al. ClinVar: improving access to variant interpretations and supporting evidence. *Nucleic Acids Res.* 2018 Jan 4;46(D1):D1062-D1067. PMID: 29165669
22. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]
23. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 3.2022]
24. NCCN Guidelines® - NCCN-Systemic Mastocytosis [Version 1.2020]
25. Coccato et al. SETBP1 dysregulation in congenital disorders and myeloid neoplasms. *Oncotarget.* 2017 Aug 1;8(31):51920-51935. PMID: 28881700
26. Hoischen et al. De novo mutations of SETBP1 cause Schinzel-Giedion syndrome. *Nat. Genet.* 2010 Jun;42(6):483-5. PMID: 20436468
27. Li et al. The myeloid leukemia-associated protein SET is a potent inhibitor of protein phosphatase 2A. *J. Biol. Chem.* 1996 May 10;271(19):11059-62. PMID: 8626647
28. Janssens et al. Protein phosphatase 2A: a highly regulated family of serine/threonine phosphatases implicated in cell growth and signalling. *Biochem. J.* 2001 Feb 1;353(Pt 3):417-39. PMID: 11171037

## References (continued)

29. Cristóbal et al. SETBP1 overexpression is a novel leukemogenic mechanism that predicts adverse outcome in elderly patients with acute myeloid leukemia. *Blood*. 2010 Jan 21;115(3):615-25. PMID: 19965692
30. Makishima et al. Somatic SETBP1 mutations in myeloid malignancies. *Nat. Genet.* 2013 Aug;45(8):942-6. PMID: 23832012
31. Inoue et al. SETBP1 mutations drive leukemic transformation in ASXL1-mutated MDS. *Leukemia*. 2015 Apr;29(4):847-57. PMID: 25306901
32. Acuna-Hidalgo et al. Overlapping SETBP1 gain-of-function mutations in Schinzel-Giedion syndrome and hematologic malignancies. *PLoS Genet.* 2017 Mar;13(3):e1006683. PMID: 28346496
33. Linder et al. SETBP1 mutations as a biomarker for myelodysplasia /myeloproliferative neoplasm overlap syndrome. *Biomark Res.* 2017 Dec 6;5:33. doi: 10.1186/s40364-017-0113-8. eCollection 2017. PMID: 29225884
34. Meggendorfer et al. SETBP1 mutations occur in 9% of MDS/MPN and in 4% of MPN cases and are strongly associated with atypical CML, monosomy 7, isochromosome i(17)(q10), ASXL1 and CBL mutations. *Leukemia*. 2013 Sep;27(9):1852-60. PMID: 23628959
35. Piazza et al. Recurrent SETBP1 mutations in atypical chronic myeloid leukemia. *Nat. Genet.* 2013 Jan;45(1):18-24. PMID: 23222956
36. Panagopoulos et al. Fusion of NUP98 and the SET binding protein 1 (SETBP1) gene in a paediatric acute T cell lymphoblastic leukaemia with t(11;18)(p15;q12). *Br. J. Haematol.* 2007 Jan;136(2):294-6. PMID: 17233820
37. Huret et al. Atlas of genetics and cytogenetics in oncology and haematology in 2013. *Nucleic Acids Res.* 2013 Jan;41(Database issue):D920-4. PMID: 23161685
38. Arber et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood*. 2016 May 19;127(20):2391-405. PMID: 27069254
39. Shou et al. Prognostic significance of SETBP1 mutations in myelodysplastic syndromes, chronic myelomonocytic leukemia, and chronic neutrophilic leukemia: A meta-analysis. *PLoS ONE*. 2017;12(2):e0171608. PMID: 28158286
40. Stieglitz et al. Subclonal mutations in SETBP1 confer a poor prognosis in juvenile myelomonocytic leukemia. *Blood*. 2015 Jan 15;125(3):516-24. PMID: 25395418
41. Yoshida et al. Frequent pathway mutations of splicing machinery in myelodysplasia. *Nature*. 2011 Sep 11;478(7367):64-9. PMID: 21909114
42. Liu et al. Mutations in the RNA Splicing Factor SF3B1 Promote Tumorigenesis through MYC Stabilization. *Cancer Discov.* 2020 Mar 18. PMID: 32188705
43. Patnaik et al. Refractory anemia with ring sideroblasts (RARS) and RARS with thrombocytosis (RARS-T): 2017 update on diagnosis, risk-stratification, and management. *Am. J. Hematol.* 2017 Mar;92(3):297-310. PMID: 28188970
44. Patnaik et al. Spliceosome mutations involving SRSF2, SF3B1, and U2AF35 in chronic myelomonocytic leukemia: prevalence, clinical correlates, and prognostic relevance. *Am. J. Hematol.* 2013 Mar;88(3):201-6. PMID: 23335386
45. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
46. Ellis et al. Whole-genome analysis informs breast cancer response to aromatase inhibition. *Nature*. 2012 Jun 10;486(7403):353-60. PMID: 22722193
47. Maguire et al. SF3B1 mutations constitute a novel therapeutic target in breast cancer. *J. Pathol.* 2015 Mar;235(4):571-80. PMID: 25424858
48. Harbour et al. Recurrent mutations at codon 625 of the splicing factor SF3B1 in uveal melanoma. *Nat. Genet.* 2013 Feb;45(2):133-5. PMID: 23313955
49. Martin et al. Exome sequencing identifies recurrent somatic mutations in EIF1AX and SF3B1 in uveal melanoma with disomy 3. *Nat. Genet.* 2013 Aug;45(8):933-6. PMID: 23793026
50. Furney et al. SF3B1 mutations are associated with alternative splicing in uveal melanoma. *Cancer Discov.* 2013 Oct;3(10):1122-1129. PMID: 23861464
51. Dono et al. Mutation frequencies of GNAQ, GNA11, BAP1, SF3B1, EIF1AX and TERT in uveal melanoma: detection of an activating mutation in the TERT gene promoter in a single case of uveal melanoma. *Br. J. Cancer*. 2014 Feb 18;110(4):1058-65. PMID: 24423917
52. DeBoever et al. Transcriptome sequencing reveals potential mechanism of cryptic 3' splice site selection in SF3B1-mutated cancers. *PLoS Comput. Biol.* 2015 Mar;11(3):e1004105. PMID: 25768983
53. Darman et al. Cancer-Associated SF3B1 Hotspot Mutations Induce Cryptic 3' Splice Site Selection through Use of a Different Branch Point. *Cell Rep.* 2015 Nov 3;13(5):1033-45. PMID: 26565915



---

## References (continued)

54. Alsafadi et al. Cancer-associated SF3B1 mutations affect alternative splicing by promoting alternative branchpoint usage. *Nat Commun.* 2016 Feb 4;7:10615. doi: 10.1038/ncomms10615. PMID: 26842708
55. Wan et al. SF3B1 mutations in chronic lymphocytic leukemia. *Blood.* 2013 Jun 6;121(23):4627-34. PMID: 23568491
56. Seiler et al. H3B-8800, an orally available small-molecule splicing modulator, induces lethality in spliceosome-mutant cancers. *Nat. Med.* 2018 May;24(4):497-504. PMID: 29457796
57. Lee et al. Therapeutic targeting of splicing in cancer. *Nat. Med.* 2016 Sep 7;22(9):976-86. PMID: 27603132