

# ACT Onco<sup>®</sup> + Report

PATIENT		
Identifier: 謝進富		Patient ID: 46354954
Date of Birth: Feb 10, 1959		Gender: Male
Diagnosis: Hepatocellular carcinoma		
ORDERING PHYSICIAN		
Name: 陳三奇醫師		Tel: 886-228712121
Facility: 臺北榮總		
Address: 臺北市北投區石牌路二段 201 號		
SPECIMEN		
Specimen ID: S11155447A	Collection site: Bone	Type: FFPE tissue
Date received: Jan 05, 2023	Lab ID: AA-23-00107	D/ID: NA

## ABOUT ACT Onco<sup>®</sup>+

The test is a next-generation sequencing (NGS)-based assay developed for efficient and comprehensive genomic profiling of cancers. This test interrogates coding regions of 440 genes associated with cancer treatment, prognosis and diagnosis. Genetic mutations detected by this test include small-scale mutations like single nucleotide variants (SNVs), small insertions and deletions (InDels) ( $\leq 15$  nucleotides) and large-scale genomic alterations like copy number alterations (CNAs). The test also includes an RNA test, detecting fusion transcripts of 13 genes.

## SUMMARY FOR ACTIONABLE VARIANTS

### VARIANTS/BIOMARKERS WITH EVIDENCE OF CLINICAL SIGNIFICANCE

Genomic Alterations/Biomarkers	Probable Effects in Patient's Cancer Type		Probable Sensitive in Other Cancer Types
	Sensitive	Resistant	
TSC1 Splice donor	-	-	Everolimus

### VARIANTS/BIOMARKERS WITH POTENTIAL CLINICAL SIGNIFICANCE

Genomic Alterations/Biomarkers	Possibly Sensitive	Possibly Resistant
ARID1A I1816fs	Dasatinib, Olaparib, Rucaparib, Talazoparib	-
TSC1 Splice donor	Temsirolimus	-
LYN Amplification	Dasatinib	-
MCL1 Amplification	Regorafenib, Sorafenib	-

#### Note:

- The above summary tables present genomic variants and biomarkers based on the three-tiered approach proposed by US FDA for reporting tumor profiling NGS testing. "Variants/biomarkers with evidence of clinical significance" refers to mutations that are widely recognized as standard-of-care biomarkers (FDA level 2/AMP tier 1). "Variants/biomarkers with potential clinical significance" refers to mutations that are not included in the standard of care but are informational for clinicians, which are commonly biomarkers used as inclusion criteria for clinical trials (FDA level 3/AMP tier 2).
- The therapeutic agents and possible effects to a given drug are based on mapping the variants/biomarkers with ACT Genomics clinical knowledge database. The mapping results only provide information for reference, but not medical recommendation.
- Please refer to corresponding sections for more detailed information about genomic alteration and clinical relevance listed above.

# ACTOnco<sup>®</sup> + Report

## TESTING RESULTS

### VARIANT(S) WITH CLINICAL RELEVANCE

#### - Single Nucleotide and Small InDel Variants

Gene	Amino Acid Change	Allele Frequency
ARID1A	I1816fs	21.4%
CTNNB1	S33Y	35.1%
TSC1	Splice donor	11.6%

#### - Copy Number Alterations

Chromosome	Gene	Variation	Copy Number
Chr1	MCL1, MDM4, NTRK1	Amplification	7
Chr5	TERT	Amplification	7
Chr6	E2F3	Amplification	7
Chr8	RECQL4	Amplification	11
Chr8	LYN, MYC, NBN	Amplification	15

#### - Fusions

Fusion Gene & Exon	Transcript ID
No fusion gene detected in this sample	

#### - Immune Checkpoint Inhibitor (ICI) Related Biomarkers

Biomarker	Results
Tumor Mutational Burden (TMB)	6.9 muts/Mb
Microsatellite Instability (MSI)	Microsatellite stable (MSS)

#### Note:

- Variant(s) enlisted in the SNV table may currently exhibit no relevance to treatment response prediction. Please refer to INTERPRETATION for more biological information and/or potential clinical impacts of the variants.
- Loss of heterozygosity (LOH) information was used to infer tumor cellularity. Copy number alteration in the tumor was determined based on 44% tumor purity.
- TMB was calculated by using the sequenced regions of ACTOnco<sup>®</sup> to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The threshold for high mutation load is set at  $\geq 7.5$  mutations per megabase. TMB, microsatellite status and gene copy number deletion cannot be determined if calculated tumor purity is  $< 30\%$ .

# ACT Onco<sup>®</sup> + Report

## THERAPEUTIC IMPLICATIONS TARGETED THERAPIES

Genomic Alterations	Therapies	Effect
<b>Level 3A</b>		
<b>TSC1</b> Splice donor	Everolimus	<b>sensitive</b>
<b>Level 3B</b>		
<b>ARID1A</b> I1816fs	Olaparib	<b>sensitive</b>
<b>TSC1</b> Splice donor	Temsirolimus	<b>sensitive</b>
<b>Level 4</b>		
<b>ARID1A</b> I1816fs	Dasatinib, Rucaparib, Talazoparib	<b>sensitive</b>
<b>LYN</b> Amplification	Dasatinib	<b>sensitive</b>
<b>MCL1</b> Amplification	Regorafenib, Sorafenib	<b>sensitive</b>

Therapies associated with benefit or lack of benefit are based on biomarkers detected in this tumor and published evidence in professional guidelines or peer-reviewed journals.

Level	Description
<b>1</b>	FDA-recognized biomarkers predictive of response or resistance to FDA approved drugs in this indication
<b>2</b>	Standard care biomarkers (recommended by the NCCN guideline) predictive of response or resistance to FDA approved drugs in this indication
<b>3A</b>	Biomarkers predictive of response or resistance to therapies approved by the FDA or NCCN guideline in a different cancer type
<b>3B</b>	Biomarkers that serve as inclusion criteria for clinical trials (minimal supportive data required)
<b>4</b>	Biomarkers that show plausible therapeutic significance based on small studies, few case reports, or preclinical studies

# ACT Onco<sup>®</sup> + Report

## IMMUNE CHECKPOINT INHIBITORS (ICIs)

No genomic alterations detected to confer sensitivity or lack of benefit to immune checkpoint therapies.

### - Other Biomarkers with Potential Clinical Effects for ICIs

Genomic Alterations	Potential Clinical Effects
MDM2/MDM4 amplification	Likely associated with WORSE response to ICIs

Note: Tumor non-genomic factors, such as patient germline genetics, PDL1 expression, tumor microenvironment, epigenetic alterations or other factors not provided by this test may affect ICI response.

## CHEMOTHERAPIES

Genomic Alterations	Therapies	Effect	Level of Evidence	Cancer Type
<b>MYC</b> Amplification	FAC CMF and P-FEC regimens	<b>Sensitive</b>	Clinical	Breast cancer
	Platinum-based regimens	<b>Sensitive</b>	Clinical	Ovarian cancer
<b>ARID1A</b> I1816fs	Platinum-based regimens	<b>Less sensitive</b>	Clinical	Ovarian cancer

## HORMONAL THERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to hormonal therapies.

## OTHERS

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to other therapies.

### Note:

Therapeutic implications provided in the test are based solely on the panel of 440 genes sequenced. Therefore, alterations in genes not covered in this panel, epigenetic and post-transcriptional and post-translational factors may also determine a patient's response to therapies. In addition, several other patient-associated clinical factors, including but not limited to, prior lines of therapies received, dosage and combinations with other therapeutic agents, patient's cancer types, sub-types, and/or stages, may also determine the patient's clinical response to therapies.

# ACT Onco<sup>®</sup> + Report

## VARIANT INTERPRETATION

### ARID1A I1816fs

#### Biological Impact

The AT-rich interactive domain 1A (ARID1A) gene encodes the BAF250A protein, a component of the SWI/SNF chromatin remodeling complex that plays a role in various cellular functions, including DNA repair, DNA synthesis, and transcription<sup>[1][2]</sup>. Haploinsufficiency of ARID1A is associated with tumor formation in some cancers<sup>[3]</sup>. Inactivation of ARID1A is commonly observed in ovarian, endometrial, uterine, and, gastric cancers<sup>[4][5][6][7][8]</sup>.

I1816fs mutation results in a change in the amino acid sequence beginning at 1816, likely to cause premature truncation of the functional ARID1A protein (UniProtKB). This mutation is predicted to lead to a loss of ARID1A protein function, despite not being characterized in the literature.

#### Therapeutic and prognostic relevance

ARID1A is the most frequently mutated genes in ovarian clear cell carcinoma and several synthetic lethality hypothesis-based therapeutic targets in ARID1A mutated cancer are in development. For examples, 1) EZH2 inhibitor<sup>[9][10]</sup>; 2) AKT-inhibitors MK-2206 and perifosine, as well as PI3K-inhibitor buparlisib<sup>[11]</sup>; 3) multiple kinase inhibitor, dasatinib<sup>[12]</sup>.

Some preclinical evidences suggested that reduced ARID1A expression confers resistance to several HER2/PI3K/mTOR signaling cascade inhibitors such as AZD8055 and trastuzumab, through activation of annexin A1 expression<sup>[13]</sup>. Loss or decreased expression of ARID1A has been reported to associate with resistance to platinum-based chemotherapies, shorter overall survival and lower complete response rate in ovarian cancer patients<sup>[14][15]</sup>.

Low expression of ARID1A is a significant and independent prognostic factor for poor disease-free and overall survival in breast cancer patients<sup>[16][17]</sup>. Besides, loss of ARID1A expression was more frequently seen in mismatch repair (MMR)-deficient colorectal cancers, predominantly in tumor with MLH1 promoter hypermethylation<sup>[18]</sup>. Positive ARID1A expression could independently predict worse overall survival in stage IV CRC patients compared with negative ARID1A expression<sup>[19]</sup>.

ARID1A mutation has been determined as an inclusion criterion for the trials evaluating olaparib efficacy in metastatic biliary tract cancer (NCT04042831), and niraparib efficacy in melanoma (NCT03925350), pancreatic cancer (NCT03553004), or any malignancy, except prostate cancer (NCT03207347).

The preclinical study discovered that ARID1A deficiency sensitized some tumors to PARP inhibitor drugs, such as olaparib, rucaparib, talazoparib, and veliparib, which block DNA damage repair pathways<sup>[20]</sup>.

### CTNNB1 S33Y

#### Biological Impact

The CTNNB1 gene encodes for the  $\beta$ -catenin, a transcriptional activator involves in the canonical Wnt signaling pathway<sup>[21][22]</sup>.  $\beta$ -catenin also regulates cyclin D1 and MYC expression, which play important roles in cancer development<sup>[23][24]</sup>. Mutations of CTNNB1 are common in a wide range of solid tumors, including liver, endometrial, colorectal, and lung cancer<sup>[25][26][27][28][29][30]</sup>. CTNNB1 mutations are more frequently found in hepatocellular carcinomas (HCCs) patients without hepatitis B virus (HBV) infection, which is mostly developed on the well-differentiated, noncirrhotic liver, and displayed cholestasis<sup>[31][32][33][34]</sup>. Of note, the majority of CTNNB1 alterations identified in cancers are missense mutations and all of which localize in the hotspot exon 3 at S33, S37, S45, T41, D32, and G34<sup>[35][36]</sup>.

CTNNB1 S33Y lies within the ubiquitination recognition motif and changes the potential GSK3 $\beta$  phosphorylation site on the  $\beta$ -catenin protein<sup>[37]</sup>. S33Y is demonstrated as a gain-of-function mutation that increases CTNNB1-dependent transcription and cell proliferation in vitro<sup>[38]</sup>.

# ACT Onco<sup>®</sup> + Report

## Therapeutic and prognostic relevance

In a retrospective study, patients with desmoid fibromatosis harboring CTNNB1 activating mutations such as S45F/N/P or T41A demonstrated a greater progression arrest rate (PAR) at 6 months compared to patients with wild-type CTNNB1 when treated with imatinib, a multi-target inhibitor of c-KIT, PDGFR, and BCR-ABL<sup>[39]</sup>.

Results from a Phase II study of temsirolimus-containing regimens in advanced endometrial cancer (EC) showed that CTNNB1 exon 3 mutations were associated with longer PFS on temsirolimus<sup>[40]</sup>. Besides, three patients with recurrent endometrial carcinoma harboring CTNNB1 mutations on exon 3 (one is D32V, another is S37Y, and the other is both H36Y and S37C) also responded well to everolimus and letrozole, based on the results of a Phase II study<sup>[41]</sup>.

Low expression of CTNNB1 has been reported to associate with longer overall survival in low-grade endometrioid endometrial carcinoma (EEC)<sup>[42]</sup>.

## TSC1 Splice donor

### Biological Impact

The tuberous sclerosis complex 1 (TSC1) gene encodes a tumor suppressor, hamartin, a key negative regulator of the mammalian target of rapamycin (mTOR) pathway<sup>[43][44]</sup>. Mutations in TSC1/TSC2 tumor suppressor genes that result in inactivation of the complex are commonly found in patients with tuberous sclerosis<sup>[45][46][47]</sup>, while LOH in TSC1/TSC2 has been identified in head and neck squamous cell carcinoma (HNSCC)<sup>[48]</sup> and endometrial cancer<sup>[49]</sup>. Loss of single TSC1 allele (haploinsufficiency) may provide a growth advantage to bladder epithelial cells, contributing to bladder cancer development<sup>[50]</sup>. Both TSC1 and TSC2 mutations cause the autosomal dominant genetic disorder tuberous sclerosis complex (TSC), in which individuals develop a variety of benign but often progressive neoplasms<sup>[51]</sup>.

TSC1 c.1997+1G>C is a variant located at the splice donor region, which may result in the exon skipping.

## Therapeutic and prognostic relevance

Genomic alterations with activating effects of the mTOR signaling pathway (including deletion/inactivation of TSC1/TSC2) have been shown to confer sensitivity to everolimus across multiple neoplasms, such as bladder tumors<sup>[52]</sup>, gastric, sarcoma, thyroid cancer, and HNSCC<sup>[53]</sup>. There were case reports demonstrated the efficacy of sirolimus in malignant uterine perivascular epithelioid cell tumors (PEComa) patients harboring mutations/deletions in TSC1 and TSC2 genes, and temsirolimus in PEComa patients with hyperactivated mTOR pathway. Genomic profiling analysis of GOG248, a Phase II study of temsirolimus or temsirolimus and alternating megestrol acetate and tamoxifen for advanced endometrial cancer showed that mutations in AKT1, TSC1 and TSC2 may predict clinical benefit from temsirolimus<sup>[40]</sup>. Recent studies indicate that there are mTORC1-independent signaling pathways downstream of hamartin-tuberin, which may represent new therapeutic targets<sup>[54]</sup>.

Everolimus has been approved by the U.S. FDA for Tuberous Sclerosis Complex (TSC)-associated renal angiomyolipoma and Tuberous Sclerosis Complex (TSC)-associated subependymal giant cell astrocytoma (SEGA). This approval is based on the results from EXIST-1, EXIST-2, and Study 2485 trials (NCT00789828, NCT00790400, and NCT00411619).

## E2F3 Amplification

### Biological Impact

The E2F3 gene encodes a transcription factor that interacts directly with the retinoblastoma protein (pRB) to regulate the expression of genes involved in the cell cycle and DNA replication<sup>[55][56][57]</sup>.

Amplification or overexpression of E2F3 has been reported in various types of cancers, including bladder cancer, hepatocellular carcinoma, retinoblastomas, and melanoma<sup>[58][59][60][61][62]</sup>.



# ACT Onco<sup>®</sup> + Report

## Therapeutic and prognostic relevance

A tissue microarray analysis indicated that amplification of the E2F3 gene is associated with increased E2F3 protein overexpression, accelerated cell proliferation, and poor prognosis in bladder cancer<sup>[59]</sup>. Besides, elevated E2F genes and E2F transcriptional targets in tumors have been linked with poor prognosis in the liver and pancreatic cancers<sup>[57]</sup>.

## LYN Amplification

### Biological Impact

The LYN Proto-Oncogene, Src Family Tyrosine Kinase (LYN) gene encodes a non-receptor tyrosine protein kinase of the Src family (SFK)<sup>[63]</sup>. LYN plays an important role in the regulation of immune responses, hematopoiesis, signal transduction of growth factors and cytokines and is activated in the cellular response to DNA damage and genotoxic agents<sup>[64][65][66][67]</sup>. LYN has been described to promote tumor growth, invasion, epithelial to mesenchymal transition (EMT) and ERK signaling in different cancer types<sup>[68][69][70]</sup>. Amplification of LYN has been identified in prostate cancer, breast cancer and ovarian cancer (cBioPortal).

## Therapeutic and prognostic relevance

LYN could be pharmacologically targeted with Src-kinase inhibitors. Results of preclinical studies showed that dasatinib, a dual-specificity tyrosine kinase inhibitor of ABL and the Src-family tyrosine kinases, exerted antitumor activity in LYN-expressing breast cancer cells<sup>[68]</sup>.

## MCL1 Amplification

### Biological Impact

The myeloid cell leukemia 1 (MCL1) gene encodes a member of the BCL2 pro-survival family<sup>[71]</sup>. MCL1 is highly regulated by various oncogenic signaling pathways, including the mitogen-activated protein kinase (MAPK) pathway<sup>[72]</sup>, the mTOR pathway<sup>[73]</sup>, and the phosphatidylinositol-3 kinase (PI3K) pathway<sup>[74]</sup>. Oncogenic roles for MCL1 have been previously suggested by the report of increased rates of lymphoma in transgenic mice<sup>[75]</sup>. Somatic amplification of MCL1 may be a common mechanism in cancer cells to increase cell survival<sup>[76]</sup>. MCL1 overexpression was observed from a retrospective analysis of parotid gland carcinomas, including adenoid cystic carcinoma<sup>[77]</sup>.

## Therapeutic and prognostic relevance

Therapies targeting MCL1 and other BCL2 family members with the pan-BCL2 family inhibitors are currently under investigation<sup>[78]</sup>. A case report has demonstrated clinical efficacy of sorafenib, when combined with vorinostat, in a metastatic triple-negative breast cancer (TNBC) patient with MCL1-amplified tumor<sup>[79]</sup>. Several in vitro studies also showed that sorafenib induces cell death via inhibition of MCL1 expression in multiple cancer types including, hepatocellular carcinoma (HCC), lung cancer, breast cancer, cholangiocarcinoma, endometrial cancer and chronic lymphocytic leukemia<sup>[80][81][82][83][84][85]</sup>. Preclinical studies have also demonstrated the efficacy of regorafenib in reducing MCL1 expression in human colorectal cancer (CRC) cell lines<sup>[86][87]</sup>. In vivo models of colon cancer showed that MCL1 expression is inhibited by targeting of the mTOR pathway using everolimus, promoting increased tumor cell killing of cancers with KRAS or BRAF mutations<sup>[88]</sup>.

# ACT Onco<sup>®</sup> + Report

## MDM4 Amplification

### Biological Impact

The MDM4 gene encodes a E3-ubiquitin ligase that exerts its oncogenic roles via blocking the transactivation and enhancing the degradation of the p53 tumor suppressor<sup>[89][90][91]</sup>. Similar to MDM2, dysregulation of MDM4 has been reported in various tumor types such as retinoblastoma<sup>[92]</sup>, Ewing sarcoma<sup>[93]</sup>, cutaneous melanoma<sup>[94]</sup> and glioma<sup>[95]</sup>.

### Therapeutic and prognostic relevance

Several small molecules that disrupt the interaction between p53 and MDM4, such as CTX1, ATSP-7041, SAH-p53-8, SJ-172550 have demonstrated p53-dependent tumor growth suppression in preclinical models<sup>[96][97][98][99]</sup>. Results from a study showed that patients with MDM2 family amplification, including MDM2 and MDM4, or EGFR aberrations has a poor clinical outcome and significantly increased rate of tumor growth (hyper-progression) after receiving immune checkpoint (PD-1/PD-L1) therapy<sup>[100]</sup>.

## MYC Amplification

### Biological Impact

The v-myc avian myelocytomatosis viral oncogene homolog, also known as c-myc (MYC) gene encodes a transcription factor involved in cellular proliferation, inhibiting exit from the cell cycle, stimulating vascularization and enhancing genomic instability<sup>[101][102][103]</sup>. Dysregulated MYC expression is implicated in a wide range of human cancers<sup>[104]</sup>.

### Therapeutic and prognostic relevance

MYC amplification was associated with better clinical outcome in breast cancer patients treated with FAC (5-fluorouracil, doxorubicin, and cyclophosphamide), CMF (cyclophosphamide, methotrexate and 5-fluorouracil) and P-FEC (paclitaxel followed by 5-fluorouracil, epirubicin and cyclophosphamide) and higher expression of MYC was also associated with a better response rate in platinum-treated ovarian cancer patients<sup>[105][106][107]</sup>.

CDK inhibition using the dinaciclib, a CDK1, 2, 5 and 9 inhibitors, exerted antitumor activity in triple-negative breast cancer (TNBC) tumor xenograft and cell lines with increased activity of the MYC pathway<sup>[108][109]</sup>.

Overexpression of MYC has been reported as a favorable prognostic biomarker in colorectal carcinoma (CRC)<sup>[110][111]</sup>. However, the favorable prognostic value of MYC in CRC is abrogated by the TP53 mutation<sup>[111]</sup>.

MYC amplification with the loss of tumor suppressor pathways such as p53 and RB has been shown to be associated with poor outcomes and was correlated with shortened disease-free survival in breast cancer with BRCA1 deficiency in TNBC<sup>[108][112]</sup>.

## NBN Amplification

### Biological Impact

The NBN gene encodes a component of the MRE11-RAD50-NBN (MRN) complex, which involves in DNA double-strand break sensing and repair<sup>[113]</sup>. NBN mutation is related to Nijmegen breakage syndrome, increased cancer incidence and ionizing radiation sensitivity<sup>[113][114]</sup>. NBN mutations have been found in various cancers, including cholangiocarcinoma, hepatocellular carcinoma<sup>[115]</sup>, prostate cancer<sup>[116]</sup>, leukemia, lymphoma<sup>[117]</sup>, and triple-negative breast cancer<sup>[118]</sup>.

### Therapeutic and prognostic relevance

In a phase II trial (ARIEL2), an ovarian cancer patient harboring a NBN germline mutation showed responses to rucaparib treatment<sup>[119]</sup>. NBN loss of function mutation has been determined as an inclusion criterion for the trial evaluating olaparib efficacy in ovarian cancer<sup>[120]</sup>; the trials evaluating rucaparib efficacy in ovarian cancer<sup>[121]</sup> or prostate



# ACT Onco<sup>®</sup> + Report

cancer<sup>[122]</sup>; the trials evaluating talazoparib efficacy in HER2-negative breast cancer (NCT02401347), prostate cancer (NCT03148795) or lung cancer (NCT03377556), and the trials evaluating niraparib efficacy in pancreatic cancer (NCT03553004) or any malignancy (except prostate) cancer (NCT03207347).

Germline and somatic mutations in homologous recombination genes, including NBN, have been suggested to be prognostic biomarkers for platinum-based treatment response and superior survival in patients with ovarian, fallopian tube, peritoneal carcinomas and pancreatic cancer<sup>[123][124]</sup>.

In a retrospective study of localized prostate cancer, NBN gene amplification has been demonstrated to associate with overall tumor genomic instability and lower biochemical relapse-free rate following image-guided radiotherapy (IGRT)<sup>[125]</sup>.

Another retrospective study showed that amplification of the NBN gene is associated with protein overexpression and mostly correlated with poor prognosis in several cancer types, including ovarian cancer, breast invasive carcinoma, uterine corpus endometrial carcinoma, and sarcoma. Besides, in vivo and in vitro assays demonstrated that amplification of the NBN gene could induce cisplatin and PARP inhibitor resistance in breast and ovarian cancer cells<sup>[126]</sup>.

## NTRK1 Amplification

### Biological Impact

The NTRK1 gene encodes the TRKA (tropomyosin receptor kinase) receptor which plays an important role in the development and function of the nervous system. Gene fusions of NTRK1 lead to constitutive activation of MAP-kinase, PI3-kinase, and PLC-γ pathways, and represent the main molecular alterations with known oncogenic and transforming potential in various malignancies, including soft tissue sarcoma, non-small cell lung cancer (NSCLC), glioblastoma multiforme (GBM), thyroid carcinoma, and pilocytic astrocytomas<sup>[127][128]</sup>. A pan-cancer study (n=1250) demonstrated that 2.2% of the metastatic cancer patients harbored NTRK amplification and NTRK protein overexpression was observed in 14.8% of NTRK-amplified tumors (doi.org/10.23838/pfm.2017.00142).

### Therapeutic and prognostic relevance

Patients with NTRK1 amplification had only limited benefit from larotrectinib treatment according to the few clinical studies. One of them had a partial response with larotrectinib of short duration (3.7 months)<sup>[129]</sup>, and the other one with metastatic NTRK1-amplified (copy number=8) esophageal carcinoma showed clinical efficacy for six weeks, and then a progressive disease of new lesions were observed<sup>[130]</sup>.

## RECQL4 Amplification

### Biological Impact

The RECQL4 gene encodes a member of the RECQ helicase family that plays an important role in DNA replication and various types of DNA repair, including double-strand break repair, nucleotide excision repair, base excision repair, and single-strand repair<sup>[131][132][133][134]</sup>.

### Therapeutic and prognostic relevance

There are currently no therapies targeting RECQL4 mutations. Expression of RECQL4 has been shown to drive cisplatin resistance in gastric cancer cell lines<sup>[135]</sup>. In contrast, RECQL4-deficient breast cancer cell lines were sensitive to cisplatin and PARP inhibitions (AG14361 and olaparib), but demonstrated resistance to taxane<sup>[136]</sup>. In a preclinical study, RECQL4-deficient colon cancer cell lines were also sensitive to cisplatin and olaparib treatment<sup>[137]</sup>. RECQL4 mutations have been selected as an inclusion criteria for the trials examining olaparib efficacy in metastatic urothelial cancer (NCT03448718) and relapsed small cell lung cancer (NCT03009682).

In breast cancers, the amplification of RECQL4 was significantly associated with aggressive tumor behavior, including

# ACT Onco<sup>®</sup> + Report

lymph node positivity, larger tumor size, HER2 overexpression and poor survival<sup>[138]</sup>

## TERT Amplification

### Biological Impact

The TERT gene encodes the catalytic subunit of telomerase, an enzyme that maintains telomere length and genomic integrity<sup>[139]</sup>. Upregulation of TERT promotes cancer development and progression via modulation of Wnt-catenin and nuclear factor kappa B signaling<sup>[140][141]</sup>, and mitochondrial RNA processing<sup>[142]</sup>. Activating mutations in the TERT promoter have been identified in a number of cancer types including melanoma, hepatocellular carcinoma, urothelial carcinoma, medulloblastoma, and glioma whereas TERT gene amplification is implicated in lung cancer, cervical cancer, breast cancer, Merkel cell carcinoma, neuroblastoma and adrenocortical carcinoma<sup>[143][144][145][146][147]</sup>.

### Therapeutic and prognostic relevance

Imetelstat (GRN163L), a telomere inhibitor which has been shown to inhibit cell proliferation in various cancer cell lines and tumor xenografts is currently in clinical trials<sup>[139]</sup>.

TERT gene amplification is an independent poor prognostic marker for disease-free survival in non-small cell lung cancer (NSCLC) and breast cancer<sup>[148][149][150]</sup>.

# ACT Onco<sup>®</sup> + Report

## US FDA-APPROVED DRUG(S)

### Dasatinib (SPRYCEL)

Dasatinib is an oral Bcr-Abl tyrosine kinase inhibitor (inhibits the "Philadelphia chromosome") and Src family tyrosine kinase inhibitor. Dasatinib is produced by Bristol-Myers Squibb and sold under the trade name SPRYCEL.

### - FDA Approval Summary of Dasatinib (SPRYCEL)

<b>DASISION</b> <sup>[151]</sup> NCT00481247	<b>Chronic myeloid leukemia</b> (Approved on 2010/10/28)
	- Dasatinib vs. Imatinib [ORR(%): 76.8 vs. 66.2]
<sup>[152]</sup> NCT00123474	<b>Chronic myeloid leukemia</b> (Approved on 2007/11/08)
	- Dasatinib [ORR(%): 63.0]
<sup>[153]</sup> NCT00123487	<b>Acute lymphocytic leukemia</b> (Approved on 2006/06/28)
	- Dasatinib [ORR(%): 38.0]

### Everolimus (AFINITOR)

Everolimus, a derivative of sirolimus, works as an inhibitor of mammalian target of rapamycin complex 1 (mTORC1) and blocks mTORC1-mediated downstream signals for cell growth, proliferation, and survival. Everolimus is developed and marketed by Novartis under the trade name AFINITOR.

### - FDA Approval Summary of Everolimus (AFINITOR)

<b>RADIANT-4</b> <sup>[154]</sup> NCT01524783	<b>Lung or gastrointestinal neuroendocrine tumor</b> (Approved on 2016/02/26)
	- Everolimus vs. Placebo [PFS(M): 11 vs. 3.9]
<b>BOLERO-2</b> <sup>[155]</sup> NCT00863655	<b>Breast cancer</b> (Approved on 2012/07/20)
	<b>ER+/HER2-</b> Everolimus + exemestane vs. Placebo + exemestane [PFS(M): 7.8 vs. 3.2]
<b>EXIST-2</b> NCT00790400	<b>Tuberous sclerosis complex (tsc)-associated renal angiomyolipoma</b> (Approved on 2012/04/26)
	- Everolimus vs. Placebo [ORR(%): 41.8 vs. 0]
<b>RADIANT-3</b> <sup>[156]</sup> NCT00510068	<b>Pancreatic neuroendocrine tumor</b> (Approved on 2011/05/05)
	- Everolimus vs. Placebo [PFS(M): 11 vs. 4.6]
<b>EXIST-1</b> <sup>[157]</sup> NCT00789828	<b>Subependymal giant cell astrocytoma</b> (Approved on 2010/10/29)
	- Everolimus vs. Placebo [ORR(%): 35.0]
<b>RECORD-1</b> <sup>[158]</sup> NCT00410124	<b>Renal cell carcinoma</b> (Approved on 2009/05/30)
	- Everolimus vs. Placebo [PFS(M): 4.9 vs. 1.9]

# ACT Onco<sup>®</sup> + Report

## Olaparib (LYNPARZA)

Olaparib is an oral, small molecule inhibitor of poly (ADP-ribose) polymerase-1, -2, and -3 (PARP-1, -2, -3). Olaparib is developed by KuDOS Pharmaceuticals and marketed by AstraZeneca under the trade name LYNPARZA.

### - FDA Approval Summary of Olaparib (LYNPARZA)

<b>Olympia</b> NCT02032823	<b>Her2-negative high-risk early breast cancer</b> (Approved on 2022/03/11)
	<b>HER2-/gBRCA mutation</b> Olaparib vs. Placebo [invasive disease-free survival (IDFS)(M): ]
<b>PROfound</b> <sup>[159]</sup> NCT02987543	<b>Prostate cancer</b> (Approved on 2020/05/19)
	<b>HRR genes mutation</b> Olaparib vs. Enzalutamide or abiraterone acetate [PFS(M): 5.8 vs. 3.5]
<b>PAOLA-1</b> <sup>[160]</sup> NCT02477644	<b>Ovarian cancer</b> (Approved on 2020/05/08)
	<b>HRD+</b> Olaparib + bevacizumab vs. Placebo + bevacizumab [PFS(M): 37.2 vs. 17.7]
<b>POLO</b> <sup>[161]</sup> NCT02184195	<b>Pancreatic adenocarcinoma</b> (Approved on 2019/12/27)
	<b>gBRCA mutation</b> Olaparib vs. Placebo [ORR(%): 23.0 vs. 12.0, PFS(M): 7.4 vs. 3.8]
<b>SOLO-1</b> <sup>[162]</sup> NCT01844986	<b>Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma</b> (Approved on 2018/12/19)
	<b>gBRCA mutation or sBRCA mutation</b> Olaparib vs. Placebo [PFS(M): NR vs. 13.8]
<b>OlympiAD</b> <sup>[163]</sup> NCT02000622	<b>Breast cancer</b> (Approved on 2018/02/06)
	<b>HER2-/gBRCA mutation</b> Olaparib vs. Chemotherapy [PFS(M): 7 vs. 4.2]
<b>SOLO-2/ENGOT-Ov21</b> <sup>[164]</sup> NCT01874353	<b>Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma</b> (Approved on 2017/08/17)
	<b>gBRCA mutation</b> Olaparib vs. Placebo [PFS(M): 19.1 vs. 5.5]
<b>Study19</b> <sup>[165]</sup> NCT00753545	<b>Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma</b> (Approved on 2017/08/17)
	- Olaparib vs. Placebo [PFS(M): 8.4 vs. 4.8]

## Regorafenib (STIVARGA)

Regorafenib is a multi-kinase inhibitor which targets angiogenic, stromal and oncogenic receptor tyrosine kinases (RTKs). Regorafenib is developed and marketed by Bayer HealthCare Pharmaceuticals under the trade name STIVARGA.

### - FDA Approval Summary of Regorafenib (STIVARGA)

<b>RESORCE</b> <sup>[166]</sup> NCT01774344	<b>Hepatocellular carcinoma, Hepatocellular carcinoma</b> (Approved on 2017/04/27)
	- Bsc vs. Placebo [OS(M): 10.6 vs. 7.8]
<b>GRID</b> <sup>[167]</sup> NCT01271712	<b>Gastrointestinal stromal tumor</b> (Approved on 2013/02/25)
	- Regorafenib vs. Placebo [PFS(M): 4.8 vs. 0.9]
<b>CORRECT</b> <sup>[168]</sup> NCT01103323	<b>Colorectal cancer</b> (Approved on 2012/09/27)
	- Regorafenib vs. Placebo [OS(M): 6.4 vs. 5]

# ACT Onco<sup>®</sup> + Report

## Rucaparib (RUBRACA)

Rucaparib is an inhibitor of the DNA repair enzyme poly (ADP-ribose) polymerase-1, -2 and -3 (PARP-1, -2, -3). Rucaparib is developed and marketed by Clovis Oncology under the trade name RUBRACA.

### - FDA Approval Summary of Rucaparib (RUBRACA)

<b>TRITON2</b> NCT02952534	<b>Prostate cancer</b> (Approved on 2020/05/15)
	<b>gBRCA mutation or sBRCA mutation</b>
	Rucaparib [ORR(%): 44.0, DOR(M): NE]
<b>ARIEL3</b> <sup>[121]</sup> NCT01968213	<b>Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma</b> (Approved on 2018/04/06)
	-
	Rucaparib vs. Placebo [PFS (All)(M): 10.8 vs. 5.4, PFS (HRD)(M): 13.6 vs. 5.4, PFS (tBRCA)(M): 16.6 vs. 5.4]

## Sorafenib (NEXAVAR)

Sorafenib is a small molecule multi-kinase inhibitor that targets multiple kinase families including VEGFR, PDGFRB, and the RAF family kinases. Sorafenib is co-developed and co-marketed by Bayer HealthCare Pharmaceuticals and Onyx Pharmaceuticals under the trade name NEXAVAR.

### - FDA Approval Summary of Sorafenib (NEXAVAR)

<b>DECISION</b> <sup>[169]</sup> NCT00984282	<b>Differentiated thyroid carcinoma</b> (Approved on 2013/11/22)
	-
	Sorafenib vs. Placebo [PFS(M): 10.8 vs. 5.8]
<b>SHARP</b> <sup>[170]</sup> NCT00105443	<b>Hepatocellular carcinoma</b> (Approved on 2007/11/16)
	-
	Sorafenib vs. Placebo [OS(M): 10.7 vs. 7.9]
<b>TARGET</b> <sup>[171]</sup> NCT00073307	<b>Renal cell carcinoma</b> (Approved on 2005/12/20)
	-
	Sorafenib vs. Placebo [PFS(D): 167 vs. 84]

## Talazoparib (TALZENNA)

Talazoparib is an inhibitor of poly (ADP-ribose) polymerase (PARP) enzymes, including PARP1 and PARP2. Talazoparib is developed and marketed by Pfizer under the trade name TALZENNA.

### - FDA Approval Summary of Talazoparib (TALZENNA)

<b>EMBRACA</b> <sup>[172]</sup> NCT01945775	<b>Breast cancer</b> (Approved on 2018/10/16)
	<b>HER2-/gBRCA mutation</b>
	Talazoparib vs. Chemotherapy [PFS(M): 8.6 vs. 5.6]

# ACT Onco<sup>®</sup> + Report

## Temsirolimus (TORISEL)

Temsirolimus is a soluble ester of sirolimus (rapamycin, brand-name drug Rapamune) and functions as an inhibitor of mammalian target of rapamycin complex (mTORC). The inhibitory molecular mechanism is similar to Everolimus. Temsirolimus is developed by Wyeth Pharmaceuticals and marketed by Pfizer under the trade name TORISEL.

## - FDA Approval Summary of Temsirolimus (TORISEL)

[173]	<b>Renal cell carcinoma</b> (Approved on 2007/05/30)
NCT00065468	-
	Temsirolimus vs. Ifn- $\alpha$ [OS(M): 10.9 vs. 7.3]

D=day; W=week; M=month



# ACT Onco<sup>®</sup> + Report

## ONGOING CLINICAL TRIALS

Trials were searched by applying filters: study status, patient's diagnosis, intervention, location and/or biomarker(s). Please visit <https://clinicaltrials.gov> to search and view for a complete list of open available and updated matched trials.

No trial has been found.

# ACT Onco<sup>®</sup> + Report


## SUPPLEMENTARY INFORMATION OF TESTING RESULTS DETAILED INFORMATION OF VARIANTS WITH CLINICAL RELEVANCE

### - Single Nucleotide and Small InDel Variants

Gene	Amino Acid Change	Exon	cDNA Change	Accession Number	COSMIC ID	Allele Frequency	Coverage
ARID1A	I1816fs	20	c.5447_5448del	NM_006015	-	21.4%	856
CTNNB1	S33Y	3	c.98C>A	NM_001904	COSM5673	35.1%	903
TSC1	Splice donor	-	c.1997+1G>C	NM_000368	-	11.6%	622

### - Copy Number Alterations

Observed copy number (CN) for each evaluated position is shown on the y-axis. Regions referred to as amplification or deletion are shown in color. Regions without significant changes are represented in gray.



# ACT Onco<sup>®</sup> + Report

## OTHER DETECTED VARIANTS

Gene	Amino Acid Change	Exon	cDNA Change	Accession Number	COSMIC ID	Allele Frequency	Coverage
ADAMTS18	Splice region	18	c.2799T>A	NM_199355	-	31.6%	792
AXL	Y409F	9	c.1226A>T	NM_021913	-	39.7%	453
CCNB2	V301G	7	c.902T>G	NM_004701	-	51.9%	1369
CYP3A4	I118V	5	c.352A>G	NM_017460	-	71.2%	59
GNAS	R645H	1	c.1934G>A	NM_080425	-	23.2%	719
GNAS	P96A	1	c.286C>G	NM_080425	-	51.8%	386
HIF1A	A593P	12	c.1777G>C	NM_001530	-	48.9%	712
JAK2	V392M	9	c.1174G>A	NM_004972	COSM5979661	52.4%	820
KIT	K199E	3	c.595A>G	NM_000222	-	23.0%	313
KMT2A	S2445Y	27	c.7334C>A	NM_001197104	-	27.2%	848
LRP1B	F1264L	24	c.3792T>A	NM_018557	-	8.2%	852
MAPK1	Splice region	-	c.119+8T>C	NM_002745	-	18.5%	108
MET	R591Q	6	c.1772G>A	NM_001127500	COSM7000463	39.1%	494
NBN	Q494L	11	c.1481A>T	NM_002485	-	40.3%	1047
PTGS2	L467H	9	c.1400T>A	NM_000963	-	18.7%	1076
RAD50	M640V	14	c.1918A>G	NM_005732	-	32.2%	881
TEK	Q677K	13	c.2029C>A	NM_000459	-	43.8%	406



### Note:

- This table enlists variants detected by the panel other than those with clinical relevance (reported in Testing Result section). The clinical impact of a genetic variant is determined according to ACT Genomics in-house clinical knowledge database. A negative result does not necessarily indicate absence of biological effect on the tumor. Some variants listed here may possibly have preclinical data or may show potential clinical relevance in the future.

# ACT Onco<sup>®</sup> + Report

## TEST DETAILS

### SPECIMEN RECEIVED AND PATHOLOGY REVIEW



- Collection date: Dec 30, 2022
- Facility retrieved: 臺北榮總
- H&E-stained section No.: S11155447A
- Collection site: Bone
- Examined by: Dr. Yun-An Chen
  1. The percentage of viable tumor cells in total cells in the whole slide (%): 75%
  2. The percentage of viable tumor cells in total cells in the encircled areas in the whole slide (%): 75%
  3. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the whole slide (%): 15%
  4. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the encircled areas in the whole slide (%): 15%
  5. Additional comment: N/A
- Manual macrodissection: Not performed
- The outline highlights the area of malignant neoplasm annotated by a pathologist.

## RUN QC

- Panel: ACTOnco<sup>®</sup>+

### DNA test

- Mean Depth: 819x
- Target Base Coverage at 100x: 95%

### RNA test

- Average unique RNA Start Sites per control GSP2: 187

# ACT Onco<sup>®</sup> + Report

## LIMITATIONS

1. This test does not provide information of variant causality and does not detect variants in non-coding regions that could affect gene expression. This report does not report polymorphisms and we do not classify whether a mutation is germline or somatic. Variants identified by this assay were not subject to validation by Sanger or other technologies.
2. The possibility cannot be excluded that certain pathogenic variants detected by other sequencing tools may not be reported in the test because of technical limitation of bioinformatics algorithm or the NGS sequencing platform, e.g. low coverage.
3. This test has been designed to detect fusions in 13 genes sequenced. Therefore, fusion in genes not covered by this test would not be reported. For novel fusions detected in this test, Sanger sequencing confirmation is recommended if residue specimen is available.

## NEXT-GENERATION SEQUENCING (NGS) METHODS

### DNA test

Extracted genomic DNA was amplified using primers targeting coding exons of analyzed genes and subjected to library construction. Barcoded libraries were subsequently conjugated with sequencing beads by emulsion PCR and enriched using Ion Chef system. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific).

Raw reads generated by the sequencer were mapped to the hg19 reference genome using the Ion Torrent Suite. Coverage depth was calculated using Torrent Coverage Analysis plug-in. Single nucleotide variants (SNVs) and short insertions/deletions (InDels) were identified using the Torrent Variant Caller plug-in. VEP (Variant Effect Predictor) was used to annotate every variant using databases from Clinvar, COSMIC and Genome Aggregation database. Variants with coverage  $\geq 20$ , allele frequency  $\geq 5\%$  and actionable variants with allele frequency  $\geq 2\%$  were retained. This test provides uniform coverage of the targeted regions, enabling target base coverage at  $100\times \geq 85\%$  with a mean coverage  $\geq 500\times$ .

Variants reported in Genome Aggregation database with  $> 1\%$  minor allele frequency (MAF) were considered as polymorphisms. ACT Genomics in-house database was used to determine technical errors. Clinically actionable and biologically significant variants were determined based on the published medical literature.

The copy number alterations (CNAs) were predicted as described below:

Amplicons with read counts in the lowest 5th percentile of all detectable amplicons and amplicons with a coefficient of variation  $\geq 0.3$  were removed. The remaining amplicons were normalized to correct the pool design bias. ONCOCNV (an established method for calculating copy number aberrations in amplicon sequencing data by Boeva et al., 2014) was applied for the normalization of total amplicon number, amplicon GC content, amplicon length, and technology-related biases, followed by segmenting the sample with a gene-aware model. The method was used as well for establishing the baseline of copy number variations.

Tumor mutational burden (TMB) was calculated by using the sequenced regions of ACTOnco<sup>®</sup>+ to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The TMB calculation predicted somatic variants and applied a machine learning model with a cancer hotspot correction. TMB may be reported as "TMB-High", "TMB-Low" or "Cannot Be Determined". TMB-High corresponds to  $\geq 7.5$  mutations per megabase (Muts/Mb); TMB-Low corresponds to  $< 7.5$  Muts/Mb. TMB is reported as "Cannot Be Determined" if the tumor purity of the sample is  $< 30\%$ .

Classification of microsatellite instability (MSI) status is determined by a machine learning prediction algorithm. The change of a number of repeats of different lengths from a pooled microsatellite stable (MSS) baseline in  $> 400$  genomic loci are used as the features for the algorithm. The final output of the results is either microsatellite Stable (MSS) or microsatellite instability high (MSI-H).

### RNA test

Extracted RNA was reverse-transcribed and subjected to library construction. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific). To ensure sequencing quality for fusion variant analysis, the average unique RNA Start Sites (SS) per control Gene Specific Primer 2 (GSP 2) should be  $\geq 10$ .

# ACT Onco<sup>®</sup> + Report

The fusion analysis pipeline aligned sequenced reads to the human reference genome, identified regions that map to noncontiguous regions of the genome, applied filters to exclude probable false-positive events and, annotated previously characterized fusion events according to Quiver Gene Fusion Database, a curated database owned and maintained by ArcherDX. In general, samples with detectable fusions need to meet the following criteria: (1) Number of unique start sites (SS) for the GSP2  $\geq 3$ ; (2) Number of supporting reads spanning the fusion junction  $\geq 5$ ; (3) Percentage of supporting reads spanning the fusion junction  $\geq 10\%$ ; (4) Fusions annotated in Quiver Gene Fusion Database.

## DATABASE USED

- Reference genome: Human genome sequence hg19
- COSMIC v.92
- Genome Aggregation database r2.1.1
- ClinVar (version 20210404)
- ACT Genomics in-house database
- Quiver Gene Fusion Database version 5.1.18

## Variant Analysis:

醫檢師陳韻鈺 博士  
Yun-Yu Chen Ph.D.  
檢字第 015647 號

Yun Yu Chen

## Sign Off

解剖病理專科醫師王業翰  
Yeh-Han Wang M.D.  
病解字第 000545 號

Yeh



# ACT Onco<sup>®</sup> + Report

## GENE LIST SNV & CNV

ABCB1*	ABCC2*	ABCG2*	ABL1	ABL2	ADAMTS1	ADAMTS13	ADAMTS15	ADAMTS16	ADAMTS18	ADAMTS6	ADAMTS9
ADAMTSL1	ADGRA2	ADH1C*	AKT1	AKT2	AKT3	ALDH1A1*	ALK	AMER1	APC	AR	ARAF
ARID1A	ARID1B	ARID2	ASXL1	ATM	ATR	ATRX	AURKA	AURKB	AXIN1	AXIN2	AXL
B2M	BAP1	BARD1	BCL10	BCL2*	BCL2L1	BCL2L2*	BCL6	BCL9	BCOR	BIRC2	BIRC3
BLM	BMPR1A	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2*	BTB	BUB1B	CALR
CANX	CARD11	CASP8	CBFB	CBL	CCNA1	CCNA	CCNB1	CCNB2	CCNB3	CCND1	CCND2
CCND3	CCNE1	CCNE2	CCNH	CD19	CD274	CD58	CD70*	CD79A	CD79B	CDC73	CDH1
CDK1	CDK12	CDK2	CDK4	CDK5	CDK6	CDK7	CDK8	CDK9	CDKN1A	CDKN1B	CDKN2A
CDKN2B	CDKN2C	CEBPA*	CHEK1	CHEK2	CIC	CREBBP	CRKL	CRLF2	CSF1R	CTCF	CTLA4
CTNNA1	CTNNB1	CUL3	CYLD	CYP1A1*	CYP2B6*	CYP2C19*	CYP2C8*	CYP2D6	CYP2E1*	CYP3A4*	CYP3A5*
DAXX	DCUN1D1	DDR2	DICER1	DNMT3A	DOT1L	DPYD	DTX1	E2F3	EGFR	EP300	EPCAM
EPHA2	EPHA3	EPHA5	EPHA7	EPHB1	ERBB2	ERBB3	ERBB4	ERCC1	ERCC2	ERCC3	ERCC4
ERCC5	ERG	ESR1	ESR2	ETV1	ETV4	EZH2	FAM46C	FANCA	FANCC	FANCD2	FANCE
FANCF	FANCG	FANCL	FAS	FAT1	FBXW7	FCGR2B	FGF1*	FGF10	FGF14	FGF19*	FGF23
FGF3	FGF4*	FGF6	FGFR1	FGFR2	FGFR3	FGFR4	FH	FLCN	FLT1	FLT3	FLT4
FOXL2*	FOXP1	FRG1	FUBP1	GATA1	GATA2	GATA3	GNA11	GNA13	GNAQ	GNAS	GREM1
GRIN2A	GSK3B	GSTP1*	GSTT1*	HGF	HIF1A	HIST1H1C*	HIST1H1E*	HNF1A	HR	HRAS*	HSP90AA1
HSP90AB1	HSPA4	HSPA5	IDH1	IDH2	IFNL3*	IGF1	IGF1R	IGF2	IKBKB	IKBKE	IKZF1
IL6	IL7R	INPP4B	INSR	IRF4	IRS1	IRS2*	JAK1	JAK2	JAK3	JUN*	KAT6A
KDM5A	KDM5C	KDM6A	KDR	KEAP1	KIT	KMT2A	KMT2C	KMT2D	KRAS	LCK	LIG1
LIG3	LMO1	LRP1B	LYN	MALT1	MAP2K1	MAP2K2	MAP2K4	MAP3K1	MAP3K7	MAPK1	MAPK3
MAX	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1	MET	MITF	MLH1	MPL	MRE11
MSH2	MSH6	MTHFR*	MTOR	MUC16	MUC4	MUC6	MUTYH	MYC	MYCL	MYCN	MYD88
NAT2*	NBN	NEFH	NF1	NF2	NFE2L2	NFKB1	NFKBIA	NKX2-1*	NOTCH1	NOTCH2	NOTCH3
NOTCH4	NPM1	NQO1*	NRAS	NSD1	NTRK1	NTRK2	NTRK3	PAK3	PALB2	PARP1	PAX5
PAX8	PBRM1	PDCD1	PDCD1LG2	PDGFRA	PDGFRB	PDIA3	PGF	PHOX2B*	PIK3C2B	PIK3C2G	PIK3C3
PIK3CA	PIK3CB	PIK3CD	PIK3CG	PIK3R1	PIK3R2	PIK3R3	PIM1	PMS1	PMS2	POLB	POLD1
POLE	PPARG	PPP2R1A	PRDM1	PRKAR1A	PRKCA	PRKCB	PRKCG	PRKCI	PRKCQ	PRKDC	PRKN
PSMB8	PSMB9	PSME1	PSME2	PSME3	PTCH1	PTEN	PTGS2	PTPN11	PTPRD	PTPRT	RAC1
RAD50	RAD51	RAD51B	RAD51C	RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	RECQL4
REL	RET	RHOA	RICTOR	RNF43	ROS1	RPPH1	RPTOR	RUNX1	RUNX1T1	RXRA	SDHA
SDHB	SDHC	SDHD	SERPINB3	SERPINB4	SETD2	SF3B1	SGK1	SH2D1A*	SLC19A1*	SLC22A2*	SLC1B1*
SLC1B3*	SMAD2	SMAD3	SMAD4	SMARCA4	SMARCB1	SMO	SOC1*	SOX2*	SOX9	SPEN	SPOP
SRC	STAG2	STAT3	STK11	SUFU	SYK	SYNE1	TAF1	TAP1	TAP2	TAPBP	TBX3
TEK	TERT	TET1	TET2	TGFB2	TMSB4X*	TNF	TNFAIP3	TNFRSF14	TNFSF11	TOP1	TP53
TPMT*	TSC1	TSC2	TSHR	TYMS	U2AF1	UBE2A*	UBE2K	UBR5	UGT1A1*	USH2A	VDR*
VEGFA	VEGFB	VHL	WT1	XIAP	XPO1	XRCC2	ZNF217				

\*Analysis of copy number alterations NOT available.

## FUSION

ALK	BRAF	EGFR	FGFR1	FGFR2	FGFR3	MET	NRG1	NTRK1	NTRK2	NTRK3	RET	ROS1
-----	------	------	-------	-------	-------	-----	------	-------	-------	-------	-----	------


# ACT Onco<sup>®</sup> + Report

## APPENDIX

### POSSIBLE THERAPEUTIC IMPLICATIONS FOR HETEROZYGOUS DELETION


Not Applicable.

### SIGNALING PATHWAYS AND MOLECULAR-TARGETED AGENTS




1: Everolimus, Temsirolimus; 2: Sorafenib

### Receptor Tyrosine Kinase/Growth Factor Signalling



1: Dasatinib, Regorafenib; 2: Dasatinib, Sorafenib, Regorafenib

# ACT Onco<sup>®</sup> + Report



1: Olaparib, Rucaparib, Talazoparib

# ACT Onco<sup>®</sup> + Report

## DISCLAIMER

### 法律聲明

本檢驗報告僅提供專業醫療參考，結果需經專業醫師解釋及判讀。基因突變資訊非必具備藥物或治療有效性指標，反之亦然。本檢驗報告提供之用藥指引不聲明或保證其臨床有效性，反之亦然。本基因檢測方法係由本公司研究開發，已經過有效性測試。

本檢驗報告非經本公司許可，不得私自變造、塗改，或以任何方式作為廣告及其他宣傳之用途。

本公司於提供檢驗報告後，即已完成本次契約義務，後續之報告解釋、判讀及用藥、治療，應自行尋求相關專業醫師協助，若需將報告移件其他醫師，本人應取得該醫師同意並填寫移件申請書，主動告知行動基因，行動基因僅能配合該醫師意願與時間提供醫師解說。

### 醫療決策需由醫師決定

任何治療與用藥需經由醫師在考慮病患所有健康狀況相關資訊包含健檢、其他檢測報告和病患意願後，依照該地區醫療照護標準由醫師獨立判斷。醫師不應僅依據單一報告結果(例如本檢測或本報告書內容)做決策。

### 基因突變與用藥資訊並非依照有效性排序

本報告中列出之生物標記變異與藥物資訊並非依照潛在治療有效性排序。

### 證據等級

藥物潛在臨床效益(或缺乏潛在臨床效益)的實證證據是依據至少一篇臨床療效個案報告或臨床前試驗做為評估。本公司盡力提供適時及準確之資料，但由於醫學科技之發展日新月異，本公司不就本報告提供的資料是否為準確、適宜或最新作保證。

### 責任

本檢驗報告僅提供專業醫療參考，本公司及其員工不對任何由使用本報告之內容引起的直接、間接、特殊、連帶或衍生的損失或損害承擔責任。

# ACT Onco<sup>®</sup> + Report

## REFERENCE

1. PMID: 10757798; 2000, Mol Cell Biol;20(9):3137-46  
The human SWI-SNF complex protein p270 is an ARID family member with non-sequence-specific DNA binding activity.
2. PMID: 25387058; 2015, Annu Rev Pathol;10():145-71  
SWI/SNF chromatin remodeling and human malignancies.
3. PMID: 23208470; 2013, Cancer Discov;3(1):35-43  
ARID1A mutations in cancer: another epigenetic tumor suppressor?
4. PMID: 20826764; 2010, Science;330(6001):228-31  
Frequent mutations of chromatin remodeling gene ARID1A in ovarian clear cell carcinoma.
5. PMID: 20942669; 2010, N Engl J Med;363(16):1532-43  
ARID1A mutations in endometriosis-associated ovarian carcinomas.
6. PMID: 21590771; 2011, J Pathol;224(3):328-33  
Loss of BAF250a (ARID1A) is frequent in high-grade endometrial carcinomas.
7. PMID: 21412130; 2011, Am J Surg Pathol;35(5):625-32  
Mutation and loss of expression of ARID1A in uterine low-grade endometrioid carcinoma.
8. PMID: 22037554; 2011, Nat Genet;43(12):1219-23  
Exome sequencing identifies frequent mutation of ARID1A in molecular subtypes of gastric cancer.
9. PMID: 26125128; 2015, Expert Opin Ther Targets;19(11):1419-22  
Potential therapeutic targets in ARID1A-mutated cancers.
10. PMID: 29093822; 2017, Gynecol Oncol Res Pract;4():17  
EZH2 inhibition in ARID1A mutated clear cell and endometrioid ovarian and endometrioid endometrial cancers.
11. PMID: 24979463; 2014, Oncotarget;5(14):5295-303  
Loss of ARID1A expression sensitizes cancer cells to PI3K- and AKT-inhibition.
12. PMID: 27364904; 2016, Mol Cancer Ther;15(7):1472-84  
Synthetic Lethal Targeting of ARID1A-Mutant Ovarian Clear Cell Tumors with Dasatinib.
13. PMID: 27172896; 2016, Clin Cancer Res;22(21):5238-5248  
Loss of ARID1A Activates ANXA1, which Serves as a Predictive Biomarker for Trastuzumab Resistance.
14. PMID: 22101352; 2012, Mod Pathol;25(2):282-8  
Loss of ARID1A expression is related to shorter progression-free survival and chemoresistance in ovarian clear cell carcinoma.
15. PMID: 24459582; 2014, J Gynecol Oncol;25(1):58-63  
Decreased ARID1A expression is correlated with chemoresistance in epithelial ovarian cancer.
16. PMID: 26770240; 2015, J Breast Cancer;18(4):339-46  
Loss of Tumor Suppressor ARID1A Protein Expression Correlates with Poor Prognosis in Patients with Primary Breast Cancer.
17. PMID: 21889920; 2012, Cancer Epidemiol;36(3):288-93  
Frequent low expression of chromatin remodeling gene ARID1A in breast cancer and its clinical significance.
18. PMID: 25311944; 2014, Hum Pathol;45(12):2430-6  
Immunohistochemical detection of ARID1A in colorectal carcinoma: loss of staining is associated with sporadic microsatellite unstable tumors with medullary histology and high TNM stage.
19. PMID: 25561809; 2014, World J Gastroenterol;20(48):18404-12  
Clinicopathologic and prognostic relevance of ARID1A protein loss in colorectal cancer.

# ACT Onco<sup>®</sup> + Report

20. PMID: 26069190; 2015, Cancer Discov;5(7):752-67  
ARID1A Deficiency Impairs the DNA Damage Checkpoint and Sensitizes Cells to PARP Inhibitors.
21. PMID: 22682243; 2012, Cell;149(6):1192-205  
Wnt/ $\beta$ -catenin signaling and disease.
22. PMID: 22617422; 2012, EMBO J;31(12):2714-36  
The many faces and functions of  $\beta$ -catenin.
23. PMID: 10201372; 1999, Nature;398(6726):422-6  
Beta-catenin regulates expression of cyclin D1 in colon carcinoma cells.
24. PMID: 9727977; 1998, Science;281(5382):1509-12  
Identification of c-MYC as a target of the APC pathway.
25. PMID: 23788652; 2013, Genome Res;23(9):1422-33  
Whole-genome sequencing identifies recurrent mutations in hepatocellular carcinoma.
26. PMID: 22634756; 2012, Nat Genet;44(7):760-4  
Whole-genome sequencing of liver cancers identifies etiological influences on mutation patterns and recurrent mutations in chromatin regulators.
27. PMID: 23636398; 2013, Nature;497(7447):67-73  
Integrated genomic characterization of endometrial carcinoma.
28. PMID: 22810696; 2012, Nature;487(7407):330-7  
Comprehensive molecular characterization of human colon and rectal cancer.
29. PMID: 25079552; 2014, Nature;511(7511):543-50  
Comprehensive molecular profiling of lung adenocarcinoma.
30. PMID: 22980975; 2012, Cell;150(6):1107-20  
Mapping the hallmarks of lung adenocarcinoma with massively parallel sequencing.
31. PMID: 17187432; 2007, Hepatology;45(1):42-52  
Transcriptome classification of HCC is related to gene alterations and to new therapeutic targets.
32. PMID: 17487939; 2007, J Pathol;212(3):345-52  
Cholestasis is a marker for hepatocellular carcinomas displaying beta-catenin mutations.
33. PMID: 19101982; 2009, Hepatology;49(3):821-31  
Unique phenotype of hepatocellular cancers with exon-3 mutations in beta-catenin gene.
34. PMID: 26171210; 2015, Mol Clin Oncol;3(4):936-940  
 $\beta$ -catenin mutation is correlated with a favorable prognosis in patients with hepatocellular carcinoma.
35. PMID: 11957146; 2002, Hum Pathol;33(2):206-12  
CTNNB1 mutations and beta-catenin expression in endometrial carcinomas.
36. PMID: 11955436; 2002, Cell;108(6):837-47  
Control of beta-catenin phosphorylation/degradation by a dual-kinase mechanism.
37. PMID: 15064718; 2004, Oncogene;23(28):4839-46  
Phosphorylation and ubiquitination of oncogenic mutants of beta-catenin containing substitutions at Asp32.
38. PMID: 18757411; 2008, Cancer Res;68(17):6969-77  
Illegitimate WNT pathway activation by beta-catenin mutation or autocrine stimulation in T-cell malignancies.
39. PMID: 26861905; 2016, Ann Surg Oncol;23(6):1924-7



# ACT Onco<sup>®</sup> + Report

Correlation of CTNNB1 Mutation Status with Progression Arrest Rate in RECIST Progressive Desmoid-Type Fibromatosis Treated with Imatinib: Translational Research Results from a Phase 2 Study of the German Interdisciplinary Sarcoma Group (GISG-01).

40. PMID: 27016228; 2016, Gynecol Oncol;141(1):43-8  
Tumor mutational analysis of GOG248, a phase II study of temsirolimus or temsirolimus and alternating megestrol acetate and tamoxifen for advanced endometrial cancer (EC): An NRG Oncology/Gynecologic Oncology Group study.
41. PMID: 25624430; 2015, J Clin Oncol;33(8):930-6  
Phase II study of everolimus and letrozole in patients with recurrent endometrial carcinoma.
42. PMID: 25214561; 2014, J Natl Cancer Inst;106(9):  
Clinical significance of CTNNB1 mutation and Wnt pathway activation in endometrioid endometrial carcinoma.
43. PMID: 21157483; 2011, Nat Rev Mol Cell Biol;12(1):21-35  
mTOR: from growth signal integration to cancer, diabetes and ageing.
44. PMID: 12271141; 2002, Proc Natl Acad Sci U S A;99(21):13571-6  
Tuberous sclerosis complex-1 and -2 gene products function together to inhibit mammalian target of rapamycin (mTOR)-mediated downstream signaling.
45. PMID: 9242607; 1997, Science;277(5327):805-8  
Identification of the tuberous sclerosis gene TSC1 on chromosome 9q34.
46. PMID: 8269512; 1993, Cell;75(7):1305-15  
Identification and characterization of the tuberous sclerosis gene on chromosome 16.
47. PMID: 1303246; 1992, Nat Genet;2(1):37-41  
Linkage of an important gene locus for tuberous sclerosis to a chromosome 16 marker for polycystic kidney disease.
48. PMID: 18538015; 2008, BMC Cancer;8():163  
Involvement of TSC genes and differential expression of other members of the mTOR signaling pathway in oral squamous cell carcinoma.
49. PMID: 28339086; 2017, Int J Oncol;50(5):1778-1784  
Identification of novel mutations in endometrial cancer patients by whole-exome sequencing.
50. PMID: 20610279; 2010, Urol Oncol;28(4):409-28  
Bladder cancer or bladder cancers? Genetically distinct malignant conditions of the urothelium.
51. PMID: 17005952; 2006, N Engl J Med;355(13):1345-56  
The tuberous sclerosis complex.
52. PMID: 22923433; 2012, Science;338(6104):221  
Genome sequencing identifies a basis for everolimus sensitivity.
53. PMID: 26859683; 2016, Oncotarget;7(9):10547-56  
Next-generation sequencing reveals somatic mutations that confer exceptional response to everolimus.
54. PMID: 26412398; 2015, Sci Rep;5():14534  
PAK2 is an effector of TSC1/2 signaling independent of mTOR and a potential therapeutic target for Tuberous Sclerosis Complex.
55. PMID: 10733529; 2000, Genes Dev;14(6):690-703  
E2f3 is critical for normal cellular proliferation.
56. PMID: 24901045; 2014, Cell Death Dis;5():e1274  
Retinoblastoma protein (RB) interacts with E2F3 to control terminal differentiation of Sertoli cells.
57. PMID: 31053804; 2019, Nat Rev Cancer;19(6):326-338  
The broken cycle: E2F dysfunction in cancer.
58. PMID: 12782593; 2003, Cancer Res;63(11):2872-80

# ACT Onco<sup>®</sup> + Report

Array-based comparative genomic hybridization for genome-wide screening of DNA copy number in bladder tumors.

59. PMID: 15122326; 2004, Oncogene;23(33):5616-23  
E2F3 amplification and overexpression is associated with invasive tumor growth and rapid tumor cell proliferation in urinary bladder cancer.
60. PMID: 28134624; 2017, J Clin Invest;127(3):830-842  
Dosage-dependent copy number gains in E2f1 and E2f3 drive hepatocellular carcinoma.
61. PMID: 16007192; 2005, Oncogene;24(42):6441-9  
Gains and overexpression identify DEK and E2F3 as targets of chromosome 6p gains in retinoblastoma.
62. PMID: 30214236; 2018, Onco Targets Ther;11():5303-5313  
E2F3 promotes cancer growth and is overexpressed through copy number variation in human melanoma.
63. PMID: 9202419; 1997, Int J Biochem Cell Biol;29(3):397-400  
Lyn, a src-like tyrosine kinase.
64. PMID: 19290919; 2009, Immunol Rev;228(1):23-40  
Multiple roles of Lyn kinase in myeloid cell signaling and function.
65. PMID: 11825908; 2002, J Biol Chem;277(16):13732-8  
Phosphatidylinositol 3-kinase and Src family kinases are required for phosphorylation and membrane recruitment of Dok-1 in c-Kit signaling.
66. PMID: 11435302; 2001, Blood;98(2):343-50  
Lyn is required for normal stem cell factor-induced proliferation and chemotaxis of primary hematopoietic cells.
67. PMID: 10574731; 1999, J Exp Biol;202 Pt 24():3541-53  
Redox control in development and evolution: evidence from colonial hydroids
68. PMID: 20215510; 2010, Cancer Res;70(6):2296-306  
LYN is a mediator of epithelial-mesenchymal transition and a target of dasatinib in breast cancer.
69. PMID: 22490227; 2012, Clin Cancer Res;18(10):2850-60  
Lyn kinase mediates cell motility and tumor growth in EGFRvIII-expressing head and neck cancer.
70. PMID: 22731636; 2012, Mol Cancer;11():43  
Lyn is involved in CD24-induced ERK1/2 activation in colorectal cancer.
71. PMID: 18955968; 2008, Oncogene;27(50):6398-406  
Bcl-2 family proteins and cancer.
72. PMID: 18676833; 2008, Cancer Res;68(15):6109-17  
Down-regulation of myeloid cell leukemia-1 through inhibiting Erk/Pin 1 pathway by sorafenib facilitates chemosensitization in breast cancer.
73. PMID: 18664580; 2008, Proc Natl Acad Sci U S A;105(31):10853-8  
mTORC1 promotes survival through translational control of Mcl-1.
74. PMID: 11314001; 2001, Oncogene;20(6):677-85  
The involvement of PI 3-K/Akt-dependent up-regulation of Mcl-1 in the prevention of apoptosis of Hep3B cells by interleukin-6.
75. PMID: 11389033; 2001, Blood;97(12):3902-9  
MCL1 transgenic mice exhibit a high incidence of B-cell lymphoma manifested as a spectrum of histologic subtypes.
76. PMID: 20164920; 2010, Nature;463(7283):899-905  
The landscape of somatic copy-number alteration across human cancers.
77. PMID: 21734342; 2011, Dis Markers;30(5):229-33  
Mcl-1 expression is up-regulated in malignancies of the parotid gland.
78. PMID: 26045609; 2015, Blood;126(3):363-72

# ACT Onco<sup>®</sup> + Report

Inhibition of Mcl-1 with the pan-Bcl-2 family inhibitor (-)BI97D6 overcomes ABT-737 resistance in acute myeloid leukemia.

79. PMID: 27293397; 2016, Case Rep Oncol;9(1):112-8  
A Combination of Targeted Therapy with Chemotherapy Backbone Induces Response in a Treatment-Resistant Triple-Negative MCL1-Amplified Metastatic Breast Cancer Patient.
80. PMID: 17178882; 2006, Cancer Res;66(24):11851-8  
Sorafenib blocks the RAF/MEK/ERK pathway, inhibits tumor angiogenesis, and induces tumor cell apoptosis in hepatocellular carcinoma model PLC/PRF/5.
81. PMID: 23392173; 2013, Cell Death Dis;4():e485  
Mcl-1-dependent activation of Beclin 1 mediates autophagic cell death induced by sorafenib and SC-59 in hepatocellular carcinoma cells.
82. PMID: 16007148; 2005, Oncogene;24(46):6861-9  
The role of Mcl-1 downregulation in the proapoptotic activity of the multikinase inhibitor BAY 43-9006.
83. PMID: 19821497; 2009, Hepatology;50(6):1861-70  
Sorafenib inhibits signal transducer and activator of transcription-3 signaling in cholangiocarcinoma cells by activating the phosphatase shatterproof 2.
84. PMID: 20071162; 2010, Eur J Cancer;46(4):836-50  
The multikinase inhibitor Sorafenib induces apoptosis and sensitises endometrial cancer cells to TRAIL by different mechanisms.
85. PMID: 21979753; 2012, Mol Med;18():19-28  
Sorafenib-induced apoptosis of chronic lymphocytic leukemia cells is associated with downregulation of RAF and myeloid cell leukemia sequence 1 (Mcl-1).
86. PMID: 24763611; 2014, Clin Cancer Res;20(13):3472-84  
Regorafenib inhibits colorectal tumor growth through PUMA-mediated apoptosis.
87. PMID: 26561209; 2015, Cancer Biol Ther;16(12):1710-9  
Regorafenib with a fluoropyrimidine for metastatic colorectal cancer after progression on multiple 5-FU-containing combination therapies and regorafenib monotherapy.
88. PMID: 24163374; 2014, Cancer Discov;4(1):42-52  
mTOR inhibition specifically sensitizes colorectal cancers with KRAS or BRAF mutations to BCL-2/BCL-XL inhibition by suppressing MCL-1.
89. PMID: 21730163; 2011, Proc Natl Acad Sci U S A;108(29):12001-6  
The p53 inhibitors MDM2/MDMX complex is required for control of p53 activity in vivo.
90. PMID: 14507994; 2003, Proc Natl Acad Sci U S A;100(21):12009-14  
HdmX stimulates Hdm2-mediated ubiquitination and degradation of p53.
91. PMID: 12162806; 2002, DNA Cell Biol;21(7):519-25  
MDMX inhibits the p300/CBP-mediated acetylation of p53.
92. PMID: 17080083; 2006, Nature;444(7115):61-6  
Inactivation of the p53 pathway in retinoblastoma.
93. PMID: 21098696; 2011, Clin Cancer Res;17(3):494-504  
Nutlin-3a is a potential therapeutic for ewing sarcoma.
94. PMID: 22820643; 2012, Nat Med;18(8):1239-47  
MDM4 is a key therapeutic target in cutaneous melanoma.
95. PMID: 12640683; 2003, Int J Cancer;104(6):752-7  
Refined mapping of 1q32 amplicons in malignant gliomas confirms MDM4 as the main amplification target.
96. PMID: 23946421; 2013, Proc Natl Acad Sci U S A;110(36):E3445-54  
Stapled  $\alpha$ -helical peptide drug development: a potent dual inhibitor of MDM2 and MDMX for p53-dependent cancer therapy.

# ACT Onco<sup>®</sup> + Report

97. PMID: 26883273; 2016, Mol Cancer Ther;15(4):574-582  
Identification of a Small Molecule That Overcomes HdmX-Mediated Suppression of p53.
98. PMID: 21075307; 2010, Cancer Cell;18(5):411-22  
A stapled p53 helix overcomes HDMX-mediated suppression of p53.
99. PMID: 20080970; 2010, J Biol Chem;285(14):10786-96  
Identification and characterization of the first small molecule inhibitor of MDMX.
100. PMID: 28351930; 2017, Clin Cancer Res;23(15):4242-4250  
Hyperprogressors after Immunotherapy: Analysis of Genomic Alterations Associated with Accelerated Growth Rate.
101. PMID: 19029958; 2008, Nat Rev Cancer;8(12):976-90  
Reflecting on 25 years with MYC.
102. PMID: 22464321; 2012, Cell;149(1):22-35  
MYC on the path to cancer.
103. PMID: 10378696; 1999, Oncogene;18(19):3004-16  
MYC oncogenes and human neoplastic disease.
104. PMID: 16934487; 2006, Semin Cancer Biol;16(4):318-30  
The Myc oncoprotein as a therapeutic target for human cancer.
105. PMID: 22113465; 2012, Clin Exp Med;12(4):217-23  
C-myc as a predictive marker for chemotherapy in metastatic breast cancer.
106. PMID: 21741827; 2011, Eur J Cancer;47(12):1779-88  
Association between c-myc amplification and pathological complete response to neoadjuvant chemotherapy in breast cancer.
107. PMID: 15132769; 2004, Cancer Sci;95(5):418-23  
Expression of the c-myc gene as a predictor of chemotherapy response and a prognostic factor in patients with ovarian cancer.
108. PMID: 22430491; 2012, J Exp Med;209(4):679-96  
MYC pathway activation in triple-negative breast cancer is synthetic lethal with CDK inhibition.
109. PMID: 27486754; 2016, Oncotarget;7(35):56864-56875  
Inhibition of cyclin dependent kinase 9 by dinaciclib suppresses cyclin B1 expression and tumor growth in triple negative breast cancer.
110. PMID: 24503701; 2014, PLoS One;9(2):e87456  
Immunohistochemistry for myc predicts survival in colorectal cancer.
111. PMID: 9816266; 1996, Clin Cancer Res;2(6):1049-53  
Overexpression of the c-myc proto-oncogene in colorectal carcinoma is associated with a reduced mortality that is abrogated by point mutation of the p53 tumor suppressor gene.
112. PMID: 23860775; 2013, Tumour Biol;34(6):3945-58  
MYC overexpression and poor prognosis in sporadic breast cancer with BRCA1 deficiency.
113. PMID: 9590181; 1998, Cell;93(3):477-86  
The hMre11/hRad50 protein complex and Nijmegen breakage syndrome: linkage of double-strand break repair to the cellular DNA damage response.
114. PMID: 9590180; 1998, Cell;93(3):467-76  
Nibrin, a novel DNA double-strand break repair protein, is mutated in Nijmegen breakage syndrome.
115. PMID: 24349281; 2013, PLoS One;8(12):e82426  
Mutation inactivation of Nijmegen breakage syndrome gene (NBS1) in hepatocellular carcinoma and intrahepatic cholangiocarcinoma.

# ACT Onco<sup>®</sup> + Report

116. PMID: 22864661; 2012, Fam Cancer;11(4):595-600  
Identification of a novel NBN truncating mutation in a family with hereditary prostate cancer.
117. PMID: 21923652; 2011, Br J Haematol;155(4):468-76  
Promising therapy results for lymphoid malignancies in children with chromosomal breakage syndromes (Ataxia teleangiectasia or Nijmegen-breakage syndrome): a retrospective survey.
118. PMID: 26083025; 2015, PLoS One;10(6):e0130393  
Prevalence of Germline Mutations in Genes Engaged in DNA Damage Repair by Homologous Recombination in Patients with Triple-Negative and Hereditary Non-Triple-Negative Breast Cancers.
119. PMID: 27908594; 2017, Lancet Oncol;18(1):75-87  
Rucaparib in relapsed, platinum-sensitive high-grade ovarian carcinoma (ARIEL2 Part 1): an international, multicentre, open-label, phase 2 trial.
120. PMID: 30353044; 2018, Br J Cancer;119(11):1401-1409  
Candidate biomarkers of PARP inhibitor sensitivity in ovarian cancer beyond the BRCA genes.
121. PMID: 28916367; 2017, Lancet;390(10106):1949-1961  
Rucaparib maintenance treatment for recurrent ovarian carcinoma after response to platinum therapy (ARIEL3): a randomised, double-blind, placebo-controlled, phase 3 trial.
122. PMID: 32086346; 2020, Clin Cancer Res;26(11):2487-2496  
Non-BRCA DNA Damage Repair Gene Alterations and Response to the PARP Inhibitor Rucaparib in Metastatic Castration-Resistant Prostate Cancer: Analysis From the Phase II TRITON2 Study.
123. PMID: 24240112; 2014, Clin Cancer Res;20(3):764-75  
Germline and somatic mutations in homologous recombination genes predict platinum response and survival in ovarian, fallopian tube, and peritoneal carcinomas.
124. PMID: 29961768; 2019, Genet Med;21(1):213-223  
Germline cancer susceptibility gene variants, somatic second hits, and survival outcomes in patients with resected pancreatic cancer.
125. PMID: 25415046; 2014, Oncotarget;5(22):11081-90  
NBN gain is predictive for adverse outcome following image-guided radiotherapy for localized prostate cancer.
126. PMID: 32226530; 2020, Theranostics;10(9):3939-3951  
Copy Number Amplification of DNA Damage Repair Pathways Potentiates Therapeutic Resistance in Cancer.
127. PMID: 24162815; 2013, Nat Med;19(11):1469-1472  
Oncogenic and drug-sensitive NTRK1 rearrangements in lung cancer.
128. PMID: 27843590; 2016, ESMO Open;1(2):e000023  
NTRK gene fusions as novel targets of cancer therapy across multiple tumour types.
129. PMID: 30624546; 2019, Ann Oncol;30(2):325-331  
Larotrectinib in adult patients with solid tumours: a multi-centre, open-label, phase I dose-escalation study.
130. PMID: 32323889; 2020, Oncologist;25(6):e881-e886  
Antitumor Activity of Larotrectinib in Esophageal Carcinoma with NTRK Gene Amplification.
131. PMID: 16141230; 2005, J Cell Sci;118(Pt 18):4261-9  
The human Rothmund-Thomson syndrome gene product, RECQL4, localizes to distinct nuclear foci that coincide with proteins involved in the maintenance of genome stability.
132. PMID: 20065033; 2010, Mol Cell Biol;30(6):1382-96  
Human RECQ1 and RECQ4 helicases play distinct roles in DNA replication initiation.
133. PMID: 20222902; 2010, Aging Cell;9(3):358-71  
The involvement of human RECQL4 in DNA double-strand break repair.

# ACT Onco<sup>®</sup> + Report

134. PMID: 19567405; 2009, Hum Mol Genet;18(18):3470-83  
Direct and indirect roles of RECQL4 in modulating base excision repair capacity.
135. PMID: 27013200; 2016, Cancer Res;76(10):3057-66  
Human Helicase RECQL4 Drives Cisplatin Resistance in Gastric Cancer by Activating an AKT-YB1-MDR1 Signaling Pathway.
136. PMID: 24072219; 2013, Oncologist;18(10):1063-73  
DNA repair gene patterns as prognostic and predictive factors in molecular breast cancer subtypes.
137. PMID: 31495919; 2020, Int J Cancer;146(11):3098-3113  
Human RECQL4 represses the RAD52-mediated single-strand annealing pathway after ionizing radiation or cisplatin treatment.
138. PMID: 26690729; 2016, J Pathol;238(4):495-501  
RECQL4 helicase has oncogenic potential in sporadic breast cancers.
139. PMID: 21332640; 2011, J Cell Mol Med;15(7):1433-42  
Targeting telomerase-expressing cancer cells.
140. PMID: 19571879; 2009, Nature;460(7251):66-72  
Telomerase modulates Wnt signalling by association with target gene chromatin.
141. PMID: 23159929; 2012, Nat Cell Biol;14(12):1270-81  
Telomerase directly regulates NF- $\kappa$ B-dependent transcription.
142. PMID: 19701182; 2009, Nature;461(7261):230-5  
An RNA-dependent RNA polymerase formed by TERT and the RMRP RNA.
143. PMID: 23348506; 2013, Science;339(6122):957-9  
Highly recurrent TERT promoter mutations in human melanoma.
144. PMID: 23530248; 2013, Proc Natl Acad Sci U S A;110(15):6021-6  
TERT promoter mutations occur frequently in gliomas and a subset of tumors derived from cells with low rates of self-renewal.
145. PMID: 11103775; 2000, Cancer Res;60(22):6230-5  
Frequent amplification of the telomerase reverse transcriptase gene in human tumors.
146. PMID: 12007187; 2002, Genes Chromosomes Cancer;34(3):269-75  
Amplification of the telomerase reverse transcriptase (hTERT) gene in cervical carcinomas.
147. PMID: 25301727; 2014, Oncotarget;5(20):10048-57  
TERT promoter mutations and gene amplification: promoting TERT expression in Merkel cell carcinoma.
148. PMID: 16641908; 2006, Br J Cancer;94(10):1452-9  
Amplification of telomerase (hTERT) gene is a poor prognostic marker in non-small-cell lung cancer.
149. PMID: 27982019; 2017, Cancer Gene Ther;24(1):20-27  
The associations of TERT-CLPTM1L variants and TERT mRNA expression with the prognosis of early stage non-small cell lung cancer.
150. PMID: 29100407; 2017, Oncotarget;8(44):77540-77551  
TERT promoter status and gene copy number gains: effect on TERT expression and association with prognosis in breast cancer.
151. PMID: 20525995; 2010, N Engl J Med;362(24):2260-70  
Dasatinib versus imatinib in newly diagnosed chronic-phase chronic myeloid leukemia.
152. PMID: 18541900; 2008, J Clin Oncol;26(19):3204-12  
Intermittent target inhibition with dasatinib 100 mg once daily preserves efficacy and improves tolerability in imatinib-resistant and -intolerant chronic-phase chronic myeloid leukemia.
153. PMID: 17496201; 2007, Blood;110(7):2309-15



# ACT Onco<sup>®</sup> + Report

Dasatinib induces rapid hematologic and cytogenetic responses in adult patients with Philadelphia chromosome positive acute lymphoblastic leukemia with resistance or intolerance to imatinib: interim results of a phase 2 study.

154. PMID: 26703889; 2016, Lancet;387(10022):968-977  
Everolimus for the treatment of advanced, non-functional neuroendocrine tumours of the lung or gastrointestinal tract (RADIANT-4): a randomised, placebo-controlled, phase 3 study.
155. PMID: 22149876; 2012, N Engl J Med;366(6):520-9  
Everolimus in postmenopausal hormone-receptor-positive advanced breast cancer.
156. PMID: 21306238; 2011, N Engl J Med;364(6):514-23  
Everolimus for advanced pancreatic neuroendocrine tumors.
157. PMID: 23158522; 2013, Lancet;381(9861):125-32  
Efficacy and safety of everolimus for subependymal giant cell astrocytomas associated with tuberous sclerosis complex (EXIST-1): a multicentre, randomised, placebo-controlled phase 3 trial.
158. PMID: 18653228; 2008, Lancet;372(9637):449-56  
Efficacy of everolimus in advanced renal cell carcinoma: a double-blind, randomised, placebo-controlled phase III trial.
159. PMID: 32343890; 2020, N Engl J Med;382(22):2091-2102  
Olaparib for Metastatic Castration-Resistant Prostate Cancer.
160. PMID: 31851799; 2019, N Engl J Med;381(25):2416-2428  
Olaparib plus Bevacizumab as First-Line Maintenance in Ovarian Cancer.
161. PMID: 31157963; 2019, N Engl J Med;381(4):317-327  
Maintenance Olaparib for Germline BRCA-Mutated Metastatic Pancreatic Cancer.
162. PMID: 30345884; 2018, N Engl J Med;379(26):2495-2505  
Maintenance Olaparib in Patients with Newly Diagnosed Advanced Ovarian Cancer.
163. PMID: 28578601; 2017, N Engl J Med;377(6):523-533  
Olaparib for Metastatic Breast Cancer in Patients with a Germline BRCA Mutation.
164. PMID: 28754483; 2017, Lancet Oncol;18(9):1274-1284  
Olaparib tablets as maintenance therapy in patients with platinum-sensitive, relapsed ovarian cancer and a BRCA1/2 mutation (SOLO2/ENGOT-Ov21): a double-blind, randomised, placebo-controlled, phase 3 trial.
165. PMID: 27617661; 2016, Lancet Oncol;17(11):1579-1589  
Overall survival in patients with platinum-sensitive recurrent serous ovarian cancer receiving olaparib maintenance monotherapy: an updated analysis from a randomised, placebo-controlled, double-blind, phase 2 trial.
166. PMID: 27932229; 2017, Lancet;389(10064):56-66  
Regorafenib for patients with hepatocellular carcinoma who progressed on sorafenib treatment (RESORCE): a randomised, double-blind, placebo-controlled, phase 3 trial.
167. PMID: 23177515; 2013, Lancet;381(9863):295-302  
Efficacy and safety of regorafenib for advanced gastrointestinal stromal tumours after failure of imatinib and sunitinib (GRID): an international, multicentre, randomised, placebo-controlled, phase 3 trial.
168. PMID: 23177514; 2013, Lancet;381(9863):303-12  
Regorafenib monotherapy for previously treated metastatic colorectal cancer (CORRECT): an international, multicentre, randomised, placebo-controlled, phase 3 trial.
169. PMID: 24768112; 2014, Lancet;384(9940):319-28  
Sorafenib in radioactive iodine-refractory, locally advanced or metastatic differentiated thyroid cancer: a randomised, double-blind, phase 3 trial.
170. PMID: 18650514; 2008, N Engl J Med;359(4):378-90

# ACT Onco<sup>®</sup> + Report

Sorafenib in advanced hepatocellular carcinoma.

171. PMID: 17189398; 2006, Clin Cancer Res;12(24):7271-8  
Sorafenib for the treatment of advanced renal cell carcinoma.
172. PMID: 30110579; 2018, N Engl J Med;379(8):753-763  
Talazoparib in Patients with Advanced Breast Cancer and a Germline BRCA Mutation.
173. PMID: 17538086; 2007, N Engl J Med;356(22):2271-81  
Temsilolimus, interferon alfa, or both for advanced renal-cell carcinoma.