



Sample Information

Patient Name: 林陳淑媛**Gender:** Female**ID No.:** P201176311**History No.:** 45445963**Age:** 68**Ordering Doctor:** DOC3153J 黃煦晴**Ordering REQ.:** D54397K**Signing in Date:** 2020/05/21**Path No.:** S109-99489**MP No.:** F2023**Assay:** Oncomine Focus Assay**Sample Type:** FFPE**Block No.:** S109-76164A**Percentage of tumor cells:** 50%**Note:**

Sample Cancer Type: Non-Small Cell Lung Cancer

Table of Contents

	Page
Variant Details	2
Biomarker Descriptions	3
Relevant Therapy Summary	4
Relevant Therapy Details	5

Report Highlights

2 Relevant Biomarkers
2 Therapies Available
21 Clinical Trials

Relevant Non-Small Cell Lung Cancer Findings

Gene	Finding	Gene	Finding
ALK	Not detected	NTRK1	Not detected
BRAF	Not detected	NTRK2	Not detected
EGFR	Not detected	NTRK3	Not detected
ERBB2	Not detected	RET	KIF5B-RET fusion
KRAS	Not detected	ROS1	Not detected
MET	Not detected		



Relevant Biomarkers

■ Indicated ■ Contraindicated

Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
KIF5B-RET fusion kinesin family member 5B - ret proto-oncogene Tier: IA	■ cabozantinib ■ vandetanib	None	18
MYC amplification MYC proto-oncogene, bHLH transcription factor Tier: IIC	None	None	3

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

Tier Reference: Li et al. Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists. J Mol Diagn. 2017 Jan;19(1):4-23.

Variant Details

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
JAK1	p.(=)	c.2199A>G	.	chr1:65310489	49.47%	NM_002227.3	synonymous	1993
ALK	p.(D1529E)	c.4587C>G	.	chr2:29416366	48.45%	NM_004304.4	missense	2000
ALK	p.(I1461V)	c.4381A>G	.	chr2:29416572	99.95%	NM_004304.4	missense	1999
ALK	p.(=)	c.3600G>C	.	chr2:29443617	50.28%	NM_004304.4	synonymous	1997
ALK	p.(=)	c.3375C>A	.	chr2:29445458	50.48%	NM_004304.4	synonymous	1995
FGFR3	p.(=)	c.1953G>A	.	chr4:1807894	99.78%	NM_000142.4	synonymous	904
PDGFRA	p.(=)	c.939T>G	.	chr4:55133726	47.94%	NM_006206.5	synonymous	1994
PDGFRA	p.(=)	c.1701A>G	.	chr4:55141055	99.90%	NM_006206.5	synonymous	1996
KIT	p.(=)	c.1638A>G	.	chr4:55593481	99.75%	NM_000222.2	synonymous	1994
FGFR4	p.(P136L)	c.407C>T	.	chr5:176517797	99.30%	NM_213647.2	missense	2000
FGFR4	p.(=)	c.483A>G	.	chr5:176517985	5.92%	NM_213647.2	synonymous	1741
RET	p.(=)	c.2307G>T	.	chr10:43613843	50.45%	NM_020975.4	synonymous	1996

Gene Fusions (RNA)

Genes	Variant ID	Locus
KIF5B-RET	KIF5B-RET.K15R12.COSF1232	chr10:32317356 - chr10:43612032

Copy Number Variations

Gene	Locus	Copy Number
MYC	chr8:128748885	8.1



Biomarker Descriptions

MYC (MYC proto-oncogene, bHLH transcription factor)

Background: The MYC gene encodes the MYC proto-oncogene (c-MYC), a basic helix-loop-helix transcription factor that regulates the expression of numerous genes that control cell cycle progression, apoptosis, metabolic pathways, and cellular transformation^{1,2,3,4}. MYC is part of the MYC oncogene family that includes related transcription factors MYCN and MYCL that regulate transcription in 10-15% of promoter regions⁵. MYC functions as a heterodimer in complex with the transcription factor MAX^{2,6}.

Alterations and prevalence: Recurrent somatic alterations are observed in both solid and hematological cancers. Recurrent somatic mutations in MYC, including codon T58, are infrequent and hypothesized to increase the stability of the MYC protein^{7,8}. MYC gene amplification is particularly common in diverse solid tumors. MYC amplification is observed in 30% of serous ovarian cancer, 20% of uterine serous carcinoma, 15% of esophageal and breast cancers, and is common (1-10%) in numerous other cancer types^{9,10,11}. MYC is the target of the t(8;14)(q24;32) chromosomal translocation in Burkitt's lymphoma that places MYC coding sequences adjacent to immunoglobulin region regulatory sequences, which results in increased MYC expression^{12,13}.

Potential relevance: Currently, no therapies are approved for MYC aberrations. Due to the high frequency of somatic MYC alterations in cancer, many approaches are being investigated in clinical trials including strategies to disrupt complex formation with MAX, including inhibition of MYC expression and synthetic lethality associated with MYC overexpression^{1,14,15,16}.

RET (ret proto-oncogene)

Background: The RET gene encodes the RET receptor tyrosine kinase which is activated by a ligand family of glial cell line-derived neurotrophic factors (GDNF)¹⁷. RET is the target of recurrent chromosomal rearrangements that generate fusion proteins containing the intact RET tyrosine kinase domain combined with several fusion partner genes. RET fusion kinases are constitutively activated and drive oncogenic transformation which can lead to activation of PI3K/AKT, RAS/RAF/MEK/ERK, and PLCγ/PKC pathways resulting in cell survival and proliferation¹⁸.

Alterations and prevalence: RET fusions occur in approximately 55% of papillary thyroid carcinomas (PTC) with even higher frequencies observed in PTC patients with radiation exposure^{19,20,21}. RET rearrangement is also present in 1-2% of non-small cell lung cancer (NSCLC)²². Point mutations in RET are relatively common in sporadic medullary thyroid cancer (MTC), with 6% of patients found to contain germline mutations²³. Somatic mutations (specifically at codon 918), which leads to increased kinase activity, have been observed in at least 25% of MTC cases²³.

Potential relevance: Currently, no therapies are approved for RET aberrations. However, the RET inhibitor, pralsetinib^{24,25}, was granted breakthrough therapy designation for RET mutation-positive medullary thyroid cancer (2019) as well as for RET fusion-positive NSCLC (2020). The FDA approved small-molecule tyrosine kinase inhibitors, vandetanib (2011) and cabozantinib (2012), are recommended for treatment of NSCLC patients with RET rearrangements²⁶. Cabozantinib has also demonstrated clinical benefit in RET mutated medullary thyroid cancer patients²⁷. Point mutations involving codons 804 and 806 have been shown to confer resistance to selective kinase inhibitors including vandetanib^{28,29}. RET mutations at codon 918 are associated with high risk and adverse prognosis in patients diagnosed with MTC³⁰.



Relevant Therapy Summary

● In this cancer type ○ In other cancer type ● In this cancer type and other cancer types ⛔ Contraindicated ⚠ Both for use and contraindicated ✕ No evidence

KIF5B-RET fusion

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
cabozantinib	✕	●	✕	✕	● (II)
vandetanib	✕	●	✕	✕	✕
alectinib	✕	✕	✕	✕	● (IV)
alectinib, crizotinib	✕	✕	✕	✕	● (III)
ipilimumab, nivolumab, radiation therapy, surgical intervention	✕	✕	✕	✕	● (III)
erdafitinib	✕	✕	✕	✕	● (II)
ponatinib	✕	✕	✕	✕	● (II)
sunitinib	✕	✕	✕	✕	● (II)
targeted therapy, chemotherapy	✕	✕	✕	✕	● (II)
pralsetinib	✕	✕	✕	✕	● (I/II)
selpercatinib	✕	✕	✕	✕	● (I/II)
TPX-0046	✕	✕	✕	✕	● (I/II)
BOS172738	✕	✕	✕	✕	● (I)

MYC amplification

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
VX-970	✕	✕	✕	✕	● (II)
entinostat, nivolumab	✕	✕	✕	✕	● (I/II)
BMS-986158	✕	✕	✕	✕	● (I)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.



Relevant Therapy Details

Current NCCN Information

- ☒ In this cancer type
 ☐ In other cancer type
 ☐ In this cancer type and other cancer types
 ☒ Contraindicated
 ☐ Not recommended
 ☐ Resistance

NCCN information is current as of 2019-11-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

KIF5B-RET fusion

cabozantinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: RET fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Non-Small Cell Lung Cancer; Emerging targeted agents

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 2.2020]

vandetanib

Cancer type: Non-Small Cell Lung Cancer

Variant class: RET fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Non-Small Cell Lung Cancer; Emerging targeted agents

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 2.2020]

Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:



References

- Chen et al. Targeting oncogenic Myc as a strategy for cancer treatment. *Signal Transduct Target Ther*. 2018 Feb 23;3:5. doi: 10.1038/s41392-018-0008-7. eCollection 2018. PMID: 29527331
- Dang. MYC on the path to cancer. *Cell*. 2012 Mar 30;149(1):22-35. PMID: 22464321
- Dominguez-Sola et al. Non-transcriptional control of DNA replication by c-Myc. *Nature*. 2007 Jul 26;448(7152):445-51. PMID: 17597761
- Wahlström et al. Impact of MYC in regulation of tumor cell metabolism. *Biochim. Biophys. Acta*. 2015 May;1849(5):563-9. PMID: 25038584
- Dang et al. The c-Myc target gene network. *Semin. Cancer Biol*. 2006 Aug;16(4):253-64. PMID: 16904903
- Blackwood et al. Myc and Max function as a nucleoprotein complex. *Curr. Opin. Genet. Dev*. 1992 Apr;2(2):227-35. PMID: 1638116
- Chakraborty et al. A common functional consequence of tumor-derived mutations within c-MYC. *Oncogene*. 2015 Apr 30;34(18):2406-9. PMID: 24998853
- Xu-Monette et al. Clinical and Biologic Significance of MYC Genetic Mutations in De Novo Diffuse Large B-cell Lymphoma. *Clin. Cancer Res*. 2016 Jul 15;22(14):3593-605. PMID: 26927665
- Kalkat et al. MYC Deregulation in Primary Human Cancers. *Genes (Basel)*. 2017 May 25;8(6). PMID: 28587062
- Beroukhi et al. The landscape of somatic copy-number alteration across human cancers. *Nature*. 2010 Feb 18;463(7283):899-905. doi: 10.1038/nature08822. PMID: 20164920
- Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov*. 2012 May;2(5):401-4. PMID: 22588877
- Taub et al. Translocation of the c-myc gene into the immunoglobulin heavy chain locus in human Burkitt lymphoma and murine plasmacytoma cells. *Proc Natl Acad Sci U S A*. 1982 Dec;79(24):7837-41. PMID: 6818551
- Ott et al. Understanding MYC-driven aggressive B-cell lymphomas: pathogenesis and classification. *Hematology Am Soc Hematol Educ Program*. 2013;2013:575-83. PMID: 24319234
- Posternak et al. Strategically targeting MYC in cancer. *F1000Res*. 2016;5. PMID: 27081479
- Carabet et al. Therapeutic Inhibition of Myc in Cancer. *Structural Bases and Computer-Aided Drug Discovery Approaches*. *Int J Mol Sci*. 2018 Dec 29;20(1). PMID: 30597997
- Shahbazi et al. The Bromodomain Inhibitor JQ1 and the Histone Deacetylase Inhibitor Panobinostat Synergistically Reduce N-Myc Expression and Induce Anticancer Effects. *Clin. Cancer Res*. 2016 May 15;22(10):2534-44. PMID: 26733615
- Knowles et al. Structure and chemical inhibition of the RET tyrosine kinase domain. *J. Biol. Chem*. 2006 Nov 3;281(44):33577-87. PMID: 16928683
- Ibáñez. Structure and physiology of the RET receptor tyrosine kinase. *Cold Spring Harb Perspect Biol*. 2013 Feb 1;5(2). PMID: 23378586
- Santoro et al. Central role of RET in thyroid cancer. *Cold Spring Harb Perspect Biol*. 2013 Dec 1;5(12):a009233. PMID: 24296167
- Elisei et al. RET/PTC rearrangements in thyroid nodules: studies in irradiated and not irradiated, malignant and benign thyroid lesions in children and adults. *J. Clin. Endocrinol. Metab*. 2001 Jul;86(7):3211-6. PMID: 11443191
- Ciampi et al. RET/PTC rearrangements and BRAF mutations in thyroid tumorigenesis. *Endocrinology*. 2007 Mar;148(3):936-41. PMID: 16946010
- Kohn et al. KIF5B-RET fusions in lung adenocarcinoma. *Nat. Med*. 2012 Feb 12;18(3):375-7. PMID: 22327624
- Wohllk et al. Relevance of RET proto-oncogene mutations in sporadic medullary thyroid carcinoma. *J. Clin. Endocrinol. Metab*. 1996 Oct;81(10):3740-5. PMID: 8855832
- <http://ir.blueprintmedicines.com/news-releases/news-release-details/blueprint-medicines-reports-first-quarter-2019-financial-results>
- Blueprint Medicines Announces Top-line Data for Pralsetinib and Initiates Rolling NDA Submission to FDA for the Treatment of Patients with RET Fusion-Positive Non-Small Cell Lung Cancer. <http://ir.blueprintmedicines.com/news-releases/news-release-details/blueprint-medicines-announces-top-line-data-pralsetinib-and>
- NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 2.2020]



References (continued)

27. Sherman et al. Correlative analyses of RET and RAS mutations in a phase 3 trial of cabozantinib in patients with progressive, metastatic medullary thyroid cancer. *Cancer*. 2016 Dec 15;122(24):3856-3864. PMID: 27525386
28. Carlomagno et al. Disease associated mutations at valine 804 in the RET receptor tyrosine kinase confer resistance to selective kinase inhibitors. *Oncogene*. 2004 Aug 12;23(36):6056-63. PMID: 15184865
29. Carlomagno et al. Identification of tyrosine 806 as a molecular determinant of RET kinase sensitivity to ZD6474. *Endocr Relat Cancer*. 2009 Mar;16(1):233-41. doi: 10.1677/ERC-08-0213. Epub 2008 Nov 24. PMID: 19029224
30. NCCN Guidelines® - NCCN-Thyroid Carcinoma [Version 2.2019]