



Sample Information

Patient Name: 褚侯霖
Gender: Male
ID No.: F120070737
History No.: 19697337
Age: 58

Ordering Doctor: DOC5388L 洪雅文
Ordering REQ.: 0BTYXPQ
Signing in Date: 2022/04/15

Path No.: S111-99032
MP No.: TM22007
Assay: Tumor Mutation Load Assay
Sample Type: FFPE
Block No.: S111-70464A
Percentage of tumor cells: 80%

Reporting Doctor: DOC5466K 葉奕成 (Phone: 8#5466)

Note:

Sample Cancer Type: Liver Cancer

Table of Contents	Page
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2
Biomarker Descriptions	3
Relevant Therapy Summary	5
Relevant Therapy Details	6
Clinical Trials Summary	16
Alert Details	16

Report Highlights
2 Relevant Biomarkers
2 Therapies Available
2 Clinical Trials

Relevant Liver Cancer Variants

Gene	Finding
NTRK1	None detected
NTRK3	None detected

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	Tumor Mutational Burden 5.92 Mut/Mb measured Prognostic significance: None Diagnostic significance: None	pembrolizumab ¹	pembrolizumab	2
IIC	ATM c.7789-2A>G ATM serine/threonine kinase Allele Frequency: 24.63% Prognostic significance: None Diagnostic significance: None	None	olaparib ¹	0

Public data sources included in relevant therapies: FDA¹, NCCN, EMA², ESMO

Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

Tier Reference: Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

Prevalent cancer biomarkers without relevant evidence based on included data sources

*TSC2 p.(M1715Ifs*106) c.5145_5160delGGCCCTGCACGCAAT*

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

DNA Sequence Variants								
Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
ATM	p.(?)	c.7789-2A>G	.	chr11:108203487	24.63%	NM_000051.3	unknown	1080
TSC2	p.(M1715Ifs*106)	c.5145_5160delGGC CCTGCACGCAAT	.	chr16:2138123	34.19%	NM_000548.5	frameshift Deletion	1094
PARP1	p.(Q694=)	c.2082G>A	.	chr1:226558207	49.06%	NM_001618.4	synonymous	1013
AKT3	p.(Q78*)	c.232C>T	.	chr1:243828126	26.91%	NM_005465.7	nonsense	1059
BCL11A	p.(S370F)	c.1109C>T	.	chr2:60688938	3.77%	NM_022893.4	missense	106
LRP1B	p.(S2410C)	c.7228A>T	.	chr2:141299507	53.07%	NM_018557.3	missense	797
NFE2L2	p.(Q527*)	c.1579C>T	.	chr2:178095752	3.77%	NM_006164.5	nonsense	53
STK36	p.(K20=)	c.60G>A	.	chr2:219537612	50.50%	NM_015690.5	synonymous	2000
MYD88	p.(S137C)	c.409A>T	.	chr3:38181435	26.21%	NM_001172567.2	missense	1984
LTF	p.(R23dup)	c.68_69insAAG	.	chr3:46501284	99.85%	NM_002343.6	nonframeshift Insertion	1976
BAP1	p.(N446=)	c.1338C>T	.	chr3:52437823	48.65%	NM_004656.4	synonymous	1998
FGFR3	p.(L164V)	c.490C>G	.	chr4:1803138	51.68%	NM_000142.4	missense	1455
PDGFRA	p.(P567=)	c.1701A>G	.	chr4:55141055	100.00%	NM_006206.6	synonymous	1986
ADGRL3	p.(T443I)	c.1328C>T	.	chr4:62758425	50.62%	NM_015236.6	missense	162
NOTCH4	p.([P271=;D272G])	c.813_815delAGAAins GGG	.	chr6:32188640	47.22%	NM_004557.4	synonymous, missense	1995
NOTCH4	p.(L13_L16del)	c.36_47delGCTGCTG CTGCT	.	chr6:32191658	73.70%	NM_004557.4	nonframeshift Deletion	

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2022.03(006).

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency) (continued)

DNA Sequence Variants (continued)

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
HSP90AB1	p.(D691G)	c.2072A>G	.	chr6:44221232	51.00%	NM_001271970.1	missense	1998
TRIM24	p.(H407=)	c.1221C>T	.	chr7:138235885	48.20%	NM_015905.3	synonymous	2000
KMT2C	p.(D219G)	c.656A>G	.	chr7:152008966	4.00%	NM_170606.3	missense	50
RECQL4	p.(C857_T858del)	c.2569_2574delTGC ACC	.	chr8:145738410	60.85%	NM_004260.4	nonframeshift Deletion	1990
TAF1L	p.(P1443S)	c.4327C>T	.	chr9:32631251	4.48%	NM_153809.2	missense	67
NFKB2	p.(P423=)	c.1269A>G	.	chr10:104159196	100.00%	NM_001077494.3	synonymous	309
NUP98	p.(I736=)	c.2208T>C	.	chr11:3741994	42.22%	NM_016320.5	synonymous	1582
EXT2	p.(?)	c.-36C>T	.	chr11:44117767	66.67%	NM_000401.3	unknown	186
NUMA1	p.(R1885=)	c.5655T>C	.	chr11:71717118	48.29%	NM_006185.4	synonymous	1847
NUMA1	p.(G1514S)	c.4540G>A	.	chr11:71724009	50.45%	NM_006185.4	missense	2000
ATM	p.(E2039V)	c.6116A>T	.	chr11:108186758	28.01%	NM_000051.3	missense	1171
ZNF384	p.(Q501Hfs*48)	c.1503delG	.	chr12:6777110	100.00%	NM_001135734.2	frameshift Deletion	604
ADAMTS20p.	(K655T)	c.1964A>C	.	chr12:43846192	30.47%	NM_025003.5	missense	279
RB1	p.(Y728=)	c.2184C>T	.	chr13:49037944	3.57%	NM_000321.2	synonymous	56
NIN	p.(K884R)	c.2651A>G	.	chr14:51225097	71.97%	NM_020921.3	missense	1998
DICER1	p.(?)	c.4204_4206+19delin . sTG TAGTTT	.	chr14:95566098	61.89%	NM_030621.4	unknown	391
DICER1	p.(?)	c.4206+9G>T	.	chr14:95566108	35.81%	NM_030621.4	unknown	391
NTRK3	p.(V21F)	c.61G>T	.	chr15:88799324	48.90%	NM_001012338.2	missense	1998
IGF1R	p.(G1328S)	c.3982G>A	.	chr15:99500549	48.12%	NM_000875.5	missense	904
ITGB3	p.(E775A)	c.2324A>C	.	chr17:45387527	29.49%	NM_000212.3	missense	1933
ZNF521	p.(A182=)	c.546G>A	.	chr18:22807336	32.27%	NM_015461.3	synonymous	1999
ERCC2	p.(G413D)	c.1238G>A	.	chr19:45860957	5.95%	NM_000400.4	missense	84

Biomarker Descriptions

ATM (ATM serine/threonine kinase)

Background: The ATM gene encodes a serine/threonine kinase that belongs to the phosphatidylinositol-3-kinase related kinases (PIKKs) family of genes that also includes ATR and PRKDC (also known as DNA-PKc)¹. ATM and ATR act as master regulators of DNA damage response. Specifically, ATM is involved in double-stranded break (DSB) repair while ATR is involved in single-stranded DNA (ssDNA) repair². ATM is recruited to the DNA damage site by the MRE11/RAD50/NBN (MRN) complex that senses DSB^{2,3}. Upon activation, ATM phosphorylates several downstream proteins such as the NBN, MDC1, BRCA1, CHK2 and TP53BP1 proteins⁴. ATM is a tumor suppressor gene and loss of function mutations in ATM are implicated in the BRCAness phenotype, which is characterized by a defect in homologous recombination repair (HRR), mimicking BRCA1 or BRCA2 loss^{5,6}. Germline mutations in ATM often result in Ataxia-telangiectasia, a hereditary disease also referred to as DNA damage response syndrome that is characterized by chromosomal instability⁷.

Biomarker Descriptions (continued)

Alterations and prevalence: Recurrent somatic mutations in ATM are observed in 17% of endometrial carcinoma, 15% of undifferentiated stomach adenocarcinoma, 13% of bladder urothelial carcinoma, 12% of colorectal adenocarcinoma, 9% of melanoma as well as esophagogastric adenocarcinoma and 8% of non-small cell lung cancer^{8,9}.

Potential relevance: The PARP inhibitor, olaparib¹⁰ is approved (2020) for metastatic castration-resistant prostate cancer (mCRPC) with deleterious or suspected deleterious, germline or somatic mutations in HRR genes that includes ATM. Consistent with other genes associated with the BRCAness phenotype, ATM mutations may aid in selecting patients likely to respond to PARP inhibitors^{5,11,12}. Specifically, in a phase II trial of metastatic, castration-resistant prostate cancer, four of six patients with germline or somatic ATM mutations demonstrated clinical responses to olaparib¹³. In 2022, the FDA granted fast track designation to the small molecule inhibitor, pidnarulex¹⁴, for BRCA1/2, PALB2, or other homologous recombination deficiency (HRD) mutations in breast and ovarian cancers.

TSC2 (TSC complex subunit 2)

Background: The TSC2 gene encodes the tuberlin protein. TSC2 and TSC1 (also known as hamartin) form a complex through their respective coiled-coil domains¹⁵. The TSC1-TSC2 complex is a negative regulator of the mTOR signaling pathway that regulates cell growth, cell proliferation, and protein and lipid synthesis¹⁶. Specifically, the TSC1-TSC2 complex acts as a GTPase activating (GAP) protein that inhibits the G-protein RHEB and keeps it in an inactivated state (RHEB-GDP). GTP bound RHEB (RHEB-GTP) is required to activate the mTOR complex 1 (mTORC1). TSC1 and TSC2 are tumor suppressor genes. Loss of function mutations in TSC1 and TSC2 lead to dysregulation of the mTOR pathway^{15,17}. Inactivating germline mutations in TSC1 and TSC2 are associated with tuberous sclerosis complex (TSC), an autosomal dominant neurocutaneous and progressive disorder that presents with multiple benign tumors in different organs¹⁵.

Alterations and prevalence: Somatic mutations are observed in up to 8% of skin cutaneous melanoma, 7% of uterine corpus endometrial carcinoma, and 4% of cervical squamous cell carcinoma^{8,9}.

Potential relevance: Currently, no therapies are approved for TSC2 aberrations.

Tumor Mutational Burden

Background: Tumor mutational burden (TMB), also known as tumor mutational load (TML), is the count of somatic mutations in the DNA of cancer cells. TMB is determined by next-generation sequencing and is expressed as the number of mutations per megabase (mut/Mb) of DNA coding sequence¹⁸. Errors in DNA repair, including mutations in the POLE gene and in mismatch repair (MMR) genes, are associated with increased TMB^{19,20,21,22,23}. High TMB is associated with increased neo-antigen burden and has been linked to response to immune checkpoint inhibitors (ICIs) that target the cytotoxic T lymphocyte antigen-4 (CTLA4), programmed death protein 1 (PD1), and programmed death-ligand 1 (PD-L1) inhibitors^{24,25,26,27}.

Alterations and prevalence: In one study of over 100,000 tumor samples, the median TMB value was 3.6 mut/Mb although TMB values vary widely across cancers²⁸. Certain childhood cancers, leukemia, glioblastoma, and neuroblastoma typically have low mutation burden and median TMB values <1 mut/Mb^{25,28}. In comparison, cancers that experience genotoxic insults including skin cancer and lung cancer have higher median TMB values of approximately 10 mut/Mb^{25,28}. For example, within non-small cell lung cancer (NSCLC), higher TMB was observed in former/current smokers (10.5 mut/Mb) relative to never smokers (0.6 mut/Mb)^{25,28,29}. There is no consensus around the definition of high and low TMB that could be applied universally to all tumor types, instead multiple sources suggest that TMB status is a cancer type specific attribute^{28,30,31}. In NSCLC, several studies have suggested establishing a threshold between low and high TMB of 10 +/- 1 mut/Mb^{32,33,34,35}.

Potential relevance: ICIs stimulate a patient's own T-cells to kill tumors and have exhibited benefits in some patients. The first ICI to be approved by the FDA was ipilimumab (2011), an anti-CTLA4 antibody indicated for the treatment of metastatic melanoma. In 2014, anti-PD-1 antibodies, nivolumab (2014) and pembrolizumab (2014), were subsequently approved for the treatment of metastatic melanoma. Pembrolizumab was also approved (2014) for advanced esophageal squamous cell carcinoma. In 2020, the indication for pembrolizumab³⁶ was expanded to include TMB-H (>= 10 mut/Mb) solid tumors that have progressed on prior therapy. Indications have been expanded for these ICIs to include several other cancer types including NSCLC, advanced renal cell carcinoma, classical Hodgkin lymphoma, recurrent or metastatic squamous cell carcinoma of the head and neck, urothelial carcinoma, microsatellite instability (MSI)-High or mismatch repair deficient (dMMR) colorectal cancer, and hepatocellular carcinoma. Atezolizumab (2016), avelumab (2017), and durvalumab (2017), that target programmed death-ligand 1 (PD-L1), were subsequently approved by the FDA. However, the predictive biomarkers that underlie the clinical benefits of these approved immunotherapies, including TMB, are under active investigation. Several published studies including the CheckMate 586 and CheckMate 817 clinical trials have concluded that high TMB was associated with improved response to FDA approved checkpoint inhibitors^{33,37,38}. In contrast, several promising previous trials failed to show an improvement in survival outcomes between high and low TMB including CheckMate 227 (ipilimumab + nivolumab vs. chemotherapy), CheckMate 026 (nivolumab vs. chemotherapy), KEYNOTE 189 (pembrolizumab vs. chemotherapy), KEYNOTE 021 (pembrolizumab vs. pembrolizumab + chemotherapy), and Lung-MAP (nivolumab + ipilimumab vs. nivolumab). In

Biomarker Descriptions (continued)

response, suggestions to combine TMB score with PD-L1 expression as a way to increase the predictive power for patient stratification have been reported³⁹. Nivolumab alone or in combination with ipilimumab is recommended for use in NSCLC with evidence of high TMB⁴⁰. Pembrolizumab is indicated for use in various cancer types with evidence of metastasis including Ewing sarcoma, salivary gland neoplasms, cervical cancer, uterine sarcoma, endometrial carcinoma, thyroid cancer, ovarian cancer, esophageal cancer, esophagogastric junction cancer, breast cancer, and germ cell tumors with high TMB^{41,42,43,44,45,46,47,48,49}. TMB score estimation is affected by the utilized assays, therefore efforts are underway to develop a standardized approach for score calculation with the aim to support consistent reporting of TMB values across laboratories^{50,51,52,53}.

Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

Tumor Mutational Burden

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
pembrolizumab	<input checked="" type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/> (II)
atezolizumab	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/> (II)

ATM c.7789-2A>G

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
olaparib	<input type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Details

Current FDA Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

FDA information is current as of 2022-02-16. For the most up-to-date information, search www.fda.gov.

Tumor Mutational Burden

● pembrolizumab

Cancer type: Solid Tumor

Label as of: 2022-02-04

Variant class: Tumor Mutational Burden

Indications and usage:

KEYTRUDA® is a programmed death receptor-1 (PD-1)-blocking antibody indicated:

Melanoma

- for the treatment of patients with unresectable or metastatic melanoma.
- for the adjuvant treatment of adult and pediatric (12 years and older) patients with Stage IIB, IIC, or III melanoma following complete resection.

Non-Small Cell Lung Cancer (NSCLC)

- in combination with pemetrexed and platinum chemotherapy, as first-line treatment of patients with metastatic nonsquamous NSCLC, with no EGFR or ALK genomic tumor aberrations.
- in combination with carboplatin and either paclitaxel or paclitaxel protein-bound, as first-line treatment of patients with metastatic squamous NSCLC.
- as a single agent for the first-line treatment of patients with NSCLC expressing PD-L1 [Tumor Proportion Score (TPS) $\geq 1\%$] as determined by an FDA-approved test, with no EGFR or ALK genomic tumor aberrations, and is:
 - stage III where patients are not candidates for surgical resection or definitive chemoradiation, or
 - metastatic.
- as a single agent for the treatment of patients with metastatic NSCLC whose tumors express PD-L1 (TPS $\geq 1\%$) as determined by an FDA-approved test, with disease progression on or after platinum-containing chemotherapy. Patients with EGFR or ALK genomic tumor aberrations should have disease progression on FDA-approved therapy for these aberrations prior to receiving KEYTRUDA®.

Head and Neck Squamous Cell Cancer (HNSCC)

- in combination with platinum and FU for the first-line treatment of patients with metastatic or with unresectable, recurrent HNSCC.
- as a single agent for the first-line treatment of patients with metastatic or with unresectable, recurrent HNSCC whose tumors express PD-L1 [Combined Positive Score (CPS) ≥ 1] as determined by an FDA-approved test.
- as a single agent for the treatment of patients with recurrent or metastatic HNSCC with disease progression on or after platinum-containing chemotherapy.

Classical Hodgkin Lymphoma (cHL)

- for the treatment of adult patients with relapsed or refractory cHL.
- for the treatment of pediatric patients with refractory cHL, or cHL that has relapsed after 2 or more lines of therapy.

Primary Mediastinal Large B-Cell Lymphoma (PMBCL)

- for the treatment of adult and pediatric patients with refractory PMBCL, or who have relapsed after 2 or more prior lines of therapy.
- Limitations of Use: KEYTRUDA® is not recommended for treatment of patients with PMBCL who require urgent cytoreductive therapy.

Urothelial Carcinoma

- for the treatment of patients with locally advanced or metastatic urothelial carcinoma who:
 - are not eligible for any platinum-containing chemotherapy, or

Tumor Mutational Burden (continued)

- who have disease progression during or following platinum-containing chemotherapy or within 12 months of neoadjuvant or adjuvant treatment with platinum-containing chemotherapy
- for the treatment of patients with Bacillus Calmette-Guerin (BCG)-unresponsive, high-risk, non-muscle invasive bladder cancer (NMIBC) with carcinoma in situ (CIS) with or without papillary tumors who are ineligible for or have elected not to undergo cystectomy.

Microsatellite Instability-High or Mismatch Repair Deficient Cancer

- for the treatment of adult and pediatric patients with unresectable or metastatic, microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) solid tumors that have progressed following prior treatment and who have no satisfactory alternative treatment options.¹
- Limitations of Use: The safety and effectiveness of KEYTRUDA® in pediatric patients with MSI-H central nervous system cancers have not been established.

Microsatellite Instability-High or Mismatch Repair Deficient Colorectal Cancer (CRC)

- for the treatment of patients with unresectable or metastatic MSI-H or dMMR colorectal cancer (CRC).

Gastric Cancer

- in combination with trastuzumab, fluoropyrimidine- and platinum-containing chemotherapy, for the first-line treatment of patients with locally advanced unresectable or metastatic HER2-positive gastric or gastroesophageal junction (GEJ) adenocarcinoma.

Esophageal Cancer

- for the treatment of patients with locally advanced or metastatic esophageal or gastroesophageal junction (GEJ) (tumors with epicenter 1 to 5 centimeters above the GEJ) carcinoma that is not amenable to surgical resection or definitive chemoradiation either:
 - in combination with platinum- and fluoropyrimidine-based chemotherapy, or
 - as a single agent after one or more prior lines of systemic therapy for patients with tumors of squamous cell histology that express PD-L1 (CPS ≥ 10) as determined by an FDA-approved test.

Cervical Cancer

- in combination with chemotherapy, with or without bevacizumab, for the treatment of patients with persistent, recurrent, or metastatic cervical cancer whose tumors express PD-L1 (CPS ≥ 1) as determined by an FDA-approved test.
- as a single agent for the treatment of patients with recurrent or metastatic cervical cancer with disease progression on or after chemotherapy whose tumors express PD-L1 (CPS ≥ 1) as determined by an FDA-approved test.

Hepatocellular Carcinoma (HCC)

- for the treatment of patients with HCC who have been previously treated with sorafenib.¹

Merkel Cell Carcinoma (MCC)

- for the treatment of adult and pediatric patients with recurrent locally advanced or metastatic Merkel cell carcinoma.¹

Renal Cell Carcinoma (RCC)

- in combination with axitinib, for the first-line treatment of adult patients with advanced RCC.
- in combination with lenvatinib, for the first-line treatment of adult patients with advanced RCC.
- for the adjuvant treatment of patients with RCC at intermediate-high or high risk of recurrence following nephrectomy, or following nephrectomy and resection of metastatic lesions.

Endometrial Carcinoma

- in combination with lenvatinib, for the treatment of patients with advanced endometrial carcinoma that is not MSI-H or dMMR, who have disease progression following prior systemic therapy in any setting and are not candidates for curative surgery or radiation.

Tumor Mutational Burden-High (TMB-H) Cancer

- for the treatment of adult and pediatric patients with unresectable or metastatic tumor mutational burden-high (TMB-H) [≥ 10 mutations/megabase (mut/Mb)] solid tumors, as determined by an FDA-approved test, that have progressed following prior treatment and who have no satisfactory alternative treatment options.¹
- Limitations of Use: The safety and effectiveness of KEYTRUDA® in pediatric patients with TMB-H central nervous system cancers have not been established.

Cutaneous Squamous Cell Carcinoma (cSCC) and description of clinical benefit in the confirmatory trial

Tumor Mutational Burden (continued)

- for the treatment of patients with recurrent or metastatic cSCC or locally advanced cSCC that is not curable by surgery or radiation.

Triple-Negative Breast Cancer (TNBC)

- for the treatment of patients with high-risk early-stage TNBC in combination with chemotherapy as neoadjuvant treatment, and then continued as a single agent as adjuvant treatment after surgery.
- in combination with chemotherapy, for the treatment of patients with locally recurrent unresectable or metastatic TNBC whose tumors express PD-L1 (CPS ≥ 10) as determined by an FDA approved test.

Adult Indications: Additional Dosing Regimen of 400 mg Every 6 Weeks

- for use at an additional recommended dosage of 400 mg every 6 weeks for all approved adult indications.²

¹ This indication is approved under accelerated approval based on tumor response rate and durability of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in the confirmatory trials.

² This indication is approved under accelerated approval based on pharmacokinetic data, the relationship of exposure to efficacy, and the relationship of exposure to safety. Continued approval for this dosing may be contingent upon verification and description of clinical benefit in the confirmatory trials.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/125514s125lbl.pdf

ATM c.7789-2A>G

○ olaparib

Cancer type: Castration-Resistant Prostate Cancer

Label as of: 2022-01-31

Variant class: ATM mutation

Indications and usage:

LYNPARZA® is a poly (ADP-ribose) polymerase (PARP) inhibitor indicated:

Ovarian cancer

- for the maintenance treatment of adult patients with deleterious or suspected deleterious germline or somatic BRCA-mutated advanced epithelial ovarian, fallopian tube or primary peritoneal cancer who are in complete or partial response to first-line platinum-based chemotherapy. Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA®.
- in combination with bevacizumab for the maintenance treatment of adult patients with advanced epithelial ovarian, fallopian tube or primary peritoneal cancer who are in complete or partial response to first-line platinum-based chemotherapy and whose cancer is associated with homologous recombination deficiency (HRD)-positive status defined by either:
 - a deleterious or suspected deleterious BRCA mutation, and/or
 - genomic instability.

Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA®.

- for the maintenance treatment of adult patients with recurrent epithelial ovarian, fallopian tube or primary peritoneal cancer, who are in complete or partial response to platinum-based chemotherapy.
- for the treatment of adult patients with deleterious or suspected deleterious germline BRCA-mutated (gBRCAm) advanced ovarian cancer who have been treated with three or more prior lines of chemotherapy. Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA®.

Breast cancer

- for the treatment of adult patients with deleterious or suspected deleterious gBRCAm, HER2-negative metastatic breast cancer who have been treated with chemotherapy in the neoadjuvant, adjuvant or metastatic setting. Patients with hormone receptor (HR)-positive breast cancer should have been treated with a prior endocrine therapy or be considered inappropriate for endocrine therapy. Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA®.

Pancreatic cancer

- for the maintenance treatment of adult patients with deleterious or suspected deleterious gBRCAm metastatic pancreatic adenocarcinoma whose disease has not progressed on at least 16 weeks of a first-line platinum-based chemotherapy regimen. Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA®.

Prostate cancer

- for the treatment of adult patients with deleterious or suspected deleterious germline or somatic homologous recombination repair (HRR) gene-mutated metastatic castration-resistant prostate cancer (mCRPC) who have progressed following prior treatment with enzalutamide or abiraterone. Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA®.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/208558s021lbl.pdf

Current NCCN Information

- ☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

NCCN information is current as of 2022-02-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

Tumor Mutational Burden

☐ pembrolizumab

Cancer type: Chondrosarcoma, Ewing Sarcoma, Osteosarcoma **Variant class:** Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Unresectable, Metastatic, Progression (Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Bone Cancer [Version 2.2022]

☐ pembrolizumab

Cancer type: Breast Cancer **Variant class:** Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Stage IV; Invasive, Unresectable, Metastatic, Progression (Line of therapy not specified); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 2.2022]

☐ pembrolizumab

Cancer type: Cervical Small Cell Neuroendocrine Carcinoma **Variant class:** Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent, Metastatic (Second-line therapy, Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Cervical Cancer [Version 1.2022]

☐ pembrolizumab

Cancer type: Cervical Cancer **Variant class:** Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Squamous Cell, Adenocarcinoma, Adenosquamous; Recurrent, Metastatic, Progression (Second-line therapy, Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Cervical Cancer [Version 1.2022]

Tumor Mutational Burden (continued)

○ pembrolizumab

Cancer type: Esophageal Cancer,
Gastroesophageal Junction Adenocarcinoma

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Squamous Cell; Unresectable, Locally Advanced, Recurrent, Metastatic (Second-line therapy, Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 1.2022]

○ pembrolizumab

Cancer type: Gastric Cancer

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Unresectable, Locally Advanced, Recurrent, Metastatic (Second-line therapy, Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2022]

○ pembrolizumab

Cancer type: Head and Neck Cancer

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Nasopharyngeal; Recurrent, Unresectable, Metastatic (Subsequent therapy); Useful in certain circumstances
- Salivary Gland Neoplasm; Recurrent, Unresectable, Metastatic (Line of therapy not specified); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Head and Neck Cancers [Version 1.2022]

○ pembrolizumab

Cancer type: Extrahepatic Cholangiocarcinoma,
Gallbladder Carcinoma, Intrahepatic
Cholangiocarcinoma

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Unresectable, Metastatic, Progression (Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Hepatobiliary Cancers [Version 5.2021]

Tumor Mutational Burden (continued)

○ pembrolizumab

Cancer type: Large Cell Neuroendocrine Carcinoma, Small Cell Neuroendocrine Carcinoma **Variant class:** Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Poorly Differentiated; Unresectable, Metastatic, Progression (Line of therapy not specified); Consider

Reference: NCCN Guidelines® - NCCN-Neuroendocrine and Adrenal Tumors [Version 4.2021]

○ pembrolizumab

Cancer type: Neuroendocrine Tumor **Variant class:** Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Well Differentiated; G3; Locally Advanced, Metastatic, Progression (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Neuroendocrine and Adrenal Tumors [Version 4.2021]

○ pembrolizumab

Cancer type: Ovarian Cancer **Variant class:** Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Epithelial, Less Common Ovarian Cancers, Fallopian Tube, Primary Peritoneal; Recurrent (Recurrence therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Ovarian Cancer [Version 1.2022]

○ pembrolizumab

Cancer type: Castration-Resistant Prostate Cancer **Variant class:** Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma; Metastatic, Progression (Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Prostate Cancer [Version 3.2022]

○ pembrolizumab

Cancer type: Testicular Cancer **Variant class:** Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Germ Cell Tumor; Metastatic, Recurrent (Third-line therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Testicular Cancer [Version 2.2022]

Tumor Mutational Burden (continued)

○ pembrolizumab

Cancer type: Thyroid Gland Follicular Carcinoma, Thyroid Gland Hurthle Cell Carcinoma, Thyroid Gland Papillary Carcinoma

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Locally Recurrent, Advanced, Metastatic (Line of therapy not specified); Consider

Reference: NCCN Guidelines® - NCCN-Thyroid Carcinoma [Version 3.2021]

○ pembrolizumab

Cancer type: Thyroid Gland Medullary Carcinoma

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent, Persistent, Local, Distant Metastases, Regional (Line of therapy not specified); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Thyroid Carcinoma [Version 3.2021]

○ pembrolizumab

Cancer type: Thyroid Gland Anaplastic Carcinoma

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Stage IVC; Metastatic (Second-line therapy); Useful in certain circumstances, Consider

Reference: NCCN Guidelines® - NCCN-Thyroid Carcinoma [Version 3.2021]

○ pembrolizumab

Cancer type: Endometrial Carcinoma, Endometrial Clear Cell Adenocarcinoma, Endometrial Serous Adenocarcinoma, Undifferentiated and Dedifferentiated Carcinomas of the Uterine Corpus, Uterine Corpus Carcinosarcoma

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Unresectable, Metastatic, Progression (Second-line therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Uterine Neoplasms [Version 1.2022]

Tumor Mutational Burden (continued)

○ pembrolizumab

Cancer type: Uterine Sarcoma

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Unresectable, Metastatic, Progression (Second-line therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Uterine Neoplasms [Version 1.2022]

○ pembrolizumab

Cancer type: Neuroendocrine Tumor

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Well Differentiated; G3; Locally Advanced, Metastatic, Progression, Unresectable (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Neuroendocrine and Adrenal Tumors [Version 4.2021]

○ pembrolizumab

Cancer type: Castration-Resistant Prostate Cancer Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Adenocarcinoma; Visceral Metastases, Progression (Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Prostate Cancer [Version 3.2022]

ATM c.7789-2A>G

○ olaparib

Cancer type: Castration-Resistant Prostate Cancer Variant class: ATM mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Adenocarcinoma; Metastatic (Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Prostate Cancer [Version 3.2022]

ATM c.7789-2A>G (continued)

○ olaparib

Cancer type: Castration-Resistant Prostate Cancer Variant class: ATM mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Adenocarcinoma; Visceral Metastases (Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Prostate Cancer [Version 3.2022]

Clinical Trials in Taiwan region:

Clinical Trials Summary

Tumor Mutational Burden

NCT ID	Title	Phase
NCT04589845	Tumor-Agnostic Precision Immunooncology and Somatic Targeting Rational for You (TAPISTRY) Phase II Platform Trial	II
NCT02628067	A Clinical Trial of Pembrolizumab (MK-3475) Evaluating Predictive Biomarkers in Subjects With Advanced Solid Tumors (KEYNOTE 158).	II

Alerts Informed By Public Data Sources

Current NCCN Information

 Contraindicated
  Not recommended
  Resistance
  Breakthrough
  Fast Track

NCCN information is current as of 2022-02-01. For the most up-to-date information, search www.nccn.org. For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

Tumor Mutational Burden

pembrolizumab

Cancer type: Giant Cell Tumor of Soft Tissue

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "NCCN does not recommend this systemic treatment for GCTB since it is not technically a malignant tumor."

Reference: NCCN Guidelines® - NCCN-Bone Cancer [Version 2.2022]

Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

References

1. Maréchal et al. DNA damage sensing by the ATM and ATR kinases. *Cold Spring Harb Perspect Biol.* 2013 Sep 1;5(9). PMID: 24003211
2. Matsuoka et al. ATM and ATR substrate analysis reveals extensive protein networks responsive to DNA damage. *Science.* 2007 May 25;316(5828):1160-6. PMID: 17525332
3. Ditch et al. The ATM protein kinase and cellular redox signaling: beyond the DNA damage response. *Trends Biochem. Sci.* 2012 Jan;37(1):15-22. PMID: 22079189
4. Kozlov et al. Autophosphorylation and ATM activation: additional sites add to the complexity. *J. Biol. Chem.* 2011 Mar 18;286(11):9107-19. PMID: 21149446
5. Lim et al. Evaluation of the methods to identify patients who may benefit from PARP inhibitor use. *Endocr. Relat. Cancer.* 2016 Jun;23(6):R267-85. PMID: 27226207
6. Lord et al. BRCAness revisited. *Nat. Rev. Cancer.* 2016 Feb;16(2):110-20. PMID: 26775620
7. Cynthia et al. Ataxia telangiectasia: a review. *Orphanet J Rare Dis.* 2016 Nov 25;11(1):159. PMID: 27884168
8. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
9. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
10. https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/208558s021lbl.pdf
11. Gilardini et al. ATM-depletion in breast cancer cells confers sensitivity to PARP inhibition. *CR.* PMID: 24252502
12. Pennington et al. Germline and somatic mutations in homologous recombination genes predict platinum response and survival in ovarian, fallopian tube, and peritoneal carcinomas. *Clin. Cancer Res.* 2014 Feb 1;20(3):764-75. PMID: 24240112
13. Mateo et al. DNA-Repair Defects and Olaparib in Metastatic Prostate Cancer. *N. Engl. J. Med.* 2015 Oct 29;373(18):1697-708. PMID: 26510020
14. <https://www.senhwabio.com/en/news/20220125>
15. Rosset et al. TSC1 and TSC2 gene mutations and their implications for treatment in Tuberous Sclerosis Complex: a review. *Genet Mol Biol.* 2017 Jan-Mar;40(1):69-79. PMID: 28222202
16. Henske et al. Tuberous sclerosis complex. *Nat Rev Dis Primers.* 2016 May 26;2:16035. PMID: 27226234
17. Santiago et al. Identification of regions critical for the integrity of the TSC1-TSC2-TBC1D7 complex. *PLoS ONE.* 2014;9(4):e93940. PMID: 24714658
18. Yuan et al. Novel technologies and emerging biomarkers for personalized cancer immunotherapy. 4:3. PMID: 26788324
19. Temko et al. Somatic POLE exonuclease domain mutations are early events in sporadic endometrial and colorectal carcinogenesis, determining driver mutational landscape, clonal neoantigen burden and immune response. *J. Pathol.* 2018 Jul;245(3):283-296. PMID: 29604063
20. Mehnert et al. Immune activation and response to pembrolizumab in POLE-mutant endometrial cancer. 126(6):2334-40. PMID: 27159395
21. Bouffet et al. Immune Checkpoint Inhibition for Hypermutant Glioblastoma Multiforme Resulting From Germline Biallelic Mismatch Repair Deficiency. *J. Clin. Oncol.* 2016 Jul 1;34(19):2206-11. PMID: 27001570
22. Humphris et al. Hypermutation In Pancreatic Cancer. *Gastroenterology.* 2017 Jan;152(1):68-74.e2. PMID: 27856273
23. Bonneville et al. Landscape of Microsatellite Instability Across 39 Cancer Types. *JCO Precis Oncol.* 2017;2017. PMID: 29850653
24. Snyder et al. Genetic basis for clinical response to CTLA-4 blockade in melanoma. *N. Engl. J. Med.* 2014 Dec 4;371(23):2189-2199. PMID: 25409260
25. Alexandrov et al. Signatures of mutational processes in human cancer. *Nature.* 2013 Aug 22;500(7463):415-21. PMID: 23945592
26. Rizvi et al. Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer. *Science.* 2015 Apr 3;348(6230):124-8. PMID: 25765070
27. Van et al. Genomic correlates of response to CTLA-4 blockade in metastatic melanoma. *Science.* 2015 Oct 9;350(6257):207-211. PMID: 26359337
28. Chalmers et al. Analysis of 100,000 human cancer genomes reveals the landscape of tumor mutational burden. 9:34. PMID: 28420421
29. Govindan et al. Genomic landscape of non-small cell lung cancer in smokers and never-smokers. *Cell.* 2012 Sep 14;150(6):1121-34. PMID:22980976
30. Endris et al. Measurement of tumor mutational burden (TMB) in routine molecular diagnostics: in silico and real-life analysis of three larger gene panels. *Int. J. Cancer.* 2019 May 1;144(9):2303-2312. PMID: 30446996

References (continued)

31. Meléndez et al. Methods of measurement for tumor mutational burden in tumor tissue. *Transl Lung Cancer Res.* 2018 Dec;7(6):661-667. PMID: 30505710
32. Hellmann et al. Nivolumab plus Ipilimumab in Lung Cancer with a High Tumor Mutational Burden. *N. Engl. J. Med.* 2018 May 31;378(22):2093-2104. PMID: 29658845
33. Ready et al. First-Line Nivolumab Plus Ipilimumab in Advanced Non-Small-Cell Lung Cancer (CheckMate 568): Outcomes by Programmed Death Ligand 1 and Tumor Mutational Burden as Biomarkers. *J. Clin. Oncol.* 2019 Apr 20;37(12):992-1000. PMID: 30785829
34. Alborelli et al. Tumor mutational burden assessed by targeted NGS predicts clinical benefit from immune checkpoint inhibitors in non-small cell lung cancer. *J. Pathol.* 2020 Jan;250(1):19-29. PMID: 31471895
35. Heeke et al. In-house Implementation of Tumor Mutational Burden Testing to Predict Durable Clinical Benefit in Non-small Cell Lung Cancer and Melanoma Patients. *Cancers (Basel).* 2019 Aug 29;11(9). PMID: 31470674
36. https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/125514s125lbl.pdf
37. F. et al. CheckMate 817: First-Line Nivolumab + Ipilimumab in Patients with ECOG PS 2 and Other Special Populations with Advanced NSCLC. Volume 14, Issue 10, Supplement, Pages S214–S215. Abstract OA04.02
38. Zhu et al. Association Between Tumor Mutation Burden (TMB) and Outcomes of Cancer Patients Treated With PD-1/PD-L1 Inhibitions: A Meta-Analysis. *Front Oncol*, 9:1161, 04 Nov 2019. PMID: 31258479
39. Castellanos et al. Tumor mutational burden (TMB) and PD-L1 expression as predictors of response to immunotherapy (IO) in NSCLC. *Journal of Clinical Oncology* 37, no. 15_suppl (May 20, 2019) 2630-2630. Abstract 2630
40. NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2022]
41. NCCN Guidelines® - NCCN-Bone Cancer [Version 2.2022]
42. NCCN Guidelines® - NCCN-Head and Neck Cancers [Version 1.2022]
43. NCCN Guidelines® - NCCN-Testicular Cancer [Version 2.2022]
44. NCCN Guidelines® - NCCN-Cervical Cancer [Version 1.2022]
45. NCCN Guidelines® - NCCN-Uterine Neoplasms [Version 1.2022]
46. NCCN Guidelines® - NCCN-Thyroid Carcinoma [Version 3.2021]
47. NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 1.2022]
48. NCCN Guidelines® - NCCN-Ovarian Cancer [Version 1.2022]
49. NCCN Guidelines® - NCCN-Breast Cancer [Version 2.2022]
50. <https://www.focr.org/tmb>
51. <http://www.iqnpaath.org/category/tmb>
52. Stenzinger et al. Tumor mutational burden standardization initiatives: Recommendations for consistent tumor mutational burden assessment in clinical samples to guide immunotherapy treatment decisions. *Genes Chromosomes Cancer.* 2019 Aug;58(8):578-588. PMID: 30664300
53. Merino et al. Establishing guidelines to harmonize tumor mutational burden (TMB): in silico assessment of variation in TMB quantification across diagnostic platforms: phase I of the Friends of Cancer Research TMB Harmonization Project. *J Immunother Cancer.* 2020 Mar;8(1). PMID: 32217756