

# ACT Onco<sup>®</sup> + Report

PATIENT		
Name: 姚政宏		Patient ID: 29095690
Date of Birth: Mar 28, 1960		Gender: Male
Diagnosis: Lung adenocarcinoma		
ORDERING PHYSICIAN		
Name: 陳育民醫師		Tel: 886-228712121
Facility: 臺北榮總		
Address: 臺北市北投區石牌路二段 201 號		
SPECIMEN		
Specimen ID: S11137673D	Collection site: Pleura	Type: FFPE tissue
Date received: Sep 28, 2022	Lab ID: AA-22-05804	D/ID: NA

## ABOUT ACT Onco<sup>®</sup>+

The test is a next-generation sequencing (NGS)-based assay developed for efficient and comprehensive genomic profiling of cancers. This test interrogates coding regions of 440 genes associated with cancer treatment, prognosis and diagnosis. Genetic mutations detected by this test include small-scale mutations like single nucleotide variants (SNVs), small insertions and deletions (InDels) ( $\leq 15$  nucleotides) and large-scale genomic alterations like copy number alterations (CNAs). The test also includes an RNA test, detecting fusion transcripts of 13 genes.

## SUMMARY FOR ACTIONABLE VARIANTS

### VARIANTS/BIOMARKERS WITH EVIDENCE OF CLINICAL SIGNIFICANCE

Genomic Alterations/Biomarkers	Probable Effects in Patient's Cancer Type		Probable Sensitive in Other Cancer Types
	Sensitive	Resistant	
EGFR L747_P753delinsS (Exon 19 deletion)	Afatinib, Dacomitinib, Erlotinib, Gefitinib, Osimertinib	-	-
TMB-High	Atezolizumab, Cemiplimab-rwlc, Dostarlimab-gxly, Durvalumab, Ipilimumab, Nivolumab, Pembrolizumab	-	Avelumab

### VARIANTS/BIOMARKERS WITH POTENTIAL CLINICAL SIGNIFICANCE

Genomic Alterations/Biomarkers	Possibly Sensitive	Possibly Resistant
EGFR L747_P753delinsS (Exon 19 deletion)	Mobocertinib	Cetuximab
EGFR Amplification	Afatinib, Erlotinib, Gefitinib, Osimertinib, Cetuximab, Panitumumab, Necitumumab	-

#### Note:

- The above summary tables present genomic variants and biomarkers based on the three-tiered approach proposed by US FDA for reporting tumor profiling NGS testing. "Variants/biomarkers with evidence of clinical significance" refers to mutations that are widely recognized as standard-of-care biomarkers (FDA level 2/AMP tier 1). "Variants/biomarkers with potential clinical significance" refers to mutations that are not included in the standard of care but are informational for clinicians, which are commonly biomarkers used as inclusion criteria for clinical trials (FDA level 3/AMP tier 2).
- The therapeutic agents and possible effects to a given drug are based on mapping the variants/biomarkers with ACT Genomics clinical knowledge database. The mapping results only provide information for reference, but not medical recommendation.
- Please refer to corresponding sections for more detailed information about genomic alteration and clinical relevance listed above.

# ACT Onco<sup>®</sup> + Report

## TESTING RESULTS

### VARIANT(S) WITH CLINICAL RELEVANCE

#### - Single Nucleotide and Small InDel Variants

Gene	Amino Acid Change	Allele Frequency
<i>EGFR</i>	L747_P753delinsS (Exon 19 deletion)	68.2%
<i>KMT2D</i>	Q2458*	18.1%

#### - Copy Number Alterations

Chromosome	Gene	Variation	Copy Number
Chr7	<i>EGFR</i>	Amplification	12
Chr11	<i>BIRC2</i>	Amplification	19
Chr8	<i>MYC</i>	Amplification	89

#### - Fusions

Fusion Gene & Exon	Transcript ID
No fusion gene detected in this sample	

#### - Immune Checkpoint Inhibitor (ICI) Related Biomarkers

Biomarker	Results
Tumor Mutational Burden (TMB)	7.5 muts/Mb (TMB-High)
Microsatellite Instability (MSI)	Microsatellite stable (MSS)

#### Note:

- Variant(s) enlisted in the SNV table may currently exhibit no relevance to treatment response prediction. Please refer to INTERPRETATION for more biological information and/or potential clinical impacts of the variants.
- Loss of heterozygosity (LOH) information was used to infer tumor cellularity. Copy number alteration in the tumor was determined based on 35% tumor purity.
- TMB was calculated by using the sequenced regions of ACTOnco<sup>®</sup> to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The threshold for high mutation load is set at  $\geq 7.5$  mutations per megabase. TMB, microsatellite status and gene copy number deletion cannot be determined if calculated tumor purity is  $< 30\%$ .

# ACT Onco<sup>®</sup> + Report

## THERAPEUTIC IMPLICATIONS TARGETED THERAPIES

Genomic Alterations	Therapies	Effect
<b>Level 1</b>		
<b>EGFR</b> L747_P753delinsS (Exon 19 deletion)	Afatinib, Dacomitinib, Erlotinib, Gefitinib, Osimertinib	<b>sensitive</b>
<b>Level 3B</b>		
<b>EGFR</b> Amplification	Afatinib, Erlotinib, Gefitinib, Osimertinib, Cetuximab, Panitumumab, Necitumumab	<b>sensitive</b>
<b>Level 4</b>		
<b>EGFR</b> L747_P753delinsS (Exon 19 deletion)	Mobocertinib	<b>sensitive</b>
<b>EGFR</b> L747_P753delinsS (Exon 19 deletion)	Cetuximab	<b>resistant</b>

Therapies associated with benefit or lack of benefit are based on biomarkers detected in this tumor and published evidence in professional guidelines or peer-reviewed journals.

Level	Description
<b>1</b>	FDA-recognized biomarkers predictive of response or resistance to FDA approved drugs in this indication
<b>2</b>	Standard care biomarkers (recommended by the NCCN guideline) predictive of response or resistance to FDA approved drugs in this indication
<b>3A</b>	Biomarkers predictive of response or resistance to therapies approved by the FDA or NCCN guideline in a different cancer type
<b>3B</b>	Biomarkers that serve as inclusion criteria for clinical trials (minimal supportive data required)
<b>4</b>	Biomarkers that show plausible therapeutic significance based on small studies, few case reports, or preclinical studies

# ACT Onco<sup>®</sup> + Report

## IMMUNE CHECKPOINT INHIBITORS (ICIs)

Genomic Alterations	Approved for Patient's Cancer Type	Approved for Other Cancer Type
<b>TMB-High</b> (7.5 muts/Mb)	Atezolizumab, Cemiplimab-rwlc, Dostarlimab-gxly, Durvalumab, Ipilimumab, Nivolumab, Pembrolizumab	Avelumab

TMB, Tumor Mutational Burden; Muts/Mb, mutations per megabase

## - Other Biomarkers with Potential Clinical Effects for ICIs

Genomic Alterations	Potential Clinical Effects
EGFR aberration	Likely associated with WORSE response to ICIs

Note: Tumor non-genomic factors, such as patient germline genetics, PDL1 expression, tumor microenvironment, epigenetic alterations or other factors not provided by this test may affect ICI response.

## CHEMOTHERAPIES

Genomic Alterations	Therapies	Effect	Level of Evidence	Cancer Type
<b>MYC</b> Amplification	FAC CMF and P-FEC regimens	<b>Sensitive</b>	Clinical	Breast cancer
	Platinum-based regimens	<b>Sensitive</b>	Clinical	Ovarian cancer

## HORMONAL THERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to hormonal therapies.

## OTHERS

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to other therapies.

### Note:

Therapeutic implications provided in the test are based solely on the panel of 440 genes sequenced. Therefore, alterations in genes not covered in this panel, epigenetic and post-transcriptional and post-translational factors may also determine a patient's response to therapies. In addition, several other patient-associated clinical factors, including but not limited to, prior lines of therapies received, dosage and combinations with other therapeutic agents, patient's cancer types, sub-types, and/or stages, may also determine the patient's clinical response to therapies.

# ACT Onco<sup>®</sup> + Report

## VARIANT INTERPRETATION

### Tumor mutational burden (TMB): High (7.5 mutations / Mb)

High TMB is a potential biomarker that predicts response to immune checkpoint inhibitors, including anti-CTLA-4 and anti-PD-1 in melanoma, anti-PD-1 in non-small cell lung cancer (NSCLC) and colorectal cancer (CRC), cutaneous squamous cell carcinoma (CSCC), and anti-PD-L1 therapy in bladder cancer<sup>[1][2][3][3][4][5][6][7][8]</sup>. Of note, the U.S. FDA has approved tumor mutational burden-high (TMB-H) as a predictive biomarker for pembrolizumab in adult and pediatric patients with unresectable or metastatic solid tumor who have progressed following prior treatment and have no satisfactory alternative treatment options. CRCs with defects in mismatch-repair (MMR) are more susceptible to PD-1 blockade<sup>[6]</sup>. High mutation load is associated with shorter overall survival in lung cancer and breast cancer patients<sup>[9][10]</sup>.

### EGFR L747\_P753delinsS (Exon 19 deletion), Amplification

#### Biological Impact

The EGFR gene encodes for the Epidermal Growth Factor Receptor, a receptor tyrosine kinase which binds to its ligands, including Epidermal Growth Factor (EGF) and Transforming Growth Factor-alpha (TGF-alpha), activates downstream signaling pathways, including the canonical oncogenic MAPK and PI3K/AKT/mTOR signaling cascades<sup>[11]</sup>. Increased EGFR activity by mutations and/or amplification of the EGFR gene has been described in a wide range of cancers, such as lung, brain, colorectal and head and neck cancer<sup>[12]</sup>. Mutations in the kinase domain of EGFR are commonly observed in non-small cell lung cancer (NSCLC), resulting in a constitutively activated form of the receptor<sup>[13]</sup>. On the other hand, in the brain and colorectal cancers, the most prevalent EGFR alteration is copy number amplification that results in receptor overexpression<sup>[14]</sup>.

EGFR L747\_P753delinsS (exon 19 deletion) lies within the tyrosine kinase domain of EGFR, resulting in a deletion of 7 amino acids from residues 747 to 753 and insertion of a serine residue (UniProtKB).

EGFR exon 19 deletions are in-frame deletions of 9–24 nucleotides in exon 19 centred around codons 746–750 of the kinase domain of EGFR. The two most common EGFR alterations, L858R mutation and exon 19 deletions can result in constitutive activation of signal transduction pathways, leading to cell proliferation or anti-apoptosis without ligand binding<sup>[15]</sup>.

#### Therapeutic and prognostic relevance

There is accumulated clinical evidence suggested that patients with MDM2/MDM4 amplification or EGFR aberrations exhibited poor clinical outcome and demonstrated a significantly increased rate of tumor growth (hyper-progression) after receiving immune checkpoint (PD-1/PD-L1) inhibitors therapies<sup>[16]</sup> (Annals of Oncology (2017) 28 (suppl\_5): v403-v427. 10.1093/annonc/mdx376).

There is accumulated clinical evidence reported that patients with lung adenocarcinoma or NSCLC harboring EGFR L747\_P753delinsS responded to EGFR TKIs<sup>[17][18][19]</sup>. In preclinical studies, cells expression EGFR L747\_P753delinsS were sensitive to mobocertinib, but resistant to cetuximab<sup>[20][21]</sup>.

The first- and second-generation EGFR tyrosine kinase inhibitors (EGFR-TKIs), dacomitinib, erlotinib, gefitinib and afatinib have been approved by the U.S. Food and Drug Administration (FDA) as the first-line treatment in non-small cell lung cancer (NSCLC) patients whose tumor carries EGFR exon 19 deletion or L858R mutation<sup>[22][23][24]</sup>, as detected by a U.S. FDA-approved test. A Phase III clinical trial (NCT01774721) show that dacomitinib significantly improved progression-free survival over gefitinib in first-line treatment of patients with EGFR-mutation-positive NSCLC<sup>[22]</sup>. Another Phase III clinical trial (NCT00949650) demonstrated that median progression-free survival (PFS) among lung cancer patients with exon 19 deletion or L858R EGFR mutation (n=308) was 13.6 months for afatinib and 6.9 months for chemotherapy. The EGFR T790M mutation has been demonstrated to confer resistance to TKIs (dacomitinib,

# ACT Onco<sup>®</sup> + Report

gefitinib, erlotinib, and afatinib) in preclinical and clinical studies<sup>[25][26][27][28]</sup>.

Osimertinib, a third-generation irreversible EGFR-TKI that selectively inhibits both EGFR-TKI-sensitizing and EGFR T790M resistance mutations, has been approved by the U.S. FDA for NSCLC patient harboring T790M mutation-positive tumor<sup>[29][30][31]</sup>. Results from a double-blind, Phase 3 trial further showed that osimertinib significantly demonstrated longer PFS than standard EGFR-TKIs (18.9 months vs. 10.2 months) in previously untreated EGFR mutation-positive (exon 19 deletion or L858R) advanced NSCLC<sup>[32]</sup>.

Increased EGFR copy number is associated with tumor response to panitumumab, an EGFR-targeted antibody, in colorectal cancer patients, based on data from a phase III study<sup>[33]</sup>. A recent Phase II trial of cetuximab (another approved anti-EGFR antibody) oxaliplatin/leucovorin/5-fluorouracil therapy in first-line setting also demonstrated an association between higher EGFR copy number and better overall survival in gastric cancer patients<sup>[34]</sup>. The addition of cetuximab to chemotherapy reduced the risk of death by 44% for advanced squamous non-small cell lung cancer (NSCLC) patients with EGFR-amplified tumor, according to clinical trial findings presented at the 2015 World Conference on Lung Cancer. Preclinical data of gastric cancer (GC)-derived xenograft also showed that EGFR amplification or overexpression is associated with response to cetuximab<sup>[35]</sup>. Besides, a phase III study of necitumumab showed squamous cell lung cancer patients with EGFR amplification had improved overall survival (14.8 versus 7.6 months,  $p = 0.033$ ) (NCT00981058)<sup>[36]</sup>.

Increased EGFR copy number has been shown to be associated with better response and survival in gefitinib or erlotinib treatment for NSCLC<sup>[37][38][39][40][41][42]</sup>, esophageal cancer<sup>[43]</sup>, and mucinous urethral adenocarcinoma<sup>[44]</sup>. Concurrent amplification of EGFR and ERBB2 is associated with response to afatinib in patients with trastuzumab-refractory esophagogastric cancer<sup>[45]</sup>. However, dacomitinib has been reported with a limited single-agent activity in recurrent glioblastoma with EGFR amplification in a phase II trial<sup>[46]</sup>. EGFR amplification has been determined as an inclusion criterion for the trials evaluating erlotinib, afatinib, and osimertinib efficacy in PDAC with co-expressing EGFR and c-Met (NCT03213626), glioblastoma (NCT03732352), urothelial tract carcinoma (NCT02780687), and brain cancer (NCT02423525).

## KMT2D Q2458\*

### Biological Impact

KMT2D (Lysine methyltransferase 2D) gene encodes the histone methyltransferase MLL2, which methylates lysine residue 4 on the tail of histone H3 (H3K4) and regulates gene expression via modulating chromatin structures<sup>[47]</sup>. KMT2D mutations have been reported in bladder cancer, diffuse large B cell lymphoma (DLBCL), non-Hodgkin lymphoma, and acute myeloid leukemia<sup>[48][49][50][51]</sup>, and deletion of KMT2D has been reported to lead to genomic instability in vitro<sup>[52]</sup>.

Q2458\* mutation results in a premature truncation of the KMT2D protein at amino acid 2458 (UniProtKB). This mutation is predicted to lead to a loss of KMT2D function, despite not having characterized in the literature.

### Therapeutic and prognostic relevance

A study of non-small cell lung cancer patients ( $n=194$ ) indicated that patients harboring mutant KMT2D had shorter overall survival and progression-free survival compared with patients with wild-type KMT2D. However, this correlation had not found in small cell lung cancer patients<sup>[53]</sup>.

Low levels of KMT2D expression was associated with better overall survival in pancreatic ductal adenocarcinoma (PDAC)<sup>[54]</sup>, esophageal squamous cell carcinoma (ESCC)<sup>[55]</sup>, and better disease-free survival in prostate cancer<sup>[56]</sup>. However, low expression of KMT2D had been reported to correlate with advanced stages and imatinib resistance in chronic myeloid leukemia (CML)<sup>[57]</sup>.



# ACT Onco<sup>®</sup> + Report

## BIRC2 Amplification

### Biological Impact

Baculoviral IAP repeat containing 2 (BIRC2) gene encodes cellular inhibitor of apoptosis protein 1 and is involved in regulating TNF and NF- $\kappa$ B activated signaling pathways<sup>[58]</sup>.

BIRC2 amplification has been identified in squamous cell carcinomas of the uterine cervix, head and neck squamous cellular carcinoma and oral squamous cell carcinoma<sup>[59][60]</sup>. Increased BIRC2 copy number was associated with higher expression level of BIRC2 protein<sup>[61]</sup>. BIRC2 is required for cell migration in vitro<sup>[59]</sup>, and inhibits non-canonical NF- $\kappa$ B activity, resulting in reduction of CXCL9 and immunosuppression<sup>[62]</sup>.

### Therapeutic and prognostic relevance

In oral squamous cell carcinoma (OSCC), BIRC2 amplification was associated with poor clinical outcome<sup>[59]</sup>. A preclinical study demonstrated BIRC2 expression impairs the sensitivity of anti-CTLA4 and/or anti-PD1 immune checkpoint inhibitors in animal models<sup>[62]</sup>.

## MYC Amplification

### Biological Impact

The v-myc avian myelocytomatosis viral oncogene homolog, also known as c-myc (MYC) gene encodes a transcription factor involved in cellular proliferation, inhibiting exit from the cell cycle, stimulating vascularization and enhancing genomic instability<sup>[63][64][65]</sup>. Dysregulated MYC expression is implicated in a wide range of human cancers<sup>[66]</sup>.

### Therapeutic and prognostic relevance

MYC amplification was associated with better clinical outcome in breast cancer patients treated with FAC (5-fluorouracil, doxorubicin, and cyclophosphamide), CMF (cyclophosphamide, methotrexate and 5-fluorouracil) and P-FEC (paclitaxel followed by 5-fluorouracil, epirubicin and cyclophosphamide) and higher expression of MYC was also associated with a better response rate in platinum-treated ovarian cancer patients<sup>[67][68][69]</sup>.

CDK inhibition using the dinaciclib, a CDK1, 2, 5 and 9 inhibitors, exerted antitumor activity in triple-negative breast cancer (TNBC) tumor xenograft and cell lines with increased activity of the MYC pathway<sup>[70][71]</sup>.

Overexpression of MYC has been reported as a favorable prognostic biomarker in colorectal carcinoma (CRC)<sup>[72][73]</sup>. However, the favorable prognostic value of MYC in CRC is abrogated by the TP53 mutation<sup>[73]</sup>.

MYC amplification with the loss of tumor suppressor pathways such as p53 and RB has been shown to be associated with poor outcomes and was correlated with shortened disease-free survival in breast cancer with BRCA1 deficiency in TNBC<sup>[70][74]</sup>.

# ACT Onco<sup>®</sup> + Report

## US FDA-APPROVED DRUG(S)

### Afatinib (GILOTRIF)

Afatinib acts as an irreversible covalent inhibitor of the ErbB family of receptor tyrosine kinases, including epidermal growth factor receptor (EGFR) and erbB-2 (HER2). Afatinib is developed and marketed by Boehringer Ingelheim under the trade name GILOTRIF (United States) and GIOTRIF (Europe).

### - FDA Approval Summary of Afatinib (GILOTRIF)

LUX-Lung 8 <sup>[75]</sup> NCT01523587	Non-small cell lung carcinoma (Approved on 2016/04/15)
	EGFR ex19del or L858R
	Afatinib vs. Erlotinib [PFS(M): 2.4 vs. 1.9]
LUX-Lung 3 <sup>[76]</sup> NCT00949650	Non-small cell lung carcinoma (Approved on 2013/07/13)
	EGFR ex19del or L858R
	Afatinib vs. Pemetrexed + cisplatin [PFS(M): 11.1 vs. 6.9]

### Atezolizumab (TECENTRIQ)

Atezolizumab is a humanized, anti-programmed cell death-ligand 1 (PD-L1) monoclonal antibody of the IgG1 isotype, which can lead to the reactivation of immune cells that might recognize and attack tumor cells. Atezolizumab is developed and marketed by Genentech/Roche under the trade name TECENTRIQ.

### - FDA Approval Summary of Atezolizumab (TECENTRIQ)

IMpower010 NCT02486718	Non-small cell lung carcinoma (Approved on 2021/10/15)
	PD-L1
	Atezolizumab vs. Best supportive care (bsc) [DFS (PD-L1 TC≥1%)(M): not reached vs. 35.3]
IMbrave150 NCT03434379	Hepatocellular carcinoma (Approved on 2020/05/29)
	-
	Atezolizumab plus bevacizumab vs. Sorafenib [PFS(M): 6.8 vs. 4.3, OS(M): NR vs. 13.2]
IMpower133 <sup>[77]</sup> NCT02763579	Small cell lung cancer (Approved on 2019/03/18)
	-
	Atezolizumab plus carboplatin and etoposide vs. Carboplatin and etoposide [PFS(M): 5.2 vs. 4.3, OS(M): 12.3 vs. 10.3]
OAK <sup>[78]</sup> NCT02008227	Non-small cell lung carcinoma (Approved on 2016/10/18)
	PD-L1
	Atezolizumab vs. Docetaxel [OS(M): 13.8 vs. 9.6]
POPLAR <sup>[79]</sup> NCT01903993	Non-small cell lung carcinoma (Approved on 2016/10/18)
	PD-L1
	Atezolizumab vs. Docetaxel [OS(M): 12.6 vs. 9.7]
IMvigor210 <sup>[80]</sup> NCT02951767	Bladder urothelial carcinoma (Approved on 2016/05/18)
	-
	Atezolizumab [ORR (PD-L1 < 5%)(%): 21.8, ORR (PD-L1 ≥ 5%)(%): 28.1]



# ACT Onco<sup>®</sup> + Report

## Avelumab (BAVENCIO)

Avelumab is fully human monoclonal programmed death ligand-1 (PD-L1) antibody, belonging to the group of immune checkpoint blockade cancer therapies. Avelumab is developed and marketed by Merck KGaA and Pfizer under the trade name BAVENCIO.

### - FDA Approval Summary of Avelumab (BAVENCIO)

<b>JAVELIN Renal 101</b> <sup>[81]</sup> NCT02684006	<b>Renal cell carcinoma</b> (Approved on 2019/05/14)
	- Avelumab plus axitinib vs. Sunitinib [ORR(%): 51.4 vs. 25.7, PFS(M): 13.8 vs. 8.4]
<b>JAVELIN Solid Tumor</b> NCT01772004	<b>Bladder urothelial carcinoma</b> (Approved on 2017/05/09)
	- Avelumab [ORR(13W)(%): 13.6, ORR(6M)(%): 16.1]
<b>JAVELIN Merkel 200</b> <sup>[82]</sup> NCT02155647	<b>Merkel cell carcinoma</b> (Approved on 2017/03/23)
	- Avelumab [ORR(%): 33.0, DOR(M): 2.8 to 23.3+]

## Cemiplimab-rwlc (LIBTAYO)

Cemiplimab-rwlc is a recombinant human IgG4 monoclonal antibody that binds to PD-1 and blocks its interaction with PD-L1 and PD-L2. Cemiplimab-rwlc is developed and marketed by Sanofi and Regeneron under the trade name LIBTAYO.

### - FDA Approval Summary of Cemiplimab-rwlc (LIBTAYO)

<b>Study 1624</b> NCT03088540	<b>Non-small lung cancer</b> (Approved on 2021/02/22)
	<b>PD-L1</b> Cemiplimab-rwlc vs. Platinum-based chemotherapy [PFS(M): 6.2 vs. 5.6, OS(M): 22.1 vs. 14.3]
<b>Study 1620</b> NCT03132636	<b>Locally advanced basal cell carcinoma (labcc)</b> (Approved on 2021/02/09)
	- Cemiplimab-rwlc [ORR(%): 29.0, DOR(M): NR]
<b>Study 1620</b> NCT03132636	<b>Metastatic basal cell carcinoma (mbcc)</b> (Approved on 2021/02/09)
	- Cemiplimab-rwlc [ORR(%): 21.0, DOR(M): NR]
<b>Study 1423, Study 1540</b> <sup>[7]</sup> NCT02383212, NCT02760498	<b>cutaneous squamous cell carcinoma</b> (Approved on 2018/09/28)
	- Cemiplimab-rwlc [ORR(%): 47.2]

## Cetuximab (ERBITUX)

Cetuximab is a recombinant, chimeric (human/mouse) monoclonal antibody that binds to the extracellular domain and inhibits epidermal growth factor receptor (EGFR). Cetuximab is developed by ImClone and marketed by Eli Lilly under the trade name ERBITUX.

### - FDA Approval Summary of Cetuximab (ERBITUX)

<b>BEACON CRC</b> <sup>[83]</sup> NCT02928224	<b>Colorectal cancer</b> (Approved on 2020/04/08)
	<b>BRAF V600E</b> Encorafenib in combination with cetuximab vs. Irinotecan or folfox with cetuximab [OS(M): 8.4 vs. 5.4]

# ACT Onco<sup>®</sup> + Report

<b>CRYSTAL</b> <sup>[84]</sup> NCT00154102	<b>Colorectal cancer</b> (Approved on 2012/07/06)
	<b>KRAS Wild-type/EGFR-expressing</b> Cetuximab + 5-fluorouracil + folinic acid + irinotecan vs. 5-fluorouracil + folinic acid + irinotecan [PFS(M): 8.9 vs. 8.1]
<b>EXTREME</b> <sup>[85]</sup> NCT00122460	<b>Head and neck cancer</b> (Approved on 2011/11/07)
	- Cetuximab + cisplatin/carboplatin + 5-fu vs. Cisplatin/carboplatin + 5-fu [OS(M): 10.1 vs. 7.4]
<sup>[86]</sup> NCT00004227	<b>Head and neck cancer</b> (Approved on 2006/03/01)
	- Cetuximab + radiation vs. Radiation [OS(M): 49 vs. 29.3]
<sup>[87]</sup> NCT00063141	<b>Colorectal cancer</b> (Approved on 2004/02/12)
	<b>KRAS Wild-type/EGFR-expressing</b> Cetuximab + irinotecan vs. Irinotecan [DOR(M): 5.7 vs. 4.2]

## Dacomitinib (VIZIMPRO)

Dacomitinib is an oral kinase inhibitor that targets EGFR. Dacomitinib is developed and marketed by Pfizer under the trade name VIZIMPRO.

### - FDA Approval Summary of Dacomitinib (VIZIMPRO)

<b>ARCHER 1050</b> <sup>[22]</sup> NCT01774721	<b>Non-small cell lung carcinoma</b> (Approved on 2018/09/27)
	<b>EGFR ex19del or L858R</b> Dacomitinib vs. Gefitinib [PFS(M): 14.7 vs. 9.2]

## Dostarlimab-gxly (JEMPERLI)

Dostarlimab-gxly is a programmed death receptor-1 (PD-1)-blocking antibody. Dostarlimab-gxly is developed and marketed by GlaxoSmithKline LLC under the trade name JEMPERLI.

### - FDA Approval Summary of Dostarlimab-gxly (JEMPERLI)

<b>GARNET</b> NCT02715284	<b>Cancer</b> (Approved on 2021/08/17)
	<b>dMMR</b> Dostarlimab [ORR(%): 41.6, DoR(M): 34.7]
<b>GARNET (Cohort A)</b> NCT02715284	<b>Endometrial carcinoma</b> (Approved on 2021/04/22)
	<b>dMMR</b> Dostarlimab-gxly [ORR(%): 42.3, DOR(M): NR]

# ACT Onco<sup>®</sup> + Report

## Durvalumab (IMFINZI)

Durvalumab is a programmed death ligand-1 (PD-L1)-blocking antibody. Durvalumab is developed and marketed by AstraZeneca under the trade name IMFINZI.

### - FDA Approval Summary of Durvalumab (IMFINZI)

<b>TOPAZ-1</b> NCT03875235	<b>Biliary tract cancer</b> (Approved on 2022/09/02)
	- Durvalumab [OS(M): 12.8 vs. 11.5]
<b>CASPIAN<sup>[88]</sup></b> NCT03043872	<b>Extensive-stage small cell lung cancer</b> (Approved on 2020/03/27)
	- Durvalumab + etoposide + carboplatin or durvalumab + etoposide + cisplatin vs. Etoposide + carboplatin or etoposide + cisplatin [OS(M): 13 vs. 10.3]
<b>PACIFIC<sup>[89]</sup></b> NCT02125461	<b>Non-small cell lung carcinoma</b> (Approved on 2018/02/16)
	- Durvalumab vs. Placebo [PFS(M): 16.8 vs. 5.6]

## Erlotinib (TARCEVA)

Erlotinib is a small molecule, reversible inhibitor of epidermal growth factor receptor (EGFR), a receptor tyrosine kinase. Erlotinib is developed by OSI Pharmaceuticals, Genentech and Roche, and marketed by Astellas Pharm Global Development under the trade name TARCEVA.

### - FDA Approval Summary of Erlotinib (TARCEVA)

<b>RELAY</b> NCT02411448	<b>Non-small cell lung carcinoma</b> (Approved on 2020/05/29)
	<b>EGFR ex19del or L858R</b> Erlotinib + ramucirumab vs. Erlotinib + placebo [PFS(M): 19.4 vs. 12.4]
<b>EURTAC<sup>[90]</sup></b> NCT00446225	<b>Non-small cell lung carcinoma</b> (Approved on 2013/05/14)
	<b>EGFR ex19del or L858R</b> Erlotinib vs. Cisplatin + gemcitabine or cisplatin + docetaxel or carboplatin + gemcitabine or carboplatin + docetaxel [PFS(M): 10.4 vs. 5.2]
<b>PA.3<sup>[91]</sup></b> NCT00026338	<b>Pancreatic cancer</b> (Approved on 2005/11/02)
	- Gemcitabine vs. Placebo [OS(M): 6.4 vs. 6]

## Gefitinib (IRESSA)

Gefitinib is a small molecule inhibitor of epidermal growth factor receptor (EGFR), a receptor tyrosine kinase. Gefitinib is developed and marketed by AstraZeneca under the trade name IRESSA.

### - FDA Approval Summary of Gefitinib (IRESSA)

<b>IFUM<sup>[92]</sup></b> NCT01203917	<b>Non-small cell lung carcinoma</b> (Approved on 2015/07/13)
	<b>EGFR ex19del or L858R</b> Gefitinib [ORR(%): 50.0]

# ACT Onco<sup>®</sup> + Report

## Ipilimumab (YERVOY)

Ipilimumab is a fully human monoclonal antibody against the cytotoxic T-lymphocyte associated protein 4 (CTLA-4), an immune checkpoint protein receptor, to reactivate the immune responses. Ipilimumab is developed by Medarex and Bristol-Myers Squibb, and marketed by the latter under the trade name YERVOY.

### - FDA Approval Summary of Ipilimumab (YERVOY)

<b>CHECKMATE-648</b> NCT03143153	<b>Esophagus squamous cell carcinoma</b> (Approved on 2022/05/27)
	- Nivolumab and ipilimumab vs. Chemotherapy [OS(M): 12.8 vs. 10.7]
<b>CHECKMATE-743</b> NCT02899299	<b>Pleural mesothelioma</b> (Approved on 2020/10/02)
	- Nivolumab + ipilimumab vs. Pemetrexed + cisplatin [OS(M): 18.1 vs. 14.1]
<b>CHECKMATE-9LA</b> NCT03215706	<b>Non-small cell lung carcinoma</b> (Approved on 2020/05/26)
	- Nivolumab + ipilimumab + platinum-doublet chemotherapy vs. Platinum-doublet chemotherapy [OS(M): 14.1 vs. 10.7]
<b>CHECKMATE-227</b> NCT02477826	<b>Non-small cell lung carcinoma</b> (Approved on 2020/05/15)
	<b>PD-L1</b> Nivolumab + ipilimumab vs. Platinum-doublet chemotherapy [OS(M): 17.1 vs. 14.9]
<b>CHECKMATE-040</b> NCT01658878	<b>Hepatocellular carcinoma</b> (Approved on 2020/03/10)
	- Nivolumab + ipilimumab [ORR(%): 33.0]
<b>CHECKMATE-142</b> <sup>[93]</sup> NCT02060188	<b>Colorectal cancer</b> (Approved on 2018/07/10)
	<b>MSI-H or dMMR</b> Ipilimumab plus nivolumab vs. Nivolumab [ORR(%): 49.0 vs. 32.0]
<b>CHECKMATE-214</b> <sup>[94]</sup> NCT02231749	<b>Renal cell carcinoma</b> (Approved on 2018/04/16)
	- Nivolumab plus ipilimumab vs. Sunitinib [OS(M): 67.1 vs. 55.5]
<b>EORTC 18071</b> <sup>[95]</sup> NCT00636168	<b>Melanoma</b> (Approved on 2015/10/28)
	- Ipilimumab vs. Placebo [RFS(M): 26 vs. 17]
<b>MDX010-20</b> <sup>[96]</sup> NCT00094653	<b>Melanoma</b> (Approved on 2011/03/25)
	- Ipilimumab vs. Peptide vaccine with incomplete freund's adjuvant (gp100) [OS(M): 10 vs. 6]

## Mobocertinib (EXKIVITY)

Mobocertinib is a first-in-class, oral tyrosine kinase inhibitor (TKI) specifically designed to selectively target epidermal growth factor receptor (EGFR) Exon 20 insertion mutations. Mobocertinib is developed and marketed by Takeda under the trade name EXKIVITY.

### - FDA Approval Summary of Mobocertinib (EXKIVITY)

<b>Study 101</b> <sup>[97]</sup> NCT02716116	<b>Non-small cell lung carcinoma</b> (Approved on 2021/09/15)
	<b>EGFR ex20ins</b> Mobocertinib [ORR(%): 28.0, DOR(M): 17.5]

# ACT Onco<sup>®</sup> + Report

## Necitumumab (PORTRAZZA)

Necitumumab is a recombinant human IgG1 monoclonal antibody against the human epidermal growth factor receptor (EGFR) and blocks the binding of EGFR to its ligands. Necitumumab is developed and marketed by Eli Lilly under the trade name PORTRAZZA.

### - FDA Approval Summary of Necitumumab (PORTRAZZA)

<b>SQUIRE<sup>[98]</sup></b> NCT00981058	<b>Lung squamous cell carcinoma</b> (Approved on 2015/11/14)
	-
	Gemcitabine + cisplatin vs. Placebo [OS(M): 11.5 vs. 9.9]

## Nivolumab (OPDIVO)

Nivolumab is a programmed death receptor-1 (PD-1)-blocking antibody. Nivolumab is developed and marketed by Bristol-Myers Squibb under the trade name OPDIVO.

### - FDA Approval Summary of Nivolumab (OPDIVO)

<b>CHECKMATE-648</b> NCT03143153	<b>Esophagus squamous cell carcinoma</b> (Approved on 2022/05/27)
	-
	Nivolumab and ipilimumab vs. Chemotherapy [OS(M): 12.8 vs. 10.7]
<b>CHECKMATE-648</b> NCT03143153	<b>Esophagus squamous cell carcinoma</b> (Approved on 2022/05/27)
	-
	Nivolumab, fluorouracil, and cisplatin vs. Chemotherapy [OS(M): 13.2 vs. 10.7]
<b>CHECKMATE-816</b> NCT02998528	<b>Non-small cell lung cancer (nscic)</b> (Approved on 2022/03/04)
	-
	Nivolumab plus platinum-doublet chemotherapy vs. Platinum-chemotherapy [EFS(M): 31.6 vs. 20.8]
<b>CHECKMATE-274</b> NCT02632409	<b>Bladder urothelial carcinoma</b> (Approved on 2021/08/19)
	-
	Nivolumab [DFS (all randomized)(M): 20.8 vs. 10.8, DFS (PD-L1 ≥ 1%)(M): NR vs. 8.4]
<b>CHECKMATE-577</b> NCT02743494	<b>Gastroesophageal junction adenocarcinoma</b> (Approved on 2021/05/20)
	-
	Nivolumab vs. Placebo every 4 weeks beginning at week 17 for up to one year of treatment [DFS(M): 22.4 vs. 11]
<b>CHECKMATE-649</b> NCT02872116	<b>Gastroesophageal junction adenocarcinoma, Gastric adenocarcinoma</b> (Approved on 2021/04/16)
	-
	Nivolumab + chemotherapy (xelox or folfox) vs. Chemotherapy (xelox or folfox) [PFS(M): 7.7 vs. 6, OS(M): 14.4 vs. 11.1]
<b>CHECKMATE-9ER</b> NCT03141177	<b>Renal cell carcinoma</b> (Approved on 2021/01/22)
	-
	Nivolumab + cabozantinib vs. Sunitinib [ORR(%): 55.7 vs. 27.1, PFS(M): 16.6 vs. 8.3, OS(M): NR vs. NR]
<b>CHECKMATE-743</b> NCT02899299	<b>Pleural mesothelioma</b> (Approved on 2020/10/02)
	-
	Nivolumab + ipilimumab vs. Pemetrexed + cisplatin [OS(M): 18.1 vs. 14.1]

# ACT Onco<sup>®</sup> + Report

<b>CHECKMATE-9LA</b> NCT03215706	<b>Non-small cell lung carcinoma</b> (Approved on 2020/05/26)
	- Nivolumab + ipilimumab + platinum-doublet chemotherapy vs. Platinum-doublet chemotherapy [OS(M): 14.1 vs. 10.7]
<b>CHECKMATE-227</b> NCT02477826	<b>Non-small cell lung carcinoma</b> (Approved on 2020/05/15)
	<b>PD-L1</b> Nivolumab + ipilimumab vs. Platinum-doublet chemotherapy [OS(M): 17.1 vs. 14.9]
<b>CheckMate 040</b> NCT01658878	<b>Hepatocellular carcinoma</b> (Approved on 2020/03/10)
	- Nivolumab + ipilimumab [ORR(%): 33.0]
<b>CheckMate 142</b> NCT02060188	<b>Colorectal cancer</b> (Approved on 2017/07/31)
	<b>MSI-H or dMMR</b> Nivolumab [ORR(%): 32.0]
<b>CheckMate 141</b> <sup>[99]</sup> NCT02105636	<b>Squamous cell carcinoma of the head and neck cancer</b> (Approved on 2016/11/10)
	- Nivolumab vs. Investigator's choice of cetuximab, methotrexate or docetaxel [OS(M): 7.5 vs. 5.1]
<b>CheckMate 205</b> <sup>[100]</sup> NCT02181738	<b>Hodgkin's lymphoma</b> (Approved on 2016/05/17)
	- Nivolumab [ORR(%): 66.0]
<b>CheckMate 039</b> <sup>[101]</sup> NCT01592370	<b>Hodgkin's lymphoma</b> (Approved on 2016/05/17)
	- Nivolumab [ORR(%): 66.0]
<b>CheckMate 067</b> <sup>[102]</sup> NCT01844505	<b>Melanoma</b> (Approved on 2016/01/23)
	- Ipilimumab vs. Placebo [PFS(M): 11.5 vs. 2.9]
<b>CheckMate 066</b> <sup>[103]</sup> NCT01721772	<b>Melanoma</b> (Approved on 2015/11/24)
	<b>BRAF V600 wild-type</b> Nivolumab vs. Dacarbazine [OS(M): Not Reached vs. 10.8]
<b>CheckMate 025</b> <sup>[104]</sup> NCT01668784	<b>Renal cell carcinoma</b> (Approved on 2015/11/23)
	- Nivolumab vs. Everolimus [OS(M): 25 vs. 19.6]
<b>CheckMate 057</b> <sup>[105]</sup> NCT01673867	<b>Non-small cell lung carcinoma</b> (Approved on 2015/10/09)
	- Nivolumab vs. Docetaxel [OS(M): 12.2 vs. 9.4]
<b>CheckMate 017</b> <sup>[106]</sup> NCT01642004	<b>Non-small cell lung carcinoma</b> (Approved on 2015/03/04)
	- Nivolumab vs. Docetaxel [OS(M): 9.2 vs. 6]
<b>CheckMate 037</b> <sup>[107]</sup> NCT01721746	<b>Melanoma</b> (Approved on 2014/12/22)
	- Nivolumab vs. Dacarbazine or carboplatin + paclitaxel [ORR(%): 32.0]



# ACT Onco<sup>®</sup> + Report

## Osimertinib (TAGRISSO)

Osimertinib is a third-generation tyrosine kinase inhibitor (TKI) for patients with tumors harboring EGFR T790M mutation. Osimertinib is developed and marketed by AstraZeneca under the trade name TAGRISSO.

### - FDA Approval Summary of Osimertinib (TAGRISSO)

<b>ADAURA</b> NCT02511106	<b>Non-small cell lung carcinoma</b> (Approved on 2020/12/18)
	<b>EGFR ex19del or L858R</b>
	Osimertinib vs. Placebo + adjuvant chemotherapy [DFS(M): NR vs. 19.6]
<b>FLAURA<sup>[32]</sup></b> NCT02296125	<b>Non-small cell lung carcinoma</b> (Approved on 2018/04/18)
	<b>EGFR ex19del or L858R</b>
	Osimertinib vs. Gefitinib or erlotinib [PFS(M): 18.9 vs. 10.2]
<b>AURA3<sup>[108]</sup></b> NCT02151981	<b>Non-small cell lung carcinoma</b> (Approved on 2017/03/30)
	<b>EGFR T790M</b>
	Osimertinib vs. Chemotherapy [PFS(M): 10.1 vs. 4.4]
<b>AURA<sup>[31]</sup></b> NCT01802632	<b>Non-small cell lung carcinoma</b> (Approved on 2015/11/13)
	<b>EGFR T790M</b>
	Osimertinib [ORR(%): 59.0]

## Panitumumab (VECTIBIX)

Panitumumab is a fully human monoclonal antibody against the human epidermal growth factor receptor (EGFR) and binds to the extracellular domain to prevent its activation. Panitumumab is developed by Abgenix and Amgen, and marketed by the latter under the trade name VECTIBIX.

### - FDA Approval Summary of Panitumumab (VECTIBIX)

<b>Study 20050203<sup>[109]</sup></b> NCT01412957	<b>Colorectal cancer</b> (Approved on 2017/06/29)
	<b>KRAS Wild-type</b>
	Panitumumab + bsc vs. Bsc [OS(M): 10 vs. 6.9]
<b>PRIME<sup>[110]</sup></b> NCT00364013	<b>Colorectal cancer</b> (Approved on 2014/05/23)
	<b>KRAS Wild-type</b>
	Panitumumab + folfox vs. Folfox [PFS(M): 9.6 vs. 8]
<b>ASPECCT<sup>[111]</sup></b> NCT01001377	<b>Colorectal cancer</b> (Approved on 2014/05/23)
	<b>KRAS Wild-type</b>
	Panitumumab vs. Cetuximab [OS(M): 10.4 vs. 10]
<b>Study 20080763<sup>[112]</sup></b> NCT00113763	<b>Colorectal cancer</b> (Approved on 2006/09/27)
	<b>KRAS Wild-type</b>
	Panitumumab + bsc vs. Bsc [PFS(M): 3.2 vs. 2]

# ACT Onco<sup>®</sup> + Report

## Pembrolizumab (KEYTRUDA)

Pembrolizumab is a programmed death receptor-1 (PD-1)-blocking antibody. Pembrolizumab is developed and marketed by Merck under the trade name KEYTRUDA.

## - FDA Approval Summary of Pembrolizumab (KEYTRUDA)

KEYNOTE-158 NCT02628067	<b>Endometrial carcinoma</b> (Approved on 2022/03/21)
	<b>MSI-H or dMMR</b> Pembrolizumab [ORR(%): 46.0, DoR(M): NR]
KEYNOTE-716 NCT03553836	<b>Melanoma</b> (Approved on 2021/12/03)
	- Pembrolizumab [RFS(M): Not reached vs. Not reached]
KEYNOTE-564 NCT03142334	<b>Renal cell carcinoma</b> (Approved on 2021/11/17)
	- Pembrolizumab vs. Placebo [DFS(M): NR vs. NR, OS(M): NR vs. NR]
KEYNOTE-826 NCT03635567	<b>Cervical cancer</b> (Approved on 2021/10/13)
	<b>PD-L1</b> Pembrolizumab + paclitaxel + cisplatin with or without bevacizumab vs. Placebo + paclitaxel + cisplatin with or without bevacizumab [OS (PD-L1, CPS ≥1)(M): Not reached vs. 16.3, PFS(M): 10.4 vs. 8.2]
CLEAR (Study 307/KEYNOTE-581) NCT02811861	<b>renal cell carcinoma</b> (Approved on 2021/08/11)
	- Pembrolizumab + lenvatinib vs. Sunitinib [PFS(M): 23.9 vs. 9.2, OS(M): NR vs. NR, ORR(%): 71.0 vs. 36.0]
KEYNOTE-522 NCT03036488	<b>Triple-receptor negative breast cancer</b> (Approved on 2021/07/26)
	- Pembrolizumab + chemotherapy as neoadjuvant treatment vs. Placebo in combination with chemotherapy [pCR(%): 63.0 vs. 56.0, EFS(): 123 vs. 93]
KEYNOTE-775 (Study 309) NCT03517449	<b>Endometrial carcinoma</b> (Approved on 2021/07/22)
	<b>MSS/pMMR</b> Pembrolizumab + lenvatinib vs. Investigator's choice of doxorubicin or paclitaxel [PFS(M): 6.6 vs. 3.8, OS(M): 17.4 vs. 12]
KEYNOTE-811 NCT03615326	<b>Gastroesophageal junction adenocarcinoma</b> (Approved on 2021/05/05)
	<b>HER2+</b> Pembrolizumab 200 mg every 3 weeks, in combination with trastuzumab and either fluorouracil plus cisplatin or capecitabine plus oxaliplatin vs. Placebo every 3 weeks, in combination with trastuzumab and either fluorouracil plus cisplatin or capecitabine plus oxaliplatin [ORR(%): 74.0 vs. 52.0, DOR(M): 10.6 vs. 9.5]
KEYNOTE-590 NCT03189719	<b>Esophageal cancer, Gastroesophageal junction adenocarcinoma</b> (Approved on 2021/03/22)
	- Pembrolizumab in combination with cisplatin and fluorouracil vs. Placebo with cisplatin and fluorouracil [PFS(M): 6.3 vs. 5.8, OS(M): 12.4 vs. 9.8]
KEYNOTE-355 NCT02819518	<b>Triple-receptor negative breast cancer</b> (Approved on 2020/11/13)
	<b>PD-L1</b> Pembrolizumab + paclitaxel protein-bound, or paclitaxel, or gemcitabine plus carboplatin vs. Placebo + paclitaxel protein-bound, or paclitaxel, or gemcitabine plus carboplatin [PFS(M): 9.7 vs. 5.6]

# ACT Onco® + Report

<b>KEYNOTE-204</b> NCT02684292	<b>Hodgkin's lymphoma</b> (Approved on 2020/10/14)
	-
<b>KEYNOTE-158</b> NCT02628067	<b>Cancer</b> (Approved on 2020/06/17)
	<b>TMB-H</b>
<b>KEYNOTE-146</b> NCT02501096	<b>Endometrial carcinoma</b> (Approved on 2019/09/17)
	<b>MSS/pMMR</b>
<b>KEYNOTE-426</b> <sup>[113]</sup> NCT02853331	<b>Renal cell carcinoma</b> (Approved on 2019/04/19)
	-
<b>KEYNOTE-017</b> <sup>[114]</sup> NCT02267603	<b>Merkel cell carcinoma</b> (Approved on 2018/12/19)
	-
<b>KEYNOTE-224</b> <sup>[115]</sup> NCT02702414	<b>Hepatocellular carcinoma</b> (Approved on 2018/11/09)
	-
<b>KEYNOTE-407</b> <sup>[116]</sup> NCT02775435	<b>Squamous non-small cell lung carcinoma</b> (Approved on 2018/10/30)
	-
<b>KEYNOTE-189</b> <sup>[116]</sup> NCT02578680	<b>Nonsquamous non-small cell lung carcinoma</b> (Approved on 2018/08/20)
	-
<b>KEYNOTE-158</b> NCT02628067	<b>Cervical cancer</b> (Approved on 2018/06/13)
	-
<b>KEYNOTE-170</b> NCT02576990	<b>Mediastinal large b-cell lymphoma</b> (Approved on 2018/06/13)
	-
<b>KEYNOTE-059</b> NCT02335411	<b>Gastric adenocarcinoma, Gastroesophageal junction adenocarcinoma</b> (Approved on 2017/09/22)
	-
<b>KEYNOTE-164</b> NCT02460198	<b>Cancer</b> (Approved on 2017/05/23)
	<b>MSI-H or dMMR</b>
<b>KEYNOTE-016</b> <sup>[6]</sup> NCT01876511	<b>Cancer</b> (Approved on 2017/05/23)
	<b>MSI-H or dMMR</b>
<b>KEYNOTE-158</b> NCT02628067	<b>Cancer</b> (Approved on 2017/05/23)
	<b>MSI-H or dMMR</b>
<b>KEYNOTE-028</b> <sup>[117][118]</sup> NCT02054806	<b>Cancer</b> (Approved on 2017/05/23)
	<b>MSI-H or dMMR</b>

# ACT Onco<sup>®</sup> + Report

<b>KEYNOTE-012</b> <sup>[119][120][121][122]</sup> NCT01848834	<b>Cancer</b> (Approved on 2017/05/23)
	<b>MSI-H or dMMR</b>
	Pembrolizumab [ORR(%): 39.6]
<b>KEYNOTE-045</b> <sup>[123]</sup> NCT02256436	<b>Urinary bladder urothelial carcinoma</b> (Approved on 2017/05/18)
	-
	Pembrolizumab vs. Chemotherapy [ORR(%): 21.0 vs. 11.0]
<b>KEYNOTE-052</b> NCT02335424	<b>Urinary bladder urothelial carcinoma</b> (Approved on 2017/05/18)
	-
	Pembrolizumab [ORR(%): 29.0]
<b>KEYNOTE-087</b> <sup>[124]</sup> NCT02453594	<b>Hodgkin's lymphoma</b> (Approved on 2017/03/14)
	-
	Pembrolizumab [ORR(%): 69.0]
<b>KEYNOTE-024</b> <sup>[125]</sup> NCT02142738	<b>Non-small cell lung carcinoma</b> (Approved on 2016/10/24)
	<b>PD-L1</b>
	Pembrolizumab vs. Chemotherapy [PFS(M): 10.3 vs. 6]
<b>KEYNOTE-012</b> <sup>[120]</sup> NCT01848834	<b>Head and neck squamous cell carcinoma</b> (Approved on 2016/08/05)
	-
	Pembrolizumab [ORR(%): 16.0]
<b>KEYNOTE-006</b> <sup>[126]</sup> NCT01866319	<b>Melanoma</b> (Approved on 2015/12/18)
	-
	Pembrolizumab vs. Ipilimumab (3mg/kg every 3 weeks) [OS(M): NR vs. 16]
<b>KEYNOTE-010</b> <sup>[127]</sup> NCT01905657	<b>Non-small cell lung carcinoma</b> (Approved on 2015/10/02)
	<b>PD-L1</b>
	Pembrolizumab [OS(M): 10.4 vs. 8.5]
<b>KEYNOTE-002</b> <sup>[128]</sup> NCT01704287	<b>Melanoma</b> (Approved on 2014/09/24)
	-
	Pembrolizumab vs. Chemotherapy [PFS(M): 2.9 vs. 2.7]

D=day; W=week; M=month

# ACT Onco<sup>®</sup> + Report

## ONGOING CLINICAL TRIALS

Trials were searched by applying filters: study status, patient's diagnosis, intervention, location and/or biomarker(s). Please visit <https://clinicaltrials.gov> to search and view for a complete list of open available and updated matched trials.

No trial has been found.

# ACT Onco<sup>®</sup> + Report


## SUPPLEMENTARY INFORMATION OF TESTING RESULTS DETAILED INFORMATION OF VARIANTS WITH CLINICAL RELEVANCE

### - Single Nucleotide and Small InDel Variants

Gene	Amino Acid Change	Exon	cDNA Change	Accession Number	COSMIC ID	Allele Frequency	Coverage
EGFR	L747_P753delinsS (Exon 19 deletion)	19	c.2240_2257del	NM_005228	COSM12370	68.2%	3867
KMT2D	Q2458*	31	c.7372C>T	NM_003482	COSM4507076	18.1%	282

### - Copy Number Alterations

Observed copy number (CN) for each evaluated position is shown on the y-axis. Regions referred to as amplification or deletion are shown in color. Regions without significant changes are represented in gray.





# ACT Onco<sup>®</sup> + Report

## OTHER DETECTED VARIANTS

Gene	Amino Acid Change	Exon	cDNA Change	Accession Number	COSMIC ID	Allele Frequency	Coverage
ADAMTS16	P966L	19	c.2897C>T	NM_139056	-	5.8%	445
ADAMTS18	R100P	3	c.299G>C	NM_199355	-	40.8%	524
ARID1A	G2283S	20	c.6847G>A	NM_006015	-	57.3%	1199
ARID1B	A1993V	19	c.5978C>T	NM_017519	-	45.4%	1404
ATR	H2277Y	40	c.6829C>T	NM_001184	-	30.0%	679
BRCA2	V2109I	11	c.6325G>A	NM_000059	-	33.5%	481
CCNE1	D218G	8	c.653A>G	NM_001238	-	67.2%	527
DPYD	G593E	14	c.1778G>A	NM_000110	-	41.8%	832
EGFR	A21T	1	c.61G>A	NM_005228	-	83.6%	1606
EP300	H2324N	31	c.6970C>A	NM_001429	-	67.7%	663
ERBB4	T926M	23	c.2777C>T	NM_005235	COSM5966952	59.7%	828
FANCF	H51Y	1	c.151C>T	NM_022725	-	50.4%	811
FANCF	S345I	1	c.1034G>T	NM_022725	-	47.9%	1169
FAT1	E1292K	5	c.3874G>A	NM_005245	-	36.8%	544
HGF	C211F	6	c.632G>T	NM_000601	-	22.8%	918
IGF1R	V1277D	21	c.3830T>A	NM_000875	-	11.0%	565
INSR	M1319I	22	c.3957G>A	NM_000208	-	31.7%	1841
MUC6	S1700C	31	c.5099C>G	NM_005961	COSM5048600	6.7%	834
NKX2-1	G322S	2	c.964G>A	NM_003317	-	35.1%	97
NSD1	T2382I	23	c.7145C>T	NM_022455	-	53.4%	328
PRKN	R256H	7	c.767G>A	NM_004562	COSM7732066	12.8%	1046
RAD51D	V66M	3	c.196G>A	NM_002878	-	67.7%	919
RUNX1	Q370R	6	c.1109A>G	NM_001001890	COSM26028	41.4%	440
RUNX1T1	A471V	11	c.1412C>T	NM_175634	COSM33136	29.4%	503
SMARCA4	Splice region	-	c.2617-4G>A	NM_001128844	-	40.6%	416
TNFAIP3	Splice region	-	c.295+8T>C	NM_006290	-	55.4%	1035
TNFAIP3	I207L	4	c.619A>C	NM_006290	-	50.9%	454
TNFAIP3	T108A	3	c.322A>G	NM_006290	COSM1581845	51.9%	2087
TP53	Splice region	6	c.672G>A	NM_000546	COSM44754	29.5%	227
USH2A	E4264K	63	c.12790G>A	NM_206933	COSM213160	54.6%	1184


Note:

- This table enlists variants detected by the panel other than those with clinical relevance (reported in Testing Result section). The clinical impact of a genetic variant is determined according to ACT Genomics in-house clinical knowledge database. A negative result does not necessarily indicate absence of biological effect on the tumor. Some variants listed here may possibly have preclinical data or may show potential clinical relevance in the future.

# ACT Onco<sup>®</sup> + Report

## TEST DETAILS

### SPECIMEN RECEIVED AND PATHOLOGY REVIEW



- Collection date: Sep 22, 2022
- Facility retrieved: 臺北榮總
- H&E-stained section No.: S11137673D
- Collection site: Pleura
- Examined by: Dr. Chien-Ta Chiang
  1. The percentage of viable tumor cells in total cells in the whole slide (%): 10%
  2. The percentage of viable tumor cells in total cells in the encircled areas in the whole slide (%): 45%
  3. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the whole slide (%): 80%
  4. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the encircled areas in the whole slide (%): 40%
  5. Additional comment: NA
- Manual macrodissection: Performed on the highlighted region
- The outline highlights the area of malignant neoplasm annotated by a pathologist.

## RUN QC

- Panel: ACTOnco<sup>®</sup>+

### DNA test

- Mean Depth: 846x
- Target Base Coverage at 100x: 94%

### RNA test

- Average unique RNA Start Sites per control GSP2: 134

# ACT Onco<sup>®</sup> + Report

## LIMITATIONS

1. This test does not provide information of variant causality and does not detect variants in non-coding regions that could affect gene expression. This report does not report polymorphisms and we do not classify whether a mutation is germline or somatic. Variants identified by this assay were not subject to validation by Sanger or other technologies.
2. The possibility cannot be excluded that certain pathogenic variants detected by other sequencing tools may not be reported in the test because of technical limitation of bioinformatics algorithm or the NGS sequencing platform, e.g. low coverage.
3. This test has been designed to detect fusions in 13 genes sequenced. Therefore, fusion in genes not covered by this test would not be reported. For novel fusions detected in this test, Sanger sequencing confirmation is recommended if residue specimen is available.

## NEXT-GENERATION SEQUENCING (NGS) METHODS

### DNA test

Extracted genomic DNA was amplified using primers targeting coding exons of analyzed genes and subjected to library construction. Barcoded libraries were subsequently conjugated with sequencing beads by emulsion PCR and enriched using Ion Chef system. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific).

Raw reads generated by the sequencer were mapped to the hg19 reference genome using the Ion Torrent Suite. Coverage depth was calculated using Torrent Coverage Analysis plug-in. Single nucleotide variants (SNVs) and short insertions/deletions (InDels) were identified using the Torrent Variant Caller plug-in. VEP (Variant Effect Predictor) was used to annotate every variant using databases from Clinvar, COSMIC and Genome Aggregation database. Variants with coverage  $\geq 20$ , allele frequency  $\geq 5\%$  and actionable variants with allele frequency  $\geq 2\%$  were retained. This test provides uniform coverage of the targeted regions, enabling target base coverage at  $100\times \geq 85\%$  with a mean coverage  $\geq 500\times$ .

Variants reported in Genome Aggregation database with  $> 1\%$  minor allele frequency (MAF) were considered as polymorphisms. ACT Genomics in-house database was used to determine technical errors. Clinically actionable and biologically significant variants were determined based on the published medical literature.

The copy number alterations (CNAs) were predicted as described below:

Amplicons with read counts in the lowest 5th percentile of all detectable amplicons and amplicons with a coefficient of variation  $\geq 0.3$  were removed. The remaining amplicons were normalized to correct the pool design bias. ONCOCNV (an established method for calculating copy number aberrations in amplicon sequencing data by Boeva et al., 2014) was applied for the normalization of total amplicon number, amplicon GC content, amplicon length, and technology-related biases, followed by segmenting the sample with a gene-aware model. The method was used as well for establishing the baseline of copy number variations.

Tumor mutational burden (TMB) was calculated by using the sequenced regions of ACTOnco<sup>®</sup> to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The TMB calculation predicted somatic variants and applied a machine learning model with a cancer hotspot correction. TMB may be reported as "TMB-High", "TMB-Low" or "Cannot Be Determined". TMB-High corresponds to  $\geq 7.5$  mutations per megabase (Muts/Mb); TMB-Low corresponds to  $< 7.5$  Muts/Mb. TMB is reported as "Cannot Be Determined" if the tumor purity of the sample is  $< 30\%$ .

Classification of microsatellite instability (MSI) status is determined by a machine learning prediction algorithm. The change of a number of repeats of different lengths from a pooled microsatellite stable (MSS) baseline in  $> 400$  genomic loci are used as the features for the algorithm. The final output of the results is either microsatellite Stable (MSS) or microsatellite instability high (MSI-H).

### RNA test

Extracted RNA was reverse-transcribed and subjected to library construction. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific). To ensure sequencing quality for fusion variant analysis, the average unique RNA Start Sites (SS) per control Gene Specific Primer 2 (GSP 2) should be  $\geq 10$ .

# ACT Onco<sup>®</sup> + Report

The fusion analysis pipeline aligned sequenced reads to the human reference genome, identified regions that map to noncontiguous regions of the genome, applied filters to exclude probable false-positive events and, annotated previously characterized fusion events according to Quiver Gene Fusion Database, a curated database owned and maintained by ArcherDX. In general, samples with detectable fusions need to meet the following criteria: (1) Number of unique start sites (SS) for the GSP2  $\geq 3$ ; (2) Number of supporting reads spanning the fusion junction  $\geq 5$ ; (3) Percentage of supporting reads spanning the fusion junction  $\geq 10\%$ ; (4) Fusions annotated in Quiver Gene Fusion Database.

## DATABASE USED

- Reference genome: Human genome sequence hg19
- COSMIC v.92
- Genome Aggregation database r2.1.1
- ClinVar (version 20210404)
- ACT Genomics in-house database
- Quiver Gene Fusion Database version 5.1.18

## Variant Analysis:

醫藥資訊研究員  
楊杭哲 博士  
Hang-Che Yang Ph.D.



## Sign Off

解剖病理專科醫師王業翰  
Yeh-Han Wang M.D.  
病解字第 000545 號



# ACT Onco<sup>®</sup> + Report

## GENE LIST SNV & CNV

ABCB1*	ABCC2*	ABCG2*	ABL1	ABL2	ADAMTS1	ADAMTS13	ADAMTS15	ADAMTS16	ADAMTS18	ADAMTS6	ADAMTS9
ADAMTS11	ADGRA2	ADH1C*	AKT1	AKT2	AKT3	ALDH1A1*	ALK	AMER1	APC	AR	ARAF
ARID1A	ARID1B	ARID2	ASXL1	ATM	ATR	ATRX	AURKA	AURKB	AXIN1	AXIN2	AXL
B2M	BAP1	BARD1	BCL10	BCL2*	BCL2L1	BCL2L2*	BCL6	BCL9	BCOR	BIRC2	BIRC3
BLM	BMPR1A	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2*	BTK	BUB1B	CALR
CANX	CARD11	CASP8	CBFB	CBL	CCNA1	CCNA	CCNB1	CCNB2	CCNB3	CCND1	CCND2
CCND3	CCNE1	CCNE2	CCNH	CD19	CD274	CD58	CD70*	CD79A	CD79B	CDC73	CDH1
CDK1	CDK12	CDK2	CDK4	CDK5	CDK6	CDK7	CDK8	CDK9	CDKN1A	CDKN1B	CDKN2A
CDKN2B	CDKN2C	CEBPA*	CHEK1	CHEK2	CIC	CREBBP	CRKL	CRLF2	CSF1R	CTCF	CTLA4
CTNNA1	CTNNB1	CUL3	CYLD	CYP1A1*	CYP2B6*	CYP2C19*	CYP2C8*	CYP2D6	CYP2E1*	CYP3A4*	CYP3A5*
DAXX	DCUN1D1	DDR2	DICER1	DNMT3A	DOT1L	DPYD	DTX1	E2F3	EGFR	EP300	EPCAM
EPHA2	EPHA3	EPHA5	EPHA7	EPHB1	ERBB2	ERBB3	ERBB4	ERCC1	ERCC2	ERCC3	ERCC4
ERCC5	ERG	ESR1	ESR2	ETV1	ETV4	EZH2	FAM46C	FANCA	FANCC	FANCD2	FANCE
FANCF	FANCG	FANCL	FAS	FAT1	FBXW7	FCGR2B	FGF1*	FGF10	FGF14	FGF19*	FGF23
FGF3	FGF4*	FGF6	FGFR1	FGFR2	FGFR3	FGFR4	FH	FLCN	FLT1	FLT3	FLT4
FOXL2*	FOXP1	FRG1	FUBP1	GATA1	GATA2	GATA3	GNA11	GNA13	GNAQ	GNAS	GREM1
GRIN2A	GSK3B	GSTP1*	GSTT1*	HGF	HIF1A	HIST1H1C*	HIST1H1E*	HNF1A	HR	HRAS*	HSP90AA1
HSP90AB1	HSPA4	HSPA5	IDH1	IDH2	IFNL3*	IGF1	IGF1R	IGF2	IKBKB	IKBKE	IKZF1
IL6	IL7R	INPP4B	INSR	IRF4	IRS1	IRS2*	JAK1	JAK2	JAK3	JUN*	KAT6A
KDM5A	KDM5C	KDM6A	KDR	KEAP1	KIT	KMT2A	KMT2C	KMT2D	KRAS	LCK	LIG1
LIG3	LMO1	LRP1B	LYN	MALT1	MAP2K1	MAP2K2	MAP2K4	MAP3K1	MAP3K7	MAPK1	MAPK3
MAX	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1	MET	MITF	MLH1	MPL	MRE11
MSH2	MSH6	MTHFR*	MTOR	MUC16	MUC4	MUC6	MUTYH	MYC	MYCL	MYCN	MYD88
NAT2*	NBN	NEFH	NF1	NF2	NFE2L2	NFKB1	NFKBIA	NKX2-1*	NOTCH1	NOTCH2	NOTCH3
NOTCH4	NPM1	NQO1*	NRAS	NSD1	NTRK1	NTRK2	NTRK3	PAK3	PALB2	PARP1	PAX5
PAX8	PBRM1	PDCD1	PDCD1LG2	PDGFRA	PDGFRB	PDIA3	PGF	PHOX2B*	PIK3C2B	PIK3C2G	PIK3C3
PIK3CA	PIK3CB	PIK3CD	PIK3CG	PIK3R1	PIK3R2	PIK3R3	PIM1	PMS1	PMS2	POLB	POLD1
POLE	PPARG	PPP2R1A	PRDM1	PRKAR1A	PRKCA	PRKCB	PRKCG	PRKCI	PRKCQ	PRKDC	PRKN
PSMB8	PSMB9	PSME1	PSME2	PSME3	PTCH1	PTEN	PTGS2	PTPN11	PTPRD	PTPRT	RAC1
RAD50	RAD51	RAD51B	RAD51C	RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	RECQL4
REL	RET	RHOA	RICTOR	RNF43	ROS1	RPPH1	RPTOR	RUNX1	RUNX1T1	RXRA	SDHA
SDHB	SDHC	SDHD	SERPINB3	SERPINB4	SETD2	SF3B1	SGK1	SH2D1A*	SLC19A1*	SLC22A2*	SLC01B1*
SLC01B3*	SMAD2	SMAD3	SMAD4	SMARCA4	SMARCB1	SMO	SOCS1*	SOX2*	SOX9	SPEN	SPOP
SRC	STAG2	STAT3	STK11	SUFU	SYK	SYNE1	TAF1	TAP1	TAP2	TAPBP	TBX3
TEK	TERT	TET1	TET2	TGFBR2	TMSB4X*	TNF	TNFAIP3	TNFRSF14	TNFSF11	TOP1	TP53
TPMT*	TSC1	TSC2	TSHR	TYMS	U2AF1	UBE2A*	UBE2K	UBR5	UGT1A1*	USH2A	VDR*
VEGFA	VEGFB	VHL	WT1	XIAP	XPO1	XRCC2	ZNF217				

\*Analysis of copy number alterations NOT available.

## FUSION

ALK	BRAF	EGFR	FGFR1	FGFR2	FGFR3	MET	NRG1	NTRK1	NTRK2	NTRK3	RET	ROS1
-----	------	------	-------	-------	-------	-----	------	-------	-------	-------	-----	------


# ACT Onco<sup>®</sup> + Report

## APPENDIX

### POSSIBLE THERAPEUTIC IMPLICATIONS FOR HETEROZYGOUS DELETION

Not Applicable.

### SIGNALING PATHWAYS AND MOLECULAR-TARGETED AGENTS



1: Gefitinib, Afatinib, Erlotinib, Osimertinib, Dacomitinib, Mobocertinib; 2: Cetuximab, Panitumumab, Necitumumab; 3:

Afatinib



# ACT Onco<sup>®</sup> + Report

## DISCLAIMER

### 法律聲明

本檢驗報告僅提供專業醫療參考，結果需經專業醫師解釋及判讀。基因突變資訊非必具備藥物或治療有效性指標，反之亦然。本檢驗報告提供之用藥指引不聲明或保證其臨床有效性，反之亦然。本基因檢測方法係由本公司研究開發，已經過有效性測試。

本檢驗報告非經本公司許可，不得私自變造、塗改，或以任何方式作為廣告及其他宣傳之用途。

本公司於提供檢驗報告後，即已完成本次契約義務，後續之報告解釋、判讀及用藥、治療，應自行尋求相關專業醫師協助，若需將報告移件其他醫師，本人應取得該醫師同意並填寫移件申請書，主動告知行動基因，行動基因僅能配合該醫師意願與時間提供醫師解說。

### 醫療決策需由醫師決定

任何治療與用藥需經由醫師在考慮病患所有健康狀況相關資訊包含健檢、其他檢測報告和病患意願後，依照該地區醫療照護標準由醫師獨立判斷。醫師不應僅依據單一報告結果(例如本檢測或本報告書內容)做決策。

### 基因突變與用藥資訊並非依照有效性排序

本報告中列出之生物標記變異與藥物資訊並非依照潛在治療有效性排序。

### 證據等級

藥物潛在臨床效益(或缺乏潛在臨床效益)的實證證據是依據至少一篇臨床療效個案報告或臨床前試驗做為評估。本公司盡力提供適時及準確之資料，但由於醫學科技之發展日新月異，本公司不就本報告提供的資料是否為準確、適宜或最新作保證。

### 責任

本檢驗報告僅提供專業醫療參考，本公司及其員工不對任何由使用本報告之內容引起的直接、間接、特殊、連帶或衍生的損失或損害承擔責任。

# ACT Onco<sup>®</sup> + Report

## REFERENCE

1. PMID: 26559592; 2015, N Engl J Med;373(20):1984  
Genetic Basis for Clinical Response to CTLA-4 Blockade in Melanoma.
2. PMID: 26359337; 2015, Science;350(6257):207-211  
Genomic correlates of response to CTLA-4 blockade in metastatic melanoma.
3. PMID: 28251903; 2017, Sci Transl Med;9(379):  
Integrated molecular analysis of tumor biopsies on sequential CTLA-4 and PD-1 blockade reveals markers of response and resistance.
4. PMID: 26997480; 2016, Cell;165(1):35-44  
Genomic and Transcriptomic Features of Response to Anti-PD-1 Therapy in Metastatic Melanoma.
5. PMID: 25765070; 2015, Science;348(6230):124-8  
Cancer immunology. Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer.
6. PMID: 26028255; 2015, N Engl J Med;372(26):2509-20  
PD-1 Blockade in Tumors with Mismatch-Repair Deficiency.
7. PMID: 29863979; 2018, N Engl J Med;379(4):341-351  
PD-1 Blockade with Cemiplimab in Advanced Cutaneous Squamous-Cell Carcinoma.
8. PMID: 26952546; 2016, Lancet;387(10031):1909-20  
Atezolizumab in patients with locally advanced and metastatic urothelial carcinoma who have progressed following treatment with platinum-based chemotherapy: a single-arm, multicentre, phase 2 trial.
9. PMID: 27009843; 2016, Oncotarget;7(16):22857-64  
Analysis of ultra-deep targeted sequencing reveals mutation burden is associated with gender and clinical outcome in lung adenocarcinoma.
10. PMID: 24839032; 2014, Breast Cancer Res Treat;146(1):211-20  
Somatic mutation load of estrogen receptor-positive breast tumors predicts overall survival: an analysis of genome sequence data.
11. PMID: 18045542; 2007, Cell;131(5):1018  
SnapShot: EGFR signaling pathway.
12. PMID: 10880430; 2000, EMBO J;19(13):3159-67  
The ErbB signaling network: receptor heterodimerization in development and cancer.
13. PMID: 15329413; 2004, Proc Natl Acad Sci U S A;101(36):13306-11  
EGF receptor gene mutations are common in lung cancers from "never smokers" and are associated with sensitivity of tumors to gefitinib and erlotinib.
14. PMID: 11426640; 2000, Oncogene;19(56):6550-65  
The EGF receptor family as targets for cancer therapy.
15. PMID: 22263017; 2010, J Thorac Dis;2(1):48-51  
Epidermal growth factor receptor (EGFR) in lung cancer: an overview and update.
16. PMID: 28351930; 2017, Clin Cancer Res;23(15):4242-4250  
Hyperprogressors after Immunotherapy: Analysis of Genomic Alterations Associated with Accelerated Growth Rate.
17. PMID: 18508816; 2008, Eur Respir J;32(4):924-30  
Frequent epidermal growth factor receptor gene mutations in malignant pleural effusion of lung adenocarcinoma.
18. PMID: 15118073; 2004, N Engl J Med;350(21):2129-39  
Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib.
19. PMID: 25179728; 2015, Biomed J;38(3):221-8

# ACT Onco<sup>®</sup> + Report

Early radiographic response to epidermal growth factor receptor-tyrosine kinase inhibitor in non-small cell lung cancer patients with epidermal growth factor receptor mutations: A prospective study.

20. PMID: 33728415; 2021, JTO Clin Res Rep;2(3):  
Preclinical characterization of mobocertinib highlights the putative therapeutic window of this novel EGFR inhibitor to EGFR exon 20 insertion mutations.
21. PMID: 29141884; 2017, Sci Transl Med;9(416):  
A method of high-throughput functional evaluation of EGFR gene variants of unknown significance in cancer.
22. PMID: 28958502; 2017, Lancet Oncol;18(11):1454-1466  
Dacomitinib versus gefitinib as first-line treatment for patients with EGFR-mutation-positive non-small-cell lung cancer (ARCHER 1050): a randomised, open-label, phase 3 trial.
23. PMID: 24868098; 2014, Oncologist;19(7):774-9  
U.S. Food and Drug Administration approval summary: Erlotinib for the first-line treatment of metastatic non-small cell lung cancer with epidermal growth factor receptor exon 19 deletions or exon 21 (L858R) substitution mutations.
24. PMID: 23982599; 2013, Drugs;73(13):1503-15  
Afatinib: first global approval.
25. PMID: 27912760; 2016, J Biomed Sci;23(1):86  
Update on recent preclinical and clinical studies of T790M mutant-specific irreversible epidermal growth factor receptor tyrosine kinase inhibitors.
26. PMID: 15728811; 2005, N Engl J Med;352(8):786-92  
EGFR mutation and resistance of non-small-cell lung cancer to gefitinib.
27. PMID: 26862733; 2016, Oncotarget;7(11):12404-13  
The mechanism of acquired resistance to irreversible EGFR tyrosine kinase inhibitor-afatinib in lung adenocarcinoma patients.
28. PMID: 29410323; 2018, J Thorac Oncol;13(5):727-731  
EGFR T790M and C797S Mutations as Mechanisms of Acquired Resistance to Dacomitinib.
29. PMID: 24893891; 2014, Cancer Discov;4(9):1046-61  
AZD9291, an irreversible EGFR TKI, overcomes T790M-mediated resistance to EGFR inhibitors in lung cancer.
30. PMID: 27071706; 2016, J Hematol Oncol;9():34  
Third-generation inhibitors targeting EGFR T790M mutation in advanced non-small cell lung cancer.
31. PMID: 25923549; 2015, N Engl J Med;372(18):1689-99  
AZD9291 in EGFR inhibitor-resistant non-small-cell lung cancer.
32. PMID: 29151359; 2018, N Engl J Med;378(2):113-125  
Osimertinib in Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer.
33. PMID: 17664472; 2007, J Clin Oncol;25(22):3238-45  
Epidermal growth factor receptor gene copy number and clinical outcome of metastatic colorectal cancer treated with panitumumab.
34. PMID: 22152101; 2011, BMC Cancer;11():509  
Biomarker analysis of cetuximab plus oxaliplatin/leucovorin/5-fluorouracil in first-line metastatic gastric and oesophago-gastric junction cancer: results from a phase II trial of the Arbeitsgemeinschaft Internistische Onkologie (AIO).
35. PMID: 24141978; 2013, Sci Rep;3():2992  
A subset of gastric cancers with EGFR amplification and overexpression respond to cetuximab therapy.
36. PMID: 29158193; 2018, J Thorac Oncol;13(2):228-236  
EGFR Gene Copy Number by FISH May Predict Outcome of Necitumumab in Squamous Lung Carcinomas: Analysis from the SQUIRE Study.
37. PMID: 19102716; 2009, Pharmacogenomics;10(1):59-68

# ACT Onco<sup>®</sup> + Report

EGFR-targeted therapies in lung cancer: predictors of response and toxicity.

38. PMID: 15870435; 2005, J Natl Cancer Inst;97(9):643-55  
Epidermal growth factor receptor gene and protein and gefitinib sensitivity in non-small-cell lung cancer.
39. PMID: 15998906; 2005, J Clin Oncol;23(28):6838-45  
Increased epidermal growth factor receptor gene copy number detected by fluorescence in situ hybridization associates with increased sensitivity to gefitinib in patients with bronchioloalveolar carcinoma subtypes: a Southwest Oncology Group Study.
40. PMID: 16204011; 2005, J Clin Oncol;23(31):8081-92  
Epidermal growth factor receptor mutations and gene amplification in non-small-cell lung cancer: molecular analysis of the IDEAL/INTACT gefitinib trials.
41. PMID: 23557218; 2013, J Transl Med;11():90  
High EGFR copy number predicts benefits from tyrosine kinase inhibitor treatment for non-small cell lung cancer patients with wild-type EGFR.
42. PMID: 26141217; 2015, Lung Cancer;89(3):337-42  
Concurrence of EGFR amplification and sensitizing mutations indicate a better survival benefit from EGFR-TKI therapy in lung adenocarcinoma patients.
43. PMID: 28537764; 2017, J Clin Oncol;35(20):2279-2287  
Gefitinib and EGFR Gene Copy Number Aberrations in Esophageal Cancer.
44. PMID: 28057415; 2017, Clin Genitourin Cancer;15(4):e727-e734  
Comprehensive Genomic Analysis of Metastatic Mucinous Urethral Adenocarcinoma Guides Precision Oncology Treatment: Targetable EGFR Amplification Leading to Successful Treatment With Erlotinib.
45. PMID: 30463996; 2019, Cancer Discov;9(2):199-209  
EGFR and MET Amplifications Determine Response to HER2 Inhibition in ERBB2-Amplified Esophagogastric Cancer.
46. PMID: 28575464; 2017, Neuro Oncol;19(11):1522-1531  
Phase II trial of dacomitinib, a pan-human EGFR tyrosine kinase inhibitor, in recurrent glioblastoma patients with EGFR amplification.
47. PMID: 25998713; 2015, Nat Rev Cancer;15(6):334-46  
Hijacked in cancer: the KMT2 (MLL) family of methyltransferases.
48. PMID: 24476821; 2014, Nature;507(7492):315-22  
Comprehensive molecular characterization of urothelial bladder carcinoma.
49. PMID: 21804550; 2011, Nat Genet;43(9):830-7  
Analysis of the coding genome of diffuse large B-cell lymphoma.
50. PMID: 21796119; 2011, Nature;476(7360):298-303  
Frequent mutation of histone-modifying genes in non-Hodgkin lymphoma.
51. PMID: 28609655; 2017, Cancer Cell;31(6):755-770.e6  
MLL2, Not MLL1, Plays a Major Role in Sustaining MLL-Rearranged Acute Myeloid Leukemia.
52. PMID: 26883360; 2016, Genes Dev;30(4):408-20  
Mutation of cancer driver MLL2 results in transcription stress and genome instability.
53. PMID: 29627316; 2018, Clin Lung Cancer;19(4):e489-e501  
KMT2D Mutation Is Associated With Poor Prognosis in Non-Small-Cell Lung Cancer.
54. PMID: 27280393; 2016, Cancer Res;76(16):4861-71  
Reduced Expression of Histone Methyltransferases KMT2C and KMT2D Correlates with Improved Outcome in Pancreatic Ductal Adenocarcinoma.
55. PMID: 29532228; 2018, J Cancer Res Clin Oncol;144(6):1025-1035  
High MLL2 expression predicts poor prognosis and promotes tumor progression by inducing EMT in esophageal squamous cell carcinoma.

# ACT Onco<sup>®</sup> + Report

56. PMID: 29269867; 2018, Oncogene;37(10):1354-1368  
Histone methyltransferase KMT2D sustains prostate carcinogenesis and metastasis via epigenetically activating LIFR and KLF4.
57. PMID: 29483845; 2018, Cancer Cell Int;18():26  
MLL2/KMT2D and MLL3/KMT2C expression correlates with disease progression and response to imatinib mesylate in chronic myeloid leukemia.
58. PMID: 21653699; 2011, J Biol Chem;286(30):26406-17  
Cellular inhibitor of apoptosis protein-1 (cIAP1) can regulate E2F1 transcription factor-mediated control of cyclin transcription.
59. PMID: 29167558; 2017, Sci Rep;7(1):16051  
Recurring Amplification at 11q22.1-q22.2 Locus Plays an Important Role in Lymph Node Metastasis and Radioresistance in OSCC.
60. PMID: 22733500; 2012, Virchows Arch;461(2):123-8  
BIRC2 amplification in squamous cell carcinomas of the uterine cervix.
61. PMID: 27469115; 2016, Cancer Res;76(18):5442-5454  
SMAC Mimetic Birinapant plus Radiation Eradicates Human Head and Neck Cancers with Genomic Amplifications of Cell Death Genes FADD and BIRC2.
62. PMID: 32846130; 2020, Cell Rep;32(8):108073  
BIRC2 Expression Impairs Anti-Cancer Immunity and Immunotherapy Efficacy.
63. PMID: 19029958; 2008, Nat Rev Cancer;8(12):976-90  
Reflecting on 25 years with MYC.
64. PMID: 22464321; 2012, Cell;149(1):22-35  
MYC on the path to cancer.
65. PMID: 10378696; 1999, Oncogene;18(19):3004-16  
MYC oncogenes and human neoplastic disease.
66. PMID: 16934487; 2006, Semin Cancer Biol;16(4):318-30  
The Myc oncoprotein as a therapeutic target for human cancer.
67. PMID: 22113465; 2012, Clin Exp Med;12(4):217-23  
C-myc as a predictive marker for chemotherapy in metastatic breast cancer.
68. PMID: 21741827; 2011, Eur J Cancer;47(12):1779-88  
Association between c-myc amplification and pathological complete response to neoadjuvant chemotherapy in breast cancer.
69. PMID: 15132769; 2004, Cancer Sci;95(5):418-23  
Expression of the c-myc gene as a predictor of chemotherapy response and a prognostic factor in patients with ovarian cancer.
70. PMID: 22430491; 2012, J Exp Med;209(4):679-96  
MYC pathway activation in triple-negative breast cancer is synthetic lethal with CDK inhibition.
71. PMID: 27486754; 2016, Oncotarget;7(35):56864-56875  
Inhibition of cyclin dependent kinase 9 by dinaciclib suppresses cyclin B1 expression and tumor growth in triple negative breast cancer.
72. PMID: 24503701; 2014, PLoS One;9(2):e87456  
Immunohistochemistry for myc predicts survival in colorectal cancer.
73. PMID: 9816266; 1996, Clin Cancer Res;2(6):1049-53  
Overexpression of the c-myc proto-oncogene in colorectal carcinoma is associated with a reduced mortality that is abrogated by point mutation of the p53 tumor suppressor gene.
74. PMID: 23860775; 2013, Tumour Biol;34(6):3945-58  
MYC overexpression and poor prognosis in sporadic breast cancer with BRCA1 deficiency.

# ACT Onco<sup>®</sup> + Report

75. PMID: 26156651; 2015, Lancet Oncol;16(8):897-907  
Afinib versus erlotinib as second-line treatment of patients with advanced squamous cell carcinoma of the lung (LUX-Lung 8): an open-label randomised controlled phase 3 trial.
76. PMID: 23816960; 2013, J Clin Oncol;31(27):3327-34  
Phase III study of afatinib or cisplatin plus pemetrexed in patients with metastatic lung adenocarcinoma with EGFR mutations.
77. PMID: 30280641; 2018, N Engl J Med;379(23):2220-2229  
First-Line Atezolizumab plus Chemotherapy in Extensive-Stage Small-Cell Lung Cancer.
78. PMID: 27979383; 2017, Lancet;389(10066):255-265  
Atezolizumab versus docetaxel in patients with previously treated non-small-cell lung cancer (OAK): a phase 3, open-label, multicentre randomised controlled trial.
79. PMID: 26970723; 2016, Lancet;387(10030):1837-46  
Atezolizumab versus docetaxel for patients with previously treated non-small-cell lung cancer (POPLAR): a multicentre, open-label, phase 2 randomised controlled trial.
80. PMID: 27939400; 2017, Lancet;389(10064):67-76  
Atezolizumab as first-line treatment in cisplatin-ineligible patients with locally advanced and metastatic urothelial carcinoma: a single-arm, multicentre, phase 2 trial.
81. PMID: 30779531; 2019, N Engl J Med;380(12):1103-1115  
Avelumab plus Axitinib versus Sunitinib for Advanced Renal-Cell Carcinoma.
82. PMID: 27592805; 2016, Lancet Oncol;17(10):1374-1385  
Avelumab in patients with chemotherapy-refractory metastatic Merkel cell carcinoma: a multicentre, single-group, open-label, phase 2 trial.
83. PMID: 33503393; 2021, J Clin Oncol;39(4):273-284  
Encorafenib Plus Cetuximab as a New Standard of Care for Previously Treated BRAF V600E-Mutant Metastatic Colorectal Cancer: Updated Survival Results and Subgroup Analyses from the BEACON Study.
84. PMID: 19339720; 2009, N Engl J Med;360(14):1408-17  
Cetuximab and chemotherapy as initial treatment for metastatic colorectal cancer.
85. PMID: 18784101; 2008, N Engl J Med;359(11):1116-27  
Platinum-based chemotherapy plus cetuximab in head and neck cancer.
86. PMID: 16467544; 2006, N Engl J Med;354(6):567-78  
Radiotherapy plus cetuximab for squamous-cell carcinoma of the head and neck.
87. PMID: 15269313; 2004, N Engl J Med;351(4):337-45  
Cetuximab monotherapy and cetuximab plus irinotecan in irinotecan-refractory metastatic colorectal cancer.
88. PMID: 31590988; 2019, Lancet;394(10212):1929-1939  
Durvalumab plus platinum-etoposide versus platinum-etoposide in first-line treatment of extensive-stage small-cell lung cancer (CASPIAN): a randomised, controlled, open-label, phase 3 trial.
89. PMID: 28885881; 2017, N Engl J Med;377(20):1919-1929  
Durvalumab after Chemoradiotherapy in Stage III Non-Small-Cell Lung Cancer.
90. PMID: 22285168; 2012, Lancet Oncol;13(3):239-46  
Erlotinib versus standard chemotherapy as first-line treatment for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EORTAC): a multicentre, open-label, randomised phase 3 trial.
91. PMID: 17452677; 2007, J Clin Oncol;25(15):1960-6  
Erlotinib plus gemcitabine compared with gemcitabine alone in patients with advanced pancreatic cancer: a phase III trial of the National Cancer Institute of Canada Clinical Trials Group.



# ACT Onco<sup>®</sup> + Report

92. PMID: 24263064; 2014, Br J Cancer;110(1):55-62  
First-line gefitinib in Caucasian EGFR mutation-positive NSCLC patients: a phase-IV, open-label, single-arm study.
93. PMID: 28734759; 2017, Lancet Oncol;18(9):1182-1191  
Nivolumab in patients with metastatic DNA mismatch repair-deficient or microsatellite instability-high colorectal cancer (CheckMate 142): an open-label, multicentre, phase 2 study.
94. PMID: 29562145; 2018, N Engl J Med;378(14):1277-1290  
Nivolumab plus Ipilimumab versus Sunitinib in Advanced Renal-Cell Carcinoma.
95. PMID: 25840693; 2015, Lancet Oncol;16(5):522-30  
Adjuvant ipilimumab versus placebo after complete resection of high-risk stage III melanoma (EORTC 18071): a randomised, double-blind, phase 3 trial.
96. PMID: 20525992; 2010, N Engl J Med;363(8):711-23  
Improved survival with ipilimumab in patients with metastatic melanoma.
97. PMID: 33632775; 2021, Cancer Discov;11(7):1688-1699  
Activity and Safety of Mobocertinib (TAK-788) in Previously Treated Non-Small Cell Lung Cancer with EGFR Exon 20 Insertion Mutations from a Phase I/II Trial.
98. PMID: 26045340; 2015, Lancet Oncol;16(7):763-74  
Necitumumab plus gemcitabine and cisplatin versus gemcitabine and cisplatin alone as first-line therapy in patients with stage IV squamous non-small-cell lung cancer (SQUIRE): an open-label, randomised, controlled phase 3 trial.
99. PMID: 27718784; 2016, N Engl J Med;375(19):1856-1867  
Nivolumab for Recurrent Squamous-Cell Carcinoma of the Head and Neck.
100. PMID: 27451390; 2016, Lancet Oncol;17(9):1283-94  
Nivolumab for classical Hodgkin's lymphoma after failure of both autologous stem-cell transplantation and brentuximab vedotin: a multicentre, multicohort, single-arm phase 2 trial.
101. PMID: 25482239; 2015, N Engl J Med;372(4):311-9  
PD-1 blockade with nivolumab in relapsed or refractory Hodgkin's lymphoma.
102. PMID: 26027431; 2015, N Engl J Med;373(1):23-34  
Combined Nivolumab and Ipilimumab or Monotherapy in Untreated Melanoma.
103. PMID: 25399552; 2015, N Engl J Med;372(4):320-30  
Nivolumab in previously untreated melanoma without BRAF mutation.
104. PMID: 26406148; 2015, N Engl J Med;373(19):1803-13  
Nivolumab versus Everolimus in Advanced Renal-Cell Carcinoma.
105. PMID: 26412456; 2015, N Engl J Med;373(17):1627-39  
Nivolumab versus Docetaxel in Advanced Nonsquamous Non-Small-Cell Lung Cancer.
106. PMID: 26028407; 2015, N Engl J Med;373(2):123-35  
Nivolumab versus Docetaxel in Advanced Squamous-Cell Non-Small-Cell Lung Cancer.
107. PMID: 25795410; 2015, Lancet Oncol;16(4):375-84  
Nivolumab versus chemotherapy in patients with advanced melanoma who progressed after anti-CTLA-4 treatment (CheckMate 037): a randomised, controlled, open-label, phase 3 trial.
108. PMID: 27959700; 2017, N Engl J Med;376(7):629-640  
Osimertinib or Platinum-Pemetrexed in EGFR T790M-Positive Lung Cancer.
109. PMID: 27736842; 2016, Br J Cancer;115(10):1206-1214  
A phase 3 trial evaluating panitumumab plus best supportive care vs best supportive care in chemorefractory wild-type KRAS or RAS metastatic colorectal cancer.

# ACT Onco<sup>®</sup> + Report

110. PMID: 24024839; 2013, N Engl J Med;369(11):1023-34  
Panitumumab-FOLFOX4 treatment and RAS mutations in colorectal cancer.
111. PMID: 24739896; 2014, Lancet Oncol;15(6):569-79  
Panitumumab versus cetuximab in patients with chemotherapy-refractory wild-type KRAS exon 2 metastatic colorectal cancer (ASPECCT): a randomised, multicentre, open-label, non-inferiority phase 3 study.
112. PMID: 17470858; 2007, J Clin Oncol;25(13):1658-64  
Open-label phase III trial of panitumumab plus best supportive care compared with best supportive care alone in patients with chemotherapy-refractory metastatic colorectal cancer.
113. PMID: 30779529; 2019, N Engl J Med;380(12):1116-1127  
Pembrolizumab plus Axitinib versus Sunitinib for Advanced Renal-Cell Carcinoma.
114. PMID: 27093365; 2016, N Engl J Med;374(26):2542-52  
PD-1 Blockade with Pembrolizumab in Advanced Merkel-Cell Carcinoma.
115. PMID: 29875066; 2018, Lancet Oncol;19(7):940-952  
Pembrolizumab in patients with advanced hepatocellular carcinoma previously treated with sorafenib (KEYNOTE-224): a non-randomised, open-label phase 2 trial.
116. PMID: 29658856; 2018, N Engl J Med;378(22):2078-2092  
Pembrolizumab plus Chemotherapy in Metastatic Non-Small-Cell Lung Cancer.
117. PMID: 28291584; 2017, Lancet Oncol;18(5):623-630  
Clinical safety and activity of pembrolizumab in patients with malignant pleural mesothelioma (KEYNOTE-028): preliminary results from a non-randomised, open-label, phase 1b trial.
118. PMID: 28489510; 2017, J Clin Oncol;35(22):2535-2541  
Safety and Antitumor Activity of Pembrolizumab in Advanced Programmed Death Ligand 1-Positive Endometrial Cancer: Results From the KEYNOTE-028 Study.
119. PMID: 27138582; 2016, J Clin Oncol;34(21):2460-7  
Pembrolizumab in Patients With Advanced Triple-Negative Breast Cancer: Phase 1b KEYNOTE-012 Study.
120. PMID: 28081914; 2017, Lancet Oncol;18(2):212-220  
Safety and activity of pembrolizumab in patients with locally advanced or metastatic urothelial cancer (KEYNOTE-012): a non-randomised, open-label, phase 1b study.
121. PMID: 27247226; 2016, Lancet Oncol;17(7):956-965  
Safety and clinical activity of pembrolizumab for treatment of recurrent or metastatic squamous cell carcinoma of the head and neck (KEYNOTE-012): an open-label, multicentre, phase 1b trial.
122. PMID: 27157491; 2016, Lancet Oncol;17(6):717-726  
Pembrolizumab for patients with PD-L1-positive advanced gastric cancer (KEYNOTE-012): a multicentre, open-label, phase 1b trial.
123. PMID: 28212060; 2017, N Engl J Med;376(11):1015-1026  
Pembrolizumab as Second-Line Therapy for Advanced Urothelial Carcinoma.
124. PMID: 28441111; 2017, J Clin Oncol;35(19):2125-2132  
Phase II Study of the Efficacy and Safety of Pembrolizumab for Relapsed/Refractory Classic Hodgkin Lymphoma.
125. PMID: 27718847; 2016, N Engl J Med;375(19):1823-1833  
Pembrolizumab versus Chemotherapy for PD-L1-Positive Non-Small-Cell Lung Cancer.
126. PMID: 25891173; 2015, N Engl J Med;372(26):2521-32  
Pembrolizumab versus Ipilimumab in Advanced Melanoma.
127. PMID: 26712084; 2016, Lancet;387(10027):1540-50

# ACT Onco<sup>®</sup> + Report

Pembrolizumab versus docetaxel for previously treated, PD-L1-positive, advanced non-small-cell lung cancer (KEYNOTE-010): a randomised controlled trial.

128. PMID: 26115796; 2015, Lancet Oncol;16(8):908-18  
Pembrolizumab versus investigator-choice chemotherapy for ipilimumab-refractory melanoma (KEYNOTE-002): a randomised, controlled, phase 2 trial.