



Sample Information

Patient Name: 張淑晶
Gender: Female
ID No.: A223978111
History No.: 46783500
Age: 47

Ordering Doctor: DOC8919E 周逸峰
Ordering REQ.: 0BZXUCG
Signing in Date: 2022/09/22

Path No.: S111-97900
MP No.: MY22027
Assay: Oncomine Myeloid Assay
Sample Type: Bone Marrow
Bone Marrow Aspirating Date: 2022/09/13

Reporting Doctor: DOC5466K 葉奕成 (Phone: 8#5466)

Note:

Sample Cancer Type: Chronic Myelomonocytic Leukemia

Table of Contents	Page	Report Highlights
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2	2 Relevant Biomarkers
Biomarker Descriptions	2	13 Therapies Available
Relevant Therapy Summary	5	0 Clinical Trials
Relevant Therapy Details	6	
Prognostic Details	11	

Relevant Chronic Myelomonocytic Leukemia Variants

Gene	Finding
ASXL1	ASXL1 p.(Y591*) c.1772_1773insAAGGTCAGCCCACTTA

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	ASXL1 p.(Y591*) c.1772_1773insAAGGTCAGCCCACTTA ASXL transcriptional regulator 1 Allele Frequency: 23.84%	None	allogeneic stem cells azacitidine cytarabine cytarabine + daunorubicin cytarabine + daunorubicin + etoposide	0

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

Relevant Biomarkers (continued)

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
			cytarabine + etoposide + idarubicin cytarabine + fludarabine + idarubicin + filgrastim cytarabine + idarubicin cytarabine + mitoxantrone decitabine gemtuzumab ozogamicin + chemotherapy venetoclax + chemotherapy	
	Prognostic significance: NCCN: Poor			
IIC	<i>SF3B1</i> p.(K666N) c.1998G>T splicing factor 3b subunit 1 Allele Frequency: 39.52%	None	luspatercept	0

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

Prevalent cancer biomarkers without relevant evidence based on included data sources

NRAS p.(Q61H) c.183A>T, *PTPN11* p.(E76K) c.226G>A, *IKZF1* p.(S308Cfs*4) c.922_923insGT, *U2AF1* p.(Q157P) c.470A>C

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
NRAS	p.(Q61H)	c.183A>T	COSM585	chr1:115256528	23.52%	NM_002524.5	missense	1998
SF3B1	p.(K666N)	c.1998G>T	COSM131557	chr2:198267359	39.52%	NM_012433.4	missense	1999
IKZF1	p.(S308Cfs*4)	c.922_923insGT	.	chr7:50467685	7.52%	NM_006060.6	frameshift Insertion	1996
PTPN11	p.(E76K)	c.226G>A	COSM13000	chr12:112888210	18.01%	NM_002834.5	missense	1999
ASXL1	p.(Y591*)	c.1772_1773insAAG GTCAGCCCACTTA	.	chr20:31022268	23.84%	NM_015338.6	nonsense	1204
U2AF1	p.(Q157P)	c.470A>C	COSM211534	chr21:44514777	39.40%	NM_006758.2	missense	2000
CSF3R	p.(Q776E)	c.2326C>G	.	chr1:36932224	42.40%	NM_156039.3	missense	2000
IKZF1	p.(H480Q)	c.1440C>G	.	chr7:50468205	18.10%	NM_006060.6	missense	2000
U2AF1L5	p.(Q84P)	c.251A>C	COSM211534	chr21:44514777	39.40%	NM_001320651.1	missense	2000
BCOR	p.(P483L)	c.1448C>T	.	chrX:39933151	48.52%	NM_001123385.2	missense	1999

Biomarker Descriptions

ASXL1 (ASXL transcriptional regulator 1)

Background: The ASXL1 gene encodes the ASXL transcriptional regulator 1 protein, a ligand-dependent co-activator and epigenetic scaffolding protein involved in transcriptional regulation^{1,2}. ASXL1 belongs to the ASXL gene family, which also includes ASXL2 and ASXL3². ASXL proteins contain a conserved c-terminal plant homeodomain (PHD) which facilitates interaction with DNA and histones^{2,3}. ASXL1 influences chromatin remodeling and transcription through interaction with BAP1 and polycomb repressive complex

Biomarker Descriptions (continued)

(PRC) proteins, as well as other transcriptional activators and repressors²⁴. In cancer, ASXL1 is the target of somatic mutations which often result in a truncated ASXL1 protein and loss of its PHD^{5,6,7}. Such mutations can lead to impaired protein function and consequent upregulation of HOXA gene expression, supporting a tumor suppressor role for ASXL1⁸.

Alterations and prevalence: Missense, nonsense, and frameshift mutations in ASXL1 are reported in 3-6% of de novo acute myeloid leukemia (AML), up to 36% of secondary AML, approximately 15% of myelodysplastic syndromes (MDS), up to 23% of myeloproliferative neoplasms (MPN), up to 30% of systemic mastocytosis (SM), and approximately 45% of chronic myelomonocytic leukemia (CMML)^{4,9,10,11,12,13,14,15,16}. The ASXL1 G646Wfs*12 mutation accounts for over 50% of ASXL1 mutated cases in myeloid malignancies^{6,11,17}. This mutation results from a single nucleotide expansion that occurs within an eight base pair guanine repeat that extends from c.1927 to c.1934. It is proposed that the high prevalence of the G646Wfs*12 variant is due to replication slippage which can occur in areas of repetitive sequence¹⁸. As a consequence, detection of G646Wfs*12 may result as an artifact of PCR and/or sequencing¹⁹. However, multiple studies observe an increase in the frequency of G646Wfs*12 in myeloid cancer relative to normal suggesting that G646Wfs*12 is a bona fide somatic mutation^{9,18,20}.

Potential relevance: The majority of frameshift and nonsense mutations in ASXL1 that result in protein truncation and removal of the PHD domain are considered pathogenic²¹. Mutations in ASXL1 confer poor/adverse risk in AML and are associated with poor outcomes and adverse risk¹⁶. Additionally, ASXL1 nonsense or frameshift mutations are independently associated with poor prognosis in MDS and CMML²². Moreover, ASXL1 mutations are independently associated with inferior overall survival (OS) in patients with MPN or SM^{23,24}.

IKZF1 (IKAROS family zinc finger 1)

Background: The IKZF1 gene encodes the IKAROS family zinc finger 1 transcription factor belonging to the family of zinc-finger DNA-binding proteins that influence gene expression through chromatin remodeling. IKZF1 was originally characterized as a lymphoid-restricted transcription factor for its critical role in lymphocyte differentiation²⁵. However, it has since been observed to be involved in myeloid differentiation as well^{26,27,28}. IKZF1 is a tumor suppressor and target of genetic aberrations including deletions and mutations that lead to loss of function and contribute to the development of lymphoid malignancies^{29,30,31}. Specifically, IKZF1 deletion is an acquired alteration associated with the transformation of chronic myeloid leukemia (CML) to lymphoid blast crisis in BCR-ABL1 positive acute lymphoblastic leukemia (ALL)^{32,33}. Similarly, IKZF1 deletion is associated with acute myeloid leukemia (AML) and transformation of myeloproliferative neoplasms (MPN) to AML^{31,34,35}.

Alterations and prevalence: IKZF1 aberrations, including deletions and mutations, occur in 15-20% of pediatric B-cell ALL and in greater than 75% of BCR-ABL1 positive ALL^{32,36,37,38}. In adults, IKZF1 aberrations occur in about 25-35% of ALL and in approximately 65% of BCR-ABL1 positive ALL^{36,39,40,41}. IKZF1 deletion has also been observed in 4% of diffuse large B-cell lymphoma (DLBCL) and 1% of pediatric AML^{12,34}. IKZF1 mutations are observed in about 2% of AML and myelodysplastic syndromes (MDS)⁴². Mutations in IKZF1 usually result in expression of a protein that lacks DNA binding ability with dominant negative activity^{37,38}.

Potential relevance: IKZF1 mutations and deletions that result in loss of function are associated with poor prognosis and a greater chance of relapse in B-cell ALL^{36,43}.

NRAS (NRAS proto-oncogene, GTPase)

Background: The NRAS proto-oncogene encodes a GTPase that functions in signal transduction and is a member of the RAS superfamily which also includes KRAS and HRAS. RAS proteins mediate the transmission of growth signals from the cell surface to the nucleus via the PI3K/AKT/MTOR and RAS/RAF/MEK/ERK pathways, which regulate cell division, differentiation, and survival^{44,45,46}.

Alterations and prevalence: Recurrent mutations in RAS oncogenes cause constitutive activation and are found in 20-30% of cancers. NRAS mutations are particularly common in melanomas (up to 25%) and are observed at frequencies of 5-10% in acute myeloid leukemia, colorectal, and thyroid cancers^{12,47}. The majority of NRAS mutations consist of point mutations at G12, G13, and Q61^{12,48}. Mutations at A59, K117, and A146 have also been observed but are less frequent^{49,50}.

Potential relevance: Currently, no therapies are approved for NRAS aberrations. The EGFR antagonists, cetuximab⁵¹ and panitumumab⁵², are contraindicated for treatment of colorectal cancer patients with NRAS mutations in exon 2 (codons 12 and 13), exon 3 (codons 59 and 61), and exon 4 (codons 117 and 146)⁵⁰. NRAS mutations are associated with poor prognosis in patients with low-risk myelodysplastic syndrome²² as well as melanoma⁵³. In a phase III clinical trial in patients with advanced NRAS-mutant melanoma, binimetinib improved progression free survival (PFS) relative to dacarbazine with median PFS of 2.8 and 1.5 months, respectively⁵⁴.

PTPN11 (protein tyrosine phosphatase non-receptor type 11)

Background: The PTPN11 gene encodes a tyrosine phosphatase non-receptor type 11 protein, and is also known as Src homology region 2 domain-containing phosphatase-2 (SHP-2)⁵⁵. PTPN11 is a member of the protein tyrosine phosphatase (PTP) family that

Biomarker Descriptions (continued)

is ubiquitously expressed and regulates cellular growth, differentiation, mitotic cycle, and oncogenic transformation. PTPN11 contains two tandem N-terminal Src homology-2 domains (N-SH2 and C-SH2), a PTP catalytic domain, and uncharacterized C-terminal domain⁵⁶. PTPN11 regulates various signaling processes including the RAS/RAF/MEK/ERK, PI3K/AKT/MTOR, and JAK/STAT pathways^{57,58}. Germline mutations in PTPN11 are associated with LEOPARD syndrome and Noonan syndrome with a predisposition to juvenile myelomonocytic leukemia (JMML) or myeloproliferative neoplasms (MPN)^{22,59}. Somatic mutations in PTPN11 are associated with JMML^{60,61} and solid tumors such as lung, colon, and thyroid^{56,62}.

Alterations and prevalence: Somatic alterations in PTPN11 include mutations and amplification^{59,63}. PTPN11 mutations occur in 6% of uterine carcinoma and 5% of acute myeloid leukemia (AML) cases⁴⁹. Mutations including E76K and D61Y result in PTPN11 activation and are associated with 30% of JMML⁵⁸.

Potential relevance: Currently, no therapies are approved for PTPN11 aberrations. Somatic mutations in PTPN11 confer drug resistance to venetoclax and azacitidine in AML^{64,65}.

SF3B1 (splicing factor 3b subunit 1)

Background: The SF3B1 gene encodes the splicing factor 3b subunit 1 protein, a core component of the U2 small nuclear ribonucleoprotein (snRNP) complex of the spliceosome responsible for RNA splicing. SF3B1 is involved in recognition of the branch point sequence during selection of the 3' splice site. Recurrent somatic mutations in SF3B1 and other components of the splicing machinery including SRSF2, U2AF1, and ZRSR2, are common in myelodysplasia. These components experience mutations in a mutually exclusive manner suggesting a common impact on RNA splicing and the pathogenesis of myelodysplasia⁶⁶. SF3B1 mutations are believed to contribute to aberrant post-translational inactivation of the regulatory complex PPP2R5A of protein phosphatase 2A (PP2A), leading to the activation and stabilization of MYC activation and impairing apoptosis⁶⁷.

Alterations and prevalence: SF3B1 mutations occur in the majority (70-80%) of myelodysplastic syndromes (MDS) with ring sideroblasts (RS) and at lower frequency in other myeloid neoplasms including MDS without RS (7%), chronic myelomonocytic leukemia (5-6%), therapy-related acute myeloid leukemia (AML) or AML with MDS features (5%), and de novo AML (3%)^{66,68,69}. Recurrent somatic SF3B1 mutations are also common in certain solid cancers including uveal melanoma (20-30%) and breast cancer (2%) and at lower frequencies in diverse cancer types^{12,49,70,71,72,73,74,75}. Cancer-associated recurrent missense mutations in SF3B1 occur within the HEAT repeat domains 5-9 at codon positions R625, K666, K700, G742, and D781⁷⁶. The functional significance of recurrent SF3B1 mutations is to alter branch point selection thus inducing cryptic 3' splice site selection^{76,77,78}.

Potential relevance: Currently, no therapies are approved for SF3B1 aberrations. SF3B1 mutations are associated with aggressive disease and shorter survival in patients diagnosed with chronic lymphocytic leukemia (CLL)⁷⁹. Investigational inhibitors of the spliceosome are in early clinical development^{80,81}.

U2AF1 (U2 small nuclear RNA auxiliary factor 1)

Background: The U2AF1 gene encodes the U2 small nuclear RNA auxiliary factor 1 protein that belongs to the splicing factor SR family of genes involved in RNA splicing^{1,82}. U2AF1, also known as U2AF35, mediates the recruitment of the U2AF complex to the 3' end of that pre-mRNA that is being spliced⁸³. U2AF1 is the smaller subunit of the U2 auxiliary factor and along with the larger subunit, U2AF65 regulates the removal of introns from pre-mRNAs to produce mature mRNAs for translation during protein synthesis⁸⁴. Mutations in U2AF1 alter the differential splicing of genes that are involved in various biological pathways, including DNMT3B in DNA methylation, ATR along with FANCA in DNA damage response, and H2AFY in X-chromosome inactivation⁸⁵. Spliceosomal genes such as U2AF1 are common targets of somatic mutations in myelodysplastic syndrome (MDS) and are associated with the progression of MDS to acute myeloid leukemia (AML)^{85,86,87}.

Alterations and prevalence: Recurrent mutations in U2AF1 occur at S34 and Q157 and are observed in 8-12% of MDS²². Somatic mutations in U2AF1 are also observed in 10% of uterine carcinoma, 4% of AML, as well as 2% of lung adenocarcinoma and stomach adenocarcinoma⁴⁹.

Potential relevance: U2AF1 mutations including S34 and Q157 are associated with poor prognosis in MDS²². U2AF1 mutations are associated with inferior overall survival in primary myelofibrosis (PMF)²³. Specifically, the Q157 mutation is associated with a significantly shorter overall survival than U2AF1 S34 mutated and U2AF1 unmutated MPN²³.

Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

ASXL1 p.(Y591*) c.1772_1773insAAGGTCAGCCCACTTA

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
Allogeneic hematopoietic stem cell transplantation	×	○	×	×	×
azacitidine	×	○	×	×	×
cytarabine	×	○	×	×	×
cytarabine + daunorubicin	×	○	×	×	×
cytarabine + daunorubicin + etoposide	×	○	×	×	×
cytarabine + etoposide + idarubicin	×	○	×	×	×
cytarabine + fludarabine + idarubicin + filgrastim	×	○	×	×	×
cytarabine + idarubicin	×	○	×	×	×
cytarabine + mitoxantrone	×	○	×	×	×
decitabine	×	○	×	×	×
gemtuzumab ozogamicin + cytarabine + daunorubicin	×	○	×	×	×
venetoclax + azacitidine	×	○	×	×	×
venetoclax + cytarabine	×	○	×	×	×
venetoclax + decitabine	×	○	×	×	×

SF3B1 p.(K666N) c.1998G>T

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
luspaterecept	×	×	×	○	×

Relevant Therapy Details

Current NCCN Information

- ☒ In this cancer type ☐ In other cancer type ☒ In this cancer type and other cancer types

NCCN information is current as of 2022-08-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

ASXL1 p.(Y591*) c.1772_1773insAAGGTCAGCCCCTTA

☐ cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

☐ cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

☐ cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

☐ cytarabine + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

ASXL1 p.(Y591*) c.1772_1773insAAGGTCAGCCCACTTA (continued)**○ Allogeneic hematopoietic stem cell transplantation**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ cytarabine

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ cytarabine + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

ASXL1 p.(Y591*) c.1772_1773insAAGGTCAGCCCACTTA (continued)**○ cytarabine + mitoxantrone**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ gemtuzumab ozogamicin + cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ venetoclax + cytarabine

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

ASXL1 p.(Y591*) c.1772_1773insAAGGTCAGCCCACTTA (continued)**○ venetoclax + decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

Current ESMO Information

☒ In this cancer type ☐ In other cancer type ☒ In this cancer type and other cancer types

ESMO information is current as of 2022-08-01. For the most up-to-date information, search www.esmo.org.

SF3B1 p.(K666N) c.1998G>T

☐ Iuspatercept

Cancer type: Myelodysplastic Syndrome

Variant class: SF3B1 mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Refractory (Second-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Myelodysplastic Syndromes [Annals of Oncology (2020), <https://doi.org/10.1016/j.annonc.2020.11.002>]

Prognostic Details

Current NCCN Information

NCCN information is current as of 2022-08-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

ASXL1 p.(Y591*) c.1772_1773insAAGGTCAGCCCCTTA

Prognostic significance: NCCN: Poor

Cancer type: Chronic Myelomonocytic Leukemia

Variant class: ASXL1 truncating mutation

NCCN Recommendation category: 2A

Summary:

- NCCN Guidelines® independently associate the biomarker with poor prognosis in MDS and CMML

Reference: NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 3.2022]

Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

References

1. O'Leary et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res.* 2016 Jan 4;44(D1):D733-45. PMID: 26553804
2. Katoh. Functional and cancer genomics of ASXL family members. *Br. J. Cancer.* 2013 Jul 23;109(2):299-306. PMID: 23736028
3. Gelsi-Boyer et al. Mutations of polycomb-associated gene ASXL1 in myelodysplastic syndromes and chronic myelomonocytic leukaemia. *Br. J. Haematol.* 2009 Jun;145(6):788-800. PMID: 19388938
4. Gelsi-Boyer et al. Mutations in ASXL1 are associated with poor prognosis across the spectrum of malignant myeloid diseases. *J Hematol Oncol.* 2012 Mar 21;5:12. doi: 10.1186/1756-8722-5-12. PMID: 22436456
5. Larsson et al. The changing mutational landscape of acute myeloid leukemia and myelodysplastic syndrome. *Mol. Cancer Res.* 2013 Aug;11(8):815-27. PMID: 23645565
6. Alvarez et al. ASXL1 mutations in myeloid neoplasms: pathogenetic considerations, impact on clinical outcomes and survival. *Curr Med Res Opin.* 2018 May;34(5):757-763. PMID: 28027687
7. Yang et al. Gain of function of ASXL1 truncating protein in the pathogenesis of myeloid malignancies. *Blood.* 2018 Jan 18;131(3):328-341. PMID: 29113963
8. Abdel-Wahab et al. ASXL1 mutations promote myeloid transformation through loss of PRC2-mediated gene repression. *Cancer Cell.* 2012 Aug 14;22(2):180-93. PMID: 22897849
9. Alberti et al. Discriminating a common somatic ASXL1 mutation (c.1934dup; p.G646Wfs*12) from artifact in myeloid malignancies using NGS. *Leukemia.* 2018 Aug;32(8):1874-1878. PMID: 29959414
10. Kakosaiou et al. ASXL1 mutations in AML are associated with specific clinical and cytogenetic characteristics. *Leuk. Lymphoma.* 2018 Oct;59(10):2439-2446. PMID: 29411666
11. Paschka et al. ASXL1 mutations in younger adult patients with acute myeloid leukemia: a study by the German-Austrian Acute Myeloid Leukemia Study Group. *Haematologica.* 2015 Mar;100(3):324-30. PMID: 25596267
12. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
13. Jawhar et al. The clinical and molecular diversity of mast cell leukemia with or without associated hematologic neoplasm. *Haematologica.* 2017 Jun;102(6):1035-1043. PMID: 28255023
14. Jawhar et al. KIT D816 mutated/CBF-negative acute myeloid leukemia: a poor-risk subtype associated with systemic mastocytosis. *Leukemia.* 2019 May;33(5):1124-1134. PMID: 30635631
15. Damaj et al. ASXL1 but not TET2 mutations adversely impact overall survival of patients suffering systemic mastocytosis with associated clonal hematologic non-mast-cell diseases. *PLoS ONE.* 2014;9(1):e85362. PMID: 24465546
16. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]
17. Boulton et al. Frequent mutation of the polycomb-associated gene ASXL1 in the myelodysplastic syndromes and in acute myeloid leukemia. *Leukemia.* 2010 May;24(5):1062-5. doi: 10.1038/leu.2010.20. Epub 2010 Feb 25. PMID: 20182461
18. Yannakou et al. ASXL1 c.1934dup;p.Gly646Trpfs*12-a true somatic alteration requiring a new approach. *Blood Cancer J.* 2017 Dec 15;7(12):656. doi: 10.1038/s41408-017-0025-8. PMID: 29242575
19. Abdel-Wahab et al. The most commonly reported variant in ASXL1 (c.1934dupG;p.Gly646TrpfsX12) is not a somatic alteration. *Leukemia.* 2010 Sep;24(9):1656-7. doi: 10.1038/leu.2010.144. Epub 2010 Jul 1. PMID: 20596031
20. Montes-Moreno et al. Clinical molecular testing for ASXL1 c.1934dupG p.Gly646fs mutation in hematologic neoplasms in the NGS era. *PLoS ONE.* 2018;13(9):e0204218. PMID: 30222780
21. Landrum et al. ClinVar: improving access to variant interpretations and supporting evidence. *Nucleic Acids Res.* 2018 Jan 4;46(D1):D1062-D1067. PMID: 29165669
22. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 3.2022]
23. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 2.2022]
24. NCCN Guidelines® - NCCN-Systemic Mastocytosis [Version 1.2020]
25. Georgopoulos et al. Ikaros, an early lymphoid-specific transcription factor and a putative mediator for T cell commitment. *Science.* 1992 Oct 30;258(5083):808-12. PMID: 1439790
26. Klug et al. Hematopoietic stem cells and lymphoid progenitors express different Ikaros isoforms, and Ikaros is localized to heterochromatin in immature lymphocytes. *Proc. Natl. Acad. Sci. U.S.A.* 1998 Jan 20;95(2):657-62. PMID: 9435248
27. Yoshida et al. Early hematopoietic lineage restrictions directed by Ikaros. *Nat. Immunol.* 2006 Apr;7(4):382-91. PMID: 16518393
28. Francis et al. Regulator of myeloid differentiation and function: The secret life of Ikaros. *World J Biol Chem.* 2011 Jun 26;2(6):119-25. PMID: 21765977
29. Payne et al. Ikaros and tumor suppression in acute lymphoblastic leukemia. *Crit Rev Oncog.* 2011;16(1-2):3-12. PMID: 22150303

References (continued)

30. Kastner et al. Role of Ikaros in T-cell acute lymphoblastic leukemia. *World J Biol Chem.* 2011 Jun 26;2(6):108-14. PMID: 21765975
31. Marke et al. The many faces of IKZF1 in B-cell precursor acute lymphoblastic leukemia. *Haematologica.* 2018 Apr;103(4):565-574. PMID: 29519871
32. Mullighan et al. BCR-ABL1 lymphoblastic leukaemia is characterized by the deletion of Ikaros. *Nature.* 2008 May 1;453(7191):110-4. PMID: 18408710
33. Kang et al. The Philadelphia chromosome in leukemogenesis. *Chin J Cancer.* 2016 May 27;35:48. doi: 10.1186/s40880-016-0108-0. PMID: 27233483
34. de et al. Recurrent deletions of IKZF1 in pediatric acute myeloid leukemia. *Haematologica.* 2015 Sep;100(9):1151-9. PMID: 26069293
35. Jäger et al. Deletions of the transcription factor Ikaros in myeloproliferative neoplasms. *Leukemia.* 2010 Jul;24(7):1290-8. PMID: 20508609
36. NCCN Guidelines® - Acute Lymphoblastic Leukemia [Version 2.2019]. 2019 May 15
37. Mullighan et al. Deletion of IKZF1 and prognosis in acute lymphoblastic leukemia. *N. Engl. J. Med.* 2009 Jan 29;360(5):470-80. PMID: 19129520
38. Caye et al. Breakpoint-specific multiplex polymerase chain reaction allows the detection of IKZF1 intragenic deletions and minimal residual disease monitoring in B-cell precursor acute lymphoblastic leukemia. *Haematologica.* 2013 Apr;98(4):597-601. PMID: 23065506
39. Dupuis et al. Biclonal and biallelic deletions occur in 20% of B-ALL cases with IKZF1 mutations. *Leukemia.* 2013 Feb;27(2):503-7. doi: 10.1038/leu.2012.204. Epub 2012 Jul 18. PMID: 22868967
40. Mi et al. Newly diagnosed acute lymphoblastic leukemia in China (II): prognosis related to genetic abnormalities in a series of 1091 cases. *Leukemia.* 2012 Jul;26(7):1507-16. PMID: 22297722
41. Ribera et al. Prognostic significance of copy number alterations in adolescent and adult patients with precursor B acute lymphoblastic leukemia enrolled in PETHEMA protocols. *Cancer.* 2015 Nov 1;121(21):3809-17. PMID: 26194343
42. Ethan et al. AACR Project GENIE: Powering Precision Medicine through an International Consortium. *Cancer Discov.* 2017 Aug;7(8):818-831. PMID: 28572459
43. Boer et al. Prognostic value of rare IKZF1 deletion in childhood B-cell precursor acute lymphoblastic leukemia: an international collaborative study. *Leukemia.* 2016 Jan;30(1):32-8. PMID: 26202931
44. Pylayeva-Gupta et al. RAS oncogenes: weaving a tumorigenic web. *Nat. Rev. Cancer.* 2011 Oct 13;11(11):761-74. PMID: 21993244
45. Karnoub et al. Ras oncogenes: split personalities. *Nat. Rev. Mol. Cell Biol.* 2008 Jul;9(7):517-31. PMID: 18568040
46. Scott et al. Therapeutic Approaches to RAS Mutation. *Cancer J.* 2016 May-Jun;22(3):165-74. doi: 10.1097/PPO.000000000000187. PMID: 27341593
47. Janku et al. PIK3CA mutations frequently coexist with RAS and BRAF mutations in patients with advanced cancers. *PLoS ONE.* 2011;6(7):e22769. PMID: 21829508
48. Ohashi et al. Characteristics of lung cancers harboring NRAS mutations. *Clin. Cancer Res.* 2013 May 1;19(9):2584-91. PMID: 23515407
49. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
50. Allegra et al. Extended RAS Gene Mutation Testing in Metastatic Colorectal Carcinoma to Predict Response to Anti-Epidermal Growth Factor Receptor Monoclonal Antibody Therapy: American Society of Clinical Oncology Provisional Clinical Opinion Update 2015. *J. Clin. Oncol.* 2016 Jan 10;34(2):179-85. PMID: 26438111
51. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125084s279lbl.pdf
52. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125147s210lbl.pdf
53. Johnson et al. Treatment of NRAS-Mutant Melanoma. *Curr Treat Options Oncol.* 2015 Apr;16(4):15. doi: 10.1007/s11864-015-0330-z. PMID: 25796376
54. Dummer et al. Binimetinib versus dacarbazine in patients with advanced NRAS-mutant melanoma (NEMO): a multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol.* 2017 Apr;18(4):435-445. PMID: 28284557
55. Tartaglia et al. Somatic PTPN11 mutations in childhood acute myeloid leukaemia. *Br. J. Haematol.* 2005 May;129(3):333-9. PMID: 15842656
56. Chan et al. The tyrosine phosphatase Shp2 (PTPN11) in cancer. *Cancer Metastasis Rev.* 2008 Jun;27(2):179-92. PMID: 18286234

References (continued)

57. Steelman et al. Roles of the Raf/MEK/ERK and PI3K/PTEN/Akt/mTOR pathways in controlling growth and sensitivity to therapy-implications for cancer and aging. *Aging (Albany NY)*. 2011 Mar;3(3):192-222. PMID: 21422497
58. Liu et al. Inhibition of the Gab2/PI3K/mTOR signaling ameliorates myeloid malignancy caused by Ptpn11 (Shp2) gain-of-function mutations. *Leukemia*. 2017 Jun;31(6):1415-1422. PMID: 27840422
59. Tartaglia et al. Somatic mutations in PTPN11 in juvenile myelomonocytic leukemia, myelodysplastic syndromes and acute myeloid leukemia. *Nat. Genet.* 2003 Jun;34(2):148-50. PMID: 12717436
60. Tartaglia et al. Mutations in PTPN11, encoding the protein tyrosine phosphatase SHP-2, cause Noonan syndrome. *Nat. Genet.* 2001 Dec;29(4):465-8. PMID: 11704759
61. Tartaglia et al. Diversity and functional consequences of germline and somatic PTPN11 mutations in human disease. *Am. J. Hum. Genet.* 2006 Feb;78(2):279-90. PMID: 16358218
62. Jongmans et al. Cancer risk in patients with Noonan syndrome carrying a PTPN11 mutation. *Eur. J. Hum. Genet.* 2011 Aug;19(8):870-4. PMID: 21407260
63. Liu et al. Gain-of-function mutations of Ptpn11 (Shp2) cause aberrant mitosis and increase susceptibility to DNA damage-induced malignancies. *Proc. Natl. Acad. Sci. U.S.A.* 2016 Jan 26;113(4):984-9. PMID: 26755576
64. Chyla et al. Genetic Biomarkers Of Sensitivity and Resistance to Venetoclax Monotherapy in Patients With Relapsed Acute Myeloid Leukemia. *Am. J. Hematol.* 2018 May 17. PMID: 29770480
65. Zhang et al. Biomarkers Predicting Venetoclax Sensitivity and Strategies for Venetoclax Combination Treatment. *Blood* 2018 132:175; doi: <https://doi.org/10.1182/blood-2018-175.132:175>
66. Yoshida et al. Frequent pathway mutations of splicing machinery in myelodysplasia. *Nature*. 2011 Sep 11;478(7367):64-9. PMID: 21909114
67. Liu et al. Mutations in the RNA Splicing Factor SF3B1 Promote Tumorigenesis through MYC Stabilization. *Cancer Discov.* 2020 Mar 18. PMID: 32188705
68. Patnaik et al. Refractory anemia with ring sideroblasts (RARS) and RARS with thrombocytosis (RARS-T): 2017 update on diagnosis, risk-stratification, and management. *Am. J. Hematol.* 2017 Mar;92(3):297-310. PMID: 28188970
69. Patnaik et al. Spliceosome mutations involving SRSF2, SF3B1, and U2AF35 in chronic myelomonocytic leukemia: prevalence, clinical correlates, and prognostic relevance. *Am. J. Hematol.* 2013 Mar;88(3):201-6. PMID: 23335386
70. Ellis et al. Whole-genome analysis informs breast cancer response to aromatase inhibition. *Nature*. 2012 Jun 10;486(7403):353-60. PMID: 22722193
71. Maguire et al. SF3B1 mutations constitute a novel therapeutic target in breast cancer. *J. Pathol.* 2015 Mar;235(4):571-80. PMID: 25424858
72. Harbour et al. Recurrent mutations at codon 625 of the splicing factor SF3B1 in uveal melanoma. *Nat. Genet.* 2013 Feb;45(2):133-5. PMID: 23313955
73. Martin et al. Exome sequencing identifies recurrent somatic mutations in EIF1AX and SF3B1 in uveal melanoma with disomy 3. *Nat. Genet.* 2013 Aug;45(8):933-6. PMID: 23793026
74. Furney et al. SF3B1 mutations are associated with alternative splicing in uveal melanoma. *Cancer Discov.* 2013 Oct;3(10):1122-1129. PMID: 23861464
75. Dono et al. Mutation frequencies of GNAQ, GNA11, BAP1, SF3B1, EIF1AX and TERT in uveal melanoma: detection of an activating mutation in the TERT gene promoter in a single case of uveal melanoma. *Br. J. Cancer.* 2014 Feb 18;110(4):1058-65. PMID: 24423917
76. DeBoever et al. Transcriptome sequencing reveals potential mechanism of cryptic 3' splice site selection in SF3B1-mutated cancers. *PLoS Comput. Biol.* 2015 Mar;11(3):e1004105. PMID: 25768983
77. Darman et al. Cancer-Associated SF3B1 Hotspot Mutations Induce Cryptic 3' Splice Site Selection through Use of a Different Branch Point. *Cell Rep.* 2015 Nov 3;13(5):1033-45. PMID: 26565915
78. Alsafadi et al. Cancer-associated SF3B1 mutations affect alternative splicing by promoting alternative branchpoint usage. *Nat Commun.* 2016 Feb 4;7:10615. doi: 10.1038/ncomms10615. PMID: 26842708
79. Wan et al. SF3B1 mutations in chronic lymphocytic leukemia. *Blood*. 2013 Jun 6;121(23):4627-34. PMID: 23568491
80. Seiler et al. H3B-8800, an orally available small-molecule splicing modulator, induces lethality in spliceosome-mutant cancers. *Nat. Med.* 2018 May;24(4):497-504. PMID: 29457796
81. Lee et al. Therapeutic targeting of splicing in cancer. *Nat. Med.* 2016 Sep 7;22(9):976-86. PMID: 27603132
82. Long et al. The SR protein family of splicing factors: master regulators of gene expression. *Biochem. J.* 2009 Jan 1;417(1):15-27. PMID: 19061484

References (continued)

83. Zuo et al. The splicing factor U2AF35 mediates critical protein-protein interactions in constitutive and enhancer-dependent splicing. *Genes Dev.* 1996 Jun 1;10(11):1356-68. PMID: 8647433
84. Ruskin et al. A factor, U2AF, is required for U2 snRNP binding and splicing complex assembly. *Cell.* 1988 Jan 29;52(2):207-19. PMID: 2963698
85. Ilagan et al. U2AF1 mutations alter splice site recognition in hematological malignancies. *Genome Res.* 2015 Jan;25(1):14-26. PMID: 25267526
86. Graubert et al. Recurrent mutations in the U2AF1 splicing factor in myelodysplastic syndromes. *Nat. Genet.* 2011 Dec 11;44(1):53-7. PMID: 22158538
87. Papaemmanuil et al. Clinical and biological implications of driver mutations in myelodysplastic syndromes. *Blood.* 2013 Nov 21;122(22):3616-27; quiz 3699. PMID: 24030381