

ABOUT THE TEST FoundationOne®CDx is a next-generation sequencing (NGS) based assay that identifies genomic findings within hundreds of cancer-related genes.

PATIENT	DISEASE Colon adenocarcinoma (CRC)	PHYSICIAN	ORDERING PHYSICIAN Yeh, Yi-Chen	SPECIMEN	SPECIMEN SITE Head and Neck
	NAME Chao, Ching Feng		MEDICAL FACILITY Taipei Veterans General Hospital		SPECIMEN ID S111-54650 A (PF23034)
	DATE OF BIRTH 09 October 1974		ADDITIONAL RECIPIENT None		SPECIMEN TYPE Slide Deck
	SEX Male		MEDICAL FACILITY ID 205872		DATE OF COLLECTION 27 December 2022
	MEDICAL RECORD # 45561066		PATHOLOGIST Not Provided		SPECIMEN RECEIVED 10 April 2023

Biomarker Findings

Microsatellite status - MS-Stable
Tumor Mutational Burden - 4 Muts/Mb

Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

BRAF V600E
KRAS wildtype
NRAS wildtype
BARD1 W218*
TET2 E1151*
TP53 C176*

2 Disease relevant genes with no reportable alterations: **KRAS**, **NRAS**

Report Highlights

- Targeted therapies with **NCCN categories of evidence** in this tumor type: **Encorafenib + Cetuximab** (p. 9)
- Targeted therapies with **potential resistance** based on this patient's genomic findings: **Cetuximab** (p. 10), **Panitumumab** (p. 11)
- Evidence-matched **clinical trial options** based on this patient's genomic findings: (p. 14)
- Variants with **prognostic implications** for this tumor type that may impact treatment decisions: **BRAF V600E** (p. 5)
- Variants that may represent **clonal hematopoiesis** and may originate from non-tumor sources: **TET2 E1151*** (p. 7)

BIOMARKER FINDINGS

Microsatellite status - MS-Stable

Tumor Mutational Burden - 4 Muts/Mb

GENOMIC FINDINGS

BRAF - V600E

10 Trials see p. 16

KRAS - wildtype

0 Trials

NRAS - wildtype

0 Trials

BARD1 - W218*

10 Trials see p. 14

THERAPY AND CLINICAL TRIAL IMPLICATIONS

No therapies or clinical trials. See Biomarker Findings section

No therapies or clinical trials. See Biomarker Findings section

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)	THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)
Encorafenib + Cetuximab 2A	Dabrafenib + Trametinib
Cetuximab ✗	Encorafenib + Binimetinib
Panitumumab ✗	Vemurafenib + Cobimetinib
Cetuximab ✗	none
Panitumumab ✗	
Cetuximab ✗	none
Panitumumab ✗	
none	none

✗ Extensive evidence showing variant(s) in this sample may confer resistance to this therapy

2A NCCN category

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS (CH)

Genomic findings below may include nontumor somatic alterations, such as CH. The efficacy of targeting such nontumor somatic alterations is unknown. This content should be interpreted based on clinical context. Refer to appendix for additional information on CH.

TET2 - E1151* p. [7](#)

GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

TET2 - E1151* p. [7](#) **TP53 - C176*** p. [8](#)

NOTE Genomic alterations detected may be associated with activity of certain approved therapies; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and exhaustive. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies.

Therapies contained in this report may have been approved by the US FDA.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1606031-01

BIOMARKER FINDINGS

BIOMARKER

Microsatellite status

RESULT

MS-Stable

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence, MSS tumors are significantly less likely than MSI-H tumors to respond to anti-PD-1 immune checkpoint inhibitors¹⁻³, including approved therapies nivolumab and pembrolizumab⁴. In a retrospective analysis of 361 patients with solid tumors treated with pembrolizumab, 3% were MSI-H and experienced a significantly higher ORR compared with non-MSI-H cases (70% vs. 12%, $p=0.001$)⁵. For patients with chemotherapy-refractory microsatellite-stable (MSS) metastatic colorectal cancer (CRC), a Phase 3 trial reported no OS advantage from the combination of the PD-L1 inhibitor atezolizumab plus cobimetinib relative to regorafenib (8.9 vs. 8.5 months, HR=1.00);

atezolizumab monotherapy similarly did not prolong OS (7.1 vs. 8.5 months, HR=1.19)⁶. For patients with MSS CRC, a Phase 2 study combining ipilimumab and nivolumab reported an overall DCR of 25% (10/40)⁷. Two Phase 1 studies for patients with MSS CRC treated with regorafenib and nivolumab reported PFSs of 7.9 months⁸ and 5.7 months⁹, and a patient with MSS CRC refractory to chemotherapy treated with the PD-1 inhibitor sintilimab and regorafenib reported a CR¹⁰.

— Nontargeted Approaches —

MSI has not been found to be a predictive biomarker for combination chemotherapy regimens, including FOLFOX¹¹⁻¹² and FOLFIRI¹³⁻¹⁴. Patients with MSS CRC are more likely to benefit from postsurgical fluorouracil (FU)-based adjuvant therapy¹⁵⁻¹⁶ but less likely to benefit from irinotecan chemotherapy¹⁷.

FREQUENCY & PROGNOSIS

MSS colorectal cancers (CRCs) make up 70-85% of CRC cases^{3,18-22}. MSS colorectal cancers are

molecularly heterogeneous, driven by diverse mechanisms such as extensive DNA methylation, oncogenic mutations in KRAS or BRAF, or chromosomal instability²². Multiple studies have shown that MSS CRCs have a worse prognosis than MSI-high tumors^{18,23-29}.

FINDING SUMMARY

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor²⁰. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2^{20,30-31}. This sample is microsatellite-stable (MSS), equivalent to the clinical definition of an MSS tumor: one with mutations in none of the tested microsatellite markers^{19,32-33}. MSS status indicates MMR proficiency and typically correlates with intact expression of all MMR family proteins^{19-20,31,33}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606031-01

BIOMARKER FINDINGS

BIOMARKER

Tumor Mutational Burden

RESULT

4 Muts/Mb

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1³⁴⁻³⁶, anti-PD-1 therapies³⁴⁻³⁷, and combination nivolumab and ipilimumab³⁸⁻⁴³. In multiple pan-tumor studies, increased tissue tumor mutational burden (TMB) was associated with sensitivity to immune checkpoint inhibitors^{34-37,44-48}. In the KEYNOTE 158 trial of pembrolizumab monotherapy for patients with solid tumors, significant improvement in ORR was observed for patients with TMB ≥ 10 Muts/Mb (as measured by this assay) compared with those with TMB < 10 Muts/Mb in a large cohort that included multiple tumor types⁴⁴; similar findings were observed in the KEYNOTE 028 and 012 trials³⁷. At the same TMB cutpoint, retrospective analysis of patients with solid tumors treated with any checkpoint inhibitor identified that tissue TMB scores ≥ 10 Muts/Mb were associated with prolonged time to treatment failure compared with scores < 10 Muts/Mb (HR=0.68)⁴⁸. For patients with solid tumors treated with nivolumab plus ipilimumab in the CheckMate 848 trial, improved responses were observed in patients with a tissue TMB ≥ 10 Muts/Mb independent of blood TMB at any cutpoint in matched samples⁴⁹. However, support for higher TMB thresholds and efficacy was observed in the prospective Phase 2 MyPathway trial of atezolizumab for patients with pan-solid tumors, where improved ORR and DCR

was seen in patients with TMB ≥ 16 Muts/Mb than those with TMB ≥ 10 and < 16 Muts/Mb⁴⁷. Similarly, analyses across several solid tumor types reported that patients with higher TMB (defined as $\geq 16-20$ Muts/Mb) achieved greater clinical benefit from PD-1 or PD-L1-targeting monotherapy compared with patients with higher TMB treated with chemotherapy⁵⁰ or those with lower TMB treated with PD-1 or PD-L1-targeting agents³⁵. In CRC specifically, a retrospective analysis of immune checkpoint inhibitor efficacy reported significantly improved OS for patients with tumors harboring TMB ≥ 9.8 Muts/MB compared with those with tumors with TMB < 9.8 Muts/Mb (~equivalency < 12 Muts/Mb as measured by this assay)³⁴. Another retrospective study reported that a TMB ≥ 12 Muts/Mb cutoff identifies $> 99\%$ of MSI-High CRC cases but only 3% of MSS cases, indicating the utility of this cutoff for identification of patients with CRC likely to benefit from treatment with immune checkpoint inhibitors⁵¹.

FREQUENCY & PROGNOSIS

Elevated tumor mutational burden (TMB) has been reported in 8-25% of colorectal cancer (CRC) samples^{21,52-53}. Multiple studies have reported that up to 90% of hypermutated CRC cases exhibit high levels of microsatellite instability (MSI-H) and mismatch repair deficiency (MMR-D)^{21,52}. Increased TMB is significantly associated with MSI-H and MMR-D, with studies reporting that 100% of MSI-H CRCs harbor elevated TMB and conversely that 100% of tumors with low TMB harbor intact MMR⁵². A subset of CRCs that harbor increased TMB but not MSI-H are driven by mutations in POLE, which leads to an "ultramutated" phenotype with especially high TMB^{21,52}. Tumors with increased TMB harbor BRAF V600E mutations more frequently than those with low TMB^{21,52}, whereas TMB-low tumors more frequently harbor mutations in TP53 and APC²¹. The prognostic value of tumor mutational burden (TMB) in colorectal cancer (CRC) is context- and therapy-dependent. A

study of tissue TMB (tTMB) in 145 CRC samples showed longer OS in TMB-high samples compared with TMB-low ones⁵⁴. Similarly, for patients with metastatic CRC treated with first-line chemotherapy combined with bevacizumab or cetuximab, high tissue TMB (tTMB-H) was associated with longer OS⁵⁵. For patients treated with adjuvant chemotherapy, tTMB-H was associated with better 5-year relapse-free survival⁵⁶. However, for patients with EGFR/BRAF-inhibitor-treated, BRAF-mutated microsatellite stable (MSS) metastatic CRC, intermediate tTMB was associated with significantly poorer PFS and OS compared with TMB-low status; patients with primary resistance to EGFR/BRAF blockage had higher TMB than those sensitive to these therapies⁵⁷. In a study for 61 patients with metastatic, MSS CRC treated with best standard of care, plasma TMB scores ≥ 28 Muts/Mb (approximately 14 Muts/Mb as measured by this assay) were associated with reduced OS compared with plasma TMB scores < 28 Muts/Mb (3.0 vs. 5.3 months, HR=0.76, $p=0.007$), whereas tTMB was not found to be prognostic in this population⁵⁸.

FINDING SUMMARY

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma⁵⁹⁻⁶⁰ and cigarette smoke in lung cancer⁶¹⁻⁶², treatment with temozolomide-based chemotherapy in glioma⁶³⁻⁶⁴, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes^{21,65-68}, and microsatellite instability (MSI)^{21,65,68}. This sample harbors a TMB below levels that would be predicted to be associated with sensitivity to PD-1- or PD-L1-targeting immune checkpoint inhibitors, alone or in combination with other agents^{34,44,51}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606031-01

GENOMIC FINDINGS

GENE

BRAF

ALTERATION

V600E

HGVS VARIANT

NM_004333.4: c.1799T>A (p.V600E)

VARIANT CHROMOSOMAL POSITION

chr7:140453136

VARIANT ALLELE FREQUENCY (% VAF)

25.9%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Significant benefit for patients with BRAF V600-mutated colorectal cancers (CRC) has been achieved with combinatorial approaches involving BRAF inhibitors, EGFR-targeting antibodies, and MEK inhibitors⁶⁹⁻⁷². In a Phase 3 study for patients with metastatic CRC on second- or third-line treatments, doublet therapy with the RAF inhibitor encorafenib and the EGFR antibody cetuximab showed superior mOS to cetuximab plus chemotherapy (9.3 vs. 5.9 months, HR=0.61, n=220 and n=221), and similar benefit was seen for a triplet therapy cohort adding the MEK inhibitor binimetinib (OS of 9.3 months, n=224)⁷³. Combinations of other RAF inhibitors such as dabrafenib or vemurafenib with EGFR antibodies such as panitumumab have also resulted in clinical benefit for similar patient populations in Phase 1 and 2 studies. A trial of dabrafenib and panitumumab with or without the MEK inhibitor trametinib reported a 21% ORR and 86% DCR (n=91) for the triplet combination and a 10% ORR and 90% DCR (n=20) for the doublet therapy⁶⁹. Multiple similar studies of vemurafenib with panitumumab or cetuximab doublet therapy have also reported a benefit⁷⁰⁻⁷¹. In a randomized Phase 2 study for patients with 0-4 previous lines of therapy, the addition of vemurafenib to cetuximab and irinotecan significantly improved ORR (17% vs. 4.2%, n=50 and n=50) and DCR (65% vs. 21%)⁷². A Phase 2 trial evaluating the investigational agent spartalizumab, an anti-PD-1 antibody, with dabrafenib and trametinib reported an ORR of 35%

(n=20) and DCR of 75%⁷⁴. Extensive clinical evidence supports a significant benefit in BRAF-inhibitor and MEK-inhibitor doublet therapy for patients with BRAF V600E-mutated metastatic CRC. A Phase 2 study of vemurafenib plus cobimetinib for patients with advanced BRAF V600E-mutated CRC reported an ORR of 29% (n=28) and DCR of 57%⁷⁵, and a similar trial of dabrafenib and trametinib reported a 12% ORR (n=43) and 67% DCR⁷⁶. A basket trial of the combination of encorafenib and binimetinib for patients with BRAF V600-mutated solid cancers elicited 1 PR and 1 SD for 3 patients with CRC⁷⁷. In 2 Phase 1 studies evaluating the MEK-pan-RAF dual inhibitor CH5126766, 3 patients harboring BRAF V600E mutations experienced PRs, including 2 patients with melanoma⁷⁸ and 1 patient with low-grade serous ovarian carcinoma⁷⁹. Based on clinical data in solid tumors, patients with tumors harboring BRAF V600 mutations may benefit from treatment with type-II RAF inhibitors such as tovorafenib, lifrafenib, and belvarafenib⁸⁰⁻⁸¹.

— Potential Resistance —

On the basis of extensive clinical data, BRAF V600 mutation does not generally associate with significant clinical benefit from addition of cetuximab or panitumumab to chemotherapy (NCCN Colon Cancer Guidelines, v3.2022)⁸²⁻⁹¹. Low response rates to cetuximab or panitumumab monotherapy or combination with chemotherapy have been frequently observed among patients with BRAF V600-mutated CRC, although similarly low response rates in this patient population were also often observed to chemotherapy alone; additionally, response rates were generally lower for patients with BRAF-mutated tumors than for those whose tumors were BRAF-wildtype^{84,87-88,91-94}. For a limited number of patients with CRC treated with cetuximab- or panitumumab-containing chemotherapy regimens, BRAF V600E was found to be present at the time of progression⁹⁵⁻¹⁰⁰, to be a mechanism of acquired¹⁰¹⁻¹⁰² or primary¹⁰³ resistance, or to be enriched in tumors of non-responders versus responders⁹⁸.

FREQUENCY & PROGNOSIS

BRAF mutations have been reported in approximately 5-19% of colorectal cancer samples^{92,104-107}. BRAF V600E is a strong adverse prognostic marker in colorectal cancer¹⁰⁸. BRAF mutations have been associated with poor prognosis and shorter survival for patients with colorectal cancer, particularly those with metastatic disease, as well as with smoking history^{12,84,86,109-115}. Analysis of individual BRAF mutations in 2127 patients with advanced colorectal cancer treated with chemotherapy with or without cetuximab revealed that BRAF V600E associated with poor prognosis (HR 2.60, P=1.0e-15, with median reduction of survival being 320 days) and distinct clinicopathological features, including correlation with increased peritoneal metastases compared to BRAF wild-type tumors (24% vs. 12%, P=0.0015), while BRAF D594G inactivating mutation was not prognostic (HR 1.30, P=0.37) and had similar clinicopathologic features as BRAF wild-type tumors¹¹⁶.

FINDING SUMMARY

BRAF encodes a member of the RAF family of protein kinases, which includes ARAF, BRAF, and CRAF. These kinases function downstream of RAS as part of the MAPK (RAF-MEK-ERK) signaling cascade that facilitates cell proliferation, survival and transformation¹¹⁷⁻¹¹⁸. BRAF mutations have been reported in up to 20% of all cancers, with the majority of mutations occurring at the V600 position¹¹⁹⁻¹²⁰. Among the V600 mutations, V600E accounts for 70-80% of observations, V600K for 10-30%, and V600R for 5-7%, with V600D comprising the majority of the rest^{119,121-122}. Mutations at V600 are Class 1 BRAF alterations that have been shown to constitutively activate BRAF kinase and hyperactivate the downstream MEK-ERK signaling, promoting oncogenic transformation^{119,123}. In multiple cancer types, multiple mutations at V600, including V600E, V600K, V600R, V600D, and V600M, exhibited sensitivity to V600-targeted therapies^{122,124-134}; other mutations at this position are predicted to behave similarly.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606031-01

GENOMIC FINDINGS
GENE
KRAS
ALTERATION
 wildtype

targeting antibodies cetuximab^{84,135-137} or panitumumab^{86,138-139} for patients with CRC. Therefore, these agents are indicated to treat patients with CRC lacking such mutations (NCCN Colon Cancer Guidelines, v3.2022, Rectal Cancer Guidelines, v4.2022).

with decreased metastasis, better clinicopathological features, and longer survival of patients with CRC^{141-144,148-149}.

FINDING SUMMARY

KRAS encodes a member of the RAS family of small GTPases. Activating mutations in RAS genes can cause uncontrolled cell proliferation and tumor formation¹⁵⁰⁻¹⁵¹. No alterations in KRAS were identified in this case.

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

Lack of mutations in KRAS or NRAS is associated with clinical benefit of treatment with EGFR-

FREQUENCY & PROGNOSIS

Approximately 50-65% of colorectal cancers (CRCs) have been reported to lack KRAS mutations^{104,140-147}. Numerous studies have reported that KRAS wild-type status is associated

GENE
NRAS
ALTERATION
 wildtype

targeting antibodies cetuximab^{84,135-137} or panitumumab^{86,138-139} for patients with CRC. Therefore, these agents are indicated to treat patients with CRC lacking such mutations (NCCN Colon Cancer Guidelines, v3.2022, Rectal Cancer Guidelines, v4.2022).

frequency of metastasis¹⁴⁷ and longer survival¹⁵⁷⁻¹⁵⁸ of patients with CRC.

FINDING SUMMARY

NRAS encodes a member of the RAS family of small GTPases that mediate transduction of growth signals. Activation of RAS signaling causes cell growth, differentiation, and survival by activating the RAF-MAPK-ERK, PI3K, and other pathways¹⁵⁰. No alterations in NRAS were identified in this case.

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

Lack of mutations in KRAS or NRAS is associated with clinical benefit of treatment with EGFR-

FREQUENCY & PROGNOSIS

The majority of colorectal cancers (CRCs) (91-98%) have been reported to lack NRAS mutations^{21,147,152-157}. NRAS wild-type status has been reported to be associated with decreased

GENE
BARD1
ALTERATION
 W218*

HGVS VARIANT
 NM_000465.2: c.654G>A (p.W218*)

VARIANT CHROMOSOMAL POSITION
 chr2:215645944

VARIANT ALLELE FREQUENCY (% VAF)
 21.7%

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

Clinical benefit from rucaparib has been observed in a patient with BARD1-mutated ovarian cancer¹⁵⁹. On the basis of preclinical evidence, tumors with BARD1 inactivation may be sensitive to PARP inhibitors¹⁶⁰⁻¹⁶³.

one study, loss of BARD1 expression was correlated with worse prognosis in colon cancer¹⁷⁰.

FINDING SUMMARY

BARD1 encodes the BRCA1-associated RING domain 1 protein, which is required for stabilization and nuclear localization of BRCA1 as well as formation of the E3 ubiquitin ligase¹⁷¹. The BARD1 ANK repeats and BRCT motifs play important roles in chromosome stability, and both these regions and the RING domain are necessary for homology-directed repair^{160,172-173}. Alterations such as seen here may disrupt BARD1 function or expression.

FREQUENCY & PROGNOSIS

BARD1 mutations have been found in 0.9-2.8% of colorectal adenocarcinomas and other colon cancers¹⁶⁴⁻¹⁶⁹. Published data investigating the prognostic implications of BARD1 alterations in colon cancer are limited (PubMed, Mar 2023). In

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1606031-01

GENOMIC FINDINGS
GENE
TET2
ALTERATION

E1151*

HGVS VARIANT

NM_017628.4: c.3451G>T (p.E1151*)

VARIANT CHROMOSOMAL POSITION

chr4:106158550

VARIANT ALLELE FREQUENCY (% VAF)

41.7%

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are no targeted therapies available to address genomic alterations in TET2 in solid tumors.

FREQUENCY & PROGNOSIS

TET2 alterations have been reported at relatively low frequencies in solid tumors and are more prevalent in hematological malignancies¹⁷⁴. Published data investigating the prognostic implications of TET2 alterations in solid tumors are limited (PubMed, Jan 2023).

FINDING SUMMARY

TET2 encodes a tumor suppressor involved in reversing DNA methylation marks, a process critical for proper gene regulation¹⁷⁵⁻¹⁷⁶. Alterations such as seen here may disrupt TET2 function or expression¹⁷⁷⁻¹⁸¹.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to

occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion¹⁸²⁻¹⁸⁷. CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy¹⁸²⁻¹⁸³. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease¹⁸⁸. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH^{186,189-190}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606031-01

GENOMIC FINDINGS

GENE

TP53

ALTERATION

C176*

HGVS VARIANT

NM_000546.4: c.528C>A (p.C176*)

VARIANT CHROMOSOMAL POSITION

chr17:7578402

VARIANT ALLELE FREQUENCY (% VAF)

21.0%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved therapies to address TP53 mutation or loss. However, tumors with TP53 loss of function alterations may be sensitive to the WEE1 inhibitor adavosertib¹⁹¹⁻¹⁹⁴ or p53 gene therapy such as SGT53¹⁹⁵⁻¹⁹⁹. In a Phase 1 study, adavosertib in combination with gemcitabine, cisplatin, or carboplatin elicited PRs in 9.7% and SDs in 53% of patients with solid tumors; the response rate was 21% (4/19) for patients with TP53 mutations versus 12% (4/33) for patients who were TP53 wildtype²⁰⁰. A Phase 2 trial of adavosertib in combination with chemotherapy (gemcitabine, carboplatin, paclitaxel, or doxorubicin) reported a 32% (30/94, 3 CR) ORR and a 73% (69/94) DCR for patients with platinum-refractory TP53-mutated ovarian, Fallopian tube, or peritoneal cancer²⁰¹. A smaller Phase 2 trial of adavosertib in combination with carboplatin achieved a 43% (9/21, 1 CR) ORR and a 76% (16/21) DCR for patients with platinum-refractory TP53-mutated ovarian cancer²⁰². The combination of adavosertib with paclitaxel and carboplatin for patients with TP53-mutated ovarian cancer also significantly increased PFS compared with paclitaxel and carboplatin alone²⁰³. In the Phase 2

VIKTORY trial, patients with TP53-mutated metastatic and/or recurrent gastric cancer experienced a 24% (6/25) ORR with adavosertib combined with paclitaxel²⁰⁴. A Phase 1 trial of neoadjuvant adavosertib in combination with cisplatin and docetaxel for head and neck squamous cell carcinoma (HNSCC) elicited a 71% (5/7) response rate for patients with TP53 alterations²⁰⁵. The Phase 2 FOCUS4-C trial for patients with TP53- and RAS-mutated colorectal cancer reported improvement in PFS (3.61 vs. 1.87 months, HR=0.35, p=0.0022), but not OS (14.0 vs 12.8 months, p=0.93), following adavosertib treatment compared with active monitoring²⁰⁶. In a Phase 1b clinical trial of SGT-53 in combination with docetaxel for patients with solid tumors, 75% (9/12) of evaluable patients experienced clinical benefit, including 2 confirmed and 1 unconfirmed PRs and 2 instances of SD with significant tumor shrinkage¹⁹⁹. Missense mutations leading to TP53 inactivation may be sensitive to therapies that reactivate mutated p53 such as eprenetapopt. In a Phase 1b trial for patients with p53-positive high-grade serous ovarian cancer, eprenetapopt combined with carboplatin and pegylated liposomal doxorubicin achieved a 52% (11/21) response rate and 100% DCR²⁰⁷. A Phase 1 trial of eprenetapopt with pembrolizumab for patients with solid tumors reported an ORR of 10% (3/29)²⁰⁸.

FREQUENCY & PROGNOSIS

TP53 mutations have been reported in up to 75% of colorectal cancer cases^{21,88,209-213}. A study reported p53 expression in 49% of analyzed colorectal cancer cases²¹⁴. TP53 mutation has not been consistently demonstrated to be a significant independent prognostic marker in the context of CRC²¹⁵.

FINDING SUMMARY

Functional loss of the tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers²¹⁶. Alterations such as seen here may disrupt TP53 function or expression²¹⁷⁻²²¹.

POTENTIAL GERMLINE IMPLICATIONS

Germline mutations in TP53 are associated with the very rare autosomal dominant disorder Li-Fraumeni syndrome and the early onset of many cancers²²²⁻²²⁴, including sarcomas²²⁵⁻²²⁶. Estimates for the prevalence of germline TP53 mutations in the general population range from 1:5,000²²⁷ to 1:20,000²²⁶. For pathogenic TP53 mutations identified during tumor sequencing, the rate of germline mutations was 1% in the overall population and 6% in tumors arising before age 30²²⁸. In the appropriate clinical context, germline testing of TP53 is recommended.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion¹⁸²⁻¹⁸⁷. CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy¹⁸²⁻¹⁸³. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease¹⁸⁸. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH^{186,189-190}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606031-01

THERAPIES WITH CLINICAL BENEFIT

IN PATIENT'S TUMOR TYPE

Encorafenib + Cetuximab

Assay findings association
BRAF
V600E

AREAS OF THERAPEUTIC USE

Encorafenib is an inhibitor of BRAF, and cetuximab is a monoclonal antibody that targets EGFR. The combination is FDA approved to treat patients with BRAF V600E-mutated colorectal cancer (CRC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

Patients with BRAF V600-mutated CRC are considered unlikely to benefit from cetuximab, alone or in combination with chemotherapy, unless combined with BRAF inhibitors (NCCN Colon Cancer Guidelines, v3.2022, NCCN Rectal Cancer Guidelines, v4.2022). Response rates to cetuximab, both as monotherapy and in combination with chemotherapy, have generally been found to be low for patients with BRAF V600-mutated CRC, independent of treatment line and chemotherapy backbone^{84,87,92-94,96,100,115,155,229-233}. However, significant clinical responses have been reported for patients with BRAF V600-mutated CRC treated with cetuximab in combination with the BRAF inhibitor vemurafenib⁷⁰, the 2 in combination with irinotecan²³⁴, or cetuximab in combination with BRAF inhibitor encorafenib²³⁵⁻²³⁶.

SUPPORTING DATA

For patients with BRAF V600E-mutated metastatic colorectal cancer (CRC), the Phase 3 BREAKWATER study evaluating encorafenib and cetuximab with mFOLFOX6 and/or FOLFIRI reported an ORR of 67-68% ([8/12]-[13/19]) for untreated patients and an ORR of 50-61% ([4/8]-[11/18]) for previously treated patients²³⁷. The Phase 3 BEACON study for previously treated patients with BRAF V600E-mutated metastatic CRC demonstrated significantly improved efficacy of encorafenib and cetuximab doublet therapy over standard irinotecan and cetuximab therapy (median OS [mOS] of 9.3 vs. 5.9 months, HR=0.61; median PFS [mPFS] of 4.3 vs. 1.5 months, HR=0.44; ORR of 20% vs. 1.8%)^{235,238}. In the same study, triplet therapy of encorafenib and cetuximab combined with the MEK inhibitor binimetinib resulted in similar efficacy as the doublet therapy, compared with standard therapy (mOS of 9.3 vs. 5.9 months, HR=0.60; mPFS of 4.5 vs. 1.5 months, HR=0.42; ORR of 27% vs. 1.8%)²³⁸. In a Phase 1/2 trial for patients with microsatellite stable and BRAF V600E-positive metastatic CRC, treatment with triplet encorafenib, cetuximab, and nivolumab resulted in an ORR of 50% (11/22 PRs), a DCR of 95% (21/22), an mPFS of 7.4 months, and an mOS of 15.1 months²³⁹.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606031-01

THERAPIES ASSOCIATED WITH RESISTANCE

IN PATIENT'S TUMOR TYPE

Cetuximab

✖ Resistance of variant(s) to associated therapy is likely

Assay findings association

BRAF

V600E

KRAS

wildtype

NRAS

wildtype

AREAS OF THERAPEUTIC USE

Cetuximab is a monoclonal antibody that targets EGFR. It is FDA approved for the treatment of head and neck squamous cell carcinoma (HNSCC) and KRAS-wild-type, EGFR-expressing metastatic colorectal cancer (CRC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

Therapies targeting EGFR, including cetuximab, have been shown to have significant clinical activity for patients with CRC^{84,135-137,240-241}; wild-type KRAS and NRAS are predictive biomarkers for the efficacy of cetuximab in metastatic CRC (NCCN Colon Cancer Guidelines, v3.2022, NCCN Rectal Cancer Guidelines, v4.2022). Patients with BRAF V600-mutated CRC are considered unlikely to benefit from cetuximab, alone or in combination with chemotherapy, unless combined with BRAF inhibitors (NCCN Colon Cancer Guidelines, v3.2022). Response rates to cetuximab, both as monotherapy and in combination with chemotherapy, have generally been found to be low for patients with BRAF V600-mutated CRC, independent of treatment line and chemotherapy backbone^{84,87,92-94,96,100,115,155,229-233}. However, significant clinical responses have been reported for patients with BRAF V600-mutated CRC treated with cetuximab in combination with the BRAF inhibitor vemurafenib⁷⁰, the 2 in combination with irinotecan²³⁴, or cetuximab in combination with BRAF inhibitor encorafenib²³⁵⁻²³⁶.

SUPPORTING DATA

Cetuximab has been shown to improve OS, PFS, and response rate for patients with KRAS-wildtype colorectal

cancer (CRC), both in combination with FOLFIRI, FOLFOX₄, or irinotecan^{84,135,240-242} and as monotherapy for chemotherapy-refractory patients^{87,137}. The Phase 3 study STRATEGIC-1 reported a similar duration of disease control (DDC) for patients with unresectable metastatic CRC (mCRC) and KRAS-, NRAS-, and BRAF-wildtype status treated with mFOLFOX-bevacizumab alternated with a cetuximab regimen in first or second line, respectively (overall DDC 22.5 vs. 23.5 months); in addition, the study reported similar OS (37.8 vs. 34.4 months) and higher numerical ORR for patients treated with cetuximab in the first line followed by mFOLFOX-bevacizumab compared with those receiving EGFR-directed antibodies in the second or third line²⁴³. A prospective study of cetuximab monotherapy for patients with KRAS-, NRAS-, and BRAF-wildtype mCRC reported 11% (2/19) PRs and 58% (11/19) SDs²⁴⁴. The Phase 2 AVETUX trial of cetuximab combined with avelumab and mFOLFOX6 for patients with RAS- and BRAF-wildtype mCRC resulted in an ORR of 81% (4 CR and 27 PRs, n=37) and a DCR of 89%²⁴⁵. In the Phase 3 ASPECCT study, panitumumab was found to be non-inferior to cetuximab with respect to median OS (10.4 vs. 10.0 months, HR=0.97) for patients with previously treated KRAS exon 2 wildtype metastatic colorectal cancer; median PFS was also similar between the two treatment groups (4.4 vs. 4.1 months, HR=1.00)²⁴⁶. In a similar patient population, a Phase 2 study of combination panitumumab and irinotecan versus combination cetuximab and irinotecan also demonstrated non-inferiority with respect to median PFS (5.4 vs. 4.3 months, HR = 0.64) and median OS (14.9 vs. 11.5 months, HR=0.66)²⁴⁷.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606031-01

THERAPIES ASSOCIATED WITH RESISTANCE

IN PATIENT'S TUMOR TYPE

Panitumumab

✖ Resistance of variant(s) to associated therapy is likely

Assay findings association

BRAF
V600E

KRAS
wildtype

NRAS
wildtype

AREAS OF THERAPEUTIC USE

Panitumumab is a monoclonal antibody that targets EGFR. It is FDA approved to treat KRAS wild-type and NRAS wild-type metastatic colorectal cancer (CRC) combined with chemotherapy or as monotherapy for patients who have progressed on prior chemotherapy. Please see the drug label for full prescribing information.

GENE ASSOCIATION

Therapies targeting EGFR, including panitumumab, have been shown to have significant clinical activity for patients with CRC^{138,246,248}; wild-type KRAS and NRAS are predictive biomarkers for the efficacy of panitumumab in metastatic CRC (NCCN Colon Cancer Guidelines v3.2022)(NCCN Rectal Cancers Guidelines, v4.2022). Patients with BRAF V600-mutated CRC are considered unlikely to benefit from panitumumab, alone or in combination with chemotherapy (NCCN Colon Cancer Guidelines, v3.2022)(NCCN Rectal Cancers Guidelines, v4.2022). Response rates to panitumumab, both as monotherapy and in combination with chemotherapy, have generally been found to be low for patients with BRAF V600-mutated CRC, independent of line of treatment and chemotherapy backbone^{88,91,95,99-100,155,231}. However, significant clinical responses have been reported for patients with BRAF V600E-mutated CRC upon treatment with panitumumab in combination with the BRAF inhibitor dabrafenib and the MEK inhibitor trametinib²⁴⁹.

SUPPORTING DATA

Panitumumab has been shown to improve OS, PFS, and ORR for patients with KRAS-wildtype colorectal cancer (CRC), both in combination with FOLFOX₄, FOLFIRI, irinotecan, or best supportive care^{89,138,250-252}, and as monotherapy for chemotherapy-refractory patients^{88,246,248}. The Phase 3 PARADIGM trial comparing panitumumab plus mFOLFOX6 versus bevacizumab plus mFOLFOX6 as first-line treatment for patients with RAS-wildtype left-sided metastatic CRC demonstrated that treatment with panitumumab significantly improved median OS (mOS; 36.2 months vs. 31.3 months) compared with bevacizumab²⁵³. A Phase 2 trial reported that, for patients with unresectable RAS-wildtype colorectal adenocarcinoma treated with panitumumab plus FOLFOX₄, maintenance with a combination of panitumumab plus fluorouracil and leucovorin was superior to panitumumab monotherapy (10-month PFS OF 59% vs. 49%)²⁵⁴. In the Phase 3 ASPECCT study, panitumumab was found to be non-inferior to cetuximab with respect to median OS (10.4 vs. 10.0 months, HR=0.97) for patients with previously treated KRAS exon 2 wildtype metastatic colorectal cancer; median PFS was also similar between the two treatment groups (4.4 vs. 4.1 months, HR=1.00)²⁴⁶. In a similar patient population, a Phase 2 study of combination panitumumab and irinotecan versus combination cetuximab and irinotecan also demonstrated non-inferiority with respect to median PFS (5.4 vs. 4.3 months, HR = 0.64) and median OS (14.9 vs. 11.5 months, HR=0.66)²⁴⁷.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606031-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Dabrafenib + Trametinib

Assay findings association
BRAF
V600E

AREAS OF THERAPEUTIC USE

Dabrafenib is a BRAF V600 selective inhibitor and trametinib is a MEK inhibitor. These 2 therapies are FDA approved in combination to treat metastatic non-small cell lung cancer (NSCLC) with BRAF V600E mutation, advanced anaplastic thyroid cancer (ATC) with BRAF V600E mutation, advanced solid tumors with BRAF V600E mutation in adult and pediatric patients 6 years of age and older, and low-grade glioma with BRAF V600E mutation in pediatric patients 1 year of age or older. This combination is also approved to treat patients with melanoma with BRAF V600E/K mutations. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical evidence in various solid tumors and hematologic malignancies, BRAF activating alterations may confer sensitivity to the combination of BRAF V600-targeted therapies and MEK inhibitors such as dabrafenib and trametinib^{76,255-265}.

SUPPORTING DATA

The combination of BRAF inhibitors with MEK inhibitors has shown clinical activity for patients with BRAF V600-mutated metastatic colorectal carcinoma (mCRC). A Phase 1/2 open-label trial combining dabrafenib and trametinib for BRAF V600-mutated mCRC reported an

ORR of 12% (5/43, including 1 CR with a response duration >36 months)⁷⁶. For patients with BRAF V600E-mutated mCRC, a combination of dabrafenib, trametinib, and the EGFR-targeting antibody panitumumab elicited a 21% (19/91) ORR, 86% (78/91) DCR, and 7.6-month estimated median duration of response (DoR)⁶⁹. A Phase 2 trial evaluating dabrafenib and trametinib in combination with the anti-PD-1 immune checkpoint inhibitor spartalizumab reported a 33% (7/21) ORR for patients with BRAF V600-mutated mCRC, with 5.6 months median DoR; within microsatellite stable patients, the ORR was 42% (5/12)⁷⁴. One case report describes a patient with colon adenocarcinoma who responded to a combination of dabrafenib, trametinib, and oxaliplatin²⁶⁶. Dabrafenib can induce adverse effects such as the development of cutaneous squamous cell carcinomas and keratoacanthomas caused by inactivation of wildtype BRAF that leads to paradoxical activation of the MAPK pathway, but it has been reported to be well tolerated in patients with BRAF V600E-mutated thyroid cancer^{124,267-268}. Patients with melanoma harboring BRAF V600E or V600K mutation treated with a combination of dabrafenib and trametinib experienced significantly lower rates of cutaneous squamous cell carcinoma and regression of established BRAF inhibitor-induced skin lesions^{256-257,269-271}.

Encorafenib + Binimetinib

Assay findings association
BRAF
V600E

AREAS OF THERAPEUTIC USE

The combination of the BRAF inhibitor encorafenib and MEK inhibitor binimetinib is FDA approved to treat patients with melanoma with BRAF V600E or BRAF V600K mutations. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical efficacy in the treatment of patients with BRAF V600-mutated melanoma²⁷²⁻²⁷⁵, and activity in colorectal, thyroid, and lung cancer²⁷⁵⁻²⁷⁷, activating alterations affecting BRAF predict sensitivity to the combination of encorafenib and binimetinib.

SUPPORTING DATA

A Phase 1/2 trial of encorafenib combined with binimetinib for patients with BRAF V600E- or BRAF V600K-mutated solid tumors reported an ORR of 18.2%

(2/11) for the subset of patients with metastatic colorectal cancer²⁷⁵. The combination of encorafenib and binimetinib has been reported to provide clinical benefit for patients with various solid tumors harboring BRAF V600 activating alterations^{272,275-277}, and has been studied primarily in the context of BRAF V600-mutated melanoma where patients treated with this combination achieved greater PFS and OS compared with encorafenib or vemurafenib monotherapy^{272-273,278}. A combination of encorafenib, binimetinib, and the CDK4/6 inhibitor ribociclib in a Phase 1b trial for patients with BRAF V600-mutant cancers elicited responses in melanoma, astrocytoma, unknown carcinoma, and in 1 of 3 patients with colorectal cancer; a Phase 2 study of this combination in V600-mutant melanoma reported an ORR of 52.4% (22/42), including 5 CRs, median PFS of 9.2 months, and median OS of 19.4 months⁷⁷.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1606031-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Vemurafenib + Cobimetinib

Assay findings association
BRAF
V600E

AREAS OF THERAPEUTIC USE

Vemurafenib is a selective inhibitor of BRAF with V600 mutations and cobimetinib is a MEK inhibitor. The combination is FDA approved to treat patients with melanoma with BRAF V600E or V600K mutations. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical evidence in melanoma and colorectal carcinoma, BRAF activating alterations may confer sensitivity to the combination of BRAF V600-targeted therapies and MEK inhibitors such as vemurafenib and cobimetinib^{75,279-280}.

SUPPORTING DATA

The Phase 2 TAPUR study of vemurafenib plus

cobimetinib for patients with advanced BRAF V600E-mutated CRC reported an ORR of 29% (8/28), median PFS of 15.8 weeks, and median OS of 38.9 weeks⁷⁵.

Vemurafenib can induce adverse effects, such as the development of cutaneous squamous cell carcinomas, keratoacanthomas, and new primary melanomas caused by inactivation of wildtype BRAF and leading to paradoxical activation of the MAPK pathway^{125,267}. In a Phase 1b trial, patients with BRAF V600E-mutated melanoma treated with a combination of vemurafenib and cobimetinib had increased RR (87%) and PFS (13.7 months) compared to the RR and PFS values previously reported for vemurafenib or MEK inhibitor monotherapy; this combination also resulted in lower rates of cutaneous SCC²⁸¹.

NOTE Genomic alterations detected may be associated with activity of certain FDA approved drugs, however, the agents listed in this report may have varied evidence in the patient's tumor type.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606031-01

CLINICAL TRIALS

NOTE Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually

updated and should be investigated by the physician or research staff. This is not a comprehensive list of all available clinical trials. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials listed here may have additional enrollment criteria that

may require medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see [clinicaltrials.gov](https://www.clinicaltrials.gov). Or, visit <https://www.foundationmedicine.com/genomic-testing#support-services>.

GENE
BARD1
ALTERATION
W218*
RATIONALE

Tumors with BARD1 inactivating mutation or loss may be sensitive to PARP inhibitors.

NCT04123366
PHASE 2

Study of Olaparib (MK-7339) in Combination With Pembrolizumab (MK-3475) in the Treatment of Homologous Recombination Repair Mutation (HRRm) and/or Homologous Recombination Deficiency (HRD)-Positive Advanced Cancer (MK-7339-007/KEYLYNK-007)

TARGETS
 PARP, PD-1

LOCATIONS: Fukuoka (Japan), Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Okayama (Japan), Nagoya (Japan), Tokyo (Japan), Kashiwa (Japan), Sapporo (Japan), Nedlands (Australia), Southport (Australia)

NCT03742895
PHASE 2

Efficacy and Safety of Olaparib (MK-7339) in Participants With Previously Treated, Homologous Recombination Repair Mutation (HRRm) or Homologous Recombination Deficiency (HRD) Positive Advanced Cancer (MK-7339-002 / LYNK-002)

TARGETS
 PARP

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Darlinghurst (Australia), Saint Petersburg (Russian Federation), Adana (Turkey), Jerusalem (Israel), Konya (Turkey), Ramat Gan (Israel), Istanbul (Turkey), Antalya (Turkey)

NCT02264678
PHASE 1/2

Ascending Doses of AZD6738 in Combination With Chemotherapy and/or Novel Anti Cancer Agents

TARGETS
 ATR, PARP, PD-L1

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Goyang-si (Korea, Republic of), Cambridge (United Kingdom), Withington (United Kingdom), Manchester (United Kingdom), London (United Kingdom), Coventry (United Kingdom), Sutton (United Kingdom), Oxford (United Kingdom)

NCT05035745
PHASE 1/2

Selinexor & Talazoparib in Advanced Refractory Solid Tumors; Advanced/Metastatic Triple Negative Breast Cancer (START)

TARGETS
 XPO1, PARP

LOCATIONS: Singapore (Singapore)

NCT03772561
PHASE 1

Phase I Study of AZD5363 + Olaparib + Durvalumab in Patients With Advanced or Metastatic Solid Tumor Malignancies

TARGETS
 PARP, AKTs, PD-L1

LOCATIONS: Singapore (Singapore)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1606031-01

CLINICAL TRIALS
NCT04801966
PHASE NULL

Safety and Oversight of the Individually Tailored Treatment Approach: A Novel Pilot Study

TARGETS
 CDK4, CDK6, PI3K-alpha, PD-L1, MEK, PARP, PD-1, BRAF

LOCATIONS: Melbourne (Australia)

NCT03127215
PHASE 2

Study of Olaparib/Trabectedin vs. Doctor's Choice in Solid Tumors

TARGETS
 FUS-DDIT3, PARP

LOCATIONS: Dresden (Germany), München (Germany), Frankfurt (Germany), Essen (Germany), Mainz (Germany), Heidelberg (Germany), Stuttgart (Germany), Tuebingen (Germany), Freiburg (Germany)

NCT03297606
PHASE 2

Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR)

TARGETS
 VEGFRs, ABL, SRC, ALK, ROS1, AXL, TRKA, MET, TRKC, DDR2, KIT, EGFR, PD-1, CTLA-4, PARP, CDK4, CDK6, FLT3, CSF1R, RET, mTOR, ERBB2, MEK, BRAF, SMO

LOCATIONS: Vancouver (Canada), Kelowna (Canada), Edmonton (Canada), Saskatoon (Canada), Regina (Canada), Ottawa (Canada), Montreal (Canada), Toronto (Canada), Kingston (Canada), London (Canada)

NCT04991480
PHASE 1/2

A Study of ART4215 for the Treatment of Advanced or Metastatic Solid Tumors

TARGETS
 PARP, Pol theta

LOCATIONS: London (United Kingdom), Oklahoma, Connecticut, New York, Pennsylvania, Tennessee, Texas, Florida

NCT05327010
PHASE 2

Testing the Combination of the Anti-cancer Drugs ZEN003694 (ZEN-3694) and Talazoparib in Patients With Advanced Solid Tumors, The ComBET Trial

TARGETS
 PARP, BRD4, BRDT, BRD2, BRD3

LOCATIONS: Illinois, Texas, North Carolina, Georgia

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 17 April 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606031-01

CLINICAL TRIALS
GENE
BRAF
ALTERATION
V600E
RATIONALE

BRAF activating alterations may predict sensitivity to inhibitors of BRAF, MEK, or ERK. Limited clinical and preclinical studies indicate

BRAF mutations may predict sensitivity to MEK-pan-RAF dual inhibitors.

NCT04607421
PHASE 3

BRAF V600E-mutant Colorectal Cancer Study of Encorafenib Taken With Cetuximab Plus or Minus Chemotherapy (BREAKWATER)

TARGETS

VEGFA, MEK, BRAF, EGFR

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Taichung (Taiwan), Fuzhou (China), Tainan (Taiwan), Kaohsiung (Taiwan), Hangzhou (China), Shanghai (China), Busan (Korea, Republic of), Nanjing (China)

NCT04913285
PHASE 1

A Study to Evaluate KIN-2787 in Subjects With BRAF Mutation Positive Solid Tumors

TARGETS

BRAF, MEK

LOCATIONS: Taipei (Taiwan), Shanghai (China), Gyeonggi-do (Korea, Republic of), Cheongju-si (Korea, Republic of), Incheon (Korea, Republic of), Seoul (Korea, Republic of), Perth (Australia), Wollstonecraft (Australia), Amsterdam (Netherlands), Villejuif (France)

NCT05004350
PHASE 2

A Study Evaluating the Combination of Encorafenib and Cetuximab Versus Irinotecan/Cetuximab or Infusional 5-fluorouracil (5-FU)/Folinic Acid (FA)/Irinotecan (FOLFIRI)/Cetuximab in Chinese Patients With BRAF V600E Mutant Metastatic Colorectal Cancer.

TARGETS

MEK, BRAF, EGFR

LOCATIONS: Fuzhou (China), Xiamen (China), Shantou (China), Hangzhou (China), Ganzhou (China), Shanghai (China), Nanchang (China), Changzhou (China), Shenzhen (China), Guangzhou (China)

NCT04984369
PHASE 2

The Efficacy of HLX208 (BRAF V600E Inhibitor) With Cetuximab for Metastatic Colorectal Cancer (mCRC) With BRAF V600E Mutation After First-line Treatment

TARGETS

EGFR, BRAF

LOCATIONS: Shanghai (China)

NCT03727763
PHASE 2

FIVC in Advanced Colorectal Cancer Patients With BRAF V600E Mutation.

TARGETS

EGFR, BRAF

LOCATIONS: Shanghai (China)

NCT03781219
PHASE 1

A Phase I Study of HL-085 Plus Vemurafenib in Solid Tumor With BRAF V600 Mutation

TARGETS

MEK, BRAF

LOCATIONS: Hangzhou (China), Zhengzhou (China), Beijing (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 17 April 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606031-01

CLINICAL TRIALS
NCT04803318
PHASE 2

Trametinib Combined With Everolimus and Lenvatinib for Recurrent/Refractory Advanced Solid Tumors

TARGETS
 mTOR, FGFRs, RET, PDGFRA, VEGFRs, KIT, MEK

LOCATIONS: Guangzhou (China)

NCT04985604
PHASE 1/2

DAY101 Monotherapy or in Combination With Other Therapies for Patients With Solid Tumors

TARGETS
 BRAF, MEK

LOCATIONS: Busan (Korea, Republic of), Seoul (Korea, Republic of), Clayton (Australia), Edegem (Belgium), Oregon, Barcelona (Spain), Madrid (Spain), California, Colorado

NCT03284502
PHASE 1

Cobimetinib and HM95573 in Patients With Locally Advanced or Metastatic Solid Tumors

TARGETS
 MEK, RAFs, NRAS

LOCATIONS: Hwasun (Korea, Republic of), Pusan (Korea, Republic of), Seongnam (Korea, Republic of), Seoul (Korea, Republic of), Goyang-si (Korea, Republic of)

NCT04999761
PHASE 1

AB122 Platform Study

TARGETS
 PD-1, HSP90, FGFRs

LOCATIONS: Ehime (Japan), Wakayama (Japan), Osaka (Japan), Aichi (Japan), Shizuoka (Japan), Kanagawa (Japan), Tokyo (Japan), Chiba (Japan), Hokkaido (Japan)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 17 April 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606031-01

APPENDIX
Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

CD22

 NM_001771.3: c.148G>A
(p.D50N)
chr19:35823563

CXCR4

 NM_001008540.1: c.803T>C
(p.F268S)
chr2:136872707

EPHB4

 NM_004444.4: c.1555G>A
(p.G519S)
chr7:100414847

FLCN

 NM_144997.5: c.1265C>T
(p.P422L)
chr17:17119729

GNAQ

 NM_002072.3: c.695T>G
(p.L232R)
chr9:80409419

MDM2

 NM_002392.3: c.91G>C
(p.E31Q)
chr12:69203064

MSH3

 NM_002439.3: c.1368A>C
(p.E456D)
chr5:80021299

MSH6

 NM_000179.2: c.1701G>C
(p.K567N)
chr2:48026823

PALB2

 NM_024675.3: c.1955G>A
(p.S652N)
chr16:23641520

PDGFRB

 NM_002609.3: c.1315C>T
(p.R439W)
chr5:149510154

ROS1

 NM_002944.2: c.977C>G
(p.T326R)
chr6:117715781

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 17 April 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606031-01

APPENDIX

Genes Assayed in FoundationOne®CDx

FoundationOne CDx is designed to include genes known to be somatically altered in human solid tumors that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay interrogates 324 genes as well as introns of 36 genes involved in rearrangements. The assay will be updated periodically to reflect new knowledge about cancer biology.

DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS

ABL1	ACVR1B	AKT1	AKT2	AKT3	ALK	ALOX12B	AMER1 (FAM123B or WTX)	
APC	AR	ARAF	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX
AURKA	AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2
BCL6	BCOR	BCORL1	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1
BTG2	BTK	CALR	CARD11	CASP8	CBFB	CBL	CCND1	CCND2
CCND3	CCNE1	CD22	CD274 (PD-L1)	CD70	CD79A	CD79B	CDC73	CDH1
CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B	CDKN2C
CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R	CTCF
CTNNA1	CTNNB1	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1	DDR2
DIS3	DNMT3A	DOT1L	EED	EGFR	EMSY (C11orf30)	EP300	EPHA3	EPHB1
EPHB4	ERBB2	ERBB3	ERBB4	ERCC4	ERG	ERRF1	ESR1	EZH2
FANCA	FANCC	FANCG	FANCL	FAS	FBXW7	FGF10	FGF12	FGF14
FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1	FGFR2	FGFR3	FGFR4
FH	FLCN	FLT1	FLT3	FOXL2	FUBP1	GABRA6	GATA3	GATA4
GATA6	GID4 (C17orf39)	GNA11	GNA13	GNAQ	GNAS	GRM3	GSK3B	H3-3A (H3F3A)
HDAC1	HGF	HNFI1A	HRAS	HSD3B1	ID3	IDH1	IDH2	IGF1R
IKBKE	IKZF1	INPP4B	IRF2	IRF4	IRS2	JAK1	JAK2	JAK3
JUN	KDM5A	KDM5C	KDM6A	KDR	KEAP1	KEL	KIT	KLHL6
KMT2A (MLL)	KMT2D (MLL2)	KRAS	LTK	LYN	MAF	MAP2K1 (MEK1)	MAP2K2 (MEK2)	MAP2K4
MAP3K1	MAP3K13	MAPK1	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1
MERTK	MET	MITF	MKNK1	MLH1	MPL	MRE11 (MRE11A)	MSH2	MSH3
MSH6	MST1R	MTAP	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN	MYD88
NBN	NF1	NF2	NFE2L2	NFKBIA	NKX2-1	NOTCH1	NOTCH2	NOTCH3
NPM1	NRAS	NSD2 (WHSC1 or MMSET)	NSD3 (WHSC1L1)	NT5C2	NTRK1	NTRK2	NTRK3	NTRK3
P2RY8	PALB2	PARP1	PARP2	PARP3	PAX5	PBRM1	PDCCD1 (PD-1)	PDCCD1LG2 (PD-L2)
PDGFRA	PDGFRB	PDK1	PIK3C2B	PIK3C2G	PIK3CA	PIK3CB	PIK3R1	PIM1
PMS2	POLD1	POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI
PRKN (PARK2)	PTCH1	PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51
RAD51B	RAD51C	RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10
REL	RET	RICTOR	RNF43	ROS1	RPTOR	SDHA	SDHB	SDHC
SDHD	SETD2	SF3B1	SGK1	SMAD2	SMAD4	SMARCA4	SMARCB1	SMO
SNCAIP	SOC1	SOX2	SOX9	SPEN	SPOP	SRC	STAG2	STAT3
STK11	SUFU	SYK	TBX3	TEK	TENT5C (FAM46C)	TET2	TET2	TGFB2
TIPARP	TNFAIP3	TNFRSF14	TP53	TSC1	TSC2	TYRO3	U2AF1	VEGFA
VHL	WT1	XPO1	XRCC2	ZNF217	ZNF703			

DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS

ALK	BCL2	BCR	BRAF	BRCA1	BRCA2	CD74	EGFR	ETV4
ETV5	ETV6	EWSR1	EZR	FGFR1	FGFR2	FGFR3	KIT	KMT2A (MLL)
MSH2	MYB	MYC	NOTCH2	NTRK1	NTRK2	NUTM1	PDGFRA	RAF1
RARA	RET	ROS1	RSP02	SDC4	SLC34A2	TERC*	TERT**	TPRSS2

*TERC is an NCRNA

**Promoter region of TERT is interrogated

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS


Homologous Recombination status
Loss of Heterozygosity (LOH) score
Microsatellite (MS) status
Tumor Mutational Burden (TMB)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1606031-01

APPENDIX

About FoundationOne®CDx

FoundationOne CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a., Ciplstraat 3, 2440 Geel, Belgium. 

ABOUT FOUNDATIONONE CDx

FoundationOne CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details:
www.rochefoundationmedicine.com/f1cdxtech.

INTENDED USE

FoundationOne®CDx (F1CDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI), tumor mutational burden (TMB), and for selected forms of ovarian cancer, loss of heterozygosity (LOH) score, using DNA isolated from formalin-fixed, paraffin-embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with therapies in accordance with approved therapeutic product labeling. Additionally, F1CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms.

TEST PRINCIPLE

FoundationOne CDx will be performed exclusively as a laboratory service using DNA extracted from formalin-fixed, paraffin-embedded (FFPE) tumor samples. The proposed assay will employ a single DNA extraction method from routine FFPE biopsy or surgical resection specimens, 50-1000 ng of which will undergo whole-genome shotgun library construction and hybridization-based capture of all coding exons from 309 cancer-related genes, one promoter region, one non-coding (ncRNA), and select intronic regions from 34 commonly rearranged genes, 21 of which also include the coding exons. The assay therefore includes

detection of alterations in a total of 324 genes.

Using an Illumina® HiSeq platform, hybrid capture-selected libraries will be sequenced to high uniform depth (targeting >500X median coverage with >99% of exons at coverage >100X). Sequence data will be processed using a customized analysis pipeline designed to accurately detect all classes of genomic alterations, including base substitutions, indels, focal copy number amplifications, homozygous gene deletions, and selected genomic rearrangements (e.g., gene fusions). Additionally, genomic signatures including loss of heterozygosity (LOH), microsatellite instability (MSI) and tumor mutational burden (TMB) will be reported.

THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. The F1CDx report may be used as an aid to inform molecular eligibility for clinical trials. Note: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

Diagnostic Significance

FoundationOne CDx identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

Qualified Alteration Calls (Equivocal and Subclonal)

An alteration denoted as "amplification – equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne CDx for identifying a copy number amplification is four (4) for *ERBB2* and six (6) for all other genes. Conversely, an alteration denoted as "loss – equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is one that the FoundationOne CDx analytical

methodology has identified as being present in <10% of the assayed tumor DNA.

Ranking of Therapies and Clinical Trials

Ranking of Therapies in Summary Table

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

Ranking of Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) (www.nccn.org). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2023. All rights reserved. To view the most recent and complete version of the guidelines, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

Limitations

1. In the fraction-based MSI algorithm, a tumor specimen will be categorized as MSI-H, MSS, or MS-Equivocal according to the fraction of microsatellite loci determined to be altered or unstable (i.e., the fraction unstable loci score). In the F1CDx assay, MSI is evaluated based on a genome-wide analysis across >2000 microsatellite loci. For a given microsatellite locus, non-somatic alleles are discarded, and the microsatellite is categorized as unstable if remaining alleles differ from the reference genome. The final fraction unstable loci score is calculated as the number of unstable microsatellite loci divided by the number of evaluable microsatellite loci. The MSI-H and MSS cut-off thresholds were determined by

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606031-01

APPENDIX

About FoundationOne®CDx

- analytical concordance to a PCR comparator assay using a pan-tumor FFPE tissue sample set. Patients with results categorized as "MS-Stable" with median exon coverage <300X, "MS-Equivocal," or "Cannot Be Determined" should receive confirmatory testing using a validated orthogonal (alternative) method.
- TMB by F1CDx is determined by counting all synonymous and non-synonymous variants present at 5% allele frequency or greater (after filtering) and the total number is reported as mutations per megabase (mut/Mb) unit. Observed TMB is dependent on characteristics of the specific tumor focus tested for a patient (e.g., primary vs. metastatic, tumor content) and the testing platform used for the detection; therefore, observed TMB results may vary between different specimens for the same patient and between detection methodologies employed on the same sample. The TMB calculation may differ from TMB calculations used by other assays depending on variables such as the amount of genome interrogated, percentage of tumor, assay limit of detection (LoD), filtering of alterations included in the score, and the read depth and other bioinformatic test specifications. Refer to the SSED for a detailed description of these variables in FMI's TMB calculation https://www.accessdata.fda.gov/cdrh_docs/pdf17/P170019B.pdf. The clinical validity of TMB defined by this panel has been established for TMB as a qualitative output for a cut-off of 10 mutations per megabase but has not been established for TMB as a quantitative score.
 - Homologous Recombination status may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas (Coleman et al., 2017; 28916367). Samples with deleterious *BRCA1/2* alteration and/or Loss of Heterozygosity (LOH) score $\geq 16\%$ will be reported as "HRD Positive" and samples with absence of these findings will be reported as "HRD Not Detected," agnostic of potential secondary *BRCA1/2* reversion alterations. Certain potentially deleterious missense or small in-frame deletions in *BRCA1/2* may not be classified as deleterious and, in the absence of an elevated LOH profile, samples with such mutations may be classified as "HRD Not Detected." A result of "HRD Not Detected" does not rule out the presence of a *BRCA1/2* alteration or an elevated LOH profile outside the assay performance characteristic limitations.
 - The LOH score is determined by analyzing SNPs spaced at 1Mb intervals across the genome on the FoundationOne CDx test and

- extrapolating an LOH profile, excluding arm- and chromosome-wide LOH segments. Detection of LOH has been verified only for ovarian cancer patients, and the LOH score result may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas. The LOH score will be reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine LOH. Performance of the LOH classification has not been established for samples below 35% tumor content. There may be potential interference of ethanol with LOH detection. The interfering effects of xylene, hemoglobin, and triglycerides on the LOH score have not been demonstrated.
- Alterations reported may include somatic (not inherited) or germline (inherited) alterations; however, the test does not distinguish between germline and somatic alterations. The test does not provide information about susceptibility.
 - Biopsy may pose a risk to the patient when archival tissue is not available for use with the assay. The patient's physician should determine whether the patient is a candidate for biopsy.
 - Reflex testing to an alternative FDA approved companion diagnostic should be performed for patients who have an *ERBB2* amplification result detected with copy number equal to 4 (baseline ploidy of tumor +2) for confirmatory testing. While this result is considered negative by FoundationOne®CDx (F1CDx), in a clinical concordance study with an FDA approved FISH test, 70% (7 out of 10 samples) were positive, and 30% (3 out of 10 samples) were negative by the FISH test with an average ratio of 2.3. The frequency of *ERBB2* copy number 4 in breast cancer is estimated to be approximately 2%. Multiple references listed in <https://www.mycancergenome.org/content/disease/breast-cancer/ERBB2/238/> report the frequency of *HER2* overexpression as 20% in breast cancer. Based on the F1CDx *HER2* CDx concordance study, approximately 10% of *HER2* amplified samples had copy number 4. Thus, total frequency is conservatively estimated to be approximately 2%.

REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic, nontargeted treatment, germline, or clonal

hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

VARIANT ALLELE FREQUENCY

Variant Allele Frequency (VAF) represents the fraction of sequencing reads in which the variant is observed. This attribute is not taken into account for therapy inclusion, clinical trial matching, or interpretive content. Caution is recommended in interpreting VAF to indicate the potential germline or somatic origin of an alteration, recognizing that tumor fraction and tumor ploidy of samples may vary.

Precision of VAF for base substitutions and indels

BASE SUBSTITUTIONS	%CV*
Repeatability	5.11 - 10.40
Reproducibility	5.95 - 12.31
INDELS	%CV*
Repeatability	6.29 - 10.00
Reproducibility	7.33 - 11.71

*Interquartile Range = 1st Quartile to 3rd Quartile

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of $>10\%$, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM*, *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *CHEK2*, *FH*, *FLCN*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *PALB2*, *PMS2*, *POLE*, *RAD51C*, *RAD51D*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *TSC2*, and *VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606031-01

APPENDIX

About FoundationOne®CDx

tumor sequencing is germline or somatic.
Interpretation should be based on clinical context.

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1*, *CBL*, *DNMT3A*, *IDH2*, *JAK2*, *KMT2D (MLL2)*, *MPL*, *MYD88*, *SF3B1*, *TET2*, and *U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne CDx.

TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking

into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. FoundationOne CDx is performed using DNA derived from tumor, and as such germline events may not be reported.

SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
mut/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

REFERENCE SEQUENCE INFORMATION

Sequence data is mapped to the human genome, Genome Reference Consortium Human Build 37 (GRCh37), also known as hg19.

MR Suite Version (RG) 7.7.0

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of the data. The median exon coverage for this sample is 848x. The suitability of use.

ORDERED TEST # **ORD-1606031-01**
APPENDIX References

1. Gatalica Z, et al. Cancer Epidemiol. Biomarkers Prev. (2014) PMID: 25392179
2. Kroemer G, et al. Oncoimmunology (2015) PMID: 26140250
3. Lal N, et al. Oncoimmunology (2015) PMID: 25949894
4. Le DT, et al. N. Engl. J. Med. (2015) PMID: 26028255
5. Ayers et al., 2016; ASCO-SITC Abstract P60
6. Ciardiello et al., 2018; ESMO Abstract LBA-004
7. Parikh et al., 2021; DOI: 10.1038/s43018-021-00269-7
8. Fukuoka S, et al. J. Clin. Oncol. (2020) PMID: 32343640
9. Kim et al., 2020; DOI: 10.1016/j.annonc.2020.04.073
10. Zhang Y, et al. BMC Gastroenterol (2021) PMID: 34688262
11. Sinicrope FA, et al. J. Clin. Oncol. (2013) PMID: 24019539
12. Gavin PG, et al. Clin. Cancer Res. (2012) PMID: 23045248
13. Bertagnolli MM, et al. J. Clin. Oncol. (2009) PMID: 19273709
14. Van Cutsem E, et al. J. Clin. Oncol. (2009) PMID: 19451425
15. Ribic CM, et al. N. Engl. J. Med. (2003) PMID: 12867608
16. Sargent DJ, et al. J. Clin. Oncol. (2010) PMID: 20498393
17. Fallik D, et al. Cancer Res. (2003) PMID: 14522894
18. Guastadisegni C, et al. Eur. J. Cancer (2010) PMID: 20627535
19. Pawlik TM, et al. Dis. Markers (2004) PMID: 15528785
20. Kocarnik JM, et al. Gastroenterol Rep (Oxf) (2015) PMID: 26337942
21. Nature (2012) PMID: 22810696
22. Histopathology (2007) PMID: 17204026
23. Samowitz WS, et al. Cancer Epidemiol. Biomarkers Prev. (2001) PMID: 11535541
24. Elsaleh H, et al. Clin Colorectal Cancer (2001) PMID: 12445368
25. Brueckl WM, et al. Anticancer Res. () PMID: 12820457
26. Guidoboni M, et al. Am. J. Pathol. (2001) PMID: 11438476
27. Gryfe R, et al. N. Engl. J. Med. (2000) PMID: 10631274
28. Sinicrope FA, et al. Gastroenterology (2006) PMID: 16952542
29. Laghi L, et al. Dig Dis (2012) PMID: 22722556
30. You JF, et al. Br. J. Cancer (2010) PMID: 21081928
31. Bairwa NK, et al. Methods Mol. Biol. (2014) PMID: 24623249
32. Boland CR, et al. Cancer Res. (1998) PMID: 9823339
33. Boland CR, et al. Gastroenterology (2010) PMID: 20420947
34. Samstein RM, et al. Nat. Genet. (2019) PMID: 30643254
35. Goodman AM, et al. Mol. Cancer Ther. (2017) PMID: 28835386
36. Goodman AM, et al. Cancer Immunol Res (2019) PMID: 31405947
37. Cristescu R, et al. Science (2018) PMID: 30309915
38. Ready N, et al. J. Clin. Oncol. (2019) PMID: 30785829
39. Hellmann MD, et al. N. Engl. J. Med. (2018) PMID: 29658845
40. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
41. Hellmann MD, et al. Cancer Cell (2018) PMID: 29731394
42. Rozeman EA, et al. Nat Med (2021) PMID: 33558721
43. Sharma P, et al. Cancer Cell (2020) PMID: 32916128
44. Marabelle A, et al. Lancet Oncol. (2020) PMID: 32919526
45. Ott PA, et al. J. Clin. Oncol. (2019) PMID: 30557521
46. Cristescu R, et al. J Immunother Cancer (2022) PMID: 35101941
47. Friedman CF, et al. Cancer Discov (2022) PMID: 34876409
48. Sturgill EG, et al. Oncologist (2022) PMID: 35274716
49. Schenker et al., 2022; AACR Abstract 7845
50. Legrand et al., 2018; ASCO Abstract 12000
51. Fabrizio DA, et al. J Gastrointest Oncol (2018) PMID: 30151257
52. Stadler ZK, et al. J. Clin. Oncol. (2016) PMID: 27022117
53. Shao C, et al. JAMA Netw Open (2020) PMID: 33119110
54. Schwartz et al., 2018; ASCO Abstract 572
55. Innocenti F, et al. J Clin Oncol (2019) PMID: 30865548
56. Lee DW, et al. Clin Cancer Res (2019) PMID: 31285374
57. Randon G, et al. Eur J Cancer (2022) PMID: 34933155
58. Chen EX, et al. JAMA Oncol (2020) PMID: 32379280
59. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
60. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
61. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
62. Rizvi NA, et al. Science (2015) PMID: 25765070
63. Johnson BE, et al. Science (2014) PMID: 24336570
64. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
65. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
66. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
67. Heitzer E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
68. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
69. Corcoran RB, et al. Cancer Discov (2018) PMID: 29431699
70. Hyman DM, et al. N. Engl. J. Med. (2015) PMID: 26287849
71. Yaeger R, et al. Clin. Cancer Res. (2015) PMID: 25589621
72. Kopetz S, et al. J Clin Oncol (2021) PMID: 33356422
73. Kopetz et al., 2020; ASCO Abstract 4001
74. Corcoran et al., 2020; ESMO GI Abstract SO-26
75. Klute et al., 2020; ASCO Abstract 122
76. Corcoran RB, et al. J. Clin. Oncol. (2015) PMID: 26392102
77. Ascierto et al., 2017; ASCO Abstract 9518
78. Martinez-Garcia M, et al. Clin. Cancer Res. (2012) PMID: 22761467
79. Guo C, et al. Lancet Oncol (2020) PMID: 33128873
80. Desai J, et al. J Clin Oncol (2020) PMID: 32182156
81. Yen I, et al. Nature (2021) PMID: 33953400
82. Pietrantonio F, et al. Eur. J. Cancer (2015) PMID: 25673558
83. Rowland A, et al. Br. J. Cancer (2015) PMID: 25989278
84. Van Cutsem E, et al. J. Clin. Oncol. (2011) PMID: 21502544
85. Smith CG, et al. Clin. Cancer Res. (2013) PMID: 23741067
86. Douillard JY, et al. N. Engl. J. Med. (2013) PMID: 24024839
87. Karapetis CS, et al. Clin. Cancer Res. (2014) PMID: 24218517
88. Peeters M, et al. Clin. Cancer Res. (2013) PMID: 23325582
89. Peeters M, et al. Clin. Cancer Res. (2015) PMID: 26341920
90. Guren TK, et al. Br. J. Cancer (2017) PMID: 28399112
91. Seymour MT, et al. Lancet Oncol. (2013) PMID: 23725851
92. Di Nicolantonio F, et al. J. Clin. Oncol. (2008) PMID: 19001320
93. Stintzing S, et al. Eur. J. Cancer (2017) PMID: 28463756
94. Tol J, et al. N. Engl. J. Med. (2009) PMID: 19571295
95. Freeman DJ, et al. Clin Colorectal Cancer (2008) PMID: 18621636
96. Gao J, et al. Chin. J. Cancer Res. (2011) PMID: 23357879
97. Soeda H, et al. Int. J. Clin. Oncol. (2013) PMID: 22638623
98. Molinari F, et al. Clin. Cancer Res. (2011) PMID: 21632860
99. André T, et al. Ann. Oncol. (2013) PMID: 23041588
100. Benvenuti S, et al. Cancer Res. (2007) PMID: 17363584
101. Arena S, et al. Clin. Cancer Res. (2015) PMID: 25623215
102. Montagut C, et al. Nat. Med. (2012) PMID: 22270724
103. Toledo RA, et al. Oncotarget (2017) PMID: 27852040
104. De Roock W, et al. Lancet Oncol. (2011) PMID: 21163703
105. Dienstmann R, et al. Mol. Cancer Ther. (2012) PMID: 22723336
106. Safaee Ardekani G, et al. PLoS ONE (2012) PMID: 23056577
107. Guedes JG, et al. BMC Cancer (2013) PMID: 23548132
108. Cervantes A, et al. Ann Oncol (2023) PMID: 36307056
109. Sinicrope et al., 2012; ASCO Abstract 3514
110. Hassabo et al., 2014; ASCO Gastrointestinal Cancers Symposium Abstract 473
111. Bokemeyer C, et al. Eur. J. Cancer (2012) PMID: 22446022
112. Laurent-Puig P, et al. J. Clin. Oncol. (2009) PMID: 19884556
113. Ogino S, et al. Clin. Cancer Res. (2012) PMID: 22147942
114. Roth AD, et al. J. Clin. Oncol. (2010) PMID: 20008640
115. Hsu HC, et al. Oncotarget (2016) PMID: 26989027
116. Summers MG, et al. Clin. Cancer Res. (2017) PMID: 27815357
117. Holderfield M, et al. Nat. Rev. Cancer (2014) PMID: 24957944
118. Burotto M, et al. Cancer (2014) PMID: 24948110
119. Davies H, et al. Nature (2002) PMID: 12068308
120. Kandath C, et al. Nature (2013) PMID: 24132290
121. Greaves WO, et al. J Mol Diagn (2013) PMID: 23273605
122. Klein O, et al. Eur. J. Cancer (2013) PMID: 23237741
123. Wellbrock C, et al. Cancer Res. (2004) PMID: 15059882
124. Hauschild A, et al. Lancet (2012) PMID: 22735384
125. McArthur GA, et al. Lancet Oncol. (2014) PMID: 24508103
126. Fisher R, et al. Cancer Manag Res (2012) PMID: 22904646
127. Yang H, et al. Cancer Res. (2010) PMID: 20551065
128. Gentilcore G, et al. BMC Cancer (2013) PMID: 23317446
129. van den Brom RR, et al. Eur. J. Cancer (2013) PMID: 23473613
130. Klein O, et al. Eur. J. Cancer (2013) PMID: 23490649
131. Ponti G, et al. J. Clin. Pathol. (2013) PMID: 23463675
132. Ponti G, et al. J Hematol Oncol (2012) PMID: 23031422
133. Parakh S, et al. J Clin Pharm Ther (2015) PMID: 25382067
134. Lee LH, et al. JCI Insight (2017) PMID: 28194436
135. Bokemeyer C, et al. Ann. Oncol. (2011) PMID: 21228335
136. Karapetis CS, et al. N. Engl. J. Med. (2008) PMID: 18946061
137. De Roock W, et al. Ann. Oncol. (2008) PMID: 17998284
138. Douillard JY, et al. Ann. Oncol. (2014) PMID: 24718886
139. Amado RG, et al. J. Clin. Oncol. (2008) PMID: 18316791
140. Lièvre A, et al. Cancer Res. (2006) PMID: 16618717
141. Chen J, et al. BMC Cancer (2014) PMID: 25367198
142. Li W, et al. BMC Cancer (2015) PMID: 25929517
143. Hu J, et al. Medicine (Baltimore) (2016) PMID: 27977612
144. Zekri J, et al. Sci. Genet. Mol. Res. (2017) PMID: 28218784
145. Staudacher JJ, et al. Clin Transl Gastroenterol (2017) PMID: 29048416
146. Wang Y, et al. Virchows Arch. (2018) PMID: 29705968
147. Guo F, et al. Sci Rep (2018) PMID: 29666387
148. Mármol I, et al. Int J Mol Sci (2017) PMID: 28106826
149. Kwak MS, et al. Medicine (Baltimore) (2017) PMID: 28106826

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1606031-01

APPENDIX References

- 28858102
150. Pylayeva-Gupta Y, et al. Nat. Rev. Cancer (2011) pmid: 21993244
151. Kahn S, et al. Anticancer Res. (2011) pmid: 3310850
152. Pentheroudakis G, et al. BMC Cancer (2013) pmid: 23374602
153. Vaughn CP, et al. Genes Chromosomes Cancer (2011) pmid: 21305640
154. Janku F, et al. Target Oncol (2013) pmid: 23400451
155. De Rooij W, et al. Lancet Oncol. (2010) pmid: 20619739
156. Irahara N, et al. Diagn. Mol. Pathol. (2010) pmid: 20736745
157. Schirripa M, et al. Int. J. Cancer (2015) pmid: 24806288
158. Cercek A, et al. Clin. Cancer Res. (2017) pmid: 28446505
159. O'Malley et al., 2017; AACR-NCI-EORTC Abstract LB-A12
160. Wu W, et al. Cancer Res. (2015) pmid: 25634209
161. Wang Y, et al. J. Clin. Invest. (2016) pmid: 27454289
162. Ozden O, et al. Sci Rep (2016) pmid: 27197561
163. Zhao W, et al. Nature (2017) pmid: 28976962
164. Seshagiri S, et al. Nature (2012) pmid: 22895193
165. Brannon AR, et al. Genome Biol. (2014) pmid: 25164765
166. Giannakis M, et al. Cell Rep (2016) pmid: 27149842
167. Yaeger R, et al. Cancer Cell (2018) pmid: 29316426
168. Cerami E, et al. Cancer Discov (2012) pmid: 22588877
169. Gao J, et al. Sci Signal (2013) pmid: 23550210
170. Sporn JC, et al. Clin. Cancer Res. (2011) pmid: 21693656
171. Gynecol. Oncol. (2010) pmid: 19959210
172. Laufer M, et al. J. Biol. Chem. (2007) pmid: 17848578
173. Westernmark UK, et al. Mol. Cell. Biol. (2003) pmid: 14560035
174. Zehir A, et al. Nat. Med. (2017) pmid: 28481359
175. Ito S, et al. Nature (2010) pmid: 20639862
176. Guo JU, et al. Cell (2011) pmid: 21496894
177. Iyer LM, et al. Cell Cycle (2009) pmid: 19411852
178. Ko M, et al. Nature (2010) pmid: 21057493
179. Yang H, et al. Oncogene (2013) pmid: 22391558
180. Hu L, et al. Cell (2013) pmid: 24315485
181. Wang Y, et al. Mol. Cell (2015) pmid: 25601757
182. Jaiswal S, et al. N. Engl. J. Med. (2014) pmid: 25426837
183. Genovese G, et al. N. Engl. J. Med. (2014) pmid: 25426838
184. Xie M, et al. Nat. Med. (2014) pmid: 25326804
185. Acuna-Hidalgo R, et al. Am. J. Hum. Genet. (2017) pmid: 28669404
186. Severson EA, et al. Blood (2018) pmid: 29678827
187. Fuster JJ, et al. Circ. Res. (2018) pmid: 29420212
188. Hematology Am Soc Hematol Educ Program (2018) pmid: 30504320
189. Chabon JJ, et al. Nature (2020) pmid: 32269342
190. Razavi P, et al. Nat. Med. (2019) pmid: 31768066
191. Hirai H, et al. Cancer Biol. Ther. (2010) pmid: 20107315
192. Bridges KA, et al. Clin. Cancer Res. (2011) pmid: 21799033
193. Rajeshkumar NV, et al. Clin. Cancer Res. (2011) pmid: 21389100
194. Osman AA, et al. Mol. Cancer Ther. (2015) pmid: 25504633
195. Xu L, et al. Mol. Cancer Ther. (2002) pmid: 12489850
196. Xu L, et al. Mol. Med. (2001) pmid: 11713371
197. Camp ER, et al. Cancer Gene Ther. (2013) pmid: 23470564
198. Kim SS, et al. Nanomedicine (2015) pmid: 25240597
199. Pirollo KF, et al. Mol. Ther. (2016) pmid: 27357628
200. Leijen S, et al. J. Clin. Oncol. (2016) pmid: 27601554
201. Moore et al., 2019; ASCO Abstract 5513
202. Leijen S, et al. J. Clin. Oncol. (2016) pmid: 27998224
203. Oza et al., 2015; ASCO Abstract 5506
204. Lee J, et al. Cancer Discov (2019) pmid: 31315834
205. Méndez E, et al. Clin. Cancer Res. (2018) pmid: 29535125
206. Seligmann JF, et al. J Clin Oncol (2021) pmid: 34538072
207. Gourley et al., 2016; ASCO Abstract 5571
208. Park H, et al. ESMO Open (2022) pmid: 36084396
209. Goh HS, et al. Cancer Res. (1995) pmid: 7585578
210. Berg M, et al. PLoS ONE (2010) pmid: 21103049
211. Han SW, et al. PLoS ONE (2013) pmid: 23700467
212. Malhotra P, et al. Tumour Biol. (2013) pmid: 23526092
213. Di Bartolomeo M, et al. Target Oncol (2014) pmid: 23821376
214. Wangefjord S, et al. Diagn Pathol (2013) pmid: 23337059
215. Russo A, et al. J. Clin. Oncol. (2005) pmid: 16172461
216. Brown CJ, et al. Nat. Rev. Cancer (2009) pmid: 19935675
217. Joergers AC, et al. Annu. Rev. Biochem. (2008) pmid: 18410249
218. Kato S, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) pmid: 12826609
219. Kamada R, et al. J. Biol. Chem. (2011) pmid: 20978130
220. Zerdoumi Y, et al. Hum. Mol. Genet. (2017) pmid: 28472496
221. Yamada H, et al. Carcinogenesis (2007) pmid: 17690113
222. Bougeard G, et al. J. Clin. Oncol. (2015) pmid: 26014290
223. Sorrell AD, et al. Mol Diagn Ther (2013) pmid: 23355100
224. Nichols KE, et al. Cancer Epidemiol. Biomarkers Prev. (2001) pmid: 11219776
225. Kleihues P, et al. Am. J. Pathol. (1997) pmid: 9006316
226. Gonzalez KD, et al. J. Clin. Oncol. (2009) pmid: 19204208
227. Lalloo F, et al. Lancet (2003) pmid: 12672316
228. Mandelker D, et al. Ann. Oncol. (2019) pmid: 31050713
229. Park JH, et al. Cancer Chemother. Pharmacol. (2011) pmid: 21340604
230. Loupakis F, et al. Br. J. Cancer (2009) pmid: 19603018
231. Lupini L, et al. BMC Cancer (2015) pmid: 26508446
232. Inno A, et al. Clin Colorectal Cancer (2011) pmid: 21729677
233. Modest DP, et al. Int. J. Cancer (2012) pmid: 21960311
234. Hong DS, et al. Cancer Discov (2016) pmid: 27729313
235. Kopetz S, et al. N. Engl. J. Med. (2019) pmid: 31566309
236. van Geel RMJM, et al. Cancer Discov (2017) pmid: 28363909
237. Tabernero et al., 2022; ESMO Abstract LBA26
238. Tabernero J, et al. J Clin Oncol (2021) pmid: 33503393
239. Morris et al., 2022; ASCO GI Abstract 12
240. Cunningham D, et al. N. Engl. J. Med. (2004) pmid: 15269313
241. Jonker DJ, et al. N. Engl. J. Med. (2007) pmid: 18003960
242. Papamichael D, et al. Eur J Cancer (2022) pmid: 35033994
243. Chibaudel et al., 2022; ASCO Abstract 3504
244. Moiseyenko VM, et al. Clin Drug Investig (2018) pmid: 29470838
245. Stein A, et al. J Immunother Cancer (2021) pmid: 34315821
246. Price TJ, et al. Lancet Oncol. (2014) pmid: 24739896
247. Sakai D, et al. Eur J Cancer (2020) pmid: 32526634
248. Van Cutsem E, et al. J. Clin. Oncol. (2007) pmid: 17470858
249. Bendell et al., 2014; ASCO Abstract 3515
250. Watanabe J, et al. Int J Cancer (2022) pmid: 35723084
251. Kim TW, et al. Clin Colorectal Cancer (2018) pmid: 29703606
252. Shitara K, et al. Cancer Sci (2016) pmid: 27712015
253. Yoshino et al., 2022; ASCO Abstract LBA1
254. Pietrantonio F, et al. JAMA Oncol (2019) pmid: 31268481
255. Long GV, et al. Ann. Oncol. (2017) pmid: 28475671
256. Long GV, et al. Lancet (2015) pmid: 26037941
257. Robert C, et al. N. Engl. J. Med. (2015) pmid: 25399551
258. Planchard D, et al. Lancet Oncol. (2017) pmid: 28919011
259. Subbiah V, et al. J. Clin. Oncol. (2018) pmid: 29072975
260. Kreitman et al., 2018; ASH Abstract 391
261. Lagana et al., 2018; DOI: 10.1200/PO.18.00019
262. Salama AKS, et al. J Clin Oncol (2020) pmid: 32758030
263. Hendifar A, et al. JCO Precis Oncol (2021) pmid: 34476331
264. Wen PY, et al. Lancet Oncol (2022) pmid: 34838156
265. Subbiah V, et al. Lancet Oncol (2020) pmid: 32818466
266. Williams CB, et al. Onco Targets Ther (2015) pmid: 26664139
267. Gibney GT, et al. Nat Rev Clin Oncol (2013) pmid: 23712190
268. Falchook GS, et al. Thyroid (2015) pmid: 25285888
269. Flaherty KT, et al. N. Engl. J. Med. (2012) pmid: 23020132
270. Long GV, et al. N. Engl. J. Med. (2014) pmid: 25265492
271. Peters S, et al. Melanoma Res. (2014) pmid: 25185693
272. Dummer R, et al. Lancet Oncol. (2018) pmid: 29573941
273. Ascierto PA, et al. Eur. J. Cancer (2020) pmid: 31901705
274. Holbrook KJ, et al. Cancer (2020) pmid: 31658370
275. Sullivan RJ, et al. Clin Cancer Res (2020) pmid: 32669376
276. Kefford et al., 2013; Melanoma Bridge Meeting Abstract P5
277. McLoughlin EM, et al. J Thorac Oncol (2019) pmid: 31757377
278. Gogas et al., 2020; ASCO Abstract 10012
279. Ascierto PA, et al. Lancet Oncol. (2016) pmid: 27480103
280. Ribas A, et al. Clin. Cancer Res. (2020) pmid: 31732523
281. Ribas A, et al. Lancet Oncol. (2014) pmid: 25037139

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.