

ABOUT THE TEST FoundationOne®CDx is a next-generation sequencing (NGS) based assay that identifies genomic findings within hundreds of cancer-related genes.

PATIENT	DISEASE	Brain glioblastoma (GBM)	PHYSICIAN	ORDERING PHYSICIAN	Yeh, Yi-Chen	SPECIMEN	SPECIMEN SITE	Brain
	NAME	Liu, Shu-Yen		MEDICAL FACILITY	Taipei Veterans General Hospital		SPECIMEN ID	S111-12619F (PF22047)
	DATE OF BIRTH	26 May 1970		ADDITIONAL RECIPIENT	None		SPECIMEN TYPE	Slide Deck
	SEX	Female		MEDICAL FACILITY ID	205872		DATE OF COLLECTION	26 March 2022
	MEDICAL RECORD #	48392161		PATHOLOGIST	Not Provided		SPECIMEN RECEIVED	11 April 2022

Biomarker Findings

Microsatellite status - MS-Stable
Tumor Mutational Burden - 1 Muts/Mb

Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

EGFR EGFR-SEPT14 rearrangement, EGFR-SEPT14 fusion, amplification

PIK3CA R88Q, P104L

CDKN2A/B CDKN2A loss, CDKN2B loss

MUTYH W142*

TERT promoter -124C>T

TP53 V272L

2 Disease relevant genes with no reportable alterations: **IDH1**, **PDGFRA**

Report Highlights

- Variants with **diagnostic implications** that may indicate a specific cancer type: **EGFR amplification** (p. 4), **TERT promoter -124C>T** (p. 7)
- Targeted therapies with potential clinical benefit **approved in another tumor type**: Cetuximab (p. 9), Everolimus (p. 10), Panitumumab (p. 10), Temsirolimus (p. 11)
- Evidence-matched **clinical trial options** based on this patient's genomic findings: (p. 12)
- Variants with **prognostic implications** for this tumor type that may impact treatment decisions: **TERT promoter -124C>T** (p. 7)
- Variants in select cancer susceptibility genes to consider for possible **follow-up germline testing** in the appropriate clinical context: **MUTYH W142*** (p. 7)

BIOMARKER FINDINGS

Microsatellite status - MS-Stable

Tumor Mutational Burden - 1 Muts/Mb

GENOMIC FINDINGS

EGFR - EGFR-SEPT14 rearrangement, EGFR-SEPT14 fusion, amplification

5 Trials see p. 12

PIK3CA - R88Q, P104L

10 Trials see p. 13

THERAPY AND CLINICAL TRIAL IMPLICATIONS

No therapies or clinical trials. see Biomarker Findings section

No therapies or clinical trials. see Biomarker Findings section

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)	THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)
none	Cetuximab
	Panitumumab
none	Everolimus
	Temsirolimus

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING IN SELECT CANCER SUSCEPTIBILITY GENES

Findings below have been previously reported as pathogenic germline in the ClinVar genomic database and were detected at an allele frequency of >10%. See appendix for details.

MUTYH - W142* p. 7

This report does not indicate whether variants listed above are germline or somatic in this patient. In the appropriate clinical context, follow-up germline testing would be needed to determine whether a finding is germline or somatic.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

CDKN2A/B - CDKN2A loss, CDKN2B loss.....	p. 6	TERT - promoter -124C>T.....	p. 7
MUTYH - W142*	p. 7	TP53 - V272L.....	p. 8

NOTE Genomic alterations detected may be associated with activity of certain approved therapies; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and exhaustive. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies.

Therapies contained in this report may have been approved by the US FDA.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

BIOMARKER FINDINGS
BIOMARKER

Microsatellite status

RESULT

MS-Stable

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

On the basis of clinical evidence, MSS tumors are significantly less likely than MSI-H tumors to respond to anti-PD-1 immune checkpoint inhibitors¹⁻³, including approved therapies nivolumab and pembrolizumab⁴. In a retrospective analysis of 361 patients with solid tumors treated with pembrolizumab, 3% were MSI-H and

experienced a significantly higher ORR compared with non-MSI-H cases (70% vs. 12%, $p=0.001$)⁵.

FREQUENCY & PROGNOSIS

Low-level MSI has been reported in 5-9% of glioblastoma (GBM) samples⁶⁻⁸. A large-scale study did not find high-level microsatellite instability (MSI-H) in any of 129 GBM samples⁶, although a small-scale study reported MSI-H in 4 of 15 pediatric GBMs and 1 of 12 adult GBMs⁹. The frequency of MSI has been reported to be increased in relapsed compared to primary GBM⁶, in GBMs with a previous lower grade astrocytoma⁷, and in giant cell GBM compared to classic GBM⁸.

FINDING SUMMARY

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor¹⁰. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2¹⁰⁻¹². This sample is microsatellite-stable (MSS), equivalent to the clinical definition of an MSS tumor: one with mutations in none of the tested microsatellite markers¹³⁻¹⁵. MSS status indicates MMR proficiency and typically correlates with intact expression of all MMR family proteins^{10,12,14-15}.

BIOMARKER

Tumor Mutational Burden

RESULT

1 Muts/Mb

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1¹⁶⁻¹⁸, anti-PD-1 therapies¹⁶⁻¹⁹, and combination nivolumab and ipilimumab²⁰⁻²⁵. In glioma, a lack of association between TMB and clinical benefit from immune checkpoint inhibitors has been reported^{16,26-27}. However, multiple case studies have reported that patients with ultramutated gliomas driven by POLE

mutations have benefited from treatment with anti-PD-1²⁸⁻²⁹ or anti-PD-L1³⁰ therapies. Therefore, although increased TMB alone may not be a strong biomarker for PD-1 or PD-L1 inhibitors in this cancer type, these agents may have efficacy for patients with glioma harboring both high TMB and POLE mutation.

FREQUENCY & PROGNOSIS

Glioblastoma (GBM) harbors a median TMB of 2.7 mutations per megabase (mut/Mb), and 4.2% of cases have high TMB (>20 muts/Mb)³¹. For pediatric patients, high TMB has been reported in a subset of high-grade gliomas, frequently in association with mutations in mismatch repair or proofreading genes and in TP53, whereas BRAF alterations or other oncogene fusions were observed more frequently in brain tumors harboring low TMB³²⁻³³. Increased TMB has been reported to correlate with higher tumor grade in glioma³⁴ and glioblastoma (GBM) tissue samples with biallelic mismatch repair deficiency

(bMMRD)²⁸, as well as with shorter OS of patients with diffuse glioma³⁵.

FINDING SUMMARY

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma³⁶⁻³⁷ and cigarette smoke in lung cancer³⁸⁻³⁹, treatment with temozolomide-based chemotherapy in glioma⁴⁰⁻⁴¹, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes⁴²⁻⁴⁶, and microsatellite instability (MSI)^{42,45-46}. This sample harbors a TMB below levels that would be predicted to be associated with sensitivity to PD-1- or PD-L1-targeting immune checkpoint inhibitors, alone or in combination with other agents^{16,26-30}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

GENOMIC FINDINGS

GENE

EGFR

ALTERATION

EGFR-SEPT14 rearrangement, EGFR-SEPT14 fusion, amplification

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

In multiple glioblastoma (GBM) studies, the presence of EGFRvIII has not predicted clinical benefit from first-generation EGFR TKIs such as erlotinib⁴⁷⁻⁵² or gefitinib^{50,53}. However, case reports have described patients with EGFRvIII-positive GBM responding to erlotinib⁵⁴⁻⁵⁷. In a retrospective study of patients with GBM treated with erlotinib or gefitinib, co-expression of EGFRvIII with PTEN protein was the strongest predictor of response ($P < 0.001$)⁵⁸, suggesting that activity in this setting is dependent on PTEN status⁵⁹⁻⁶⁰. However, a prospective Phase 2 trial testing erlotinib monotherapy for patients with EGFRvIII and PTEN-positive recurrent glioblastoma reported minimal efficacy and was terminated⁵². The second-generation EGFR TKIs afatinib and dacomitinib have shown minimal efficacy for patients with EGFRvIII glioblastoma (GBM)⁶¹⁻⁶⁴. A Phase 1/2 study of afatinib, temozolomide, or the combination for patients with GBM reported clinical benefit, including for patients with EGFRvIII; however, temozolomide alone and in combination exhibited better responses than afatinib monotherapy⁶¹⁻⁶². A Phase 2 trial of dacomitinib for patients with EGFR-amplified GBM reported a DCR of 26% (5/19) among patients with EGFR amplification and EGFRvIII; however, the trial failed to meet its primary endpoint of 6-month PFS⁶³. A retrospective biomarker analysis of another Phase 2 study of dacomitinib for patients with GBM found no association between EGFRvIII and clinical benefit⁶⁴. A patient with multiple glioblastoma (GBM) tumors, one of which harbored EGFRvIII, experienced progression of the EGFRvIII-positive tumor during treatment

with osimertinib⁶⁵. Clinical studies of the second-generation EGFR TKIs afatinib and dacomitinib for patients with EGFR-amplified gliomas have shown limited efficacy^{61,63-64,66-67}; however, a small subset of patients has experienced clinical benefit^{63-64,66}. Multiple studies have failed to find a positive association between increased EGFR expression and clinical benefit from erlotinib or gefitinib for patients with glioblastoma^{58,68-70}. There are conflicting data on the efficacy of anti-EGFR antibodies for the treatment of EGFR-amplified tumors. A meta-analysis of colorectal cancer patients treated with second-line or higher cetuximab or panitumumab observed an association between EGFR copy number gain and increased OS and PFS⁷¹. However, studies in head and neck squamous cell carcinoma and gastric cancer found either no association or a negative association between EGFR copy number gain and survival after treatment with first-line cetuximab or panitumumab in combination with chemotherapy⁷²⁻⁷³. The Phase 3 INTELLANCE trial of depatuxizumab mafodotin (ABT-414), an EGFR-targeted antibody-drug conjugate with a toxic payload, in patients with EGFR-amplified glioblastoma (GBM) was stopped for futility. Interim analysis demonstrated improved median PFS (mPFS) of ABT-414 monotherapy compared with placebo (HR=0.84); however, no OS benefit was observed (HR=1.01). Improved mPFS was also observed in patients harboring EGFRvIII (HR=0.73) but without an OS improvement (HR=0.95)⁷⁴. The Phase 2 INTELLANCE trial demonstrated clinical benefit for EGFR-amplified GBM for the combination of ABT-414, temozolomide, and radiotherapy (HR=0.66, $p=0.017$), but there was no evidence of efficacy for ABT-414 monotherapy (HR=1.04, $p=0.83$)⁷⁵.

FREQUENCY & PROGNOSIS

Across several genomic studies of CNS tumors, EGFR amplification has been reported in 16.9% of anaplastic astrocytomas, and 39.7% of glioblastoma multiformes (GBMs)⁷⁶⁻⁷⁹. EGFR alterations have been reported in 13.2% of anaplastic astrocytomas, 5.3-15.9% of glioblastoma multiformes (GBMs), and 0% of pilocytic astrocytomas in several genomic

studies of CNS tumors⁷⁶⁻⁷⁹. In GBMs, Missense mutations in the EGFR extracellular domain have been found in 10-15% of cases and approximately half have a low-level amplification of the mutated allele⁸⁰⁻⁸¹. In a study of IDH-wildtype GBM samples, EGFR alterations were detected in 50% (117/232) of IDH-wildtype GBM samples analyzed, including 41% (95/232) with a co-occurring EGFR amplification and mutation, 26% (61/232) with an EGFR domain truncation event, such as EGFRvIII, and 2.2% (5/232) with an EGFR fusion event⁸². No definitive correlation has been identified between EGFR amplification and length of survival in patients with GBM⁸³⁻⁸⁴; however, EGFR amplification has been associated with prolonged survival in patients over the age of 60 with GBM⁸⁵.

FINDING SUMMARY

EGFR encodes the epidermal growth factor receptor, which belongs to a class of proteins called receptor tyrosine kinases. In response to signals from the environment, EGFR passes biochemical messages to the cell that stimulate it to grow and divide⁸⁶. Amplification of EGFR has been associated with increased expression of EGFR mRNA and protein in several cancer types⁸⁷⁻⁸⁹. Disruption of the EGFR C-terminal region through deletion⁹⁰⁻⁹², truncation^{90-91,93}, splicing errors^{90,93}, or gene fusion⁹⁴⁻⁹⁵, has been demonstrated to be activating and is likely to be oncogenic. These mutations have been shown to cause cellular transformation and tumor formation and to be sensitive to EGFR-targeting therapies, including erlotinib, lapatinib, and cetuximab^{90-92,94-95}. One or more of the alterations observed here are predicted to be activating.

POTENTIAL DIAGNOSTIC IMPLICATIONS

The presence of EGFR gene amplification or TERT promoter mutations are indicative of diffuse astrocytic glioma with molecular features of glioblastoma, WHO grade 4 in IDH1/2-wildtype tumors (NCCN CNS Cancers Guidelines, v2.2021)⁹⁶.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

GENOMIC FINDINGS

GENE

PIK3CA

ALTERATION

R88Q, P104L

TRANSCRIPT ID

NM_006218, NM_006218

CODING SEQUENCE EFFECT

263G>A, 311C>T

VARIANT ALLELE FREQUENCY (% VAF)

26.4%, 59.3%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Clinical and preclinical data in various tumor types indicate that PIK3CA activating alterations may predict sensitivity to therapies targeting PI3K⁹⁷⁻¹⁰⁰, AKT¹⁰¹⁻¹⁰², or mTOR¹⁰³⁻¹¹⁰. In the Phase 2 MATCH trial for patients with PIK3CA-mutated solid tumors, 28% (18/65) of patients experienced PFS lasting at least 6 months after treatment with tasisib; however, no ORs were observed in this study¹¹¹. A separate Phase 1b study of tasisib in combination with the CDK4/6 inhibitor palbociclib for patients with PIK3CA-mutated solid tumors reported an ORR of 0% (n=12) and a DCR of 17% (2/12)¹¹². In a Phase 1 trial of the dual PI3K/mTOR kinase inhibitor apitolisib, 79% (11/

14) of patients with PIK3CA-mutated advanced solid tumors experienced disease control (3 PRs, 8 SDs)¹¹³. The PI3K inhibitor alpelisib is approved as a single agent for the treatment of patients with PIK3CA-related overgrowth spectrum (PROS)¹¹⁴ but has shown limited activity as monotherapy for PIK3CA-mutated solid tumors with a Phase 1a study reporting an ORR of 6.0% (8/134) and a DCR of 58% (78/134)¹¹⁵. The PI3K inhibitor copanlisib exhibited limited efficacy in PIK3CA-mutated tumors¹¹⁶⁻¹¹⁷.

FREQUENCY & PROGNOSIS

PIK3CA mutations have been reported in 9% of glioblastoma (GBM) samples analyzed in the TCGA dataset⁷⁷, and other studies report the incidence of PIK3CA mutations in primary GBMs as 5-18%¹¹⁸⁻¹²⁰. One study detected PIK3CA mutation in 16% (36/232) of IDH-wildtype GBM samples analyzed⁸². PIK3CA mutations have been reported in 5-23% of high-grade gliomas (including glioblastomas, anaplastic astrocytomas, and anaplastic oligodendrogliomas)¹¹⁸⁻¹²². While another study did not observe PIK3CA mutations in low-grade astrocytomas or in anaplastic astrocytomas, it did report high ERK and AKT activity¹²⁰. One study found that PIK3CA mutation in glioblastoma (GBM) was associated with shorter median PFS in both a discovery cohort (6.9 vs. 12.4 months, HR=2.89, p=0.01) and

in the TCGA cohort (6.1 vs. 9 months, p=0.008), but was not consistently associated with median OS¹²³. In a study of IDH-wildtype GBM, patients with alterations in PI3K class I genes (PIK3CA, PIK3R1, PIK3CG, and PIK3R2) had significantly longer OS (20.0 months altered vs. 16.9 months wildtype, HR=0.62, p=0.002) and PFS (11.0 months altered vs. 7.4 months wildtype, p=0.0043); patients with PIK3CA alterations experienced an improved OS but this association was not highly significant (20.0 months altered vs. 18.1 months wildtype, p=0.0407)⁸².

FINDING SUMMARY

PIK3CA encodes p110-alpha, which is the catalytic subunit of phosphatidylinositol 3-kinase (PI3K). The PI3K pathway is involved in cell signaling that regulates a number of critical cellular functions, including cell growth, proliferation, differentiation, motility, and survival¹²⁴⁻¹²⁵. PIK3CA P104L, E110del, E453del and G914R have been reported to increase PI3K activity and have been reported in individuals with megalencephaly syndromes, suggesting they promote cell growth¹²⁶. In addition, missense mutations of E453 and a short deletion including this residue have been shown to be activating¹²⁷⁻¹²⁹. PIK3CA alterations that have been characterized as activating, such as observed here, are predicted to be oncogenic^{126-127,130-149}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

GENOMIC FINDINGS

GENE

CDKN2A/B

ALTERATION

CDKN2A loss, CDKN2B loss

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Preclinical data suggest that tumors with loss of p16INK4a function may be sensitive to CDK4/6 inhibitors, such as abemaciclib, ribociclib, and palbociclib¹⁵⁰⁻¹⁵³. Although case studies have reported that patients with breast cancer or uterine leiomyosarcoma harboring CDKN2A loss responded to palbociclib treatment¹⁵⁴⁻¹⁵⁵, multiple other clinical studies have shown no significant correlation between p16INK4a loss or inactivation and therapeutic benefit of these agents¹⁵⁶⁻¹⁶²; it is not known whether CDK4/6 inhibitors would be beneficial in this case. Although preclinical studies have suggested that loss of p14ARF function may be associated with reduced sensitivity to MDM2 inhibitors¹⁶³⁻¹⁶⁴, the clinical relevance of p14ARF as a predictive biomarker is not clear. There are no drugs that directly target the mutation or loss of CDKN2B in cancer. Because the p15INK4b protein encoded by CDKN2B is known to inhibit CDK4, tumors with CDKN2B mutation or loss may predict sensitivity to CDK4/6 inhibitors, such as ribociclib, abemaciclib, and palbociclib^{157,159-160,165-167}.

FREQUENCY & PROGNOSIS

Concurrent putative homozygous deletion of CDKN2A and CDKN2B has been reported in 35% of patients with gliomas⁷⁸ and detected more frequently in patients with glioblastoma multiforme (GBM; 58%)⁷⁷ than in those with lower grade gliomas (13%) (cBioPortal, Sep 2021)¹⁶⁸⁻¹⁶⁹. In other studies, loss of CDKN2A/B by deletion has been reported in up to 78% of astrocytomas (including anaplastic astrocytomas and GBM)¹⁷⁰⁻¹⁷². A study found homozygous deletion of both p16INK4a and p14ARF in 26% (13/50) of glioblastomas (GBMs); 18% (9/50) of cases showed homozygous deletion of the p14ARF-encoding locus alone¹⁷³. One study detected CDKN2A/B loss in 69% (161/232) and mutation in 2.6% (6/232) of IDH-wildtype GBM samples analyzed⁸². Decreased p14ARF and p16INK4a expression levels were found to be tightly associated in a study of glioma samples¹⁷⁴. Homozygous deletion of the genomic region including CDKN2A and CDKN2B has been found to be associated with poor prognosis in GBM and likely serves as an early event in GBM progression^{171,175}. In addition, expression of p16INK4a has been found to be lower in patients with high grade malignant gliomas compared to patients with low grade gliomas, and loss of p16INK4a expression has been associated with shorter overall survival in pilocytic astrocytomas¹⁷⁶⁻¹⁷⁷.

FINDING SUMMARY

CDKN2A encodes two different, unrelated tumor suppressor proteins, p16INK4a and p14ARF, whereas CDKN2B encodes the tumor suppressor

p15INK4b¹⁷⁸⁻¹⁷⁹. Both p15INK4b and p16INK4a bind to and inhibit CDK4 and CDK6, thereby maintaining the growth-suppressive activity of the Rb tumor suppressor; loss or inactivation of either p15INK4b or p16INK4a contributes to dysregulation of the CDK4/6-cyclin-Rb pathway and loss of cell cycle control¹⁸⁰⁻¹⁸¹. The tumor suppressive functions of p14ARF involve stabilization and activation of p53, via a mechanism of MDM2 inhibition¹⁸²⁻¹⁸³. One or more alterations observed here are predicted to result in p16INK4a loss of function¹⁸⁴⁻²⁰⁵. One or more alterations seen here are predicted to result in p14ARF loss of function^{188,205-208}. CDKN2B alterations such as seen here are predicted to inactivate p15INK4b²⁰⁹.

POTENTIAL GERMLINE IMPLICATIONS

Germline CDKN2A mutation is associated with melanoma-pancreatic cancer syndrome, a condition marked by increased risk of developing malignant melanoma and/or pancreatic cancer²¹⁰. Mutation carriers within families may develop either or both types of cancer, and melanoma cases may be referred to as familial or hereditary melanoma²¹¹⁻²¹². CDKN2A is the most implicated gene in familial melanoma, with germline mutations present in 16% to 20% of familial melanoma cases²¹³⁻²¹⁵. CDKN2A alteration has also been implicated in familial melanoma-astrocytoma syndrome, an extremely rare tumor association characterized by dual predisposition to melanoma and nervous system tumors²¹⁶⁻²¹⁸. In the appropriate clinical context, germline testing of CDKN2A is recommended.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

GENOMIC FINDINGS
GENE

MUTYH

ALTERATION

W142*

TRANSCRIPT ID

NM_001048171

CODING SEQUENCE EFFECT

425G>A

VARIANT ALLELE FREQUENCY (% VAF)

48.4%

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are no therapies or clinical trials available to address MUTYH alterations in cancer.

FREQUENCY & PROGNOSIS

In general, somatic MUTYH mutations are infrequently reported across cancer types (COSMIC, 2022)²¹⁹. Monoallelic MUTYH mutation

occurs in 1-2% of the general population²²⁰⁻²²¹. There is conflicting data regarding the impact of monoallelic mutations on the risk of developing CRC²²²⁻²²⁴. Patients with MUTYH-mutant CRC were reported to have significantly improved overall survival compared to patients without MUTYH mutation²²⁵.

FINDING SUMMARY

MUTYH (also known as MYH) encodes an enzyme involved in DNA base excision repair, and loss of function mutations in MUTYH result in increased rates of mutagenesis and promotion of tumorigenesis²²⁶. The two most frequently reported MUTYH loss of function mutations are G382D (also referred to as G396D) and Y165C (also referred to as Y179C)^{220-221,227-229}. Numerous other MUTYH mutations have also been shown to result in loss of function²²⁷⁻²³⁰.

POTENTIAL GERMLINE IMPLICATIONS

One or more of the MUTYH variants observed here has been described in the ClinVar database as

a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters) associated with MUTYH-associated polyposis (ClinVar, Sep 2021)²³¹. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. Germline biallelic MUTYH mutation causes MUTYH-associated polyposis (also known as MYH-associated polyposis or MAP), an autosomal recessive condition characterized by multiple colorectal adenomas and increased lifetime risk of colorectal cancer (CRC)^{220,232-234}. MAP accounts for approximately 0.7% of all CRC cases and 2% of early-onset CRC cases²²⁰. In contrast to CRC, the role of MUTYH mutation in the context of other cancer types is not well established²³⁵⁻²³⁹. Estimates for the prevalence of MAP in the general population range from 1:5,000-1:10,000²²¹. Therefore, in the appropriate clinical context, germline testing of MUTYH is recommended.

GENE

TERT

ALTERATION

promoter -124C>T

TRANSCRIPT ID

NM_198253

CODING SEQUENCE EFFECT

-124C>T

VARIANT ALLELE FREQUENCY (% VAF)

48.7%

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

Therapeutic options for targeting tumors with TERT mutations are limited, although a variety of approaches have been investigated, including immunotherapies using TERT as a tumor-associated antigen and antisense oligonucleotide- or peptide-based therapies. TERT peptide vaccines showed limited anticancer efficacy in clinical trials²⁴⁰; however, in one preclinical study, the combination of a TERT peptide vaccine and anti-CTLA-4 therapy suppressed tumor growth²⁴¹. A Phase 2 study of the TERT inhibitor imetelstat for patients with advanced non-small cell lung cancer

reported no improvement in PFS or OS²⁴².

FREQUENCY & PROGNOSIS

TERT promoter mutations have been reported in 51-59% of gliomas²⁴³⁻²⁴⁴, most frequently in glioblastoma (GBM, 54-84%), gliosarcoma (81%), oligodendroglioma (78%), and historically in oligoastrocytomas (25-31%) but less frequently in lower grade astrocytomas (10-18%) and in only 1% of ependymomas²⁴³⁻²⁴⁷. In patients with glioblastoma (GBM), the prevalence of TERT promoter mutation is lower in pediatric primary GBM (11%) and adult secondary GBM (28%) compared with adult primary GBM (58-83%)^{243,245}. One study detected TERT promoter mutations in 78% (181/232) of IDH-wildtype GBM samples analyzed⁸². TERT promoter mutation has been shown to be significantly associated with increased TERT gene expression in astrocytoma, oligodendroglioma, and GBM²⁴⁸. TERT promoter mutations significantly associate with poor prognosis in patients with GBM, although this correlation may be due to the association with primary GBM as opposed to IDH-positive secondary GBM^{243,245,248-249}. In the context of IDH-wildtype glioma, TERT mutations are associated with reduced OS (NCCN CNS Cancers Guidelines, v2.2021).

FINDING SUMMARY

Telomerase reverse transcriptase (TERT, or hTERT) is a catalytic subunit of the telomerase complex, which is required to maintain appropriate chromosomal length²⁵⁰. Activation of TERT is a hallmark of cancer, being detected in up to 80-90% of malignancies and absent in quiescent cells²⁵¹⁻²⁵³. Mutations within the promoter region of TERT that confer enhanced TERT promoter activity have been reported in two hotspots, located at -124 bp and -146 bp upstream of the transcriptional start site (also termed C228T and C250T, respectively)²⁵⁴⁻²⁵⁶, as well as tandem mutations at positions -124/-125 bp and -138/-139 bp²⁵⁴.

POTENTIAL DIAGNOSTIC IMPLICATIONS

TERT mutations are associated with 1p/19q co-deletion in oligodendrogliomas, and are highly recurrent in IDH/ATRX-wildtype glioblastoma (GBM) (NCCN CNS Cancers Guidelines, v2.2021)²⁵⁷. The presence of EGFR gene amplification or TERT promoter mutations are indicative of diffuse astrocytic glioma with molecular features of glioblastoma, WHO grade 4 in IDH1/2-wildtype tumors (NCCN CNS Cancers Guidelines, v2.2021)⁹⁶.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

GENOMIC FINDINGS

GENE

TP53

ALTERATION

V272L

TRANSCRIPT ID

NM_000546

CODING SEQUENCE EFFECT

814G>T

VARIANT ALLELE FREQUENCY (% VAF)

65.4%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved therapies to address TP53 mutation or loss. However, tumors with TP53 loss of function alterations may be sensitive to the WEE1 inhibitor adavosertib²⁵⁸⁻²⁶¹, or p53 gene therapy and immunotherapeutics such as SGT-53²⁶²⁻²⁶⁶ and ALT-801²⁶⁷. In a Phase 1 study, adavosertib in combination with gemcitabine, cisplatin, or carboplatin elicited PRs in 9.7% and SDs in 53% of patients with solid tumors; the response rate was 21% (4/19) for patients with TP53 mutations versus 12% (4/33) for patients who were TP53 wildtype²⁶⁸. A Phase 2 trial of adavosertib in combination with chemotherapy (gemcitabine, carboplatin, paclitaxel, or doxorubicin) reported a 32% (30/94, 3 CR) ORR and a 73% (69/94) DCR for patients with platinum-refractory TP53-mutated ovarian, Fallopian tube, or peritoneal cancer²⁶⁹. A smaller Phase 2 trial of adavosertib in combination with carboplatin achieved a 43% (9/21, 1 CR) ORR and a 76% (16/21) DCR for patients with platinum-refractory TP53-mutated ovarian cancer²⁷⁰. The combination of adavosertib with paclitaxel and carboplatin for patients with TP53-mutated ovarian cancer also significantly increased PFS compared with paclitaxel and carboplatin alone²⁷¹. In the Phase 2 VIKTORY trial, patients with TP53-mutated metastatic and/or recurrent gastric cancer experienced a 24% (6/25) ORR with adavosertib combined with paclitaxel²⁷². A Phase 1 trial of neoadjuvant adavosertib in combination with cisplatin and docetaxel for head and neck squamous cell carcinoma (HNSCC) elicited a 71% (5/7) response rate for patients with TP53

alterations²⁷³. The Phase 2 FOCUS4-C trial for patients with TP53- and RAS-mutated colorectal cancer reported improvement in PFS (3.61 vs. 1.87 months, HR=0.35, p=0.0022), but not OS (14.0 vs 12.8 months, p=0.93), following adavosertib treatment compared with active monitoring²⁷⁴. In a Phase 1b clinical trial of SGT-53 in combination with docetaxel for patients with solid tumors, 75% (9/12) of evaluable patients experienced clinical benefit, including 2 confirmed and 1 unconfirmed PRs and 2 instances of SD with significant tumor shrinkage²⁶⁶. Missense mutations leading to TP53 inactivation may also be sensitive to therapies that reactivate mutated p53 such as APR-246²⁷⁵⁻²⁷⁷. In a Phase 1b trial for patients with p53-positive high-grade serous ovarian cancer, APR-246 combined with carboplatin and pegylated liposomal doxorubicin achieved a 52% (11/21) response rate and 100% DCR²⁷⁸. ATR inhibitor treatment of chronic lymphocytic leukemia (CLL) cells with biallelic inactivation of TP53 suppressed cell viability, promoted DNA damage, and attenuated xenograft growth in preclinical studies²⁷⁹⁻²⁸⁰; however, ATR inhibitors as monotherapy had little effect on these parameters in solid tumor models in other preclinical studies²⁸¹⁻²⁸². Therefore, it is unclear whether TP53 inactivation predicts sensitivity to ATR inhibition.

FREQUENCY & PROGNOSIS

In the TCGA dataset, TP53 alterations have been reported in 35% of glioblastomas (GBMs), with a high incidence in pediatric and secondary GBMs and a low incidence in primary GBMs^{122,283}. One study detected TP53 alterations in 31% (73/232) of IDH-wildtype GBM samples analyzed, with most of the events being mutations⁸². TP53 mutations have been reported in 18-40% of astrocytoma samples, and preferentially in anaplastic astrocytoma; one study reported TP53 loss of function and partially/fully functional mutations in 15% and 25% of anaplastic astrocytomas, respectively²⁸⁴⁻²⁸⁹. Some studies suggest that the presence of a TP53 mutation is correlated with a favorable prognosis in patients with glioblastoma (GBM)²⁹⁰. One study reported that TP53 alterations were associated with poorer OS (12.9 months altered vs. 19.7 months wildtype, HR=1.58, p=0.0054) in IDH-wildtype GBM⁸². Mutation of TP53 is thought to be an early step in the

tumorigenesis of astrocytomas, which can progress into anaplastic astrocytoma and then glioblastoma through gain of other genetic abnormalities such as loss of CDKN2A or RB1, followed by loss of PTEN²⁹¹.

FINDING SUMMARY

Functional loss of the tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers²⁹². Alterations such as seen here may disrupt TP53 function or expression²⁹³⁻²⁹⁷.

POTENTIAL GERMLINE IMPLICATIONS

Germline mutations in TP53 are associated with the very rare autosomal dominant disorder Li-Fraumeni syndrome and the early onset of many cancers²⁹⁸⁻³⁰⁰, including sarcomas³⁰¹⁻³⁰². Estimates for the prevalence of germline TP53 mutations in the general population range from 1:5,000³⁰³ to 1:20,000³⁰². For pathogenic TP53 mutations identified during tumor sequencing, the rate of germline mutations was 1% in the overall population and 6% in tumors arising before age 30³⁰⁴. In the appropriate clinical context, germline testing of TP53 is recommended.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion³⁰⁵⁻³¹⁰. CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy³⁰⁵⁻³⁰⁶. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease³¹¹. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH^{309,312-313}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Cetuximab

Assay findings association

EGFR

EGFR-SEPT14 rearrangement,
EGFR-SEPT14 fusion,
amplification

AREAS OF THERAPEUTIC USE

Cetuximab is a monoclonal antibody that targets EGFR. It is FDA approved for the treatment of head and neck squamous cell carcinoma (HNSCC) and KRAS-wild-type, EGFR-expressing metastatic colorectal cancer (CRC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

For patients with metastatic CRC receiving cetuximab or panitumumab as mono- or combination therapy, increased EGFR copy number associated with improved OS (HR=0.62) in a meta-analysis, although increased survival was not seen in populations that received first-line treatment with EGFR antibodies⁷¹.

SUPPORTING DATA

A clinical trial of cetuximab with bevacizumab (an anti-VEGF monoclonal antibody) in patients with glioblastoma (GBM) did not show improved efficacy compared with bevacizumab alone³¹⁴. In preclinical trials, cetuximab, matuzumab, and panitumumab were reported to be ineffective at blocking EGFR dimerization and activation in GBM cells expressing EGFR extracellular

domain mutations³¹⁵. However, another study demonstrated that in patients with GBM harboring EGFR amplification but lacking expression of the EGFRvIII variant, treatment with cetuximab resulted in significantly better progression-free survival (PFS) and numerical (although not statistically significant) improvement in overall survival (OS)³¹⁶. A Phase 3 trial of combined cetuximab and platinum/5-FU in patients with HNSCC demonstrated improved response compared to platinum/5-FU alone, but EGFR amplification was not shown to predict response to this treatment⁷². A Phase 3 study of patients with pancreatic adenocarcinoma did not report any improved outcome in patients treated with a combination of cetuximab plus gemcitabine vs gemcitabine alone³¹⁷. In a Phase 1/2 trial of 36 patients with metastatic castration-resistant prostate cancer (mCRPC) treated with cetuximab in combination with doxorubicin, stable disease was reported in approximately 63% of patients³¹⁸. A Phase 1 study of the combination therapy of cetuximab, erlotinib, and bevacizumab reported stable disease in 21% (7/34) of patients with non-small cell lung cancer (NSCLC)³¹⁹.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Everolimus

Assay findings association

PIK3CA

R88Q, P104L

AREAS OF THERAPEUTIC USE

Everolimus is an orally available mTOR inhibitor that is FDA approved to treat renal cell carcinoma (RCC) following antiangiogenic therapy; pancreatic neuroendocrine tumors; and well-differentiated non-functional neuroendocrine tumors of the lung or gastrointestinal tract. Everolimus is also approved to treat either renal angiomyolipoma or subependymal giant cell astrocytoma in association with tuberous sclerosis complex (TSC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical evidence¹⁰³⁻¹¹⁰, PIK3CA activating mutations may predict sensitivity to mTOR inhibitors such as everolimus and temsirolimus. Studies have reported modest activity of these therapies as single agents (ORRs of 0-4%), but improved activity has been observed when they are combined with other agents such as bevacizumab and doxorubicin (ORRs of 25-44%), for patients with PIK3CA-mutated solid tumors^{107-110,320-324}.

SUPPORTING DATA

Case reports have described 2 children with PIK3CA-mutated diffuse glioma or glioneuronal tumor who benefited from treatment with everolimus alone or in combination with temozolomide³²⁵⁻³²⁶, and 1 adult with glioblastoma (GBM) harboring PIK3CA mutation and KRAS amplification who experienced disease progression

with single-agent everolimus⁶⁷. A Phase 2 trial of radiotherapy (RT), temozolomide (TMZ), and bevacizumab followed by everolimus and bevacizumab reported that 61% (31/51) of patients with newly diagnosed glioblastoma had objective responses with a median progression-free survival (PFS) of 11.3 months and median overall survival (OS) of 13.9 months³²⁷. A Phase 2 study of everolimus combined with TMZ and RT for the treatment of newly diagnosed glioblastoma reported a median PFS of 6.4 months and median OS of 15.8 months³²⁸. A Phase 1 trial of everolimus plus TMZ for patients with newly diagnosed or progressive glioblastoma reported partial responses (PR) in 11% (3/28) and stable disease (SD) in 57% (16/28) of cases³²⁹. A pilot study of everolimus with gefitinib in patients with recurrent glioblastoma reported 14% (3/22) PRs, 36% (8/22) SDs, and median PFS and OS of 2.6 months and 5.8 months, respectively³³⁰. Everolimus treatment achieved SD in 45% (5/11) of pediatric patients with heavily pretreated low-grade CNS tumors; median PFS of these responses was 14 months³³¹. Whereas frequent adverse events precluded a recommended Phase 2 dose and schedule for the combination of trametinib and everolimus in a Phase 1b trial for solid tumors³³², a retrospective study for heavily pretreated patients with solid tumors reported tolerable regimens of the combination for 23/31 patients, with 16 patients treated >3 months and evaluable patients achieving a median PFS of 6.5 months³³³.

Panitumumab

Assay findings association

EGFR

EGFR-SEPT14 rearrangement,
EGFR-SEPT14 fusion,
amplification

AREAS OF THERAPEUTIC USE

Panitumumab is a monoclonal antibody that targets EGFR. It is FDA approved to treat KRAS wild-type and NRAS wild-type metastatic colorectal cancer (CRC) combined with chemotherapy or as monotherapy for patients who have progressed on prior chemotherapy. Please see the drug label for full prescribing information.

GENE ASSOCIATION

For patients with metastatic CRC receiving cetuximab or panitumumab as mono- or combination therapy, increased EGFR copy number associated with improved OS (HR=0.62) in a meta-analysis, although increased survival was not seen in populations that received first-line treatment with EGFR antibodies⁷¹.

SUPPORTING DATA

A Phase 1 trial of EnGeneIC delivery vehicle (EDV) targeting EGFR with panitumumab in combination with

doxorubicin for 14 patients with glioblastoma (GBM) reported no responses and 28% (4/14) SDs³³⁴. Panitumumab has shown efficacy as monotherapy or in combination with chemotherapy for patients with KRAS-wildtype colorectal cancer³³⁵⁻³³⁷ and has been investigated in a variety of other tumor types. For patients with head and neck squamous cell carcinoma (HNSCC), data are conflicting; some trials of panitumumab in various lines and with different chemotherapy combinations have shown modest benefit³³⁸⁻³⁴⁰ and others have reported no benefit³⁴¹⁻³⁴³. A Phase 3 study of chemotherapy with or without panitumumab for patients with advanced gastroesophageal cancer was terminated for futility³⁴⁴. Trials in a variety of tumor types have failed to show significant benefit for patients, including non-small cell lung cancer (NSCLC)³⁴⁵⁻³⁴⁶; biliary tract cancers, including cholangiocarcinoma³⁴⁷⁻³⁴⁸; and renal cell carcinoma (RCC)³⁴⁹.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Temsirolimus

Assay findings association

PIK3CA

R88Q, P104L

AREAS OF THERAPEUTIC USE

Temsirolimus is an intravenous mTOR inhibitor that is FDA approved for the treatment of advanced renal cell carcinoma. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical evidence¹⁰³⁻¹¹⁰, PIK3CA activating mutations may predict sensitivity to mTOR inhibitors such as everolimus and temsirolimus. Studies have reported modest activity of these therapies as single agents (ORRs of 0-4%), but improved activity has been observed when they are combined with other agents such as bevacizumab and doxorubicin (ORRs of 25-44%), for patients with PIK3CA-mutated solid tumors^{107-110,320-324}.

SUPPORTING DATA

A Phase 1, dose-escalation trial combining temsirolimus and radiation/temozolomide therapy, with or without adjuvant temozolomide monotherapy, in patients with

newly diagnosed glioblastoma reported no clinical responses but 24/25 patients experienced a period of stable disease; increased infection rates were noted with this regimen³⁵⁰. A Phase 1/2 trial of temsirolimus in combination with sorafenib in glioblastoma was terminated at the Phase 2 interim analysis after patients failed to meet the primary endpoint of 6 month progression-free survival; significant toxicity was also observed in the combination therapy, even at low doses of temsirolimus³⁵¹. A Phase 2 study showed that addition of temsirolimus to bevacizumab therapy in patients with recurrent glioblastoma did not add clinical benefit³⁵². A Phase 2 clinical trial of temsirolimus in pediatric glioma reported disease stabilization in 7/17 patients including one patient with anaplastic astrocytoma³⁵³. A Phase 1/2 study of temsirolimus in combination with erlotinib reported 6% (1/16) complete responses, 6% (1/16) partial responses, and 12.5% (2/16) instances of stable disease in patients with anaplastic glioma⁵¹.

NOTE Genomic alterations detected may be associated with activity of certain FDA approved drugs, however, the agents listed in this report may have varied evidence in the patient's tumor type.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

CLINICAL TRIALS

NOTE Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually updated and

should be investigated by the physician or research staff. This is not a comprehensive list of all available clinical trials. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials listed here may have additional enrollment criteria that may require

medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see clinicaltrials.gov. Or visit <https://www.foundationmedicine.com/genomic-testing#support-services>.

GENE
EGFR

ALTERATION
 EGFR-SEPT14 rearrangement, EGFR-SEPT14 fusion, amplification

RATIONALE
 EGFR activating mutations, rearrangements, or amplification may predict sensitivity to EGFR-targeted therapies. Strategies to overcome

resistance to current agents include next-generation EGFR inhibitors and combination therapies.

NCT03618667
PHASE 2

GC1118 in Recurrent Glioblastoma Patients With High EGFR Amplification

TARGETS
 EGFR

LOCATIONS: Seoul (Korea, Republic of)

NCT04720976
PHASE 1/2

JAB-3312 Activity in Adult Patients With Advanced Solid Tumors

TARGETS
 MEK, SHP2, PD-1, EGFR, KRAS

LOCATIONS: Utah

NCT02800486
PHASE 2

Super Selective Intra-arterial Repeated Infusion of Cetuximab (Erbix) With Reirradiation for Treatment of Relapsed/Refractory GBM, AA, and AOA

TARGETS
 EGFR

LOCATIONS: New York

NCT02861898
PHASE 1/2

Super-selective Intra-arterial Repeated Infusion of Cetuximab for the Treatment of Newly Diagnosed Glioblastoma

TARGETS
 EGFR

LOCATIONS: New York

NCT03783403
PHASE 1

 A Study of CC-95251, a Monoclonal Antibody Directed Against SIRP α , in Subjects With Advanced Solid and Hematologic Cancers

TARGETS
 CD20, EGFR, SIRP-alpha

LOCATIONS: Heidelberg (Australia), Melbourne (Australia), Edmonton (Canada), Oregon, California, Arizona, Toronto (Canada), Oklahoma, Missouri, Pennsylvania

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

CLINICAL TRIALS

GENE

PIK3CA

ALTERATION

R88Q, P104L

RATIONALE

PIK3CA activating mutations may lead to activation of the PI3K-AKT-mTOR pathway and may therefore indicate sensitivity to inhibitors of

this pathway. Strong clinical data support sensitivity of PIK3CA-mutated solid tumors to the PI3K-alpha inhibitor alpelisib.

NCT04589845

PHASE 2

Tumor-Agnostic Precision Immuno-Oncology and Somatic Targeting Rational for You (TAPISTRY) Platform Study

TARGETS

TRKB, ALK, TRKC, ROS1, TRKA, RET, PD-L1, AKTs, ERBB2, MDM2, PI3K-alpha

LOCATIONS: Taipei City (Taiwan), Taoyuan County (Taiwan), Tainan (Taiwan), Hong Kong (Hong Kong), Seoul (Korea, Republic of), Xi'an (China), Tianjin (China), Beijing (China), Chengdu City (China), Changchun (China)

NCT03239015

PHASE 2

Efficacy and Safety of Targeted Precision Therapy in Refractory Tumor With Druggable Molecular Event

TARGETS

EGFR, ERBB4, ERBB2, PARP, mTOR, MET, ROS1, RET, VEGFRs, BRAF, CDK4, CDK6

LOCATIONS: Shanghai (China)

NCT04337463

PHASE NULL

ATG-008 Combined With Toripalimab in Advanced Solid Tumors

TARGETS

mTORC1, mTORC2, PD-1

LOCATIONS: Chongqing (China), Chengdu (China)

NCT04803318

PHASE 2

Trametinib Combined With Everolimus and Lenvatinib for Recurrent/Refractory Advanced Solid Tumors

TARGETS

mTOR, FGFRs, RET, PDGFRA, VEGFRs, KIT, MEK

LOCATIONS: Guangzhou (China)

NCT05125523

PHASE 1

A Study of Sirolimus for Injection (Albumin Bound) in Patients With Advanced Solid Tumors

TARGETS

mTOR

LOCATIONS: Tianjin (China)

NCT03772561

PHASE 1

Phase I Study of AZD5363 + Olaparib + Durvalumab in Patients With Advanced or Metastatic Solid Tumor Malignancies

TARGETS

PARP, AKTs, PD-L1

LOCATIONS: Singapore (Singapore)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 18 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

CLINICAL TRIALS
NCT04801966
PHASE NULL

Safety and Oversight of the Individually Tailored Treatment Approach: A Novel Pilot Study

TARGETS

CDK4, CDK6, PI3K-alpha, PD-L1, MEK, PARP, PD-1, BRAF

LOCATIONS: Melbourne (Australia)

NCT04632992
PHASE 2

A Study Evaluating Targeted Therapies in Participants Who Have Advanced Solid Tumors With Genomic Alterations or Protein Expression Patterns Predictive of Response

TARGETS

TRKB, ALK, TRKC, ROS1, TRKA, PD-L1, ERBB2, PI3K-alpha, RET, AKTs

LOCATIONS: Alaska, Washington, Oregon, California, Idaho

NCT03994796
PHASE 2

Genetic Testing in Guiding Treatment for Patients With Brain Metastases

TARGETS

TRKB, ALK, TRKC, ROS1, TRKA, CDK4, CDK6, PI3K, mTOR

LOCATIONS: Washington, Oregon, Idaho, Montana

NCT03673787
PHASE 1/2

A Trial of Ipatasertib in Combination With Atezolizumab

TARGETS

AKTs, PD-L1

LOCATIONS: Sutton (United Kingdom)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

APPENDIX
Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

ARID1A
A183T

ASXL1
V1092M

CDK6
rearrangement

EGFR
rearrangement and
rearrangement

LTK
G213_A214insGGG and
R575W

MSH6
K1358fs*2

PTCH1
E44G and S827G

RAD51D
G265R

SPEN
N2999K

ZNF217
I211V

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

APPENDIX
Genes Assayed in FoundationOne®CDx

FoundationOne CDx is designed to include genes known to be somatically altered in human solid tumors that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay interrogates 324 genes as well as introns of 36 genes involved in rearrangements. The assay will be updated periodically to reflect new knowledge about cancer biology.

DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS

ABL1	ACVR1B	AKT1	AKT2	AKT3	ALK	ALOX12B	AMER1 (FAM123B)	APC
AR	ARAF	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX	AURKA
AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2	BCL6
BCOR	BCORL1	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2
BTK	C11orf30 (EMSY)	C17orf39 (GID4)	CALR	CARD11	CASP8	CBFB	CBL	CCND1
CCND2	CCND3	CCNE1	CD22	CD274 (PD-L1)	CD70	CD79A	CD79B	CDC73
CDH1	CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B
CDKN2C	CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R
CTCF	CTNNA1	CTNNB1	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1
DDR2	DIS3	DNMT3A	DOT1L	EED	EGFR	EP300	EPHA3	EPHB1
EPHB4	ERBB2	ERBB3	ERBB4	ERCC4	ERG	ERRF1	ESR1	EZH2
FAM46C	FANCA	FANCC	FANCG	FANCL	FAS	FBXW7	FGF10	FGF12
FGF14	FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1	FGFR2	FGFR3
FGFR4	FH	FLCN	FLT1	FLT3	FOXL2	FUBP1	GABRA6	GATA3
GATA4	GATA6	GNA11	GNA13	GNAQ	GNAS	GRM3	GSK3B	H3F3A
HDAC1	HGF	HNF1A	HRAS	HSD3B1	ID3	IDH1	IDH2	IGF1R
IKBKE	IKZF1	INPP4B	IRF2	IRF4	IRS2	JAK1	JAK2	JAK3
JUN	KDM5A	KDM5C	KDM6A	KDR	KEAP1	KEL	KIT	KLHL6
KMT2A (MLL)	KMT2D (MLL2)	KRAS	LTK	LYN	MAF	MAP2K1 (MEK1)	MAP2K2 (MEK2)	MAP2K4
MAP3K1	MAP3K13	MAPK1	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1
MERTK	MET	MITF	MKKN1	MLH1	MPL	MRE11A	MSH2	MSH3
MSH6	MST1R	MTAP	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN	MYD88
NBN	NF1	NF2	NFE2L2	NFKB1A	NKX2-1	NOTCH1	NOTCH2	NOTCH3
NPM1	NRAS	NSD3 (WHSC1L1)	NTSC2	NTRK1	NTRK2	NTRK3	P2RY8	PALB2
PARK2	PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)	PDGFRA
PDGFRB	PDK1	PIK3C2B	PIK3C2G	PIK3CA	PIK3CB	PIK3R1	PIM1	PMS2
POLD1	POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI	PTCH1
PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51	RAD51B	RAD51C
RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	REL	RET
RICTOR	RNF43	ROS1	RPTOR	SDHA	SDHB	SDHC	SDHD	SETD2
SF3B1	SGK1	SMAD2	SMAD4	SMARCA4	SMARCB1	SMO	SNCAIP	SOC3
SOX2	SOX9	SPEN	SPOP	SRC	STAG2	STAT3	STK11	SUFU
SYK	TBX3	TEK	TET2	TGFBR2	TIPARP	TNFAIP3	TNFRSF14	TP53
TSC1	TSC2	TYRO3	U2AF1	VEGFA	VHL	WHSC1	WT1	XPO1
XRCC2	ZNF217	ZNF703						

DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS

ALK	BCL2	BCR	BRAF	BRCA1	BRCA2	CD74	EGFR	ETV1
ETV4	ETV5	ETV6	EWSR1	EZR	FGFR1	FGFR2	FGFR3	KIT
KMT2A (MLL)	MSH2	MYB	MYC	NOTCH2	NTRK1	NTRK2	NUTM1	PDGFRA
RAF1	RARA	RET	ROS1	RSPO2	SDC4	SLC34A2	TERC*	TERT**
TPRSS2								

*TERC is an NCRNA

**Promoter region of TERT is interrogated

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS

Loss of Heterozygosity (LOH) score

Microsatellite (MS) status

Tumor Mutational Burden (TMB)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

APPENDIX
About FoundationOne®CDx

FoundationOne CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Cipalstraat 3, 2440 Geel, Belgium.


ABOUT FOUNDATIONONE CDx

FoundationOne CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details:
www.rochefoundationmedicine.com/f1cdxtech.

INTENDED USE

FoundationOne®CDx (F1CDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI), tumor mutational burden (TMB), and for selected forms of ovarian cancer, loss of heterozygosity (LOH) score, using DNA isolated from formalin-fixed, paraffin-embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with therapies in accordance with approved therapeutic product labeling. Additionally, F1CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms.

TEST PRINCIPLES

FoundationOne CDx will be performed exclusively as a laboratory service using DNA extracted from formalin-fixed, paraffin-embedded (FFPE) tumor samples. The proposed assay will employ a single DNA extraction method from routine FFPE biopsy or surgical resection specimens, 50-1000 ng of which will undergo whole-genome shotgun library construction and hybridization-based capture of all coding exons from 309 cancer-related genes, one promoter region, one non-coding (ncRNA), and select intronic regions from 34 commonly rearranged genes, 21 of which also include the coding exons. The assay therefore includes detection of alterations in a total of 324 genes.

Using an Illumina® HiSeq platform, hybrid capture-selected libraries will be sequenced to high uniform depth (targeting >500X median coverage with >99% of exons at coverage >100X). Sequence data will be processed using a customized analysis pipeline designed to accurately detect all classes of genomic alterations, including base substitutions, indels, focal copy number amplifications, homozygous gene deletions, and selected genomic rearrangements (e.g., gene fusions). Additionally, genomic signatures including loss of heterozygosity (LOH), microsatellite instability (MSI) and tumor mutational burden (TMB) will be reported.

THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. The F1CDx report may be used as an aid to inform molecular eligibility for clinical trials. Note: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

Diagnostic Significance

FoundationOne CDx identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

Qualified Alteration Calls (Equivocal and Subclonal)

An alteration denoted as "amplification - equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne CDx for identifying a copy number amplification is four (4) for ERBB2 and six (6) for all other genes. Conversely, an alteration denoted as "loss - equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is one that the FoundationOne CDx analytical methodology has identified as being present in <10% of the assayed tumor DNA.

Ranking of Therapies and Clinical Trials
Ranking of Therapies in Summary Table

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

Ranking of Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) (www.nccn.org). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2022. All rights reserved. To view the most recent and complete version of the guidelines, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

Limitations

1. In the fractional-based MSI algorithm, a tumor specimen will be categorized as MSI-H, MSS, or MS-Equivocal according to the fraction of microsatellite loci determined to be altered or unstable (i.e., the fraction unstable loci score). In the F1CDx assay, MSI is evaluated based on a genome-wide analysis across >2000 microsatellite loci. For a given microsatellite locus, non-somatic alleles are discarded, and the microsatellite is categorized as unstable if remaining alleles differ from the reference genome. The final fraction unstable loci score is calculated as the number of unstable microsatellite loci divided by the number of evaluable microsatellite loci. The MSI-H and MSS cut-off thresholds were determined by analytical concordance to a PCR comparator assay using a pan-tumor FFPE tissue sample set. Patients with results categorized as "MS-

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

APPENDIX

About FoundationOne®CDx

- Stable" with median exon coverage <300X, "MS-Equivocal," or "Cannot Be Determined" should receive confirmatory testing using a validated orthogonal (alternative) method.
- TMB by F1CDx is determined by counting all synonymous and non-synonymous variants present at 5% allele frequency or greater (after filtering) and the total number is reported as mutations per megabase (mut/Mb) unit. Observed TMB is dependent on characteristics of the specific tumor focus tested for a patient (e.g., primary vs. metastatic, tumor content) and the testing platform used for the detection; therefore, observed TMB results may vary between different specimens for the same patient and between detection methodologies employed on the same sample. The TMB calculation may differ from TMB calculations used by other assays depending on variables such as the amount of genome interrogated, percentage of tumor, assay limit of detection (LoD), filtering of alterations included in the score, and the read depth and other bioinformatic test specifications. Refer to the SSED for a detailed description of these variables in FMI's TMB calculation https://www.accessdata.fda.gov/cdrh_docs/pdf17/P170019B.pdf. The clinical validity of TMB defined by this panel has been established for TMB as a qualitative output for a cut-off of 10 mutations per megabase but has not been established for TMB as a quantitative score.
 - The LOH score is determined by analyzing SNPs spaced at 1Mb intervals across the genome on the FoundationOne CDx test and extrapolating an LOH profile, excluding arm- and chromosome-wide LOH segments. Detection of LOH has been verified only for ovarian cancer patients, and the LOH score result may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas. The LOH score will be reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine LOH. Performance of the LOH classification has not been established for samples below 35% tumor content. There may be potential interference of ethanol with LOH detection. The interfering effects of xylene, hemoglobin, and triglycerides on the LOH score have not been demonstrated.
 - Alterations reported may include somatic (not inherited) or germline (inherited) alterations; however, the test does not distinguish between germline and somatic alterations. The test does not provide information about susceptibility.
 - Biopsy may pose a risk to the patient when archival tissue is not available for use with the assay. The patient's physician should determine

whether the patient is a candidate for biopsy.

- Reflex testing to an alternative FDA approved companion diagnostic should be performed for patients who have an *ERBB2* amplification result detected with copy number equal to 4 (baseline ploidy of tumor +2) for confirmatory testing. While this result is considered negative by FoundationOne®CDx (F1CDx), in a clinical concordance study with an FDA approved FISH test, 70% (7 out of 10 samples) were positive, and 30% (3 out of 10 samples) were negative by the FISH test with an average ratio of 2.3. The frequency of *ERBB2* copy number 4 in breast cancer is estimated to be approximately 2%. Multiple references listed in <https://www.mycancergenome.org/content/disease/breast-cancer/ERBB2/238/> report the frequency of *HER2* overexpression as 20% in breast cancer. Based on the F1CDx *HER2* CDx concordance study, approximately 10% of *HER2* amplified samples had copy number 4. Thus, total frequency is conservatively estimated to be approximately 2%.

REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic, nontargeted treatment, germline, or clonal hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

VARIANT ALLELE FREQUENCY

Variant Allele Frequency (VAF) represents the fraction of sequencing reads in which the variant is observed. This attribute is not taken into account for therapy inclusion, clinical trial matching, or interpretive content. Caution is recommended in interpreting VAF to indicate the potential germline or somatic origin of an alteration, recognizing that tumor fraction and tumor ploidy of samples may vary.

Precision of VAF for base substitutions and indels

BASE SUBSTITUTIONS	%CV*
Repeatability	5.11 - 10.40
Reproducibility	5.95 - 12.31
INDELS	%CV*
Repeatability	6.29 - 10.00
Reproducibility	7.33 - 11.71

*Interquartile Range = 1st Quartile to 3rd Quartile

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 291656669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >10%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM*, *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *CHEK2*, *FH*, *FLCN*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *PALB2*, *PMS2*, *POLE*, *RAD51C*, *RAD51D*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *TSC2*, and *VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1*, *CBL*, *DNMT3A*, *IDH2*, *JAK2*, *KMT2D* (*MLL2*), *MPL*, *MYD88*, *SF3B1*, *TET2*, and *U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

APPENDIX

About FoundationOne®CDx

cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne CDx.

TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. FoundationOne CDx is performed using DNA derived from tumor, and as such germline events may not be reported.

SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
mut/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

MR Suite Version 6.1.0

The median exon coverage for this sample is 697x

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # **ORD-1341404-01**
APPENDIX **References**

1. Gatalica Z, et al. Cancer Epidemiol. Biomarkers Prev. (2014) PMID: 25392179
2. Kroemer G, et al. Oncoimmunology (2015) PMID: 26140250
3. Lal N, et al. Oncoimmunology (2015) PMID: 25949894
4. Le DT, et al. N. Engl. J. Med. (2015) PMID: 26028255
5. Ayers et al., 2016; ASCO-SITC Abstract P60
6. Martinez R, et al. Oncology (2004) PMID: 15331927
7. Martinez R, et al. J. Cancer Res. Clin. Oncol. (2005) PMID: 15672285
8. Martinez R, et al. Cancer Genet. Cytogenet. (2007) PMID: 17498554
9. Szybka M, et al. Clin. Neuropathol. () PMID: 12908754
10. Kocarnik JM, et al. Gastroenterol Rep (Oxf) (2015) PMID: 26337942
11. You JF, et al. Br. J. Cancer (2010) PMID: 21081928
12. Bairwa NK, et al. Methods Mol. Biol. (2014) PMID: 24623249
13. Boland CR, et al. Cancer Res. (1998) PMID: 9823339
14. Pawlik TM, et al. Dis. Markers (2004) PMID: 15528785
15. Boland CR, et al. Gastroenterology (2010) PMID: 20420947
16. Samstein RM, et al. Nat. Genet. (2019) PMID: 30643254
17. Goodman AM, et al. Mol. Cancer Ther. (2017) PMID: 28835386
18. Goodman AM, et al. Cancer Immunol Res (2019) PMID: 31405947
19. Cristescu R, et al. Science (2018) PMID: 30309915
20. Ready N, et al. J. Clin. Oncol. (2019) PMID: 30785829
21. Hellmann MD, et al. N. Engl. J. Med. (2018) PMID: 29658845
22. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
23. Hellmann MD, et al. Cancer Cell (2018) PMID: 29731394
24. Rozeman EA, et al. Nat Med (2021) PMID: 33558721
25. Sharma P, et al. Cancer Cell (2020) PMID: 32916128
26. Zhao J, et al. Nat. Med. (2019) PMID: 30742119
27. Thout M, et al. Nature (2020) PMID: 32322066
28. Bouffett E, et al. J. Clin. Oncol. (2016) PMID: 27001570
29. Johanns TM, et al. Cancer Discov (2016) PMID: 27683556
30. Lukas RV, et al. J. Neurooncol. (2018) PMID: 30073642
31. Chalmers ZR, et al. Genome Med (2017) PMID: 28420421
32. Patel RR, et al. Pediatr Blood Cancer (2020) PMID: 32386112
33. Johnson A, et al. Oncologist (2017) PMID: 28912153
34. Draaisma K, et al. Acta Neuropathol Commun (2015) PMID: 26699864
35. Wang L, et al. BMC Cancer (2020) PMID: 32164609
36. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
37. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
38. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
39. Rizvi NA, et al. Science (2015) PMID: 25765070
40. Johnson BE, et al. Science (2014) PMID: 24336570
41. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
42. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
43. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
44. Heitzer E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
45. Nature (2012) PMID: 22810696
46. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
47. van den Bent MJ, et al. J Clin Oncol (2009) PMID: 19204207
48. Haas-Kogan DA, et al. J Natl Cancer Inst (2005) PMID: 15956649
49. Brown PD, et al. J Clin Oncol (2008) PMID: 18955445
50. Preusser M, et al. J Neurooncol (2008) PMID: 18458820
51. Wen PY, et al. Neuro-oncology (2014) PMID: 24470557
52. Gallego O, et al. J Neurooncol (2014) PMID: 24352766
53. Uhm JH, et al. Int J Radiat Oncol Biol Phys (2011) PMID: 20510539
54. Doyle SP, et al. Oxf Med Case Reports (2018) PMID: 30410775
55. D'Alessandris QG, et al. Acta Neurochir (Wien) (2013) PMID: 23132371
56. Custodio A, et al. Clin Transl Oncol (2010) PMID: 20462843
57. D'Alessandris QG, et al. Acta Neurochir (Wien) (2018) PMID: 30306271
58. Mellinghoff IK, et al. N. Engl. J. Med. (2005) PMID: 16282176
59. Mellinghoff IK, et al. Clin Cancer Res (2007) PMID: 17255257
60. Arif SH, et al. Asian J Neurosurg () PMID: 29492119
61. Reardon DA, et al. Neuro-oncology (2015) PMID: 25140039
62. Alshami J, et al. Oncotarget (2015) PMID: 26423602
63. Sepúlveda-Sánchez JM, et al. Neuro-oncology (2017) PMID: 28575464
64. Chi AS, et al. JCO Precis Oncol (2020) PMID: 32923886
65. Makhlin I, et al. CNS Oncol (2019) PMID: 31769726
66. Tanaka S, et al. Sci Rep (2019) PMID: 30644426
67. Blumenthal DT, et al. J. Neurooncol. (2016) PMID: 27531351
68. Franceschi E, et al. Br. J. Cancer (2007) PMID: 17353924
69. Chakravarti A, et al. Int. J. Radiat. Oncol. Biol. Phys. (2013) PMID: 23182702
70. Hegi ME, et al. Mol. Cancer Ther. (2011) PMID: 21471286
71. Jiang Z, et al. PLoS ONE (2013) PMID: 23441167
72. Licitra L, et al. Ann. Oncol. (2011) PMID: 21048039
73. Smyth EC, et al. Gut (2021) PMID: 33199443
74. Lassman et al., 2019; Neuro-Oncology Abstract ACTR-21
75. Van Den Bent M, et al. Neuro Oncol (2020) PMID: 31747009
76. Jonsson P, et al. Clin. Cancer Res. (2019) PMID: 31263031
77. Brennan CW, et al. Cell (2013) PMID: 24120142
78. Ceccarelli M, et al. Cell (2016) PMID: 26824661
79. Thomas AA, et al. Neuro-oncology (2017) PMID: 28472509
80. Lee JC, et al. PLoS Med. (2006) PMID: 17177598
81. Vivanco I, et al. Cancer Discov (2012) PMID: 22588883
82. Yan et al. 2020; DOI:10.1200/PO.19.00385
83. Srividya MR, et al. J. Clin. Pathol. (2010) PMID: 20702468
84. Das P, et al. J Clin Neurosci (2011) PMID: 20888234
85. Smith JS, et al. J. Natl. Cancer Inst. (2001) PMID: 11504770
86. Ciardiello F, et al. N. Engl. J. Med. (2008) PMID: 18337605
87. Liang Z, et al. BMC Cancer (2010) PMID: 20637128
88. Bhargava R, et al. Mod. Pathol. (2005) PMID: 15920544
89. Yang YL, et al. Chin. Med. J. (2012) PMID: 22490401
90. Cho J, et al. Cancer Res. (2011) PMID: 22001862
91. Imielinski M, et al. Cell (2012) PMID: 22980975
92. Pines G, et al. Oncogene (2010) PMID: 20676128
93. Wang Q, et al. Exp. Cell Res. (2007) PMID: 17643422
94. Frattini V, et al. Nat. Genet. (2013) PMID: 23917401
95. Konduri K, et al. Cancer Discov (2016) PMID: 27102076
96. Louis DN, et al. Neuro Oncol (2021) PMID: 34185076
97. Fritsch C, et al. Mol. Cancer Ther. (2014) PMID: 24608574
98. Juric D, et al. J. Clin. Oncol. (2018) PMID: 29401002
99. Gallant JN, et al. NPJ Precis Oncol (2019) PMID: 30793038
100. Delestre F, et al. Sci Transl Med (2021) PMID: 34613809
101. André F, et al. N. Engl. J. Med. (2019) PMID: 31091374
102. Smyth LM, et al. NPJ Breast Cancer (2021) PMID: 33863913
103. Park HS, et al. PLoS ONE (2016) PMID: 27105424
104. Lim SM, et al. Oncotarget (2016) PMID: 26859683
105. Hou MM, et al. Oncotarget (2014) PMID: 25426553
106. Varnier R, et al. Eur J Cancer (2019) PMID: 31351267
107. Janku F, et al. Cell Rep (2014) PMID: 24440717
108. Moroney J, et al. Clin. Cancer Res. (2012) PMID: 22927482
109. Basho RK, et al. JAMA Oncol (2017) PMID: 27893038
110. Moroney JW, et al. Clin. Cancer Res. (2011) PMID: 21890452
111. Krop et al., 2018; ASCO Abstract 101
112. Pascual J, et al. Cancer Discov (2021) PMID: 32958578
113. Dolly SO, et al. Clin. Cancer Res. (2016) PMID: 26787751
114. Canaud et al., 2021; ESMO Abstract LBA23
115. Aust Fam Physician (1986) PMID: 2941002
116. Santin AD, et al. Gynecol Oncol Rep (2020) PMID: 31934607
117. Patnaik A, et al. Ann. Oncol. (2016) PMID: 27672108
118. Gallia GL, et al. Mol. Cancer Res. (2006) PMID: 17050665
119. Broderick DK, et al. Cancer Res. (2004) PMID: 15289301
120. El-Habr EA, et al. Clin. Neuropathol. () PMID: 20569675
121. Derakhshandeh-Peykar P, et al. J. Neurogenet. (2011) PMID: 22026810
122. Nature (2008) PMID: 18772890
123. Tanaka S, et al. Acta Neuropathol Commun (2019) PMID: 31036078
124. Samuels Y, et al. Cancer Cell (2005) PMID: 15950905
125. Nat. Rev. Cancer (2009) PMID: 19629070
126. Rivière JB, et al. Nat. Genet. (2012) PMID: 22729224
127. Rudd ML, et al. Clin. Cancer Res. (2011) PMID: 21266528
128. Huchtagowder V, et al. Clin. Genet. (2017) PMID: 27307077
129. Kuentz P, et al. Genet. Med. (2017) PMID: 28151489
130. Kang S, et al. Proc. Natl. Acad. Sci. U.S.A. (2005) PMID: 15647370
131. Ikenoue T, et al. Cancer Res. (2005) PMID: 15930273
132. Gymnopoulou M, et al. Proc. Natl. Acad. Sci. U.S.A. (2007) PMID: 17376864
133. Horn S, et al. Oncogene (2008) PMID: 18317450
134. Hon WC, et al. Oncogene (2012) PMID: 22120714
135. Burke JE, et al. Proc. Natl. Acad. Sci. U.S.A. (2012) PMID: 22949682
136. Wu H, et al. Proc. Natl. Acad. Sci. U.S.A. (2009) PMID: 19915146
137. Laurenti R, et al. Rev Saude Publica (1990) PMID: 2103068
138. Dan S, et al. Cancer Res. (2010) PMID: 20530683
139. Oda K, et al. Cancer Res. (2008) PMID: 18829572
140. Zhao L, et al. Oncogene (2008) PMID: 18794883
141. Lui VW, et al. Cancer Discov (2013) PMID: 23619167
142. Ross RL, et al. Oncogene (2013) PMID: 22430209
143. Shibata T, et al. Cancer Lett. (2009) PMID: 19394761
144. Dogruluk T, et al. Cancer Res. (2015) PMID: 26627007
145. Croessmann S, et al. Clin. Cancer Res. (2018) PMID: 29284706
146. Ng PK, et al. Cancer Cell (2018) PMID: 29533785
147. Spangle JM, et al. (2020) PMID: 32929011

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

APPENDIX
References

148. Chen L, et al. Nat Commun (2018) PMID: 29636477
149. Jin N, et al. J Clin Invest (2021) PMID: 34779417
150. Konecny GE, et al. Clin. Cancer Res. (2011) PMID: 21278246
151. Katsumi Y, et al. Biochem. Biophys. Res. Commun. (2011) PMID: 21871868
152. Cen L, et al. Neuro-oncology (2012) PMID: 22711607
153. Logan JE, et al. Anticancer Res. (2013) PMID: 23898052
154. Elvin JA, et al. Oncologist (2017) PMID: 28283584
155. Gao J, et al. Curr Oncol (2015) PMID: 26715889
156. Gopalan et al., 2014; ASCO Abstract 8077
157. Peguero et al., 2016; ASCO Abstract 2528
158. Konecny et al., 2016; ASCO Abstract 5557
159. DeMichele A, et al. Clin. Cancer Res. (2015) PMID: 25501126
160. Finn RS, et al. Lancet Oncol. (2015) PMID: 25524798
161. Infante JR, et al. Clin. Cancer Res. (2016) PMID: 27542767
162. Johnson DB, et al. Oncologist (2014) PMID: 24797823
163. Van Maerken T, et al. Mol. Cancer Ther. (2011) PMID: 21460101
164. Gamble LD, et al. Oncogene (2012) PMID: 21725357
165. Shapiro et al., 2013; ASCO Abstract 2500
166. Flaherty KT, et al. Clin. Cancer Res. (2012) PMID: 22090362
167. Dickson MA, et al. J. Clin. Oncol. (2013) PMID: 23569312
168. Cerami E, et al. Cancer Discov (2012) PMID: 22588877
169. Gao J, et al. Sci Signal (2013) PMID: 23550210
170. Verhaak RG, et al. Cancer Cell (2010) PMID: 20129251
171. Sottoriva A, et al. Proc. Natl. Acad. Sci. U.S.A. (2013) PMID: 23412337
172. Weber RG, et al. Oncogene (2007) PMID: 16909113
173. Nakamura M, et al. Brain Pathol. (2001) PMID: 11303791
174. Chakravarti A, et al. Clin. Cancer Res. (2001) PMID: 11489817
175. Feng J, et al. Cancer (2012) PMID: 21713760
176. Raabe EH, et al. Clin. Cancer Res. (2011) PMID: 21636552
177. Liu W, et al. J. Exp. Clin. Cancer Res. (2011) PMID: 21843312
178. Quelle DE, et al. Cell (1995) PMID: 8521522
179. Mutat. Res. (2005) PMID: 15878778
180. Gazzieri S, et al. Oncogene (1998) PMID: 9484839
181. Oncogene (1999) PMID: 10498883
182. Sherr CJ, et al. Cold Spring Harb. Symp. Quant. Biol. (2005) PMID: 16869746
183. Ozanne P, et al. Int. J. Cancer (2010) PMID: 20549699
184. Ruas M, et al. Oncogene (1999) PMID: 10498896
185. Jones R, et al. Cancer Res. (2007) PMID: 17909018
186. Haferkamp S, et al. Aging Cell (2008) PMID: 18843795
187. Huot TJ, et al. Mol. Cell. Biol. (2002) PMID: 12417717
188. Rizos H, et al. J. Biol. Chem. (2001) PMID: 11518711
189. Gombart AF, et al. Leukemia (1997) PMID: 9324288
190. Yang R, et al. Cancer Res. (1995) PMID: 7780957
191. Parry D, et al. Mol. Cell. Biol. (1996) PMID: 8668202
192. Greenblatt MS, et al. Oncogene (2003) PMID: 12606942
193. Yarbrough WG, et al. J. Natl. Cancer Inst. (1999) PMID: 10491434
194. Poi MJ, et al. Mol. Carcinog. (2001) PMID: 11255261
195. Byeon IJ, et al. Mol. Cell (1998) PMID: 9660926
196. Kannengiesser C, et al. Hum. Mutat. (2009) PMID: 19260062
197. Lal G, et al. Genes Chromosomes Cancer (2000) PMID: 10719365
198. Koh J, et al. Nature (1995) PMID: 7777061
199. McKenzie HA, et al. Hum. Mutat. (2010) PMID: 20340136
200. Miller PJ, et al. Hum. Mutat. (2011) PMID: 21462282
201. Kutscher CL, et al. Physiol. Behav. (1977) PMID: 905385
202. Scaini MC, et al. Hum. Mutat. (2014) PMID: 24659262
203. Jenkins NC, et al. J. Invest. Dermatol. (2013) PMID: 23190892
204. Walker GJ, et al. Int. J. Cancer (1999) PMID: 10389768
205. Rutter JL, et al. Oncogene (2003) PMID: 12853981
206. Itahana K, et al. Cancer Cell (2008) PMID: 18538737
207. Zhang Y, et al. Mol. Cell (1999) PMID: 10360174
208. Zhang Y, et al. Cell (1998) PMID: 9529249
209. Jafri M, et al. Cancer Discov (2015) PMID: 25873077
210. Whelan AJ, et al. N Engl J Med (1995) PMID: 7666917
211. Adv Exp Med Biol (2010) PMID: 20687502
212. Hogg D, et al. J Cutan Med Surg (1998) PMID: 9479083
213. De Unamuno B, et al. Melanoma Res (2018) PMID: 29543703
214. Soura E, et al. J Am Acad Dermatol (2016) PMID: 26892650
215. Huerta C, et al. Acta Derm Venereol (2018) PMID: 29405243
216. Kaufman DK, et al. Neurology (1993) PMID: 8414022
217. Bahau M, et al. Cancer Res (1998) PMID: 9622062
218. Chan AK, et al. Clin Neuropathol () PMID: 28699883
219. Tate JG, et al. Nucleic Acids Res. (2019) PMID: 30371878
220. Hegde M, et al. Genet. Med. (2014) PMID: 24310308
221. Aretz S, et al. Eur. J. Hum. Genet. (2013) PMID: 22872101
222. Win AK, et al. Gastroenterology (2014) PMID: 24044654
223. Lubbe SJ, et al. J. Clin. Oncol. (2009) PMID: 19620482
224. Jones N, et al. Gastroenterology (2009) PMID: 19394335
225. Nielsen M, et al. J. Natl. Cancer Inst. (2010) PMID: 21044966
226. David SS, et al. Nature (2007) PMID: 17581577
227. Molatore S, et al. Hum. Mutat. (2010) PMID: 19953527
228. Kundu S, et al. DNA Repair (Amst.) (2009) PMID: 19836313
229. D'Agostino VG, et al. DNA Repair (Amst.) (2010) PMID: 20418187
230. Ali M, et al. Gastroenterology (2008) PMID: 18534194
231. Landrum MJ, et al. Nucleic Acids Res. (2018) PMID: 29165669
232. Sampson JR, et al. Lancet (2003) PMID: 12853198
233. Sieber OM, et al. N. Engl. J. Med. (2003) PMID: 12606733
234. Al-Tassan N, et al. Nat. Genet. (2002) PMID: 11818965
235. Rennert G, et al. Cancer (2012) PMID: 21952991
236. Zhang Y, et al. Cancer Epidemiol. Biomarkers Prev. (2006) PMID: 16492928
237. von der Thüsen JH, et al. J. Clin. Oncol. (2011) PMID: 21189386
238. Casper M, et al. Fam. Cancer (2014) PMID: 24420788
239. Smith LM, et al. Pancreatol (2009) PMID: 20110747
240. Nat Rev Clin Oncol (2017) PMID: 27245281
241. Duperret EK, et al. Mol Ther (2018) PMID: 29249395
242. Chiappori AA, et al. Ann Oncol (2015) PMID: 25467017
243. Killela PJ, et al. Proc. Natl. Acad. Sci. U.S.A. (2013) PMID: 23530248
244. Killela PJ, et al. Oncotarget (2014) PMID: 24722048
245. Nonoguchi N, et al. Acta Neuropathol. (2013) PMID: 23955565
246. Liu X, et al. Cell Cycle (2013) PMID: 23603989
247. Koelsche C, et al. Acta Neuropathol. (2013) PMID: 24154961
248. Arita H, et al. Acta Neuropathol. (2013) PMID: 23764841
249. Reitman ZJ, et al. Acta Neuropathol. (2013) PMID: 24217890
250. Shay JW, et al. Semin. Cancer Biol. (2011) PMID: 22015685
251. Shay JW, et al. Eur. J. Cancer (1997) PMID: 9282118
252. Kim NW, et al. Science (1994) PMID: 7605428
253. Hanahan D, et al. Cell (2000) PMID: 10647931
254. Horn S, et al. Science (2013) PMID: 23348503
255. Huang FW, et al. Science (2013) PMID: 23348506
256. Vinagre J, et al. Nat Commun (2013) PMID: 23887589
257. Weller M, et al. Nat Rev Clin Oncol (2021) PMID: 33293629
258. Hirai H, et al. Cancer Biol. Ther. (2010) PMID: 20107315
259. Bridges KA, et al. Clin. Cancer Res. (2011) PMID: 21799033
260. Rajeshkumar NV, et al. Clin. Cancer Res. (2011) PMID: 21389100
261. Osman AA, et al. Mol. Cancer Ther. (2015) PMID: 25504633
262. Xu L, et al. Mol. Cancer Ther. (2002) PMID: 12489850
263. Xu L, et al. Mol. Med. (2001) PMID: 11713371
264. Camp ER, et al. Cancer Gene Ther. (2013) PMID: 23470564
265. Kim SS, et al. Nanomedicine (2015) PMID: 25240597
266. Pirolo KF, et al. Mol. Ther. (2016) PMID: 27357628
267. Hajdenberg et al., 2012; ASCO Abstract e15010
268. Leijen S, et al. J. Clin. Oncol. (2016) PMID: 27601554
269. Moore et al., 2019; ASCO Abstract 5513
270. Leijen S, et al. J. Clin. Oncol. (2016) PMID: 27998224
271. Oza et al., 2015; ASCO Abstract 5506
272. Lee J, et al. Cancer Discov (2019) PMID: 31315834
273. Méndez E, et al. Clin. Cancer Res. (2018) PMID: 29535125
274. Seligmann JF, et al. J Clin Oncol (2021) PMID: 34538072
275. Lehmann S, et al. J. Clin. Oncol. (2012) PMID: 22965953
276. Mohell N, et al. Cell Death Dis (2015) PMID: 26086967
277. Fransson Å, et al. J Ovarian Res (2016) PMID: 27179933
278. Gourley et al., 2016; ASCO Abstract 5571
279. Kwok M, et al. Blood (2016) PMID: 26563332
280. Boudny M, et al. Haematologica (2019) PMID: 30975914
281. Dillon MT, et al. Mol. Cancer Ther. (2017) PMID: 28062704
282. Middleton FK, et al. Cancers (Basel) (2018) PMID: 30127241
283. Jha P, et al. Diagn. Mol. Pathol. (2011) PMID: 22089350
284. Uno M, et al. Cancer Lett. (2005) PMID: 15914282
285. Uno M, et al. Int. J. Biol. Markers () PMID: 16711514
286. Lass U, et al. PLoS ONE (2012) PMID: 22844452
287. Faria MH, et al. APMS (2012) PMID: 23009112
288. Milinkovic V, et al. PLoS ONE (2013) PMID: 24358143
289. Galat TF, et al. PLoS ONE (2013) PMID: 23613880
290. Schmidt MC, et al. J. Neuropathol. Exp. Neurol. (2002) PMID: 11939587
291. Nozaki M, et al. Neuro-oncology (1999) PMID: 11550308
292. Brown CJ, et al. Nat. Rev. Cancer (2009) PMID: 19935675
293. Joerger AC, et al. Annu. Rev. Biochem. (2008) PMID: 18410249
294. Kato S, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) PMID: 12826609
295. Kamada R, et al. J. Biol. Chem. (2011) PMID: 20978130
296. Zerdoumy Y, et al. Hum. Mol. Genet. (2017) PMID: 28472496
297. Yamada H, et al. Carcinogenesis (2007) PMID: 17690113
298. Bougeard G, et al. J. Clin. Oncol. (2015) PMID: 26014290
299. Sorrell AD, et al. Mol Diagn Ther (2013) PMID: 23355100
300. Nichols KE, et al. Cancer Epidemiol. Biomarkers Prev. (2001) PMID: 11219776
301. Kleihues P, et al. Am. J. Pathol. (1997) PMID: 9006316

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1341404-01

APPENDIX
References

302. Gonzalez KD, et al. J. Clin. Oncol. (2009) pmid: 19204208
303. Lalloo F, et al. Lancet (2003) pmid: 12672316
304. Mandelker D, et al. Ann. Oncol. (2019) pmid: 31050713
305. Jaiswal S, et al. N. Engl. J. Med. (2014) pmid: 25426837
306. Genovese G, et al. N. Engl. J. Med. (2014) pmid: 25426838
307. Xie M, et al. Nat. Med. (2014) pmid: 25326804
308. Acuna-Hidalgo R, et al. Am. J. Hum. Genet. (2017) pmid: 28669404
309. Severson EA, et al. Blood (2018) pmid: 29678827
310. Fuster JJ, et al. Circ. Res. (2018) pmid: 29420212
311. Hematology Am Soc Hematol Educ Program (2018) pmid: 30504320
312. Chabon JJ, et al. Nature (2020) pmid: 32269342
313. Razavi P, et al. Nat. Med. (2019) pmid: 31768066
314. Hasselbalch B, et al. Neuro-oncology (2010) pmid: 20406901
315. Gajadhar AS, et al. Mol. Cancer Res. (2012) pmid: 22232519
316. Lv S, et al. Int. J. Oncol. (2012) pmid: 22752145
317. Philip PA, et al. J. Clin. Oncol. (2010) pmid: 20606093
318. Slovin SF, et al. Clin Genitourin Cancer (2009) pmid: 19815486
319. Falchook GS, et al. Oncotarget (2013) pmid: 23435217
320. Janku F, et al. Cancer Res. (2013) pmid: 23066039
321. Janku F, et al. J. Clin. Oncol. (2012) pmid: 22271473
322. Janku F, et al. Mol. Cancer Ther. (2011) pmid: 21216929
323. Moulder S, et al. Ann. Oncol. (2015) pmid: 25878190
324. Byeon et al., 2020; doi: 10.21037/tcr.2020.04.07
325. Gojo J, et al. Front Oncol (2019) pmid: 31998633
326. McNall-Knapp et al., 2020; SNO Abstract LGG-47
327. Hainsworth JD, et al. Clin Adv Hematol Oncol (2012) pmid: 22706484
328. Ma DJ, et al. Neuro-oncology (2015) pmid: 25526733
329. Mason WP, et al. Invest New Drugs (2012) pmid: 22160854
330. Kreisl TN, et al. J. Neurooncol. (2009) pmid: 19018475
331. Segal et al., 2016; ISPNO Abstract EPT-21
332. Tolcher AW, et al. Ann. Oncol. (2015) pmid: 25344362
333. Patterson et al., 2018; AACR Abstract 3891
334. Whittle JR, et al. J Clin Neurosci (2015) pmid: 26279503
335. Douillard JY, et al. Ann. Oncol. (2014) pmid: 24718886
336. Price TJ, et al. Lancet Oncol. (2014) pmid: 24739896
337. Van Cutsem E, et al. J. Clin. Oncol. (2007) pmid: 17470858
338. Vermorken JB, et al. Lancet Oncol (2013) pmid: 23746666
339. Wirth LJ, et al. Ann. Oncol. (2010) pmid: 19892746
340. Siano M, et al. Oncologist (2017) pmid: 28592616
341. Mesia R, et al. Lancet Oncol (2015) pmid: 25596660
342. Giral J, et al. Lancet Oncol (2015) pmid: 25596659
343. Siu LL, et al. JAMA Oncol (2016) pmid: 27930762
344. Waddell T, et al. Lancet Oncol. (2013) pmid: 23594787
345. Crawford J, et al. J Thorac Oncol (2013) pmid: 24389433
346. Schuette W, et al. Clin Lung Cancer (2015) pmid: 26094080
347. Leone F, et al. Cancer (2016) pmid: 26540314
348. Vogel A, et al. Eur J Cancer (2018) pmid: 29413685
349. Rowinsky EK, et al. J. Clin. Oncol. (2004) pmid: 15210739
350. Sarkaria JN, et al. Clin. Cancer Res. (2010) pmid: 20921209
351. Lee EQ, et al. Neuro-oncology (2012) pmid: 23099651
352. Lassen U, et al. Anticancer Res. (2013) pmid: 23564811
353. Georger B, et al. Eur. J. Cancer (2012) pmid: 22033322

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 18 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531