



Sample Information

Patient Name: 林徐秀英
Gender: Female
ID No.: G200335525
History No.: 6547213
Age: 71

Ordering Doctor: DOC4205A 柯博仲
Ordering REQ.: H45N3FC
Signing in Date: 2023/02/21

Path No.: M112-00032
MP No.: MY23009
Assay: Oncomine Myeloid Assay
Sample Type: Bone Marrow
Bone Marrow Aspirating Date: 2023/02/16

Reporting Doctor: DOC5466K 葉奕成 (Phone: 8#5466)

Note:

Sample Cancer Type: Chronic Myelomonocytic Leukemia

| | | |
|--|-------------|--------------------------|
| Table of Contents | Page | Report Highlights |
| Variants (Exclude variant in Taiwan BioBank with >1% allele frequency) | 2 | 0 Relevant Biomarkers |
| Biomarker Descriptions | 2 | 0 Therapies Available |
| | | 0 Clinical Trials |

Relevant Chronic Myelomonocytic Leukemia Variants

| Gene | Finding |
|-------|---------------|
| ASXL1 | None detected |

Relevant Biomarkers

No clinically significant biomarkers found in this sample.

Prevalent cancer biomarkers without relevant evidence based on included data sources

CSF3R p.(T618I) c.1853C>T

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

DNA Sequence Variants

| Gene | Amino Acid Change | Coding | Variant ID | Locus | Allele Frequency | Transcript | Variant Effect | Coverage |
|-------|-------------------|-----------------------|-------------|---------------|------------------|-------------|----------------------|----------|
| CSF3R | p.(T618I) | c.1853C>T | COSM1737962 | chr1:36933434 | 41.55% | NM_156039.3 | missense | 2000 |
| CSF3R | p.(L827=;V828I) | c.2481_2482delGGinsAA | . | chr1:36932068 | 38.10% | NM_156039.3 | synonymous, missense | 1995 |
| CSF3R | p.(G825=) | c.2475G>A | . | chr1:36932075 | 38.59% | NM_156039.3 | synonymous | 1998 |
| CSF3R | p.(W818*) | c.2453G>A | . | chr1:36932097 | 39.45% | NM_156039.3 | nonsense | 2000 |

Biomarker Descriptions

CSF3R (colony stimulating factor 3 receptor)

Background: The CSF3R gene encodes the colony stimulating factor 3 trans-membrane receptor for the granulocyte colony-stimulating factor (G-CSF) ligand. CSF3R is a class I membrane-bound cytokine receptor, which lacks intrinsic kinase activity and therefore must interact with downstream proteins for activation¹. Upon ligand activation, CSF3R activates downstream oncogenic pathways by interacting with intracellular signaling proteins through its cytoplasmic tyrosine residues, including proteins from the JAK/STAT, MAPK/ERK, and PI3K/AKT pathways^{1,2}. Nonsense and frameshift mutations in CSF3R target and truncate its cytoplasmic tail, subsequently impairing the internalization signal of the receptor and leading to an overexpression CSF3R on the cell surface^{3,4,5}. Missense mutations in the proximal membrane lead to increased dimerization of the receptor, independent of G-CSF binding^{3,4,5}. These mutations promote constitutive oncogenic JAK-STAT signaling, and increase granulocyte proliferation and survival signaling of hematopoietic progenitor cells^{1,6}.

Alterations and prevalence: CSF3R activating mutations are observed in up to 80% of patients with chronic neutrophilic leukemia (CNL), and in up to 59% of patients with atypical chronic myelogenous leukemia (aCML)^{3,7}. CSF3R mutations occur in 0.5-1% of adult acute myeloid leukemia (AML) and in 2.4% of pediatric AML⁷. In solid malignancies, CSF3R mutations are observed in up to 5% of uterine carcinosarcoma and skin cutaneous melanoma^{8,9}. Somatic mutations identified in CNL and AML include T615A, T618I, and T640N missense mutations, as well as truncating mutations at Q749, Q754, Y767, S783, Y787, and P820⁷. T618I mutation is the most frequent variant observed in CNL¹⁰.

Potential relevance: CSF3R activating mutations including T618I are a diagnostic criteria for CNL as defined by the World Health Organization (WHO)¹¹. Mutations in CSF3R are observed in patients with severe congenital neutropenia, which can progress to AML. CSF3R mutations frequently co-occur with CEBPA, and these co-mutations are associated with unfavorable prognosis in AML^{7,12,13}. In independent reports, a CNL patient and an aCML patient with the proximal membrane T618I mutation demonstrated sensitivity to JAK1/2 inhibitor, ruxolitinib^{3,14}.

Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

References

1. Zhang et al. Gain-of-function mutations in granulocyte colony-stimulating factor receptor (CSF3R) reveal distinct mechanisms of CSF3R activation. *J. Biol. Chem.* 2018 May 11;293(19):7387-7396. PMID: 29572350
2. Nicholson et al. Distinct regions of the granulocyte colony-stimulating factor receptor are required for tyrosine phosphorylation of the signaling molecules JAK2, Stat3, and p42, p44MAPK. *Blood.* 1995 Nov 15;86(10):3698-704. PMID: 7579336
3. Maxson et al. Oncogenic CSF3R mutations in chronic neutrophilic leukemia and atypical CML. *N. Engl. J. Med.* 2013 May 9;368(19):1781-90. PMID: 23656643
4. Dong et al. The carboxyl terminus of the granulocyte colony-stimulating factor receptor, truncated in patients with severe congenital neutropenia/acute myeloid leukemia, is required for SH2-containing phosphatase-1 suppression of Stat activation. *J. Immunol.* 2001 Dec 1;167(11):6447-52. PMID: 11714811
5. van et al. G-CSF receptor truncations found in SCN/AML relieve SOCS3-controlled inhibition of STAT5 but leave suppression of STAT3 intact. *Blood.* 2004 Aug 1;104(3):667-74. PMID: 15069015
6. Panopoulos et al. Granulocyte colony-stimulating factor: molecular mechanisms of action during steady state and 'emergency' hematopoiesis. *Cytokine.* 2008 Jun;42(3):277-88. PMID: 18400509
7. Maxson et al. CSF3R mutations have a high degree of overlap with CEBPA mutations in pediatric AML. *Blood.* 2016 Jun 16;127(24):3094-8. PMID: 27143256
8. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
9. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
10. Pardanani et al. CSF3R T618I is a highly prevalent and specific mutation in chronic neutrophilic leukemia. *Leukemia.* 2013 Sep;27(9):1870-3. PMID: 23604229
11. Arber et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood.* 2016 May 19;127(20):2391-405. PMID: 27069254
12. Dong et al. Mutations in the gene for the granulocyte colony-stimulating-factor receptor in patients with acute myeloid leukemia preceded by severe congenital neutropenia. *N. Engl. J. Med.* 1995 Aug 24;333(8):487-93. PMID: 7542747
13. Su et al. CSF3R mutations were associated with an unfavorable prognosis in patients with acute myeloid leukemia with CEBPA double mutations. *Ann. Hematol.* 2019 Jul;98(7):1641-1646. PMID: 31041512
14. Dao et al. Significant clinical response to JAK1/2 inhibition in a patient with CSF3R-T618I-positive atypical chronic myeloid leukemia. *Leuk Res Rep.* 2014;3(2):67-9. PMID: 25180155