



## Sample Information

**Patient Name:** 黃坤雄

**Gender:** Male

**ID No.:** A121626265

**History No.:** 21391140

**Age:** 58

**Ordering Doctor:** DOC3072G 吳佳儒

**Ordering REQ.:** 0AQRKFR

**Signing in Date:** 2020/03/31

**Path No.:** S109-99289

**MP No.:** F2007

**Assay:** Oncomine Focus Assay

**Sample Type:** FFPE

**Block No.:** S109-75343E

**Percentage of tumor cells:** 90%

**Note:**

## Sample Cancer Type: Non-Small Cell Lung Cancer

### Table of Contents

	Page
Variant Details	2
Biomarker Descriptions	3
Relevant Therapy Summary	4
Relevant Therapy Details	9

### Report Highlights

2 Clinically Significant Biomarkers  
 7 Therapies Available  
 114 Clinical Trials

## Relevant Non-Small Cell Lung Cancer Findings

Gene	Finding	Gene	Finding
ALK	Not detected	NTRK1	Not detected
BRAF	Not detected	NTRK2	Not detected
EGFR	<b>EGFR exon 19 deletion, EGFR p.(T790M) c.2369C&gt;T</b>	NTRK3	Not detected
ERBB2	Not detected	RET	Not detected
KRAS	Not detected	ROS1	Not detected
MET	Not detected		



## Clinically Significant Biomarkers

■ Indicated ■ Contraindicated

Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
<b>EGFR exon 19 deletion</b> epidermal growth factor receptor Tier: IA Allele Frequency: 43.98%	<span style="color: green;">■</span> osimertinib <sup>1, 2</sup> <span style="color: green;">■</span> afatinib + cetuximab <span style="color: green;">■</span> bevacizumab* + erlotinib <sup>2</sup> <span style="color: green;">■</span> gefitinib + chemotherapy <span style="color: green;">■</span> bevacizumab + gefitinib <span style="color: green;">■</span> atezolizumab + bevacizumab + chemotherapy	None	94
<b>EGFR p.(T790M) c.2369C&gt;T</b> epidermal growth factor receptor Tier: IA Allele Frequency: 28.11%	<span style="color: green;">■</span> osimertinib <sup>1, 2</sup> <span style="color: green;">■</span> afatinib + cetuximab <span style="color: red;">■</span> gefitinib <sup>2</sup>	None	74

Sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

\* Includes biosimilars

## Tier Criteria Met

Genomic Alteration	Tier Classification for Non-Small Cell Lung Cancer
<b>EGFR exon 19 deletion</b> Tier: IA	IA: Biomarker predicts response or resistance to FDA or EMA approved therapies in this cancer type IA: Biomarker is included in NCCN or ESMO guidelines that predict response or resistance to therapies in this cancer type IIC: Biomarker is an inclusion criteria for clinical trials
<b>EGFR p.(T790M) c.2369C&gt;T</b> Tier: IA	IA: Biomarker predicts response or resistance to FDA or EMA approved therapies in this cancer type IA: Biomarker is included in NCCN or ESMO guidelines that predict response or resistance to therapies in this cancer type IIC: Biomarker is an inclusion criteria for clinical trials

**Reference:** Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

## Variant Details

### DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
EGFR	p.(L747_S752del)	c.2239_2256delTTAA GAGAAGCAACATCT	COSM6255	chr7:55242468	43.98%	NM_005228.4	nonframeshift Deletion	1951
EGFR	p.(T790M)	c.2369C>T	COSM6240	chr7:55249071	28.11%	NM_005228.4	missense	1996
ALK	p.(D1529E)	c.4587C>G	.	chr2:29416366	99.80%	NM_004304.4	missense	1996
ALK	p.(I1461V)	c.4381A>G	.	chr2:29416572	99.95%	NM_004304.4	missense	1994



## Variant Details (continued)

### DNA Sequence Variants (continued)

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
ALK	p.(=)	c.3375C>A	.	chr2:29445458	100.00%	NM_004304.4	synonymous	1994
FGFR3	p.(=)	c.1953G>A	.	chr4:1807894	99.80%	NM_000142.4	synonymous	1995
PDGFRA	p.(=)	c.1701A>G	.	chr4:55141055	99.90%	NM_006206.5	synonymous	2000
PDGFRA	p.(=)	c.2472C>T	.	chr4:55152040	9.60%	NM_006206.5	synonymous	2000
FGFR4	p.(P136L)	c.407C>T	.	chr5:176517797	99.35%	NM_213647.2	missense	2000
FGFR4	p.(=)	c.483A>G	.	chr5:176517985	24.27%	NM_213647.2	synonymous	1998
RET	p.(=)	c.2307G>T	.	chr10:43613843	9.56%	NM_020975.4	synonymous	1997

## Biomarker Descriptions

### EGFR (epidermal growth factor receptor)

**Background:** The EGFR gene encodes the epidermal growth factor receptor (EGFR) tyrosine kinase, a member of the human epidermal growth factor receptor (HER) family. Along with EGFR/ERBB1/HER1, ERBB2/HER2, ERBB3/HER3, and ERBB4/HER4 make up the HER protein family<sup>1</sup>. EGFR ligand induced dimerization results in kinase activation and leads to stimulation of oncogenic signaling pathways including the PI3K/AKT/MTOR and RAS/RAF/MEK/ERK pathways. Activation of these pathways promote cell proliferation, differentiation, and survival<sup>2,3</sup>.

**Alterations and prevalence:** Recurrent somatic mutations in the tyrosine kinase domain of EGFR are observed in approximately 10-20% of lung adenocarcinoma and at higher frequencies in never-smoker, female, and in Asian populations with lung cancer<sup>4,5,6,7</sup>. The most common mutations occur near the ATP-binding pocket of the kinase domain and include short in-frame deletions in exon 19 (EGFR exon 19 deletion) and the L858R amino acid substitution in exon 21<sup>8</sup>. These mutations constitutively activate the EGFR kinase resulting in downstream signaling and represent 80% of the EGFR mutations observed in lung cancer. A second group of recurrent activating mutations that are less common include E709K, G719X, S768I, L861Q, and short in-frame insertions in exon 20<sup>9,10,11,12</sup>. EGFR activating mutations in lung cancer tend to be mutually exclusive to KRAS activating mutations<sup>13</sup>. Although these variants are common in lung cancer, they are rare in other cancer types. In glioblastoma, recurrent activating EGFR mutations in the extracellular domain include R108K, A289V and G598V<sup>8,14</sup>. The recurrent focal amplification of the EGFR gene leads to an increase in expression in several cancer types. EGFR is amplified in up to 30% of glioblastoma, 12% of esophageal cancer, 10% of head and neck cancer, 5% of bladder cancer, and 5% of lung squamous cell carcinoma<sup>5,6,7,14,15</sup>. Deletion of exons 2-7 encoding the extracellular domain of EGFR (EGFRvIII) results in overexpression of a ligand-independent constitutively active protein which is frequently observed in glioblastoma and has been shown to lead to lung cancer development as well as sensitivity to TKIs<sup>16,17,18</sup>.

**Potential clinical relevance:** Erlotinib<sup>19</sup> (2004), afatinib<sup>20</sup> (2013), gefitinib<sup>21</sup> (2015), osimertinib<sup>22</sup> (2015), and dacomitinib<sup>23</sup> (2018) are small molecule TKIs that are FDA approved for non-small cell lung cancer (NSCLC) patients with sensitizing exon 19 deletions and exon 21 L858R mutations. Acquired secondary mutations often confer resistance to first line TKI therapy with the T790M amino acid substitution accounting for 50-60% of cases<sup>8</sup>. Osimertinib is also indicated for NSCLC patients harboring EGFR T790M mutations whose disease has progressed on or after treatment with a first line TKI. EGFR targeting antibodies including cetuximab<sup>24</sup> (2004), panitumumab<sup>25</sup> (2006), and necitumumab<sup>26</sup> (2016) are also under investigation in combination with EGFR-targeting TKIs for efficacy against EGFR mutations. The use of cetuximab in combination with afatinib is currently recommended by the NCCN for patients who have progressed after receiving erlotinib, afatinib, dacomitinib, or gefitinib and chemotherapy<sup>27</sup>.



## Relevant Therapy Summary

● In this cancer type    ○ In other cancer type    ● In this cancer type and other cancer types    ⛔ Contraindicated    ⚠ Both for use and contraindicated    ✕ No evidence

### EGFR exon 19 deletion

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
osimertinib	●	●	●	●	● (IV)
afatinib + cetuximab	✕	●	✕	✕	✕
bevacizumab + erlotinib	✕	✕	●	●	● (II)
bevacizumab (Allergan) + erlotinib	✕	✕	●	✕	✕
atezolizumab + bevacizumab + carboplatin + paclitaxel	✕	✕	✕	●	✕
bevacizumab + gefitinib	✕	✕	✕	●	✕
gefitinib + carboplatin + pemetrexed	✕	✕	✕	●	✕
bevacizumab + osimertinib, osimertinib	✕	✕	✕	✕	● (IV)
icotinib hydrochloride	✕	✕	✕	✕	● (IV)
icotinib hydrochloride, icotinib hydrochloride + chemotherapy	✕	✕	✕	✕	● (IV)
icotinib hydrochloride, radiation therapy	✕	✕	✕	✕	● (IV)
atezolizumab, bevacizumab, chemotherapy	✕	✕	✕	✕	● (III)
bevacizumab + chemotherapy, bevacizumab (Shanghai Hengrui Pharmaceutical) + chemotherapy	✕	✕	✕	✕	● (III)
chemotherapy, durvalumab	✕	✕	✕	✕	● (III)
chemotherapy, icotinib hydrochloride	✕	✕	✕	✕	● (III)
chemotherapy, osimertinib	✕	✕	✕	✕	● (III)
chemotherapy, pembrolizumab	✕	✕	✕	✕	● (III)
icotinib hydrochloride, icotinib hydrochloride + radiation therapy	✕	✕	✕	✕	● (III)
anlotinib hydrochloride + icotinib hydrochloride	✕	✕	✕	✕	● (II)
anlotinib hydrochloride, osimertinib	✕	✕	✕	✕	● (II)
atezolizumab, chemotherapy	✕	✕	✕	✕	● (II)
bevacizumab, osimertinib	✕	✕	✕	✕	● (II)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.



## Relevant Therapy Summary (continued)

● In this cancer type    ○ In other cancer type    ● In this cancer type and other cancer types    ⛔ Contraindicated    ⚠ Both for use and contraindicated    ✕ No evidence

### EGFR exon 19 deletion (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
bintrafusp alfa, chemoradiation therapy, durvalumab	✕	✕	✕	✕	● (II)
BPI-7711	✕	✕	✕	✕	● (II)
chemotherapy, durvalumab, tremelimumab	✕	✕	✕	✕	● (II)
chemotherapy, ramucirumab	✕	✕	✕	✕	● (II)
crizotinib + chemotherapy	✕	✕	✕	✕	● (II)
icotinib hydrochloride + radiation therapy	✕	✕	✕	✕	● (II)
ipilimumab + nivolumab, nivolumab + chemotherapy	✕	✕	✕	✕	● (II)
ipilimumab, nivolumab	✕	✕	✕	✕	● (II)
maiHuatinib	✕	✕	✕	✕	● (II)
osimertinib + radiation therapy	✕	✕	✕	✕	● (II)
osimertinib, osimertinib + chemotherapy	✕	✕	✕	✕	● (II)
osimertinib, radiation therapy	✕	✕	✕	✕	● (II)
osimertinib, savolitinib	✕	✕	✕	✕	● (II)
pembrolizumab + chemotherapy	✕	✕	✕	✕	● (II)
radiation therapy, tyrosine kinase inhibitors	✕	✕	✕	✕	● (II)
sunitinib	✕	✕	✕	✕	● (II)
alflutinib	✕	✕	✕	✕	● (I/II)
anlotinib hydrochloride, CBT-502	✕	✕	✕	✕	● (I/II)
ASK120067	✕	✕	✕	✕	● (I/II)
AZD4205, AZD4205 + osimertinib	✕	✕	✕	✕	● (I/II)
AZD4635 + oleclumab	✕	✕	✕	✕	● (I/II)
CK-101	✕	✕	✕	✕	● (I/II)
DZD-9008	✕	✕	✕	✕	● (I/II)
EMB01	✕	✕	✕	✕	● (I/II)
icotinib hydrochloride + chemotherapy + radiation therapy	✕	✕	✕	✕	● (I/II)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.



## Relevant Therapy Summary (continued)

● In this cancer type    ○ In other cancer type    ● In this cancer type and other cancer types    ⛔ Contraindicated    ⚠ Both for use and contraindicated    ✕ No evidence

### EGFR exon 19 deletion (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
KP-673	✕	✕	✕	✕	● (I/II)
lazertinib	✕	✕	✕	✕	● (I/II)
lerociclib, osimertinib	✕	✕	✕	✕	● (I/II)
TAK788	✕	✕	✕	✕	● (I/II)
U3-1402	✕	✕	✕	✕	● (I/II)
cetuximab, FATE-NK100	✕	✕	✕	✕	● (I)
durvalumab + oleclumab, oleclumab	✕	✕	✕	✕	● (I)
everolimus + neratinib, neratinib + palbociclib, neratinib + trametinib	✕	✕	✕	✕	● (I)
FCN-411	✕	✕	✕	✕	● (I)
JNJ-61186372	✕	✕	✕	✕	● (I)
navitoclax, osimertinib	✕	✕	✕	✕	● (I)
nazartinib, trametinib	✕	✕	✕	✕	● (I)
necitumumab, osimertinib	✕	✕	✕	✕	● (I)
osimertinib + ramucirumab	✕	✕	✕	✕	● (I)
osimertinib, osimertinib + radiation therapy	✕	✕	✕	✕	● (I)
osimertinib, telisotuzumab vedotin	✕	✕	✕	✕	● (I)
pirotinib	✕	✕	✕	✕	● (I)
SH-1028	✕	✕	✕	✕	● (I)
TNO-155	✕	✕	✕	✕	● (I)
TP-0903	✕	✕	✕	✕	● (I)
tyrosine kinase inhibitors, tyrosine kinase inhibitors + chemotherapy	✕	✕	✕	✕	● (I)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.



## Relevant Therapy Summary (continued)

● In this cancer type    ○ In other cancer type    ⓘ In this cancer type and other cancer types    ⚡ Contraindicated    ⚠ Both for use and contraindicated    ✕ No evidence

### EGFR p.(T790M) c.2369C>T

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
osimertinib	●	●	●	●	● (IV)
afatinib + cetuximab	✕	●	✕	✕	✕
gefitinib	✕	✕	⚡	✕	✕
anlotinib hydrochloride, osimertinib	✕	✕	✕	✕	● (IV)
icotinib hydrochloride, radiation therapy	✕	✕	✕	✕	● (IV)
atezolizumab, bevacizumab, chemotherapy	✕	✕	✕	✕	● (III)
bevacizumab (Innovent Biologics), chemotherapy, sintilimab	✕	✕	✕	✕	● (III)
bevacizumab + chemotherapy, bevacizumab (Shanghai Hengrui Pharmaceutical) + chemotherapy	✕	✕	✕	✕	● (III)
chemotherapy, durvalumab	✕	✕	✕	✕	● (III)
chemotherapy, icotinib hydrochloride	✕	✕	✕	✕	● (III)
chemotherapy, osimertinib	✕	✕	✕	✕	● (III)
chemotherapy, toripalimab	✕	✕	✕	✕	● (III)
apatinib + chemotherapy	✕	✕	✕	✕	● (II)
avitinib	✕	✕	✕	✕	● (II)
bevacizumab, osimertinib	✕	✕	✕	✕	● (II)
BPI-7711	✕	✕	✕	✕	● (II)
chemotherapy, durvalumab, tremelimumab	✕	✕	✕	✕	● (II)
chemotherapy, ramucirumab	✕	✕	✕	✕	● (II)
D-0316	✕	✕	✕	✕	● (II)
ipilimumab + nivolumab, nivolumab + chemotherapy	✕	✕	✕	✕	● (II)
ipilimumab, nivolumab	✕	✕	✕	✕	● (II)
KN046	✕	✕	✕	✕	● (II)
osimertinib, osimertinib + chemotherapy	✕	✕	✕	✕	● (II)
radiation therapy, tyrosine kinase inhibitors	✕	✕	✕	✕	● (II)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.



## Relevant Therapy Summary (continued)

● In this cancer type    ○ In other cancer type    ● In this cancer type and other cancer types    ⛔ Contraindicated    ⚠ Both for use and contraindicated    ✕ No evidence

### EGFR p.(T790M) c.2369C>T (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
sunitinib	✕	✕	✕	✕	● (II)
alflutinib	✕	✕	✕	✕	● (I/II)
ASK120067	✕	✕	✕	✕	● (I/II)
AZD4635 + oleclumab	✕	✕	✕	✕	● (I/II)
CK-101	✕	✕	✕	✕	● (I/II)
DZD-9008	✕	✕	✕	✕	● (I/II)
EMB01	✕	✕	✕	✕	● (I/II)
icotinib hydrochloride + chemotherapy + radiation therapy	✕	✕	✕	✕	● (I/II)
KP-673	✕	✕	✕	✕	● (I/II)
lazertinib	✕	✕	✕	✕	● (I/II)
lerociclib, osimertinib	✕	✕	✕	✕	● (I/II)
TAK788	✕	✕	✕	✕	● (I/II)
U3-1402	✕	✕	✕	✕	● (I/II)
APG-1252, osimertinib	✕	✕	✕	✕	● (I)
cetuximab, FATE-NK100	✕	✕	✕	✕	● (I)
durvalumab + oleclumab, oleclumab	✕	✕	✕	✕	● (I)
ES-072	✕	✕	✕	✕	● (I)
everolimus + neratinib, neratinib + palbociclib, neratinib + trametinib	✕	✕	✕	✕	● (I)
FCN-411	✕	✕	✕	✕	● (I)
JNJ-61186372	✕	✕	✕	✕	● (I)
navitoclax, osimertinib	✕	✕	✕	✕	● (I)
nazartinib, trametinib	✕	✕	✕	✕	● (I)
necitumumab, osimertinib	✕	✕	✕	✕	● (I)
osimertinib + ramucirumab	✕	✕	✕	✕	● (I)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.





## Relevant Therapy Summary (continued)

● In this cancer type    ○ In other cancer type    ⓘ In this cancer type and other cancer types    ⚡ Contraindicated    ⚠ Both for use and contraindicated    ✕ No evidence

### EGFR p.(T790M) c.2369C>T (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
osimertinib, osimertinib + radiation therapy	✕	✕	✕	✕	● (I)
pirotinib	✕	✕	✕	✕	● (I)
SH-1028	✕	✕	✕	✕	● (I)
TP-0903	✕	✕	✕	✕	● (I)
TQB3456	✕	✕	✕	✕	● (I)
tyrosine kinase inhibitors, tyrosine kinase inhibitors + chemotherapy	✕	✕	✕	✕	● (I)
YK-029A	✕	✕	✕	✕	● (I)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Details

### Current FDA Information

● In this cancer type    ○ In other cancer type    ⓘ In this cancer type and other cancer types    ⚡ Contraindicated    🚫 Not recommended    🛡 Resistance

FDA information is current as of 2019-11-25. For the most up-to-date information, search [www.fda.gov](http://www.fda.gov).

### EGFR exon 19 deletion

#### ● osimertinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2018-08-28

Variant class: EGFR exon 19 deletion

#### Indications and usage:

TAGRISSO® is a kinase inhibitor indicated for

- the first-line treatment of patients with metastatic NSCLC whose tumors have epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test.
- the treatment of patients with metastatic EGFR T790M mutation-positive NSCLC, as detected by an FDA-approved test, whose disease has progressed on or after EGFR TKI therapy.

#### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/208065s011lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/208065s011lbl.pdf)



## EGFR p.(T790M) c.2369C>T

### ● osimertinib

**Cancer type:** Non-Small Cell Lung Cancer

**Label as of:** 2018-08-28

**Variant class:** EGFR T790M mutation

#### Indications and usage:

TAGRISSO® is a kinase inhibitor indicated for

- the first-line treatment of patients with metastatic NSCLC whose tumors have epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test.
- the treatment of patients with metastatic EGFR T790M mutation-positive NSCLC, as detected by an FDA-approved test, whose disease has progressed on or after EGFR TKI therapy.

#### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/208065s011lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/208065s011lbl.pdf)



## Current NCCN Information

- ☒ In this cancer type  
 ☐ In other cancer type  
 ☒ In this cancer type and other cancer types  
 ☒ Contraindicated  
 ☒ Not recommended  
 ☒ Resistance

NCCN information is current as of 2019-08-15. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org).  
 For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

### EGFR exon 19 deletion + EGFR p.(T790M) c.2369C>T

#### ● afatinib + cetuximab

Cancer type: Non-Small Cell Lung Cancer

Variant classes: EGFR T790M mutation & EGFRi sensitizing mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Non-Small Cell Lung Cancer; Progression after receiving erlotinib, afatinib, dacomitinib, or gefitinib and systemic therapy (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 7.2019]

### EGFR exon 19 deletion

#### ● osimertinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR exon 19 deletion

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Non-Small Cell Lung Cancer (NOS), Squamous Cell Carcinoma; Sensitizing EGFR mutation discovered prior to first-line systemic therapy (First-line therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 7.2019]

#### ● osimertinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR exon 19 deletion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Non-Small Cell Lung Cancer (NOS), Squamous Cell Carcinoma; Sensitizing EGFR mutation discovered during first-line systemic therapy; Interrupt or complete planned systemic therapy, including maintenance therapy (First-line therapy) (Preferred)
- Progression on osimertinib (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 7.2019]



## EGFR exon 19 deletion (continued)

### ● osimertinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR mutation

**NCCN Recommendation category:** 2A

**Population segment (Line of therapy):**

- Non-Small Cell Lung Cancer; Brain metastases; Newly diagnosed (Not specified)
- Non-Small Cell Lung Cancer; Leptomeningeal and Spine metastases (Not specified)

**Reference:** NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 1.2019]

### 🗨 alectinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFRi sensitizing mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Likewise, crizotinib, ceritinib, alectinib, brigatinib, or lorlatinib are not recommended for patients with sensitizing EGFR mutations who relapse on EGFR TKI therapy."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 7.2019]

### 🗨 brigatinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFRi sensitizing mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Likewise, crizotinib, ceritinib, alectinib, brigatinib, or lorlatinib are not recommended for patients with sensitizing EGFR mutations who relapse on EGFR TKI therapy."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 7.2019]

### 🗨 ceritinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFRi sensitizing mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Likewise, crizotinib, ceritinib, alectinib, brigatinib, or lorlatinib are not recommended for patients with sensitizing EGFR mutations who relapse on EGFR TKI therapy."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 7.2019]



## EGFR exon 19 deletion (continued)

### crizotinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFRi sensitizing mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Likewise, crizotinib, ceritinib, alectinib, brigatinib, or lorlatinib are not recommended for patients with sensitizing EGFR mutations who relapse on EGFR TKI therapy."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 7.2019]

### lorlatinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFRi sensitizing mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Likewise, crizotinib, ceritinib, alectinib, brigatinib, or lorlatinib are not recommended for patients with sensitizing EGFR mutations who relapse on EGFR TKI therapy."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 7.2019]

### pembrolizumab

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR mutation

**Other criteria:** CD274 overexpression

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "A small study suggests that single-agent pembrolizumab is not effective as first-line therapy in patients with metastatic NSCLC and EGFR mutations, even those with PD-L1 levels more than 50%."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 7.2019]



## EGFR p.(T790M) c.2369C>T

### ● osimertinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR T790M mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Non-Small Cell Lung Cancer (NOS), Squamous Cell Carcinoma; Progression on erlotinib, afatinib, gefitinib or dacomitinib (Subsequent therapy)
- Adenocarcinoma, Large Cell, Non-Small Cell Lung Cancer (NOS), Squamous Cell Carcinoma; Progression with symptomatic brain metastases (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 7.2019]

### ● osimertinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR T790M mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Non-Small Cell Lung Cancer; Brain metastases; Recurrent disease; Use agents active against primary tumor (Not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 1.2019]

### ● osimertinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Non-Small Cell Lung Cancer; Brain metastases; Newly diagnosed (Not specified)
- Non-Small Cell Lung Cancer; Leptomeningeal and Spine metastases (Not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 1.2019]

### 🗨️ pembrolizumab

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR mutation

Other criteria: CD274 overexpression

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "A small study suggests that single-agent pembrolizumab is not effective as first-line therapy in patients with metastatic NSCLC and EGFR mutations, even those with PD-L1 levels more than 50%."

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 7.2019]



## EGFR p.(T790M) c.2369C>T (continued)

### afatinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR T790M mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "The most common known mechanism is the acquisition of T790M (which is a secondary mutation in EGFR), which renders the kinase resistant to erlotinib, gefitinib, dacomitinib or afatinib."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 7.2019]

### dacomitinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR T790M mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "The most common known mechanism is the acquisition of T790M (which is a secondary mutation in EGFR), which renders the kinase resistant to erlotinib, gefitinib, dacomitinib or afatinib."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 7.2019]

### EGFR tyrosine kinase inhibitor

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR T790M mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "EGFR p.Thr790Met (T790M) is a mutation associated with acquired resistance to EGFR TKI therapy and has been reported in about 60% of patients with disease progression after initial response to erlotinib, gefitinib, or afatinib."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 7.2019]

### erlotinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR T790M mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "The most common known mechanism is the acquisition of T790M (which is a secondary mutation in EGFR), which renders the kinase resistant to erlotinib, gefitinib, dacomitinib or afatinib."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 7.2019]



## EGFR p.(T790M) c.2369C>T (continued)

### gefitinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR T790M mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "The most common known mechanism is the acquisition of T790M (which is a secondary mutation in EGFR), which renders the kinase resistant to erlotinib, gefitinib, dacomitinib or afatinib."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 7.2019]





## Current EMA Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ Contraindicated
 ☒ Not recommended
 ☒ Resistance

EMA information is current as of 2019-11-25. For the most up-to-date information, search [www.ema.europa.eu/ema](http://www.ema.europa.eu/ema).

### EGFR exon 19 deletion

#### ● bevacizumab (Allergan) + erlotinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2019-11-12

Variant class: EGFR exon 19 deletion

Reference:

[https://www.ema.europa.eu/en/documents/product-information/mvasi-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/mvasi-epar-product-information_en.pdf)

#### ● bevacizumab + erlotinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2019-11-12

Variant class: EGFR exon 19 deletion

Reference:

[https://www.ema.europa.eu/en/documents/product-information/avastin-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/avastin-epar-product-information_en.pdf)

#### ● osimertinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2019-10-23

Variant class: EGFR exon 19 deletion

Reference:

[https://www.ema.europa.eu/en/documents/product-information/tagrisso-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/tagrisso-epar-product-information_en.pdf)

### EGFR p.(T790M) c.2369C>T

#### ● osimertinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2019-10-23

Variant class: EGFR T790M mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/tagrisso-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/tagrisso-epar-product-information_en.pdf)

#### ⊘ gefitinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2019-05-28

Variant class: EGFR T790M mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/iressa-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/iressa-epar-product-information_en.pdf)



## Current ESMO Information

☒ In this cancer type  
 ☐ In other cancer type  
 ☒ In this cancer type and other cancer types  
 ☒ Contraindicated  
 ☒ Not recommended  
 ☒ Resistance

ESMO information is current as of 2019-08-15. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

### EGFR exon 19 deletion

#### ● osimertinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFRi sensitizing mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Stage IV Non-Small Cell Lung Cancer; ESMO-Magnitude of Clinical Benefit Scale Version 1.1 Score: 4 (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Ann Oncol (2018) 29 (suppl 4): iv192–iv237. (Corrigendum: 30 January 2019)]

#### ● gefitinib + carboplatin + pemetrexed

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Advanced Non-Small Cell Lung Cancer (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Ann Oncol (2018) 29 (suppl 4): iv192–iv237. (Corrigendum: 30 January 2019)]

#### ● bevacizumab + erlotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: II / B

Population segment (Line of therapy):

- Stage IV Non-Squamous Non-Small Cell Lung Cancer; ESMO-Magnitude of Clinical Benefit Scale Version 1.1 Score: 3 (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Ann Oncol (2018) 29 (suppl 4): iv192–iv237. (Corrigendum: 30 January 2019)]



## EGFR exon 19 deletion (continued)

### ● bevacizumab + gefitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: II / B

Population segment (Line of therapy):

- Stage IV Non-Squamous Non-Small Cell Lung Cancer; ESMO-Magnitude of Clinical Benefit Scale Version 1.1 Score: 3 (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Ann Oncol (2018) 29 (suppl 4): iv192–iv237. (Corrigendum: 30 January 2019)]

### ● atezolizumab + bevacizumab + carboplatin + paclitaxel

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Metastatic Non-Squamous Non-Small Cell Lung Cancer; Without contraindications to immunotherapy after targeted therapies have been exploited (Second-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Ann Oncol (2018) 29 (suppl 4): iv192–iv237. (Corrigendum: 30 January 2019)]

## EGFR p.(T790M) c.2369C>T

### ● osimertinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR T790M mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Stage IV Non-Squamous Non-Small Cell Lung Cancer; Resistance to first/second generation EGFR TKI and T790M positive; If not received previously; ESMO-Magnitude of Clinical Benefit Scale Version 1.1 Score: 4 (Second-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Ann Oncol (2018) 29 (suppl 4): iv192–iv237. (Corrigendum: 30 January 2019)]

## Signatures

Testing Personnel:

Laboratory Supervisor:



**Pathologist:**



## References

1. King et al. Amplification of a novel v-erbB-related gene in a human mammary carcinoma. *Science*. 1985 Sep 6;229(4717):974-6. PMID: 2992089
2. ErbB Receptors and Cancer. *Methods Mol. Biol.* 2017;1652:3-35. PMID: 28791631
3. Gutierrez et al. HER2: biology, detection, and clinical implications. *Arch. Pathol. Lab. Med.* 2011 Jan;135(1):55-62. PMID: 21204711
4. Pines et al. Oncogenic mutant forms of EGFR: lessons in signal transduction and targets for cancer therapy. *FEBS Lett.* 2010 Jun 18;584(12):2699-706. PMID: 20388509
5. Cancer Genome Atlas Research Network. Comprehensive molecular profiling of lung adenocarcinoma. *Nature*. 2014 Jul 31;511(7511):543-50. doi: 10.1038/nature13385. Epub 2014 Jul 9. PMID: 25079552
6. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
7. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
8. da et al. EGFR mutations and lung cancer. *Annu Rev Pathol.* 2011;6:49-69. doi: 10.1146/annurev-pathol-011110-130206. PMID: 20887192
9. Arcila et al. EGFR exon 20 insertion mutations in lung adenocarcinomas: prevalence, molecular heterogeneity, and clinicopathologic characteristics. *Mol. Cancer Ther.* 2013 Feb;12(2):220-9. PMID: 23371856
10. Kobayashi et al. EGFR Exon 18 Mutations in Lung Cancer: Molecular Predictors of Augmented Sensitivity to Afatinib or Neratinib as Compared with First- or Third-Generation TKIs. *Clin Cancer Res.* 2015 Dec 1;21(23):5305-13. doi: 10.1158/1078-0432.CCR-15-1046. Epub 2015 Jul 23. PMID: 26206867
11. Yasuda et al. Structural, biochemical, and clinical characterization of epidermal growth factor receptor (EGFR) exon 20 insertion mutations in lung cancer. *Sci Transl Med.* 2013 Dec 18;5(216):216ra177. PMID: 24353160
12. Chiu et al. Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitor Treatment Response in Advanced Lung Adenocarcinomas with G719X/L861Q/S768I Mutations. *J Thorac Oncol.* 2015 May;10(5):793-9. PMID: 25668120
13. Karachaliou et al. KRAS mutations in lung cancer. *Clin Lung Cancer.* 2013 May;14(3):205-14. PMID: 23122493
14. Brennan et al. The somatic genomic landscape of glioblastoma. *Cell.* 2013 Oct 10;155(2):462-77. PMID: 24120142
15. Cancer Genome Atlas Network. Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature*. 2015 Jan 29;517(7536):576-82. PMID: 25631445
16. Mitsudomi et al. Epidermal growth factor receptor in relation to tumor development: EGFR gene and cancer. *FEBS J.* 2010 Jan;277(2):301-8. PMID: 19922469
17. Ji et al. Epidermal growth factor receptor variant III mutations in lung tumorigenesis and sensitivity to tyrosine kinase inhibitors. *Proc. Natl. Acad. Sci. U.S.A.* 2006 May 16;103(20):7817-22. PMID: 16672372
18. Gazdar. Activating and resistance mutations of EGFR in non-small-cell lung cancer: role in clinical response to EGFR tyrosine kinase inhibitors. *Oncogene*. 2009 Aug;28 Suppl 1:S24-31. PMID: 19680293
19. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2016/021743s025lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2016/021743s025lbl.pdf)
20. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2019/201292s015lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/201292s015lbl.pdf)
21. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/206995s003lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/206995s003lbl.pdf)
22. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/208065s011lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/208065s011lbl.pdf)
23. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/211288s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/211288s000lbl.pdf)
24. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2019/125084s273lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/125084s273lbl.pdf)
25. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2017/125147s207lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/125147s207lbl.pdf)
26. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2015/125547s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2015/125547s000lbl.pdf)
27. NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 7.2019]