

ABOUT THE TEST FoundationOne®CDx is a next-generation sequencing (NGS) based assay that identifies genomic findings within hundreds of cancer-related genes.

PATIENT	DISEASE	Uterus carcinosarcoma	PHYSICIAN	ORDERING PHYSICIAN	Yeh, Yi-Chen	SPECIMEN	SPECIMEN SITE	Uterus
	NAME	Chang, Shu-Mei		MEDICAL FACILITY	Taipei Veterans General Hospital		SPECIMEN ID	S110-22362 A (PF21066)
	DATE OF BIRTH	08 August 1947		ADDITIONAL RECIPIENT	None		SPECIMEN TYPE	Slide Deck
	SEX	Female		MEDICAL FACILITY ID	205872		DATE OF COLLECTION	05 August 2021
	MEDICAL RECORD #	22542025		PATHOLOGIST	Not Provided		SPECIMEN RECEIVED	13 December 2021

Biomarker Findings

Microsatellite status - MS-Stable
Tumor Mutational Burden - 0 Muts/Mb

Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

PIK3CA H1047L
ARID1A T1649fs*49, Y762fs*1
PTEN R130G
CTCF E239fs*2
MLL2 Q3910_Q3911del

† See About the Test in appendix for details.

Report Highlights

- Targeted therapies with NCCN categories of evidence in this tumor type: Temsirolimus (p. 10)
- Evidence-matched clinical trial options based on this patient's genomic findings: (p. 11)
- Variants that may represent clonal hematopoiesis and may originate from non-tumor sources: **MLL2** Q3910_Q3911del (p. 8)

BIOMARKER FINDINGS

Microsatellite status - MS-Stable

Tumor Mutational Burden - 0 Muts/Mb

GENOMIC FINDINGS

PIK3CA - H1047L

10 Trials see p. 13

ARID1A - T1649fs*49, Y762fs*1

8 Trials see p. 11

PTEN - R130G

10 Trials see p. 15

THERAPY AND CLINICAL TRIAL IMPLICATIONS

No therapies or clinical trials. see Biomarker Findings section

No therapies or clinical trials. see Biomarker Findings section

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)	THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)
none	Temsirolimus 2A
	Everolimus
none	none
none	none

2A NCCN category

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS (CH)

Genomic findings below may include nontumor somatic alterations, such as CH. The efficacy of targeting such nontumor somatic alterations is unknown. This content should be interpreted based on clinical context. Refer to appendix for additional information on CH.

MLL2 - Q3910_Q3911del p. 8

GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

CTCF - E239fs*2 **p. 7** **MLL2 - Q3910_Q3911del** **p. 8**

NOTE Genomic alterations detected may be associated with activity of certain approved therapies; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and exhaustive. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies.

Therapies contained in this report may have been approved by the US FDA.

ORDERED TEST # ORD-1257210-01

BIOMARKER FINDINGS

BIOMARKER

Microsatellite status

RESULT

MS-Stable

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence, MSS tumors are significantly less likely than MSI-H tumors to respond to anti-PD-1 immune checkpoint inhibitors¹⁻³, including approved therapies nivolumab and pembrolizumab⁴. In a retrospective analysis of 361 patients with solid tumors treated with pembrolizumab, 3% were MSI-H and experienced a significantly higher ORR compared with non-MSI-H cases (70% vs. 12%, $p=0.001$)⁵.

FREQUENCY & PROGNOSIS

In uterine carcinosarcoma, MSI at any level has been reported in 9.5% (2/21) of cases⁶, and high MSI (MSI-H) has been observed in 5% (1/21) to 21% (6/28) of cases⁶⁻⁹. MSS has been reported in 73-89% of endometrial cancers¹⁰⁻¹⁷. A genomic profiling study of gynecologic carcinosarcoma (N=109) reported that MSI was predictive of improved PFS (HR=0.1937) and OS (HR=0.0937) in multivariate analysis⁹. Data regarding the role of MSI status on prognosis and survival in endometrial cancer are conflicting, with most studies finding no relationship between MSI-H endometrial cancers and survival^{12-13,15,18-20}, and one study predicting improved disease-free and disease-specific survival¹¹. However, these studies often evaluated endometrial cancers of all FIGO stages together. Studies specifically analyzing early stage endometrial cancer have reported a correlation between MSI-H and decreased survival^{12,16,21-22}, thereby suggesting that MSI-H

predicts for poor prognosis in this subset of endometrial tumors.

FINDING SUMMARY

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor²³. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2²³⁻²⁵. This sample is microsatellite-stable (MSS), equivalent to the clinical definition of an MSS tumor: one with mutations in none of the tested microsatellite markers²⁶⁻²⁸. MSS status indicates MMR proficiency and typically correlates with intact expression of all MMR family proteins^{23,25,27-28}.

ORDERED TEST # ORD-1257210-01

BIOMARKER FINDINGS

BIOMARKER

Tumor Mutational Burden

RESULT

0 Muts/Mb

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1²⁹⁻³¹, anti-PD-1 therapies²⁹⁻³², and combination nivolumab and ipilimumab³³⁻³⁸. In multiple pan-tumor studies, higher TMB has been reported to be associated with increased ORR and OS from treatment with immune checkpoint inhibitors^{29-32,39}. Higher TMB was found to be significantly associated with improved OS upon immune checkpoint inhibitor treatment for patients with 9 types of advanced tumors²⁹. Analyses across several solid tumor types reported that patients with higher TMB (defined as ≥ 16 -20 Muts/Mb) achieved greater clinical benefit from PD-1 or PD-L1-targeting monotherapy, compared with patients with higher TMB treated with chemotherapy⁴⁰ or those with lower TMB treated with PD-1 or PD-L1-targeting agents³⁰. However, the KEYNOTE 158 trial of pembrolizumab

monotherapy for patients with solid tumors found significant improvement in ORR for patients with TMB ≥ 10 Muts/Mb (based on this assay or others) compared to those with TMB < 10 Muts/Mb, in a large cohort that included multiple tumor types; similar findings were observed in the KEYNOTE 028 and 012 trials^{32,39}. Together, these studies suggest that patients with TMB ≥ 10 Muts/Mb may derive clinical benefit from PD-1 or PD-L1 inhibitors.

FREQUENCY & PROGNOSIS

In one study of 22 gynecologic carcinosarcomas, the average mutation burden was 1.4 mutations/Mb; the four tumors with mutation in either MLH1 or MSH6 had the highest mutation burden in this study ranging from 29-191 mutations/Mb⁴¹. In the TCGA Uterine Corpus Endometrioid Carcinoma dataset, 73% of cases were ultramutated, 28% were hypermutated, and 65% of samples were considered to have a low mutation rate¹⁰. Another study evaluating TMB in endometrial adenocarcinomas reported that 24% of tumors had a mutational burden of 10.4-54.1 mut/Mb, whereas 76% had 0-10.3 mut/Mb⁴². One study found TMB level was not significantly associated with OS for patients with uterine carcinosarcoma⁴³. Increased tumor mutational burden (TMB) in endometrial carcinoma has been correlated with POLE mutation and advanced high-grade endometrioid subtypes^{10,17,44-45}. Ultramutated endometrial tumors (elevated TMB

with POLE mutations) have also been associated with improved PFS¹⁰. The same study associated lower mutational burden, independent of PD-L1 status, in endometrial carcinomas with poorer prognosis¹⁰. For patients with advanced microsatellite-stable endometrial carcinoma not treated with immunotherapy, OS did not significantly differ between patients with TMB-high (≥ 10 Muts/Mb) and TMB-low (11.4 vs. 13.5 months, adjusted HR=1.15) in 1 study⁴⁶.

FINDING SUMMARY

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma⁴⁷⁻⁴⁸ and cigarette smoke in lung cancer⁴⁹⁻⁵⁰, treatment with temozolomide-based chemotherapy in glioma⁵¹⁻⁵², mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes^{10,53-56}, and microsatellite instability (MSI)^{10,55-56}. This sample harbors a TMB level associated with lower rates of clinical benefit from treatment with PD-1- or PD-L1-targeting immune checkpoint inhibitors compared with patients with tumors harboring higher TMB levels, based on several studies in multiple solid tumor types^{30-31,39}.

ORDERED TEST # ORD-1257210-01

GENOMIC FINDINGS

GENE

PIK3CA

ALTERATION

H1047L

TRANSCRIPT ID

NM_006218

CODING SEQUENCE EFFECT

3140A>T

VARIANT ALLELE FREQUENCY (% VAF)

48.4%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Clinical and preclinical data in various tumor types indicate that PIK3CA activating alterations may predict sensitivity to therapies targeting PI3K⁵⁷⁻⁵⁹, AKT⁶⁰⁻⁶¹, or mTOR⁶²⁻⁶⁹. In the Phase 2 MATCH trial for patients with PIK3CA-mutated

solid tumors, 28% (18/65) of patients experienced PFS lasting at least 6 months after treatment with taselisib; however, no ORs were observed in this study⁷⁰. A separate Phase 1b study of taselisib in combination with the CDK4/6 inhibitor palbociclib for patients with PIK3CA-mutated solid tumors reported an ORR of 0% (n=12) and a DCR of 17% (2/12)⁷¹. In a Phase 1 trial of the dual PI3K/mTOR kinase inhibitor apitolisib, 79% (11/14) of patients with PIK3CA-mutated advanced solid tumors experienced disease control (3 PRs, 8 SDs)⁷². The PI3K inhibitor alpelisib demonstrated an ORR of 6.0% (8/134) and a DCR of 58% (78/134) in a study for patients with PIK3CA-mutated solid tumors⁷³. However, the PI3K inhibitor copanlisib exhibited limited efficacy in PIK3CA-mutated tumors⁷⁴⁻⁷⁵.

FREQUENCY & PROGNOSIS

PI3K pathway alterations are frequent in gynecologic carcinosarcomas and have been

reported in greater than half of all cases in one study⁴¹. PIK3CA mutations have been reported in 19-41% of gynecologic carcinosarcomas^{41,76} and 8% of endometrial carcinosarcoma⁷⁷. Published data investigating the prognostic implications of PIK3CA alterations in uterine carcinosarcoma are limited (PubMed, Jul 2021).

FINDING SUMMARY

PIK3CA encodes p110-alpha, which is the catalytic subunit of phosphatidylinositol 3-kinase (PI3K). The PI3K pathway is involved in cell signaling that regulates a number of critical cellular functions, including cell growth, proliferation, differentiation, motility, and survival⁷⁸⁻⁷⁹. PIK3CA alterations that have been characterized as activating, such as observed here, are predicted to be oncogenic⁸⁰⁻¹⁰⁰.

ORDERED TEST # ORD-1257210-01

GENOMIC FINDINGS

GENE

ARID1A

ALTERATION

T1649fs*49, Y762fs*1

TRANSCRIPT ID

NM_006015, NM_006015

CODING SEQUENCE EFFECT

4945_4946insA, 2286delC

VARIANT ALLELE FREQUENCY (% VAF)

44.0%, 43.5%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no therapies approved to address the mutation or loss of ARID1A in cancer. However, on the basis of limited clinical and preclinical evidence, ARID1A inactivating mutations may lead to sensitivity to ATR inhibitors such as M6620 and ceralasertib¹⁰¹. In a Phase 2 study of ceralasertib in solid tumors, 2 patients with endometrial carcinoma in the cohort with loss of ARID1A expression achieved CRs on ceralasertib monotherapy; at least 1 of these 2 patients carried an inactivating ARID1A mutation. In contrast, no responses were observed for patients with normal ARID1A expression treated with ceralasertib combined with olaparib¹⁰². One patient with small cell lung cancer harboring an ARID1A mutation

experienced a PR when treated with M6620 combined with topotecan¹⁰³. In a Phase 1 trial, a patient with metastatic colorectal cancer harboring both an ARID1A mutation and ATM loss treated with single-agent M6620 achieved a CR that was ongoing at 29 months¹⁰⁴. On the basis of limited preclinical evidence from studies in ovarian cancer, ARID1A inactivation may predict sensitivity to EZH2 inhibitors¹⁰⁵⁻¹⁰⁶, which are under investigation in clinical trials. Other studies have reported that the loss of ARID1A may activate the PI3K-AKT pathway and be linked with sensitivity to inhibitors of this pathway¹⁰⁷⁻¹⁰⁹. Patients with ARID1A alterations in advanced or metastatic solid tumors may derive benefit from treatment with anti-PD-1 or anti-PD-L1 immunotherapy¹¹⁰. Loss of ARID1A expression has been associated with chemoresistance to platinum-based therapy for patients with ovarian clear cell carcinoma¹¹¹⁻¹¹² and to 5-fluorouracil in colorectal cancer cell lines¹¹³.

FREQUENCY & PROGNOSIS

ARID1A alterations are particularly prevalent in ovarian clear cell carcinoma (46-50%), ovarian and uterine endometrioid carcinomas (24-44%), and cholangiocarcinoma (27%); they are also reported in up to 27% of gastric carcinoma, esophageal adenocarcinoma, Waldenstrom macroglobulinemia, pediatric Burkitt lymphoma, hepatocellular carcinoma, colorectal carcinoma,

and urothelial carcinoma samples analyzed (COSMIC, cBioPortal, 2021)¹¹⁴⁻¹²². ARID1A loss is associated with microsatellite instability in ovarian and endometrial endometrioid adenocarcinomas^{45,110,123-125}, CRC^{110,126-128}, and gastric cancer^{110,129-133}. ARID1A protein loss is associated with tumors of poor histological grade for many tumor types, including colorectal cancer (CRC)¹²⁶⁻¹²⁸, cervical cancer¹³⁴⁻¹³⁵, gastric cancer¹²⁹⁻¹³³, urothelial carcinoma¹³⁶⁻¹³⁸, ovarian and endometrial cancers^{45,112,123-125,139-143}, breast carcinoma¹⁴⁴⁻¹⁴⁶, and clear cell renal cell carcinoma¹⁴⁷; ARID1A mutation has been associated with poor outcomes for patients with cholangiocarcinoma¹⁴⁸⁻¹⁵¹. However, prognostic data regarding patient survival are often mixed and conflicting.

FINDING SUMMARY

ARID1A encodes the AT-rich interactive domain-containing protein 1A, also known as Baf250a, a member of the SWI/SNF chromatin remodeling complex. Mutation, loss, or inactivation of ARID1A has been reported in many cancers, and the gene is considered a tumor suppressor^{118,132,145,152-157}. ARID1A mutations, which are mostly truncating, have been identified along the entire gene and often correlate with ARID1A protein loss^{118,130,153-154,158}, whereas ARID1A missense mutations are mostly uncharacterized.

ORDERED TEST # ORD-1257210-01

GENOMIC FINDINGS

GENE

PTEN

ALTERATION

R130G

TRANSCRIPT ID

NM_000314

CODING SEQUENCE EFFECT

388C>G

VARIANT ALLELE FREQUENCY (% VAF)

95.2%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

PTEN loss or mutation leads to activation of the PI3K-AKT-mTOR pathway and may predict sensitivity to inhibitors of this pathway^{75,159-161}. Across various tissues, most clinical studies have not observed an association between PTEN deficiency and response to inhibitors of the PI3K-AKT-mTOR pathway. However, limited studies in prostate cancer¹⁶²⁻¹⁶⁵, renal cell carcinoma¹⁶⁶, breast cancer¹⁶⁷⁻¹⁶⁸, and colorectal cancer¹⁶⁹ have reported an association between PTEN deficiency and response to inhibitors targeting the PI3K-AKT-mTOR pathway. Preclinical data indicate that PTEN loss or inactivation may predict sensitivity to PARP inhibitors¹⁷⁰⁻¹⁷⁴, and clinical benefit has been observed for patients with PTEN-altered breast cancer including triple negative breast cancer¹⁷⁵, ovarian cancer¹⁷⁶, uterine leiomyosarcoma¹⁷⁷, and endometrial cancer¹⁷⁴.

treated with PARP inhibitors. However, some studies have reported a lack of association between PTEN mutation and PARP inhibitor sensitivity¹⁷⁸⁻¹⁷⁹.

FREQUENCY & PROGNOSIS

PTEN alterations have been reported in 12.9% of endometrial carcinosarcomas and 6.7% (3/45) of ovarian carcinosarcoma cases analyzed in COSMIC (Apr 2021)¹¹⁴. PTEN mutations have been reported in up to 40% of uterine carcinosarcomas in the literature¹⁸⁰⁻¹⁸¹. A study of gynecological carcinosarcomas, including 17 uterine cases and 5 ovarian cases, identified alterations activating the PI3K pathway in over half of the samples analyzed, including PTEN mutations in 41% (9/22) of cases⁴¹. PTEN mutation has been associated with endometrioid-type but not serous-type uterine carcinosarcomas in multiple studies¹⁸²⁻¹⁸⁴. Loss of PTEN expression has been reported in 39% (12/31) of primary uterine carcinosarcomas, and specifically in 64% (20/37) of the epithelial component and 47% (17/33) of the mesenchymal component¹⁸⁵. In addition, loss of PTEN expression has also been found in 53% (10/17) of the epithelial component of metastatic tissue of uterine carcinosarcomas, but not in any of the five mesenchymal components studied¹⁸⁵. Published data investigating the prognostic implications of PTEN alteration in carcinosarcomas are limited (PubMed, Apr 2021).

functions as a tumor suppressor by negatively regulating the PI3K-AKT-mTOR pathway; loss of PTEN can lead to uncontrolled cell growth and suppression of apoptosis¹⁶⁰. Alterations such as seen here may disrupt PTEN function or expression¹⁸⁶⁻²²⁷.

POTENTIAL GERMLINE IMPLICATIONS

One or more of the PTEN variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters) associated with hamartoma tumor syndrome (ClinVar, Sep 2021)²²⁸. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. PTEN mutations underlie several inherited disorders, collectively termed PTEN hamartoma tumor syndrome (PHTS), which include Cowden syndrome (CS) and its variant Lhermitte-Duclos disease (LD), Bannayan-Riley-Ruvalcaba syndrome (BRRS), PTEN-related Proteus syndrome (PS), and Proteus-like syndrome²²⁹⁻²³⁰. The mutation rate for PTEN in these disorders ranges from 20 to 85% of patients^{229,231}. The estimated incidence of Cowden syndrome is 1/200,000, which may be an underestimate due to the high variability of this disorder²²⁹. Given the association between PTEN and these inherited syndromes, in the appropriate clinical context, germline testing for mutations affecting PTEN is recommended.

FINDING SUMMARY

PTEN encodes an inositol phosphatase that

GENE

CTCF

ALTERATION

E239fs*2

TRANSCRIPT ID

NM_006565

CODING SEQUENCE EFFECT

716_717delAG

VARIANT ALLELE FREQUENCY (% VAF)

48.1%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted therapies available to address genomic alterations in CTCF.

FREQUENCY & PROGNOSIS

Somatic mutations in CTCF are infrequently reported in most cancers, but have been observed more commonly (24%) in uterine corpus endometrial carcinoma (cBioPortal, 2021)¹¹⁵⁻¹¹⁶; nearly half of the observed mutations were truncating, suggesting a tumor suppressor role for CTCF in this disease. In addition, CTCF has been found to act as a tumor suppressor in breast cancer cell line studies²³²⁻²³³.

FINDING SUMMARY

CTCF encodes an 11-zinc-finger protein that is implicated in a number of regulatory roles, including gene activation and repression, imprinting, insulation, methylation, and X chromosome inactivation²³⁴. CTCF plays a role in transcriptional regulation of a number of key cancer-associated genes, including the oncogene MYC²³⁵ and tumor suppressor TP53²³⁶, via maintenance of local DNA methylation status. The decreased expression levels of CTCF and/or BORIS, another 11-zinc-finger transcriptional regulator, were reported to be closely associated with global DNA methylation variability and decreased overall survival in epithelial ovarian cancer²³⁷⁻²³⁸.

ORDERED TEST # ORD-1257210-01

GENOMIC FINDINGS

GENE

MLL2

ALTERATION

Q3910_Q3911del

TRANSCRIPT ID

NM_003482

CODING SEQUENCE EFFECT

11729_11734delAGCAAC

VARIANT ALLELE FREQUENCY (% VAF)

40.8%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted therapies available to address genomic alterations in MLL2.

FREQUENCY & PROGNOSIS

MLL2 alterations are observed in a number of solid tumor contexts (COSMIC, 2021)¹¹⁴, and are especially prevalent in lung squamous cell carcinoma (SCC)²³⁹ and small cell lung carcinoma (SCLC)²⁴⁰. MLL2 mutation was found to be an independent prognostic factor of poor PFS and OS in non-small cell lung cancer, but not in SCLC²⁴¹. One study reported that MLL2 truncating mutations were more common in recurrent ovary granulosa cell tumors (GCT) compared with primary GCTs (24% [10/42] vs. 3.0% [1/32])²⁴². In a study of esophageal SCC, high MLL2 expression positively correlated with tumor stage, differentiation, and size, and negatively correlated with OS²⁴³.

FINDING SUMMARY

MLL2 encodes an H3K4-specific histone methyltransferase that is involved in the transcriptional response to progesterone

signaling²⁴⁴. Germline de novo mutations of MLL2 are responsible for the majority of cases of Kabuki syndrome, a complex and phenotypically distinctive developmental disorder²⁴⁵. A significant number of inactivating MLL2 alterations have been observed in multiple tumor types, suggesting a tumor suppressor role²⁴⁶.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion²⁴⁷⁻²⁵². Comprehensive genomic profiling of solid tumors may detect nontumor alterations that are due to CH^{251,253-254}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

ORDERED TEST # ORD-1257210-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Everolimus

Assay findings association
PIK3CA
H1047L

AREAS OF THERAPEUTIC USE

Everolimus is an orally available mTOR inhibitor that is FDA approved to treat renal cell carcinoma (RCC) following antiangiogenic therapy; pancreatic neuroendocrine tumors; and well-differentiated non-functional neuroendocrine tumors of the lung or gastrointestinal tract. Everolimus is also approved to treat either renal angiomyolipoma or subependymal giant cell astrocytoma in association with tuberous sclerosis complex (TSC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical evidence⁶²⁻⁶⁹, PIK3CA activating mutations may predict sensitivity to mTOR inhibitors such as everolimus and temsirolimus. Studies have reported modest activity of these therapies as single agents (ORRs of 0-4%), but improved activity has been observed when they are combined with other agents such as bevacizumab and doxorubicin (ORRs of 25-44%), for patients with PIK3CA-mutated solid tumors^{66-69,255-259}.

SUPPORTING DATA

Of 2 patients with PIK3CA-mutated uterine malignant mixed Mullerian tumor (MMMT) who received everolimus monotherapy in the ProFiLER study, 1 achieved PR for 4.6 months and 1 experienced progressive disease⁶⁵. In a Phase 1 study of everolimus in combination with sorafenib, of the 22 enrolled patients with advanced solid tumors, the best response was SD lasting 168 days in a patient with uterine carcinosarcoma²⁶⁰. A patient with a mixed Mullerian tumor exhibited a PR in a Phase 1 trial of a rapamycin analog, deforolimus²⁶¹. A study of the mTOR inhibitor ridaforolimus as a single agent reported

no clinical response in 5 patients with uterine carcinosarcoma²⁶². In a Phase 2 clinical trial of recurrent endometrial cancer, 43% (12/28) of patients reported SD at 8 weeks and 21% (6/28) of patients achieved clinical benefit at 20 weeks upon administration of everolimus monotherapy²⁶³. Combination with the aromatase inhibitor letrozole for the same disease population achieved an ORR of 31% (11/35), with 9 CRs²⁶⁴. Further addition of metformin to this regimen led to a clinical benefit rate (CR+PR+SD) of 67% (32/48), including PR in 29% (14/48) of cases; no significant difference was observed between cases with and without KRAS mutation²⁶⁵. Everolimus achieved PR or SD in 35% of patients with recurrent endometrial carcinoma; KRAS mutation was associated with reduced median PFS (3.1 vs. 1.0 months) and median OS (9.3 vs. 2.3 months)²⁶⁶. Another study investigating estrogen and/or progesterone receptor-positive gynecologic or breast malignancies featuring mutation or loss of genes in the PI3K-AKT-mTOR pathway, including PIK3CA, AKT1, or PTEN, observed SD in 17% (1/6) of patients with endometrial cancer following combined treatment with everolimus and anastrozole²⁶⁷. No response was seen in a patient with endometrial stromal sarcoma and Peutz-Jeghers Syndrome associated with a germline STK11 mutation treated with a combination of everolimus and anastrozole²⁶⁸. Whereas frequent adverse events precluded a recommended Phase 2 dose and schedule for the combination of trametinib and everolimus in a Phase 1b trial for solid tumors²⁶⁹, a retrospective study for heavily pretreated patients with solid tumors reported tolerable regimens of the combination for 23/31 patients, with 16 patients treated >3 months and evaluable patients achieving a median PFS of 6.5 months²⁷⁰.

© 2021 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 27 December 2021
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Shakti Ramkissoon, M.D., Ph.D., M.M. Sc, Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1257210-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Temsirolimus

Assay findings association
PIK3CA
H1047L

AREAS OF THERAPEUTIC USE

Temsirolimus is an intravenous mTOR inhibitor that is FDA approved for the treatment of advanced renal cell carcinoma. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical evidence⁶²⁻⁶⁹, PIK3CA activating mutations may predict sensitivity to mTOR inhibitors such as everolimus and temsirolimus. Studies have reported modest activity of these therapies as single agents (ORRs of 0-4%), but improved activity has been observed when they are combined with other agents such as bevacizumab and doxorubicin (ORRs of 25-44%), for patients with PIK3CA-mutated solid tumors^{66-69,255-259}.

SUPPORTING DATA

A study of the mTOR inhibitor ridaforolimus as a single agent reported no clinical response in 5 patients with uterine carcinosarcoma²⁶². A study of the combination of temsirolimus and topotecan was investigated in endometrial cancers, including 3 patients with

carcinosarcoma, with 9/15 patients experiencing stable disease; however, this regimen was not well tolerated in patients who had previously received radiation therapy²⁷¹. A Phase 2 clinical trial of temsirolimus in recurrent or metastatic endometrial cancer reported PR in 4/29 (14%) chemotherapy-naïve patients and 4% (1/25) of chemotherapy-treated patients, with SD reported in 69% (20/29) of chemotherapy-naïve patients and 48% (12/25) of chemotherapy-treated patients; however, response in this study was found to be independent of molecular markers of PI3K-AKT-mTOR pathway activation²⁷². Another Phase 2 study of temsirolimus in patients with endometrial cancer reported PFS of >15 months in 6 patients and associated clinical benefit and longer PFS with mutation of AKT1 or CTNNB1, respectively²⁷³. Temsirolimus combined with carboplatin and paclitaxel achieved objective partial responses in 82% (9/11) of patients with endometrial cancer²⁷⁴. A Phase 2 trial of temsirolimus in combination with bevacizumab in patients with endometrial carcinoma reported clinical response in 25% of patients²⁷⁵.

NOTE Genomic alterations detected may be associated with activity of certain FDA approved drugs, however, the agents listed in this report may have varied evidence in the patient's tumor type.

ORDERED TEST # ORD-1257210-01

CLINICAL TRIALS

NOTE Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually updated and

should be investigated by the physician or research staff. This is not a comprehensive list of all available clinical trials. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials listed here may have additional enrollment criteria that may require

medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see [clinicaltrials.gov](https://www.foundationmedicine.com/genomic-testing#support-services). Or visit <https://www.foundationmedicine.com/genomic-testing#support-services>.

GENE
ARID1A
RATIONALE
 ARID1A loss or inactivation may predict

sensitivity to ATR inhibitors.

ALTERATION
 T1649fs*49, Y762fs*1

NCT02264678
PHASE 1/2

Ascending Doses of AZD6738 in Combination With Chemotherapy and/or Novel Anti Cancer Agents

TARGETS
 ATR, PARP, PD-L1

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Cambridge (United Kingdom), Withington (United Kingdom), London (United Kingdom), Sutton (United Kingdom), Villejuif (France), Saint Herblain (France), California

NCT02630199
PHASE 1

Study of AZD6738, DNA Damage Repair/Novel Anti-cancer Agent, in Combination With Paclitaxel, in Refractory Cancer

TARGETS
 ATR

LOCATIONS: Seoul (Korea, Republic of)

NCT04266912
PHASE 1/2

Avelumab and M6620 for the Treatment of DDR Deficient Metastatic or Unresectable Solid Tumors

TARGETS
 ATR, PD-L1

LOCATIONS: Texas

NCT04497116
PHASE 1/2

Study of RP-3500 in Advanced Solid Tumors

TARGETS
 ATR, PARP

LOCATIONS: Copenhagen (Denmark), Newcastle Upon Tyne (United Kingdom), Manchester (United Kingdom), London (United Kingdom), Toronto (Canada), Massachusetts, Rhode Island, New York, Tennessee, Texas

NCT04514497
PHASE 1

Testing the Addition of an Anti-cancer Drug, BAY 1895344, to Usual Chemotherapy for Advanced Stage Solid Tumors, With a Specific Focus on Patients With Small Cell Lung Cancer, Poorly Differentiated Neuroendocrine Cancer, and Pancreatic Cancer

TARGETS
 ATR, TOP1

LOCATIONS: Arizona, Oklahoma, Missouri, Connecticut, Tennessee

ORDERED TEST # ORD-1257210-01

CLINICAL TRIALS

NCT03641547

PHASE 1

M6620 Plus Standard Treatment in Oesophageal and Other Cancer

TARGETS
 ATR

LOCATIONS: Glasgow (United Kingdom), Manchester (United Kingdom), Oxford (United Kingdom), Cardiff (United Kingdom)

NCT03669601

PHASE 1

AZD6738 & Gemcitabine as Combination Therapy

TARGETS
 ATR

LOCATIONS: Cambridge (United Kingdom)

NCT02595931

PHASE 1

ATR Kinase Inhibitor VX-970 and Irinotecan Hydrochloride in Treating Patients With Solid Tumors That Are Metastatic or Cannot Be Removed by Surgery

TARGETS
 ATR

LOCATIONS: California, Missouri, Pennsylvania, Massachusetts, Connecticut, Tennessee, Florida

ORDERED TEST # ORD-1257210-01

CLINICAL TRIALS

GENE
PIK3CA
ALTERATION
 H1047L

RATIONALE
 PIK3CA activating mutations may lead to activation of the PI3K-AKT-mTOR pathway and may therefore indicate sensitivity to inhibitors of

this pathway. Strong clinical data support sensitivity of PIK3CA-mutated solid tumors to the PI3K-alpha inhibitor alpelisib.

NCT04589845
PHASE 2

Tumor-Agnostic Precision Immuno-Oncology and Somatic Targeting Rational for You (TAPISTRY) Platform Study

TARGETS
 ALK, ROS1, TRKA, TRKB, TRKC, RET, PD-L1, AKTs, ERBB2, MDM2, PI3K-alpha

LOCATIONS: Zhongzheng Dist. (Taiwan), Taipei City (Taiwan), Tainan (Taiwan), Seoul (Korea, Republic of), Beijing (China), Woolloongabba (Australia), Darlinghurst (Australia), Randwick (Australia), Melbourne (Australia), Haifa (Israel)

NCT03239015
PHASE 2

Efficacy and Safety of Targeted Precision Therapy in Refractory Tumor With Druggable Molecular Event

TARGETS
 EGFR, ERBB2, ERBB4, PARP, mTOR, MET, RET, ROS1, VEGFRs, BRAF, CDK4, CDK6

LOCATIONS: Shanghai (China)

NCT02688881
PHASE 4

Study to Evaluate the Safety and Efficacy of Sirolimus, in Subject With Refractory Solid Tumors

TARGETS
 mTOR

LOCATIONS: Seoul (Korea, Republic of)

NCT04803318
PHASE 2

Trametinib Combined With Everolimus and Lenvatinib for Recurrent/Refractory Advanced Solid Tumors

TARGETS
 mTOR, FGFRs, KIT, PDGFRA, RET, VEGFRs, MEK

LOCATIONS: Guangzhou (China)

NCT03772561
PHASE 1

Phase I Study of AZD5363 + Olaparib + Durvalumab in Patients With Advanced or Metastatic Solid Tumor Malignancies

TARGETS
 PARP, AKTs, PD-L1

LOCATIONS: Singapore (Singapore)

ORDERED TEST # ORD-1257210-01

CLINICAL TRIALS

NCT04801966
PHASE NULL

Safety and Oversight of the Individually Tailored Treatment Approach: A Novel Pilot Study

TARGETS
CDK4, CDK6, PI3K-alpha, PD-L1, MEK,
PARP, PD-1, BRAF

LOCATIONS: Melbourne (Australia)

NCT03994796
PHASE 2

Genetic Testing in Guiding Treatment for Patients With Brain Metastases

TARGETS
ALK, ROS1, TRKA, TRKB, TRKC, CDK4,
CDK6, PI3K, mTOR

LOCATIONS: Alaska, Washington

NCT04632992
PHASE 2

A Study Evaluating Targeted Therapies in Participants Who Have Advanced Solid Tumors With Genomic Alterations or Protein Expression Patterns Predictive of Response

TARGETS
ALK, ROS1, TRKA, TRKB, TRKC, PD-L1,
ERBB2, ERBB3, PI3K-alpha, RET, AKTs

LOCATIONS: Alaska, Washington, Oregon, California, Montana

NCT03297606
PHASE 2

Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR)

TARGETS
VEGFRs, ABL, SRC, ALK, AXL, MET,
ROS1, TRKA, TRKC, DDR2, KIT, EGFR,
PD-1, CTLA-4, PARP, CDK4, CDK6,
CSF1R, FLT3, RET, mTOR, ERBB2,
ERBB3, MEK, BRAF, SMO

LOCATIONS: Vancouver (Canada), Edmonton (Canada), Saskatoon (Canada), Regina (Canada), Ottawa (Canada), Montreal (Canada), Toronto (Canada), Kingston (Canada), London (Canada)

NCT03673787
PHASE 1/2

A Trial of Ipatasertib in Combination With Atezolizumab

TARGETS
AKTs, PD-L1

LOCATIONS: Sutton (United Kingdom)

ORDERED TEST # ORD-1257210-01

CLINICAL TRIALS
GENE
PTEN
ALTERATION
R130G

RATIONALE
PTEN loss or inactivating mutations may lead to increased activation of the PI3K-AKT-mTOR pathway and may indicate sensitivity to inhibitors

of this pathway. PTEN loss or inactivation may also predict sensitivity to PARP inhibitors.

NCT04269200
PHASE 3

Durvalumab With or Without Olaparib as Maintenance Therapy After First-Line Treatment of Advanced and Recurrent Endometrial Cancer

TARGETS
PD-L1, PARP

LOCATIONS: Nakagami-gun (Japan), Shanghai (China), Guangdong (China), Hong Kong (Hong Kong), HKG (Hong Kong), Changchun (China), Wuhan (China), Kurume-shi (Japan), Gyeongsangnam-do (Korea, Republic of), Suwon (Korea, Republic of)

NCT04716686
PHASE 2

Niraparib Monotherapy as Maintain and Recurrent Treatment of Endometrial Serous Carcinoma

TARGETS
PARP

LOCATIONS: Jinan (China)

NCT04001569
PHASE 1/2

AZD8186 and Paclitaxel in Advanced Gastric Cancer

TARGETS
PI3K-beta

LOCATIONS: Seongnam-si (Korea, Republic of)

NCT02264678
PHASE 1/2

Ascending Doses of AZD6738 in Combination With Chemotherapy and/or Novel Anti Cancer Agents

TARGETS
ATR, PARP, PD-L1

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Cambridge (United Kingdom), Withington (United Kingdom), London (United Kingdom), Sutton (United Kingdom), Villejuif (France), Saint Herblain (France), California

NCT03772561
PHASE 1

Phase I Study of AZD5363 + Olaparib + Durvalumab in Patients With Advanced or Metastatic Solid Tumor Malignancies

TARGETS
PARP, AKTs, PD-L1

LOCATIONS: Singapore (Singapore)

NCT04801966
PHASE NULL

Safety and Oversight of the Individually Tailored Treatment Approach: A Novel Pilot Study

TARGETS
CDK4, CDK6, PI3K-alpha, PD-L1, MEK, PARP, PD-1, BRAF

LOCATIONS: Melbourne (Australia)

ORDERED TEST # ORD-1257210-01

CLINICAL TRIALS

NCT03651206

PHASE 2/3

Recurrent Ovarian CarcinoSarcoma Anti-pd-1 Niraparib

TARGETS
 PARP, PD-1

LOCATIONS: Lille (France), Besançon (France), Paris (France), Villejuif (France), Lyon (France), Caen (France), Marseille (France), Montpellier (France), Angers (France), Poitiers (France)

NCT03994796

PHASE 2

Genetic Testing in Guiding Treatment for Patients With Brain Metastases

TARGETS
 ALK, ROS1, TRKA, TRKB, TRKC, CDK4, CDK6, PI3K, mTOR

LOCATIONS: Alaska, Washington

NCT04632992

PHASE 2

A Study Evaluating Targeted Therapies in Participants Who Have Advanced Solid Tumors With Genomic Alterations or Protein Expression Patterns Predictive of Response

TARGETS
 ALK, ROS1, TRKA, TRKB, TRKC, PD-L1, ERBB2, ERBB3, PI3K-alpha, RET, AKTs

LOCATIONS: Alaska, Washington, Oregon, California, Montana

NCT04497116

PHASE 1/2

Study of RP-3500 in Advanced Solid Tumors

TARGETS
 ATR, PARP

LOCATIONS: Copenhagen (Denmark), Newcastle Upon Tyne (United Kingdom), Manchester (United Kingdom), London (United Kingdom), Toronto (Canada), Massachusetts, Rhode Island, New York, Tennessee, Texas

ORDERED TEST # ORD-1257210-01

APPENDIX
Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

CREBBP

V1924M

KDMSA

M1489T

MEN1

A163T

MSH3

R1061G

NOTCH1

R955H

ORDERED TEST # ORD-1257210-01

APPENDIX

Genes Assayed in FoundationOne®CDx

FoundationOne CDx is designed to include genes known to be somatically altered in human solid tumors that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay interrogates 324 genes as well as introns of 36 genes involved in rearrangements. The assay will be updated periodically to reflect new knowledge about cancer biology.

DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS

ABL1	ACVR1B	AKT1	AKT2	AKT3	ALK	ALOX12B	AMER1 (FAM123B)	APC
AR	ARAF	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX	AURKA
AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2	BCL6
BCOR	BCORL1	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2
BTB	C11orf30 (EMSY)	C17orf39 (GID4)	CALR	CARD11	CASP8	CBFB	CBL	CCND1
CCND2	CCND3	CCNE1	CD22	CD274 (PD-L1)	CD70	CD79A	CD79B	CDC73
CDH1	CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B
CDKN2C	CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R
CTCF	CTNNA1	CTNNB1	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1
DDR2	DIS3	DNMT3A	DOT1L	EED	EGFR	EP300	EPHA3	EPHB1
EPHB4	ERBB2	ERBB3	ERBB4	ERCC4	ERG	ERRF1	ESR1	EZH2
FAM46C	FANCA	FANCC	FANCG	FANCL	FAS	FBXW7	FGF10	FGF12
FGF14	FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1	FGFR2	FGFR3
FGFR4	FH	FLCN	FLT1	FLT3	FOXO2	FUBP1	GABRA6	GATA3
GATA4	GATA6	GNAI1	GNAI3	GNAQ	GNAS	GRM3	GSK3B	H3F3A
HDAC1	HGF	HNF1A	HRAS	HSD3B1	ID3	IDH1	IDH2	IGF1R
IKBKE	IKZF1	INPP4B	IRF2	IRF4	IRS2	JAK1	JAK2	JAK3
JUN	KDMSA	KDMS5C	KDM6A	KDR	KEAP1	KEL	KIT	KLHL6
KMT2A (MLL)	KMT2D (MLL2)	KRAS	LTK	LYN	MAF	MAP2K1 (MEK1)	MAP2K2 (MEK2)	MAP2K4
MAP3K1	MAP3K13	MAPK1	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1
MERTK	MET	MITF	MKNK1	MLH1	MPL	MRE11A	MSH2	MSH3
MSH6	MST1R	MTAP	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN	MYD88
NBN	NF1	NF2	NFE2L2	NFKB1A	NKX2-1	NOTCH1	NOTCH2	NOTCH3
NPM1	NRAS	NSD3 (WHSC1L1)	NT5C2	NTRK1	NTRK2	NTRK3	P2RY8	PALB2
PARK2	PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)	PDGFRA
PDGFRB	PDK1	PIK3C2B	PIK3C2G	PIK3CA	PIK3CB	PIK3R1	PIM1	PMS2
POLD1	POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI	PTCH1
PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51	RAD51B	RAD51C
RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	REL	RET
RICTOR	RNF43	ROS1	RPTOR	SDHA	SDHB	SDHC	SDHD	SETD2
SF3B1	SGK1	SMAD2	SMAD4	SMARCA4	SMARCB1	SMO	SNCAIP	SOC3
SOX2	SOX9	SPEN	SPOP	SRC	STAG2	STAT3	STK11	SUFU
SYK	TBX3	TEK	TET2	TGFBR2	TIPARP	TNFAIP3	TNFRSF14	TP53
TSC1	TSC2	TYRO3	U2AF1	VEGFA	VHL	WHSC1	WT1	XPO1
XRCC2	ZNF217	ZNF703						

DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS

ALK	BCL2	BCR	BRAF	BRCA1	BRCA2	CD74	EGFR	ETV4
ETV5	ETV6	EWSR1	EZR	FGFR1	FGFR2	FGFR3	KIT	KMT2A (MLL)
MSH2	MYB	MYC	NOTCH2	NTRK1	NTRK2	NUTM1	PDGFRA	RAF1
RARA	RET	ROS1	RSP02	SDC4	SLC34A2	TERC*	TERT**	TPRPS2

*TERC is an NCRNA

**Promoter region of TERT is interrogated

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS

Loss of Heterozygosity (LOH) score

Microsatellite (MS) status

Tumor Mutational Burden (TMB)

ORDERED TEST # ORD-1257210-01

APPENDIX

About FoundationOne®CDx

FoundationOne CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Cipalstraat 3, 2440 Geel, Belgium.


ABOUT FOUNDATIONONE CDx

FoundationOne CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details:
www.rochefoundationmedicine.com/f1cdxtech.

INTENDED USE

FoundationOne®CDx (F1CDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI), tumor mutational burden (TMB), and for selected forms of ovarian cancer, loss of heterozygosity (LOH) score, using DNA isolated from formalin-fixed, paraffin-embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with therapies in accordance with approved therapeutic product labeling. Additionally, F1CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms.

TEST PRINCIPLES

FoundationOne CDx will be performed exclusively as a laboratory service using DNA extracted from formalin-fixed, paraffin-embedded (FFPE) tumor samples. The proposed assay will employ a single DNA extraction method from routine FFPE biopsy or surgical resection specimens, 50-1000 ng of which will undergo whole-genome shotgun library construction and hybridization-based capture of all coding exons from 309 cancer-related genes, one promoter region, one non-coding (ncRNA), and select intronic regions from 34 commonly rearranged genes, 21 of which also include the coding exons. The assay therefore includes detection of alterations in a total of 324 genes.

Using an Illumina® HiSeq platform, hybrid capture-selected libraries will be sequenced to high uniform depth (targeting >500X median coverage with >99% of exons at coverage >100X). Sequence data will be processed using a customized analysis pipeline designed to accurately detect all classes of genomic alterations, including base substitutions, indels, focal copy number amplifications, homozygous gene deletions, and selected genomic rearrangements (e.g., gene fusions). Additionally, genomic signatures including loss of heterozygosity (LOH), microsatellite instability (MSI) and tumor mutational burden (TMB) will be reported.

THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. The F1CDx report may be used as an aid to inform molecular eligibility for clinical trials. Note: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

Diagnostic Significance

FoundationOne CDx identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

Qualified Alteration Calls (Equivocal and Subclonal)

An alteration denoted as "amplification – equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne CDx for identifying a copy number amplification is four (4) for ERBB2 and six (6) for all other genes. Conversely, an alteration denoted as "loss – equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is one that the FoundationOne CDx analytical methodology has identified as being present in <10% of the assayed tumor DNA.

Ranking of Therapies and Clinical Trials
Ranking of Therapies in Summary Table

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

Ranking of Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) (www.nccn.org). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2021. All rights reserved. To view the most recent and complete version of the guidelines, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

Limitations

1. In the fractional-based MSI algorithm, a tumor specimen will be categorized as MSI-H, MSS, or MS-Equivocal according to the fraction of microsatellite loci determined to be altered or unstable (i.e., the fraction unstable loci score). In the F1CDx assay, MSI is evaluated based on a genome-wide analysis across >2000 microsatellite loci. For a given microsatellite locus, non-somatic alleles are discarded, and the microsatellite is categorized as unstable if remaining alleles differ from the reference genome. The final fraction unstable loci score is calculated as the number of unstable microsatellite loci divided by the number of evaluable microsatellite loci. The MSI-H and MSS cut-off thresholds were determined by analytical concordance to a PCR comparator assay using a pan-tumor FFPE tissue sample set. Patients with results categorized as "MS-

© 2021 Foundation Medicine, Inc. All rights reserved.

ORDERED TEST # ORD-1257210-01

APPENDIX

About FoundationOne®CDx

- Stable" with median exon coverage <300X, "MS-Equivocal," or "Cannot Be Determined" should receive confirmatory testing using a validated orthogonal (alternative) method.
2. TMB by F1CDx is determined by counting all synonymous and non-synonymous variants present at 5% allele frequency or greater (after filtering) and the total number is reported as mutations per megabase (mut/Mb) unit. Observed TMB is dependent on characteristics of the specific tumor focus tested for a patient (e.g., primary vs. metastatic, tumor content) and the testing platform used for the detection; therefore, observed TMB results may vary between different specimens for the same patient and between detection methodologies employed on the same sample. The TMB calculation may differ from TMB calculations used by other assays depending on variables such as the amount of genome interrogated, percentage of tumor, assay limit of detection (LoD), filtering of alterations included in the score, and the read depth and other bioinformatic test specifications. Refer to the SSED for a detailed description of these variables in FMI's TMB calculation https://www.accessdata.fda.gov/cdrh_docs/pdf17/P170019B.pdf. The clinical validity of TMB defined by this panel has been established for TMB as a qualitative output for a cut-off of 10 mutations per megabase but has not been established for TMB as a quantitative score.
 3. The LOH score is determined by analyzing SNPs spaced at 1Mb intervals across the genome on the FoundationOne CDx test and extrapolating an LOH profile, excluding arm- and chromosome-wide LOH segments. Detection of LOH has been verified only for ovarian cancer patients, and the LOH score result may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas. The LOH score will be reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine LOH. Performance of the LOH classification has not been established for samples below 35% tumor content. There may be potential interference of ethanol with LOH detection. The interfering effects of xylene, hemoglobin, and triglycerides on the LOH score have not been demonstrated.
 4. Alterations reported may include somatic (not inherited) or germline (inherited) alterations; however, the test does not distinguish between germline and somatic alterations. The test does not provide information about susceptibility.
 5. Biopsy may pose a risk to the patient when archival tissue is not available for use with the assay. The patient's physician should determine

whether the patient is a candidate for biopsy.

6. Reflex testing to an alternative FDA approved companion diagnostic should be performed for patients who have an *ERBB2* amplification result detected with copy number equal to 4 (baseline ploidy of tumor +2) for confirmatory testing. While this result is considered negative by FoundationOne®CDx (F1CDx), in a clinical concordance study with an FDA approved FISH test, 70% (7 out of 10 samples) were positive, and 30% (3 out of 10 samples) were negative by the FISH test with an average ratio of 2.3. The frequency of *ERBB2* copy number 4 in breast cancer is estimated to be approximately 2%. Multiple references listed in <https://www.mycancergenome.org/content/disease/breast-cancer/ERBB2/238/> report the frequency of *HER2* overexpression as 20% in breast cancer. Based on the F1CDx *HER2* CDx concordance study, approximately 10% of *HER2* amplified samples had copy number 4. Thus, total frequency is conservatively estimated to be approximately 2%.

REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic, nontargeted treatment, germline, or clonal hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

VARIANT ALLELE FREQUENCY

Variant Allele Frequency (VAF) represents the fraction of sequencing reads in which the variant is observed. This attribute is not taken into account for therapy inclusion, clinical trial matching, or interpretive content. Caution is recommended in interpreting VAF to indicate the potential germline or somatic origin of an alteration, recognizing that tumor fraction and tumor ploidy of samples may vary.

Precision of VAF for base substitutions and indels

BASE SUBSTITUTIONS	%CV*
Repeatability	5.11 - 10.40
Reproducibility	5.95 - 12.31
INDELS	%CV*
Repeatability	6.29 - 10.00
Reproducibility	7.33 - 11.71

*Interquartile Range = 1st Quartile to 3rd Quartile

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >10%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM*, *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *CHEK2*, *FH*, *FLCN*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *PALB2*, *PMS2*, *POLE*, *RAD51C*, *RAD51D*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *TSC2*, and *VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1*, *CBL*, *DNMT3A*, *IDH2*, *JAK2*, *KMT2D* (*MLL2*), *MPL*, *MYD88*, *SF3B1*, *TET2*, and *U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear

ORDERED TEST # ORD-1257210-01

APPENDIX

About FoundationOne®CDx

cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne CDx.

TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. FoundationOne CDx is performed using DNA derived from tumor, and as such germline events may not be reported.

SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
mut/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

MR Suite Version 5.2.0

The median exon coverage for this sample is 965x

ORDERED TEST # ORD-1257210-01

APPENDIX
References

1. Gatalica Z, et al. Cancer Epidemiol. Biomarkers Prev. (2014) PMID: 25392179
2. Kroemer G, et al. Oncoimmunology (2015) PMID: 26140250
3. Lal N, et al. Oncoimmunology (2015) PMID: 25949894
4. Le DT, et al. N. Engl. J. Med. (2015) PMID: 26028255
5. Ayers et al., 2016; ASCO-SITC Abstract P60
6. Amant F, et al. Int. J. Gynecol. Cancer () PMID: 11437928
7. Taylor NP, et al. Mod. Pathol. (2006) PMID: 16810312
8. Nilbert M, et al. Fam. Cancer (2009) PMID: 19130300
9. Gotoh O, et al. Nat Commun (2019) PMID: 31672974
10. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
11. Black D, et al. J. Clin. Oncol. (2006) PMID: 16549821
12. Mackay HJ, et al. Eur. J. Cancer (2010) PMID: 20304627
13. Kanopienė D, et al. Medicina (Kaunas) (2014) PMID: 25458958
14. Hampel H, et al. Cancer Res. (2006) PMID: 16885385
15. Steinbakk A, et al. Cell Oncol (Dordr) (2011) PMID: 21547578
16. Bilbao C, et al. Int. J. Radiat. Oncol. Biol. Phys. (2010) PMID: 20005452
17. Church DN, et al. Hum. Mol. Genet. (2013) PMID: 23528559
18. Zigelboim I, et al. J. Clin. Oncol. (2007) PMID: 17513808
19. Bilbao-Sieyro C, et al. Oncotarget (2014) PMID: 25026289
20. Arabi H, et al. Gynecol. Oncol. (2009) PMID: 19275958
21. Stelloo E, et al. Clin. Cancer Res. (2016) PMID: 27006490
22. Nout RA, et al. Gynecol. Oncol. (2012) PMID: 22609107
23. Kocarnik JM, et al. Gastroenterol Rep (Oxf) (2015) PMID: 26337942
24. You JF, et al. Br. J. Cancer (2010) PMID: 21081928
25. Bairwa NK, et al. Methods Mol. Biol. (2014) PMID: 24623249
26. Boland CR, et al. Cancer Res. (1998) PMID: 9823339
27. Pawlik TM, et al. Dis. Markers (2004) PMID: 15528785
28. Boland CR, et al. Gastroenterology (2010) PMID: 20420947
29. Samstein RM, et al. Nat. Genet. (2019) PMID: 30643254
30. Goodman AM, et al. Mol. Cancer Ther. (2017) PMID: 28835386
31. Goodman AM, et al. Cancer Immunol Res (2019) PMID: 31405947
32. Cristescu R, et al. Science (2018) PMID: 30309915
33. Ready N, et al. J. Clin. Oncol. (2019) PMID: 30785829
34. Hellmann MD, et al. N. Engl. J. Med. (2018) PMID: 29658845
35. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
36. Hellmann MD, et al. Cancer Cell (2018) PMID: 29731394
37. Rozeman EA, et al. Nat Med (2021) PMID: 33558721
38. Sharma P, et al. Cancer Cell (2020) PMID: 32916128
39. Marabelle A, et al. Lancet Oncol. (2020) PMID: 32919526
40. Legrand et al., 2018; ASCO Abstract 12000
41. Jones S, et al. Nat Commun (2014) PMID: 25233892
42. Santin et al., 2016; ASCO Abstract 5591
43. Wu et al., 2019; DOI: 10.21037/atm.2019.10.116
44. Mehnert JM, et al. J. Clin. Invest. (2016) PMID: 27159395
45. Hussein YR, et al. Mod. Pathol. (2015) PMID: 25394778
46. Shao C, et al. JAMA Netw Open (2020) PMID: 33119110
47. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
48. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
49. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
50. Rizvi NA, et al. Science (2015) PMID: 25765070
51. Johnson BE, et al. Science (2014) PMID: 24336570
52. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
53. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
54. Heitzer E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
55. Nature (2012) PMID: 22810696
56. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
57. Fritsch C, et al. Mol. Cancer Ther. (2014) PMID: 24608574
58. Juric D, et al. J. Clin. Oncol. (2018) PMID: 29401002
59. Gallant JN, et al. NPJ Precis Oncol (2019) PMID: 30793038
60. André F, et al. N. Engl. J. Med. (2019) PMID: 31091374
61. Smyth LM, et al. NPJ Breast Cancer (2021) PMID: 33863913
62. Park HS, et al. PLoS ONE (2016) PMID: 27105424
63. Lim SM, et al. Oncotarget (2016) PMID: 26859683
64. Hou MM, et al. Oncotarget (2014) PMID: 25426553
65. Varnier R, et al. Eur J Cancer (2019) PMID: 31351267
66. Janku F, et al. Cell Rep (2014) PMID: 24440717
67. Moroney J, et al. Clin. Cancer Res. (2012) PMID: 22927482
68. Basho RK, et al. JAMA Oncol (2017) PMID: 27893038
69. Moroney JW, et al. Clin. Cancer Res. (2011) PMID: 21890452
70. Krop et al., 2018; ASCO Abstract 101
71. Pascual J, et al. Cancer Discov (2021) PMID: 32958578
72. Dolly SO, et al. Clin. Cancer Res. (2016) PMID: 26787751
73. Aust Fam Physician (1986) PMID: 2941002
74. Santin AD, et al. Gynecol Oncol Rep (2020) PMID: 31934607
75. Patnaik A, et al. Ann. Oncol. (2016) PMID: 27672108
76. Growdon WB, et al. Gynecol. Oncol. (2011) PMID: 21168197
77. Biscuola M, et al. Hum. Pathol. (2013) PMID: 23199529
78. Samuels Y, et al. Cancer Cell (2005) PMID: 15950905
79. Nat. Rev. Cancer (2009) PMID: 19629070
80. Kang S, et al. Proc. Natl. Acad. Sci. U.S.A. (2005) PMID: 15647370
81. Ikenoue T, et al. Cancer Res. (2005) PMID: 15930273
82. Gymnopoulos M, et al. Proc. Natl. Acad. Sci. U.S.A. (2007) PMID: 17376864
83. Horn S, et al. Oncogene (2008) PMID: 18317450
84. Rudd ML, et al. Clin. Cancer Res. (2011) PMID: 21266528
85. Hon WC, et al. Oncogene (2012) PMID: 22120714
86. Burke JE, et al. Proc. Natl. Acad. Sci. U.S.A. (2012) PMID: 22949682
87. Wu H, et al. Proc. Natl. Acad. Sci. U.S.A. (2009) PMID: 19915146
88. Laurenti R, et al. Rev Saude Publica (1990) PMID: 2103068
89. Dan S, et al. Cancer Res. (2010) PMID: 20530683
90. Oda K, et al. Cancer Res. (2008) PMID: 18829572
91. Zhao L, et al. Oncogene (2008) PMID: 18794883
92. Lui VW, et al. Cancer Discov (2013) PMID: 23619167
93. Ross RL, et al. Oncogene (2013) PMID: 22430209
94. Rivière JB, et al. Nat. Genet. (2012) PMID: 22729224
95. Shibata T, et al. Cancer Lett. (2009) PMID: 19394761
96. Dogruluk T, et al. Cancer Res. (2015) PMID: 26627007
97. Croessmann S, et al. Clin. Cancer Res. (2018) PMID: 29284706
98. Ng PK, et al. Cancer Cell (2018) PMID: 29533785
99. Spangle JM, et al. (2020) PMID: 32929011
100. Chen L, et al. Nat Commun (2018) PMID: 29636477
101. Williamson CT, et al. Nat Commun (2016) PMID: 27958275
102. Aggarwal et al., 2021; ESMO Abstract 5120
103. Thomas A, et al. J. Clin. Oncol. (2018) PMID: 29252124
104. Yap TA, et al. J Clin Oncol (2020) PMID: 32568634
105. Bitler BG, et al. Nat. Med. (2015) PMID: 25686104
106. Kim KH, et al. Nat. Med. (2015) PMID: 26552009
107. Wiegand KC, et al. BMC Cancer (2014) PMID: 24559118
108. Huang HN, et al. Mod. Pathol. (2014) PMID: 24336158
109. Samartzis EP, et al. Oncotarget (2014) PMID: 24979463
110. Okamura R, et al. J Immunother Cancer (2020) PMID: 32111729
111. Yokoyama Y, et al. J Gynecol Oncol (2014) PMID: 24459582
112. Katagiri A, et al. Mod. Pathol. (2012) PMID: 22101352
113. Xie C, et al. Tumour Biol. (2014) PMID: 24833095
114. Tate JG, et al. Nucleic Acids Res. (2019) PMID: 30371878
115. Cerami E, et al. Cancer Discov (2012) PMID: 22588877
116. Gao J, et al. Sci Signal (2013) PMID: 23550210
117. Wu RC, et al. Cancer Biol. Ther. (2014) PMID: 24618703
118. Jones S, et al. Hum. Mutat. (2012) PMID: 22009941
119. Dulak AM, et al. Nat. Genet. (2013) PMID: 23525077
120. Streppe MM, et al. Oncogene (2014) PMID: 23318448
121. Jiao Y, et al. J. Pathol. (2014) PMID: 24293293
122. Ross JS, et al. Oncologist (2014) PMID: 24563076
123. Huang HN, et al. Histopathology (2015) PMID: 25195947
124. Bosse T, et al. Mod. Pathol. (2013) PMID: 23702729
125. Allo G, et al. Mod. Pathol. (2014) PMID: 23887303
126. Chou A, et al. Hum. Pathol. (2014) PMID: 24925223
127. Ye J, et al. Hum. Pathol. (2014) PMID: 25311944
128. Wei XL, et al. World J. Gastroenterol. (2014) PMID: 25561809
129. Chen K, et al. Proc. Natl. Acad. Sci. U.S.A. (2015) PMID: 25583476
130. Wang K, et al. Nat. Genet. (2011) PMID: 22037554
131. Abe H, et al. Virchows Arch. (2012) PMID: 22915242
132. Wang DD, et al. PLoS ONE (2012) PMID: 22808142
133. Wiegand KC, et al. Hum. Pathol. (2014) PMID: 24767857
134. Katagiri A, et al. Int. J. Gynecol. Cancer (2012) PMID: 22274316
135. Cho H, et al. Hum. Pathol. (2013) PMID: 23427874
136. Gui Y, et al. Nat. Genet. (2011) PMID: 21822268
137. Balbás-Martínez C, et al. PLoS ONE (2013) PMID: 23650517
138. Faraj SF, et al. Hum. Pathol. (2014) PMID: 25175170
139. Rahman M, et al. Hum. Pathol. (2013) PMID: 22939958
140. Maeda D, et al. Int J Mol Sci (2010) PMID: 21614196
141. Lowery WJ, et al. Int. J. Gynecol. Cancer (2012) PMID: 22193641
142. Fadare O, et al. Mod. Pathol. (2013) PMID: 23524907
143. Mao TL, et al. Am. J. Surg. Pathol. (2013) PMID: 24076775
144. Zhang X, et al. Cancer Epidemiol (2012) PMID: 21889920
145. Mamo A, et al. Oncogene (2012) PMID: 21892209
146. Zhao J, et al. Tumour Biol. (2014) PMID: 24430365
147. Lichner Z, et al. Am. J. Pathol. (2013) PMID: 23416164
148. Feng F, et al. Int J Clin Oncol (2021) PMID: 33387086
149. Conci S, et al. Updates Surg (2020) PMID: 32020551
150. Simbolo M, et al. Sci Rep (2018) PMID: 29740198
151. Ruzzenente A, et al. Ann. Surg. Oncol. (2016) PMID: 26717940
152. Guan B, et al. Cancer Res. (2011) PMID: 21900401
153. Wiegand KC, et al. N. Engl. J. Med. (2010) PMID: 20942669

© 2021 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 27 December 2021
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Shakti Ramkissoon, M.D., Ph.D., M.M. Sc, Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1257210-01

APPENDIX **References**

154. Jones S, et al. Science (2010) PMID: 20826764
155. Yan HB, et al. Carcinogenesis (2014) PMID: 24293408
156. Huang J, et al. Nat. Genet. (2012) PMID: 22922871
157. Chan-On W, et al. Nat. Genet. (2013) PMID: 24185513
158. Zang ZJ, et al. Nat. Genet. (2012) PMID: 22484628
159. Courtney KD, et al. J. Clin. Oncol. (2010) PMID: 20085938
160. Simpson L, et al. Exp. Cell Res. (2001) PMID: 11237521
161. Milella M, et al. Sci Rep (2017) PMID: 28220839
162. Templeton AJ, et al. Eur. Urol. (2013) PMID: 23582881
163. Sweeney C, et al. Lancet (2021) PMID: 34246347
164. de Bono JS, et al. Clin. Cancer Res. (2019) PMID: 30037818
165. Saura C, et al. Cancer Discov (2017) PMID: 27872130
166. Voss MH, et al. Clin. Cancer Res. (2018) PMID: 30327302
167. André F, et al. J. Clin. Oncol. (2016) PMID: 27091708
168. Schmid P, et al. J. Clin. Oncol. (2019) PMID: 31841354
169. Weldon Gilcrease G, et al. Invest New Drugs (2019) PMID: 30302599
170. Mendes-Pereira AM, et al. EMBO Mol Med (2009) PMID: 20049735
171. Shen Y, et al. Clin. Cancer Res. (2013) PMID: 23881923
172. Chatterjee P, et al. PLoS ONE (2013) PMID: 23565244
173. McCormick A, et al. Int. J. Gynecol. Cancer (2016) PMID: 26905328
174. Forster MD, et al. Nat Rev Clin Oncol (2011) PMID: 21468130
175. Eikesdal HP, et al. Ann Oncol (2021) PMID: 33242536
176. Dougherty et al., 2014; ASCO Abstract 5536
177. Pan M, et al. Perm J (2021) PMID: 33970096
178. Sandhu SK, et al. Lancet Oncol. (2013) PMID: 23810788
179. Romero I, et al. Gynecol Oncol (2020) PMID: 32988624
180. Chorniack AD, et al. Cancer Cell (2017) PMID: 28292439
181. Lu X, et al. Cancer Biol. Ther. (2019) PMID: 30359167
182. Amant F, et al. Gynecol. Oncol. (2002) PMID: 11925138
183. McConechy MK, et al. J. Pathol. (2012) PMID: 22653804
184. Zhao S, et al. Proc. Natl. Acad. Sci. U.S.A. (2016) PMID: 27791010
185. de Jong RA, et al. Mod. Pathol. (2011) PMID: 21572397
186. Campbell RB, et al. J. Biol. Chem. (2003) PMID: 12857747
187. Rodríguez-Escudero I, et al. Hum. Mol. Genet. (2011) PMID: 21828076
188. He X, et al. Cancer Res. (2013) PMID: 23475934
189. Han SY, et al. Cancer Res. (2000) PMID: 10866302
190. Myers MP, et al. Proc. Natl. Acad. Sci. U.S.A. (1998) PMID: 9811831
191. Pradella LM, et al. BMC Cancer (2014) PMID: 24498881
192. Kim JS, et al. Mol. Cell. Biol. (2011) PMID: 21536651
193. Denning G, et al. Oncogene (2007) PMID: 17213812
194. Hlobilkova A, et al. Anticancer Res. (2016) PMID: 16619501
195. Redfern RE, et al. Protein Sci. (2010) PMID: 20718038
196. Shenoy S, et al. PLoS ONE (2012) PMID: 22505997
197. Wang Y, et al. Proc. Natl. Acad. Sci. U.S.A. (2009) PMID: 19329485
198. Okumura K, et al. J. Biol. Chem. (2006) PMID: 16829519
199. Lee JO, et al. Cell (1999) PMID: 10555148
200. Maxwell GL, et al. Cancer Res. (1998) PMID: 9635567
201. Risinger JJ, et al. Clin. Cancer Res. (1998) PMID: 9865913
202. Kato H, et al. Clin. Cancer Res. (2000) PMID: 11051241
203. Fenton TR, et al. Proc. Natl. Acad. Sci. U.S.A. (2012) PMID: 22891331
204. Ngeow J, et al. J. Clin. Endocrinol. Metab. (2012) PMID: 23066114
205. Lobo GP, et al. Hum. Mol. Genet. (2009) PMID: 19457929
206. Liu J, et al. Oncogene (2014) PMID: 23995781
207. Maehama T, et al. Annu. Rev. Biochem. (2001) PMID: 11395408
208. De Vivo I, et al. J. Med. Genet. (2000) PMID: 10807691
209. Ramaswamy S, et al. Proc. Natl. Acad. Sci. U.S.A. (1999) PMID: 10051603
210. Liu JL, et al. Mol. Cell. Biol. (2005) PMID: 15988030
211. Karoui M, et al. Br. J. Cancer (2004) PMID: 15026806
212. Gil A, et al. PLoS ONE (2015) PMID: 25875300
213. Furnari FB, et al. Cancer Res. (1998) PMID: 9823298
214. Spinelli L, et al. J. Med. Genet. (2015) PMID: 25527629
215. Mingo J, et al. Eur. J. Hum. Genet. (2018) PMID: 29706633
216. Wang Q, et al. J. Mol. Graph. Model. (2010) PMID: 20538496
217. Andrés-Pons A, et al. Cancer Res. (2007) PMID: 17942903
218. Butler MG, et al. J. Med. Genet. (2005) PMID: 15805158
219. Georgescu MM, et al. Proc. Natl. Acad. Sci. U.S.A. (1999) PMID: 10468583
220. Staal FJ, et al. Br. J. Cancer (2002) PMID: 12085208
221. Nguyen HN, et al. Oncogene (2014) PMID: 24292679
222. Rahdar M, et al. Proc. Natl. Acad. Sci. U.S.A. (2009) PMID: 19114656
223. Das S, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) PMID: 12808147
224. Wang X, et al. Biochem. J. (2008) PMID: 18498243
225. Valiente M, et al. J. Biol. Chem. (2005) PMID: 15951562
226. Nguyen HN, et al. Oncogene (2015) PMID: 25263454
227. Shan L, et al. Cell Discov (2020) PMID: 32704382
228. Landrum MJ, et al. Nucleic Acids Res. (2018) PMID: 29165669
229. Blumenthal GM, et al. Eur. J. Hum. Genet. (2008) PMID: 18781191
230. Orloff MS, et al. Oncogene (2008) PMID: 18794875
231. Zbuk KM, et al. Nat. Rev. Cancer (2007) PMID: 17167516
232. Méndez-Catalá CF, et al. Neoplasia (2013) PMID: 23908591
233. Tiffen JC, et al. Int. J. Cancer (2013) PMID: 23553099
234. Phillips JE, et al. Cell (2009) PMID: 19563753
235. Gombert WM, et al. PLoS ONE (2009) PMID: 19568426
236. Soto-Reyes E, et al. Oncogene (2010) PMID: 20101205
237. Woloszynska-Read A, et al. Clin. Cancer Res. (2011) PMID: 21296871
238. Kemp CJ, et al. Cell Rep (2014) PMID: 24794443
239. Nature (2012) PMID: 22960745
240. Augert A, et al. J Thorac Oncol (2017) PMID: 28007623
241. Ardeshtir-Larijani F, et al. Clin Lung Cancer (2018) PMID: 29627316
242. Hillman RT, et al. Nat Commun (2018) PMID: 29950560
243. Abudurehman A, et al. J. Cancer Res. Clin. Oncol. (2018) PMID: 29532228
244. Vicent GP, et al. Genes Dev. (2011) PMID: 21447625
245. Hannibal MC, et al. Am. J. Med. Genet. A (2011) PMID: 21671394
246. Fagan RJ, et al. Cancer Lett. (2019) PMID: 31128216
247. Jaiswal S, et al. N. Engl. J. Med. (2014) PMID: 25426837
248. Genovese G, et al. N. Engl. J. Med. (2014) PMID: 25426838
249. Xie M, et al. Nat. Med. (2014) PMID: 25326804
250. Acuna-Hidalgo R, et al. Am. J. Hum. Genet. (2017) PMID: 28669404
251. Severson EA, et al. Blood (2018) PMID: 29678827
252. Fuster JJ, et al. Circ. Res. (2018) PMID: 29420212
253. Chabon JJ, et al. Nature (2020) PMID: 32269342
254. Razavi P, et al. Nat. Med. (2019) PMID: 31768066
255. Janku F, et al. Cancer Res. (2013) PMID: 23066039
256. Janku F, et al. J. Clin. Oncol. (2012) PMID: 22271473
257. Janku F, et al. Mol. Cancer Ther. (2011) PMID: 21216929
258. Moulder S, et al. Ann. Oncol. (2015) PMID: 25878190
259. Byeon et al., 2020; doi: 10.21037/tcr.2020.04.07
260. Ma et al., 2012; ESMO Congress Abstract 447PD
261. Mita MM, et al. J. Clin. Oncol. (2008) PMID: 18202410
262. Colombo N, et al. Br. J. Cancer (2013) PMID: 23403817
263. Slomovitz BM, et al. Cancer (2010) PMID: 20681032
264. Slomovitz BM, et al. J. Clin. Oncol. (2015) PMID: 25624430
265. Soliman et al., 2016; ASCO Abstract 5506
266. Trédan O, et al. Target Oncol (2013) PMID: 23238879
267. Wheler JJ, et al. Oncotarget (2014) PMID: 24912489
268. Noriega-Iriondo MF, et al. Hered Cancer Clin Pract (2015) PMID: 25649062
269. Tolcher AW, et al. Ann. Oncol. (2015) PMID: 25344362
270. Patterson et al., 2018; AACR Abstract 3891
271. Temkin SM, et al. Gynecol. Oncol. (2010) PMID: 20347480
272. Oza AM, et al. J. Clin. Oncol. (2011) PMID: 21788564
273. Myers et al., 2015; ASCO Annual Meeting Abstract 5592
274. Kollmannsberger C, et al. Ann. Oncol. (2012) PMID: 21447615
275. Alvarez EA, et al. Gynecol. Oncol. (2013) PMID: 23262204

© 2021 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 27 December 2021
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Shakti Ramkissoon, M.D., Ph.D., M.M. Sc, Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531