



Sample Information

Patient Name: 王香谷**Gender:** Female**ID No.:** M220161721**History No.:** 9830722**Age:** 57**Ordering Doctor:** DOC6297D 陳庭安**Ordering REQ.:** 0BXVLKP**Signing in Date:** 2022/07/28**Path No.:** S111-97808**MP No.:** F22074**Assay:** Oncomine Focus Assay**Sample Type:** FFPE**Block No.:** S111-27173A+B**Percentage of tumor cells:** 30%**Reporting Doctor:** DOC5466K 葉奕成 (Phone: 8#5466)**Note:**

Sample Cancer Type: Non-Small Cell Lung Cancer

Table of Contents	Page
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2
Biomarker Descriptions	2
Relevant Therapy Summary	3
Relevant Therapy Details	3
Clinical Trials Summary	10
Alert Details	10

Report Highlights

1 Relevant Biomarkers
5 Therapies Available
2 Clinical Trials

Relevant Non-Small Cell Lung Cancer Variants

Gene	Finding	Gene	Finding
ALK	None detected	NTRK1	None detected
BRAF	None detected	NTRK2	None detected
EGFR	None detected	NTRK3	None detected
ERBB2	None detected	RET	None detected
KRAS	None detected	ROS1	SLC34A2-ROS1 fusion
MET	None detected		

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	SLC34A2-ROS1 fusion solute carrier family 34 member 2 - ROS proto-oncogene 1, receptor tyrosine kinase	crizotinib ^{1, 2} entrectinib ^{1, 2} ceritinib lorlatinib repotrectinib	None	2

Public data sources included in relevant therapies: FDA¹, NCCN, EMA², ESMO

Tier Reference: Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

Gene Fusions (RNA)

Genes	Variant ID	Locus	Read Count
SLC34A2-ROS1	SLC34A2-ROS1.S13R32.COSF1259	chr4:25678324 - chr6:117650609	75384
SLC34A2-ROS1	SLC34A2-ROS1.S13R33	chr4:25678324 - chr6:117647577	253
SLC34A2-ROS1	SLC34A2-ROS1.S13R34.COSF1261	chr4:25678324 - chr6:117645578	20792
SLC34A2-ROS1	SLC34A2-ROS1.S13R35.Non-Targeted	chr4:25678324 - chr6:117642557	600

Biomarker Descriptions

ROS1 (ROS proto-oncogene 1, receptor tyrosine kinase)

Background: The ROS1 gene encodes the ROS proto-oncogene receptor tyrosine kinase 1 which exhibits structural similarity to anaplastic lymphoma kinase (ALK)^{1,2}. Like ALK, ROS1 is the target of recurrent chromosomal rearrangements that generate fusion proteins containing the intact ROS1 tyrosine kinase domain combined with numerous fusion partner genes³. ROS1 fusion kinases are constitutively activated and drive oncogenic transformation⁴.

Alterations and prevalence: ROS1 fusions occur in approximately 1-2% of patients with non-small cell lung cancer (NSCLC) and are also observed in cholangiocarcinoma, gastric cancer, ovarian cancer, and glioblastoma^{1,5,6,7,8,9}.

Potential relevance: The tyrosine kinase inhibitor, entrectinib¹⁰, is approved (2019) for the treatment of ROS1 fusion positive metastatic NSCLC. Crizotinib¹¹, originally approved for the treatment of ALK positive NSCLC (2011), is also approved (2016) for the treatment of ROS1 positive NSCLC¹². Acquired resistance to crizotinib in ROS1 positive NSCLC is associated with kinase domain mutations S1986F/Y, G2032R, D2033N, and L2155S^{13,14,15}. The ROS1 tyrosine kinase inhibitor, repotrectinib¹⁶, was granted fast track and breakthrough designations (2020) for ROS1 positive NSCLC. Ceritinib is a second generation ALK inhibitor approved (2017) for ALK positive NSCLC that has also shown efficacy in ROS1 positive NSCLC. In a phase II study, ceritinib demonstrated systemic and intra-cranial activity with an objective response rate (ORR) of 62% in patients with advanced ROS1 positive NSCLC¹⁷. In addition to crizotinib and entrectinib, ceritinib is recommended for first-line treatment of ROS1-positive NSCLC¹⁸. Lorlatinib is a CNS-penetrant third-generation ALK and ROS1 inhibitor with preclinical activity against almost all known ALK and ROS1 resistance mutations^{19,20}. Lorlatinib is currently FDA approved (2018) for ALK positive metastatic NSCLC. In a phase I study testing lorlatinib in advanced ROS1-positive NSCLC, objective response was observed in 6/12 (50%) of patients²¹. Lorlatinib is recommended for subsequent therapy in ROS1 fusion-positive NSCLC in patients who have progressed after treatment with crizotinib, entrectinib, or ceritinib¹⁸.

Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

SLC34A2-ROS1 fusion

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
crizotinib	●	●	●	●	×
entrectinib	●	●	●	●	×
ceritinib	×	●	×	●	×
lorlatinib	×	●	×	●	×
repotrectinib	×	×	×	●	● (I/II)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Details

Current FDA Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

FDA information is current as of 2022-06-15. For the most up-to-date information, search www.fda.gov.

SLC34A2-ROS1 fusion

● crizotinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2021-09-22

Variant class: ROS1 fusion

Indications and usage:

XALKORI® is a kinase inhibitor indicated for the treatment of

- patients with metastatic non-small cell lung cancer (NSCLC) whose tumors are anaplastic lymphoma kinase (ALK) or ROS1-positive as detected by an FDA-approved test.
- pediatric patients 1 year of age and older and young adults with relapsed or refractory, systemic anaplastic large cell lymphoma (ALCL) that is ALK-positive.
 - Limitations of Use: The safety and efficacy of XALKORI® have not been established in older adults with relapsed or refractory, systemic ALK-positive ALCL.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/202570s031lbl.pdf

SLC34A2-ROS1 fusion (continued)

● entrectinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2021-11-05

Variant class: ROS1 fusion

Indications and usage:

ROZLYTREK® is a kinase inhibitor indicated for the treatment of:

- Adult patients with metastatic non-small cell lung cancer (NSCLC) whose tumors are ROS1-positive.
- Adult and pediatric patients 12 years of age and older with solid tumors that:
 - have a neurotrophic tyrosine receptor kinase (NTRK) gene fusion without a known acquired resistance mutation,
 - are metastatic or where surgical resection is likely to result in severe morbidity, and
 - have progressed following treatment or have no satisfactory alternative therapy

This indication is approved under accelerated approval based on tumor response rate and durability of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in the confirmatory trials.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/212725s005lbl.pdf

Current NCCN Information

- ☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

NCCN information is current as of 2022-06-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

SLC34A2-ROS1 fusion

● ceritinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ROS1 fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Other recommended intervention
- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

● crizotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ROS1 fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Preferred intervention
- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (Subsequent therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

● entrectinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ROS1 fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Preferred intervention
- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (Subsequent therapy); Preferred intervention
- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Progression, Brain Metastases (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

SLC34A2-ROS1 fusion (continued)

● lorlatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ROS1 fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic, Progression (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

● crizotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ROS1 fusion

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Brain Metastases (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 2.2021]

Current EMA Information

- ☒ In this cancer type ☐ In other cancer type ☐ In this cancer type and other cancer types

EMA information is current as of 2022-06-15. For the most up-to-date information, search www.ema.europa.eu/ema.

SLC34A2-ROS1 fusion

☒ crizotinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2022-02-21

Variant class: ROS1 fusion

Reference:

https://www.ema.europa.eu/en/documents/product-information/xalkori-epar-product-information_en.pdf

☒ entrectinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2022-06-08

Variant class: ROS1 fusion

Reference:

https://www.ema.europa.eu/en/documents/product-information/rozlytrek-epar-product-information_en.pdf

Current ESMO Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

ESMO information is current as of 2022-06-01. For the most up-to-date information, search www.esmo.org.

SLC34A2-ROS1 fusion

● entrectinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ROS1 fusion

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

● lorlatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ROS1 fusion

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

● repotrectinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ROS1 fusion

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

SLC34A2-ROS1 fusion (continued)

● crizotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ROS1 fusion

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Stage IV (First-line therapy); ESMO-MCBS v1.1 score: 3
- Stage IV (Second-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

● ceritinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ROS1 fusion

ESMO Level of Evidence/Grade of Recommendation: III / C

Population segment (Line of therapy):

- (Line of therapy not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

Clinical Trials in Taiwan region:

Clinical Trials Summary

SLC34A2-ROS1 fusion

NCT ID	Title	Phase
NCT03093116	A Phase I/II, Open-Label, Multi-Center, First-in-Human Study of the Safety, Tolerability, Pharmacokinetics, and Anti-Tumor Activity of TPX-0005 in Patients With Advanced Solid Tumors Harboring ALK, ROS1, or NTRK1-3 Rearrangements (TRIDENT-1)	I/II
NCT04094610	A Phase I/II, Open-Label, Safety, Tolerability, Pharmacokinetics, and Anti-Tumor Activity Study of Repotrectinib in Pediatric and Young Adult Subjects With Advanced or Metastatic Malignancies Harboring ALK, ROS1, NTRK1-3 Alterations	I/II

Alerts Informed By Public Data Sources

Current FDA Information

 Contraindicated
  Not recommended
  Resistance
  Breakthrough
  Fast Track

FDA information is current as of 2022-06-15. For the most up-to-date information, search www.fda.gov.

SLC34A2-ROS1 fusion

repotrectinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ROS1 positive

Supporting Statement:

The FDA has granted Breakthrough Designation to the ALK/ROS1/TRK inhibitor, repotrectinib, for the treatment of ROS1-positive metastatic non-small cell lung cancer (NSCLC) that has not been treated with a ROS1 tyrosine kinase inhibitor (TKI).

Reference:

<https://ir.tptherapeutics.com/news-releases/news-release-details/turning-point-therapeutics-granted-fda-breakthrough-therapy>

repotrectinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ROS1 positive

Supporting Statement:

The FDA has granted Fast Track Designation to the ALK/ROS1/TRK inhibitor, repotrectinib, for:

- ROS1-positive advanced non-small cell lung cancer (NSCLC) previously treated with one prior platinum chemotherapy and one prior ROS1 TKI.
- ROS1-positive advanced non-small cell lung cancer (NSCLC) without prior ROS1 TKI treatment.
- NTRK fusion positive advanced solid tumors that have progressed following treatment with at least one prior line of chemotherapy and one or two prior TRK TKIs.

Reference:

<https://ir.tptherapeutics.com/news-releases/news-release-details/turning-point-therapeutics-granted-fast-track-designation>

Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

References

1. Bergethson et al. ROS1 rearrangements define a unique molecular class of lung cancers. *J Clin Oncol*. 2012 Mar 10;30(8):863-70. doi: 10.1200/JCO.2011.35.6345. Epub 2012 Jan 3. PMID: 22215748
2. Davare et al. Structural insight into selectivity and resistance profiles of ROS1 tyrosine kinase inhibitors. *Proc Natl Acad Sci U S A*. 2015 Sep 29;112(39):E5381-90. doi: 10.1073/pnas.1515281112. Epub 2015 Sep 8. PMID: 26372962
3. Kohno et al. Beyond ALK-RET, ROS1 and other oncogene fusions in lung cancer. *Transl Lung Cancer Res*. 2015 Apr;4(2):156-64. PMID: 25870798
4. Lin et al. Recent Advances in Targeting ROS1 in Lung Cancer. *J Thorac Oncol*. 2017 Nov;12(11):1611-1625. PMID: 28818606
5. Shaw et al. Crizotinib in ROS1-rearranged non-small-cell lung cancer. *N Engl J Med*. 2014 Nov 20;371(21):1963-71. doi: 10.1056/NEJMoa1406766. Epub 2014 Sep 27. PMID: 25264305
6. Gu et al. Survey of tyrosine kinase signaling reveals ROS kinase fusions in human cholangiocarcinoma. *PLoS ONE*. 2011 Jan 6;6(1):e15640. PMID: 21253578
7. Charest et al. Fusion of FIG to the receptor tyrosine kinase ROS in a glioblastoma with an interstitial del(6)(q21q21). *Genes Chromosomes Cancer*. 2003 May;37(1):58-71. PMID: 12661006
8. Birch et al. Chromosome 3 anomalies investigated by genome wide SNP analysis of benign, low malignant potential and low grade ovarian serous tumours. *PLoS ONE*. 2011;6(12):e28250. PMID: 22163003
9. Lee et al. Identification of ROS1 rearrangement in gastric adenocarcinoma. *Cancer*. 2013 May 1;119(9):1627-35. PMID: 23400546
10. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/212725s005lbl.pdf
11. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/202570s031lbl.pdf
12. Kazandjian et al. Benefit-Risk Summary of Crizotinib for the Treatment of Patients With ROS1 Alteration-Positive, Metastatic Non-Small Cell Lung Cancer. *Oncologist*. 2016 Aug;21(8):974-80. doi: 10.1634/theoncologist.2016-0101. Epub 2016 Jun 21. PMID: 27328934
13. Song et al. Molecular Changes Associated with Acquired Resistance to Crizotinib in ROS1-Rearranged Non-Small Cell Lung Cancer. *Clin Cancer Res*. 2015 May 15;21(10):2379-87. doi: 10.1158/1078-0432.CCR-14-1350. Epub 2015 Feb 16. PMID: 25688157
14. Drilon et al. A Novel Crizotinib-Resistant Solvent-Front Mutation Responsive to Cabozantinib Therapy in a Patient with ROS1-Rearranged Lung Cancer. *Clin Cancer Res*. 2016 May 15;22(10):2351-8. doi: 10.1158/1078-0432.CCR-15-2013. Epub 2015 Dec 16. PMID: 26673800
15. Facchinetti et al. Crizotinib-Resistant ROS1 Mutations Reveal a Predictive Kinase Inhibitor Sensitivity Model for ROS1- and ALK-Rearranged Lung Cancers. *Clin Cancer Res*. 2016 Dec 15;22(24):5983-5991. Epub 2016 Jul 11. PMID: 27401242
16. <https://ir.tptherapeutics.com/news-releases/news-release-details/turning-point-therapeutics-granted-fda-breakthrough-therapy>
17. Lim et al. Open-Label, Multicenter, Phase II Study of Ceritinib in Patients With Non-Small-Cell Lung Cancer Harboring ROS1 Rearrangement. *J Clin Oncol*. 2017 Aug 10;35(23):2613-2618. doi: 10.1200/JCO.2016.71.3701. Epub 2017 May 18. PMID: 28520527
18. NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]
19. Zou et al. PF-06463922, an ALK/ROS1 Inhibitor, Overcomes Resistance to First and Second Generation ALK Inhibitors in Preclinical Models. *Cancer Cell*. 2015 Jul 13;28(1):70-81. PMID: 26144315
20. Zou et al. PF-06463922 is a potent and selective next-generation ROS1/ALK inhibitor capable of blocking crizotinib-resistant ROS1 mutations. *Proc. Natl. Acad. Sci. U.S.A.* 2015 Mar 17;112(11):3493-8. PMID: 25733882
21. Shaw et al. Lorlatinib in non-small-cell lung cancer with ALK or ROS1 rearrangement: an international, multicentre, open-label, single-arm first-in-man phase 1 trial. *Lancet Oncol*. 2017 Dec;18(12):1590-1599. PMID: 29074098