



Sample Information

Patient Name: 陳榮德
Gender: Male
ID No.: F120510469
History No.: 49211554
Age: 55

Ordering Doctor: DOC8549H 楊金蓮
Ordering REQ.: 0CERTWK
Signing in Date: 2023/01/12

Path No.: M112-00005
MP No.: MY23003
Assay: Oncomine Myeloid Assay
Sample Type: Bone Marrow
Bone Marrow Aspirating Date: 2023/01/06

Reporting Doctor: DOC5466K 葉奕成 (Phone: 8#5466)

Note:

Sample Cancer Type: Acute Myeloid Leukemia

Table of Contents	Page	Report Highlights
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2	1 Relevant Biomarkers
Biomarker Descriptions	2	4 Therapies Available
Relevant Therapy Summary	4	0 Clinical Trials
Relevant Therapy Details	4	
Prognostic Details	7	
Diagnostic Details	8	

Relevant Acute Myeloid Leukemia Variants

Gene	Finding	Gene	Finding
ABL1	None detected	MECOM	None detected
ASXL1	None detected	MLLT3	None detected
CEBPA	CEBPA p.(H24Afs*84) c.68_69insC	MYH11	None detected
CREBBP	None detected	NPM1	None detected
FLT3	None detected	NUP214	None detected
IDH1	None detected	RARA	None detected
IDH2	None detected	RUNX1	None detected
KMT2A	None detected	TP53	None detected

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	CEBPA p.(H24Afs*84) c.68_69insC CCAAT enhancer binding protein alpha Allele Frequency: 42.90% Prognostic significance: ELN 2017: Favorable Diagnostic significance: Acute Myeloid Leukemia	cytarabine + daunorubicin cytarabine + idarubicin cytarabine + mitoxantrone gemtuzumab ozogamicin + chemotherapy	None	0

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

Prevalent cancer biomarkers without relevant evidence based on included data sources

DNMT3A p.(I471Lfs*180) c.1410delT, TET2 p.(P554Afs*13) c.1659_1660insG, CSF3R p.(T618I) c.1853C>T

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
CSF3R	p.(T618I)	c.1853C>T	COSM1737962	chr1:36933434	45.10%	NM_156039.3	missense	2000
DNMT3A	p.(I471Lfs*180)	c.1410delT	.	chr2:25469047	47.86%	NM_022552.4	frameshift Deletion	1987
TET2	p.(P554Afs*13)	c.1659_1660insG	.	chr4:106156757	49.00%	NM_001127208.2	frameshift Insertion	1992
CEBPA	p.(H24Afs*84)	c.68_69insC	COSM18922	chr19:33793252	42.90%	NM_004364.4	frameshift Insertion	1979
TET2	p.(A1876T)	c.5626G>A	.	chr4:106197293	47.90%	NM_001127208.2	missense	1998
CEBPA	p.(R297P)	c.890_891delG<insC T	.	chr19:33792430	47.12%	NM_004364.4	missense	1995

Biomarker Descriptions

CEBPA (CCAAT enhancer binding protein alpha)

Background: The CEBPA gene encodes the enhancer binding protein alpha, a member of the basic region leucine zipper family of transcription factors that recognizes the CCAAT promoter¹. CEBPA gives rise to two protein isoforms— p42 and p30, where p30 is the shorter isoform lacking the N-terminal 117 amino acids that is present in p42. Both isoforms contain the basic leucine zipper (bZip) domain involved in hetero/homo-dimerization with other CEBP family members and are required for DNA binding¹. CEBPA is a tumor suppressor gene that plays a critical role in the development of granulocytes¹. Specifically, CEBPA can influence the expression of granulocyte colony-stimulating factor (G-CSF) and interleukin 6 (IL-6), which are required for neutrophil maturation^{2,3}. CEBPA also directly interacts and inhibits cell cycle kinases, including CDK2 and CDK4, thereby hindering cell proliferation⁴. CEBPA is the target of monoallelic or biallelic mutations leading to a loss of function, which can promote the development of cancers such as acute myeloid leukemia (AML)⁵. Germline mutations in CEBPA are also frequent among AML patients and are associated with predisposition to the disease^{6,7}.

Alterations and prevalence: Mutations in CEBPA are reported in 6-18% of all AML cases^{8,9,10,11}. In AML, CEBPA mutations are observed to occur as either monoallelic (single mutant) or bi-allelic (double mutant)^{11,12,13}. Biallelic CEBPA mutations are heterozygous and occur as a specific combination of an N-terminal frameshift on one allele and a C-terminal in frame mutation on the other, referred to as an N/C mutant^{13,14}. Frameshift mutations result in the N-terminal truncation of approximately 120 amino acids while preserving the remaining 300 amino acids that are initiated further downstream¹⁴. C-terminal in-frame mutations disrupt the bZip domain which

Biomarker Descriptions (continued)

interferes with DNA binding and hetero/homo-dimerization with other CEBP family members. Specifically, N/C mutants possess one N-terminal truncated allele coding for the p30 isoform while the other allele codes for either p30 or p42 isoforms harboring C-terminal mutations¹³.

Potential relevance: Biallelic CEBPA mutations are recognized as a diagnostic entity for AML by the World Health Organization (WHO)¹⁵. Biallelic CEBPA mutations are associated with favorable risk and improved prognosis in AML^{11,15,16,17,18}.

CSF3R (colony stimulating factor 3 receptor)

Background: The CSF3R gene encodes the colony stimulating factor 3 trans-membrane receptor for the granulocyte colony-stimulating factor (G-CSF) ligand. CSF3R is a class I membrane-bound cytokine receptor, which lacks intrinsic kinase activity and therefore must interact with downstream proteins for activation¹⁹. Upon ligand activation, CSF3R activates downstream oncogenic pathways by interacting with intracellular signaling proteins through its cytoplasmic tyrosine residues, including proteins from the JAK/STAT, MAPK/ERK, and PI3K/AKT pathways^{19,20}. Nonsense and frameshift mutations in CSF3R target and truncate its cytoplasmic tail, subsequently impairing the internalization signal of the receptor and leading to an overexpression CSF3R on the cell surface^{21,22,23}. Missense mutations in the proximal membrane lead to increased dimerization of the receptor, independent of G-CSF binding^{21,22,23}. These mutations promote constitutive oncogenic JAK-STAT signaling, and increase granulocyte proliferation and survival signaling of hematopoietic progenitor cells^{19,24}.

Alterations and prevalence: CSF3R activating mutations are observed in up to 80% of patients with chronic neutrophilic leukemia (CNL), and in up to 59% of patients with atypical chronic myelogenous leukemia (aCML)^{21,25}. CSF3R mutations occur in 0.5-1% of adult acute myeloid leukemia (AML) and in 2.4% of pediatric AML²⁵. In solid malignancies, CSF3R mutations are observed in up to 5% of uterine carcinosarcoma and skin cutaneous melanoma^{8,26}. Somatic mutations identified in CNL and AML include T615A, T618I, and T640N missense mutations, as well as truncating mutations at Q749, Q754, Y767, S783, Y787, and P820²⁵. T618I mutation is the most frequent variant observed in CNL²⁷.

Potential relevance: CSF3R activating mutations including T618I are a diagnostic criteria for CNL as defined by the World Health Organization (WHO)¹⁵. Mutations in CSF3R are observed in patients with severe congenital neutropenia, which can progress to AML. CSF3R mutations frequently co-occur with CEBPA, and these co-mutations are associated with unfavorable prognosis in AML^{25,28,29}. In independent reports, a CNL patient and an aCML patient with the proximal membrane T618I mutation demonstrated sensitivity to JAK1/2 inhibitor, ruxolitinib^{21,30}.

DNMT3A (DNA methyltransferase 3 alpha)

Background: The DNMT3A gene encodes the DNA methyltransferase 3 alpha which functions as a de novo methyltransferase (DNMT) with equal methylation efficiency for unmethylated and hemimethylated DNA³¹. Methylation of DNA occurs at CpG islands, a region of DNA consisting of sequential cytosine/guanine dinucleotide pairs. CpG island methylation plays an important role in development as well as stem cell regulation. Alterations to global DNA methylation patterns are dependent on DNMTs, which are associated with cancer initiation and progression^{32,33}.

Alterations and prevalence: DNMT3A mutations are observed in approximately 25% of all acute myeloid leukemia (AML) including 29-34% of AML with normal karyotype (NK-AML)^{8,18,34,35,36,37,38}. Mutations in DNMT3A are also reported in 12-18% of myelodysplastic syndromes (MDS) as well as 4-6% of melanoma, lung adenocarcinoma, and uterine cancer^{8,39}. The majority of mutations in DNMT3A are missense however, frameshift, nonsense, and splice site mutations have also been reported^{8,34}. Missense mutations at R882 are most prevalent and are observed to coexist with NPM1 and FLT3 mutations^{40,41}. The R882 mutations occur at the dimer/tetramer interface within the catalytic domain, which leads to disruption of DNMT3A tetramerization and loss of CpG methylation^{42,43}. However, DNMT3A mutations observed in AML at positions other than R882 also contribute to pathogenesis by mechanisms that do not involve methyltransferase activity⁴⁴.

Potential relevance: DNMT3A mutations confer shorter overall survival (OS) in patients with AML including those with NK-AML^{34,37,38,41}. DNMT3A mutations are a useful in the diagnosis of angioimmunoblastic T-cell lymphoma (AITCL) when trying to differentiate from other peripheral T-cell lymphomas (PTCL)⁴⁵.

TET2 (tet methylcytosine dioxygenase 2)

Background: TET2 encodes the tet methylcytosine dioxygenase 2 protein and belongs to a family of ten-eleven translocation (TET) proteins that also includes TET1 and TET3⁴⁶. TET2 is involved in DNA methylation, specifically in the conversion of 5-methylcytosine to 5-hydroxymethylcytosine^{47,48}. The TET proteins contain a C-terminal core catalytic domain that contains a cysteine-rich domain and a double stranded β -helix domain (DSBH)⁴⁹. TET2 is a tumor suppressor gene. Loss of function mutations in TET2 are associated with loss of catalytic activity and transformation to hematological malignancies^{46,47,48}.

Biomarker Descriptions (continued)

Alterations and prevalence: Somatic TET2 mutations, including nonsense, frameshift, splice site, and missense, are observed in 20-25% of myelodysplastic syndrome (MDS) associated diseases, including 40%-60% chronic myelomonocytic leukemia (CMML)³⁹. TET2 mutations at H1881 and R1896 are frequently observed in myeloid malignancies^{47,50}. TET2 mutations are also observed in 9% of uterine, 8% of melanoma and acute myeloid leukemia (AML), as well as 6% of diffuse large B-cell lymphoma (DLBCL).

Potential relevance: The presence of TET2 mutations may be used as one of the major diagnostic criteria in pre-primary myelofibrosis (pre-PMF) and overt PMF in the absence of JAK2/CALR/MPL mutations^{15,51}. TET2 mutations are associated with poor prognosis in PMF and increased rate of transformation to leukemia^{51,52}

Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

CEBPA p.(H24Afs*84) c.68_69insC

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
cytarabine + daunorubicin	×	●	×	×	×
cytarabine + idarubicin	×	●	×	×	×
cytarabine + mitoxantrone	×	●	×	×	×
gemtuzumab ozogamicin + cytarabine + daunorubicin	×	●	×	×	×
gemtuzumab ozogamicin + cytarabine + fludarabine + idarubicin + filgrastim	×	●	×	×	×

Relevant Therapy Details

Current NCCN Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

NCCN information is current as of 2022-11-01. For the most up-to-date information, search www.nccn.org. For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

CEBPA p.(H24Afs*84) c.68_69insC

● cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

CEBPA p.(H24Afs*84) c.68_69insC (continued)**● cytarabine + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

● cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

● cytarabine + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

● cytarabine + mitoxantrone

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

● gemtuzumab ozogamicin + cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

CEBPA p.(H24Afs*84) c.68_69insC (continued)**● gemtuzumab ozogamicin + cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

● gemtuzumab ozogamicin + cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

Prognostic Details

Current NCCN Information

NCCN information is current as of 2022-11-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

CEBPA p.(H24Afs*84) c.68_69insC

Prognostic significance: ELN 2017: Favorable

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 2A

Summary:

- Genetic Abnormality: Biallelic mutated CEBPA.

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

Current ESMO Information

ESMO information is current as of 2022-11-01. For the most up-to-date information, search www.esmo.org.

CEBPA p.(H24Afs*84) c.68_69insC

Prognostic significance: ELN 2017: Favorable

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

Summary:

- Double-mutant CEBPA

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

Diagnostic Details

Current ESMO Information

ESMO information is current as of 2022-11-01. For the most up-to-date information, search www.esmo.org.

CEBPA p.(H24Afs*84) c.68_69insC

Diagnostic significance: Acute Myeloid Leukemia

Variant class: CEBPA mutation

Diagnostic notes:

- CEBPA biallelic mutations; AML with recurrent genetic abnormalities; WHO classification of AML

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

References

1. Pabst et al. Complexity of CEBPA dysregulation in human acute myeloid leukemia. *Clin. Cancer Res.* 2009 Sep 1;15(17):5303-7. PMID: 19706798
2. Zhang et al. Upregulation of interleukin 6 and granulocyte colony-stimulating factor receptors by transcription factor CCAAT enhancer binding protein alpha (C/EBP alpha) is critical for granulopoiesis. *J. Exp. Med.* 1998 Sep 21;188(6):1173-84. PMID: 9743535
3. Heath et al. C/EBPalpha deficiency results in hyperproliferation of hematopoietic progenitor cells and disrupts macrophage development in vitro and in vivo. *Blood.* 2004 Sep 15;104(6):1639-47. PMID: 15073037
4. Wang et al. C/EBPalpha arrests cell proliferation through direct inhibition of Cdk2 and Cdk4. *Mol. Cell.* 2001 Oct;8(4):817-28. PMID: 11684017
5. Lourenço et al. A tumor suppressor role for C/EBPa in solid tumors: more than fat and blood. *Oncogene.* 2017 Sep 14;36(37):5221-5230. PMID: 28504718
6. Pabst et al. Somatic CEBPA mutations are a frequent second event in families with germline CEBPA mutations and familial acute myeloid leukemia. *J. Clin. Oncol.* 2008 Nov 1;26(31):5088-93. PMID: 18768433
7. Tawana et al. Disease evolution and outcomes in familial AML with germline CEBPA mutations. *Blood.* 2015 Sep 3;126(10):1214-23. PMID: 26162409
8. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
9. Benthous et al. Rapid and sensitive screening for CEBPA mutations in acute myeloid leukaemia. *Br. J. Haematol.* 2008 Oct;143(2):230-9. PMID: 18752591
10. Su et al. Mutational spectrum of acute myeloid leukemia patients with double CEBPA mutations based on next-generation sequencing and its prognostic significance. *Oncotarget.* 2018 May 18;9(38):24970-24979. PMID: 29861846
11. Dufour et al. Acute myeloid leukemia with biallelic CEBPA gene mutations and normal karyotype represents a distinct genetic entity associated with a favorable clinical outcome. *J. Clin. Oncol.* 2010 Feb 1;28(4):570-7. PMID: 20038735
12. Green et al. Prognostic significance of CEBPA mutations in a large cohort of younger adult patients with acute myeloid leukemia: impact of double CEBPA mutations and the interaction with FLT3 and NPM1 mutations. *J. Clin. Oncol.* 2010 Jun 1;28(16):2739-47. PMID: 20439648
13. Avellino et al. Expression and regulation of C/EBPa in normal myelopoiesis and in malignant transformation. *Blood.* 2017 Apr 13;129(15):2083-2091. PMID: 28179278
14. Pabst et al. Transcriptional dysregulation during myeloid transformation in AML. *Oncogene.* 2007 Oct 15;26(47):6829-37. PMID: 17934489
15. Arber et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood.* 2016 May 19;127(20):2391-405. PMID: 27069254
16. Kühnl et al. Molecular markers in acute myeloid leukaemia. *Int. J. Hematol.* 2012 Aug;96(2):153-63. PMID: 22791509
17. Barjesteh et al. Biallelic mutations in the CEBPA gene and low CEBPA expression levels as prognostic markers in intermediate-risk AML. *Hematol. J.* 2003;4(1):31-40. PMID: 12692518
18. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]
19. Zhang et al. Gain-of-function mutations in granulocyte colony-stimulating factor receptor (CSF3R) reveal distinct mechanisms of CSF3R activation. *J. Biol. Chem.* 2018 May 11;293(19):7387-7396. PMID: 29572350
20. Nicholson et al. Distinct regions of the granulocyte colony-stimulating factor receptor are required for tyrosine phosphorylation of the signaling molecules JAK2, Stat3, and p42, p44MAPK. *Blood.* 1995 Nov 15;86(10):3698-704. PMID: 7579336
21. Maxson et al. Oncogenic CSF3R mutations in chronic neutrophilic leukemia and atypical CML. *N. Engl. J. Med.* 2013 May 9;368(19):1781-90. PMID: 23656643
22. Dong et al. The carboxyl terminus of the granulocyte colony-stimulating factor receptor, truncated in patients with severe congenital neutropenia/acute myeloid leukemia, is required for SH2-containing phosphatase-1 suppression of Stat activation. *J. Immunol.* 2001 Dec 1;167(11):6447-52. PMID: 11714811
23. van et al. G-CSF receptor truncations found in SCN/AML relieve SOCS3-controlled inhibition of STAT5 but leave suppression of STAT3 intact. *Blood.* 2004 Aug 1;104(3):667-74. PMID: 15069015
24. Panopoulos et al. Granulocyte colony-stimulating factor: molecular mechanisms of action during steady state and 'emergency' hematopoiesis. *Cytokine.* 2008 Jun;42(3):277-88. PMID: 18400509
25. Maxson et al. CSF3R mutations have a high degree of overlap with CEBPA mutations in pediatric AML. *Blood.* 2016 Jun 16;127(24):3094-8. PMID: 27143256

References (continued)

26. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
27. Pardanani et al. CSF3R T618I is a highly prevalent and specific mutation in chronic neutrophilic leukemia. *Leukemia.* 2013 Sep;27(9):1870-3. PMID: 23604229
28. Dong et al. Mutations in the gene for the granulocyte colony-stimulating-factor receptor in patients with acute myeloid leukemia preceded by severe congenital neutropenia. *N. Engl. J. Med.* 1995 Aug 24;333(8):487-93. PMID: 7542747
29. Su et al. CSF3R mutations were associated with an unfavorable prognosis in patients with acute myeloid leukemia with CEBPA double mutations. *Ann. Hematol.* 2019 Jul;98(7):1641-1646. PMID: 31041512
30. Dao et al. Significant clinical response to JAK1/2 inhibition in a patient with CSF3R-T618I-positive atypical chronic myeloid leukemia. *Leuk Res Rep.* 2014;3(2):67-9. PMID: 25180155
31. Okano et al. Cloning and characterization of a family of novel mammalian DNA (cytosine-5) methyltransferases. *Nat Genet.* 1998 Jul;19(3):219-20. PMID: 9662389
32. Fernandez et al. A DNA methylation fingerprint of 1628 human samples. *Genome Res.* 2012 Feb;22(2):407-19. PMID: 21613409
33. Jones et al. The epigenomics of cancer. *Cell.* 2007 Feb 23;128(4):683-92. PMID: 17320506
34. Ley et al. DNMT3A mutations in acute myeloid leukemia. *N. Engl. J. Med.* 2010 Dec 16;363(25):2424-33. PMID: 21067377
35. Marková et al. Prognostic impact of DNMT3A mutations in patients with intermediate cytogenetic risk profile acute myeloid leukemia. *Eur. J. Haematol.* 2012 Feb;88(2):128-35. PMID: 21967546
36. Yang et al. DNMT3A in haematological malignancies. *Nat. Rev. Cancer.* 2015 Mar;15(3):152-65. PMID: 25693834
37. Renneville et al. Prognostic significance of DNA methyltransferase 3A mutations in cytogenetically normal acute myeloid leukemia: a study by the Acute Leukemia French Association. *Leukemia.* 2012 Jun;26(6):1247-54. PMID: 22289988
38. Marcucci et al. Age-related prognostic impact of different types of DNMT3A mutations in adults with primary cytogenetically normal acute myeloid leukemia. *J. Clin. Oncol.* 2012 Mar 1;30(7):742-50. PMID: 22291079
39. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]
40. Kumar et al. DNMT3A (R882) mutation features and prognostic effect in acute myeloid leukemia in Coexistent with NPM1 and FLT3 mutations. *Hematol Oncol Stem Cell Ther.* 2018 Jun;11(2):82-89. PMID: 29079128
41. Thol et al. Incidence and prognostic influence of DNMT3A mutations in acute myeloid leukemia. *J. Clin. Oncol.* 2011 Jul 20;29(21):2889-96. PMID: 21670448
42. Sandoval et al. Mutations in the DNMT3A DNA methyltransferase in acute myeloid leukemia patients cause both loss and gain of function and differential regulation by protein partners. *J. Biol. Chem.* 2019 Mar 29;294(13):4898-4910. PMID: 30705090
43. Holz-Schietinger et al. Mutations in DNA methyltransferase (DNMT3A) observed in acute myeloid leukemia patients disrupt processive methylation. *J. Biol. Chem.* 2012 Sep 7;287(37):30941-51. PMID: 22722925
44. Russler-Germain et al. The R882H DNMT3A mutation associated with AML dominantly inhibits wild-type DNMT3A by blocking its ability to form active tetramers. *Cancer Cell.* 2014 Apr 14;25(4):442-54. PMID: 24656771
45. NCCN Guidelines® - NCCN-T-Cell Lymphomas [Version 2.2022]
46. Pan et al. The TET2 interactors and their links to hematological malignancies. *IUBMB Life.* 2015 Jun;67(6):438-45. PMID: 26099018
47. Ko et al. Impaired hydroxylation of 5-methylcytosine in myeloid cancers with mutant TET2. *Nature.* 2010 Dec 9;468(7325):839-43. PMID: 21057493
48. Solary et al. The Ten-Eleven Translocation-2 (TET2) gene in hematopoiesis and hematopoietic diseases. *Leukemia.* 2014 Mar;28(3):485-96. PMID: 24220273
49. An et al. TET family dioxygenases and DNA demethylation in stem cells and cancers. *Exp. Mol. Med.* 2017 Apr 28;49(4):e323. PMID: 28450733
50. Kosmider et al. TET2 mutation is an independent favorable prognostic factor in myelodysplastic syndromes (MDSs). *Blood.* 2009 Oct 8;114(15):3285-91. PMID: 19666869
51. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 3.2022]
52. Lundberg et al. Clonal evolution and clinical correlates of somatic mutations in myeloproliferative neoplasms. *Blood.* 2014 Apr 3;123(14):2220-8. PMID: 24478400