



## Sample Information

**Patient Name:** 洪玟臻  
**Gender:** Female  
**ID No.:** A224879886  
**History No.:** 46338005  
**Age:** 43

**Ordering Doctor:** DOC5636D 吳紋綺  
**Ordering REQ.:** 0BWTMGW  
**Signing in Date:** 2022/06/23

**Path No.:** S111-99612  
**MP No.:** F22064  
**Assay:** Oncomine Focus Assay  
**Sample Type:** FFPE  
**Block No.:** S111-76392B  
**Percentage of tumor cells:** 90%

**Reporting Doctor:** DOC5466K 葉奕成 (Phone: 8#5466)

**Note:**

## Sample Cancer Type: Melanoma

Table of Contents	Page
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2
Biomarker Descriptions	2
Relevant Therapy Summary	3
Relevant Therapy Details	4
Clinical Trials Summary	5
Alert Details	5

**Report Highlights**  
2 Relevant Biomarkers  
1 Therapies Available  
1 Clinical Trials

## Relevant Melanoma Variants

Gene	Finding
BRAF	None detected
KIT	<b>KIT p.(L576P) c.1727T&gt;C, KIT amplification</b>
NRAS	None detected
NTRK1	None detected
NTRK2	None detected
NTRK3	None detected

## Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	<b>KIT p.(L576P) c.1727T&gt;C</b> KIT proto-oncogene, receptor tyrosine kinase Allele Frequency: 43.38%	imatinib	imatinib	0
IIC	<b>BAG4-FGFR1 fusion</b> BAG cochaperone 4 - fibroblast growth factor receptor 1	None	None	1

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

Tier Reference: Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

### Prevalent cancer biomarkers without relevant evidence based on included data sources

*KIT amplification*

## Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

### DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
KIT	p.(L576P)	c.1727T>C	COSM1290	chr4:55593661	43.38%	NM_000222.3	missense	1948

### Gene Fusions (RNA)

Genes	Variant ID	Locus	Read Count
BAG4-FGFR1	BAG4-FGFR1.B1F2	chr8:38034657 - chr8:38315052	753

### Copy Number Variations

Gene	Locus	Copy Number
KIT	chr4:55529117	7.41

## Biomarker Descriptions

### FGFR1 (fibroblast growth factor receptor 1)

**Background:** The FGFR1 gene encodes fibroblast growth receptor 1, a member of the fibroblast growth factor receptor (FGFR) family that also includes FGFR2, 3, and 4. These proteins are single transmembrane receptors composed of three extracellular immunoglobulin (Ig)-type domains and an intracellular kinase domain. Upon FGF-mediated stimulation, FGFRs activate several oncogenic signaling pathways, including the RAS/RAF/MEK/ERK, PI3K/AKT/MTOR, PLC/PKC, and JAK/STAT pathways influencing cell proliferation, migration, and survival<sup>1,2,3</sup>.

**Alterations and prevalence:** Recurrent somatic alterations common to the FGFR family include gene amplification, mutation, and chromosomal translocations leading to FGFR fusions<sup>4</sup>. Amplification of FGFR1 is observed in 15-20% of squamous lung cancer, 10-15% of breast cancer, 8% of bladder cancer, and 2-5% of uterine cancer cases<sup>5,6,7,8,9</sup>. The most common recurrent mutations, N546K and K656E, are relatively infrequent (<1%); they activate mutations in the kinase domain and are distributed in diverse cancer types<sup>10</sup>. FGFR1 translocations giving rise to expressed fusions are common in certain hematological cancers, but less common in solid tumors<sup>11,12,13</sup>.

**Potential relevance:** The FDA has granted fast-track designation (2018) to Debio 1347<sup>14</sup> for solid tumors harboring aberrations in FGFR1, FGFR2, or FGFR3. FDA has approved multi-kinase inhibitors, including regorafenib, ponatinib, lenvatinib, nintedanib, and

## Biomarker Descriptions (continued)

pazopanib, that are known to inhibit FGFR family members. These inhibitors have demonstrated anti-tumor activity in select cancer types with FGFR alterations<sup>15,16,17,18,19,20,21</sup>. In a phase II clinical trial, dovitinib, a multi-tyrosine kinase inhibitor (TKI), exhibited an overall response rate (ORR) of 11.5% and a disease control rate (DCR) of 50% in patients with advanced squamous cell lung cancer possessing FGFR1 amplification. The patients had a median overall survival (OS) of 5 months and progression-free survival (PFS) of 2.9 months<sup>22</sup>. Likewise, in a phase Ib study testing the FGFR inhibitor AZD4547, median OS was 4.9 months in patients with FGFR1-amplified advanced squamous cell lung cancer. One of 13 (8%) patients achieved a partial response, 4 (31%) exhibited stable disease, and 2 (13.3%) demonstrated PFS at 12 weeks<sup>23</sup>.

### KIT (KIT proto-oncogene, receptor tyrosine kinase)

**Background:** The KIT gene, also known as CD117, encodes the KIT proto-oncogene receptor tyrosine kinase (c-KIT), a member of the PDGF receptor type III receptor tyrosine kinase family, which includes PDGFRA, PDGFRB, CSF1R, FLT1, FLT3, FLT4 and KDR<sup>24,25</sup>. KIT is a receptor for stem cell factor, important in regulating growth and development of hematopoietic cells<sup>26</sup>. The KIT gene is flanked by the PDGFRA and KDR genes on chromosome 4q12. Ligand binding to KIT results in kinase activation and stimulation of downstream pathways including the RAS/RAF/MEK/ERK and PI3K/AKT/MTOR pathways promoting cell proliferation and survival<sup>27</sup>.

**Alterations and prevalence:** Recurrent somatic KIT alterations are observed in both solid and hematological cancers and include activating mutations such as single nucleotide variants, small duplications, and complex in-frame insertions or deletions (indels). Mutations in KIT exons 8, 9, 11, and 17 disrupt auto-inhibitory mechanisms and lead to constitutive activity<sup>28</sup>. Gain of function mutations are found in up to 70% of mast cell tumors, 17% of nasal T-cell lymphomas, and 9% of dysgerminoma<sup>29</sup>. Somatic mutations in exon 11 occur in 60-70% of all gastrointestinal stromal tumor (GIST), whereas alterations in exons 8 and 17 are more common in myeloid cancers<sup>9,28,29</sup>. A common kinase domain mutation that causes ligand-independent constitutive activation, D816V, occurs in 80-93% of aggressive forms of mastocytosis<sup>30,31</sup>.

**Potential relevance:** Imatinib<sup>32</sup> (2001) is approved for KIT positive malignant GIST and adult patients with aggressive systemic mastocytosis (SM) harboring D816V mutations. Imatinib is also recommended for KIT activating mutations in melanoma and exon 9 and 11 mutations in GIST<sup>33,34,35</sup>. Mutations in exon 17 have been identified to confer resistance to imatinib and sunitinib<sup>36</sup>. Patients with acute myeloid leukemia (AML) that harbor KIT activating mutations with t(8;21) and inv(16) have an increased risk of relapse<sup>37</sup>. KIT D816V mutation is associated with the diagnosis of SM and aggressiveness of the disease<sup>38,39</sup>.

## Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

### KIT p.(L576P) c.1727T>C

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
imatinib	×	🕒	×	×	×

### BAG4-FGFR1 fusion

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
erdafitinib	×	×	×	×	● (II)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Details

### Current NCCN Information

- ☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types

NCCN information is current as of 2022-03-31. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org).  
For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

#### KIT p.(L576P) c.1727T>C

##### ☒ imatinib

Cancer type: Cutaneous Melanoma

Variant class: KIT L576P mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Metastatic, Unresectable, Progression (Second-line therapy, Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Cutaneous Melanoma [Version 2.2022]

##### ☐ imatinib

Cancer type: Gastrointestinal Stromal Tumor

Variant class: KIT exon 11 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Resectable (Neoadjuvant therapy); Preferred intervention
- Resected (Adjuvant therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Gastrointestinal Stromal Tumor [Version 1.2022]

## Clinical Trials in Taiwan region:

### Clinical Trials Summary

#### BAG4-FGFR1 fusion

NCT ID	Title	Phase
NCT04083976	A Phase II Study of Erdafitinib in Subjects With Advanced Solid Tumors and FGFR Gene Alterations.	II

### Alerts Informed By Public Data Sources

#### Current FDA Information

 Contraindicated  Not recommended  Resistance  Breakthrough  Fast Track

FDA information is current as of 2022-04-13. For the most up-to-date information, search [www.fda.gov](https://www.fda.gov).

#### BAG4-FGFR1 fusion

#### Debio 1347

Cancer type: Solid Tumor

Variant class: FGFR1 aberration

##### Supporting Statement:

The FDA has granted Fast Track Designation to the FGFR 1-3 inhibitor, debio 1347, for FGFR1/2/3 alterations in unresectable or metastatic solid tumors.

##### Reference:

<https://www.debiopharm.com/drug-development/press-releases/fda-grants-fast-track-designation-to-debiopharm-internationals-debio-1347-for-the-treatment-of-patients-with-unresectable-or-metastatic-tumors-with-a-specific-fgfr-gene-alteration/>

## Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

## References

1. Babina et al. Advances and challenges in targeting FGFR signalling in cancer. *Nat. Rev. Cancer*. 2017 May;17(5):318-332. PMID: 28303906
2. Ahmad et al. Mechanisms of FGFR-mediated carcinogenesis. *Biochim. Biophys. Acta*. 2012 Apr;1823(4):850-60. PMID: 22273505
3. Sarabipour et al. Mechanism of FGF receptor dimerization and activation. *Nat Commun*. 2016 Jan 4;7:10262. doi: 10.1038/ncomms10262. PMID: 26725515
4. Helsten et al. The FGFR Landscape in Cancer: Analysis of 4,853 Tumors by Next-Generation Sequencing. *Clin. Cancer Res*. 2016 Jan 1;22(1):259-67. PMID: 26373574
5. Peter et al. Comprehensive genomic characterization of squamous cell lung cancers. *Nature*. 2012 Sep 27;489(7417):519-25. PMID: 22960745
6. Ciriello et al. Comprehensive Molecular Portraits of Invasive Lobular Breast Cancer. *Cell*. 2015 Oct 8;163(2):506-19. PMID: 26451490
7. Cancer et al. Integrated genomic characterization of endometrial carcinoma. *Nature*. 2013 May 2;497(7447):67-73. PMID: 23636398
8. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet*. 2013 Oct;45(10):1113-20. PMID: 24071849
9. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov*. 2012 May;2(5):401-4. PMID: 22588877
10. Lew et al. The precise sequence of FGF receptor autophosphorylation is kinetically driven and is disrupted by oncogenic mutations. *Sci Signal*. 2009 Feb 17;2(58):ra6. PMID: 19224897
11. Jackson et al. 8p11 myeloproliferative syndrome: a review. *Hum. Pathol*. 2010 Apr;41(4):461-76. PMID: 20226962
12. Li et al. Identification of a novel partner gene, TPR, fused to FGFR1 in 8p11 myeloproliferative syndrome. *Genes Chromosomes Cancer*. 2012 Sep;51(9):890-7. PMID: 22619110
13. Wasag et al. The kinase inhibitor TKI258 is active against the novel CUX1-FGFR1 fusion detected in a patient with T-lymphoblastic leukemia/lymphoma and t(7;8)(q22;p11). *Haematologica*. 2011 Jun;96(6):922-6. PMID: 21330321
14. <https://www.debiopharm.com/drug-development/press-releases/fda-grants-fast-track-designation-to-debiopharm-internationals-debio-1347-for-the-treatment-of-patients-with-unresectable-or-metastatic-tumors-with-a-specific-fgfr-gene-alteration/>
15. Cha et al. FGFR2 amplification is predictive of sensitivity to regorafenib in gastric and colorectal cancers in vitro. *Mol Oncol*. 2018 Jun;12(7):993-1003. PMID: 29573334
16. Chae et al. Inhibition of the fibroblast growth factor receptor (FGFR) pathway: the current landscape and barriers to clinical application. *Oncotarget*. 2017 Feb 28;8(9):16052-16074. PMID: 28030802
17. Porta et al. FGFR a promising druggable target in cancer: Molecular biology and new drugs. *Crit. Rev. Oncol. Hematol*. 2017 May;113:256-267. PMID: 28427515
18. Gozgit et al. Ponatinib (AP24534), a multitargeted pan-FGFR inhibitor with activity in multiple FGFR-amplified or mutated cancer models. *Mol. Cancer Ther*. 2012 Mar;11(3):690-9. PMID: 22238366
19. Yamamoto et al. Lenvatinib, an angiogenesis inhibitor targeting VEGFR/FGFR, shows broad antitumor activity in human tumor xenograft models associated with microvessel density and pericyte coverage. *Vasc Cell*. 2014 Sep 6;6:18. doi: 10.1186/2045-824X-6-18. eCollection 2014. PMID: 25197551
20. Kim et al. Pazopanib, a novel multitargeted kinase inhibitor, shows potent in vitro antitumor activity in gastric cancer cell lines with FGFR2 amplification. *Mol. Cancer Ther*. 2014 Nov;13(11):2527-36. PMID: 25249557
21. Hibi et al. FGFR gene alterations in lung squamous cell carcinoma are potential targets for the multikinase inhibitor nintedanib. *Cancer Sci*. 2016 Nov;107(11):1667-1676. PMID: 27581340
22. Lim et al. Efficacy and safety of dovitinib in pretreated patients with advanced squamous non-small cell lung cancer with FGFR1 amplification: A single-arm, phase 2 study. *Cancer*. 2016 Oct;122(19):3024-31. PMID: 27315356
23. Paik et al. A Phase Ib Open-Label Multicenter Study of AZD4547 in Patients with Advanced Squamous Cell Lung Cancers. *Clin. Cancer Res*. 2017 Sep 15;23(18):5366-5373. PMID: 28615371
24. Ségaliny et al. Receptor tyrosine kinases: Characterisation, mechanism of action and therapeutic interests for bone cancers. *J Bone Oncol*. 2015 Mar;4(1):1-12. PMID: 26579483
25. Berenstein. Class III Receptor Tyrosine Kinases in Acute Leukemia - Biological Functions and Modern Laboratory Analysis. *Biomark Insights*. 2015;10(Suppl 3):1-14. PMID: 26309392
26. Ashman. The biology of stem cell factor and its receptor C-kit. *Int. J. Biochem. Cell Biol*. 1999 Oct;31(10):1037-51. PMID: 10582338

## References (continued)

27. Cardoso et al. The SCF/c-KIT system in the male: Survival strategies in fertility and cancer. *Mol. Reprod. Dev.* 2014 Dec;81(12):1064-79. PMID: 25359157
28. Abbaspour et al. Receptor tyrosine kinase (c-Kit) inhibitors: a potential therapeutic target in cancer cells. *Drug Des Devel Ther.* 2016;10:2443-59. PMID: 27536065
29. Liang et al. The C-kit receptor-mediated signal transduction and tumor-related diseases. *Int. J. Biol. Sci.* 2013;9(5):435-43. PMID: 23678293
30. Garcia-Montero et al. KIT mutation in mast cells and other bone marrow hematopoietic cell lineages in systemic mast cell disorders: a prospective study of the Spanish Network on Mastocytosis (REMA) in a series of 113 patients. *Blood.* 2006 Oct 1;108(7):2366-72. PMID: 16741248
31. Chatterjee et al. Mastocytosis: a mutated KIT receptor induced myeloproliferative disorder. *Oncotarget.* 2015 Jul 30;6(21):18250-64. PMID: 26158763
32. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/021588s060lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/021588s060lbl.pdf)
33. NCCN Guidelines® - NCCN-Cutaneous Melanoma [Version 2.2022]
34. NCCN Guidelines® - NCCN-Soft Tissue Sarcoma [Version 1.2022]
35. Casali et al. Gastrointestinal stromal tumours: ESMO-EURACAN Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann. Oncol.* 2018 Oct 1;29(Supplement\_4):iv68-iv78. PMID: 29846513
36. Jonathan. KIT Oncogenic Mutations: Biologic Insights, Therapeutic Advances, and Future Directions. *Cancer Res.* 2016 Nov 1;76(21):6140-6142. PMID: 27803101
37. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 1.2022]
38. Lim et al. Systemic mastocytosis in 342 consecutive adults: survival studies and prognostic factors. *Blood.* 2009 Jun 4;113(23):5727-36. PMID: 19363219
39. Verstovsek. Advanced systemic mastocytosis: the impact of KIT mutations in diagnosis, treatment, and progression. *Eur. J. Haematol.* 2013 Feb;90(2):89-98. PMID: 23181448