

ACT Onco[®] + Report

PATIENT		
Identifier: 歐陽國華		Patient ID: 49497941
Date of Birth: Sep 28, 1963		Gender: Male
Diagnosis: Pancreatic cancer		
ORDERING PHYSICIAN		
Name: 姜乃榕醫師		Tel: 886-228712121
Facility: 臺北榮總		
Address: 臺北市北投區石牌路二段 201 號		
SPECIMEN		
Specimen ID: S11291425A	Collection site: Pancreas	Type: FFPE tissue
Date received: May 16, 2023	Lab ID: AA-23-03087	D/ID: NA

ABOUT ACT Onco[®]+

The test is a next-generation sequencing (NGS)-based assay developed for efficient and comprehensive genomic profiling of cancers. This test interrogates coding regions of 440 genes associated with cancer treatment, prognosis and diagnosis. Genetic mutations detected by this test include small-scale mutations like single nucleotide variants (SNVs), small insertions and deletions (InDels) (≤ 15 nucleotides) and large-scale genomic alterations like copy number alterations (CNAs). The test also includes an RNA test, detecting fusion transcripts of 13 genes.

SUMMARY FOR ACTIONABLE VARIANTS

VARIANTS/BIOMARKERS WITH EVIDENCE OF CLINICAL SIGNIFICANCE

Genomic Alterations/Biomarkers	Probable Effects in Patient's Cancer Type		Probable Sensitive in Other Cancer Types
	Sensitive	Resistant	
BRCA2 S1882*	Olaparib, Rucaparib	-	Niraparib, Talazoparib

VARIANTS/BIOMARKERS WITH POTENTIAL CLINICAL SIGNIFICANCE

Genomic Alterations/Biomarkers	Possibly Sensitive		Possibly Resistant
KRAS G12R	-		Cetuximab, Panitumumab

Note:

- The above summary tables present genomic variants and biomarkers based on the three-tiered approach proposed by US FDA for reporting tumor profiling NGS testing. "Variants/biomarkers with evidence of clinical significance" refers to mutations that are widely recognized as standard-of-care biomarkers (FDA level 2/AMP tier 1). "Variants/biomarkers with potential clinical significance" refers to mutations that are not included in the standard of care but are informational for clinicians, which are commonly biomarkers used as inclusion criteria for clinical trials (FDA level 3/AMP tier 2).
- The therapeutic agents and possible effects to a given drug are based on mapping the variants/biomarkers with ACT Genomics clinical knowledge database. The mapping results only provide information for reference, but not medical recommendation.
- Please refer to corresponding sections for more detailed information about genomic alteration and clinical relevance listed above.

ACTOnco® + Report

TESTING RESULTS

VARIANT(S) WITH CLINICAL RELEVANCE

- Single Nucleotide and Small InDel Variants

Gene	Amino Acid Change	Allele Frequency
<i>BRCA2</i>	S1882*	59.5%
<i>KRAS</i>	G12R	18.7%
<i>SERPINB3</i>	T155R	35.8%
<i>TP53</i>	P278S	22.3%

- Copy Number Alterations

Chromosome	Gene	Variation	Copy Number
Chr9	<i>CDKN2A</i>	Homozygous deletion	0
Chr10	<i>PTEN</i>	Heterozygous deletion	1
Chr17	<i>TP53</i>	Heterozygous deletion	1
Chr18	<i>SMAD4</i>	Heterozygous deletion	1
Chr4	<i>FBXW7</i>	Heterozygous deletion	1

- Fusions

Fusion Gene & Exon	Transcript ID
No fusion gene detected in this sample	

- Immune Checkpoint Inhibitor (ICI) Related Biomarkers

Biomarker	Results
Tumor Mutational Burden (TMB)	6.3 muts/Mb
Microsatellite Instability (MSI)	Microsatellite stable (MSS)

Note:

- Variant(s) enlisted in the SNV table may currently exhibit no relevance to treatment response prediction. Please refer to INTERPRETATION for more biological information and/or potential clinical impacts of the variants.
- Loss of heterozygosity (LOH) information was used to infer tumor cellularity. Copy number alteration in the tumor was determined based on 36% tumor purity.
- For more therapeutic agents which are possibly respond to heterozygous deletion of genes listed above, please refer to APPENDIX for more information.
- TMB was calculated by using the sequenced regions of ACTOnco®+ to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The threshold for high mutation load is set at ≥ 7.5 mutations per megabase. TMB, microsatellite status and gene copy number deletion cannot be determined if calculated tumor purity is $< 30\%$.

ACT Onco[®] + Report

THERAPEUTIC IMPLICATIONS TARGETED THERAPIES

Genomic Alterations	Therapies	Effect
Level 1		
BRCA2 S1882*	Olaparib	sensitive
Level 2		
BRCA2 S1882*	Rucaparib	sensitive
Level 3A		
BRCA2 S1882*	Niraparib, Talazoparib	sensitive
KRAS G12R	Cetuximab, Panitumumab	resistant

Therapies associated with benefit or lack of benefit are based on biomarkers detected in this tumor and published evidence in professional guidelines or peer-reviewed journals.

Level	Description
1	FDA-recognized biomarkers predictive of response or resistance to FDA approved drugs in this indication
2	Standard care biomarkers (recommended by the NCCN guideline) predictive of response or resistance to FDA approved drugs in this indication
3A	Biomarkers predictive of response or resistance to therapies approved by the FDA or NCCN guideline in a different cancer type
3B	Biomarkers that serve as inclusion criteria for clinical trials (minimal supportive data required)
4	Biomarkers that show plausible therapeutic significance based on small studies, few case reports, or preclinical studies

ACT Onco[®] + Report

IMMUNE CHECKPOINT INHIBITORS (ICIs)

No genomic alterations detected to confer sensitivity or lack of benefit to immune checkpoint therapies.

- Other Biomarkers with Potential Clinical Effects for ICIs

Genomic Alterations	Potential Clinical Effects
Not detected	

Note: Tumor non-genomic factors, such as patient germline genetics, PDL1 expression, tumor microenvironment, epigenetic alterations or other factors not provided by this test may affect ICI response.

CHEMOTHERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to chemotherapies.

HORMONAL THERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to hormonal therapies.

OTHERS

Pharmacogenomic implication

Gene	Detection Site	Genotype	Drug Impact	Level of Evidence*
UGT1A1	rs4148323	AG	Irinotecan-based regimens	Level 1B

Clinical Interpretation:

Patients with the AG genotype and cancer who are treated with irinotecan-based regimens may have an increased risk of diarrhea and neutropenia as compared to patients with the GG genotype, or a decreased risk of diarrhea and neutropenia compared to patients with the AA genotype. Other genetic and clinical factors may also influence a patient's risk of diarrhea and neutropenia.

* Level of evidence was defined by PharmGKB (<https://www.pharmgkb.org/page/clinAnnLevels>)

Level 1A: Clinical annotations describe variant-drug combinations that have variant-specific prescribing guidance available in a current clinical guideline annotation or an FDA-approved drug label annotation.

Level 1B: Clinical annotations describe variant-drug combinations with a high level of evidence supporting the association but no variant-specific prescribing guidance in an annotated clinical guideline or FDA drug label.

Level 2A: Variants in Level 2A clinical annotations are found in PharmGKB's Tier 1 Very Important Pharmacogenes (VIPs). These variants are in known pharmacogenes, implying causation of drug phenotype is more likely.

Note:

Therapeutic implications provided in the test are based solely on the panel of 440 genes sequenced. Therefore, alterations in genes not covered in this panel, epigenetic and post-transcriptional and post-translational factors may also determine a patient's response to therapies. In addition, several other patient-associated clinical factors, including but not limited to, prior lines of therapies received, dosage and combinations with other therapeutic agents, patient's cancer types, sub-types, and/or stages, may also determine the patient's clinical response to therapies.

ACT Onco[®] + Report

VARIANT INTERPRETATION

BRCA2 S1882*

Biological Impact

The BRCA2 gene encodes a tumor suppressor involved in the homologous recombination pathway for double-strand DNA repair^[1]. BRCA2 has been implicated as a haploinsufficient gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions^[2]. BRCA2 germline mutations confer an increased lifetime risk of developing breast, ovarian, prostate and pancreatic cancer, limited reports of related gastric cancer, and Fanconi anemia subtype D1-associated risk of brain cancer, medulloblastoma, pharyngeal cancer, chronic lymphocytic leukemia and acute myeloid leukemia^[3]. Somatic mutations in BRCA2 are highest in colorectal, non-small cell lung cancer (NSCLC), and ovarian cancers^[4].

S1882* mutation results in a premature truncation of the BRCA2 protein at amino acid 1882 (UniProtKB). This mutation is predicted to lead to a loss of BRCA2 function, despite not having characterized in the literature.

Therapeutic and prognostic relevance

Multiple PARP inhibitors, including olaparib, rucaparib, niraparib, and talazoparib, have been approved by the U.S. FDA for the treatment of cancer. Olaparib is approved for multiple settings in advanced ovarian cancer, metastatic breast cancer with BRCA mutations, metastatic pancreatic cancer, and mCRPC with HRR gene mutations, including BRCA mutations. Rucaparib is approved for maintenance treatment of recurrent ovarian cancer with BRCA mutations and mCRPC with BRCA mutations. Niraparib is approved for maintenance treatment of advanced ovarian cancer and recurrent ovarian cancer with BRCA mutations. Talazoparib is approved for locally advanced or metastatic breast cancer with BRCA mutations.

According to the NCCN guidelines, rucaparib is recommended as recurrence therapy for patients with BRCA-mutated ovarian cancer who have been treated with multiple lines of chemotherapy. It is also recommended as maintenance therapy for patients with metastatic pancreatic cancer who have undergone prior platinum-based therapy and harbor germline or somatic BRCA mutations. Additionally, niraparib is recommended as maintenance therapy for ovarian cancer patients with BRCA mutations.

KRAS G12R

Biological Impact

The V-Ki-Ras2 Kirsten Rat Sarcoma 2 Viral Oncogene Homolog (KRAS) gene encodes a small GTPase protein, a member of the RAS family of small GTPases, which catalyze the hydrolysis of GTP to GDP. RAS proteins cycle between an active (GTP-bound) and an inactive (GDP-bound) state, to activate the downstream oncogenic pathways, including the PI3K/AKT/mTOR and MAPK pathways^[5]. KRAS mutations occur primarily in three hotspots G12, G13 and Q61, and less frequently in codon A146^{[5][6]}. These are activating mutations that lead to constitutive activation and persistent stimulation of the downstream signaling pathways^{[7][8]}. Mutations in KRAS have been reported in a diverse spectrum of human malignancies, including pancreatic carcinomas (>80%)^{[5][9]}, colon carcinomas (40-50%)^{[10][11]}, and lung carcinomas (30-50%)^{[12][13]}, but are also present in biliary tract malignancies, endometrial cancer, cervical cancer, bladder cancer, liver cancer, myeloid leukemia and breast cancer^[6].

KRAS G12R is a hotspot mutation that lies within a GTP binding region of the KRAS protein (UniProtKB). G12R results in decreased KRAS GTPase activity and increased activation of downstream signaling in vitro^{[14][15]}.

Therapeutic and prognostic relevance

Cetuximab and panitumumab are FDA-approved for treating RAS wild-type metastatic colorectal cancer. The NCCN for CRC recommends cetuximab and panitumumab use only if both KRAS and NRAS genes are normal.

ACT Onco[®] + Report

KRAS mutation has been determined as an inclusion criterion for the trials evaluating MEK inhibitors efficacies in various types of solid tumors (NCT03704688, NCT02399943, NCT02285439, NCT03637491, NCT04214418).

KRAS mutations are associated with a lack of efficacy of EGFR TKIs^{[16][17][18]}. Some case reports suggest that MEK inhibitors may benefit patients with KRAS mutations, as shown in cervical and ovarian cancer cases (Am J Clin Exp Obstet Gynecol 2015;2(3):140-143)^{[19][20]}. However, a randomized Phase II study did not find trametinib to be superior to docetaxel in KRAS-mutant non-small cell lung cancer patients^[21]. MEK inhibitors as a monotherapy have limited response^[22].

Combining MEK and mTOR inhibitors is being evaluated as a potential strategy in RAS-mutant CRC^{[23][24]}. The combination of trametinib and palbociclib has resulted in objective responses in KRAS mutant models^[25].

Sorafenib has been shown to be beneficial in KRAS-mutant CRC/NSCLC, and KRAS-amplified melanoma^{[26][27][28]}. KRAS mutations in exon 2 (codon 12 or 13) and codon 61 have been associated with poor prognosis in CRC^[29].

Patients with KRAS or BRAF mutations in low-grade serous carcinoma of the ovary or peritoneum had better overall survival than those with wild-type genes^[30]. In ovarian serous borderline tumor, KRAS G12V mutation was linked to shorter survival time^[31].

In 242 patients with unresectable pancreatic cancer, the presence of KRAS G12R (N=17) or G12D (N=92) was associated with poorer prognosis (overall survival HR 1.6; 95% confidence interval 1.11-2.28)^[32].

In a Phase II trial, a patient with Erdheim-Chester disease harboring KRAS G12R and ARAF P216A achieved a complete response to cobimetinib treatment^[33]. An exploratory study demonstrated that pancreatic cancer patients with KRAS G12R mutation had better response to the combination of gemcitabine and cobimetinib than patients with KRAS G12D and G12V. One of six patients with KRAS G12R had partial response and five had stable disease and median PFS was 6 months^[34].

SERPINB3 T155R

Biological Impact

SERPINB3 encodes a protein of the serpin family of serine protease inhibitors. SERPINB3 is a close human homolog of SERPINB4 with which shares 92% protein sequence identity. SERPINB3 and SERPINB4 proteins have overlapping functions and are involved in both oncogenesis and immunity^{[35][36]}.

Therapeutic and prognostic relevance

Results from a clinical study showed that somatic mutations in SERPINB3 and SERPINB4 predicted improved survival from treatment with anti-CTLA4 therapy in two independent cohorts of patients with melanoma (n=174)^[37].

TP53 P278S, Heterozygous deletion

Biological Impact

TP53 encodes the p53 protein, a crucial tumor suppressor that orchestrates essential cellular processes including cell cycle arrest, senescence and apoptosis^[38]. TP53 is a proto-typical haploinsufficient gene, such that loss of a single copy of TP53 can result in tumor formation^[39].

P278S is a missense mutation located in the DNA binding domain (DBD) of the p53 protein (UniProtKB)^[40]. This mutation results in a loss of p53 transactivation activity in vitro, and promotes tumor growth in xenograft models^[41]. Loss of the second wild-type allele resulted in the biallelic inactivation of the gene.

ACT Onco[®] + Report

Therapeutic and prognostic relevance

Despite having a high mutation rate in cancers, there are currently no approved targeted therapies for TP53 mutations. A phase II trial demonstrated that Wee1 inhibitor (AZD1775) in combination with carboplatin was well tolerated and showed promising anti-tumor activity in TP53-mutated ovarian cancer refractory or resistant (< 3 months) to standard first-line therapy (NCT01164995)^[42].

In a retrospective study (n=19), advanced sarcoma patients with TP53 loss-of-function mutations displayed improved progression-free survival (208 days versus 136 days) relative to patients with wild-type TP53 when treated with pazopanib^[43]. Results from another Phase I trial of advanced solid tumors (n=78) demonstrated that TP53 hotspot mutations are associated with better clinical response to the combination of pazopanib and vorinostat^[44].

Advanced solid tumor and colorectal cancer patients harboring a TP53 mutation have been shown to be more sensitive to bevacizumab when compared with patients harboring wild-type TP53^{[45][46][47]}. In a pilot trial (n=21), TP53-negative breast cancer patients demonstrated increased survival following treatment with bevacizumab in combination with chemotherapy agents, Adriamycin (doxorubicin) and Taxotere (docetaxel)^[48]. TP53 mutations were correlated with poor survival of advanced breast cancer patients receiving tamoxifen or primary chemotherapy^{[49][50]}. In a retrospective study of non-small cell lung cancer (NSCLC), TP53 mutations were associated with high expression of VEGF-A, the primary target of bevacizumab, offering a mechanistic explanation for why patients exhibit improved outcomes after bevacizumab treatment when their tumors harbor mutant TP53 versus wild-type TP53^[51].

CDKN2A Homozygous deletion

Biological Impact

The Cyclin-Dependent Kinase Inhibitor 2A (CDKN2A) gene encodes the p16 (p16INK4a) and p14 (ARF) proteins. p16INK4a binds to CDK4 and CDK6, inhibiting these CDKs from binding D-type cyclins and phosphorylating the retinoblastoma (RB) protein whereas p14 (ARF) blocks the oncogenic activity of MDM2 by inhibiting MDM2-induced degradation of p53^{[52][53][54]}. CDKN2A has been reported as a haploinsufficient tumor suppressor with one copy loss that may lead to weak protein expression and is insufficient to execute its original physiological functions^[55]. Loss of CDKN2A has been frequently found in human tumors that result in uncontrolled cell proliferation^{[56][57]}.

Therapeutic and prognostic relevance

Intact p16-Cdk4-Rb axis is known to be associated with sensitivity to cyclin-dependent kinase inhibitors^{[58][59]}. Several case reports also revealed that patients with CDKN2A-deleted tumors respond to the CDK4/6-specific inhibitor treatments^{[60][61][62]}. However, there are clinical studies that demonstrated CDKN2A nuclear expression, CDKN2A/CDKN2B co-deletion, or CDKN2A inactivating mutation was not associated with clinical benefit from CDK4/6 inhibitors, such as palbociclib and ribociclib, in RB-positive patients^{[63][64][65]}. CDKN2A loss or mutation has been determined as an inclusion criterion for the trial evaluating CDK4/6 inhibitors efficacy in different types of solid tumors (NCT02693535, NCT02187783).

The phase II TAPUR trial demonstrated clinical benefits to palbociclib monotherapy in advanced NSCLC or head and neck cancer harboring a CDKN2A mutation or copy number loss. However, pancreatic and biliary cancer patients harboring a CDKN2A mutation or copy number loss did not demonstrate an objective response or stable disease when treated with palbociclib monotherapy for 16 weeks (DOI: 10.1200/JCO.2021.39.15_suppl.6043)^{[66][67]}.

Notably, the addition of several CDK4/6 inhibitors to hormone therapies, including palbociclib in combination with letrozole, ribociclib plus letrozole, and abemaciclib combines with fulvestrant, have been approved by the U.S. FDA for the treatment of ER+ and HER2- breast cancer^{[59][68][69]}.

In a Phase I trial, a KRAS wild-type squamous non-small cell lung cancer (NSCLC) patient with CDKN2A loss had a partial response when treated with CDK4/6 inhibitor abemaciclib^[61]. Administration of combined palbociclib and MEK

ACT Onco[®] + Report

inhibitor PD-0325901 yield promising progression-free survival among patients with KRAS mutant non-small cell lung cancer (NSCLC) (AACR 2017, Abstract CT046). Moreover, MEK inhibitor in combination with CDK4/6 inhibitor demonstrates significant anti-KRAS-mutant NSCLC activity and radiosensitizing effect in preclinical models^[70].

A retrospective analysis demonstrated that concurrent deletion of CDKN2A with EGFR mutation in patients with non-small cell lung cancer (NSCLC), predicts worse overall survival after EGFR-TKI treatment^[71].

FBXW7 Heterozygous deletion

Biological Impact

The F-box/WD repeat-containing protein 7 (FBXW7) gene encodes a protein that belongs to the SCF (SKP1-CUL1-F-box protein) E3 ligase complex. FBXW7 is recognized as a tumor suppressor which is involved in the negative regulation of oncogenes such as c-Myc^{[72][73]}, c-Jun^[74], cyclin E^[75], Notch family members^{[76][77]}, Aurora-A^[78], mTOR^[79], KLF5^[80], and MCL-1^[81]. Inactivating FBXW7 mutation or copy number loss may result in the accumulation of oncoproteins and therefore lead to malignant transformation^[82]. FBXW7 is a haploinsufficient tumor suppressor gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions^{[80][81][83]}.

Therapeutic and prognostic relevance

Clinical efficacy of mTOR inhibitors was seen in patients harboring aberrations in the FBXW7 gene (one patient with refractory fibrolamellar hepatocellular carcinoma, and one patient with lung adenocarcinoma)^{[84][85]}. Moreover, in vitro assay also suggested that loss or inactivation of FBXW7 may confer sensitivity to mTOR inhibitor^[79].

Preclinical studies suggested that mutations or loss of FBXW7 were associated with regorafenib and oxaliplatin resistance in CRC cell lines and gefitinib resistance in lung cancer cells^{[86][87][88][89]}.

Retrospective studies have indicated that a relatively low expression level of FBXW7 is an independent prognostic marker of poor survival for patients with hepatocellular carcinoma, lung adenocarcinoma and squamous cell carcinoma^{[90][88]}.

PTEN Heterozygous deletion

Biological Impact

The phosphatase and tensin homolog deleted on chromosome ten (PTEN) gene encodes a lipid/protein phosphatase that is important for the regulation of cell proliferation, survival, homologous recombination and maintenance of genomic integrity^{[91][92]}. PTEN acts as an essential tumor suppressor by antagonizing the PI3K/AKT/mTOR signaling pathway^[93]. PTEN is a haploinsufficient tumor suppressor gene, in which having only one copy of the wild-type allele does not produce enough protein product to execute wild-type functions^{[39][94][95]}. Germline loss-of-function PTEN mutations are found in approximately 80% of patients with Cowden syndrome, a disorder that is associated with high-penetrance breast and thyroid cancer^{[96][97][98]}. Somatic mutations or monoallelic loss of PTEN is regularly observed in a significant fraction of human cancers, including sporadic breast cancer, colon cancer, endometrial cancer, prostate cancer, and glioblastoma^{[99][100][101][102][103]}.

Therapeutic and prognostic relevance

Somatic loss of PTEN results in aberrant activation of PI3K/AKT/mTOR signaling pathway and provides a mechanistic rationale for PI3K pathway inhibitors treatment^{[104][105]}. Preclinical studies demonstrated that PTEN deficiency was associated with increased sensitivity to PI3K pathway inhibitors in selected cancer subtypes^{[106][107][108][109][110][111]}. Although early clinical data indicated that PTEN loss was associated with improved response and survival in solid tumor patients treated with mTORC1 inhibitor, everolimus^{[112][113][114]}, several phase II trials showed no clinical benefit of everolimus or temsirolimus treatment in patients with advanced solid tumors harboring PTEN loss^{[115][116][117]}.

ACT Onco[®] + Report

Several groups found that PTEN loss was generally associated with poor response to trastuzumab therapy, whether this agent was administered in the neoadjuvant, adjuvant, or metastatic settings^{[118][119][120][121][122]}. Also, loss of PTEN expression in advanced colorectal cancer (CRC) has been linked with resistance to anti-EGFR mAbs like cetuximab and panitumumab^{[123][124][125][126][127][128]}. Preclinical studies showed that loss of PTEN expression in EGFR mutant cells was associated with decreased sensitivity to EGFR TKIs, erlotinib and gefitinib^{[129][130]}. Inhibition of the PI3K/AKT/mTOR signal pathway has been shown to be an effective strategy to radiosensitize NSCLC cells harboring the EGFR activating mutation that acquires resistance to both TKIs due to PTEN loss or inactivation mutations^[131]. Loss or biallelic inactivation of PTEN is associated with resistance to anti-PD-1 checkpoint blockade therapies, including pembrolizumab and nivolumab in melanoma and leiomyosarcoma patients^{[132][133][134]}.

PTEN loss of function mutation has been determined as an inclusion criterion for the trial evaluating olaparib efficacy in metastatic biliary tract cancer (NCT04042831); talazoparib efficacy in HER2-negative solid tumors (NCT02401347); rucaparib efficacy in prostate cancer (NCT02952534, NCT03533946), and niraparib efficacy in breast cancer (NCT04508803) or any malignancy (except prostate) cancer (NCT03207347). Clinical data also suggested that PTEN deficient cancers may be sensitive to olaparib^[135]. However, in a phase II trial (NCT02286687), 13 patients with advanced solid tumors harboring PTEN mutation or loss (by IHC) had limited response to talazoparib treatment; only one patient with PTEN mutation had prolonged SD (Mol Cancer Ther 2018;17(1 Suppl):Abstract nr A096; NCT02286687). Besides, in a phase I trial (NCT00749502), no association between loss of PTEN expression and the efficacy of niraparib was identified in patients with castration-resistant prostate cancer^[136].

In a preclinical study, PTEN null cancer cells were sensitive to rucaparib treatment in vitro^[137].

SMAD4 Heterozygous deletion

Biological Impact

The SMAD family member 4 (SMAD4) gene encodes a transcription factor that acts as a downstream effector in the TGF- β signaling pathway. Upon phosphorylated and activated by serine-threonine receptor kinase, Smad4 is the Co-Smad which recruits other activated R-Smad proteins to the Smad transcriptional complex and regulate TGF- β -targeted genes^[138]. Smad4 has been identified as a haploinsufficient gene with one copy loss may lead to a weak protein expression and is insufficient to execute its original physiological function^[139]. SMAD4 germline mutations are associated with juvenile polyposis syndrome (JPS)^{[140][141][142][143]}. Somatic mutations of SMAD4 are commonly observed in pancreatic cancer^[144], colorectal cancer (CRC)^{[142][145][146]}, and less frequently seen in other cancers such as lung adenocarcinoma^[147], head and neck cancer^{[148][149]}, and cutaneous squamous cell carcinoma^[150].

Therapeutic and prognostic relevance

In Chinese patients with metastatic colorectal cancer, SMAD4 or NF1 mutations are suggested as a potential biomarker for poor prognosis to cetuximab-based therapy^[151]. Preclinical data demonstrated that depletion of SMAD4 by shRNA knockdown increased clonogenic survival and cetuximab resistance in HPV-negative head and neck squamous cell carcinoma cells^[152].

SMAD4 is also suggested as a predictive marker for 5-fluorouracil-based chemotherapy in colorectal cancer (CRC)^{[153][154]}. CRC patients with normal SMAD4 diploidy exhibited three-fold higher benefit of 5-FU/mitomycin-based adjuvant therapy when compared with those with SMAD4 deletion^[155].

Results from clinical and meta-analyses showed that loss of SMAD4 in CRC, pancreatic cancer was correlated with poor prognosis^{[156][157][158][159][160][161][162][163]}. In cervical cancer patients, weak cytoplasmic SMAD4 expression and absent nuclear SMAD4 expression were shown to be significantly associated with poor disease-free and overall 5-year survival^[164].

ACT Onco[®] + Report

US FDA-APPROVED DRUG(S)

Everolimus (AFINITOR)

Everolimus, a derivative of sirolimus, works as an inhibitor of mammalian target of rapamycin complex 1 (mTORC1) and blocks mTORC1-mediated downstream signals for cell growth, proliferation, and survival. Everolimus is developed and marketed by Novartis under the trade name AFINITOR.

- FDA Approval Summary of Everolimus (AFINITOR)

RADIANT-4 ^[165] NCT01524783	Lung or gastrointestinal neuroendocrine tumor (Approved on 2016/02/26)
	- Everolimus vs. Placebo [PFS(M): 11 vs. 3.9]
BOLERO-2 ^[166] NCT00863655	Breast cancer (Approved on 2012/07/20)
	ER+/HER2- Everolimus + exemestane vs. Placebo + exemestane [PFS(M): 7.8 vs. 3.2]
EXIST-2 NCT00790400	Tuberous sclerosis complex (tsc)-associated renal angiomyolipoma (Approved on 2012/04/26)
	- Everolimus vs. Placebo [ORR(%): 41.8 vs. 0]
RADIANT-3 ^[113] NCT00510068	Pancreatic neuroendocrine tumor (Approved on 2011/05/05)
	- Everolimus vs. Placebo [PFS(M): 11 vs. 4.6]
EXIST-1 ^[167] NCT00789828	Subependymal giant cell astrocytoma (Approved on 2010/10/29)
	- Everolimus vs. Placebo [ORR(%): 35.0]
RECORD-1 ^[168] NCT00410124	Renal cell carcinoma (Approved on 2009/05/30)
	- Everolimus vs. Placebo [PFS(M): 4.9 vs. 1.9]

Niraparib (ZEJULA)

Niraparib is an oral, small molecule inhibitor of the DNA repair enzyme poly (ADP-ribose) polymerase-1 and -2 (PARP-1, -2). Niraparib is developed and marketed by Tesaro under the trade name ZEJULA.

- FDA Approval Summary of Niraparib (ZEJULA)

PRIMA NCT02655016	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2020/04/29)
	- Niraparib vs. Placebo [PFS (overall population)(M): 13.8 vs. 8.2]
NOVA ^[169] NCT01847274	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/03/27)
	- Niraparib vs. Placebo [PFS (overall population)(M): 11.3 vs. 4.7]

ACT Onco® + Report

Olaparib (LYNPARZA)

Olaparib is an oral, small molecule inhibitor of poly (ADP-ribose) polymerase-1, -2, and -3 (PARP-1, -2, -3). Olaparib is developed by KuDOS Pharmaceuticals and marketed by AstraZeneca under the trade name LYNPARZA.

- FDA Approval Summary of Olaparib (LYNPARZA)

OlympiA NCT02032823	Her2-negative high-risk early breast cancer (Approved on 2022/03/11)
	HER2-/gBRCA mutation Olaparib vs. Placebo [invasive disease-free survival (IDFS)(M):]
PROfound ^[170] NCT02987543	Prostate cancer (Approved on 2020/05/19)
	HRR genes mutation Olaparib vs. Enzalutamide or abiraterone acetate [PFS(M): 5.8 vs. 3.5]
PAOLA-1 ^[171] NCT02477644	Ovarian cancer (Approved on 2020/05/08)
	HRD+ Olaparib + bevacizumab vs. Placebo + bevacizumab [PFS(M): 37.2 vs. 17.7]
POLO ^[172] NCT02184195	Pancreatic adenocarcinoma (Approved on 2019/12/27)
	gBRCA mutation Olaparib vs. Placebo [ORR(%): 23.0 vs. 12.0, PFS(M): 7.4 vs. 3.8]
SOLO-1 ^[173] NCT01844986	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2018/12/19)
	gBRCA mutation or sBRCA mutation Olaparib vs. Placebo [PFS(M): NR vs. 13.8]
OlympiAD ^[174] NCT02000622	Breast cancer (Approved on 2018/02/06)
	HER2-/gBRCA mutation Olaparib vs. Chemotherapy [PFS(M): 7 vs. 4.2]
SOLO-2/ENGOT-Ov21 ^[175] NCT01874353	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/08/17)
	gBRCA mutation Olaparib vs. Placebo [PFS(M): 19.1 vs. 5.5]
Study19 ^[176] NCT00753545	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/08/17)
	- Olaparib vs. Placebo [PFS(M): 8.4 vs. 4.8]

Rucaparib (RUBRACA)

Rucaparib is an inhibitor of the DNA repair enzyme poly (ADP-ribose) polymerase-1, -2 and -3 (PARP-1, -2, -3). Rucaparib is developed and marketed by Clovis Oncology under the trade name RUBRACA.

- FDA Approval Summary of Rucaparib (RUBRACA)

TRITON2 NCT02952534	Prostate cancer (Approved on 2020/05/15)
	gBRCA mutation or sBRCA mutation Rucaparib [ORR(%): 44.0, DOR(M): NE]
ARIEL3 ^[177] NCT01968213	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2018/04/06)
	- Rucaparib vs. Placebo [PFS (All)(M): 10.8 vs. 5.4, PFS (HRD)(M): 13.6 vs. 5.4, PFS (tBRCA)(M): 16.6 vs. 5.4]

ACT Onco[®] + Report

Talazoparib (TALZENNA)

Talazoparib is an inhibitor of poly (ADP-ribose) polymerase (PARP) enzymes, including PARP1 and PARP2. Talazoparib is developed and marketed by Pfizer under the trade name TALZENNA.

- FDA Approval Summary of Talazoparib (TALZENNA)

EMBRACA ^[178] NCT01945775	Breast cancer (Approved on 2018/10/16)
	HER2-/gBRCA mutation
	Talazoparib vs. Chemotherapy [PFS(M): 8.6 vs. 5.6]

Temsirolimus (TORISEL)

Temsirolimus is a soluble ester of sirolimus (rapamycin, brand-name drug Rapamune) and functions as an inhibitor of mammalian target of rapamycin complex (mTORC). The inhibitory molecular mechanism is similar to Everolimus. Temsirolimus is developed by Wyeth Pharmaceuticals and marketed by Pfizer under the trade name TORISEL.

- FDA Approval Summary of Temsirolimus (TORISEL)

[179] NCT00065468	Renal cell carcinoma (Approved on 2007/05/30)
	-
	Temsirolimus vs. Ifn-α [OS(M): 10.9 vs. 7.3]

D=day; W=week; M=month

ACT Onco[®] + Report

ONGOING CLINICAL TRIALS

Trials were searched by applying filters: study status, patient's diagnosis, intervention, location and/or biomarker(s). Please visit <https://clinicaltrials.gov> to search and view for a complete list of open available and updated matched trials.

No trial has been found.

ACT Onco[®] + Report


SUPPLEMENTARY INFORMATION OF TESTING RESULTS DETAILED INFORMATION OF VARIANTS WITH CLINICAL RELEVANCE

- Single Nucleotide and Small InDel Variants

Gene	Amino Acid Change	Exon	cDNA Change	Accession Number	COSMIC ID	Allele Frequency	Coverage
BRCA2	S1882*	11	c.5645C>A	NM_000059	COSM78887	59.5%	440
KRAS	G12R	2	c.34G>C	NM_004985	COSM518	18.7%	1792
SERPINB3	T155R	5	c.464C>G	NM_006919	-	35.8%	773
TP53	P278S	8	c.832C>T	NM_000546	COSM10939	22.3%	512

- Copy Number Alterations

Observed copy number (CN) for each evaluated position is shown on the y-axis. Regions referred to as amplification or deletion are shown in color. Regions without significant changes are represented in gray.



ACT Onco[®] + Report

OTHER DETECTED VARIANTS

Gene	Amino Acid Change	Exon	cDNA Change	Accession Number	COSMIC ID	Allele Frequency	Coverage
ATR	G1342V	22	c.4025G>T	NM_001184	COSM6930215	43.9%	660
CALR	P243L	6	c.728C>T	NM_004343	-	47.4%	1087
CARD11	S694L	16	c.2081C>T	NM_032415	COSM5505215	52.6%	565
EPCAM	R153T	4	c.458G>C	NM_002354	-	31.2%	791
ERCC4	A109V	2	c.326C>T	NM_005236	COSM2143614	49.6%	1021
ERCC5	D703G	9	c.2108A>G	NM_000123	-	68.3%	1076
FGF6	R188W	3	c.562C>T	NM_020996	-	17.1%	1234
HIST1H1C	L82F	1	c.244C>T	NM_005319	COSM6942690	52.1%	946
JAK1	N973K	21	c.2919C>A	NM_002227	-	35.5%	1214
LRP1B	A811D	15	c.2432C>A	NM_018557	-	14.2%	1736
MED12	T344N	7	c.1031C>A	NM_005120	COSM6302051	99.5%	427
MUC16	T9972I	3	c.29915C>T	NM_024690	-	51.8%	737
PDCD1 (PD-1)	Splice region	-	c.77-3C>G	NM_005018	-	16.4%	494
PIK3C3	Splice region	-	c.2523+6T>A	NM_002647	-	54.0%	311
PIK3C3	P714A	20	c.2140C>G	NM_002647	-	21.2%	623
RECQL4	Splice region	-	c.2200+7C>G	NM_004260	-	10.9%	496


Note:

- This table enlists variants detected by the panel other than those with clinical relevance (reported in Testing Result section). The clinical impact of a genetic variant is determined according to ACT Genomics in-house clinical knowledge database. A negative result does not necessarily indicate absence of biological effect on the tumor. Some variants listed here may possibly have preclinical data or may show potential clinical relevance in the future.

ACT Onco[®] + Report

TEST DETAILS

SPECIMEN RECEIVED AND PATHOLOGY REVIEW



- Collection date: May 10, 2023
- Facility retrieved: 臺北榮總
- H&E-stained section No.: S11291425A
- Collection site: Pancreas
- Examined by: Dr. Yeh-Han Wang
- 1. The percentage of viable tumor cells in total cells in the whole slide (%): 45%
- 2. The percentage of viable tumor cells in total cells in the encircled areas in the whole slide (%): 45%
- 3. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the whole slide (%): 0%
- 4. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the encircled areas in the whole slide (%): 0%
- 5. Additional comment: NA
- Manual macrodissection: Not performed
- The outline highlights the area of malignant neoplasm annotated by a pathologist.

RUN QC

- Panel: ACTOnco[®]+

DNA test

- Mean Depth: 763x
- Target Base Coverage at 100x: 95%

RNA test

- Average unique RNA Start Sites per control GSP2: 126

ACT Onco[®] + Report

LIMITATIONS

1. This test does not provide information of variant causality and does not detect variants in non-coding regions that could affect gene expression. This report does not report polymorphisms and we do not classify whether a mutation is germline or somatic. Variants identified by this assay were not subject to validation by Sanger or other technologies.
2. The possibility cannot be excluded that certain pathogenic variants detected by other sequencing tools may not be reported in the test because of technical limitation of bioinformatics algorithm or the NGS sequencing platform, e.g. low coverage.
3. This test has been designed to detect fusions in 13 genes sequenced. Therefore, fusion in genes not covered by this test would not be reported. For novel fusions detected in this test, Sanger sequencing confirmation is recommended if residue specimen is available.

NEXT-GENERATION SEQUENCING (NGS) METHODS

DNA test

Extracted genomic DNA was amplified using primers targeting coding exons of analyzed genes and subjected to library construction. Barcoded libraries were subsequently conjugated with sequencing beads by emulsion PCR and enriched using Ion Chef system. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific).

Raw reads generated by the sequencer were mapped to the hg19 reference genome using the Ion Torrent Suite. Coverage depth was calculated using Torrent Coverage Analysis plug-in. Single nucleotide variants (SNVs) and short insertions/deletions (InDels) were identified using the Torrent Variant Caller plug-in. VEP (Variant Effect Predictor) was used to annotate every variant using databases from Clinvar, COSMIC and Genome Aggregation database. Variants with coverage ≥ 20 , allele frequency $\geq 5\%$ and actionable variants with allele frequency $\geq 2\%$ were retained. This test provides uniform coverage of the targeted regions, enabling target base coverage at $100\times \geq 85\%$ with a mean coverage $\geq 500\times$.

Variants reported in Genome Aggregation database with $> 1\%$ minor allele frequency (MAF) were considered as polymorphisms. ACT Genomics in-house database was used to determine technical errors. Clinically actionable and biologically significant variants were determined based on the published medical literature.

The copy number alterations (CNAs) were predicted as described below:

Amplicons with read counts in the lowest 5th percentile of all detectable amplicons and amplicons with a coefficient of variation ≥ 0.3 were removed. The remaining amplicons were normalized to correct the pool design bias. ONCOCNV (an established method for calculating copy number aberrations in amplicon sequencing data by Boeva et al., 2014) was applied for the normalization of total amplicon number, amplicon GC content, amplicon length, and technology-related biases, followed by segmenting the sample with a gene-aware model. The method was used as well for establishing the baseline of copy number variations.

Tumor mutational burden (TMB) was calculated by using the sequenced regions of ACTOnco[®] to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The TMB calculation predicted somatic variants and applied a machine learning model with a cancer hotspot correction. TMB may be reported as "TMB-High", "TMB-Low" or "Cannot Be Determined". TMB-High corresponds to ≥ 7.5 mutations per megabase (Muts/Mb); TMB-Low corresponds to < 7.5 Muts/Mb. TMB is reported as "Cannot Be Determined" if the tumor purity of the sample is $< 30\%$.

Classification of microsatellite instability (MSI) status is determined by a machine learning prediction algorithm. The change of a number of repeats of different lengths from a pooled microsatellite stable (MSS) baseline in > 400 genomic loci are used as the features for the algorithm. The final output of the results is either microsatellite Stable (MSS) or microsatellite instability high (MSI-H).

ACT Onco[®] + Report

RNA test

Extracted RNA was reverse-transcribed and subjected to library construction. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific). To ensure sequencing quality for fusion variant analysis, the average unique RNA Start Sites (SS) per control Gene Specific Primer 2 (GSP 2) should be ≥ 10 .

The fusion analysis pipeline aligned sequenced reads to the human reference genome, identified regions that map to noncontiguous regions of the genome, applied filters to exclude probable false-positive events and, annotated previously characterized fusion events according to Quiver Gene Fusion Database, a curated database owned and maintained by ArcherDX. In general, samples with detectable fusions need to meet the following criteria: (1) Number of unique start sites (SS) for the GSP2 ≥ 3 ; (2) Number of supporting reads spanning the fusion junction ≥ 5 ; (3) Percentage of supporting reads spanning the fusion junction $\geq 10\%$; (4) Fusions annotated in Quiver Gene Fusion Database.

DATABASE USED

- Reference genome: Human genome sequence hg19
- COSMIC v.92
- Genome Aggregation database r2.1.1
- ClinVar (version 20210404)
- ACT Genomics in-house database
- Quiver Gene Fusion Database version 5.1.18

Variant Analysis:

醫檢師張筑芃 博士
Chu-Yuan Chang Ph.D.
檢字第 020115 號



Sign Off

解剖病理專科醫師王業翰
Yeh-Han Wang M.D.
病解字第 000545 號



ACT Onco[®] + Report

GENE LIST SNV & CNV

ABCB1*	ABCC2*	ABCG2*	ABL1	ABL2	ADAMTS1	ADAMTS13	ADAMTS15	ADAMTS16	ADAMTS18	ADAMTS6	ADAMTS9
ADAMTSL1	ADGRA2	ADH1C*	AKT1	AKT2	AKT3	ALDH1A1*	ALK	AMER1	APC	AR	ARAF
ARID1A	ARID1B	ARID2	ASXL1	ATM	ATR	ATRX	AURKA	AURKB	AXIN1	AXIN2	AXL
B2M	BAP1	BARD1	BCL10	BCL2*	BCL2L1	BCL2L2*	BCL6	BCL9	BCOR	BIRC2	BIRC3
BLM	BMPR1A	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2*	BTB	BUB1B	CALR
CANX	CARD11	CASP8	CBFB	CBL	CCNA1	CCNA	CCNB1	CCNB2	CCNB3	CCND1	CCND2
CCND3	CCNE1	CCNE2	CCNH	CD19	CD274	CD58	CD70*	CD79A	CD79B	CDC73	CDH1
CDK1	CDK12	CDK2	CDK4	CDK5	CDK6	CDK7	CDK8	CDK9	CDKN1A	CDKN1B	CDKN2A
CDKN2B	CDKN2C	CEBPA*	CHEK1	CHEK2	CIC	CREBBP	CRKL	CRLF2	CSF1R	CTCF	CTLA4
CTNNA1	CTNNB1	CUL3	CYLD	CYP1A1*	CYP2B6*	CYP2C19*	CYP2C8*	CYP2D6	CYP2E1*	CYP3A4*	CYP3A5*
DAXX	DCUN1D1	DDR2	DICER1	DNMT3A	DOT1L	DPYD	DTX1	E2F3	EGFR	EP300	EPCAM
EPHA2	EPHA3	EPHA5	EPHA7	EPHB1	ERBB2	ERBB3	ERBB4	ERCC1	ERCC2	ERCC3	ERCC4
ERCC5	ERG	ESR1	ESR2	ETV1	ETV4	EZH2	FAM46C	FANCA	FANCC	FANCD2	FANCE
FANCF	FANCG	FANCL	FAS	FAT1	FBXW7	FCGR2B	FGF1*	FGF10	FGF14	FGF19*	FGF23
FGF3	FGF4*	FGF6	FGFR1	FGFR2	FGFR3	FGFR4	FH	FLCN	FLT1	FLT3	FLT4
FOXL2*	FOXP1	FRG1	FUBP1	GATA1	GATA2	GATA3	GNA11	GNA13	GNAQ	GNAS	GREM1
GRIN2A	GSK3B	GSTP1*	GSTT1*	HGF	HIF1A	HIST1H1C*	HIST1H1E*	HNF1A	HR	HRAS*	HSP90AA1
HSP90AB1	HSPA4	HSPA5	IDH1	IDH2	IFNL3*	IGF1	IGF1R	IGF2	IKBKB	IKBKE	IKZF1
IL6	IL7R	INPP4B	INSR	IRF4	IRS1	IRS2*	JAK1	JAK2	JAK3	JUN*	KAT6A
KDM5A	KDM5C	KDM6A	KDR	KEAP1	KIT	KMT2A	KMT2C	KMT2D	KRAS	LCK	LIG1
LIG3	LMO1	LRP1B	LYN	MALT1	MAP2K1	MAP2K2	MAP2K4	MAP3K1	MAP3K7	MAPK1	MAPK3
MAX	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1	MET	MITF	MLH1	MPL	MRE11
MSH2	MSH6	MTHFR*	MTOR	MUC16	MUC4	MUC6	MUTYH	MYC	MYCL	MYCN	MYD88
NAT2*	NBN	NEFH	NF1	NF2	NFE2L2	NFKB1	NFKBIA	NKX2-1*	NOTCH1	NOTCH2	NOTCH3
NOTCH4	NPM1	NQO1*	NRAS	NSD1	NTRK1	NTRK2	NTRK3	PAK3	PALB2	PARP1	PAX5
PAX8	PBRM1	PDCD1	PDCD1LG2	PDGFRA	PDGFRB	PDIA3	PGF	PHOX2B*	PIK3C2B	PIK3C2G	PIK3C3
PIK3CA	PIK3CB	PIK3CD	PIK3CG	PIK3R1	PIK3R2	PIK3R3	PIM1	PMS1	PMS2	POLB	POLD1
POLE	PPARG	PPP2R1A	PRDM1	PRKAR1A	PRKCA	PRKCB	PRKCG	PRKCI	PRKCQ	PRKDC	PRKN
PSMB8	PSMB9	PSME1	PSME2	PSME3	PTCH1	PTEN	PTGS2	PTPN11	PTPRD	PTPRT	RAC1
RAD50	RAD51	RAD51B	RAD51C	RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	RECQL4
REL	RET	RHOA	RICTOR	RNF43	ROS1	RPPH1	RPTOR	RUNX1	RUNX1T1	RXRA	SDHA
SDHB	SDHC	SDHD	SERPINB3	SERPINB4	SETD2	SF3B1	SGK1	SH2D1A*	SLC19A1*	SLC22A2*	SLC01B1*
SLC01B3*	SMAD2	SMAD3	SMAD4	SMARCA4	SMARCB1	SMO	SOC1*	SOX2*	SOX9	SPEN	SPOP
SRC	STAG2	STAT3	STK11	SUFU	SYK	SYNE1	TAF1	TAP1	TAP2	TAPBP	TBX3
TEK	TERT	TET1	TET2	TGFBR2	TMSB4X*	TNF	TNFAIP3	TNFRSF14	TNFSF11	TOP1	TP53
TPMT*	TSC1	TSC2	TSHR	TYMS	U2AF1	UBE2A*	UBE2K	UBR5	UGT1A1*	USH2A	VDR*
VEGFA	VEGFB	VHL	WT1	XIAP	XPO1	XRCC2	ZNF217				

*Analysis of copy number alterations NOT available.

FUSION

ALK	BRAF	EGFR	FGFR1	FGFR2	FGFR3	MET	NRG1	NTRK1	NTRK2	NTRK3	RET	ROS1
-----	------	------	-------	-------	-------	-----	------	-------	-------	-------	-----	------


ACT Onco[®] + Report

APPENDIX

POSSIBLE THERAPEUTIC IMPLICATIONS FOR HETEROZYGOUS DELETION


Gene	Therapies	Possible effect
FBXW7	Everolimus, Temsirolimus	sensitive
PTEN	Niraparib, Olaparib, Rucaparib, Talazoparib	sensitive
SMAD4	Cetuximab	resistant
PTEN	Erlotinib, Gefitinib, Panitumumab, Cetuximab, Trastuzumab	resistant
FBXW7	Gefitinib, Regorafenib	resistant

SIGNALING PATHWAYS AND MOLECULAR-TARGETED AGENTS



1: Olaparib, Niraparib, Rucaparib, Talazoparib

ACT Onco[®] + Report



1: Everolimus, Temsirolimus

ACT Onco[®] + Report

DISCLAIMER

法律聲明

本檢驗報告僅提供專業醫療參考，結果需經專業醫師解釋及判讀。基因突變資訊非必具備藥物或治療有效性指標，反之亦然。本檢驗報告提供之用藥指引不聲明或保證其臨床有效性，反之亦然。本基因檢測方法係由本公司研究開發，已經過有效性測試。

本檢驗報告非經本公司許可，不得私自變造、塗改，或以任何方式作為廣告及其他宣傳之用途。

本公司於提供檢驗報告後，即已完成本次契約義務，後續之報告解釋、判讀及用藥、治療，應自行尋求相關專業醫師協助，若需將報告移件其他醫師，本人應取得該醫師同意並填寫移件申請書，主動告知行動基因，行動基因僅能配合該醫師意願與時間提供醫師解說。

醫療決策需由醫師決定

任何治療與用藥需經由醫師在考慮病患所有健康狀況相關資訊包含健檢、其他檢測報告和病患意願後，依照該地區醫療照護標準由醫師獨立判斷。醫師不應僅依據單一報告結果(例如本檢測或本報告書內容)做決策。

基因突變與用藥資訊並非依照有效性排序

本報告中列出之生物標記變異與藥物資訊並非依照潛在治療有效性排序。

證據等級

藥物潛在臨床效益(或缺乏潛在臨床效益)的實證證據是依據至少一篇臨床療效個案報告或臨床前試驗做為評估。本公司盡力提供適時及準確之資料，但由於醫學科技之發展日新月異，本公司不就本報告提供的資料是否為準確、適宜或最新作保證。

責任

本檢驗報告僅提供專業醫療參考，本公司及其員工不對任何由使用本報告之內容引起的直接、間接、特殊、連帶或衍生的損失或損害承擔責任。

ACT Onco[®] + Report

REFERENCE

1. PMID: 11239455; 2001, Mol Cell;7(2):263-72
BRCA2 is required for homology-directed repair of chromosomal breaks.
2. PMID: 17597348; 2007, Ann Surg Oncol;14(9):2510-8
Heterogenic loss of the wild-type BRCA allele in human breast tumorigenesis.
3. PMID: 22193408; 2011, Nat Rev Cancer;12(1):68-78
BRCA1 and BRCA2: different roles in a common pathway of genome protection.
4. PMID: 27283171; 2016, J Natl Compr Canc Netw;14(6):795-806
The Relevance of Hereditary Cancer Risks to Precision Oncology: What Should Providers Consider When Conducting Tumor Genomic Profiling?
5. PMID: 2453289; 1988, Cell;53(4):549-54
Most human carcinomas of the exocrine pancreas contain mutant c-K-ras genes.
6. PMID: 2114981; 1990, Eur J Clin Invest;20(3):225-35
ras oncogenes: their role in neoplasia.
7. PMID: 20617134; 2010, J Biomed Biotechnol;2010():150960
Clinical relevance of KRAS in human cancers.
8. PMID: 21993244; 2011, Nat Rev Cancer;11(11):761-74
RAS oncogenes: weaving a tumorigenic web.
9. PMID: 3047672; 1988, Nucleic Acids Res;16(16):7773-82
KRAS codon 12 mutations occur very frequently in pancreatic adenocarcinomas.
10. PMID: 3587348; 1987, Nature;327(6120):293-7
Prevalence of ras gene mutations in human colorectal cancers.
11. PMID: 1942608; 1991, Nihon Shokakibyo Gakkai Zasshi;88(8):1539-44
[Prevalence of K-ras gene mutations in human colorectal cancers].
12. PMID: 2252272; 1990, Am Rev Respir Dis;142(6 Pt 2):S27-30
The ras oncogenes in human lung cancer.
13. PMID: 1486840; 1992, Environ Health Perspect;98():13-24
Role of proto-oncogene activation in carcinogenesis.
14. PMID: 23455880; 2013, J Cancer Res Clin Oncol;139(6):953-61
KRAS allele-specific activity of sunitinib in an isogenic disease model of colorectal cancer.
15. PMID: 26037647; 2015, Mol Cancer Res;13(9):1325-35
Biochemical and Structural Analysis of Common Cancer-Associated KRAS Mutations.
16. PMID: 18349398; 2008, J Clin Oncol;26(9):1472-8
Molecular characteristics of bronchioloalveolar carcinoma and adenocarcinoma, bronchioloalveolar carcinoma subtype, predict response to erlotinib.
17. PMID: 23401440; 2013, J Clin Oncol;31(8):1112-21
KRAS mutation: should we test for it, and does it matter?
18. PMID: 18024870; 2007, J Clin Oncol;25(33):5240-7
Prognostic and predictive importance of p53 and RAS for adjuvant chemotherapy in non small-cell lung cancer.
19. PMID: 29946554; 2018, Gynecol Oncol Rep;25():41-44

ACT Onco[®] + Report

Binimetinib (MEK162) in recurrent low-grade serous ovarian cancer resistant to chemotherapy and hormonal treatment.

20. PMID: 26075998; 2014, Gynecol Oncol Rep;10():28-9
Response to MEK inhibitor in small cell neuroendocrine carcinoma of the cervix with a KRAS mutation.
21. PMID: 25722381; 2015, Ann Oncol;26(5):894-901
A randomized phase II study of the MEK1/MEK2 inhibitor trametinib (GSK1120212) compared with docetaxel in KRAS-mutant advanced non-small-cell lung cancer (NSCLC)†.
22. PMID: 24947927; 2014, Clin Cancer Res;20(16):4251-61
Phase I expansion and pharmacodynamic study of the oral MEK inhibitor RO4987655 (CH4987655) in selected patients with advanced cancer with RAS-RAF mutations.
23. PMID: 27340376; 2016, Curr Colorectal Cancer Rep;12():141-150
Molecular Subtypes and Personalized Therapy in Metastatic Colorectal Cancer.
24. PMID: 22392911; 2012, Clin Cancer Res;18(9):2515-25
Inhibition of MEK and PI3K/mTOR suppresses tumor growth but does not cause tumor regression in patient-derived xenografts of RAS-mutant colorectal carcinomas.
25. PMID: 26369631; 2016, Clin Cancer Res;22(2):405-14
Sensitivity of KRAS-Mutant Colorectal Cancers to Combination Therapy That Cotargets MEK and CDK4/6.
26. PMID: 24407191; 2014, Br J Cancer;110(5):1148-54
Sorafenib and irinotecan (NEXIRI) as second- or later-line treatment for patients with metastatic colorectal cancer and KRAS-mutated tumours: a multicentre Phase I/II trial.
27. PMID: 23224737; 2013, Clin Cancer Res;19(3):743-51
A phase II study of sorafenib in patients with platinum-pretreated, advanced (Stage IIIB or IV) non-small cell lung cancer with a KRAS mutation.
28. PMID: 26307133; 2016, Clin Cancer Res;22(2):374-82
Copy Number Changes Are Associated with Response to Treatment with Carboplatin, Paclitaxel, and Sorafenib in Melanoma.
29. PMID: 15923428; 2005, Ann Oncol;16 Suppl 4():iv44-49
Prognostic and predictive factors in colorectal cancer: Kirsten Ras in CRC (RASCAL) and TP53CRC collaborative studies.
30. PMID: 26484411; 2015, Br J Cancer;113(9):1254-8
Impact of mutational status on survival in low-grade serous carcinoma of the ovary or peritoneum.
31. PMID: 24549645; 2013, J Pathol;231(4):449-56
KRAS (but not BRAF) mutations in ovarian serous borderline tumour are associated with recurrent low-grade serous carcinoma.
32. PMID: 22983505; 2013, J Gastroenterol;48(5):640-6
Prognostic value of K-ras mutation status and subtypes in endoscopic ultrasound-guided fine-needle aspiration specimens from patients with unresectable pancreatic cancer.
33. PMID: 30867592; 2019, Nature;567(7749):521-524
Efficacy of MEK inhibition in patients with histiocytic neoplasms.
34. PMID: 34901697; 2021, J Pancreat Cancer;7(1):65-70
Cobimetinib Plus Gemcitabine: An Active Combination in KRAS G12R-Mutated Pancreatic Ductal Adenocarcinoma Patients in Previously Treated and Failed Multiple Chemotherapies.
35. PMID: 25111616; 2015, J Invest Dermatol;135(1):160-169
SERPINB3/B4 contributes to early inflammation and barrier dysfunction in an experimental murine model of atopic dermatitis.
36. PMID: 24759783; 2014, Nat Commun;5():3729
Oncogenic Ras induces inflammatory cytokine production by upregulating the squamous cell carcinoma antigens SerpinB3/B4.
37. PMID: 27668655; 2016, Nat Genet;48(11):1327-1329

ACT Onco[®] + Report

Recurrent SERPINB3 and SERPINB4 mutations in patients who respond to anti-CTLA4 immunotherapy.

38. PMID: 24739573; 2014, Nat Rev Cancer;14(5):359-70
Unravelling mechanisms of p53-mediated tumour suppression.
39. PMID: 21125671; 2011, J Pathol;223(2):137-46
Haplo-insufficiency: a driving force in cancer.
40. PMID: 22713868; 2012, Genes Dev;26(12):1268-86
Mutant p53: one name, many proteins.
41. PMID: 22989750; 2012, Biochem Biophys Res Commun;428(1):6-10
Allele specific gain-of-function activity of p53 mutants in lung cancer cells.
42. PMID: 27998224; 2016, J Clin Oncol;34(36):4354-4361
Phase II Study of WEE1 Inhibitor AZD1775 Plus Carboplatin in Patients With TP53-Mutated Ovarian Cancer Refractory or Resistant to First-Line Therapy Within 3 Months.
43. PMID: 26646755; 2016, Ann Oncol;27(3):539-43
TP53 mutational status is predictive of pazopanib response in advanced sarcomas.
44. PMID: 25669829; 2015, Ann Oncol;26(5):1012-1018
Phase I study of pazopanib and vorinostat: a therapeutic approach for inhibiting mutant p53-mediated angiogenesis and facilitating mutant p53 degradation.
45. PMID: 27466356; 2016, Mol Cancer Ther;15(10):2475-2485
TP53 Alterations Correlate with Response to VEGF/VEGFR Inhibitors: Implications for Targeted Therapeutics.
46. PMID: 23670029; 2013, Oncotarget;4(5):705-14
P53 mutations in advanced cancers: clinical characteristics, outcomes, and correlation between progression-free survival and bevacizumab-containing therapy.
47. PMID: 17145525; 2006, Semin Oncol;33(5 Suppl 10):S8-14
Bevacizumab in combination with chemotherapy: first-line treatment of patients with metastatic colorectal cancer.
48. PMID: 21399868; 2011, Int J Oncol;38(5):1445-52
p53, HER2 and tumor cell apoptosis correlate with clinical outcome after neoadjuvant bevacizumab plus chemotherapy in breast cancer.
49. PMID: 20549698; 2011, Int J Cancer;128(8):1813-21
p53 status influences response to tamoxifen but not to fulvestrant in breast cancer cell lines.
50. PMID: 10786679; 2000, Cancer Res;60(8):2155-62
Complete sequencing of TP53 predicts poor response to systemic therapy of advanced breast cancer.
51. PMID: 25672981; 2015, Cancer Res;75(7):1187-90
VEGF-A Expression Correlates with TP53 Mutations in Non-Small Cell Lung Cancer: Implications for Antiangiogenesis Therapy.
52. PMID: 17055429; 2006, Cell;127(2):265-75
The regulation of INK4/ARF in cancer and aging.
53. PMID: 8521522; 1995, Cell;83(6):993-1000
Alternative reading frames of the INK4a tumor suppressor gene encode two unrelated proteins capable of inducing cell cycle arrest.
54. PMID: 9529249; 1998, Cell;92(6):725-34
ARF promotes MDM2 degradation and stabilizes p53: ARF-INK4a locus deletion impairs both the Rb and p53 tumor suppression pathways.
55. PMID: 16115911; 2005, Clin Cancer Res;11(16):5740-7
Comprehensive analysis of CDKN2A status in microdissected urothelial cell carcinoma reveals potential haploinsufficiency, a high frequency of homozygous co-deletion and associations with clinical phenotype.

ACT Onco[®] + Report

56. PMID: 7550353; 1995, Nat Genet;11(2):210-2
Frequency of homozygous deletion at p16/CDKN2 in primary human tumours.
57. PMID: 24089445; 2013, Clin Cancer Res;19(19):5320-8
The cell-cycle regulator CDK4: an emerging therapeutic target in melanoma.
58. PMID: 27849562; 2017, Gut;66(7):1286-1296
Palbociclib (PD-0332991), a selective CDK4/6 inhibitor, restricts tumour growth in preclinical models of hepatocellular carcinoma.
59. PMID: 25524798; 2015, Lancet Oncol;16(1):25-35
The cyclin-dependent kinase 4/6 inhibitor palbociclib in combination with letrozole versus letrozole alone as first-line treatment of oestrogen receptor-positive, HER2-negative, advanced breast cancer (PALOMA-1/TRIO-18): a randomised phase 2 study.
60. PMID: 28283584; 2017, Oncologist;22(4):416-421
Clinical Benefit in Response to Palbociclib Treatment in Refractory Uterine Leiomyosarcomas with a Common CDKN2A Alteration.
61. PMID: 27217383; 2016, Cancer Discov;6(7):740-53
Efficacy and Safety of Abemaciclib, an Inhibitor of CDK4 and CDK6, for Patients with Breast Cancer, Non-Small Cell Lung Cancer, and Other Solid Tumors.
62. PMID: 26715889; 2015, Curr Oncol;22(6):e498-501
Does CDKN2A loss predict palbociclib benefit?
63. PMID: 25501126; 2015, Clin Cancer Res;21(5):995-1001
CDK 4/6 inhibitor palbociclib (PD0332991) in Rb+ advanced breast cancer: phase II activity, safety, and predictive biomarker assessment.
64. PMID: 27542767; 2016, Clin Cancer Res;22(23):5696-5705
A Phase I Study of the Cyclin-Dependent Kinase 4/6 Inhibitor Ribociclib (LEE011) in Patients with Advanced Solid Tumors and Lymphomas.
65. PMID: 24797823; 2014, Oncologist;19(6):616-22
Enabling a genetically informed approach to cancer medicine: a retrospective evaluation of the impact of comprehensive tumor profiling using a targeted next-generation sequencing panel.
66. PMID: 35050752; 2020, JCO Precis Oncol;4():757-766
Palbociclib in Patients With Non-Small-Cell Lung Cancer With CDKN2A Alterations: Results From the Targeted Agent and Profiling Utilization Registry Study.
67. PMID: 35100714; 2019, JCO Precis Oncol;3():1-8
Palbociclib in Patients With Pancreatic and Biliary Cancer With CDKN2A Alterations: Results From the Targeted Agent and Profiling Utilization Registry Study.
68. PMID: 27717303; 2016, N Engl J Med;375(18):1738-1748
Ribociclib as First-Line Therapy for HR-Positive, Advanced Breast Cancer.
69. PMID: 28580882; 2017, J Clin Oncol;35(25):2875-2884
MONARCH 2: Abemaciclib in Combination With Fulvestrant in Women With HR+/HER2- Advanced Breast Cancer Who Had Progressed While Receiving Endocrine Therapy.
70. PMID: 26728409; 2016, Clin Cancer Res;22(1):122-33
Coadministration of Trametinib and Palbociclib Radiosensitizes KRAS-Mutant Non-Small Cell Lung Cancers In Vitro and In Vivo.
71. PMID: 31401335; 2019, Transl Oncol;12(11):1425-1431
Concomitant Genetic Alterations are Associated with Worse Clinical Outcome in EGFR Mutant NSCLC Patients Treated with Tyrosine Kinase Inhibitors.
72. PMID: 15498494; 2004, Curr Biol;14(20):1852-7
A nucleolar isoform of the Fbw7 ubiquitin ligase regulates c-Myc and cell size.
73. PMID: 15103331; 2004, EMBO J;23(10):2116-25
Phosphorylation-dependent degradation of c-Myc is mediated by the F-box protein Fbw7.

ACT Onco[®] + Report

74. PMID: 16023596; 2005, Cancer Cell;8(1):25-33
The v-Jun point mutation allows c-Jun to escape GSK3-dependent recognition and destruction by the Fbw7 ubiquitin ligase.
75. PMID: 11533444; 2001, Science;294(5540):173-7
Phosphorylation-dependent ubiquitination of cyclin E by the SCFFbw7 ubiquitin ligase.
76. PMID: 11461910; 2001, J Biol Chem;276(38):35847-53
The Notch intracellular domain is ubiquitinated and negatively regulated by the mammalian Sel-10 homolog.
77. PMID: 11425854; 2001, J Biol Chem;276(37):34371-8
Functional interaction between SEL-10, an F-box protein, and the nuclear form of activated Notch1 receptor.
78. PMID: 16863506; 2006, Cancer Sci;97(8):729-36
Fbw7 contributes to tumor suppression by targeting multiple proteins for ubiquitin-dependent degradation.
79. PMID: 18787170; 2008, Science;321(5895):1499-502
FBXW7 targets mTOR for degradation and cooperates with PTEN in tumor suppression.
80. PMID: 20484041; 2010, Cancer Res;70(11):4728-38
The Fbw7 tumor suppressor targets KLF5 for ubiquitin-mediated degradation and suppresses breast cell proliferation.
81. PMID: 21368833; 2011, Nature;471(7336):104-9
SCF(FBW7) regulates cellular apoptosis by targeting MCL1 for ubiquitylation and destruction.
82. PMID: 18094723; 2008, Nat Rev Cancer;8(2):83-93
FBW7 ubiquitin ligase: a tumour suppressor at the crossroads of cell division, growth and differentiation.
83. PMID: 23032637; 2012, Cancer Inform;11():157-71
Haploinsufficiency of Tumor Suppressor Genes is Driven by the Cumulative Effect of microRNAs, microRNA Binding Site Polymorphisms and microRNA Polymorphisms: An In silico Approach.
84. PMID: 24586741; 2014, PLoS One;9(2):e89388
FBXW7 mutations in patients with advanced cancers: clinical and molecular characteristics and outcomes with mTOR inhibitors.
85. PMID: 24360397; 2014, Lung Cancer;83(2):300-1
Temozolomide therapy in a patient with lung adenocarcinoma harboring an FBXW7 mutation.
86. PMID: 27399335; 2017, Oncogene;36(6):787-796
FBW7 mutations mediate resistance of colorectal cancer to targeted therapies by blocking Mcl-1 degradation.
87. PMID: 25860929; 2015, Oncotarget;6(11):9240-56
FBXW7-mutated colorectal cancer cells exhibit aberrant expression of phosphorylated-p53 at Serine-15.
88. PMID: 29633504; 2018, Mol Oncol;12(6):883-895
FBXW7 deletion contributes to lung tumor development and confers resistance to gefitinib therapy.
89. PMID: 28522751; 2017, Cancer Res;77(13):3527-3539
Targeting FBW7 as a Strategy to Overcome Resistance to Targeted Therapy in Non-Small Cell Lung Cancer.
90. PMID: 24884509; 2014, Mol Cancer;13():110
Fbw7 is an independent prognostic marker and induces apoptosis and growth arrest by regulating YAP abundance in hepatocellular carcinoma.
91. PMID: 17218262; 2007, Cell;128(1):157-70
Essential role for nuclear PTEN in maintaining chromosomal integrity.
92. PMID: 18794879; 2008, Oncogene;27(41):5443-53
PTEN: a new guardian of the genome.

ACT Onco[®] + Report

93. PMID: 18767981; 2009, Annu Rev Pathol;4():127-50
PTEN and the PI3-kinase pathway in cancer.
94. PMID: 11553783; 2001, Proc Natl Acad Sci U S A;98(20):11563-8
Haploinsufficiency of the Pten tumor suppressor gene promotes prostate cancer progression.
95. PMID: 20400965; 2010, Nat Genet;42(5):454-8
Subtle variations in Pten dose determine cancer susceptibility.
96. PMID: 9467011; 1998, Hum Mol Genet;7(3):507-15
Mutation spectrum and genotype-phenotype analyses in Cowden disease and Bannayan-Zonana syndrome, two hamartoma syndromes with germline PTEN mutation.
97. PMID: 24136893; 2013, J Natl Cancer Inst;105(21):1607-16
Cowden syndrome and the PTEN hamartoma tumor syndrome: systematic review and revised diagnostic criteria.
98. PMID: 21430697; 2011, Nat Rev Cancer;11(4):289-301
PTEN loss in the continuum of common cancers, rare syndromes and mouse models.
99. PMID: 18455982; 2008, Cell;133(3):403-14
Tenets of PTEN tumor suppression.
100. PMID: 9393738; 1997, Cancer Res;57(23):5221-5
MMAC1/PTEN mutations in primary tumor specimens and tumor cell lines.
101. PMID: 9829719; 1998, Clin Cancer Res;4(11):2577-83
Loss of heterozygosity and mutational analysis of the PTEN/MMAC1 gene in synchronous endometrial and ovarian carcinomas.
102. PMID: 9582022; 1998, Oncogene;16(13):1743-8
Analysis of PTEN and the 10q23 region in primary prostate carcinomas.
103. PMID: 9671321; 1998, Oncogene;17(1):123-7
Allelic loss of chromosome 10q23 is associated with tumor progression in breast carcinomas.
104. PMID: 11504908; 2001, Proc Natl Acad Sci U S A;98(18):10314-9
Enhanced sensitivity of PTEN-deficient tumors to inhibition of FRAP/mTOR.
105. PMID: 23714559; 2013, Am Soc Clin Oncol Educ Book;():
Targeting the PI3K/AKT/mTOR pathway: biomarkers of success and tribulation.
106. PMID: 20231295; 2010, J Biol Chem;285(20):14980-14989
Phosphoinositide 3-kinase pathway activation in phosphate and tensin homolog (PTEN)-deficient prostate cancer cells is independent of receptor tyrosine kinases and mediated by the p110beta and p110delta catalytic subunits.
107. PMID: 23287563; 2013, Clin Cancer Res;19(7):1760-72
Targeting activated Akt with GDC-0068, a novel selective Akt inhibitor that is efficacious in multiple tumor models.
108. PMID: 17047067; 2006, Cancer Res;66(20):10040-7
Inhibition of mammalian target of rapamycin or apoptotic pathway induces autophagy and radiosensitizes PTEN null prostate cancer cells.
109. PMID: 22422409; 2012, Clin Cancer Res;18(6):1777-89
PIK3CA/PTEN mutations and Akt activation as markers of sensitivity to allosteric mTOR inhibitors.
110. PMID: 22662154; 2012, PLoS One;7(5):e37431
Genotype-dependent efficacy of a dual PI3K/mTOR inhibitor, NVP-BEZ235, and an mTOR inhibitor, RAD001, in endometrial carcinomas.
111. PMID: 23136191; 2012, Clin Cancer Res;18(24):6771-83
Phosphoinositide 3-kinase (PI3K) pathway alterations are associated with histologic subtypes and are predictive of sensitivity to PI3K inhibitors in lung cancer preclinical models.

ACT Onco[®] + Report

112. PMID: 27091708; 2016, J Clin Oncol;34(18):2115-24
Molecular Alterations and Everolimus Efficacy in Human Epidermal Growth Factor Receptor 2-Overexpressing Metastatic Breast Cancers: Combined Exploratory Biomarker Analysis From BOLERO-1 and BOLERO-3.
113. PMID: 21306238; 2011, N Engl J Med;364(6):514-23
Everolimus for advanced pancreatic neuroendocrine tumors.
114. PMID: 23582881; 2013, Eur Urol;64(1):150-8
Phase 2 trial of single-agent everolimus in chemotherapy-naïve patients with castration-resistant prostate cancer (SAKK 08/08).
115. PMID: 28330462; 2017, BMC Cancer;17(1):211
Prospective phase II trial of everolimus in PIK3CA amplification/mutation and/or PTEN loss patients with advanced solid tumors refractory to standard therapy.
116. PMID: 27016228; 2016, Gynecol Oncol;141(1):43-8
Tumor mutational analysis of GOG248, a phase II study of temsirolimus or temsirolimus and alternating megestrol acetate and tamoxifen for advanced endometrial cancer (EC): An NRG Oncology/Gynecologic Oncology Group study.
117. PMID: 26951309; 2016, J Clin Oncol;34(14):1660-8
Randomized Open-Label Phase II Trial of Apatolisib (GDC-0980), a Novel Inhibitor of the PI3K/Mammalian Target of Rapamycin Pathway, Versus Everolimus in Patients With Metastatic Renal Cell Carcinoma.
118. PMID: 15324695; 2004, Cancer Cell;6(2):117-27
PTEN activation contributes to tumor inhibition by trastuzumab, and loss of PTEN predicts trastuzumab resistance in patients.
119. PMID: 20813970; 2010, Am J Pathol;177(4):1647-56
PTEN, PIK3CA, p-AKT, and p-p70S6K status: association with trastuzumab response and survival in patients with HER2-positive metastatic breast cancer.
120. PMID: 21135276; 2011, J Clin Oncol;29(2):166-73
Loss of phosphatase and tensin homolog or phosphoinositol-3 kinase activation and response to trastuzumab or lapatinib in human epidermal growth factor receptor 2-overexpressing locally advanced breast cancers.
121. PMID: 21594665; 2011, Breast Cancer Res Treat;128(2):447-56
Evaluation of the association of PIK3CA mutations and PTEN loss with efficacy of trastuzumab therapy in metastatic breast cancer.
122. PMID: 17936563; 2007, Cancer Cell;12(4):395-402
A functional genetic approach identifies the PI3K pathway as a major determinant of trastuzumab resistance in breast cancer.
123. PMID: 18700047; 2008, BMC Cancer;8():234
Potential value of PTEN in predicting cetuximab response in colorectal cancer: an exploratory study.
124. PMID: 17940504; 2007, Br J Cancer;97(8):1139-45
PTEN loss of expression predicts cetuximab efficacy in metastatic colorectal cancer patients.
125. PMID: 19398573; 2009, J Clin Oncol;27(16):2622-9
PTEN expression and KRAS mutations on primary tumors and metastases in the prediction of benefit from cetuximab plus irinotecan for patients with metastatic colorectal cancer.
126. PMID: 19953097; 2010, Br J Cancer;102(1):162-4
PTEN status in advanced colorectal cancer treated with cetuximab.
127. PMID: 27605871; 2016, World J Gastroenterol;22(28):6345-61
Mechanisms of resistance to anti-epidermal growth factor receptor inhibitors in metastatic colorectal cancer.
128. PMID: 24666267; 2014, Acta Oncol;53(7):852-64
The predictive value of KRAS, NRAS, BRAF, PIK3CA and PTEN for anti-EGFR treatment in metastatic colorectal cancer: A systematic review and meta-analysis.
129. PMID: 19351834; 2009, Cancer Res;69(8):3256-61

ACT Onco[®] + Report

PTEN loss contributes to erlotinib resistance in EGFR-mutant lung cancer by activation of Akt and EGFR.

130. PMID: 23133538; 2012, PLoS One;7(10):e48004
Modeling of tumor progression in NSCLC and intrinsic resistance to TKI in loss of PTEN expression.
131. PMID: 23592446; 2013, J Cell Biochem;114(6):1248-56
mTOR inhibitors radiosensitize PTEN-deficient non-small-cell lung cancer cells harboring an EGFR activating mutation by inducing autophagy.
132. PMID: 26645196; 2016, Cancer Discov;6(2):202-16
Loss of PTEN Promotes Resistance to T Cell-Mediated Immunotherapy.
133. PMID: 28228279; 2017, Immunity;46(2):197-204
Loss of PTEN Is Associated with Resistance to Anti-PD-1 Checkpoint Blockade Therapy in Metastatic Uterine Leiomyosarcoma.
134. PMID: 30150660; 2018, Nat Genet;50(9):1271-1281
Genomic correlates of response to immune checkpoint blockade in microsatellite-stable solid tumors.
135. PMID: 21468130; 2011, Nat Rev Clin Oncol;8(5):302-6
Treatment with olaparib in a patient with PTEN-deficient endometrioid endometrial cancer.
136. PMID: 23810788; 2013, Lancet Oncol;14(9):882-92
The poly(ADP-ribose) polymerase inhibitor niraparib (MK4827) in BRCA mutation carriers and patients with sporadic cancer: a phase 1 dose-escalation trial.
137. PMID: 23565244; 2013, PLoS One;8(4):e60408
PARP inhibition sensitizes to low dose-rate radiation TMPRSS2-ERG fusion gene-expressing and PTEN-deficient prostate cancer cells.
138. PMID: 25935112; 2015, Trends Biochem Sci;40(6):296-308
Structural determinants of Smad function in TGF- β signaling.
139. PMID: 19014666; 2008, Pathogenetics;1(1):2
Smad4 haploinsufficiency: a matter of dosage.
140. PMID: 9545410; 1998, Am J Hum Genet;62(5):1129-36
A gene for familial juvenile polyposis maps to chromosome 18q21.1.
141. PMID: 8553070; 1996, Science;271(5247):350-3
DPC4, a candidate tumor suppressor gene at human chromosome 18q21.1.
142. PMID: 8673134; 1996, Nat Genet;13(3):343-6
Evaluation of candidate tumour suppressor genes on chromosome 18 in colorectal cancers.
143. PMID: 18662538; 2008, Cell;134(2):215-30
TGFbeta in Cancer.
144. PMID: 9135016; 1997, Cancer Res;57(9):1731-4
Tumor-suppressive pathways in pancreatic carcinoma.
145. PMID: 23139211; 2013, Cancer Res;73(2):725-35
SMAD2, SMAD3 and SMAD4 mutations in colorectal cancer.
146. PMID: 22810696; 2012, Nature;487(7407):330-7
Comprehensive molecular characterization of human colon and rectal cancer.
147. PMID: 25890228; 2015, World J Surg Oncol;13():128
Clinical outcome and expression of mutant P53, P16, and Smad4 in lung adenocarcinoma: a prospective study.
148. PMID: 19841540; 2009, J Clin Invest;119(11):3208-11
Smad4: gatekeeper gene in head and neck squamous cell carcinoma.

ACT Onco[®] + Report

149. PMID: 15867212; 2005, Clin Cancer Res;11(9):3191-7
Differences in Smad4 expression in human papillomavirus type 16-positive and human papillomavirus type 16-negative head and neck squamous cell carcinoma.
150. PMID: 25589618; 2015, Clin Cancer Res;21(6):1447-56
Genomic analysis of metastatic cutaneous squamous cell carcinoma.
151. PMID: 29703253; 2018, BMC Cancer;18(1):479
SMAD4 and NF1 mutations as potential biomarkers for poor prognosis to cetuximab-based therapy in Chinese metastatic colorectal cancer patients.
152. PMID: 28522603; 2017, Clin Cancer Res;23(17):5162-5175
SMAD4 Loss Is Associated with Cetuximab Resistance and Induction of MAPK/JNK Activation in Head and Neck Cancer Cells.
153. PMID: 16144935; 2005, Clin Cancer Res;11(17):6311-6
SMAD4 levels and response to 5-fluorouracil in colorectal cancer.
154. PMID: 24384683; 2014, Br J Cancer;110(4):946-57
Loss of Smad4 in colorectal cancer induces resistance to 5-fluorouracil through activating Akt pathway.
155. PMID: 12237773; 2002, Br J Cancer;87(6):630-4
SMAD4 is a predictive marker for 5-fluorouracil-based chemotherapy in patients with colorectal cancer.
156. PMID: 25749173; 2015, Transl Oncol;8(1):18-24
A Meta-Analysis of SMAD4 Immunohistochemistry as a Prognostic Marker in Colorectal Cancer.
157. PMID: 19478385; 2009, Cell Oncol;31(3):169-78
Presence of a high amount of stroma and downregulation of SMAD4 predict for worse survival for stage I-II colon cancer patients.
158. PMID: 25681512; 2015, J Clin Pathol;68(5):341-5
Smad4 inactivation predicts for worse prognosis and response to fluorouracil-based treatment in colorectal cancer.
159. PMID: 26861460; 2016, Clin Cancer Res;22(12):3037-47
Reduced Expression of SMAD4 Is Associated with Poor Survival in Colon Cancer.
160. PMID: 26947875; 2016, Transl Oncol;9(1):1-7
Prognostic Value of SMAD4 in Pancreatic Cancer: A Meta-Analysis.
161. PMID: 25760429; 2015, Pancreas;44(4):660-4
SMAD4 expression predicts local spread and treatment failure in resected pancreatic cancer.
162. PMID: 22504380; 2012, Pancreas;41(4):541-6
SMAD4 genetic alterations predict a worse prognosis in patients with pancreatic ductal adenocarcinoma.
163. PMID: 19584151; 2009, Clin Cancer Res;15(14):4674-9
SMAD4 gene mutations are associated with poor prognosis in pancreatic cancer.
164. PMID: 18425078; 2008, Mod Pathol;21(7):866-75
Expression of Smad2 and Smad4 in cervical cancer: absent nuclear Smad4 expression correlates with poor survival.
165. PMID: 26703889; 2016, Lancet;387(10022):968-977
Everolimus for the treatment of advanced, non-functional neuroendocrine tumours of the lung or gastrointestinal tract (RADIANT-4): a randomised, placebo-controlled, phase 3 study.
166. PMID: 22149876; 2012, N Engl J Med;366(6):520-9
Everolimus in postmenopausal hormone-receptor-positive advanced breast cancer.
167. PMID: 23158522; 2013, Lancet;381(9861):125-32
Efficacy and safety of everolimus for subependymal giant cell astrocytomas associated with tuberous sclerosis complex (EXIST-1): a multicentre, randomised, placebo-controlled phase 3 trial.

ACT Onco[®] + Report

168. PMID: 18653228; 2008, Lancet;372(9637):449-56
Efficacy of everolimus in advanced renal cell carcinoma: a double-blind, randomised, placebo-controlled phase III trial.
169. PMID: 27717299; 2016, N Engl J Med;375(22):2154-2164
Niraparib Maintenance Therapy in Platinum-Sensitive, Recurrent Ovarian Cancer.
170. PMID: 32343890; 2020, N Engl J Med;382(22):2091-2102
Olaparib for Metastatic Castration-Resistant Prostate Cancer.
171. PMID: 31851799; 2019, N Engl J Med;381(25):2416-2428
Olaparib plus Bevacizumab as First-Line Maintenance in Ovarian Cancer.
172. PMID: 31157963; 2019, N Engl J Med;381(4):317-327
Maintenance Olaparib for Germline BRCA-Mutated Metastatic Pancreatic Cancer.
173. PMID: 30345884; 2018, N Engl J Med;379(26):2495-2505
Maintenance Olaparib in Patients with Newly Diagnosed Advanced Ovarian Cancer.
174. PMID: 28578601; 2017, N Engl J Med;377(6):523-533
Olaparib for Metastatic Breast Cancer in Patients with a Germline BRCA Mutation.
175. PMID: 28754483; 2017, Lancet Oncol;18(9):1274-1284
Olaparib tablets as maintenance therapy in patients with platinum-sensitive, relapsed ovarian cancer and a BRCA1/2 mutation (SOLO2/ENGOT-Ov21): a double-blind, randomised, placebo-controlled, phase 3 trial.
176. PMID: 27617661; 2016, Lancet Oncol;17(11):1579-1589
Overall survival in patients with platinum-sensitive recurrent serous ovarian cancer receiving olaparib maintenance monotherapy: an updated analysis from a randomised, placebo-controlled, double-blind, phase 2 trial.
177. PMID: 28916367; 2017, Lancet;390(10106):1949-1961
Rucaparib maintenance treatment for recurrent ovarian carcinoma after response to platinum therapy (ARIEL3): a randomised, double-blind, placebo-controlled, phase 3 trial.
178. PMID: 30110579; 2018, N Engl J Med;379(8):753-763
Talazoparib in Patients with Advanced Breast Cancer and a Germline BRCA Mutation.
179. PMID: 17538086; 2007, N Engl J Med;356(22):2271-81
Temsilolimus, interferon alfa, or both for advanced renal-cell carcinoma.

ACT Onco[®] + Report

PATIENT		
Identifier: 歐陽國華		Patient ID: 49497941
Date of Birth: Sep 28, 1963		Gender: Male
Diagnosis: Pancreatic cancer		
ORDERING PHYSICIAN		
Name: 姜乃榕醫師		Tel: 886-228712121
Facility: 臺北榮總		
Address: 臺北市北投區石牌路二段 201 號		
SPECIMEN		
Specimen ID: S11291425A	Collection site: Pancreas	Type: FFPE tissue
Date received: May 16, 2023	Lab ID: AA-23-03087	D/ID: NA
Date reported: May 29, 2023	Project ID: C23-M001-01474	Testing panel: ACTOnco+

VARIANT(S) WITH CLINICAL RELEVANCE

- Single Nucleotide and Small InDel Variants

Gene Alterations	Allele Frequency	Classification
BRCA2 c.5645C>A (S1882*)	59.5%	Deleterious

Notes:

The addendum report further provides the AMP/ACMG classification of deleterious and suspected deleterious BRCA1/2 mutations if detected by ACTOnco assay based on the reimbursement requirement of Taiwan National Health Insurance.



王業翰 醫師

Yeh-Han Wang M.D.

病解專醫字第 000545 號