



Sample Information

Patient Name: 楊桑滿
Gender: Female
ID No.: R222228590
History No.: 17606198
Age: 63

Ordering Doctor: DOC1901H 高志平
Ordering REQ.: OBRWUBN
Signing in Date: 2022/02/11

Path No.: S111-98363
MP No.: MY22005
Assay: Oncomine Myeloid Assay
Sample Type: Bone Marrow
Bone Marrow Aspirating Date: 2022/02/07

Reporting Doctor: DOC5466K 葉奕成 (Phone: 8#5466)

Note:

Sample Cancer Type: Acute Myeloid Leukemia

Table of Contents	Page
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2
Biomarker Descriptions	2
Relevant Therapy Summary	4
Relevant Therapy Details	4
Prognostic Details	8
Clinical Trials Summary	8
Alert Details	9

Report Highlights
 3 Relevant Biomarkers
 4 Therapies Available
 1 Clinical Trials

Relevant Acute Myeloid Leukemia Variants

Gene	Finding	Gene	Finding
ABL1	None detected	MECOM	None detected
ASXL1	ASXL1 p.(R693*) c.2077C>T, ASXL1 p.(E635Rfs*15) c.1900_1922delAGAGAGGCGGCCACCACT GCCAT	MLLT3	None detected
CEBPA	None detected	MYH11	None detected
CREBBP	None detected	NPM1	None detected
FLT3	None detected	NUP214	None detected
IDH1	IDH1 p.(R132C) c.394C>T	RARA	None detected
IDH2	None detected	RUNX1	None detected
KMT2A	None detected	TP53	None detected

Relevant Biomarkers

Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IDH1 p.(R132C) c.394C>T isocitrate dehydrogenase (NADP(+)) 1 Allele Frequency: 30.83% Prognostic significance: None Diagnostic significance: None	ivosidenib ¹ azacitidine decitabine venetoclax + chemotherapy	None	1
ASXL1 p.(R693*) c.2077C>T, ASXL1 p.(E635Rfs*15) c.1900_1922delAGAGAGGCGGC CACCACTGCCAT ASXL transcriptional regulator 1 Allele Frequency: 14.85%, 10.48% (2 variants) Prognostic significance: ELN 2017: Adverse Diagnostic significance: None	None	None	0

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

Prevalent cancer biomarkers without relevant evidence based on included data sources

PHF6 p.(Y103*) c.308_309insA, PHF6 p.(R274*) c.820C>T

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
IDH1	p.(R132C)	c.394C>T	COSM28747	chr2:209113113	30.83%	NM_005896.3	missense	1998
ASXL1	p.(E635Rfs*15)	c.1900_1922delAGA GAGGCGGCCACCACTGCCAT	.	chr20:31022403	10.48%	NM_015338.6	frameshift Deletion	1889
ASXL1	p.(R693*)	c.2077C>T	COSM51388	chr20:31022592	14.85%	NM_015338.6	nonsense	2000
PHF6	p.(Y103*)	c.308_309insA	.	chrX:133527597	16.94%	NM_032458.3	nonsense	1995
PHF6	p.(R274*)	c.820C>T	COSM144567	chrX:133549136	12.56%	NM_032458.3	nonsense	1999
BCOR	p.(A1527=)	c.4581C>T	.	chrX:39916422	49.90%	NM_001123385.2	synonymous	2000

Biomarker Descriptions

ASXL1 (ASXL transcriptional regulator 1)

Background: The ASXL1 gene encodes the ASXL transcriptional regulator 1 protein, a ligand-dependent co-activator and epigenetic scaffolding protein involved in transcriptional regulation^{1,2}. ASXL1 belongs to the ASXL gene family, which also includes ASXL2 and ASXL3². ASXL proteins contain a conserved c-terminal plant homeodomain (PHD) which facilitates interaction with DNA and histones^{2,3}. ASXL1 influences chromatin remodeling and transcription through interaction with BAP1 and polycomb repressive complex (PRC) proteins, as well as other transcriptional activators and repressors^{2,4}. In cancer, ASXL1 is the target of somatic mutations which often result in a truncated ASXL1 protein and loss of its PHD^{5,6,7}. Such mutations can lead to impaired protein function and consequent upregulation of HOXA gene expression, supporting a tumor suppressor role for ASXL1⁸.

Biomarker Descriptions (continued)

Alterations and prevalence: Missense, nonsense, and frameshift mutations in ASXL1 are reported in 3-6% of de novo acute myeloid leukemia (AML), up to 36% of secondary AML, approximately 15% of myelodysplastic syndromes (MDS), up to 23% of myeloproliferative neoplasms (MPN), up to 30% of systemic mastocytosis (SM), and approximately 45% of chronic myelomonocytic leukemia (CMML)^{4,9,10,11,12,13,14,15,16}. The ASXL1 G646Wfs*12 mutation accounts for over 50% of ASXL1 mutated cases in myeloid malignancies^{6,11,17}. This mutation results from a single nucleotide expansion that occurs within an eight base pair guanine repeat that extends from c.1927 to c.1934. It is proposed that the high prevalence of the G646Wfs*12 variant is due to replication slippage which can occur in areas of repetitive sequence¹⁸. As a consequence, detection of G646Wfs*12 may result as an artifact of PCR and/or sequencing¹⁹. However, multiple studies observe an increase in the frequency of G646Wfs*12 in myeloid cancer relative to normal suggesting that G646Wfs*12 is a bona fide somatic mutation^{9,18,20}.

Potential relevance: The majority of frameshift and nonsense mutations in ASXL1 that result in protein truncation and removal of the PHD domain are considered pathogenic²¹. Mutations in ASXL1 confer poor/adverse risk in AML and are associated with poor outcomes and adverse risk¹⁶. Additionally, ASXL1 nonsense or frameshift mutations are independently associated with poor prognosis in MDS and CMML²². Moreover, ASXL1 mutations are independently associated with inferior overall survival (OS) in patients with MPN or SM^{23,24}.

IDH1 (isocitrate dehydrogenase (NADP(+)) 1)

Background: The IDH1 and IDH2 genes encode homologous isocitrate dehydrogenase enzymes that catalyze the conversion of isocitrate to α -ketoglutarate (α -KG)²⁵. The IDH1 gene encodes the NADP+ dependent cytoplasmic isocitrate dehydrogenase enzyme; IDH2 encodes the mitochondrial isoform.

Alterations and prevalence: Recurrent somatic mutations in IDH1 and IDH2 are mutually exclusive and observed in several malignancies including glioma, chondrosarcoma, intrahepatic cholangiocarcinoma, acute myeloid leukemia (AML), and myelodysplastic syndrome (MDS)²⁶. Recurrent IDH1 variants include predominately R132H/C plus other substitutions at lower frequencies. These gain of function variants confer neomorphic enzyme activity²⁷. Although wild-type enzymatic activity is ablated, recurrent IDH1 variants catalyze the conversion of α -KG to D-2-hydroxyglutarate, an oncometabolite with diverse effects on cellular metabolism, epigenetic regulation, redox states, and DNA repair^{25,28}. Recurrent IDH1 mutations are present in 5-10% of patients with AML and 5% of patients with MDS^{29,30,31}. Recurrent IDH1 mutations are present in nearly 80% of lower grade diffuse gliomas^{12,32}.

Potential relevance: Ivosidenib³³ is FDA approved (2018) for the treatment of AML patients with IDH1 R132C/G/H/L/S variants³⁴. Ivosidenib has also been granted breakthrough designation (2020) for IDH1 mutated relapsed or refractory myelodysplastic syndrome (MDS)³⁵. IDH1 mutations are associated with inferior leukemia-free survival in primary myelofibrosis (PMF) and inferior overall survival in polycythemia vera (PV) but have been shown to confer improved prognosis in lower grade gliomas^{23,36,37}.

PHF6 (PHD finger protein 6)

Background: The PHF6 gene encodes the plant homeodomain (PHD) finger protein 6 which contains four nuclear localization signals and two imperfect PHD zinc finger domains. PHF6 is a tumor suppressor that interacts with the nucleosome remodeling deacetylase (NuRD) complex, which regulates nucleosome positioning and transcription of genes involved in development and cell-cycle progression^{38,39}.

Alterations and prevalence: The majority of PHF6 aberrations are nonsense, frameshift (70%), or missense (30%) mutations, which result in complete loss of protein expression^{38,40,41,42}. Truncating or missense mutations in PHF6 are observed in 38% of adult and 16% of pediatric T-cell acute lymphoblastic leukemia (T-ALL), 20-25% of mixed phenotype acute leukemias (MPAL), and 3% of AML, and 2.6% of hepatocellular carcinoma (HCC)^{40,42}. Missense mutations recurrently involve codon C215 and the second zinc finger domain of PHF6⁴⁰. PHF6 mutations are frequently observed in hematologic malignancies from male patients^{38,40}.

Potential relevance: Somatic mutations in PHF6 are associated with reduced overall survival in AML patients treated with high-dose induction chemotherapy⁴³.

Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

IDH1 p.(R132C) c.394C>T

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
ivosidenib	●	●	×	●	×
azacitidine	×	●	×	×	×
decitabine	×	●	×	×	×
venetoclax + azacitidine	×	●	×	×	×
venetoclax + cytarabine	×	●	×	×	×
venetoclax + decitabine	×	●	×	×	×
LY-3410738	×	×	×	×	● (I)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Details

Current FDA Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

FDA information is current as of 2021-11-17. For the most up-to-date information, search www.fda.gov.

IDH1 p.(R132C) c.394C>T

● ivosidenib

Cancer type: Acute Myeloid Leukemia

Label as of: 2021-08-25

Variant class: IDH1 R132C mutation

Indications and usage:

TIBSOVO® is an isocitrate dehydrogenase-1 (IDH1) inhibitor indicated for the treatment of adult patients with a susceptible IDH1 mutation as detected by an FDA-approved test with:

Acute Myeloid Leukemia (AML)

- Newly-diagnosed AML who are ≥ 75 years old or who have comorbidities that preclude use of intensive induction chemotherapy.
- Relapsed or refractory AML.

Locally Advanced or Metastatic Cholangiocarcinoma

- Locally advanced or metastatic cholangiocarcinoma who have been previously treated.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/211192_s008lbl.pdf

Current NCCN Information

- ☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

NCCN information is current as of 2021-11-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

IDH1 p.(R132C) c.394C>T

● venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2021]

● azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2021]

● decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2021]

● ivosidenib

Cancer type: Acute Myeloid Leukemia

Variant class: IDH1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention
- Relapsed, Refractory (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2021]

IDH1 p.(R132C) c.394C>T (continued)**● venetoclax + cytarabine**

Cancer type: Acute Myeloid Leukemia

Variant class: IDH1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2021]

● venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2021]

Current ESMO Information

☒ In this cancer type ☐ In other cancer type ☐ In this cancer type and other cancer types

ESMO information is current as of 2021-11-01. For the most up-to-date information, search www.esmo.org.

IDH1 p.(R132C) c.394C>T

☒ ivosidenib

Cancer type: Acute Myeloid Leukemia

Variant class: IDH1 mutation

ESMO Level of Evidence/Grade of Recommendation: IV / B

Population segment (Line of therapy):

- Relapsed, Refractory (Second-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

Prognostic Details

Current NCCN Information

NCCN information is current as of 2021-11-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

**ASXL1 p.(R693*) c.2077C>T, ASXL1 p.(E635Rfs*15)
c.1900_1922delAGAGAGGCGGCCACCACTGCCAT**

Prognostic significance: ELN 2017: Adverse

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

NCCN Recommendation category: 2A

Summary:

- Do not use as an adverse prognostic marker if it co-occurs with favorable-risk AML subtypes.

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2021]

Current ESMO Information

ESMO information is current as of 2021-11-01. For the most up-to-date information, search www.esmo.org.

**ASXL1 p.(R693*) c.2077C>T, ASXL1 p.(E635Rfs*15)
c.1900_1922delAGAGAGGCGGCCACCACTGCCAT**

Prognostic significance: ELN 2017: Adverse

Cancer type: Acute Myeloid Leukemia

Variant class: ASXL1 mutation

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]


Clinical Trials Summary

IDH1 p.(R132C) c.394C>T

NCT ID	Title	Phase
NCT04603001	A Phase I Study of Oral LY3410738 in Patients With Advanced Hematologic Malignancies With IDH1 or IDH2 Mutations.	I

Alerts Informed By Public Data Sources


Current FDA Information

 Contraindicated

 Not recommended

 Resistance

 Breakthrough

 Fast Track

FDA information is current as of 2021-11-17. For the most up-to-date information, search www.fda.gov.

IDH1 p.(R132C) c.394C>T

ivosidenib

Cancer type: Myelodysplastic Syndrome

Variant class: IDH1 mutation

Supporting Statement:

The FDA has granted Breakthrough Designation to the isocitrate dehydrogenase-1 inhibitor, ivosidenib, for the treatment of adult patients with relapsed or refractory myelodysplastic syndrome (MDS) with a susceptible IDH1 mutation as detected by an FDA-approved test.

Reference:

<https://investor.agios.com/news-releases/news-release-details/agios-receives-fda-breakthrough-therapy-designation-tibsovor-0>

Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

References

1. O'Leary et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res.* 2016 Jan 4;44(D1):D733-45. PMID: 26553804
2. Katoh. Functional and cancer genomics of ASXL family members. *Br. J. Cancer.* 2013 Jul 23;109(2):299-306. PMID: 23736028
3. Gelsi-Boyer et al. Mutations of polycomb-associated gene ASXL1 in myelodysplastic syndromes and chronic myelomonocytic leukaemia. *Br. J. Haematol.* 2009 Jun;145(6):788-800. PMID: 19388938
4. Gelsi-Boyer et al. Mutations in ASXL1 are associated with poor prognosis across the spectrum of malignant myeloid diseases. *J Hematol Oncol.* 2012 Mar 21;5:12. doi: 10.1186/1756-8722-5-12. PMID: 22436456
5. Larsson et al. The changing mutational landscape of acute myeloid leukemia and myelodysplastic syndrome. *Mol. Cancer Res.* 2013 Aug;11(8):815-27. PMID: 23645565
6. Alvarez et al. ASXL1 mutations in myeloid neoplasms: pathogenetic considerations, impact on clinical outcomes and survival. *Curr Med Res Opin.* 2018 May;34(5):757-763. PMID: 28027687
7. Yang et al. Gain of function of ASXL1 truncating protein in the pathogenesis of myeloid malignancies. *Blood.* 2018 Jan 18;131(3):328-341. PMID: 29113963
8. Abdel-Wahab et al. ASXL1 mutations promote myeloid transformation through loss of PRC2-mediated gene repression. *Cancer Cell.* 2012 Aug 14;22(2):180-93. PMID: 22897849
9. Alberti et al. Discriminating a common somatic ASXL1 mutation (c.1934dup; p.G646Wfs*12) from artifact in myeloid malignancies using NGS. *Leukemia.* 2018 Aug;32(8):1874-1878. PMID: 29959414
10. Kakosaiou et al. ASXL1 mutations in AML are associated with specific clinical and cytogenetic characteristics. *Leuk. Lymphoma.* 2018 Oct;59(10):2439-2446. PMID: 29411666
11. Paschka et al. ASXL1 mutations in younger adult patients with acute myeloid leukemia: a study by the German-Austrian Acute Myeloid Leukemia Study Group. *Haematologica.* 2015 Mar;100(3):324-30. PMID: 25596267
12. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
13. Jawhar et al. The clinical and molecular diversity of mast cell leukemia with or without associated hematologic neoplasm. *Haematologica.* 2017 Jun;102(6):1035-1043. PMID: 28255023
14. Jawhar et al. KIT D816 mutated/CBF-negative acute myeloid leukemia: a poor-risk subtype associated with systemic mastocytosis. *Leukemia.* 2019 May;33(5):1124-1134. PMID: 30635631
15. Damaj et al. ASXL1 but not TET2 mutations adversely impact overall survival of patients suffering systemic mastocytosis with associated clonal hematologic non-mast-cell diseases. *PLoS ONE.* 2014;9(1):e85362. PMID: 24465546
16. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2021]
17. Boulton et al. Frequent mutation of the polycomb-associated gene ASXL1 in the myelodysplastic syndromes and in acute myeloid leukemia. *Leukemia.* 2010 May;24(5):1062-5. doi: 10.1038/leu.2010.20. Epub 2010 Feb 25. PMID: 20182461
18. Yannakou et al. ASXL1 c.1934dup;p.Gly646Trpfs*12-a true somatic alteration requiring a new approach. *Blood Cancer J.* 2017 Dec 15;7(12):656. doi: 10.1038/s41408-017-0025-8. PMID: 29242575
19. Abdel-Wahab et al. The most commonly reported variant in ASXL1 (c.1934dupG;p.Gly646TrpfsX12) is not a somatic alteration. *Leukemia.* 2010 Sep;24(9):1656-7. doi: 10.1038/leu.2010.144. Epub 2010 Jul 1. PMID: 20596031
20. Montes-Moreno et al. Clinical molecular testing for ASXL1 c.1934dupG p.Gly646fs mutation in hematologic neoplasms in the NGS era. *PLoS ONE.* 2018;13(9):e0204218. PMID: 30222780
21. Landrum et al. ClinVar: improving access to variant interpretations and supporting evidence. *Nucleic Acids Res.* 2018 Jan 4;46(D1):D1062-D1067. PMID: 29165669
22. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2022]
23. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 2.2021]
24. NCCN Guidelines® - NCCN-Systemic Mastocytosis [Version 1.2020]
25. Molenaar et al. Wild-type and mutated IDH1/2 enzymes and therapy responses. *Oncogene.* 2018 Apr;37(15):1949-1960. PMID: 29367755
26. Yan et al. IDH1 and IDH2 mutations in gliomas. *N. Engl. J. Med.* 2009 Feb 19;360(8):765-73. PMID: 19228619
27. Dang et al. Cancer-associated IDH1 mutations produce 2-hydroxyglutarate. *Nature.* 2009 Dec 10;462(7274):739-44. PMID: 19935646
28. Ward et al. The common feature of leukemia-associated IDH1 and IDH2 mutations is a neomorphic enzyme activity converting alpha-ketoglutarate to 2-hydroxyglutarate. *Cancer Cell.* 2010 Mar 16;17(3):225-34. PMID: 20171147

References (continued)

29. Paschka et al. IDH1 and IDH2 mutations are frequent genetic alterations in acute myeloid leukemia and confer adverse prognosis in cytogenetically normal acute myeloid leukemia with NPM1 mutation without FLT3 internal tandem duplication. *J. Clin. Oncol.* 2010 Aug 1;28(22):3636-43. PMID: 20567020
30. Chou et al. The prognostic impact and stability of Isocitrate dehydrogenase 2 mutation in adult patients with acute myeloid leukemia. *Leukemia.* 2011 Feb;25(2):246-53. PMID: 21079611
31. Marcucci et al. IDH1 and IDH2 gene mutations identify novel molecular subsets within de novo cytogenetically normal acute myeloid leukemia: a Cancer and Leukemia Group B study. *J. Clin. Oncol.* 2010 May 10;28(14):2348-55. PMID: 20368543
32. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
33. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/211192_s008lbl.pdf
34. Abou et al. The role of enasidenib in the treatment of mutant IDH2 acute myeloid leukemia. *Ther Adv Hematol.* 2018 Jul;9(7):163-173. PMID: 30013764
35. <https://investor.agios.com/news-releases/news-release-details/agios-receives-fda-breakthrough-therapy-designation-tibsovor-0>
36. Cancer Genome Atlas Research Network. Comprehensive, Integrative Genomic Analysis of Diffuse Lower-Grade Gliomas. *N Engl J Med.* 2015 Jun 25;372(26):2481-98. doi: 10.1056/NEJMoa1402121. Epub 2015 Jun 10. PMID: 26061751
37. Houillier et al. IDH1 or IDH2 mutations predict longer survival and response to temozolomide in low-grade gliomas. *Neurology.* 2010 Oct 26;75(17):1560-6. PMID: 20975057
38. Wendorff et al. Phf6 Loss Enhances HSC Self-Renewal Driving Tumor Initiation and Leukemia Stem Cell Activity in T-ALL. *Cancer Discov.* 2019 Mar;9(3):436-451. PMID: 30567843
39. Lower et al. Mutations in PHF6 are associated with Börjeson-Forssman-Lehmann syndrome. *Nat. Genet.* 2002 Dec;32(4):661-5. PMID: 12415272
40. Van et al. PHF6 mutations in T-cell acute lymphoblastic leukemia. *Nat. Genet.* 2010 Apr;42(4):338-42. PMID: 20228800
41. Van et al. PHF6 mutations in adult acute myeloid leukemia. *Leukemia.* 2011 Jan;25(1):130-4. PMID: 21030981
42. Yoo et al. Somatic mutation of PHF6 gene in T-cell acute lymphoblastic leukemia, acute myelogenous leukemia and hepatocellular carcinoma. *Acta Oncol.* 2012 Jan;51(1):107-11. PMID: 21736506
43. Patel et al. Prognostic relevance of integrated genetic profiling in acute myeloid leukemia. *N. Engl. J. Med.* 2012 Mar 22;366(12):1079-89. PMID: 22417203