



Sample Information

Patient Name: 譚許秋華**Gender:** Female**ID No.:** H200374238**History No.:** 14030069**Age:** 77**Ordering Doctor:** DOC1878G 沈佳儀**Ordering REQ.:** 0BYDLCS**Signing in Date:** 2022/08/03**Path No.:** S111-97822**MP No.:** F22077**Assay:** Oncomine Focus Assay**Sample Type:** FFPE**Block No.:** S111-92042A**Percentage of tumor cells:** 10%**Reporting Doctor:** DOC5466K 葉奕成 (Phone: 8#5466)**Note:**

Sample Cancer Type: Non-Small Cell Lung Cancer

Table of Contents	Page
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2
Biomarker Descriptions	2
Relevant Therapy Summary	4
Relevant Therapy Details	5
Clinical Trials Summary	6
Alert Details	6

Report Highlights

- 2 Relevant Biomarkers
- 2 Therapies Available
- 4 Clinical Trials

Relevant Non-Small Cell Lung Cancer Variants

Gene	Finding	Gene	Finding
ALK	None detected	NTRK1	None detected
BRAF	None detected	NTRK2	None detected
EGFR	None detected	NTRK3	None detected
ERBB2	ERBB2 p.(L755_E757delinsPQ) c.2263_2269delTTGAGGGinsCCTC	RET	None detected
KRAS	None detected	ROS1	None detected
MET	None detected		

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	<i>ERBB2 p.(L755_E757delinsPQ) c.2263_2269delTTGAGGGinsCCT C</i> erb-b2 receptor tyrosine kinase 2 Allele Frequency: 6.92%	trastuzumab deruxtecan	None	4
IIC	<i>AR amplification</i> androgen receptor	None	hormone therapy	0

Public data sources included in relevant therapies: FDA¹, NCCN, EMA², ESMO

Tier Reference: Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

Prevalent cancer biomarkers without relevant evidence based on included data sources

MYCN amplification

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
ERBB2	p.(L755_E757delinsPQ)	c.2263_2269delTTGAGGGinsCCTC	.	chr17:37880219	6.92%	NM_004448.3	nonframeshift Block Substitution	1980

Copy Number Variations

Gene	Locus	Copy Number
MYCN	chr2:16080663	11.4
AR	chrX:66776186	12

Biomarker Descriptions

AR (androgen receptor)

Background: The AR gene encodes the androgen receptor protein (AR), a ligand-activated transcription factor regulated by the binding of the hormones testosterone and dihydrotestosterone^{1,2}. Hormone binding to AR results in receptor dimerization, nuclear translocation, and target gene transcription, thus activating the RAS/RAF/MEK/ERK and PI3K/AKT/MTOR signaling pathways, which promote cell proliferation and survival^{2,3,4}.

Alterations and prevalence: Alterations in AR function can result from overexpression, gene amplification, or mutations. AR mutations, including L702H, W742C/L, H875Y, and T878A, are commonly observed in 10-30% of castration-resistant prostate cancer and result in decreased ligand specificity, allowing other nuclear hormones to activate AR⁵. Androgen receptor splice variants have been reported in castration resistant prostate cancer^{6,7}. The androgen receptor splice variant 7 (AR-V7) is a result of aberrant mRNA splicing of AR exons 1-3 and a cryptic exon 3, resulting in the expression of a constitutively active protein⁷.

Potential relevance: The FDA has granted fast track designation (2022) to the selective androgen receptor targeting agonist, enobosarm, for the treatment of patients with androgen AR-positive, estrogen receptor (ER)-positive, HER2-negative metastatic breast cancer⁸. The FDA also granted fast track designation (2016) to the small-molecule CYP17 lyase-selective inhibitor, seviteronel, for AR-positive triple-negative breast cancer (TNBC) patients⁹. Androgen deprivation therapy (ADT) such as abiraterone¹⁰ (2011) and enzalutamide¹¹ (2011) are FDA approved for use in locally advanced and metastatic prostate cancers. Although many men initially respond to ADT, most will develop hormone resistance. Resistance to ADT is also associated with other aberrations of the AR gene including mutations within the ligand binding domain and gene amplification^{5,12,13}. The androgen receptor splice variant, AR-V7, lacks

Biomarker Descriptions (continued)

the ligand binding domain, resulting in constitutive activation and is associated with resistance to androgen deprivation therapy (ADT) in advanced prostate cancer⁶.

ERBB2 (erb-b2 receptor tyrosine kinase 2)

Background: The ERBB2 gene encodes the erb-b2 receptor tyrosine kinase 2, a member of the human epidermal growth factor receptor (HER) family. Along with ERBB2/HER2, EGFR/ERBB1/HER1, ERBB3/HER3, and ERBB4/HER4 make up the HER protein family¹⁴. All ERBB/HER proteins encode transmembrane receptor tyrosine kinases. However, ERBB2/HER2 is an orphan receptor with no known ligand. ERBB2 preferentially binds other ligand bound ERBB/HER family members to form hetero-dimers resulting in the activation of ERBB2 tyrosine kinase activity and subsequent activation of the PI3K/AKT/MTOR and RAS/RAF/MAPK/ERK signaling pathways which promote cell proliferation, differentiation, and survival¹⁵. Recurrent focal amplification of the ERBB2 gene leads to increased expression in several cancer types. ERBB2 overexpression in immortalized cell lines is oncogenic and leads to ERBB2 homo-dimerization and activation without ligand binding^{16,17,18}.

Alterations and prevalence: ERBB2 gene amplification occurs in 10-20% of breast, esophageal, and gastric cancers, 5-10% of bladder, cervical, pancreas, and uterine cancers, and 1-5% of colorectal, lung, and ovarian cancers^{19,20,21,22,23,24,25,26}. Recurrent somatic activating mutations in ERBB2/HER2 occur at low frequencies (<1%) in diverse cancer types^{26,27,28}. In breast, bladder, and colorectal cancers, the most common recurrent ERBB2 activating mutations include kinase domain mutations L755S and V777L and the extracellular domain mutation S310F. In lung cancer, the most common recurrent ERBB2 activating mutations include in-frame exon 20 insertions, particularly Y772_A775dup.

Potential relevance: The discovery of ERBB2/HER2 as an important driver of breast cancer in 1987 led to the development of trastuzumab, a humanized monoclonal antibody with specificity to the extracellular domain of HER2^{29,30}. Trastuzumab³¹ was FDA approved for the treatment of HER2 positive breast cancer in 1998, and subsequently in HER2 positive metastatic gastric and gastroesophageal junction adenocarcinoma in 2010. Additional monoclonal antibody therapies have been approved by the FDA for HER2-positive breast cancer including pertuzumab³² (2012), a humanized monoclonal antibody that inhibits HER2 dimerization, and ado-trastuzumab emtansine³³ (2013), a conjugate of trastuzumab and a potent antimicrotubule agent. The combination of pertuzumab, trastuzumab, and a taxane is the preferred front-line regimen for HER2-positive metastatic breast cancer³⁴. In addition to monoclonal antibodies, the small molecule inhibitor lapatinib³⁵, with specificity for both EGFR and ERBB2, was FDA approved (2007) for the treatment of patients with advanced HER2-positive breast cancer who have received prior therapy including trastuzumab. In 2017, the FDA approved the use of neratinib³⁶, an irreversible kinase inhibitor of EGFR, ERBB2/HER2, and ERBB4, for the extended adjuvant treatment of adult patients with early stage HER2-positive breast cancer. In 2020, the FDA approved neratinib³⁶ in combination with capecitabine for HER2-positive advanced or metastatic patients after two or more prior HER2-directed therapies. Also in 2020, the TKI irbinetinib³⁷ was FDA approved for HER2 overexpressing or amplified breast cancer in combination with trastuzumab and capecitabine. In 2021, the PD-1 blocking antibody, pembrolizumab, in combination with trastuzumab, fluoropyrimidine- and platinum-based chemotherapy, was approved for HER2 amplified gastric or gastroesophageal (GEJ) adenocarcinoma in the first line³⁸. The vaccine, nelipecimut-S³⁹, was granted fast-track designation by the FDA (2016) in patients with low to intermediate HER2 expressing (IHC score 1+ or 2+) breast cancer. In 2018 fast-track designation was granted to the monoclonal antibody margetuximab⁴⁰ in patients with ERBB2 positive breast cancer previously treated with an anti-HER2 therapy. In 2019, fast track designation was granted to the HER2-targeting antibody drug conjugate, amcenestrant⁴¹, for HER2-positive advanced or metastatic breast cancer after one or more prior anti-HER2 based regimens. Additionally, in 2019, the novel bispecific antibody, zanidatamab⁴², received fast-track designation in combination with standard chemotherapy for patients with HER2-overexpressing gastroesophageal adenocarcinoma (GEA) and breakthrough therapy designation (2020) as a monotherapy for patients with HER2-amplified biliary tract cancer⁴³. In 2020, BDTX-189⁴⁴ received fast-track designation for adult patients with solid tumors harboring an allosteric human ERBB2 mutation or exon 20 insertion, and the humanized anti-HER2 antibody drug conjugate disitamab vedotin received breakthrough designation for adult patients with HER2-positive urothelial cancer after previous platinum-chemotherapy treatment⁴⁵. In 2021, the antibody-drug conjugate ARX788⁴⁶ received fast-track designation as a monotherapy for advanced or metastatic HER2-positive breast cancer that have progressed on one or more anti-HER2 regimens. Additionally, in 2021, fast track designation was granted to HER2 targeted chimeric antigen receptor macrophage (CAR-M), CT-0508⁴⁷, for HER2-overexpressing solid tumors. Certain activating mutations have been observed to impart sensitivity to neratinib, afatinib, lapatinib, and trastuzumab, or dacomitinib in early and ongoing clinical studies^{48,49,50,51,52}. ERBB2 kinase domain mutations R896G and V659E both showed response to afatinib in two NSCLC case studies^{53,54}. Additionally, acquired HER2 mutations in estrogen receptor-positive (ER+) breast cancer have been shown to confer resistance to hormone therapy⁵⁵. However, this was shown to be overcome by neratinib in combination with therapies targeting ER⁵⁵.

MYCN (MYCN proto-oncogene, bHLH transcription factor)

Background: The MYCN gene encodes the MYCN proto-oncogene (n-MYC), a basic helix-loop-helix transcription factor. MYCN is a member of the MYC oncogene family that regulates the expression of numerous genes that control cell cycle progression, apoptosis, metabolic pathways, and cellular transformation^{56,57,58,59,60}. MYCN amplification is correlated with failure of cells to arrest in G1 phase

Biomarker Descriptions (continued)

of the cell cycle, leading to uncontrolled proliferation⁶¹. Like MYC, MYCN functions as a heterodimer in complex with the transcription factor MAX^{58,62}.

Alterations and prevalence: Recurrent somatic mutations in MYCN, including codon P44, are observed in approximately 2% of neuroblastoma with the same variants observed in Wilms' tumor^{63,64}. MYCN amplification is found in 20 to 30% of neuroblastoma⁶⁵. Amplification and overexpression of MYCN is also common in other central nervous system cancers as well as prostate cancer, hematologic malignancies, and pancreatic cancer⁶⁶.

Potential relevance: Currently, no therapies are approved for MYCN aberrations. Dysregulation of MYCN is associated with poor prognosis of several pediatric tumor types including neuroblastoma, Wilms' tumor, retinoblastoma, medulloblastoma, and rhabdomyosarcoma⁶¹. In neuroblastoma, increased MYCN signaling is directly related to tumor aggressiveness and increased metastatic potential^{61,67}. Strategies targeting MYCN driven cancers currently focus on targeting MYCN expression, transcription, and synthetic lethality associated with MYCN overexpression⁶⁶.

Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

ERBB2 p.(L755_E757delinsPQ) c.2263_2269delTTGAGGGinsCCTC

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
trastuzumab deruxtecan	×	●	×	×	×
trastuzumab deruxtecan, pembrolizumab, chemotherapy	×	×	×	×	● (III)
ado-trastuzumab emtansine	×	×	×	×	● (II)
SAR-443216	×	×	×	×	● (I)
SHR-A1811	×	×	×	×	● (I)

AR amplification

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
bicalutamide	×	○	×	×	×
leuprorelin	×	○	×	×	×

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Details

Current NCCN Information

- ☒ In this cancer type ☐ In other cancer type ☒ In this cancer type and other cancer types

NCCN information is current as of 2022-06-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

ERBB2 p.(L755_E757delinsPQ) c.2263_2269delTTGAGGGinsCCTC

☒ trastuzumab deruxtecan

Cancer type: Non-Small Cell Lung Cancer

Variant class: ERBB2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Metastatic (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

AR amplification

☐ bicalutamide

Cancer type: Head and Neck Cancer

Variant class: AR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Salivary Gland Neoplasm; Recurrent, Unresectable, Distant Metastases (Line of therapy not specified); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Head and Neck Cancers [Version 2.2022]

☐ leuprorelin

Cancer type: Head and Neck Cancer

Variant class: AR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Salivary Gland Neoplasm; Recurrent, Unresectable, Distant Metastases (Line of therapy not specified); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Head and Neck Cancers [Version 2.2022]

Clinical Trials in Taiwan region:

Clinical Trials Summary

ERBB2 p.(L755_E757delinsPQ) c.2263_2269delTTGAGGGinsCCTC

NCT ID	Title	Phase
NCT05048797	An Open-label, Randomized, Multicenter, Phase III Study to Assess the Efficacy and Safety of Trastuzumab Deruxtecan as First-line Treatment of Unresectable, Locally Advanced, or Metastatic NSCLC Harboring HER2 Exon 19 or 20 Mutations (DESTINY-Lung04)	III
NCT05013554	A Phase I/Ib Open-label, First-in-human, Single Agent, Dose Escalation and Expansion Study for the Evaluation of Safety, Pharmacokinetics, Pharmacodynamics, and Anti-tumor Activity of SAR443216 in Participants with Relapsed/Refractory HER2 Expressing Solid Tumors.	I
NCT04589845	Tumor-Agnostic Precision Immunooncology and Somatic Targeting Rational for You (TAPISTRY) Phase II Platform Trial	II
NCT04446260	A Phase I Multi-Country, Multi-Center, Open-Label Study to Evaluate the Safety, Tolerability, Pharmacokinetics and Efficacy of SHR-A1811 in HER2 Expressing or Mutated Advanced Malignant Solid Tumor Subjects	I

Alerts Informed By Public Data Sources

Current FDA Information

 Contraindicated
  Not recommended
  Resistance
  Breakthrough
  Fast Track

FDA information is current as of 2022-06-15. For the most up-to-date information, search www.fda.gov.

ERBB2 p.(L755_E757delinsPQ) c.2263_2269delTTGAGGGinsCCTC

trastuzumab deruxtecan

Cancer type: Non-Small Cell Lung Cancer

Variant class: ERBB2 mutation

Supporting Statement:

The FDA has granted Breakthrough Designation for the HER2-directed antibody drug conjugate, Enhertu (trastuzumab deruxtecan), for the treatment of HER2 mutated metastatic non-small cell lung cancer (NSCLC) with disease progression on or after platinum-based therapy.

Reference:

<https://www.astrazeneca.com/media-centre/press-releases/2020/enhertu-granted-breakthrough-therapy-designation-in-the-us-for-her2-mutant-metastatic-non-small-cell-lung-cancer.html>

ERBB2 p.(L755_E757delinsPQ) c.2263_2269delTTGAGGGinsCCTC (continued)**A BDTX-189**

Cancer type: Solid Tumor

Variant class: ERBB2 mutation

Supporting Statement:

The FDA has granted Fast Track Designation to BDTX-189 for solid tumors harboring a HER2 mutation or an EGFR or HER2 exon 20 insertion after progression on prior therapy.

Reference:

<https://investors.blackdiamondtherapeutics.com/news-releases/news-release-details/black-diamond-therapeutics-granted-fast-track-designation-fda>

AR amplification**A enobosarm**

Cancer type: Breast Cancer

Variant class: AR positive

Other criteria: ERBB2 negative, ER positive

Supporting Statement:

The FDA has granted Fast Track Designation to enobosarm for AR+/ER+/HER2- in metastatic breast cancer.

Reference:

<https://www.cancernetwork.com/view/fda-grants-fast-track-designation-to-enobosarm-in-ar-er-her2--metastatic-breast-cancer>

A seviteronel

Cancer type: Triple Negative Breast Cancer

Variant class: AR positive

Supporting Statement:

The FDA has granted Fast Track Designation to the small-molecule CYP17 lyase-selective inhibitor, seviteronel, for:

- Androgen receptor (AR) positive advanced triple negative breast cancer (TNBC).
- Estrogen receptor (ER) positive advanced breast cancer.

Reference:

<https://www.businesswire.com/news/home/20160106006206/en/Innocrin-Pharmaceuticals-Granted-Fast-Track-Designation-FDA>

Current NCCN Information

 Contraindicated  Not recommended  Resistance  Breakthrough  Fast Track

NCCN information is current as of 2022-06-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

ERBB2 p.(L755_E757delinsPQ) c.2263_2269delTTGAGGGinsCCTC

afatinib

Cancer type: Non-Small Cell Lung Cancer **Variant class:** ERBB2 mutation

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "The NCCN NSCLC Panel does not recommend single-agent therapy with trastuzumab or afatinib for patients with ERBB2 mutations, because response rates are lower and treatment is less effective with these agents."

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

trastuzumab

Cancer type: Non-Small Cell Lung Cancer **Variant class:** ERBB2 mutation

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "The NCCN NSCLC Panel does not recommend single-agent therapy with trastuzumab or afatinib for patients with ERBB2 mutations, because response rates are lower and treatment is less effective with these agents."

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

References

1. Lu et al. International Union of Pharmacology. LXV. The pharmacology and classification of the nuclear receptor superfamily: glucocorticoid, mineralocorticoid, progesterone, and androgen receptors. *Pharmacol. Rev.* 2006 Dec;58(4):782-97. PMID: 17132855
2. Davey et al. Androgen Receptor Structure, Function and Biology: From Bench to Bedside. *Clin Biochem Rev.* 2016 Feb;37(1):3-15. PMID: 27057074
3. Crumbaker et al. AR Signaling and the PI3K Pathway in Prostate Cancer. *Cancers (Basel).* 2017 Apr 15;9(4). PMID: 28420128
4. Leung et al. Non-Genomic Actions of the Androgen Receptor in Prostate Cancer. *Front Endocrinol (Lausanne).* 2017 Jan 17;8:2. PMID: 28144231
5. Waltering et al. Androgen receptor (AR) aberrations in castration-resistant prostate cancer. *Mol. Cell. Endocrinol.* 2012 Sep 5;360(1-2):38-43. PMID: 22245783
6. Antonarakis et al. AR-V7 and resistance to enzalutamide and abiraterone in prostate cancer. *N. Engl. J. Med.* 2014 Sep 11;371(11):1028-38. PMID: 25184630
7. Zhu et al. Novel Junction-specific and Quantifiable In Situ Detection of AR-V7 and its Clinical Correlates in Metastatic Castration-resistant Prostate Cancer. *Eur. Urol.* 2018 May;73(5):727-735. PMID: 28866255
8. <https://www.cancernetwork.com/view/fda-grants-fast-track-designation-to-enobosarm-in-ar-er-her2-metastatic-breast-cancer>
9. <https://www.businesswire.com/news/home/20160106006206/en/Innocrin-Pharmaceuticals-Granted-Fast-Track-Designation-FDA>
10. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/202379s035lbl.pdf
11. https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/203415s018,213674s005lbl.pdf
12. Lallous et al. Functional analysis of androgen receptor mutations that confer anti-androgen resistance identified in circulating cell-free DNA from prostate cancer patients. *Genome Biol.* 2016 Jan 26;17:10. PMID: 26813233
13. Robinson et al. Integrative clinical genomics of advanced prostate cancer. *Cell.* 2015 May 21;161(5):1215-1228. PMID: 26000489
14. King et al. Amplification of a novel v-erbB-related gene in a human mammary carcinoma. *Science.* 1985 Sep 6;229(4717):974-6. PMID: 2992089
15. Gutierrez et al. HER2: biology, detection, and clinical implications. *Arch. Pathol. Lab. Med.* 2011 Jan;135(1):55-62. PMID: 21204711
16. Di et al. erbB-2 is a potent oncogene when overexpressed in NIH/3T3 cells. *Science.* 1987 Jul 10;237(4811):178-82. PMID: 2885917
17. Hudziak et al. Increased expression of the putative growth factor receptor p185HER2 causes transformation and tumorigenesis of NIH 3T3 cells. *Proc. Natl. Acad. Sci. U.S.A.* 1987 Oct;84(20):7159-63. PMID: 2890160
18. Lonardo et al. The normal erbB-2 product is an atypical receptor-like tyrosine kinase with constitutive activity in the absence of ligand. *New Biol.* 1990 Nov;2(11):992-1003. PMID: 1983208
19. Ciriello et al. Comprehensive Molecular Portraits of Invasive Lobular Breast Cancer. *Cell.* 2015 Oct 8;163(2):506-19. PMID: 26451490
20. Cancer Genome Atlas Research Network. Comprehensive molecular characterization of gastric adenocarcinoma. *Nature.* 2014 Sep 11;513(7517):202-9. doi: 10.1038/nature13480. Epub 2014 Jul 23. PMID: 25079317
21. Cancer Genome Atlas Research Network. Comprehensive molecular characterization of urothelial bladder carcinoma. *Nature.* 2014 Mar 20;507(7492):315-22. doi: 10.1038/nature12965. Epub 2014 Jan 29. PMID: 24476821
22. Donna et al. Comprehensive molecular characterization of human colon and rectal cancer. *Nature.* 2012 Jul 18;487(7407):330-7. PMID: 22810696
23. Cancer Genome Atlas Research Network. Comprehensive molecular profiling of lung adenocarcinoma. *Nature.* 2014 Jul 31;511(7511):543-50. doi: 10.1038/nature13385. Epub 2014 Jul 9. PMID: 25079552
24. Cancer Genome Atlas Research Network. Integrated genomic analyses of ovarian carcinoma. *Nature.* 2011 Jun 29;474(7353):609-15. PMID: 21720365
25. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
26. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
27. Petrelli et al. Clinical and pathological characterization of HER2 mutations in human breast cancer: a systematic review of the literature. *Breast Cancer Res. Treat.* 2017 Nov;166(2):339-349. PMID: 28762010

References (continued)

28. Bose et al. Activating HER2 mutations in HER2 gene amplification negative breast cancer. *Cancer Discov.* 2013 Feb;3(2):224-37. doi: 10.1158/2159-8290.CD-12-0349. Epub 2012 Dec 7. PMID: 23220880
29. Hudis. Trastuzumab--mechanism of action and use in clinical practice. *N. Engl. J. Med.* 2007 Jul 5;357(1):39-51. PMID: 17611206
30. Slamon et al. Human breast cancer: correlation of relapse and survival with amplification of the HER-2/neu oncogene. *Science.* 1987 Jan 9;235(4785):177-82. PMID: 3798106
31. https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/103792s5345lbl.pdf
32. https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/125409s124lbl.pdf
33. https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/125427s111lbl.pdf
34. NCCN Guidelines® - NCCN-Breast Cancer [Version 3.2022]
35. https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/022059s031lbl.pdf
36. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/208051s009lbl.pdf
37. https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/213411s000lbl.pdf
38. https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/125514s123lbl.pdf
39. <https://www.globenewswire.com/news-release/2016/06/01/845166/0/en/Galena-Biopharma-Receives-Fast-Track-Designation-for-NeuVax-nelipepimut-S-PRESENT-Clinical-Trial.html>
40. https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/761150s000lbl.pdf
41. <https://www.prnewswire.com/news-releases/fda-grants-arx788-fast-track-designation-for-her2-positive-metastatic-breast-cancer-301199951.html>
42. <https://www.targetedonc.com/view/her2targeted-antibody-zw25-earns-fda-fast-track-designation-in-gea>
43. <https://www.targetedonc.com/view/fda-grants-breakthrough-designation-to-zanidatamab-for-her2-amplified-biliary-tract-cancer>
44. <https://investors.blackdiamondtherapeutics.com/news-releases/news-release-details/black-diamond-therapeutics-granted-fast-track-designation-fda>
45. <https://www.prnewswire.com/news-releases/remegen-announces-us-fda-has-granted-breakthrough-therapy-designation-for-disitamab-vedotin-rc48-in-urothelial-cancer-301138315.html>
46. <http://ambrx.com/fda-grants-arx788-fast-track-designation-for-her2-positive-metastatic-breast-cancer>
47. <https://www.prnewswire.com/news-releases/carisma-therapeutics-announces-us-food-and-drug-administration-grants-fast-track-designation-to-ct-0508-for-the-treatment-of-patients-with-solid-tumors-301381843.html>
48. Ma et al. Neratinib Efficacy and Circulating Tumor DNA Detection of HER2 Mutations in HER2 Nonamplified Metastatic Breast Cancer. *Clin. Cancer Res.* 2017 Oct 1;23(19):5687-5695. PMID: 28679771
49. De et al. Clinical activity of afatinib (BIBW 2992) in patients with lung adenocarcinoma with mutations in the kinase domain of HER2/neu. *Lung Cancer.* 2012 Apr;76(1):123-7. PMID: 22325357
50. Kris et al. Targeting HER2 aberrations as actionable drivers in lung cancers: phase II trial of the pan-HER tyrosine kinase inhibitor dacomitinib in patients with HER2-mutant or amplified tumors. *Ann. Oncol.* 2015 Jul;26(7):1421-7. PMID: 25899785
51. Falchook et al. Non-small-cell lung cancer with HER2 exon 20 mutation: regression with dual HER2 inhibition and anti-VEGF combination treatment. *J Thorac Oncol.* 2013 Feb;8(2):e19-20. PMID: 23328556
52. David et al. Neratinib in HER2- or HER3-mutant solid tumors: SUMMIT, a global, multi-histology, open-label, phase 2 'basket' study. AACR 2017. Abstract CT001
53. Lin et al. Response to Afatinib in a Patient with Non-Small Cell Lung Cancer Harboring HER2 R896G Mutation: A Case Report. *Onco Targets Ther.* 2019;12:10897-10902. PMID: 31849493
54. Chang et al. Sustained Partial Response to Afatinib in a Patient With Lung Adenocarcinoma Harboring HER2V659E Mutation. *JCO Precis Oncol.* 2020 Aug; 912-915. DOI: 10.1200/PO.20.00114
55. Nayar et al. Acquired HER2 mutations in ER+ metastatic breast cancer confer resistance to estrogen receptor-directed therapies. *Nat. Genet.* 2019 Feb;51(2):207-216. PMID: 30531871
56. Dang et al. The c-Myc target gene network. *Semin. Cancer Biol.* 2006 Aug;16(4):253-64. PMID: 16904903
57. Chen et al. Targeting oncogenic Myc as a strategy for cancer treatment. *Signal Transduct Target Ther.* 2018 Feb 23;3:5. doi: 10.1038/s41392-018-0008-7. eCollection 2018. PMID: 29527331
58. Dang. MYC on the path to cancer. *Cell.* 2012 Mar 30;149(1):22-35. PMID: 22464321
59. Dominguez-Sola et al. Non-transcriptional control of DNA replication by c-Myc. *Nature.* 2007 Jul 26;448(7152):445-51. PMID: 17597761

References (continued)

60. Wahlström et al. Impact of MYC in regulation of tumor cell metabolism. *Biochim. Biophys. Acta*. 2015 May;1849(5):563-9. PMID: 25038584
61. Ruiz-Pérez et al. The MYCN Protein in Health and Disease. *Genes (Basel)*. 2017 Mar 30;8(4). PMID: 28358317
62. Blackwood et al. Myc and Max function as a nucleoprotein complex. *Curr. Opin. Genet. Dev.* 1992 Apr;2(2):227-35. PMID: 1638116
63. Pugh et al. The genetic landscape of high-risk neuroblastoma. *Nat. Genet.* 2013 Mar;45(3):279-84. PMID: 23334666
64. Williams et al. Multiple mechanisms of MYCN dysregulation in Wilms tumour. *Oncotarget*. 2015 Mar 30;6(9):7232-43. PMID: 25749049
65. Huang et al. Neuroblastoma and MYCN. *Cold Spring Harb Perspect Med*. 2013 Oct 1;3(10):a014415. PMID: 24086065
66. Rickman et al. The Expanding World of N-MYC-Driven Tumors. *Cancer Discov*. 2018 Feb;8(2):150-163. PMID: 29358508
67. Seeger et al. Association of multiple copies of the N-myc oncogene with rapid progression of neuroblastomas. *N Engl J Med*. 1985 Oct 31;313(18):1111-6. PMID: 4047115